The ancestral shape of the access proton path of mitochondrial ATP synthases revealed by a split subunit-a

Jonathan E. Wong[1,2], Alena Zíková[1,2], Ondřej Gahura[1*]

[1]Institute of Parasitology, Biology Centre, Czech Academy of Sciences, 37005 České Budějovice, Czech Republic

[2]Faculty of Science, University of South Bohemia, 37005 České Budějovice, Czech Republic

[*]Corresponding author: gahura@paru.cas.cz

Abstract

The passage of protons across membranes through F_1F_o-ATP synthases spins their rotors and drives synthesis of ATP. While the principle of torque generation by proton transfer is known, the mechanisms and routes of proton access and release and their evolution are not fully understood. Here, we show that the entry site and path of protons in the lumenal half-channel of mitochondrial ATP synthases are largely defined by a short N-terminal α-helix of subunit-a. In *Trypanosoma brucei* and other Euglenozoa, the α-helix is part of another polypeptide chain that is a product of subunit-a gene fragmentation. This α-helix and other elements forming the proton pathway are widely conserved across eukaryotes and in Alphaproteobacteria, the closest extant relatives of mitochondria, but not in other bacteria. Thus, the proton entry site, the shape of the access half-channel, and the proton transfer mechanism involving a chain of ordered water molecules, predate eukaryotes and possibly originated in the lineage from which mitochondria evolved by endosymbiosis.

1 Introduction

F_1F_o-ATP synthases are rotary molecular machines that convert the proton-motive force across

energy-transducing membranes of bacteria, chloroplasts and mitochondria to the chemical

energy of ATP. The flow of protons down their electrochemical gradient through the

membrane-intrinsic F_o subcomplex spins the enzyme's asymmetric rotor, whose rotation

changes the conformation of three catalytic sites on the membrane-extrinsic F_1-ATPase

subcomplex, resulting in synthesis of ATP. The transmembrane part of the rotor, the c-ring,

consists of several copies of subunit-c. Proton translocation is based on sequential protonation

of a glutamate residue of each protomer of the c-ring, rotation of the c-ring with the neutralized

glutamates exposed to the hydrophobic environment of the phospholipid bilayer, and release of

protons on the other side of the membrane. The second element crucial for proton passage is

the stator subunit-a (also referred to as ATP6 or subunit-6). Canonical subunits-a contain six α-

helices (h_1-h_6), five of which are largely embedded in the membrane. While h_1 is a vertical

transmembrane helix, h_3-h_6 are horizontal, positioned at an angle ~20° relative to the plane of

the membrane. Helices h_5-h_6 form a hairpin bending around the c-ring. The sites of proton

binding and release from the c-ring glutamates are separated by a universally conserved

arginine of h_5. The positively charged arginine prevents protons from leaking without c-ring

rotation and induces their dissociation at the release site (reviewed in Walker 2017; Kuhlbrandt

2019; Guo and Rubinstein 2022). The proton-binding glutamates on the c-ring are accessible

from the surface of the complex by two offset solvent-filled half-channels. In mitochondrial

ATP synthases, the access half-channel opens into the lumen of cristae, hence it is called the

lumenal half-channel. It is shaped by subunit-a and adjacent membrane elements, including

lipids (Klusch, et al. 2017). Recently, it was proposed that delivery of protons to the c-ring is

facilitated by a Grotthuss mechanism through a chain of coordinated water molecules and

adjacent amino acid residues in the inner part of the channel in mammalian (Spikes, et al. 2020)

and trypanosomal (Gahura, et al. 2022) mitochondrial ATP synthases.

Here, we explore the lumenal half-channel forming elements across species and show that

the N-terminal region of subunits-a form a short α-helix common to alphaproteobacterial and

vast majority of mitochondrial ATP synthases. In Euglenozoa, this helix is a part of a different

polypeptide chain encoded by a nuclear gene that is presumably the product of fragmentation

of the mitochondrial gene for subunit-a. We document that the helix defines the proton entry

site and contributes to the shaping of the inner part of the half-channel in modern eukaryotes

and their common ancestors.

Results and discussion

The short N-terminal α-helix of subunit-a is a widely conserved structural element of

mitochondrial ATP synthases

When examining the proton translocation elements of mitochondrial ATP synthases of two

divergent eukaryotes, *Saccharomyces cerevisiae* (Guo, et al. 2017) and *Trypanosoma brucei*

(Gahura, et al. 2022), we noticed a remarkable similarity between the N-terminal regions of

subunit-a in the yeast and subunit ATPEG3 in the parasite. The N-termini of both proteins form

a short single-turn α-helix, hereafter referred to as h_N, which is located at the layer

corresponding to the lumenal surface of the membrane and interacts with the horizontal helix

h_5 of subunit-a in the same manner in both structures. Downstream of h_N, yeast subunit-a and

trypanosomal ATPEG3 follow a similar path; however, after ten residues the yeast chain turns

back towards the first transmembrane helix h_1, thereby diverging from ATPEG3, which

continues to the periphery of the complex. The canonical trypanosomal subunit-a, which

contains h_1 and four characteristic horizontal membrane-intrinsic helices h_3-h_6, begins nine

amino acid residues upstream of h_1 in close proximity to the site of divergence between yeast

subunit-a and ATPEG3 (Fig. 1A). Based on the striking structural and sequence (see below) similarity, we conclude that subunit-a in *T. brucei* has been split and ATPEG3 represents its N-terminal fragment. To acknowledge the origin of ATPEG3, we introduce an alternative name subunit-a_N.

The split of subunit-a in *T. brucei* occurred in the region corresponding to the loop involved in ATP synthase dimerization in *S. cerevisiae* (Guo, et al. 2017). In *T. brucei*, the loop is structurally replaced by a two-stranded β-sheet, formed by the N-terminus of subunit-a and a β-strand of subunit-a_N (Fig. 1A). Unlike in yeast, this structure does not contribute to the ATP synthase dimerization interface, which is provided by subunits g and e in this organism (Gahura, et al. 2022). After splitting, subunit-a_N was extended at its C-terminus. The extension interacts exclusively with lineage-specific subunits (ATPTB1, ATPTB6, ATPTB11, ATPTB12) or with lineage-specific extensions of subunits shared with other ATP synthases (subunit-i/j and k; Gahura, et al. 2022). It extends to the periphery of the membrane part of the F_o-moiety, where it co-constitutes a euglenozoan-specific subcomplex, separated from the conserved part by a lipid-filled cavity (Muhleip, et al. 2019; Gahura, et al. 2022). Consistently, fragmented subunit-a is also discernible in the structure of ATP synthase from *Euglena gracilis* (Muhleip, et al. 2019; Gahura, et al. 2021).

The sequence of h_N and flanking residues in yeast is highly similar to the N-terminal sequence of subunit-a in Rickettsiales and other Alphaproteobacteria (Fig. 2, Supplementary table S1 and Supplementary data 1), the closest extant relatives to mitochondria (Roger, et al. 2017). We predicted the structure of subunit-a from *Rickettsia rickettsii* using AlphaFold (Jumper, et al. 2021). The model resembles structures of bacterial and mitochondrial homologs resolved experimentally and includes h_N located at exactly the same position as the h_N in *T. brucei* and *S. cerevisiae* (Fig. 1C). Similarity of this region in ATP synthases from mitochondria of two deeply branching eukaryotic lineages and the group of bacteria from which mitochondria

evolved strongly indicates that the observed architecture of the N-terminal segment of subunit-

a is ancestral.

We compared subunit-a sequences from 88 species covering all major eukaryotic

lineages, and structures from selected species determined by cryoEM or predicted by

AlphaFold. The sequence comparison documented that the split of subunit-a observed in *T.*

brucei and *E. gracilis* is restricted to Euglenozoa, because subunits-a from all representatives

of the sister lineage Heterolobosea contain a conserved sequence of h_N. Further, the analysis

revealed high sequence similarity in the N-terminal region, with most sequences showing a

degenerate sequence motif SPLEQF (Fig. 2, Supplementary table S1), supporting the notion

that presence of h_N represents an ancestral state. The only organisms without the SPLEQF motif

were most alveolates, chlorophycean algae *Polytomella* and *Chlamydomonas*, and all

metazoans (Fig. 2, Supplementary table S1). In alveolates and chlorophycean algae, the N-

terminal region including h_1 is divergent or absent, as documeted by cryoEM structures of ATP

synthases from *Tetrahymena* (Flygaard, et al. 2020), *Polytomella* (Murphy, et al. 2019) and

Toxoplasma (Muhleip, et al. 2021). Subunit-a in *Tetrahymena* features a highly extended and

structurally divergent N-terminus, which interacts predominantly with ciliate-specific ATP

synthase components. On the contrary, subunit-a in *Polytomella* lacks the entire N-terminal

region including the transmembrane helix h_1, while subunit-a in *Toxoplasma* is even more

reduced and contains horizontal helices h_5 and h_6 only (Fig. 1C). The only alveolate with a

canonical N-terminus of subunit-a is *Colponema* (Fig. 2), which supports the view that it

represents a lineage sister to all other Alveolata (Tikhonenkov, et al. 2020).

In Metazoa, the N-terminal sequence of subunit-a differs from h_N in other groups, but

invariably contains the motif FxxF (Fig. 2 and Supplementary table S1 and Supplementary Data

1). In experimentally determined and predicted structures of mammalian (Pinke, et al. 2020;

Spikes, et al. 2020) and *Drosophila* subunit-a, respectively, this motif adopts the canonical fold

of h_N (Fig. 1C), indicating that the ancestral architecture is retained despite sequence divergence. The only exception is the group Nematoda, represented by *Caenorhabditis elegans*, whose subunit-a begins with helix h_1, like subunit-a in trypanosomes. This notable exception is in line with a recent cryoelectron tomography study that showed that the nematode ATP synthase diverged architecturally from canonical ATP synthases from yeasts and mammals and might lack subunits 8, k and i/j (Buzzard, et al. 2023), which are otherwise widely conserved in eukaryotes (Sinha and Wideman 2022). Due to high conservancy of h_N across eukaryotes, we wondered whether an equivalent of h_N could be retained in nematodes, possibly as a product of subunit-a gene split analogous to that observed in *T. brucei*. Such evolutionary independent event of fission of a mitochondrial gene would not be unprecedented. The gene encoding the mitoribosomal subunit uS3m has been fragmented in Ciliata (Swart, et al. 2012; Tobiasson and Amunts 2020) and some Heterolobosea (Fu, et al. 2014), and the gene encoding Cox2, a subunit of cytochrome oxidase, has been independently split at least three times in evolution, specifically in alveolates, chlorophycean algae (Perez-Martinez, et al. 2001; Waller and Keeling 2006; Rodriguez-Salinas, et al. 2012), and the hymenopteran insect *Campsomeris* (Szafranski 2017a). Therefore, we searched the proteomes of Nematoda using several approaches (blast, motif search, structure similarity search in AlphaFold database) to identify the protein constituting the missing region including h_N, but we found no candidate, presumably due to the short length of the region and limited sequence conservancy.

In summary, the vast majority of eukaryotes with the exception of most Alveolata and a group of chlorophycean algae have retained the ancestral architecture of the region corresponding to the N-terminus of subunit-a.

Loss of transmembrane helix h_1 predetermines the transfer of subunit-a from the mitochondrial to the nuclear genome

While the gene encoding subunit-a_N in Euglenozoa was transferred to the nuclear genome, the

subunit-a gene has been retained in the mitochondrial genome. Likewise, subunit-a in

nematodes, also starting with the first transmembrane helix h_1, is encoded mitochondrially. In

contrast, the gene of subunit-a in *Polytomella* and *Chlamydomonas*, which lacks h_1, has been

transferred to the nuclear genome. This suggests that while loss of h_1 allows the transfer of the

gene from the mitochondrial to the nuclear genome, the removal of the N-terminal fragment

upstream of h_1 is not sufficient for the relocation. Consistently, the reduced subunit-a in

Toxoplasma and related species are also nuclear encoded (Huet, et al. 2018; Muhleip, et al.

2021; Fig. 2). Together, these observations indicate that barriers associated with the insertion

of h_1-containing subunit-a to the inner mitochondrial membrane from the intermembrane space

(or with translocation of the protein into mitochondria prior its insertion from inside) may be a

factor limiting the transfer of the subunit-a gene from mitochondrial DNA to the nuclear

genome. We expanded a previous analysis of hydropathy of subunit-a (Muhleip, et al. 2021) to

all species included in our dataset. It revealed that some subunits-a (e.g., from *Nicotiana* or

Tetrahymena) retained in mitochondrial genomes show relatively low hydrophobicity similar

to those transferred to the nuclear genome (Fig. 2, Supplementary table S1), suggesting that the

reduced hydrophobicity is not the only prerequisite for the intracellular subunit-a gene transfer.

Regardless of the factor that allowed the subunit-a gene to be transferred, this is very rare event

that has most likely occurred only twice in the evolution of eukaryotes described to date, namely

in the ancestors of the present-day myzozoans and Chlamydomonadales, an order of

chlorophycean algae including *Polytomella* and *Chlamydomonas* (Fig. 2).

The entry point of the proton access half-channel in mitochondrial and

alphaproteobacterial ATP synthases is defined by the N-terminal helix h_N of subunit-a

The wide conservancy of h_N and its flanking residues (Fig. 2), whether within subunit-a or as

part of a fission product, implies the functional importance of this region. Indeed, h_N contributes

to the lining of the inner constricted portion of the lumenal proton half-channel in structurally

characterized ATP synthases from *Trypanosoma* (Gahura, et al. 2022), *S. cerevisiae* (Guo, et

al. 2017) and mammals (Pinke, et al. 2020; Spikes, et al. 2020; Fig. 1B) and, as can be inferred

from its conserved nature, presumably also in most other eukaryotic lineages and

Alphaproteobacteria. Thus, the segment of the half-channel proximal to the c-ring most likely

does not differ between these organisms. Indeed, several residues proposed to be involved

directly in the proton transfer or to coordinate ordered proton-transporting water molecules

(Spikes, et al. 2020; Gahura, et al. 2022) are largely conserved in the h_N-containing subunits

including those from Alphaproteobacteria. These include a protonatable glutamate or aspartate

residue in helix h_6 (Glu203, Glu233 and Asp202 in bovine, *S. cerevisiae* and *T. brucei*

numbering, respectively) and a histidine in h_5 (His169/195/155 in bovine/*S. cerevisiae*/*T.*

brucei; Fig. 2, Supplementary table S1). In addition, the position of histidine 19 in h_N in *T.*

brucei subunit-a_N, which coordinates one of the molecules in the Grotthuss water chain, is

occupied by another hydrogen bonding residue, glutamine, in subunits-a of vast majority of

mitochondrial ATP synthases (Gln15 in *S. cerevisiae*; Fig. 1B&2 and Supplementary table S1).

Therefore, we reason that the Grotthuss mechanism represents an ancestral mode of proton

transfer in mitochondrial ATP synthases that has been conserved in most, if not all, eukaryotic

lineages.

Notably, ATP synthase in bacteria from various groups outside Alphaproteobacteria

(*Escherichia coli;* Sobti, et al. 2019, *Mycobacterium*; Guo, et al. 2021, Montgomery, et al. 2021,

and *Acinetobacter*; Demmer, et al. 2022) contain an α-helix at the N-terminus of the subunit-a,

which is longer and sequentially dissimilar to h_N in mitochondrial ATP synthases and

positionally shifted further from the loop connecting helices h_3 and h_4 (Fig. 1C). As proposed

based on the structure of the *Acinetobacter* ATP synthase, the position of the N-terminal helix

determines whether protons enter to the half-channel using the "front" or "rear" entry site

(Demmer, et al. 2022). In *Acinetobacter* and some other bacteria, the protons enter from the

side facing the c-ring through the "front" entry located between the N-terminal helix and h_5 of

subunit-a. In yeast and bovine mitochondrial ATP synthases protons use the "rear" opening, a

cleft between h_N and the C-terminal tail of subunit-f (Spikes, et al. 2020; Demmer, et al. 2022;

Fig. 3A,B). In the latter two structures, as well as in the structure from *T. brucei* and the

predicted structure from *Rickettsia*, the position of the front entry is blocked by h_N (Fig 3B,C),

specifically by the phenylalanine residue of the SPLEQF motif that is nearly invariably

conserved in all h_N sequences included in our analysis (the only exception is *Fonticula*, where

it is substituted by other bulky and aromatic residue, tyrosine; Fig 2 and Supplementary table

S1). Therefore, we reason that the rear entry is common to all h_N-containing ATP synthases.

Taken together, h_N contributes to shaping of both the surface-exposed and the inner portions of

the access proton half-channel, underlying its almost universal occurrence in mitochondrial

ATP synthases.

Concluding remarks

By combination of analyses of available ATP synthase structures with sequence comparison

and structure prediction across species, we documented that the proton access path and proton

transfer mechanism in the access half-channel is conserved in majority of mitochondrial ATP

synthases. The key structural determinant, so far largely overlooked, is h_N of subunit-a, which

defines the entry to the access half channel and contributes to hydrogen bonding of water

molecules in the Grotthuss chain (Fig. 1C). Although the shape of the inner access half-channel

in bacteria resembles that of mitochondrial ATP synthase (Montgomery, et al. 2021), water

molecules have not been resolved in cryoEM maps of bacterial ATP synthases. In addition, the

key residues involved in water coordination and proton transfer are not present in bacteria

outside Alphaproteobacteria (columns H and D/E, Fig. 2). Therefore, the mechanism of the

proton transfer to the c-ring in bacterial complexes has not been unambiguously determined

yet. We speculate that the use of Grotthuss mechanism for the proton transfer emerged in

Alphaproteobacteria and may represent a preadaptation to eukaryogenesis.

204   Uniquely, F_1F_o-ATP synthase in *T. brucei* mitochondria contains two functionally central

components, the subunits α and a, that are both constituted of two separate polypeptide chains.

While the subunit-α is cleaved into two fragments posttranslationally (Gahura, et al. 2018;

Montgomery, et al. 2018), the subunit-a is split on the gene level. Our finding of a split subunit-

a in *T. brucei* complements recently reported cases of mitochondrial gene fragmentation

revealed by structures of mitochondrial ribosomes (uS4m and uS7m in *Polytomella*; Tobiasson,

et al. 2022) and respiratory chain supercomplexes (Cox3 and Nad5 in *Tetrahymena*; Mühleip,

et al. 2022; Zhou, et al. 2022), and several occurrences reported earlier (reviewed in Szafranski

2017b). The transfer of the same gene from the mitochondrial to the nuclear genome

independently in several lineages facilitates the inference of properties of encoded proteins that

are incompatible with such events. We inferred that the transmembrane helix h_1 might represent

such an obstacle in the case of subunit-a.

Material and Methods

Subunit-a sequences were searched by hmmsearch using the alignment of subunit-a sequences

from *Bos taurus*, *Saccharomyces cerevisiae*, *Rickettsia rickettsii* and *Colponema vietnamica* as

a query. The sequences from Ciliata and Apicomplexa were searched by phmmer using

sequences from *Tetrahymena* and *Toxoplasma*, respectively, as queries. Both searches were

performed on the HMMER web server (Potter, et al. 2018) in the Uniprot database. Amino acid

sequence alignments were constructed by MUSCLE (Edgar 2004) and corrected manually

using available structures as references. Structures were predicted by AlphaFold (Jumper, et al.

2021), and superposed by the Matchmaker tool and visualized in ChimeraX (Goddard, et al.

2018). Hydrophobicity was calculated as grand average of hydropathy using hydropathy indices

from Kyte and Doolittle 1982.

Acknowledgements

230 This work was supported by Czech Science Foundation grant 20-04150Y to O.G. and by

231 European Regional Development Fund (ERDF) and Ministry of Education, Youth and Sport

232 (MEYS) project CZ.02.1.01/0.0/0.0/16_019/0000759 and by the European Research Council

233 (ERC) under the European Union's Horizon 2020 Research and Innovation Program (grant

234 agreement no. 101044951) to A.Z.

Author contributions

237 Conceptualization: O.G. and A.Z.; investigation and data analyses: J.E.W. and O.G.; writing –

238 original draft: O.G.; writing – review and editing: J.E.W, A.Z., O.G.; funding acquisition: O.G.

239 and A.Z.

Declaration of interests

242 They authors declare they have no competing interests.

References

Burki F, Roger AJ, Brown MW, Simpson AGB. 2020. The New Tree of Eukaryotes. Trends
Ecol Evol 35:43-55.

Buzzard E, McLaren M, Bragoszewski P, Brancaccio A, Ford H, Daum B, Kuwabara P,
Collinson I, Gold, VAM, unpublished data,
https://www.biorxiv.org/content/10.1101/2023.02.02.526626v2

Demmer JK, Phillips BP, Uhrig OL, Filloux A, Allsopp LP, Bublitz M, Meier T. 2022.
Structure of ATP synthase from ESKAPE pathogen *Acinetobacter baumannii*. Sci Adv
8:eabl5966.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res 32:1792-1797.

Flygaard RK, Muhleip A, Tobiasson V, Amunts A. 2020. Type III ATP synthase is a symmetry-
deviated dimer that induces membrane curvature through tetramerization. Nat Commun
11:5342.

Fu CJ, Sheikh S, Miao W, Andersson SG, Baldauf SL. 2014. Missing genes, multiple ORFs,
and C-to-U type RNA editing in *Acrasis kona* (Heterolobosea, Excavata) mitochondrial DNA.
Genome Biol Evol 6:2240-2257.

Gahura O, Hierro-Yap C, Zikova A. 2021. Redesigned and reversed: architectural and
functional oddities of the trypanosomal ATP synthase. Parasitology 148:1151-1160.

Gahura O, Muhleip A, Hierro-Yap C, Panicucci B, Jain M, Hollaus D, Slapnickova M, Zikova
A, Amunts A. 2022. An ancestral interaction module promotes oligomerization in divergent
mitochondrial ATP synthases. Nat Commun 13:5989.

Gahura O, Subrtova K, Vachova H, Panicucci B, Fearnley IM, Harbour ME, Walker JE, Zikova
A. 2018. The F_1-ATPase from *Trypanosoma brucei* is elaborated by three copies of an
additional p18-subunit. FEBS J 285:614-628.

Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, Ferrin TE. 2018.
UCSF ChimeraX: Meeting modern challenges in visualization and analysis. Protein Sci 27:14-
25.

Guo H, Bueler SA, Rubinstein JL. 2017. Atomic model for the dimeric F_o region of
mitochondrial ATP synthase. Science 358:936-940.

Guo H, Courbon GM, Bueler SA, Mai J, Liu J, Rubinstein JL. 2021. Structure of mycobacterial
ATP synthase bound to the tuberculosis drug bedaquiline. Nature 589:143-147.

Guo H, Rubinstein JL. 2022. Structure of ATP synthase under strain during catalysis. Nat
Commun 13:2232.

Huet D, Rajendran E, van Dooren GG, Lourido S. 2018. Identification of cryptic subunits from
an apicomplexan ATP synthase. Elife 7.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates
R, Zidek A, Potapenko A, et al. 2021. Highly accurate protein structure prediction with
AlphaFold. Nature 596:583-589.

Klusch N, Murphy BJ, Mills DJ, Yildiz O, Kuhlbrandt W. 2017. Structural basis of proton
translocation and force generation in mitochondrial ATP synthase. Elife 6.

Kuhlbrandt W. 2019. Structure and Mechanisms of F-Type ATP Synthases. Annu Rev Biochem
88:515-549.

Kyte J, Doolittle RF. 1982. A simple method for displaying the hydropathic character of a
protein. J Mol Biol 157:105-132.

Montgomery MG, Gahura O, Leslie AGW, Zikova A, Walker JE. 2018. ATP synthase from
Trypanosoma brucei has an elaborated canonical F_1-domain and conventional catalytic sites.
Proc Natl Acad Sci U S A 115:2102-2107.

Montgomery MG, Petri J, Spikes TE, Walker JE. 2021. Structure of the ATP synthase from
Mycobacterium smegmatis provides targets for treating tuberculosis. Proc Natl Acad Sci U S A
118.

Mühleip A, Flygaard RK, Haapanen O, Baradaran R, Gruhl T, Tobiasson V, Maréchal A,
Sharma   V,   Amunts   A,   unpublished   data,
https://www.biorxiv.org/content/10.1101/2022.06.26.497646v2

Muhleip A, Kock Flygaard R, Ovciarikova J, Lacombe A, Fernandes P, Sheiner L, Amunts A.
2021. ATP synthase hexamer assemblies shape cristae of *Toxoplasma mitochondria*. Nat
Commun 12:120.

Muhleip A, McComas SE, Amunts A. 2019. Structure of a mitochondrial ATP synthase with
bound native cardiolipin. Elife 8.

Murphy BJ, Klusch N, Langer J, Mills DJ, Yildiz O, Kuhlbrandt W. 2019. Rotary substates of
mitochondrial ATP synthase reveal the basis of flexible F_1-F_o coupling. Science 364.

Perez-Martinez X, Antaramian A, Vazquez-Acevedo M, Funes S, Tolkunova E, d'Alayer J,
Claros MG, Davidson E, King MP, Gonzalez-Halphen D. 2001. Subunit II of cytochrome c
oxidase in Chlamydomonad algae is a heterodimer encoded by two independent nuclear genes.
J Biol Chem 276:11302-11309.

Pinke G, Zhou L, Sazanov LA. 2020. Cryo-EM structure of the entire mammalian F-type ATP
synthase. Nat Struct Mol Biol 27:1077-1085.

Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. 2018. HMMER web server: 2018
update. Nucleic Acids Res 46:W200-W204.

Rodriguez-Salinas E, Riveros-Rosas H, Li Z, Fucikova K, Brand JJ, Lewis LA, Gonzalez-
Halphen D. 2012. Lineage-specific fragmentation and nuclear relocation of the mitochondrial
cox2 gene in chlorophycean green algae (Chlorophyta). Mol Phylogenet Evol 64:166-176.

Roger AJ, Munoz-Gomez SA, Kamikawa R. 2017. The Origin and Diversification of
Mitochondria. Curr Biol 27:R1177-R1192.

Sinha   SD,   Wideman   JG.,   unpublished   data,
https://www.biorxiv.org/content/10.1101/2022.09.13.506888v1.full

Sobti M, Ishmukhametov R, Bouwer JC, Ayer A, Suarna C, Smith NJ, Christie M, Stocker R,
Duncan TM, Stewart AG. 2019. Cryo-EM reveals distinct conformations of *E. coli* ATP
synthase on exposure to ATP. Elife 8.

Spikes TE, Montgomery MG, Walker JE. 2020. Structure of the dimeric ATP synthase from
bovine mitochondria. Proc Natl Acad Sci U S A 117:23519-23526.

Swart EC, Nowacki M, Shum J, Stiles H, Higgins BP, Doak TG, Schotanus K, Magrini VJ,
Minx P, Mardis ER, et al. 2012. The *Oxytricha trifallax* mitochondrial genome. Genome Biol
Evol 4:136-154.

Szafranski P. 2017a. Evolutionarily recent, insertional fission of mitochondrial cox2 into
complementary genes in bilaterian Metazoa. BMC Genomics 18:269.

Szafranski P. 2017b. Intercompartmental Piecewise Gene Transfer. Genes (Basel) 8.

Tikhonenkov DV, Mikhailov KV, Gawryluk RMR, Belyaev AO, Mathur V, Karpov SA,
Zagumyonnyi DG, Borodina AS, Prokina KI, Mylnikov AP, et al. 2022. Microbial predators
form a new supergroup of eukaryotes. Nature 612:714-719.

Tikhonenkov DV, Strassert JFH, Janouskovec J, Mylnikov AP, Aleoshin VV, Burki F, Keeling
PJ. 2020. Predatory colponemids are the sister group to all other alveolates. Mol Phylogenet
Evol 149:106839.

Tobiasson V, Amunts A. 2020. Ciliate mitoribosome illuminates evolutionary steps of
mitochondrial translation. Elife 9.

Tobiasson V, Berzina I, Amunts A. 2022. Structure of a mitochondrial ribosome with
fragmented rRNA in complex with membrane-targeting elements. Nat Commun 13:6132.

Walker JE. 2017. Structure, Mechanism and Regulation of ATP Synthases. In: Wikstrom M,
editor: The Royal Society of Chemistry.

Waller RF, Keeling PJ. 2006. Alveolate and chlorophycean mitochondrial cox2 genes split
twice independently. Gene 383:33-37.

Zhou L, Maldonado M, Padavannil A, Guo F, Letts JA. 2022. Structures of Tetrahymena's
respiratory chain reveal the diversity of eukaryotic core metabolism. Science 376:831-839.

 Figures and legends

Figure 1. Split subunit-a of mitochondrial ATP synthase in *Trypanosoma brucei* retains

ancestral structure and shapes the access proton half-channel (**A**) Comparison of the

structures of subunit-a in *S. cerevisiae* and subunits a and a_N in *T. brucei*. The c-rings are shown

as transparent surface. In the structure from *S. cerevisiae*, a fragment of the subunit-a from the

other ATP synthase protomer is shown (su-a'). (**B**) The close-up view of the structures in (A).

Conserved residues of h_N and other selected conserved residues are shown as sticks. The

residues labeled in red are directly involved in proton translocation or coordinate water

molecules in the Grotthus chain. Water molecules inferred from cryoEM density are shown as

yellow balls in the *T. brucei* structure. (**C**) Structures of subunit-a of mitochondrial and bacterial

ATP synthases from selected species. The structure from *Rickettsia* was predicted by AlphaFold

Figure 2. Helix h$_N$ is conserved in most mitochondrial ATP synthases and Alphaproteobacteria. For individual lineages mapped on the phylogenetic tree based on Burki, et al. 2020 and Tikhonenkov, et al. 2022 following features are shown: number of species included in the analyses (in parentheses); genome, in which subunit-a/a$_N$ is encoded ("mt" – mitochondrial, "n" – nuclear); presence or absence of helices h$_N$ and h$_1$ of subunit-a/a$_N$; selected representative species used in the analyses presented in the following columns (for species shown in bold a cryoEM structure of ATP synthase has been reported); sequence of h$_N$ and flanking residues (universally conserved phenylalanine is shown in red, other conserved residues of the SPLEQF motif, or the FXXF motif in Metazoa, are shown in bold); average hydrophobicity of the entire subunit-a (FL), region h$_1$-h$_6$, and region h$_5$-h$_6$ (blue – more

hydrophobic, red – less hydrophobic); presence or absence of the histidine residue shown to

coordinate a water molecule in the Grotthuss chain in trypanosomal (His155) and bovine

(His168) ATP synthase (column H), and of the acidic residue (aspartate or glutamate) involved

in proton translocation (Asp202 in *T. brucei*; column D/E).

Figure 3. The proton entry point and path in the access half-channel is co-determined by

h_N. (A) Comparison of proton paths in the access half-channel of the *Acinetobacter baumanii*

and bovine mitochondrial ATP synthases. The c-ring is shown as white surface. **(B)**

Superposition of the structures from (A) shows that hN is positionally shifted compared to the

N-terminal helix of subunit-a in *A. baumanii*. **(C)** The position of the proposed "front" proton

entry to the access half-channel in *A. baumanii* ATP synthase (Demmer, et al. 2022) is blocked

by universally conserved phenylalanine in h_N of mitochondrial and alphaproteobacterial ATP

synthases. A view of the entry (red arrowhead) to the half channel in the *A. baumanii* subunit-

a (surface representation) from the c-ring shown individually and superposed with subunit-a

from other ATP synthases (cartoon representation): *S. cerevisiae*, the AlphaFold model of

Rickettsia, and subunits a and a_N from *T. brucei*. The conserved phenylalanine is shown as

sticks.