Light inducible protein degradation in *E. coli* with the LOVdeg tag

Nathan Tague[1,2], Cristian Coriano-Ortiz[1,2], Michael B. Sheets[1,2], Mary J. Dunlop[1,2*]

[1] Department of Biomedical Engineering, Boston University, Boston, MA, USA 02215
[2] Biological Design Center, Boston University, Boston, MA, USA 02215
* Corresponding author: mjdunlop@bu.edu

Abstract
Molecular tools for optogenetic control allow for spatial and temporal regulation of cell behavior. In particular, light controlled protein degradation is a valuable mechanism of regulation because it can be highly modular, used in tandem with other control mechanisms, and maintain functionality throughout growth phases. Here, we engineered LOVdeg, a tag that can be appended to a protein of interest for inducible degradation in *Escherichia coli* using blue light. We demonstrate the modularity of LOVdeg by using it to tag a range of proteins, including the LacI repressor, CRISPRa activator, and the AcrB efflux pump. Additionally, we demonstrate the utility of pairing the LOVdeg tag with existing optogenetic tools to enhance performance by developing a combined EL222 and LOVdeg system. Finally, we use the LOVdeg tag in a metabolic engineering application to demonstrate post-translational control of metabolism. Together, our results highlight the modularity and functionality of the LOVdeg tag system, and introduce a powerful new tool for bacterial optogenetics.

Introduction
Currently, the most widely used optogenetic systems in bacterial synthetic biology rely on transcription to modulate gene expression. Light responsive transcription factors allow genes of interest to be controlled modularly without having to re-engineer promoters. These tools have provided a valuable means to probe regulatory networks (Dessauges et al., 2022; Harrigan et al., 2018; Lugagne and Dunlop, 2019; Olson et al., 2014). However, these networks can be governed by complex control, where post-transcriptional and post-translational mechanisms work in concert with transcriptional regulation (Chubukov et al., 2014; Link et al., 2013; Mahmoud and Chien, 2018). Light responsive regulators that act beyond transcription are necessary to more fully mimic natural biology and have the potential to improve upon current deficits of transcriptional control. In synthetic contexts, transcriptional regulation suffers from limited response dynamics. For example, in the case of gene deactivation, decreasing protein abundance is limited by cell division-based dilution or natural protein half-lives, which are in the range of 5-20 hours for the majority of proteins in *E. coli* (Maurizi, 1992). When cells are growing in exponential phase the effective degradation rate of these long-lived proteins is controlled by the cell cycle time, which is on the order of 0.5 hours. However, at high cell densities, such as during stationary phase, slow growth rates result in protein half-lives on the scale of tens of hours (Maier et al., 2011). Subpopulations of cells in stationary phase have been reported to proliferate, however these growing subpopulations are small compared to non-growing cells and add to heterogeneous protein half-lives within the population (Jõers et al., 2020), making it challenging to reliably remove proteins. It should be noted that a proportion of proteins are actively degraded during slow growth, but these represent the minority in *E. coli* (Gupta et al., 2022). This is a critical issue for metabolic

engineering where chemical production is typically carried out at stationary phase in two-stage fermentations (Lalwani et al., 2018).

Post-translational optogenetic control has the potential to address some of the shortcomings of transcriptional regulation because post-translational control mechanisms can function independently of growth-based dilution. Existing approaches to post-translational optogenetic control include the use of light inducible dimers in split protein systems (Baumschlager et al., 2017; Kawano et al., 2015; Nihongaki et al., 2015), domain insertions with light controlled allosteric domains (Dagliyan et al., 2016; Gil et al., 2020; Tague et al., 2023), and membrane-confined functional control (Strickland et al., 2012; Wang et al., 2016). These approaches, however, suffer from a lack of modularity, which is a key benefit of the transcriptional control approaches. Split protein systems and domain insertions can require significant engineering for each protein of interest and solutions for a given protein are unlikely to map to proteins with different structures. Likewise, localization-based approaches are only applicable to a small subset of proteins that have position dependent function, such as transcription factors that require nuclear localization (Yumerefendi et al., 2018).

Protein degradation offers potential as a modular post-translational mechanism of control. Endogenous proteolytic machinery is necessary for proteome homeostasis and acts as a global regulatory system (Mahmoud and Chien, 2018). Targeted proteolysis has proven useful in many synthetic biology applications in both eukaryotic and prokaryotic systems (Andersen et al., 1998; Cameron and Collins, 2014; McGinness et al., 2006; Morreale et al., 2022; Trauth et al., 2019). In eukaryotic cells, light dependent protein degradation has been demonstrated using various mechanisms (Bonger et al., 2014; Deng et al., 2020; Liu et al., 2020; Renicke et al., 2013; Xue et al., 2019). For example, Bonger *et al.* utilized the LOV2 domain of *Avena sativa* phototropin 1 (*As*LOV2), to achieve light dependent protein degradation in mammalian cells (Bonger et al., 2014). LOV2 is a blue-light responsive protein that is widely used in optogenetic tools (Pudasaini et al., 2015). LOV2 contains a core Per-Arnt-Sim (PAS) domain surrounded by N- and C-terminal α-helices. Upon blue light illumination, the LOV2 protein undergoes a conformational change where the C-terminal Jα helix reversibly unfolds and becomes unstructured (Halavaty and Moffat, 2007; Harper et al., 2003; Yamamoto et al., 2009). Incorporation of a degradation-targeted peptide sequence into the C-terminal results in light dependent protein degradation in yeast (Renicke et al., 2013).

However, the bacterial proteasome differs from the eukaryotic proteasome in many ways (Mahmoud and Chien, 2018; Schrader et al., 2009). The bacterial proteasome includes several proteases with divergent targeting behaviors and does not utilize ubiquitin as a generalized modification to trigger degradation (Finley, 2009). In bacteria, protein targeting for degradation is predominantly dependent on primary amino acid sequence as opposed to a ubiquitin-like appendage and, apart from certain well-studied cases, the rules governing sequence recognition of bacterial degrons are not fully understood (Baker and Sauer, 2006; Striebel et al., 2009). Because of these complexities, light dependent degradation systems for bacteria have lagged behind their eukaryotic counterparts, where interaction of a ubiquitin ligase with a defined degron can be used as a control mechanism (Bondeson et al., 2022). Nevertheless, optogenetic degradation remains a key target for bacterial synthetic biology applications. In principle, an optogenetic degradation system could be fast-acting, modular, and interface with endogenous proteasome machinery.

These features would provide a straightforward way of adding dynamic control to a protein of interest, complementing existing transcriptional tools for optogenetic control of gene expression. One recent system developed by Komera *et al.* utilizes a light responsive split TEV protease to expose or remove constitutively active degradation tags in *E. coli* (Komera et al., 2022). Although this system achieves protein degradation in response to light, it acts indirectly through activation of an exogenous protease. A simpler approach where the degradation tag itself is light responsive would streamline this by eliminating the need for multiple exogenous components.

Here, we develop LOVdeg, a modular protein tag based on the *As*LOV2 protein that is conditionally degraded in response to blue light in *E. coli*. We demonstrate that attaching this tag to a protein of interest confers light dependent protein instability. We show the modularity of the LOVdeg tag by incorporating it into multiple proteins with widely varying function, converting them all into optogenetically controlled systems. In addition, through photocycle stabilizing mutations, we create a version of the LOVdeg tag that responds to infrequent exposure to blue light. We also demonstrate that our degradation tag can be used in concert with other optogenetic systems for multi-layer control by using EL222, a blue light responsive system for transcriptional control, together with LOVdeg. Lastly, we incorporate optogenetic degradation into a metabolically engineered strain to control production of octanoic acid. Overall, this work introduces a new bacterial optogenetic tool that overcomes several drawbacks of transcriptional control by providing post-translational degradation, while avoiding the need for substantial protein engineering that can be required for alternative post-translational control mechanisms.

Results

Design and characterization of the AsLOV2-based degradation tag

The *E. coli* proteasome consists of five AAA+ proteases and is continuously active, either degrading misfolded proteins for quality control or balancing regulatory protein levels (Mahmoud and Chien, 2018). We set out to exploit the endogenous proteasome activity in order to design a light responsive protein tag. To do this, we took insight from studies related to native protein quality control. In bacteria, peptides from stalled ribosomes are targeted for degradation through interaction with a tmRNA that appends a short amino acid sequence, known as an SsrA tag, to the C-terminal end of the incomplete protein (Keiler, 2015). The *E. coli* SsrA tag has been studied extensively and is known to interact with the unfoldases ClpX and ClpA (Flynn et al., 2001; Gottesman et al., 1998). Addition of the SsrA peptide sequence to exogenous proteins targets them for degradation by the host proteasome. SsrA mediated degradation has proven useful in synthetic gene circuit function and biochemical production (Elowitz, 2000; Gurbatri et al., 2020; Stricker et al., 2008; Torella et al., 2013; Ye et al., 2021). A recent structural study of ClpX interacting with the SsrA tag from Fei *et al.* demonstrated that in order for ClpX to unfold a tagged protein, the C-terminal tail needs to be unstructured and sufficiently long to fit into the ClpX pore (Fei et al., 2020). We reasoned that the mechanism of *As*LOV2, in which the C-terminal Jα helix becomes unstructured upon blue light absorption, could be utilized to provide light inducible protein degradation.

Biochemical studies have probed the amino acid sequence of the SsrA tag and its role in degradation targeting. Flynn *et al.* demonstrated that ClpA and ClpX unfoldases interact with

overlapping residues within the SsrA sequence and that the last three amino acids (L-A-A) are particularly important for successful degradation (Flynn et al., 2001). Importantly, in addition they showed that mutation of the leucine in the C-terminal 'L-A-A' lowers unfoldase affinity but does not hinder degradation completely. We noticed that the C-terminal amino acid sequence of the AsLOV2 domain, comprised of residues 404-546 of *Avena sativa* phototropin 1, are 'E-A-A' at positions 541-543 (Fig. 1a). The dark state structure of the native AsLOV2 domain (PDB: 2V1A) shows that these three amino acids and K544 complete the folded Jα helix (Fig. 1b). A truncation of residues 544-546 leaves the Jα helix largely intact and the resulting C-terminal 'E-A-A' remains caged as part of the folded helix as seen upon examination of the dark state structure (Fig. 1b). We hypothesized that this truncation would be stable in the dark state as a consequence of 'E-A-A' caging and unstable in the light state due to Jα helix unfolding and exposure of an unstructured degradation tag.

Optogenetic systems have used the AsLOV2 domain in *E. coli*, however, it is often appended N-terminally or internally to another protein (Li et al., 2022; Strickland et al., 2008). To the best of our knowledge, wild type or truncated AsLOV2, with its C-terminal end exposed, have not been used for proteolytic degradation in bacterial cells. To test the stability of C-terminal AsLOV2 in *E. coli* in response to light, we constructed a plasmid where we used an IPTG-inducible promoter to control the expression of mCherry translationally fused to AsLOV2 (Fig. 1c). In this construct, the full C-terminal sequence is intact so unfoldases are not expected to have good access for protein degradation. Consistent with this, induction of mCherry-AsLOV2(546) with IPTG increased expression of the fusion construct, and 465 nm blue light induction resulted in only a modest decrease in expression (Fig. 1d). Next, we tested a version of the construct where three amino acids were truncated to expose the C-terminal 'E-A-A', AsLOV2(543). This construct destabilized the protein fusion as predicted, resulting in significantly lower protein expression compared to the non-truncated version (Fig. 1d). However, counter to our expectations, we observed only a modest decrease in protein levels upon blue light induction.

While AsLOV2 switches from mostly folded in the dark state to mostly unfolded in the light state, both states are present at an equilibrium with or without light (Yao et al., 2008). Due to the suboptimal equilibrium of native AsLOV2, dark state undocking and unfolding of the Jα helix is a common issue in AsLOV2-based optogenetic tools, and many studies have aimed at improving the dynamic range of AsLOV2 (Guntas et al., 2015; Lungu et al., 2012; Strickland et al., 2012, 2010; Wang et al., 2016). Guntas *et al*., performed phage display on an AsLOV2-based protein that incorporates a caged peptide sequence in the Jα helix. They identified a variant with 11 amino acid substitutions with much tighter dark state caging, called iLID. Interestingly, several of these mutations do not have a direct interaction with the caged peptide, but instead are located at the hinge loop connecting the PAS domain to the Jα helix or in the PAS domain itself (Fig. S1). Guntas *et al*. further characterized amino acid substitutions and determined several that did not affect caging or dynamic range if reverted (F502Q, H521R, and C530M). We focused on the mutations found to result in tighter dark state caging (L493V, H519R, V520L, D522G, G528A, E537F, N538Q, and D540A), and reasoned that this set should stabilize the Jα helix irrespective of the caged peptide. Thus, we next constructed a variant of AsLOV2 with these eight mutations, which we denote as AsLOV2*.

To test whether these mutations are beneficial in the context of protein stabilization, we fused *As*LOV2*(546) or *As*LOV2*(543) to mCherry (Fig. 1c). mCherry with non-truncated *As*LOV2*(546) expressed highly with the addition of IPTG, but blue light did not result in a decrease in mCherry levels, consistent with lack of access by proteases (Fig. 1d). In contrast, the truncated version containing the mutations, AsLOV2*(543), displayed the desired light induced degradation, expressing highly in the dark and exhibiting 11.5x lower expression in response to blue light (Fig. 1d). We also confirmed that *As*LOV2*(543) expression was decreased in response to light without any IPTG induction (Fig. S2). Interestingly, *As*LOV2*(543) is degraded more in response to light relative to *As*LOV2(543). We conducted experiments to identify the mechanism underlying *As*LOV2*(543) degradation and found evidence that ClpA is involved, and other proteases and unfoldases play complementary roles (Supplementary Text, Fig. S3-6). We also verified that light exposure itself did not affect mCherry protein levels or growth by conducting experiments with an IPTG inducible mCherry without any *As*LOV2 variant fused and exposing cells to blue light (Fig. S7). We chose to move forward with the *As*LOV2*(543) variant due to its light inducible properties and we denoted this variant as the 'LOVdeg' tag.

Next, we used the LOVdeg tag to test the impact of light intensity and the timing of light exposure. We found that the LOVdeg tag is destabilized in proportion to light intensity (Fig. 1e), making it straightforward to tune degradation by adjusting light levels. In addition, a key advantage of protein degradation-based mechanisms is that they should function at a range of growth phases. We tested this by varying when we applied blue light, ranging from early exponential to stationary phase. We found that light-dependent decreases in protein levels can be achieved at various stages of growth (Fig. 1f, Fig. S8).

To benchmark the degradation capacity of the LOVdeg tag, we compared it to protein levels of mCherry subject to constitutive degradation via addition of an SsrA tag. In order to maintain comparable transcription and translation, we appended an iLID *As*LOV2 derivative, which was modified to contain a full length SsrA sequence, to the IPTG inducible mCherry. As expected, mCherry-iLID-SsrA expressed poorly independent of IPTG induction (Fig. S9). To quantify the light-dependent degradation of the LOVdeg tag, we compared its dark state expression level to that of mCherry-iLID-SsrA. Using this comparison, light induced degradation of the LOVdeg tag reached 6x degradation compared to 14x for the SsrA tag. However, the comparison between these two is convoluted by the time required to degrade the protein in response to light, making it challenging to directly compare these numbers.

A potential concern is that light induced disorder of the Jα helix could result in a decrease in solubility and aggregation of the LOVdeg tag. To rule this out as the cause of the fluorescence decrease, we captured microscopy images of cells in dark and light conditions. The imaging confirmed a light-dependent decrease in mCherry expression without the formation of visible protein aggregates (Fig. S10). Thus, the LOVdeg tag variant provides blue light dependent protein degradation.

Modularity of the LOVdeg tag

Post-translational control of protein function can require significant protein engineering for each use case (Sheets et al., 2020; Tague et al., 2023; Zhu et al., 2023). Degradation tags, by contrast,

offer post-translational control that theoretically requires little to no protein engineering and is protein agnostic. To test the modularity of the LOVdeg tag, we incorporated optogenetic control into three systems with highly diverse functions and relevance to synthetic biology and biotechnology applications: the LacI repressor, CRISPRa activation, and the AcrB efflux pump.

First, we sought to test whether the LOVdeg tag could be fused to transcription factors to enable light dependent regulation. The LacI repressor is a widely used chemically inducible system in synthetic biology. We translationally fused the LOVdeg tag to LacI and paired it with a reporter where the LacUV5 promoter controls expression of mCherry (Fig. 2a). Our results show that light exposure successfully increased mCherry expression (Fig. 2b). Light induced mCherry expression did not achieve the full levels provided with saturating IPTG induction, however we still observed a notable increase. We tested an alternative strategy for further improving de-repression, which suggested that the discrepancy between IPTG versus light dependent induction likely stems from the delay in LacI degradation compared to the rapid allosteric action of IPTG (Supplementary Text, Fig. S11).

Next, we incorporated the LOVdeg tag into the SoxS-based bacterial CRISPRa activation system (Dong et al., 2018). In this system, a scaffold RNA, which is a modified gRNA containing an MS2 stem loop, is used to localize dCas9 and the transcriptional activator SoxS, which is fused to an MS2 coat protein (MCP). We translationally fused the LOVdeg tag to the MCP-SoxS protein, such that in the dark CRISPRa will be active and light exposure relieves activation (Fig. 2c). In the original system, MCP-SoxS expression is anhydrotetracycline (aTc) inducible. This induction system in not amenable to blue light stimulation because aTc is photosensitive (Baumschlager et al., 2020). Thus, we changed the MCP-SoxS construct to an IPTG inducible promoter prior to blue light experiments (Fig. S12a, Table S1, Table S2). We confirmed that the promoter switch maintained CRISPRa activity (Fig. S12b). Fusing the LOVdeg tag to the activator component indeed relieved CRISPRa activity, resulting in a decrease in expression under blue light stimulation (Fig. 2d). However, reversal of activator activity was not complete and expression during light stimulation remained above the baseline levels from the reporter only control, which could be due to the presence of low levels of the MCP-SoxS activator.

We also added the LOVdeg tag to the endogenous membrane protein AcrB. This represents a challenging test case for degrading a native protein. The AcrAB-TolC complex is a multidrug efflux pump with clinical relevance due to its role in antibiotic tolerance and resistance acquisition (El Meouche and Dunlop, 2018; Lizarralde-Guerrero and Taraveau, 2021; Okusu et al., 1996). AcrAB-TolC has also been utilized in metabolic engineering as a mechanism to pump out toxic chemical products and boost strain performance (Dunlop et al., 2011; Fisher et al., 2014). However, inducible control of AcrB is challenging because cell viability is sensitive to overexpression (Turner and Dunlop, 2015). Optogenetic transcriptional control systems have high dynamic ranges but do not operate in the low expression ranges relevant to very potent protein complexes such as AcrAB-TolC. Light based degradation is well suited for this challenge because it works by decreasing protein levels, allowing the upper bound of expression to be determined by the promoter.

Previous work from Chai *et al*. demonstrated that AcrB can be targeted for proteolysis by fusing an SsrA tag to the C-terminal (Chai et al., 2016). Degradation is possible because the C-terminal

end of AcrB is on the cytoplasmic side of the inner membrane and can interact with cytoplasmic unfoldases (Du et al., 2014). Therefore, we reasoned that fusing the LOVdeg tag to AcrB would result in light inducible degradation that disrupts activity of the AcrAB-TolC complex (Fig. 2e). To determine if activity of the AcrAB-TolC complex was successfully disrupted by light, we performed an antibiotic sensitivity test using chloramphenicol, which is a known substrate of the AcrAB-TolC pump (Okusu et al., 1996). We transformed a construct containing *acrB-LOVdeg* into cells lacking the endogenous *acrB* gene (Δ*acrB*). When kept in the dark, Δ*acrB* cells with *acrB-LOVdeg* showed comparable chloramphenicol tolerance to wild type cells (Fig. 2f-g). Blue light stimulation, in contrast, sensitized the Δ*acrB* + AcrB-LOVdeg strain to chloramphenicol compared to both wild type and Δ*acrB* + AcrB-LOVdeg kept in the dark. This suggests that the LOVdeg tag successfully targets AcrB for degradation in a light dependent fashion. The blue light exposed cells still retain modest levels of chloramphenicol sensitivity when compared to Δ*acrB* cells without any AcrB complementation, likely due to basal levels of efflux pump expression from AcrB-LOVdeg relative to the knockout. As a point of comparison, we also tested chloramphenicol sensitivity of Δ*acrB* cells with IPTG-inducible AcrB expression (Δ*acrB* + AcrB) without IPTG. With no induction, this represents the lowest expression levels that can be achieved with a traditional IPTG-inducible system. The uninduced Δ*acrB* + AcrB cells displayed significantly higher tolerance to chloramphenicol when compared to wild type and other experimental groups. This underscores the challenge of controlling potent complexes like AcrAB-TolC through the use of chemical inducers alone and demonstrates how post-translational control, such as that provided by LOVdeg, is a viable and necessary strategy to decrease expression to levels approaching those in the Δ*acrB* knock out strain.

Tuning frequency response of the LOVdeg tag

Light inducible systems have the potential to respond to the frequency of light exposure. Frequency dependent tools can allow optogenetic circuits to be multiplexed beyond limited wavelength options or to add a layer of logic to optogenetic circuits (Benzinger et al., 2022). With added logic operations, optogenetic circuits could perform complex signal processing, analogous to those demonstrated with multiplexed chemically inducible circuits (Shin et al., 2020), while allowing dynamic light inputs. Additionally, higher sensitivity LOVdeg tags would also be useful in bioreactor settings, where poor light penetration into dense cultures is a feasibility concern. With these use cases in mind, we sought to characterize and alter the LOVdeg frequency response.

LOV domains utilize a flavin cofactor to absorb light. In the case of *As*LOV2, the cofactor responsible for light absorption is flavin mononucleotide (FMN). A cysteine residue in *As*LOV2 forms a reversible covalent bond with FMN, which initiates broader conformational change. A full photocycle of *As*LOV2 consists of absorption of a photon, covalent bond formation with cysteine 450, Jα helix destabilization and unfolding, decay of the cysteine-FMN bond, and Jα helix refolding (Swartz et al., 2001). Previous studies have determined the dynamics of bond formation and the time delay of Jα helix refolding. Further, mutations have been found that stabilize or destabilize the light state conformation (Christie et al., 2007; Kawano et al., 2013; Zayner et al., 2013). In the context of the LOVdeg tag, the time needed for the Jα helix to refold, known as the reversion time, likely determines degradation characteristics (Fig. 3a). In principle, the time spent in the light state dictates the amount of time the protein is susceptible to degradation and, therefore, would impact the frequency response given variable light inputs.

To test the effect of LOVdeg tag refolding dynamics, we illuminated cells with 5 second pulses of blue light followed by variable length dark periods to allow Jα helix refolding (Fig. 3b). While holding the blue light duration fixed, we tested dark periods ranging from 475 seconds (5:475 seconds on:off, frequency of 0.002 seconds^{-1} since there is 1 pulse every 480 seconds) to 5 seconds (5:5 seconds on:off, frequency of 0.1 seconds^{-1}) (Fig. 3b). We used 5 seconds for the pulse length because it is markedly shorter than the overall degradation dynamics for the LOVdeg tag, which ensures that a single pulse is not long enough to induce significant degradation. We first tested the frequency response of the original LOVdeg tag (Fig. 3c). The response is in line with known refolding dynamics of *As*LOV2 (Li et al., 2020), where over 50% degradation is only achieved at high frequencies (0.1 seconds^{-1}). Next, we tested a LOVdeg tag variant that contains a slow-photocycle mutation, V416I (Zoltowski et al., 2009). This amino acid substitution has been shown to increase the dark state reversion time from 8 seconds to 84 seconds *in situ* (Li et al., 2020). Indeed, for LOVdeg tag (V416I) over 50% degradation was achieved at medium frequencies (0.008 seconds^{-1}, which corresponds to 5:120 seconds on:off) (Fig. 3c). This variant offers the potential for better performance in settings where increased light sensitivity is preferred, such as within bioreactors.

Integrating LOVdeg with EL222 for multi-layer control

Another attractive aspect of post-translational optogenetic control is that it can integrate with existing systems that act at the transcriptional or translational level. Adding control at multiple layers has been shown to enhance the performance and robustness of natural and synthetic systems (Alon, 2007; Hasenjäger et al., 2019; Szydlo et al., 2022). One commonly used system for optogenetic transcriptional control is EL222. EL222 is a blue light responsive LOV protein that dimerizes and binds DNA upon light exposure (Zoltowski et al., 2013). In bacteria, EL222 can be used as a transcriptional repressor or activator depending on the placement of its binding site in the promoter (Ding et al., 2020; Jayaraman et al., 2016). We chose to combine the transcriptional repression of EL222 with the LOVdeg tag. In this arrangement, the systems work synergistically to decrease gene expression in response to blue light using simultaneous transcriptional repression and protein degradation (Fig. 4a).

To test the performance of this combined optogenetic circuit, we created a construct in which the mCherry-LOVdeg fusion protein is driven by a promoter containing an EL222 binding site (P$_{EL222}$). We tested the light response of mCherry-LOVdeg expression with and without EL222 present, as well as including EL222 control of mCherry without the LOVdeg tag fused (Fig. 4b). As expected, mCherry-LOVdeg expression decreased in response to light even without EL222 present, representing the sole action of the degradation tag. Similarly, EL222-only mCherry decreased in response to light due to repression by EL222. However, multi-layer control resulted in a faster decrease in expression in response to light and reached lower levels compared to either LOVdeg or EL222 alone. The fold change decrease in expression was improved from 7x and 6x for just the LOVdeg or EL222, respectively, to 23x when both systems were combined.

Optogenetic control of octanoic acid production

We next sought to apply the LOVdeg tag to a metabolic engineering task, as we envision that post-translational control will be especially advantageous in these biotechnology applications. Transcriptional control alone is particularly problematic in a metabolic engineering setting because chemical production is typically carried out at stationary phase with slow growth rates, meaning that proteins expressed at basal levels will accumulate in production settings. Therefore, dynamic control using transcriptional optogenetic systems alone will only allow protein levels to increase or plateau. However, a prerequisite for dynamic control of metabolic pathways is that enzyme levels can be modulated to turn off production.

As a proof of concept, we chose to control the enzyme CpFatB1 with LOVdeg, EL222, or the EL222-LOVdeg circuit. CpFatB1 is an acyl-ACP thioesterase from *Cuphea palustris* that primarily catalyzes octanoyl-ACP to produce octanoic acid. Octanoic acid is a valuable medium chain oleochemical with limited natural sources (Sarria et al., 2017). Specifically, we expressed the catalytically enhanced mutant from Hernandez-Lozada, *et al.*, CpFatB1.2-M4-287, which we denote here as CpFatB1* (Hernández Lozada et al., 2018). This enzyme interfaces with the endogenous fatty acid synthesis pathway in *E. coli* to produce free fatty acids (Fig. 5a). In this pathway, the carbon tail of an acyl-ACP moiety is elongated two carbons at a time. CpFatB1* specifically catalyzes C8-ACPs, which results in production of free octanoic acid. Importantly, since CpFatB1* is very catalytically active, low levels of expression are optimal for production and strains with high expression exhibit a growth defect (Hernández Lozada et al., 2018). Therefore, to dynamically regulate CpFatB1* activity in cells, expression must be controlled in a low range, which is particularly challenging in stationary phase.

A first step towards optogenetically controlling this metabolic pathway is to demonstrate that enzyme levels can be modulated between protein levels that are relevant to endpoint titers. If expression can only oscillate between high and medium protein levels, dynamic light control will not be effective. To test whether we could effectively stunt CpFatB1* activity, we controlled enzyme expression using only LOVdeg tag degradation, only EL222 repression, or combined LOVdeg and EL222 throughout at 24 hour fermentation period (Fig. 5b). LOVdeg alone successfully decreased octanoic acid in the light condition. The EL222-only condition did not significantly decrease octanoic acid production with light compared to conditions in the dark, demonstrating the shortcomings of solely transcriptional control at stationary phase. In contrast, the LOVdeg tag with EL222 resulted in a significant decrease in octanoic acid in the continuous light condition. Thus, protein degradation is needed to effectively shunt the metabolic pathway during stationary phase.

Interestingly, the LOVdeg-only strain, in which CpFatB1*-LOVdeg is expressed through the IPTG-inducible LacUV5 promoter, exhibits higher dark state octanoic acid production compared to CpFatB1* expressed under the EL222 promoter. The dark state production of the LOVdeg-only strain is similar to that of CpFatB1* without the LOVdeg tag, both expressed via the same LacUV5 promoter. This is likely due to limited constitutive expression from P_{EL222}. This limitation further demonstrates the utility of post-translational control where the promoter is not involved in the light response. Previous efforts aimed at increasing the strength of EL222 responsive promoters have been carried out (Ding et al., 2020), and further promoter engineering would be needed to increase EL222 promoter strength to increase octanoic acid production in the dark state. Clearly, decreasing titers is not the overall goal in metabolic engineering. However, the ability to control expression

at relevant ranges throughout stationary phase represents an important stepping stone to investigate dynamic control schemes or feedback control that may lead to enhanced strain performance.

Discussion

Here, we developed and characterized the LOVdeg tag, which provides blue light inducible protein degradation, offering unique advantages as a tool for bacterial synthetic biology. Protein degradation offers a mode of control that is currently limited in the bacterial optogenetic toolkit. The bacterial light responsive degradation system from Komera *et al.* accomplishes protein degradation indirectly through an exogenous split protease (Komera et al., 2022). In comparison, in our approach the degradation tag itself is light responsive and does not require extra components, making it straightforward to incorporate. In this study, we showed that the truncated *As*LOV2 protein can be fused to the C-terminal end of various proteins to destabilize them and then target them for degradation. The addition of stabilizing mutations from iLID (to create *As*LOV2*) result in low basal activity in the dark, with strong switching in response to light. Aside from creating a light-responsive degradation tag, the effectiveness of these stabilizing mutations may serve to improve the switch-like behavior in many other *As*LOV2-based systems.

We demonstrated the modularity of the LOVdeg tag by incorporating it into three distinct contexts with little to no fine tuning. Each system, the LacI repressor, CRISPRa activation, and the AcrB efflux pump, was converted to optogenetically controlled by simply tagging the protein with LOVdeg. This modularity mimics the ease of use of transcriptional optogenetic systems, which only require a change of the promoter. However, incorporating light control at the protein level simplifies this process because protein levels can remain in their native, or previously fine-tuned, context. This configuration is especially beneficial for proteins that require low expression, as transcriptional optogenetic systems often suffer from leaky basal expression. The LOVdeg tag provides a design that can work with endogenous protein levels, circumventing this issue. In this study, we focused on proteins encoded on plasmids expressed from synthetic promoters. However, we anticipate the LOVdeg tag can be useful in systems biology contexts where natural promoters linked to endogenous gene networks are best left untouched. In this case, the LOVdeg tag can add the ability to optogenetically manipulate genes while keeping them in their native gene regulatory context.

Since degradation occurs post-translationally, the LOVdeg tag can easily be integrated with existing optogenetic systems to enhance their function. We demonstrated this by combining the LOVdeg tag with the EL222 repression system, showing that the synergistic action of transcriptional repression and degradation results in an increase in the dynamic range, owing to the lower off-state under light illumination. In this configuration, EL222 and the LOVdeg tag work coherently to decease expression. However, the LOVdeg tag can potentially be incorporated into activating systems incoherently to create dynamic functionality, such as pulse generators and inverters (Benzinger et al., 2022).

Throughout this study, we performed experiments using low copy number plasmids that result in moderate levels of a given protein of interest. While this is a common use case for synthetic biology, degradation as the sole mode of gene expression control may be limiting when proteins are at very high levels. As an ATP-dependent process that utilizes a finite pool of proteolytic

machinery, degradation rates can saturate at sufficiently high protein levels (Cookson et al., 2011). Furthermore, using protein degradation as the sole mode of gene expression control is wasteful in some contexts. Constitutive transcription and translation of a gene followed by degradation utilizes valuable cellular resources, which is an important consideration in metabolic engineering. To avoid this type of energetic waste, we envision the LOVdeg tag could be used in concert with other modes of regulation, as we demonstrated with EL222. In addition, it is unclear whether the LOVdeg tag is compatible with other prokaryotic (including mitochondrial) proteasomes. Future studies focused on the portability of the LOVdeg tag may address this question and potentially lead to further mechanistic insight.

In summary, the LOVdeg tag offers a straightforward route for introducing optogenetic control of protein degradation in *E. coli*. By lowering the barrier to entry for incorporating light responsiveness into a protein of interest, we envision that systems typically studied with chemical induction or constitutive expression can now be controlled optogenetically without extensive fine-tuning. Furthermore, the LOVdeg tag can act as a circuit enhancer when incorporated into existing optogenetic systems to increase functionality and robustness.

Methods

Strains and plasmids
We used *E. coli* BW25113 as the wild type strain. All knockout strains are from the Keio collection (Baba et al., 2006), which were derived from BW25113. We used golden gate cloning to create all plasmid constructs (Engler et al., 2008) (Table S1). The IPTG inducible constructs were derived from pBbS5c-mRFP1 from the BglBrick plasmid library (Lee et al., 2011). In the constitutive version of the mCherry-AsLOV2 variants, we swapped the IPTG inducible promoter with a constitutive promoter, P_{W7} (5'-ttatcaaaaagagtattgaaataaagtctaacctataggaagattacagccatcgagagggacacggcgaa-3'). We used this constitutive version for microscopy and the protease knockout studies; all other experiments used the IPTG inducible promoter P_{lacUV5}.

In all cases of protein fusions to *As*LOV2 variants, a five amino acid GS linker of 'S-S-G-S-G' was used between the protein of interest and the LOVdeg tag. In the LacI-LOVdeg experiments, pBbS5c-mCherry was also used as the backbone, but with the LOVdeg sequence cloned after the *lacI* gene instead of *mCherry*. In the LacI-LOVdeg experiment to test the impact of basal LacI, pBbS5c-LacI-LOVdeg-mCherry was co-transformed with pBbAa-LacI-Decoy from Wang and Tague *et al.* (Wang et al., 2021). CRISPRa constructs were based on the MS2-SoxS system from Dong *et al.* (Dong et al., 2018). The J106 gRNA sequence from Dong *et al.* was used to target dCas9 upstream mRFP1 under control of the minimal J1 promoter. This gRNA plasmid was constructed using the golden gate assembly method to replace the targeting sequence in the pCD061 backbone (Addgene #113315). The mRFP1 reporter plasmid was derived from pJF076Sa (Addgene #113322) by replacing the ampicillin resistance gene with a kanamycin resistance gene from the BglBrick library. MS2-SoxS-LOVdeg was expressed from a variant of pJF093 (Addgene #113323). TetR and its corresponding promoter driving expression of MS2-SoxS were replaced with the LacI-P_{Trc} inducible system from pBbA1c-mRFP1 from the BglBrick library (Lee et al., 2011). The inducer was changed to IPTG because aTc is sensitive to blue light (Baumschlager et

al., 2020). The LOVdeg tag was added to this construct with a C-terminal fusion to SoxS. Plasmids containing *acrAB* were built from pBbA5k-acrAB from El Meouche *et al.* (El Meouche and Dunlop, 2018). The FLP recombination protocol from Datsenko and Wanner was used to cure the *kanR* cassette from the genome of the Δ*acrB* strain (Keio collection, JW0451) (Datsenko and Wanner, 2000).

The slow photocycle variant of the LOVdeg tag, V416I, was constructed using site directed mutagenesis of Valine at amino acid position 416.

EL222 was synthesized by IDT (Coralville, IA) and plasmids were constructed to mimic EL222 repression systems from Jayaraman *et al.* (Table S1) (Jayaraman et al., 2016). A variant of the promoter $P_{BLrep-v1}$ from Ding *et al.*, P_{raB} was used in all EL222 experiments and we refer to it in figures as P_{EL222} (Ding et al., 2020). Plasmid pBbE5k-P_{EL222}-mCherry-LOVdeg was co-transformed with pEL222, which constitutively expresses EL222 (Table S2).

For octanoic acid production experiments, the coding sequence of *Cp*FatB1.2-M4-287 derived from Hernández Lozada *et al.* was synthetized by Twist Biosciences (South San Francisco, CA) and cloned after the P_{EL222} or P_{lacUV5} promoter (Hernández Lozada et al., 2018). Plasmid pBbE5k-P_{EL222}-*Cp*FatB1*-LOVdeg was co-transformed with pEL222 for LOVdeg + EL222 production control (Table S2).

Plasmids expressing ClpA, ClpX, and HslU were constructed by amplifying the unfoldase gene from the wild type genome and inserting it into the pBbA8k backbone from the BglBrick library (Lee et al., 2011).

Constructs from this work are available on AddGene: https://www.addgene.org/Mary_Dunlop/

Blue light stimulation
Unless otherwise noted, bacteria were cultured in Luria broth (LB) with appropriate antibiotics for plasmid maintenance at 37°C with 200 rpm shaking. Antibiotic concentrations used for plasmid maintenance were 30 μg/mL for kanamycin, 100 μg/mL for carbenicillin, and 25 μg/mL for chloramphenicol. All light exposure experiments were carried out with a light plate apparatus (LPA) (Gerhardt et al., 2016) using 465 nm blue light. Overnight cultures of light sensitive strains were diluted 1:50 and precultured in the dark for 2 hours. For IPTG inducible constructs, 1 mM IPTG was added when the cells were diluted. After 2 hours in the dark, cells were exposed to blue light in the LPA at a setpoint of 100 μW/cm². Red fluorescence (excitation 560 nm, emission 600 nm) and optical density (OD) readings were taken using a BioTek Synergy H1m plate reader (BioTek, Winooski, VT) after 5 hours of incubation unless otherwise noted. For experiments testing degradation at various growth phases, a 4 hour light window was introduced at variable times, as shown in Fig. 1. For CRISPRa experiments, light stimulation was continued for 8 hours prior to RFP and OD readings. For frequency response experiments, the LPA was programmed using its Iris software (https://taborlab.github.io/Iris/) to pulse blue light at varying frequencies. A 5 second light pulse was kept constant for each experiment while the time between pulses was varied (5s, 55s, 85s, 115s, 235s, and 480s).

Chloramphenicol sensitivity testing

Chloramphenicol sensitivity experiments were performed in M9 minimal media (M9 salts, 2 mM MgSO$_4$, 100 μM CaCl$_2$) with 1% glucose at 37°C with 200 rpm shaking. Overnight cultures in LB were initially diluted 1:50 into M9 media for 4 hours. The M9 conditioned cultures were then diluted again 1:20 into 24 well plates containing M9 media with varying levels of chloramphenicol (0, 0.3125, 0.625, 1.25, 2.5, 5, and 10 μg/ml) and grown for 20 hours before measuring optical density (OD) using a BioTek Synergy H1m plate reader (BioTek, Winooski, VT). We also conducted experiments with the *E. coli* BW25113 Δ*acrB* strain (referred to as Δ*acrB*). For these, Δ*acrB* was transformed with pBbA5k-AcrAB-LOVdeg (Table S2) and chloramphenicol sensitivity experiments were carried out either in the LPA with constant light illumination or kept in the dark for the duration of growth. No IPTG was added to the Δ*acrB* + AcrAB-LOVdeg cultures because basal expression is enough to recover wild type resistance. The same chloramphenicol sensitivity protocol was performed in the dark using BW25113, Δ*acrB* + AcrB without induction, and Δ*acrB* as controls.

Microscopy
Strains were grown overnight in LB medium. Cultures were refreshed 1:100 in M9 minimal media for microscopy (M9 salts supplemented with 2 mM MgSO4, 0.1 mM CaCl2) with 0.4% glucose for two hours. Samples were then placed on 1.5% low melting agarose pads made with M9 minimal media for microscopy with 0.4% glucose. Samples were grown at 30°C. Cells were imaged at 100x using a Nikon Ti-E microscope. Blue light exposure was provided by a LED ring (Adafruit NeoPixel 1586), which was fixed above the microscope stage and controlled by an Arduino using a custom Matlab script. Blue light was kept constant except during image acquisitions.

Octanoic acid production experiment
For octanoic acid production experiments, strains expressing *Cp*FatB1* under various modes of control (Table S2) were cultured in LB overnight with light illumination to maintain low *Cp*FatB1* expression. Overnight cultures were diluted 1:20 into M9 minimal media with 2% glucose and kept in the light until they reached early stationary phase (OD$_{600}$ of 0.6) unless otherwise noted. The LPA was then programmed to either maintain light for low octanoic acid production or turn off light exposure to induce octanoic acid production for 24 hours prior to fatty acid extraction and quantification.

Fatty acid quantification
Samples for GC-MS quantification were taken at 24 hours. 400 μL of vortexed culture was used for fatty acid extraction and derivatization into fatty acid methyl esters as described by Sarria *et al.* (Sarria et al., 2018) with the following minor modifications: An internal standard of nonanoic acid (C9) was added to the 400 μL sample at a final concentration of 88.8 mg/L and vortexed for 5 sec. The following was then added to the sample for fatty acid extraction and vortexed for 30 sec: 50 μL 10% NaCl, 50 μL glacial acetic acid, and 200 μL ethyl acetate. The sample was then centrifuged at 12,000 g for 10 mins. After centrifugation, 100 μL of the ethyl acetate layer was mixed with 900 μL of a 30:1 mixture of methanol:HCl (12N) in a 2 mL microcentrifuge tube. The solution was vortexed for 30 sec followed by an incubation at 50°C for 60 mins for methyl ester derivatization. Once cooled to room temperature, 500 μL hexanes and 500 μL water were added to the 2 mL microcentrifuge tube, vortexed for 10 sec, and allowed to settle. 250 μL of the hexane layer was mixed with 250 μL ethyl acetate in a GC-MS vial for quantification.

The samples were analyzed with an Agilent 6890N/Agilent 5973 MS detector using a DB-5MS column. The inlet temperature was set to 300°C with flow at 4 mL/min. The oven heating program was initially set to 70°C for 1 min, followed by a ramp to 290°C at 30°C/min, and a final hold at 290°C for 1 min. GLC-20 and GLC-30 FAME standard mixes (Sigma) were tested using this protocol to ensure proper capture of all chain lengths and to gauge retention times. Internal standards were used for quantification, with chain lengths C8-C12 quantified with the nonanoic acid internal standard and C14-C18 quantified with the pentadecanoic internal standard.

Acknowledgements
We thank members of the Dunlop Lab for helpful discussions. This work was supported by DOE grant DE-SC0019387, NSF grant CBET-1804096, and NIH grant R01AI102922. CCO and MBS received support through the NIH training grants T32 GM130546 and T32 EB006359, respectively.

Conflict of Interest
The authors declare no competing interests.

References

Alon U. 2007. Network motifs: Theory and experimental approaches. *Nature Reviews Genetics* **8**:450–461. doi:10.1038/nrg2102

Andersen JB, Sternberg C, Poulsen LK, Bjørn SP, Givskov M, Molin S. 1998. New unstable variants of green fluorescent protein for studies of transient gene expression in bacteria. *Applied and Environmental Microbiology* **64**:2240–2246. doi:10.1128/aem.64.6.2240-2246.1998

Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants: The Keio collection. *Molecular Systems Biology* **2**. doi:10.1038/msb4100050

Baker TA, Sauer RT. 2006. ATP-dependent proteases of bacteria: recognition logic and operating principles. *Trends in Biochemical Sciences* **31**:647–653. doi:10.1016/j.tibs.2006.10.006

Baumschlager A, Aoki SK, Khammash M. 2017. Dynamic Blue Light-Inducible T7 RNA Polymerases (Opto-T7RNAPs) for Precise Spatiotemporal Gene Expression Control. *ACS Synthetic Biology* **6**:2157–2167. doi:10.1021/acssynbio.7b00169

Baumschlager A, Rullan M, Khammash M. 2020. Exploiting natural chemical photosensitivity of anhydrotetracycline and tetracycline for dynamic and setpoint chemo-optogenetic control. *Nature Communications* **11**:1–10. doi:10.1038/s41467-020-17677-5

Benzinger D, Ovinnikov S, Khammash M. 2022. Synthetic gene networks recapitulate dynamic signal decoding and differential gene expression. *Cell Systems* **13**:353-364.e6. doi:10.1016/j.cels.2022.02.004

Bondeson DP, Mullin-Bernstein Z, Oliver S, Skipper TA, Atack TC, Bick N, Ching M, Guirguis AA, Kwon J, Langan C, Millson D, Paolella BR, Tran K, Wie SJ, Vazquez F, Tothova Z, Golub TR, Sellers WR, Ianari A. 2022. Systematic profiling of conditional degron tag technologies for target validation studies. *Nature Communications* **13**. doi:10.1038/s41467-022-33246-4

Bonger KM, Rakhit R, Payumo AY, Chen JK, Wandless TJ. 2014. General method for regulating protein stability with light. *ACS Chemical Biology* **9**:111–115. doi:10.1021/cb400755b

Cameron DE, Collins JJ. 2014. Tunable protein degradation in bacteria. *Nature Biotechnology* **32**:1276–1281. doi:10.1038/nbt.3053

Chai Q, Wang Z, Webb SR, Dutch RE, Wei Y. 2016. The ssrA-Tag Facilitated Degradation of an Integral Membrane Protein. *Biochemistry* **55**:2301–2304. doi:10.1021/acs.biochem.6b00038

Christie JM, Corchnoy SB, Swartz TE, Hokenson M, Han IS, Briggs WR, Bogomolni RA. 2007. Steric interactions stabilize the signaling state of the LOV2 domain of phototropin 1. *Biochemistry* **46**:9310–9319. doi:10.1021/bi700852w

Chubukov V, Gerosa L, Kochanowski K, Sauer U. 2014. Coordination of microbial metabolism. *Nature Reviews Microbiology* **12**:327–340. doi:10.1038/nrmicro3238

Cookson NA, Mather WH, Danino T, Mondragõn-Palomino O, Williams RJ, Tsimring LS, Hasty J. 2011. Queueing up for enzymatic processing: Correlated signaling through coupled degradation. *Molecular Systems Biology* **7**. doi:10.1038/msb.2011.94

Dagliyan O, Tarnawski M, Chu PH, Shirvanyants D, Schlichting I, Dokholyan N V., Hahn KM. 2016. Engineering extrinsic disorder to control protein activity in living cells. *Science*

354:1441–1444. doi:10.1126/science.aah3404

Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proceedings of the National Academy of Sciences of the United States of America* **97**:6640–6645. doi:10.1073/pnas.120163297

Deng W, Bates JA, Wei H, Bartoschek MD, Conradt B, Leonhardt H. 2020. Tunable light and drug induced depletion of target proteins. *Nature Communications* **11**:1–13. doi:10.1038/s41467-019-14160-8

Dessauges C, Mikelson J, Dobrzyński M, Jacques M, Frismantiene A, Gagliardi PA, Khammash M, Pertz O. 2022. Optogenetic actuator – ERK biosensor circuits identify MAPK network nodes that shape ERK dynamics. *Molecular Systems Biology* **18**:1–22. doi:10.15252/msb.202110670

Ding Q, Ma D, Liu GQ, Li Y, Guo L, Gao C, Hu G, Ye C, Liu J, Liu L, Chen X. 2020. Light-powered Escherichia coli cell division for chemical production. *Nature Communications* **11**. doi:10.1038/s41467-020-16154-3

Dong C, Fontana J, Patel A, Carothers JM, Zalatan JG. 2018. Synthetic CRISPR-Cas gene activators for transcriptional reprogramming in bacteria. *Nature Communications*. doi:10.1038/s41467-018-04901-6

Du D, Wang Z, James NR, Voss JE, Klimont E, Ohene-Agyei T, Venter H, Chiu W, Luisi BF. 2014. Structure of the AcrAB-TolC multidrug efflux pump. *Nature* **509**:512–515. doi:10.1038/nature13205

Dunlop MJ, Dossani ZY, Szmidt HL, Chu HC, Lee TS, Keasling JD, Hadi MZ, Mukhopadhyay A. 2011. Engineering microbial biofuel tolerance and export using efflux pumps. *Molecular Systems Biology* **7**:1–7. doi:10.1038/msb.2011.21

El Meouche I, Dunlop MJ. 2018. Heterogeneity in efflux pump expression predisposes antibiotic-resistant cells to mutation. *Science* **362**:686–690. doi:10.1126/science.aar7981

Elowitz. 2000. A synthetic oscillatory network repressilator. *Nature* **403**:335–338.

Engler C, Kandzia R, Marillonnet S. 2008. A one pot, one step, precision cloning method with high throughput capability. *PLoS ONE* **3**:e3647. doi:10.1371/journal.pone.0003647

Fei X, Bell TA, Barkow SR, Baker TA, Sauer RT. 2020. Structural basis of clpxp recognition and unfolding of ssra-tagged substrates. *eLife* **9**:1–39. doi:10.7554/eLife.61496

Finley D. 2009. Recognition and processing of ubiquitin-protein conjugates by the proteasome. *Annual Review of Biochemistry* **78**:477–513. doi:10.1146/annurev.biochem.78.081507.101607

Fisher MA, Boyarskiy S, Yamada MR, Kong N, Bauer S, Tullman-Ercek D. 2014. Enhancing tolerance to short-chain alcohols by engineering the Escherichia coli AcrB efflux pump to secrete the non-native substrate n-butanol. *ACS Synthetic Biology* **3**:30–40. doi:10.1021/sb400065q

Flynn JM, Levchenko I, Seidel M, Wickner SH, Sauer RT, Baker TA. 2001. Overlapping recognition determinants within the ssrA degradation tag allow modulation of proteolysis. *Proceedings of the National Academy of Sciences of the United States of America* **98**:10584–10589. doi:10.1073/pnas.191375298

Gerhardt KP, Olson EJ, Castillo-Hair SM, Hartsough LA, Landry BP, Ekness F, Yokoo R, Gomez EJ, Ramakrishnan P, Suh J, Savage DF, Tabor JJ. 2016. An open-hardware platform for optogenetics and photobiology. *Scientific Reports* **6**:1–13. doi:10.1038/srep35363

Gil AA, Carrasco-López C, Zhu L, Zhao EM, Ravindran PT, Wilson MZ, Goglia AG, Avalos JL, Toettcher JE. 2020. Optogenetic control of protein binding using light-switchable

nanobodies. *Nature Communications* **11**:1–12. doi:10.1038/s41467-020-17836-8

Golding I, Paulsson J, Zawilski SM, Cox EC. 2005. Real-time kinetics of gene activity in individual bacteria. *Cell* **123**:1025–1036. doi:10.1016/j.cell.2005.09.031

Gottesman S, Roche E, Zhou YN, Sauer RT. 1998. The ClpXP and ClpAP proteases degrade proteins with carboxy-terminal peptide tails added by the SsrA-tagging system. *Genes and Development* **12**:1338–1347. doi:10.1101/gad.12.9.1338

Guntas G, Hallett RA, Zimmerman SP, Williams T, Yumerefendi H, Bear JE, Kuhlman B. 2015. Engineering an improved light-induced dimer (iLID) for controlling the localization and activity of signaling proteins. *Proceedings of the National Academy of Sciences of the United States of America* **112**:112–117. doi:10.1073/pnas.1417910112

Gupta M, Johnson A, Cruz E, Costa E, Guest R, Li SH-J, Hart EM, Nguyen T, Stadlmeier M, Bratton BP, Silhavy TJ, Wingreen NS, Gitai Z, Wühr M. 2022. Global Protein-Turnover Quantification in Escherichia coli Reveals Cytoplasmic Recycling under Nitrogen-Limitation. *bioRxiv* 2022.08.01.502339.

Gurbatri CR, Lia I, Vincent R, Coker C, Castro S, Treuting PM, Hinchliffe TE, Arpaia N, Danino T. 2020. Engineered probiotics for local tumor delivery of checkpoint blockade nanobodies. *Science Translational Medicine* **12**. doi:10.1126/scitranslmed.aax0876

Halavaty AS, Moffat K. 2007. N- and C-terminal flanking regions modulate light-induced signal transduction in the LOV2 domain of the blue light sensor phototropin 1 from Avena sativa. *Biochemistry* **46**:14001–14009. doi:10.1021/bi701543e

Harper SM, Neil LC, Gardner KH. 2003. Structural basis of a phototropin light switch. *Science* **301**:1541–1544. doi:10.1126/science.1086810

Harrigan P, Madhani HD, El-Samad H. 2018. Real-Time Genetic Compensation Defines the Dynamic Demands of Feedback Control. *Cell* **175**:877-886.e10. doi:10.1016/j.cell.2018.09.044

Hasenjäger S, Trauth J, Hepp S, Goenrich J, Essen LO, Taxis C. 2019. Optogenetic Downregulation of Protein Levels with an Ultrasensitive Switch. *ACS Synthetic Biology* **8**:1026–1036. doi:10.1021/acssynbio.8b00471

Hernández Lozada NJ, Lai RY, Simmons TR, Thomas KA, Chowdhury R, Maranas CD, Pfleger BF. 2018. Highly Active C8 -Acyl-ACP Thioesterase Variant Isolated by a Synthetic Selection Strategy. *ACS Synthetic Biology* **7**:2205–2215. doi:10.1021/acssynbio.8b00215

Jayaraman P, Devarajan K, Chua TK, Zhang H, Gunawan E, Poh CL. 2016. Blue light-mediated transcriptional activation and repression of gene expression in bacteria. *Nucleic Acids Research* **44**:6994–7005. doi:10.1093/nar/gkw548

Jõers A, Liske E, Tenson T. 2020. Dividing subpopulation of Escherichia coli in stationary phase. *Research in Microbiology* **171**:153–157. doi:10.1016/j.resmic.2020.02.002

Kawano F, Aono Y, Suzuki H, Sato M. 2013. Fluorescence imaging-based high-throughput screening of fast- and slow-cycling LOV proteins. *PLoS ONE* **8**. doi:10.1371/journal.pone.0082693

Kawano F, Suzuki H, Furuya A, Sato M. 2015. Engineered pairs of distinct photoswitches for optogenetic control of cellular proteins. *Nature Communications*. doi:10.1038/ncomms7256

Keiler KC. 2015. Mechanisms of ribosome rescue in bacteria. *Nature Reviews Microbiology* **13**:285–297. doi:10.1038/nrmicro3438

Komera I, Gao C, Guo L, Hu G, Chen X, Liu L. 2022. Bifunctional optogenetic switch for improving shikimic acid production in E. coli. *Biotechnology for Biofuels and Bioproducts* **15**:1–14. doi:10.1186/s13068-022-02111-3

Lalwani MA, Zhao EM, Avalos JL. 2018. Current and future modalities of dynamic control in metabolic engineering. *Current Opinion in Biotechnology* **52**:56–65. doi:10.1016/j.copbio.2018.02.007

Lee TS, Krupa RA, Zhang F, Hajimorad M, Holtz WJ, Prasad N, Lee SK, Keasling JD. 2011. BglBrick vectors and datasheets: A synthetic biology platform for gene expression. *Journal of Biological Engineering* **5**:12. doi:10.1186/1754-1611-5-12

Li LL, Klein FM, Li Greci L, Popinigis A, Freudenberg F, Courtney MJ. 2020. Resonance energy transfer sensitises and monitors in situ switching of LOV2-based optogenetic actuators. *Nature Communications* **11**:1–18. doi:10.1038/s41467-020-18816-8

Li Y, Zhao M, Wei D, Zhang J, Ren Y. 2022. Photocontrol of Itaconic Acid Synthesis in Escherichia coli. *ACS Synthetic Biology* **11**:2080–2088. doi:10.1021/acssynbio.2c00014

Link H, Kochanowski K, Sauer U. 2013. Systematic identification of allosteric protein-metabolite interactions that control enzyme activity in vivo. *Nature Biotechnology* **31**:357–361. doi:10.1038/nbt.2489

Liu J, Chen H, Ma L, He Z, Wang D, Liu Y, Lin Q, Zhang T, Gray N, Kaniskan HÜ, Jin J, Wei W. 2020. Light-induced control of protein destruction by opto-PROTAC. *Science Advances* **6**:1–12. doi:10.1126/sciadv.aay5154

Lizarralde-Guerrero M, Taraveau F. 2021. Role of AcrAB-TolC multidrug efflux pump in drug-resistance acquisition by plasmid transfer. *Medecine/Sciences* **37**:945–947. doi:10.1051/medsci/2021159

Lugagne JB, Dunlop MJ. 2019. Cell-machine interfaces for characterizing gene regulatory network dynamics. *Current Opinion in Systems Biology* **14**:1–8. doi:10.1016/j.coisb.2019.01.001

Lungu OI, Hallett RA, Choi EJ, Aiken MJ, Hahn KM, Kuhlman B. 2012. Designing Photoswitchable Peptides Using the AsLOV2 Domain. *Chemistry and Biology* **19**:507–517. doi:10.1016/j.chembiol.2012.02.006

Mahmoud SA, Chien P. 2018. Regulated Proteolysis in Bacteria. *Annual Review of Biochemistry* **87**:677–696. doi:10.1146/annurev-biochem-062917-012848

Maier T, Schmidt A, Güell M, Kühner S, Gavin AC, Aebersold R, Serrano L. 2011. Quantification of mRNA and protein and integration with protein turnover in a bacterium. *Molecular Systems Biology* **7**:1–12. doi:10.1038/msb.2011.38

Maurizi MR. 1992. Proteases and protein degradation inEscherichia coli. *Experientia* **48**:178–201. doi:10.1007/BF01923511

McGinness KE, Baker TA, Sauer RT. 2006. Engineering Controllable Protein Degradation. *Molecular Cell* **22**:701–707. doi:10.1016/j.molcel.2006.04.027

Morreale FE, Kleine S, Leodolter J, Junker S, Hoi DM, Ovchinnikov S, Okun A, Kley J, Kurzbauer R, Junk L, Guha S, Podlesainski D, Kazmaier U, Boehmelt G, Weinstabl H, Rumpel K, Schmiedel VM, Hartl M, Haselbach D, Meinhart A, Kaiser M, Clausen T. 2022. BacPROTACs mediate targeted protein degradation in bacteria. *Cell* **185**:2338-2353.e18. doi:10.1016/j.cell.2022.05.009

Nihongaki Y, Kawano F, Nakajima T, Sato M. 2015. Photoactivatable CRISPR-Cas9 for optogenetic genome editing. *Nature Biotechnology* **33**:755–760. doi:10.1038/nbt.3245

Okusu H, Ma D, Nikaido H. 1996. AcrAB efflux pump plays a major role in the antibiotic resistance phenotype of Escherichia coli multiple-antibiotic-resistance (Mar) mutants. *Journal of Bacteriology* **178**:306–308. doi:10.1128/jb.178.1.306-308.1996

Olson EJ, Hartsough LA, Landry BP, Shroff R, Tabor JJ. 2014. Characterizing bacterial gene

circuit dynamics with optically programmed gene expression signals. *Nature Methods* **11**:449–455. doi:10.1038/nmeth.2884

Pudasaini A, El-Arab KK, Zoltowski BD. 2015. LOV-based optogenetic devices: Light-driven modules to impart photoregulated control of cellular signaling. *Frontiers in Molecular Biosciences* **2**:1–15. doi:10.3389/fmolb.2015.00018

Renicke C, Schuster D, Usherenko S, Essen LO, Taxis C. 2013. A LOV2 domain-based optogenetic tool to control protein degradation and cellular function. *Chemistry and Biology* **20**:619–626. doi:10.1016/j.chembiol.2013.03.005

Sarria S, Bartholow TG, Verga A, Burkart MD, Peralta-Yahya P. 2018. Matching Protein Interfaces for Improved Medium-Chain Fatty Acid Production. *ACS Synthetic Biology* **7**:1179–1187. doi:10.1021/acssynbio.7b00334

Sarria S, Kruyer NS, Peralta-Yahya P. 2017. Microbial synthesis of medium-chain chemicals from renewables. *Nature Biotechnology* **35**:1158–1166. doi:10.1038/nbt.4022

Schrader EK, Harstad KG, Matouschek A. 2009. Targeting proteins for degradation. *Nature Chemical Biology* **5**:815–822. doi:10.1038/nchembio.250

Sheets MB, Wong WW, Dunlop MJ. 2020. Light-Inducible Recombinases for Bacterial Optogenetics. *ACS Synthetic Biology* **9**:227–235. doi:10.1021/acssynbio.9b00395

Shin J, Zhang S, Der BS, Nielsen AA, Voigt CA. 2020. Programming Escherichia coli to function as a digital display . *Molecular Systems Biology* **16**:1–12. doi:10.15252/msb.20199401

Stricker J, Cookson S, Bennett MR, Mather WH, Tsimring LS, Hasty J. 2008. A fast, robust and tunable synthetic gene oscillator. *Nature* **456**:516–519. doi:10.1038/nature07389

Strickland D, Lin Y, Wagner E, Hope CM, Zayner J, Antoniou C, Sosnick TR, Weiss EL, Glotzer M. 2012. TULIPs: Tunable, light-controlled interacting protein tags for cell biology. *Nature Methods* **9**:379–384. doi:10.1038/nmeth.1904

Strickland D, Moffat K, Sosnick TR. 2008. Light-activated DNA binding in a designed allosteric protein. *Proceedings of the National Academy of Sciences of the United States of America* **105**:10709–10714. doi:10.1073/pnas.0709610105

Strickland D, Yao X, Gawlak G, Rosen MK, Gardner KH, Sosnick TR. 2010. Rationally improving LOV domain-based photoswitches. *Nature Methods* **7**:623–626. doi:10.1038/nmeth.1473

Striebel F, Kress W, Weber-Ban E. 2009. Controlled destruction: AAA+ ATPases in protein degradation from bacteria to eukaryotes. *Current Opinion in Structural Biology* **19**:209–217. doi:10.1016/j.sbi.2009.02.006

Swartz TE, Corchnoy SB, Christie JM, Lewis JW, Szundi I, Briggs WR, Bogomolni RA. 2001. The Photocycle of a Flavin-binding Domain of the Blue Light Photoreceptor Phototropin. *Journal of Biological Chemistry* **276**:36493–36500. doi:10.1074/jbc.M103114200

Szydlo K, Ignatova Z, Gorochowski TE. 2022. Improving the Robustness of Engineered Bacteria to Nutrient Stress Using Programmed Proteolysis. *ACS Synthetic Biology* **11**:1049–1059. doi:10.1021/acssynbio.1c00490

Tague N, Andreani V, Fan Y, Timp W, Dunlop MJ. 2023. Comprehensive screening of a light-inducible split Cre recombinase with domain insertion profiling. *bioRxiv*. doi:https://doi.org/10.1101/2023.05.26.542511

Torella JP, Ford TJ, Kim SN, Chen AM, Way JC, Silver PA. 2013. Tailored fatty acid synthesis via dynamic control of fatty acid elongation. *Proceedings of the National Academy of Sciences of the United States of America* **110**:11290–11295. doi:10.1073/pnas.1307129110

Trauth J, Scheffer J, Hasenjäger S, Taxis C. 2019. Synthetic Control of Protein Degradation during Cell Proliferation and Developmental Processes. *ACS Omega* **4**:2766–2778. doi:10.1021/acsomega.8b03011

Turner WJ, Dunlop MJ. 2015. Trade-Offs in Improving Biofuel Tolerance Using Combinations of Efflux Pumps. *ACS Synthetic Biology* **4**:1056–1063. doi:10.1021/sb500307w

Wang H, Vilela M, Winkler A, Tarnawski M, Schlichting I, Yumerefendi H, Kuhlman B, Liu R, Danuser G, Hahn KM. 2016. LOVTRAP: An optogenetic system for photoinduced protein dissociation. *Nature Methods* **13**:755–758. doi:10.1038/nmeth.3926

Wang T, Tague N, Whelan SA, Dunlop MJ. 2021. Programmable gene regulation for metabolic engineering using decoy transcription factor binding sites. *Nucleic Acids Research* **49**:1163–1172. doi:10.1093/nar/gkaa1234

Xue G, Wang K, Zhou D, Zhong H, Pan Z. 2019. Light-Induced Protein Degradation with Photocaged PROTACs. *Journal of the American Chemical Society* **141**:18370–18374. doi:10.1021/jacs.9b06422

Yamamoto A, Iwata T, Sato Y, Matsuoka D, Tokutomi S, Kandori H. 2009. Light signal transduction pathway from flavin chromophore to the Jα helix of Arabidopsis phototropin1. *Biophysical Journal* **96**:2771–2778. doi:10.1016/j.bpj.2008.12.3924

Yao X, Rosen MK, Gardner KH. 2008. Estimation of the available free energy in a LOV2-Jα photoswitch. *Nature Chemical Biology* **4**:491–497. doi:10.1038/nchembio.99

Ye Z, Li S, Hennigan JN, Lebeau J, Moreb EA, Wolf J, Lynch MD. 2021. Two-stage dynamic deregulation of metabolism improves process robustness & scalability in engineered E. coli. *Metabolic Engineering* **68**:106–118. doi:10.1016/j.ymben.2021.09.009

Yumerefendi H, Wang H, Dickinson DJ, Lerner AM, Malkus P, Goldstein B, Hahn K, Kuhlman B. 2018. Light-Dependent Cytoplasmic Recruitment Enhances the Dynamic Range of a Nuclear Import Photoswitch. *ChemBioChem* **19**:1319–1325. doi:10.1002/cbic.201700681

Zayner JP, Antoniou C, French AR, Hause RJ, Sosnick TR. 2013. Investigating models of protein function and allostery with a widespread mutational analysis of a light-activated protein. *Biophysical Journal* **105**:1027–1036. doi:10.1016/j.bpj.2013.07.010

Zhu L, McNamara HM, Toettcher JE. 2023. Light-switchable transcription factors obtained by direct screening in mammalian cells. *Nature Communications* **14**. doi:10.1038/s41467-023-38993-6

Zoltowski BD, Motta-Mena LB, Gardner KH. 2013. Blue light-induced dimerization of a bacterial LOV-HTH DNA-binding protein. *Biochemistry* **52**:6653–6661. doi:10.1021/bi401040m

Zoltowski BD, Vaccaro B, Crane BR. 2009. Mechanism-based tuning of a LOV domain photoreceptor. *Nature Chemical Biology* **5**. doi:10.1038/nchembio.210

Figures

Figure 1. Design of *As*LOV2-based degradation tag. **(a)** Primary sequence of *As*LOV2(546) C-terminal sequence. A three amino acid truncation exposes E-A-A. **(b)** Structure of *As*LOV2

(aa404-546, PDB: 2V1A). Amino acids 541-543 (E-A-A) are red and 544-546 (K-E-L) are gray at the C-terminal of the Jα helix. **(c)** Construct used to characterize optogenetic control using *As*LOV2 variants. Each variant is translationally fused to mCherry expressed from an IPTG inducible promoter. Variants include wild type *As*LOV2 (light blue) and a dark state stabilized version, AsLOV2* (dark blue), with and without the three amino acid truncation. **(d)** mCherry protein levels in response to 465 nm blue light for wild type *As*LOV2, and mutated *As*LOV2* fusions with and without truncation. *As*LOV2*(543) is the variant we denote the 'LOVdeg' tag. (***p < 0.0001; **p < 0.001; *p<0.01; n.s., not significant; two tailed unpaired t-test; n = 3 biological replicates). **(e)** mCherry-LOVdeg in response to variable light intensities. **(f)** mCherry fluorescence levels and optical density of mCherry-LOVdeg with 4 hours of 465 nm blue light exposure applied at different points in the growth cycle. Light exposure programs are plotted above each subplot and are staggered 2 hours apart (starting at 2, 4, 6, or 8 hours), all lasting 4 hours. Expression levels are normalized to the dark state control (Fig. S8). Error bars show standard deviation around the mean.

Figure 2. Incorporating light responsiveness into diverse proteins with the LOVdeg tag. **(a)** Control of mCherry repression using a LacI-LOVdeg fusion. **(b)** mCherry expression in response to light exposure for strains with LacI-LOVdeg compared to IPTG induction (**$p < 0.001$, two tailed unpaired t-test). **(c)** Schematic of SoxS-based CRISPRa activation with a LOVdeg tag appended to the MCP-SoxS activator. **(d)** CRISPRa control of mRFP1 expression in response to light (***$p < 0.0001$, two tailed unpaired t-test). **(e)** Schematic of the LOVdeg tag appended to AcrB of the AcrAB-TolC efflux pump. IM, inner membrane; OM, outer membrane. **(f)** Chloramphenicol sensitivity tests. Wild type cells (BW25113) are compared to a $\Delta acrB$ (BW25113 $\Delta acrB$) strain, $\Delta acrB$ complemented with AcrB-LOVdeg ($\Delta acrB$ + AcrB-LOVdeg) exposed to light or kept in the dark, and $\Delta acrB$ strain complemented with an IPTG-inducible AcrB ($\Delta acrB$ + AcrB). No IPTG was added to $\Delta acrB$ + AcrA or $\Delta acrB$ + AcrB-LOVdeg. **(g)** OD_{600} of strains shown in (f) at 2.5 µg/ml chloramphenicol (**$p < 0.001$; *$p < 0.05$; ns, not significant; two tailed unpaired t-test). Error bars show standard deviation around the mean (n = 3 biological replicates).

Figure 3. Modulating LOVdeg frequency response with photocycle mutations. **(a)** Photocycle of *As*LOV2. Upon light absorption, the Jα helix unfolds for a period of time dictated by the stability of the light state conformation. If not degraded, the Jα helix refolds, blocking degradation. **(b)** The light program used to test frequency responses of LOVdeg photocycle variants in (c). A constant pulse of 5 sec is followed by a variable dark time that allows for Jα helix refolding. **(c)** Expression of mCherry-LOVdeg and variant mCherry-LOVdeg (V416I) in response to different light exposure frequencies. Fluorescence values are normalized to dark state expression. Error bars show standard deviation around the mean (n = 3 biological replicates).

Figure 4. Enhanced light response using EL222 transcriptional control together with LOVdeg. **(a)** mCherry-LOVdeg expressed from an EL222 responsive promoter that is constitutively active in the absence of EL222. Addition of EL222 represses mCherry production. These two forms of regulation are combined when mCherry-LOVdeg is expressed from an EL222 responsive promoter, resulting in a circuit that both degrades and represses in response to light. **(b)** Light and dark expression of mCherry in the 'degradation only' (closed circles), 'repression only' (open circles), or 'repression + degradation' (squares) strains. Error bars show standard deviation around the mean (n = 3 biological replicates).

Figure 5. Optogenetic control of octanoic acid production. **(a)** Schematic of fatty acid synthesis in *E. coli*. *Cp*FatB1* catalyzes elongating C8-ACP molecules from this pathway to produce free octanoic acid. *Cp*FatB1* is tagged with a LOVdeg tag to create optogenetic control. **(b)** Octanoic acid titer from strains that express *Cp*FatB1* only, *Cp*FatB1*-LOVdeg only, EL222 regulated *Cp*FatB1* only, or *Cp*FatB1*-LOVdeg + EL222. Octanoic acid is quantified by GC-MS. Strains were kept either in the dark or with continuous blue light exposure for the duration of the production period. Error bars show standard deviation around the mean (****$p < 0.0001$; ***$p < 0.0001$; ns, not significant; two tailed unpaired t-test; $n = 3$ biological replicates).