Kiwa is a bacterial membrane-embedded defence supercomplex activated by phage-induced membrane changes

Zhiying Zhang[1,8], Thomas C. Todeschini[2,8,9], Yi Wu[2,8], Roman Kogay[3], Ameena Naji[2,10], Joaquin Cardenas Rodriguez[2], Rupavidhya Mondi[2,11], Daniel Kaganovich[2], David W. Taylor[4-7], Jack P. K. Bravo[4,12], Marianna Teplova[1], Triana Amen[2], Eugene V. Koonin[3], Dinshaw J. Patel[1,*], Franklin L. Nobrega[2,*]

[1]Structural Biology Program, Memorial Sloan Kettering Cancer Center, New York, NY, USA

[2]School of Biological Sciences, University of Southampton, SO17 1BJ Southampton, UK

[3]National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, Maryland, USA.

[4]Department of Molecular Biosciences, University of Texas at Austin, Austin, TX, 78712, USA

[5]Interdisciplinary Life Sciences Graduate Programs, Austin, TX 78712, USA

[6]Center for Systems and Synthetic Biology, University of Texas at Austin, Austin, TX, 78712, USA

[7]LIVESTRONG Cancer Institutes, Dell Medical School, Austin, TX, 78712, USA

[8]Shared first authors

[9]Current address: RNA Therapeutics Institute, University of Massachusetts Chan Medical School, Worcester, MA, USA

[10]Current address: School of Molecular and Cellular Biology, University of Leeds, Leeds LS2 9JT, UK

[11]Current address: The William Harvey Research Institute, Barts and The London

School of Medicine, Queen Mary University of London, Charterhouse Square,

London EC1M 6BQ, UK

[12]Current address: Institute of Science and Technology Austria (ISTA),

Klosterneuburg, Austria

*Correspondence: pateld@MSKCC.ORG, F.Nobrega@soton.ac.uk

Summary

Bacteria and archaea deploy diverse, sophisticated defence systems to counter virus

infection, yet many immunity mechanisms remain poorly understood. Here, we

characterise the Kiwa defence system as a membrane-associated supercomplex

that senses changes in the membrane induced by phage infection and plasmid

conjugation. This supercomplex, comprising KwaA tetramers bound to KwaB dimers,

as its basic repeating unit, detects structural stress via KwaA, activating KwaB,

which binds ejected phage DNA through its DUF4868 domain, stalling phage DNA

replication forks and thus disrupting replication and late transcription. We show that

phage-encoded DNA mimic protein Gam, which inhibits RecBCD, also targets Kiwa

through KwaB recognition. However, Gam binding to one defence system precludes

its inhibition of the other. These findings reveal a distinct mechanism of bacterial

immune coordination, where sensing of membrane disruptions and inhibitor

partitioning enhance protection against phages and plasmids.

Keywords: Kiwa, transmembrane supercomplex, RecBCD, membrane disruptions,

DNA binding, replication fork, late transcription, phage defence, DNA mimic

Introduction

Bacteria and archaea have evolved an enormously diverse repertoire of mechanisms to defend against virus predation[1-3]. Among these defence mechanisms, detailed analysis of widespread defence systems, such as CRISPR-Cas and restriction modification not only illuminated fundamental biological processes but allowed adaption of these systems for transformative applications in biotechnology, such as gene editing, expression regulation, and RNA knockdown exploiting CRISPR effectors Cas9, Cas12, and Cas13[4].

However, the molecular mechanisms of many prokaryotic defence systems, particularly those discovered recently[5-13], remain poorly (if at all) understood. This knowledge gap is especially pronounced for membrane-associated systems. The Kiwa defence system is one such example. Kiwa consists of two components: KwaA, an integral membrane protein with four transmembrane helices, and KwaB, which contains a DUF4868 domain with an unknown function. Studies on the Kiwa system from *Escherichia coli* O55:H7 RM12579 have demonstrated its ability to protect against phages such as Lambda and SECphi18[6]. Notably, Kiwa is among the 20 most abundant defence systems across sequenced bacterial genomes[1], highlighting its widespread utility and hence evolutionary success. However, the molecular mechanisms through which Kiwa detects phage infection and provides protection remain elusive.

In this work, we provide mechanistic insights into the Kiwa defence system, revealing that it operates as a transmembrane supercomplex comprising KwaA tetramer bound to KwaB dimers, as its basic repeating unit. This supercomplex detects membrane disruption caused by phage infection and plasmid conjugation,

specifically during or prior to phage DNA ejection. Upon activation, KwaB binds

phage DNA via its DUF4868 domain, stalling phage replication forks and disrupting

phage replication and late transcription. Although DNA binding by KwaB is not

sequence-specific *in vitro,* its membrane localisation through KwaA enables a

proximity-driven mechanism that minimises off-target effects and prevents host self-

damage. Additionally, we demonstrate that Kiwa and RecBCD cooperate to

counteract phage-encoded inhibitors, such as the DNA mimic protein Gam, which

can only inhibit one of these systems at a time. This interplay underscores the

versatility and resilience of bacterial immune networks, offering a perspective on how

bacteria thwart phage-encoded inhibitors while safeguarding their cellular functions.

Results

Kiwa provides broad protection against phages and plasmids

To investigate the diversity and abundance of Kiwa, we first searched for Kiwa

components in 223,143 prokaryotic genomes from the precomputed annotation at

the PADLOC web server[13,14], which is based on RefSeq v209. Kiwa was identified in

9,582 genomes, including 1,713 *E. coli* genomes. We then dereplicated KwaA and

KwaB protein sets with a 90% sequence identity threshold and reconstructed the

phylogenies of each of these proteins. We found that, although deep branches were

not fully resolved, both KwaA and KwaB could be subdivided into three distinct

clusters (**Figure 1A**), where the previously described Kiwa operon from *E. coli*

O55:H7 RM12579 falls into cluster 1 (purple). The taxonomic assignments of the

dereplicated dataset suggest that Kiwa is particularly prevalent in species of the

orders *Bacillales* and *Enterobacterales* (**Figure 1B**). Notably, the phylogenetic

history of Kiwa substantially deviates from that of their host species, indicating a

major role of horizontal gene transfer (HGT) in shaping its evolution (**Figure 1A**).

Furthermore, the partial incongruency between the phylogenies of KwaA and KwaB

suggests that, on some occasions, *kwaA* and *kwaB* genes were transferred

independently (**Figure 1B**).

To identify mobile genetic elements (MGE) that might drive HGT of Kiwa operons,

we systematically analysed their genomic neighbourhoods. We found that 673

dereplicated *kwaA* (53.5%) and 676 dereplicated *kwaB* (53.6%) genes co-localised

with MGEs including 197/198 prophages, 105/110 plasmids, and 304/302 genomic

islands for KwaA and KwaB, respectively (**Figure 1A**). In 70 of the 9,797 Kiwa loci,

including 19 of the 1,262 in the dereplicated dataset, KwaB is represented by two

distinct proteins. The phylogenetic tree topology indicates that KwaB was split into

two genes on multiple, independent occasions (**Figure 1A**, right).

Prior studies suggest that diverse variants of the same defence system can have

different phage specificities[5,15,16]. To determine if the diversity of Kiwa impacts

defence specificity, we cloned five Kiwa systems from different phylogenetic clusters

into *E. coli* BL21-AI, a strain naturally lacking Kiwa (**Figure 1C**, **Figure S1A**). When

we exposed these strains to a panel of eight phages, all Kiwa systems demonstrated

anti-phage activity, each with unique phage specificity and varying levels of

protection (**Figure 1D**, **Figure S1B**). We also observed that Kiwa reduced plasmid

conjugation efficiency more than twofold, with the exception of Kiwa from *Ralstonia*

mannitolilytica strain SN82F48 (**Figure 1E**).

Overall, these findings indicate that Kiwa broadly, albeit with varying specificity,

defends bacteria against both phage infections and plasmid conjugation.

Figure 1 Diversity of Kiwa systems and patterns of defence against phage infection and plasmid conjugation.

(A) Phylogenetic tree of KwaA (left) and KwaB (right) proteins from the dereplicated dataset. Each cluster is identified by a specific colour shading and numbering. The tree includes annotations for Kiwa operons that were tested in this study, and Kiwa operons that have previously been confirmed to provide anti-phage activity are indicated with a shield symbol. Instances of KwaB split into two genes are annotated in red. The tree scale is in number of substitutions per site and nodes with bootstrap support of at least 80% are highlighted in cyan. Co-localisation of KwaA and KwaB with various classes of MGEs is shown in the out circles. The category "Other mobilome" denotes Kiwa genes that

are encoded in close proximity to at least 2 hallmark mobilome genes, as identified by COG

annotations[17], but were not classified as plasmids, prophages, or genomic islands.

(B) Distribution of KwaA and KwaB proteins across 20 bacterial orders containing the largest number

of dereplicated Kiwa components. The bacterial species tree was retrieved from GTDB database[18],

and the scale is in substitutions per site. The barplot was coloured based on phylogenic clusters

shown in (A). Numbers in the KwaB barplot indicate instances of KwaB split into two genes.

(C) Schematic representation of the Kiwa operon obtained from *Escherichia coli* strain ECOR49. TM,

transmembrane domain. DUF, domain of unknown function.

(D) Anti-phage activity of five Kiwa operons shown as fold protection of efficiency of platting assays

versus control cells expressing mVenusYFP (YFP). The Kiwa operons were obtained from *E. coli*

strains ECOR8, ECOR12, ECOR49, and D9, as well as *Ralstonia mannitolilytica* strain SN82F48. m,

plaque morphology change relative to the control strain.

(E) Anti-conjugation activity of five Kiwa operons shown as percentage of conjugants in relation to

total cell count that have uptaken plasmid pSEVA637 from S17-1 donor cells. Bar graphs represent

the average of at least four replicates, with individual data points overlaid.

KwaA forms a structurally variable homotetramer

To gain insight into the potential functions of Kiwa components, we first

characterised the structure of the transmembrane protein KwaA. Gel filtration

indicated that KwaA form an oligomer (**Figure S2A**). Subsequent investigation using

cryogenic electron microscopy (cryo-EM) in Lauryl Maltose Neopentyl Glycol

(LMNG) detergent revealed two distinct structural states for KwaA, with overall

resolutions of 3.7 Å and 4.2 Å, respectively (**Figures S2B,C**; cryo-EM statistics in

Table S1). These maps allowed us to build a model of the full-length KwaA protein.

Side chains were well-defined in the higher resolution 3.7 Å map but less so in the

4.2 Å map (**Figure S2D**).

In both structural states, KwaA forms a homotetramer with an hourglass-like shape

(**Figures 2A,B**), which shares similarities with the chloride ion channel TMEM16A. In

the absence of calcium ions, TMEM16A homodimers adopt an hourglass-shaped

configuration, forming a closed channel. Calcium ion binding near the channel

induces a conformational change that opens the channel, allowing negatively

charged chloride ions to pass through the membrane[19]. This observation suggests

that Kiwa, like TMEM16A, could function as a channel that opens in response to a

specific trigger.

Each KwaA protomer contains four transmembrane helices (TM1-4), two

extracellular loops (ECL1-2), one intracellular loop (ICL1), and the intracellular C-

terminus, with ICL1 and the C-terminus forming a beta-barrel structure consisting of

six beta strands (**Figure 2C**). The two structural states of KwaA differ in their

symmetry: the 3.7-Å state exhibits C2 symmetry (**Figure 2A**), whereas the 4.2-Å

state has C4 symmetry (**Figure 2B**). In the C2 symmetric structure, the four KwaA

protomers adopt two distinct conformations. One conformation resembles the

protomers in the C4 symmetric structure (**Figure S2E**, right panel), whereas the

other conformation shows notable differences, particularly in the ECL2 and the TM4

regions. In this distinct conformation, TM4 shifts closer to TM3, with a rotation of 30

Å (**Figure S2E**, left panel). This structural rearrangement widens the central pore

and repositions the surrounding helices. In the C2-symmetric structure (**Figure 2A**),

the pore is formed by TM3 and TM4 from two protomers, along with TM3 from the

other two protomers (**Figure 2D**), whereas in the C4 symmetric structure (**Figure**

2B), the pore is exclusively formed by TM3 from all protomers (**Figure 2E**).

Structural alignment of the two states reveals a common interface, designated

interface I, which involves interactions between TM1 and TM3 of one protomer and

TM3 and TM4 of an adjacent protomer, stabilised by hydrophobic interactions and π-

π stacking (**Figure 2F, Figure S2F**). The C2 symmetric structure features an

additional unique interface, referred to as interface II (**Figure 2D**), where TM4 from

one protomer interacts extensively with TM1 and TM3 of another protomer (**Figure**

S2F). This interface exhibits similar hydrophobic and π-π interactions as observed in

interface I, emphasizing a conserved assembly mechanism of the KwaA tetramer.

The reorganization of ECL2 in the C2-symmetric structure enables connection

between two diagonal protomers.

In the expanded view of the central channel in the C2-symmetric structure (**Figure**

S2G), we observed a LMNG detergent molecule within the central pore (**Figure 2A**).

The two hydrophilic maltoside groups extend outward from the cell membrane,

making no distinct interactions with the surrounding KwaA residues, whereas the two

hydrophobic chains are deeply embedded into the pore, interacting with nearby

hydrophobic KwaA residues (**Figure S2G**).

In summary, the transmembrane KwaA proteins of Kiwa form a homotetramer with

two distinct symmetries.

Figure 2. Structures of KwaA and KwaB.

(A) Two views of the cryo-EM structure of the KwaA tetramer (monomers in purple and orange) adopting C2 symmetry, with an LMNG (boxed segment) inserted into the central channel.

(B) Two views of the cryo-EM structure of the KwaA tetramer (monomers in purple and orange) adopting C4 symmetry.

(C) Schematic alignment of transmembrane helices 1 to 4, extracellular loops (ECL1, ECL2), intracellular loop (ICL), and intracellular beta-barrel of a KwaA monomer.

(D) Helical segments lining the central channel in KwaA tetramer with C2 symmetry, with labelling of interfaces I and II.

(E) Helical segments lining the central channel in KwaA tetramer with C4 symmetry, with labelling of interface I.

(F) Structural superposition of helical segments of KwaA tetramers with C2 and C4 symmetry.

(G) The crystal structure of KwaB. KwaB adopts a dimer of dimers topology with individual dimers

coloured in shades of magenta and blue. The expanded boxed segment shows the interface between

the two dimers primarily mediated by the C-terminal domain (CTD) of one protomer (in magenta) and

the N-terminal α2 helix of another monomer (in blue).

(H) Ribbon representation of the KwaB dimer (monomers in blue and green) observed in the crystal.

KwaB forms a dimer of dimers

We next determined the structure of KwaB. The crystal structure at 3.9 Å resolution

revealed that KwaB adopts a dimer of dimers topology (**Figure 2G**, **Table S2**). In this

configuration, extensive interactions occur between the C-terminal region of one

monomer from one dimer (shown in magenta) and the α2 helix of one monomer from

another dimer (shown in blue) (**Figure 2G**, expanded view).

KwaB consists of two distinct structural domains: the N-terminal region (residues 1-

185), which forms a Rossmann fold-like structure, and the C-terminal region

(residues 186-315) that contains a functionally uncharacterised DUF4868 domain

(**Figure 2G**, **Figure S3A**). The Rossmann fold-like structure is characterised by a

repeating pattern of β-strands alternating with α-helices, forming a three-layered

sandwich with a central parallel β-sheet in a βαβ motif. This fold is commonly found

in proteins that bind nucleotides including NAD, FAD, oligoA and others, and is one

of the most prevalent protein folds, capable of accommodating diverse ligands[20,21].

Both domains contribute to the formation of the KwaB dimer (**Figure 2H**, interactions

shown in detail in **Figure S3A,B**).

Our phylogenetic analysis showed that in some lineages, KwaB is split into two

distinct genes (**Figure S1A**). To investigate how this affects the functional

architecture of KwaB, we predicted the structures of the split proteins using

AlphaFold2 and compared them to the resolved structure of single-protein KwaB.

This comparison revealed that the split KwaB proteins are closely similar to the two

domains of the single-protein KwaB (Rossmann-like fold and DUF4868) (**Figure**

S3C-E). Additionally, AlphaFold2 modelling of multimeric structures showed that the

tetrameric assembly of split KwaB closely resembles the dimeric architecture of

single-protein KwaB (**Figure S3F-H)**. This conservation suggests that the split KwaB

proteins evolved via multiple, independent fissions of the ancestral single-protein

counterparts, while retaining their domain folds and functional equivalence.

In summary, KwaB forms a robust dimer of dimers structure with multiple stabilising

interactions across both the N- and C-terminal regions.

KwaA and KwaB form a fence-like transmembrane supercomplex

To investigate the roles of KwaA and KwaB and their possible interaction in anti-

phage defence by Kiwa, we tested the effect of single-gene deletions. Deletion of

either gene resulted in loss of defence, indicating that both are essential for

protection (**Figure 3A**). Additionally, truncating the TM domains in KwaA abolished

defence, highlighting the importance of membrane localization of KwaA (**Figure 3A**).

To test complex formation between KwaA and KwaB, we co-expressed both proteins

and detected low- and high-molecular weight complexes in sizing column fractions

(**Figure S4A**), indicating that KwaA and KwaB interact to form a complex. Using

cryo-EM, we visualised the high molecular weight KwaAB complex (**Figure S4B**) and found that it adopts a fence-like, higher-order supercomplex structure (**Figure S4C**). Following multiple classification rounds and symmetry analysis, we resolved the minimal structural unit of KwaAB supercomplex at a 3.7 Å resolution (**Figures S4C,D**; cryo-EM statistics in **Table S1**). This unit features a KwaB homodimer (in green and blue) associated with a tetrameric KwaA (in orange), where each KwaA tetramer binds four KwaB dimers (**Figure 3B, S4E**). Our structural analysis revealed that the homodimeric structure of KwaB in the KwaAB complex closely resembles its standalone crystal structure (RMSD = 0.63 Å, **Figure S4F**). The KwaA monomers in the KwaAB complex all adopt a similar conformation to their C4 symmetric structure (**Figures 3B, S4C,** right panels), except in regions interacting with KwaB, where localised rearrangements occur (RMSD = 1.24 Å; **Figure S4G**).

In the KwaAB supercomplex, the dimer of dimers configuration of KwaB was not observed. Instead, each KwaB dimer was found to interact with two KwaA tetramers (**Figure 3C**). Each KwaA monomer engages both protomers of a KwaB dimer through two main interfaces. In the smaller interface (blowup box in **Figure S4H**), the flexible DE loop (residues 134-174, between β-sheets D and E) of KwaB interacts with TM2 of KwaA, where Y157 of the DE loop forms a hydrogen bond with K75 of TM2 (**Figure S4H,I**). At the lateral of the C-terminal beta-barrel, α helix 2 (α2) of KwaB lies adjacent to ICL1 of KwaA. Here, Q79 of KwaA potentially forms a hydrogen bond with N42 of KwaB, while N77 interacts with the main chain of α2, further stabilizing the interface. Triple mutations in KwaA residues involved in this interaction (K75/V76/N77, K76A/V77A/N78A in ECOR49 (71.6% similarity to KwaA from O55:H7)) abolished defence, likely by disrupting the proper KwaAB interactions (**Figure 3D**).

In the larger interface (blowup box in **Figure S4I**), extensive contacts form between

the KwaB monomer and the C-terminal beta-barrel region of KwaA, whereas a loop

near β-sheet C (residues I96-P108) of KwaB is positioned above the beta-barrel

channel. In this area, P100 from KwaB is the deepest-embedded residue, influencing

the direction of the loop. This residue is surrounded by H156, Y181 and F183 from

KwaA, forming intricate π-π interactions. H156 engages in a cation-π interaction with

R21 from beta-sheet A of KwaB, the latter likely forming a hydrogen bond with Y181

and a salt bridge with D80 of KwaA. Additionally, two charged residues, N101 and

Q103, within the same loop also contribute polar interactions, N101 with T162 and

Q103 with the main chain of KwaA. Mutations in critical KwaA (C-terminal beta-

barrel; Y181/F183/K194, Y182A/F184A/K195A in ECOR49) and KwaB

(R21/R35/Y157, R21A/K38A/Y160A in ECOR49; P100/N101/Q103,

P102A/N103A/Q104A in ECOR49) residues within this larger interface also

eliminated defence, demonstrating the key role of the interface (**Figure 3D**).

The KwaB homodimer serves as the structural core for assembling the KwaAB

supercomplex (**Figure 3C,** left panel). Aligning the basic units through KwaB dimers

surprisingly revealed that the KwaAB supercomplex adopts a bent conformation in its

transmembrane region, opposite to the curvature of the native cell membrane

(**Figure 3C**, right panel). This organization explains why the supercomplex maintains

a C4 symmetry which, unlike a C2 symmetry, is conducive to forming the higher-

order complexes required for the assembly of the membrane-spanning KwaAB

structures.

An additional band coeluting with the high-molecular weight KwaAB complex on the

gel (**Figure S4A**) corresponded to the heptameric, closed-state K^+ sensitive

mechanosensitive channel MscK[22] (**Figure S4J**). However, MscK was not required

for Kiwa-mediated defence against phage infection, as demonstrated by unchanged

activity of Kiwa in MscK-deficient cells (**Figure S4K**).

In conclusion, our findings indicate that Kiwa forms a transmembrane supercomplex

composed of KwaA tetramer binding of KwaB dimers, as its basic repeating unit,

where KwaB homodimers serve as structural anchors for the KwaAB assembly.

Figure 3. KwaA and KwaB form a supercomplex.

(A) Impact of deletions and truncations on Kiwa ECOR8 defence against phages T3 and T4 infection.

Data represents phage PFU/mL on tested strains, with control expressing YFP. Asterisks indicate a

statistically significant increase in PFU/ml compared to Kiwa (Two-way ANOVA, p <0.05). The bars

represent the average of three biological replicates with individual data points overlaid.

(B) Two views of the higher molecular weight structure of the KwaAB complex showing four KwaB

dimers (monomers in blue and green) bound to a KwaA tetramer (in orange).

(C) Structural modelling of a fence-like KwaAB supercomplex on a curved cell membrane with KwaB

dimers (in blue) bound to KwaA tetramers (in orange).

(D) Impact of point mutations on KwaA and KwaB ECOR49 defence against phage T2 infection. Data

represents phage PFU/mL on tested strains, with control expressing YFP. Asterisks indicate a

statistically significant decrease in PFU/ml compared to control (Two-way ANOVA, $p < 0.05$). The bars

represent the average of three biological replicates with individual data points overlaid.

Kiwa localises to phage infection sites and detects phage-induced membrane disturbance

To elucidate the protective mechanism of Kiwa, we investigated whether it induces

an abortive infection (whereby the defence system actively causes death of the

infected cell to block phage propagation) or mutual destruction (whereby the infected

cell dies, blocking phage propagation, not due to the activity of the defence system

but as a result of the phage life cycle) phenotype during phage infection[23,24]. We

monitored phage and cell concentrations in liquid cultures at high phage pressure

(MOI = 10) and found that Kiwa substantially reduced the amount of phage progeny

while preserving viability of most cells (**Figure 4A**). In contrast, control cells lacking

Kiwa were reduced to below the detection limit of our assay. These findings indicate

that Kiwa does not trigger an abortive infection or mutual destruction but rather

appears to protect individual cells from phage infection.

To further explore the mechanism of Kiwa defence, we isolated phage mutants

escaping protection by Kiwa (**Figure 4B**). Sequencing of these escape mutants, from

both T3 and T7Select (T7s) phages, revealed mutations in two types of genes: (1)

inhibitors of the stationary phase σ^S subunit (RpoS) of host RNA polymerase (RNAP)

(T7s gp5.7 and T3 T3p30)[25,26] and (2) tail fibre genes (gp17 in T7s and T3p48 in T3)

(**Figure 4B**, **Table S3**). To validate the role of these genes in Kiwa activation, we

expressed the respective proteins in *trans*. The RpoS inhibitors were found to restore

Kiwa protection against escape phages, but the tail fibre genes did not (**Figure 4C**).

We hypothesised that the tail fibre proteins of infecting phage particles might

externally activate Kiwa. To test this proposition, we propagated the escape mutant

T3 phages in cells expressing wild-type T3p48 *in trans*, enabling some newly

produced phage particles to incorporate the wild-type T3p48 protein. Mutants T3 n4

and n5 were chosen for their high efficiency in escaping Kiwa defence (**Figure 4B**).

Despite low tail fibre incorporation efficiency, we expected that differences in

infectivity could be detected. When these mutant T3 populations, likely containing a

small fraction of phages with wild-type T3p48, were used to challenge Kiwa-

expressing cells, mutant n5 triggered Kiwa defence. This finding confirmed that tail

fibre proteins can activate Kiwa from outside the cell (**Figure 4D**).

Given that the KwaAB transmembrane supercomplex adopts a curvature opposite to

the normal membrane curvature (**Figure 3C**) and appears to be activated externally,

we hypothesised that Kiwa sensed changes at the membrane level. For well-

characterised phages like T4, DNA ejection is known to involve the formation of a

membrane-bound channel by the phage tail, causing outward bulging of the

cytoplasmic membrane[27]. The large size and mesh-like structure of the KwaAB

supercomplex, which spans a broad membrane area, could allow Kiwa to detect

membrane curvature changes caused by phage or pilus attachment. If this

hypothesis is correct, Kiwa should co-localise with the site of phage infection. To test

this, we performed co-localisation assays using a Kiwa system with fluorescently

tagged KwaB (mLemon) and phage particles stained with DNA dye SYTOX orange.

KwaB foci consistently formed near the site of phage binding, at the cell poles,

before DNA ejection occurred, as indicated by the DNA dye remaining within the

phage capsid (**Figure 4E-G**, **Figure S5**). Quantification of phage particles on cells in

proximity to KwaB foci showed that 26 ± 3% of the phages are proximal to KwaB

(defined by the partial overlap of fluorescent intensity profiles). Given the dynamic

nature of this process, this percentage aligns with expectations. These results further

support the idea that phage binding or the initiation of DNA ejection triggers Kiwa

defence.

In summary, these findings indicate that Kiwa defence is initiated at the site of phage

attachment, likely triggered by early structural changes such as membrane curvature

during phage binding or DNA ejection.

Figure 4. Kiwa localises to phage infection sites before phage DNA ejection occurs.

(A) Phage and cell titres (PFU/ml and CFU/ml, respectively) measured at different time points post infection of control cells and Kiwa cells with phage at a multiplicity of infection (MOI) of 10. The average of three biological replicates is shown with individual data points overlaid. Asterisks indicate a

statistically significant difference in phage or cell titre compared to the titres obtained in control cells

(Two-way ANOVA, $p < 0.05$).

(B) Phage titres (PFU/ml) of wild type (T3, T7s) and mutant phages (n) in control cells expressing

YFP and in cells expressing Kiwa. The average of three biological replicates is shown, with individual

data points overlaid. Asterisks indicate a statistically significant decrease in phage titre compared to

the titre obtained against the control cells (Two-way ANOVA, $p < 0.05$). Mutations found in the escape

phages are depicted below the graphics. The full list of mutations and deletions is detailed in Table

S3.

(C) Reconstitution of T3p30 and T7s gp5.7 recovers Kiwa defence activity against phage escape

phages, while reconstitution of T3p48 and T7s gp17 did not. Asterisks indicate a statistically

significant decrease in phage titres relative to control cells (GFP) (Two-way ANOVA, $p < 0.05$). Bars

represent the average of three biological replicates with individual data points overlaid.

(D) Phage titres of T3 phage escape mutants and T3 escape mutants propagated in cells expressing

wild type T3p48 in *trans*, in control and Kiwa cells. Asterisks indicate a statistically significant

decrease in phage titres relative to control cells (Two-way ANOVA, $p < 0.05$). Bars represent the

average of three biological replicates with individual data points overlaid.

(E) Confocal microscopy images of Kiwa-expressing cells infected with phage. Fluorescently tagged

KwaB (mLemon, depicted in cyan) and SYTOX Orange-stained phage particle DNA (red) are shown.

White arrows highlight instances where KwaB foci are proximal to the phage.

(F) Enlarged view of regions showing KwaB foci forming near phage attachment sites. Observed by

confocal microscopy as in (E), these foci appear before phage DNA is (fully) ejected into the cell.

(G) Fluorescence intensity profiles of KwaB (depicted in blue) and phage (red), demonstrating

overlapping signals that indicate spatial proximity.

KwaB inhibits phage late transcription

To elucidate the role of KwaB in defence, we examined the structure of this protein

more closely. Structural analysis of KwaB (**Figure 2G,H**) revealed an N-terminal

Rossmann fold, a common structural domain that typically binds nucleotide second

messengers, triggering a conformational change in the respective proteins that

activates an effector domain[28]. Homology-based functional predictions using

SPROF-GO[29] suggested that KwaB binds nucleic acids, which is supported by

electrostatic analysis of the KwaB dimer. This analysis revealed a positively-charged

central channel between the two KwaB monomers (**Figure 5A**), suggesting that this

site might bind negatively charged molecules, such as DNA.

To test this hypothesis, we examined the DNA-binding capacity of KwaB using

electrophoretic mobility shift assays (EMSA). Band shifts confirmed that KwaB binds

both linear ssDNA and linear dsDNA (**Figure 5B**). Moreover, KwaB also binds

circular plasmid dsDNA, bacterial genomic dsDNA (gDNA), and phage gDNA

(**Figure 5C**), indicating a broad, non-specific DNA binding affinity *in vitro*.

Importantly, we observed no degradation products from either dsDNA or ssDNA

targets (**Figure S6A**), indicating that KwaB binds but does not degrade nucleic acid

substrates.

For a closer examination of KwaB-DNA interactions, we used AlphaFold3 (AF3) to

predict the structure of KwaB dimer bound to dsDNA. As anticipated, the model

demonstrated dsDNA threading through the positively charged central channel of the

KwaB dimer, where interactions with surrounding residues (R233, N134, D169,

R142, R167, K187) stabilise binding (**Figure 5D**). We observed a conformational

shift between KwaB monomers when transitioning from the apo state (**Figures 2H,**

3B) to the dsDNA-bound state (**Figures 5D**), reflecting opening of the channel to

accommodate the bound dsDNA. The residues predicted to interact with dsDNA are

located in the DUF4868 domain of KwaB, whose function remains unclear. Here, we

show that mutations of all DNA-contacting residues (m6:

R142A/R167A/K187A/R233A/N134A/D169A) severely reduced KwaB binding to

ssDNA and dsDNA (**Figure 5E**, **Figure S6B**), indicating a critical role of DUF4868 in

DNA binding.

We hypothesised that KwaB captures the ejected phage DNA at the bended

membrane, preventing access to the DNA replication machinery. To test this

hypothesis, we used confocal microscopy which confirmed that KwaB remains

membrane-associated, at the cell poles, via its interaction with KwaA throughout

phage infection (**Figure 5F**, **Figure S6C**). As the localised foci formed by

fluorescently tagged KwaB in close proximity to the site of phage DNA entry, these

findings suggest that KwaB may interact with the ejected phage DNA, interfering with

its transcription and/or replication (**Figure 4E-G**).

To explore this further, we performed RNA-seq of T2-infected Kiwa and control cells

at 15 minutes post-infection. Of the 255 phage genes, 187 showed unchanged

expression levels, whereas 18 late genes were substantially under-expressed (log2-

fold change < -1) in Kiwa-containing cells. These under-expressed genes encode

components critical for phage morphogenesis (e.g. structural proteins, terminase),

indicating specific failure of late transcription (**Figure 5G**, **Table S4**). Conversely, 59

phage genes were over-expressed in Kiwa cells. These included genes involved in

DNA replication (e.g., helicase loader, clamp loader), DNA repair and recombination

(UvsY, thymidylate synthase, RecA), DNA modification (Dam methyltransferase),

and late transcription activation (e.g., GP45.1, RpbA, RegA). For T4-like phages

such as T2, late transcription depends on ongoing DNA replication, whereas RNA

polymerase transcribes newly replicated DNA, tightly coupling replication to late

gene expression[30]. Therefore, late transcription in Kiwa-expressing cells likely fails

because KwaB DNA binding stalls replication forks, preventing the synthesis of the

DNA templates required for late-stage transcription. The local, transient ssDNA-

dsDNA junctions at replication forks may serve as specific interaction targets for

KwaB given that this protein shows affinity to both DNA forms, with a preference for

ssDNA (**Figure 5B,H**).

To directly test for replication defects, we quantified phage DNA levels in Kiwa and

control cells. Phage DNA levels were reduced by approximately 25% in Kiwa-

expressing cells compared to controls (**Figure S6D**), consistent with stalled

replication. This defect coincides with the overexpression of phage DNA repair and

recombination genes, which are essential for reassembling functional replication

forks[31]. The upregulation of these pathways suggests that phage genomes

experience replication stress, likely caused by KwaB binding to DNA which interferes

with fork progression. This replication block explains the failure to produce late

transcripts, despite the upregulation of late transcription activators and proteins

promoting replication.

In summary, these results demonstrate that KwaB binds ejected phage DNA via the

DUF4868 domain and disrupts replication fork progression, stalling phage DNA

replication, preventing late transcription, and ultimately blocking phage

morphogenesis.

Figure 5. KwaB binds to phage DNA and reduces late gene transcription.

(A) Electrostatic surface representation of KwaB dimer.

(B) Electrophoretic mobility shift assays with KwaB and a 72 nucleotide ssDNA and dsDNA target. Increasing concentrations of KwaB (μM, show on top of the gel) were incubated with 10 nM ssDNA or dsDNA.

(C) Electrophoretic mobility shift assays with KwaB and circular plasmid DNA, genomic bacterial DNA, and genomic phage DNA. 500 ng of KwaB were incubated with 50 ng DNA.

(D) Alphafold 3 (AF3) model of the KwaB dimer (monomers in blue and green) complexed with a 16-mer dsDNA, highlighting positioning of the dsDNA between two KwaB monomers. Protein-DNA contacts are highlighted in the expanded boxed segment.

(E) Impact of mutations in residues involved in KwaB-DNA contacts on KwaB binding to dsDNA.

Electrophoretic mobility shift assays were performed with mutant versions of KwaB (2.56 μM) with a

72 bp ssDNA or dsDNA target (10 nM), as in (B).

(F) KwaB forms membrane-localised foci throughout phage infection. On the left, confocal microscopy

images on the left show Kiwa cells with mLemon tagged KwaB (depicted in blue) uninfected and

infected for 30 and 60 minutes. Membrane is stained with FM4-64 (red). On the right, quantification of

KwaB foci per cell. Bars represent the average with individual points overlaid.

(G) Log fold change in gene transcription in Kiwa-expressing cells compared to control cells infected

with phage T2 at 15 min. Genes overtranscribed in Kiwa cells are shown in red and those

undertranscribed are shown in green.

(H) Fluorescence anisotropy binding assays of KwaB to a 20-bp ss- or ds-DNA substrate. Data

represents the bound fraction of substrate relative to KwaB (nM). Data shows the average and

standard deviation of three independent replicates.

Kiwa is antagonised by phage DNA mimic protein Gam

Phages often encode proteins that counteract bacterial defence systems, allowing

successful phage propagation[32,33]. Among these proteins are DNA mimics that

prevent DNA-targeting defence mechanisms from accessing phage genomes[34-36].

Because we observed DNA binding by KwaB, we hypothesised that DNA mimics

might interfere with the anti-phage activity of Kiwa.

To investigate this possibility, we tested the effect of two well-known DNA mimic

proteins – Ocr from phage T7[36] and Gam from phage lambda[37] – on Kiwa defence

against phage T4, which lacks these inhibitors. Infection assays demonstrated that

Gam inhibited Kiwa, as indicated by an approximately 100-fold decrease in

protection (**Figure 6A**). This was further validated by assays using a Lambda phage

with an inactive Gam protein[38], against which Kiwa was approximately 100-fold more

effective than against wild-type Lambda (**Figure S7A**).

To explore the interaction between Gam and KwaB, we performed BACTH

(bacterial adenylate cyclase-based two-hybrid) assays with KwaB-T18 and T25-Gam

(**Figure S7B**) and co-purification experiments (**Figure S7C**), both of which

suggested potential interaction. Although we could not resolve the structure of Gam

bound to the KwaB dimer due to sample aggregation and precipitation, AF3

modelling predicted a binding mode for the KwaB-Gam complex resembling the

interaction of KwaB with DNA (**Figure 5D**). The model shows a pair of Gam

homodimers (in yellow and magenta) inserting into the central positively charged

channel of KwaB (**Figure 6B**). As with dsDNA binding (**Figure 5D**), AF3 models

predict that Gam binding induces a conformational shift in KwaB, changing its shape

from a V- to U-shaped topology (**Figure 2H**, **Figure 6B**).

To determine if Gam competes with DNA for KwaB binding, we conducted

competition assays using KwaB, ss- or dsDNA, and Gam. For both DNA substrates,

Gam binding to KwaB abolished KwaB binding to DNA, as indicated by decreased

polarisation of the DNA target upon Gam addition (**Figure 6C**).

Together, these findings indicate that the Gam protein inhibits Kiwa defence by

binding to KwaB and blocking its access to DNA.

Figure 6. Kiwa is antagonised by phage DNA mimic protein Gam.

(A) Effect of phage T7 Ocr and Lambda Gam protein expression on Kiwa anti-phage activity against phage T4. Data on the left represents phage PFU/mL on control (YFP (Ctrl)) or Kiwa strains with indicated proteins expressed in *trans*. Asterisks indicate a statistically significant decrease of phage concentration (PFU/mL) relative to control cells expressing the same protein in *trans* (Two-way ANOVA, $p < 0.05$). Bars represent the average of three biological replicates, with individual data points overlaid. Open points indicate instances where it was not possible to determine individual phage plaques, hence a value of 1 was assumed at the corresponding dilution. On the right, individual representative images are shown of spot assays of phage T4 dilutions on Ctrl or Kiwa strains with or without Gam. Indicated is the dilution where individual plaques are observed on Kiwa Gam+, whereas they were not on Kiwa Gam-.

(B) Two views of AlphaFold3 (AF3) predicted model of the KwaB dimer (monomers in blue and green) complexed with a GamS dimer (in magenta and yellow).

(C) Fluorescence anisotropy binding of KwaB to a 20 bp ssDNA or dsDNA substate in the presence or absence of Gam. Data represents polarisation of substrate in millipolarisation (mP), showing three representatives with individual points overlaid. Experiments were

performed using 1 μM KwaB with 50 nM ss- or dsDNA and incubated with or without 10 μM

of Gam protein.

(D) Effect of RecB subunit deletion and Gam expression on Kiwa anti-phage activity against

phage T4. Data on the left represents phage PFU/mL on control or Kiwa cells with intact

RecBCD or deleted RecB (ΔB), in the absence and presence of Gam. Asterisks indicate a

statistically significant decrease of phage concentration (PFU/mL) relative to control cells on

the same host strain (Two-way ANOVA, $p < 0.05$). Bars represent the average of three

biological replicates, with individual data points overlaid. Open points indicate instances

where it was not possible to determine individual phage plaques, hence a value of 1 was

assumed at the corresponding dilution. On the right, individual representative images of spot

assays of phage T4 dilutions on ΔRecB Ctrl or Kiwa strains with or without Gam.

(E) Effect of ΔRecB and Gam expression on Kiwa anti-phage activity against phage T4Δgp2.

Data on the left represents phage PFU/mL on control or Kiwa strains with complete RecBCD

or RecB deletion (ΔB), in the presence or absence of Gam expression. Asterisks indicate a

statistically significant decrease of phage concentration (PFU/mL) relative to control cells on

the same host strain (Two-way ANOVA, $p < 0.05$). Bars represent the average of three

biological replicates, with individual data points overlaid. Open points indicate instances

where it was not possible to determine individual phage plaques, hence a value of 1 was

assumed at the corresponding dilution. On the right, individual representative images of spot

assays of phage T4Δgp2 on Ctrl or Kiwa strains with or without Gam, and with or without

RecB. Indicated is the dilution where individual plaques are observed on Kiwa Gam+ RecB-

but not on Kiwa Gam+ RecB+.

Co-occurrence of Kiwa and RecBCD enables phage defence in the presence of

Gam

The Gam protein of phage Lambda inhibits the bacterial RecBCD complex by

directly binding to the RecB subunit[39]. RecBCD plays a critical role in bacterial

recombination and immunity, degrading phage DNA[40,41], and supporting other

defence systems such as CRISPR-Cas[42] and prokaryotic Argonaute[43]. Given that

RecBCD is present in cells used to assay the inhibition of Kiwa by Gam, we

hypothesized that Kiwa was only partially inhibited due to competition between Kiwa

and RecBCD for Gam binding. To investigate this possibility, we tested the effect of

Gam on Kiwa on a ΔRecB strain, effectively removing one of Gam's targets.

Whereas Gam only partially inhibited Kiwa in RecB+ cells, it completely suppressed

Kiwa in ΔRecB cells (**Figure 6D**), indicating that the inhibitory effect of Gam on Kiwa

is enhanced when a competing target is absent. This observation was reinforced by

infection assays using a wild type Lambda and a variant with an inactive Gam[38],

where Kiwa activity remained robust in the absence of RecB against the Gam-

inactive variant but was reduced against the Gam-containing phage (**Figure S7A**).

To further assess the contribution of RecBCD to Kiwa defence, we used a T4 mutant

lacking the DNA-end protecting protein gp2, which prevents RecBCD-mediated DNA

degradation[44]. Infection assays showed that RecBCD targets T4Δgp2 as efficiently

as Kiwa does (**Figure 6E**). Furthermore, although Gam inhibited RecBCD in control

cells, Kiwa (+RecBCD) retained activity despite the presence of Gam (**Figure 6E**).

By contrast, in the absence of RecB, Gam completely abolished Kiwa protection

against T4Δgp2 (**Figure 6E**). Importantly, Gam inhibition of RecBCD and Kiwa is

structurally exclusive so that a single Gam dimer cannot inhibit both systems

simultaneously (**Figure S7D**).

Overall, our results show that, although RecBCD and Kiwa are each individually

susceptible to Gam inhibition, their coexistence confers resilience, preventing Gam

from fully suppressing their functions.

603   Discussion

In this study, we describe the structure and detailed mechanism of the Kiwa defence

system, a membrane-embedded KwaAB supercomplex found in a broad range of

bacterial genomes. Our data indicate that Kiwa appears to operate by sensing

changes in the membrane caused by phage attachment or plasmid conjugation. The

detection of membrane change, mediated by the transmembrane protein KwaA,

triggers KwaB, which binds ejected phage DNA, apparently disrupting phage DNA

replication and consequently late transcription.

Sensing of membrane changes by KwaA is central to Kiwa activation, reflecting a

broader role of membrane sensing in both prokaryotic and eukaryotic cellular

responses to structural stress[45]. The DNA-binding of KwaB, mediated by its

DUF4868 domain, is not sequence-specific *in vitro*, yet its localisation to the

membrane via KwaA enables a targeted, versatile anti-phage response that prevents

phage DNA replication and late gene transcription. This proximity-driven defence

strategy seems to limit off-target effects, sparing the host from potential self-damage.

In addition, DNA-binding by KwaB might be specifically activated by conformational

changes in the KwaAB complex, triggered by membrane distortion caused by phage

or plasmid, which would further safeguard the host against self-targeting.

A key finding of our study is the susceptibility of Kiwa to phage-encoded DNA

mimicry proteins, such as Gam from phage lambda[46], which inhibits Kiwa by binding

to KwaB. We observed that, whereas both Kiwa and RecBCD defences are

individually vulnerable to Gam inhibition, their co-existence in *E. coli* provides a layer

of resilience. Specifically, Gam only partially inhibits Kiwa in the presence of

RecBCD, suggesting that Gam proteins are functionally partitioned between these

two defence targets. In RecBCD-deficient cells, however, Gam fully suppresses

Kiwa, indicating that these overlapping defences help to prevent phage evasion of

immunity by conferring redundancy and hence robustness on the immune repertoire

of the bacterial cell.

This redundancy underscores the adaptive advantage of the presence of multiple

defence systems in bacterial cells. Overlapping defences protect against single-point

failures in the presence of phage inhibitors, preventing a single inhibitor from entirely

bypassing host immunity. While harbouring multipurpose inhibitors, such as DNA

mimics[35,47-49], might benefit phages by conserving genomic resources, the

simultaneous presence of targeted defence systems, such as RecBCD and Kiwa, in

the host cell maintains an effective barrier against such inhibitors and thus against

phage reproduction.

Altogether, our findings reveal how bacteria have evolved to detect and respond to

changes in the membrane caused by phage infection or plasmid conjugation.

Furthermore, our findings support the view that bacterial defence systems operate as

interconnected, multilayered networks rather than isolated modules, leveraging

redundancy and versatility to maintain defence efficacy amidst rapidly evolving

phage threats.

Acknowledgements

We thank Rotem Sorek from the Weizmann Institute of Science for kindly providing a Lambda phage Gam mutant, and Ian Molineux from the University of Texas for kindly providing phage T4Δgp2. We also thank YouYu from Zhejiang University-University of Edinburgh Institute and J. De La Cruz (MSK) for assistance with cryo-EM data collection, and Lyuqin Zheng (MSK) for discussions on structural analysis. We thank the Imaging and Microscopy centre (IMC) facility at the University of Southampton. This work was supported by grant RGS\R2\222312 from the Royal Society to F.L.N., Welch Foundation grant F-1938 to D.W.T., and the National Institutes of Health R35GM138348 to D.W.T. T.A. is supported by Wessex Medical Research Innovation Grant (AE06). D.J.P. is supported by NIH grant GM145888, the Maloris Foundation and Memorial Sloan-Kettering Core grant (P30-CA008748). In addition to access of the cryo-EM resources at MSKCC, some of this work was performed at the National Center for CryoEM Access and Training (NCCAT) and the Simons Electron Microscopy Center located at the New York Structural Biology Center, supported by the NIH Common Fund Transformative High Resolution Cryo-Electron Microscopy program (U24 GM129539,) and by grants from the Simons Foundation (SF349247) and NY State Assembly. This research used NSLS-II MX x-ray User Resources (FMX) of the National Synchrotron Light Source II, a U.S. Department of Energy (DOE) Office of Science User Facility operated for the DOE Office of Science by Brookhaven National Laboratory under Contract No. DE-SC0012704. The Center for BioMolecular Structure (CBMS) is primarily supported by the National Institutes of Health, National Institute of General Medical Sciences (NIGMS) through a Center Core P30 Grant (P30GM133893), and by the DOE Office

670 of Biological and Environmental Research (KP1605010). R.K. and E.V.K. are

671 supported by the Intramural Research Program of the National Institutes of Health of

672 the USA (National Library of Medicine).

674 **Author contributions**

Conceptualization, F.L.N.; Methodology; Z.Z., T.C.T., Y.W., R.K., D.K., E.V.K., D.P.,

and F.L.N.; Formal Analysis, T.C.T., T.A., R.K., F.L.N., J.P.K.B, and D.W.T.;

Investigation, Z.Z., T.C.T., Y.W., J.C.R., M.T., T.A., A.N., R.M., and J.P.K.B;

Visualization, Z.Z., T.C.T., T.A., Y.W., R.K., and T.A.; Data Curation, T.C.T., Y.W.,

F.L.N.; Writing – Original Draft, Z.Z., T.C.T., Y.W., R.K., E.V.K., D.P., and F.L.N.;

Writing – Review & Editing, D.P., E.V.K., and F.L.N.; Resources, D.K., D.W.T., D.P.,

E.V.K., and F.L.N.; Funding Acquisition, F.L.N..

683 **Declaration of interests**

684 The authors declare no competing interests.

Methods

Data availability

All unique bacterial strains, phages, and plasmids generated in this study are available from the lead contact without restriction. Raw data have been deposited at Figshare, Protein Data Base (PDB) and the Electron Microscopy Data Bank (EMDB) and are publicly available as of the date of publication.

Bacterial strains and phages

E. coli Dh5α, BL21-AI, BL21-AIΔRecB, BL21-AIΔRecC, BL21-AIΔRecD[42], KEIO BW25113 and JW0454[50] were grown at 37 °C in Lysogeny Broth (LB) for liquid cultures or LB agar (LBA, 1.5 % (w/v) agar) for solid cultures. Whenever applicable, LB was supplemented with chloramphenicol (25 µg/mL), spectinomycin (50 µg/mL), Isopropyl ß-D-1-thiogalactopyranoside (IPTG, 1 mM) or L-Arabinose (0.2 %), to ensure maintenance of plasmids or induce protein expression. Phage infection was performed in LB at 37 °C. Phages were propagated by infecting *E. coli* BL21-AI grown to an optical density at 600 nm (OD_{600}) of 0.2-0.4 with 5-10 µL of a phage lysate. Infected cultures were grown for at least 4 hours or until culture collapse. The culture was then centrifuged at 11,000 × g, 4 °C for 15 min to remove cell debris, and the supernatant was filter sterilized through a 0.22 µm filter. Phage lambda with a frameshift mutation in Gam[38] was produced in BL21-AI ΔRecB. All strains and phages used in this study are listed in **Table S5**.

Plasmid and strain construction

Kiwa systems from *E. coli* ECOR8, ECOR12, ECOR49, and D9 were amplified by Q5 polymerase (New England Biolabs) using primers indicated in **Table S6**. Kiwa from *Ralstonia mannitolilytica* SN82F48 was ordered synthetically as a gBlock

(Integrated DNA Technologies) (**Table S6**). Kiwa systems under with native

promoters were amplified from the respective host strain using primers listed in

Table S6, in reactions that added regions of homology for cloning into pUOS016

using NEBuilder Hifi DNA Assembly Mastermix. When appropriate, deletions,

mutations and truncations of Kiwa operons were engineered by around-the-horn

PCR with primers listed in **Table S6**. For protein purification, His-tagged KwaB of

ECOR8 or ECOR12 Kiwa was constructed by around the horn PCR on pUOS017 or

pUOS018 respectively, introducing a 9xHis tag at the N-terminus of KwaB with

primers listed in **Table S6**. Phage proteins gp5.7/T3p30 and gp17/T3p48 were

amplified from the genome of T7s or T3 respectively using the primers indicated in

Table S6. These proteins were cloned into pCDF1-b (Novagen) under the control of

a T7 promoter. Lambda GamL was amplified from Lambda genomic DNA and cloned

into pCDF1-b by Gibson assembly. The GamS version was constructed by around

the horn PCR. Phage T7 Ocr was cloned into pCOLA (Novagen) by around the horn

PCR on pUOS040 using primers listed in **Table S6**. All plasmids were transformed

into *E. coli* Dh5α, extracted using the GeneJET Plasmid Miniprep kit (Thermo

Fisher), verified by Sanger sequencing (Eurofins Genomics) and transformed into

electrocompetent *E. coli* BL21-AI. Plasmids built for this study are listed in **Table S7**.

Plaque assays

Phages were ten-fold serially diluted in LB and spotted on soft agar plates (0.7 %

(w/v) agar) of *E. coli* BL21-AI containing a control plasmid or individual Kiwa

systems. For spot assays in strains expressing phage proteins *in trans*, induction

was performed with 0.2% or 0.02% L-arabinose. Plates were left to dry at room

temperature and incubated overnight at 37 °C. Plaque forming units (PFUs) were

determined by counting the plaques after overnight incubation. Whenever plaques

were too small to be counted, a clear lysis area was set to be equal to 1 PFU or indicated by an open dot on the bar graph. Fold defence was determined by calculating the ratio of phage concentration of Kiwa expressing strain relative to control strains.

Time post infection assays

Overnight bacterial cultures of *E. coli* BL21-AI with a control plasmid or individual Kiwa systems were diluted 1:100 in LB medium and grown at 37 °C with agitation until an OD_{600} of ≈0.3. Cultures were normalized and infected with phage T4 at an MOI of 3. A sample was taken at different time points post infection, serially diluted, and total phages were quantified by plaque assay on a phage-sensitive bacterial lawn (*E. coli* BL21-AI + pUOS016). For colony forming units, samples were washed twice with PBS 1x, serially diluted and spotted onto LBA plates. PFUs and CFUs were counted after overnight incubation at 37 °C.

Conjugation assays

S17 cells containing plasmid pSEVA637 (donor) and BL21-AI cells containing the Kiwa systems or control plasmid (recipient) were grown overnight with antibiotics (gentamycin and chloramphenicol, respectively). Recipient cells were diluted 1:2 in LB and grown for at least 3h, while donor cells were diluted 1:100 in LB and grown until an OD_{600} of 0.3 at 37 °C, 180 rpm. 750 µl of donor cells were mixed with 250 µl of recipient cells, centrifuged at 9000 × g for 5 min, and resuspended in 25 µl of LB. The cell mixture was spotted onto LBA plates and incubated overnight at 30 °C. The cells were recovered from the plates and resuspended in 1 ml of PBS 1x. Two or ten-fold dilutions of this cell suspension were spotted onto LBA plates supplemented with chloramphenicol (for total recipient cell count) and LBA plates supplemented

759 with chloramphenicol and gentamycin (for conjugant cell count). The plates were

760 incubated at 37 °C for 18-24h and cells counted. Conjugation efficiency was

761 calculated as the ratio (%) of conjugants per total recipient cells.

DNA replication assay

Overnight bacterial cultures of *E. coli* BL21-AI with a control plasmid or individual

Kiwa systems were diluted 1:100 in LB medium and grown at 37 °C with agitation

until cultures reached an OD_{600} of ≈0.3. Cultures were normalized and infected with

phage T3 or T4 at an MOI of 3. At 0 and 30 minutes post infection, a 5 ml sample

was taken and centrifuged (15,000 × *g*, 5 min, 4 °C). The cell pellets were snap

frozen using liquid nitrogen and stored at -80 °C. Total DNA was extracted using the

GeneJET Genomic DNA Isolation kit (Thermo Fisher), using the Gram-negative

protocol. Samples were sequenced at SeqCenter (Pittsburgh, PA, USA), where

sample libraries were prepared using the Illumina DNA Prep kit and IDT 10bp UDI

indices, and sequenced on an Illumina NextSeq 2000, producing 2×151bp reads.

Demultiplexing, quality control and adapter trimming was performed with bcl-convert

(v3.9.3). Reads were aligned to *E. coli* BL21-AI and phage T3 or T4 reference

genomes (CP047231, NC_003298 and NC_000866) using MiniMap2[51] in Geneious

prime 2023.0.1. Replication efficiency was determined by calculating the increase of

phage genomes to the number of cells at time 0. The percentage was calculated by

comparison of Kiwa infected cells to control cells over time.

Bacterial two-hybrid assay

Expression plasmids were cloned by fusing the T18 or T25 fragments of adenylate

cyclase (CyaA) of *Bordetella pertussis* to either end of KwaB or the Lambda Gam

protein, using primers listed in **Table S6**. *E. coli* BTH101 cells (F- , *cya-99*, *araD139*,

galE15, *galK16*, *rpsL1* (*Str r*), *hsdR2*, *mcrA1*, *mcrB1*) were co-transformed with pairs

of T18 and T25 plasmids, or control plasmids containing the leucine zipper motif of

GCN4. Co-transformants were grown in LB supplemented with IPTG (0.5 mM),

Ampicillin (100 μg/mL) and Kanamycin (50 μg/mL) at 30 °C with agitation. Overnight

cultures were then spotted on LB agar plates supplemented with IPTG (0.5 mM), 5-

bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal, 40 μg/mL) and antibiotics.

Plates were incubated at 30°C up to 48 hours before imaging.

RecBCD mutant assays

Strains with individual deleted RecBCD subunits were transformed by

electroporation with a control or Kiwa plasmids and recovered on LBA supplemented

with kanamycin (50 μg/mL), chloramphenicol (25 μg/mL) and spectinomycin (50

μg/mL). Colonies were picked and grown overnight at 37 °C. Spot assays were

performed with phages T4, T4Δgp2, Lambda$_{vir}$, and Lambda$_{vir}$ with a frameshift

mutation in Gam.

Isolation and sequencing of escape phages

Escape phages were isolated by picking single plaques at the lowest dilution of a

spot assay on a lawn of *E. coli* cells expression Kiwa. The resistant phenotype of

each escape phage was determined by comparing the EOP of the escape phage to

the EOP of the wild type phage in the presence of Kiwa. Isolated phages were

further propagated by infecting a liquid culture of Kiwa as described above. Phage

lysates were pelleted following the NaCl/PEG8000 precipitation protocol as

previously described[52]. The phage pellets were re-suspended in LB, and treated with

DNase I and RNase A (1 μg/mL each) for 30 min at room temperature. EDTA (20

mM), proteinase K (50 μg/mL) and SDS (0.5%) were added to the suspension and

incubated overnight at 56 °C. DNA was extracted using the Zymo Clean &

Concentrator kit (Zymo Research). Samples were sequenced at SeqCenter

(Pittsburgh, PA, USA), where sample libraries were prepared using the Illumina DNA

Prep kit and IDT 10bp UDI indices, and sequenced on an Illumina NextSeq 2000,

producing 2×151bp reads. Demultiplexing, quality control and adapter trimming was

performed with bcl-convert (v3.9.3). Mutations were identified by mapping

sequencing reads to the respective reference phage genomes (NC_003298 for T3

and **Table S3** for T7s) using Breseq[53] v0.37.0 with default parameters. Only

mutations that occurred in the isolated mutant but not in the wild type phage were

considered. Silent mutations within protein-coding regions were also neglected.

Amplification of mutant phages in cells expressing wild type tail fibres

Escape mutant T3 phages were propagated in cells expressing wild type T3p48 in

trans for 4h at 37 °C, 180 rpm. The cultures were centrifuged at 9,000 × *g*, the

supernatant recovered, filtered (0.2 µm PES), and stored at 4 °C until further use.

The recovered phages were ten-fold diluted and spotted onto double layer agar with

control or Kiwa-expressing cells. The plates were incubated overnight at 37 °C for

determining phage titres.

Protein expression and purification

Overnight cultures of *E. coli* BL21-AI with N-terminal 9xHis KwaB from ECOR8 or

ECOR12 were diluted in LB supplemented with antibiotics, and grown at 37 °C with

agitation until an OD_{600} of ≈0.6. Grown cultures were incubated on ice for 1 hour and

protein expression was induced with L-arabinose (0.2 %) and IPTG (1 mM), followed

by overnight incubation at 20 °C and 150 rpm. The overnight cultures were

harvested by centrifugation (8,000 × g, 30 min, 4 °C). The supernatant was

discarded and the cell pellets were re-suspended in lysis/wash buffer (50 mL per 1 L

of initial culture, 100 mM Tris-HCl, 500 mM NaCl, 1 mM DTT, 5% glycerol, 40 mM

imidazole, pH 8.0) supplemented with Pierce™ Protease Inhibitor Tablets EDTA-free

(Thermo Fisher). Cell lysis was performed three times in a cooled French press

(15Gpa). The cell lysate was centrifuged (17,000 × g, 30 min, 4 °C) and debris were

removed by filtration (0.45 µm PES). The filtered supernatant was incubated with

HisPur™ Ni-NTA Resin (Thermo Fisher) (500 µL/50 mL lysate) pre-washed with 20

mL of lysis/wash buffer for 30 min at 4 °C. The sample was loaded onto a 5 mL

Pierce™   Disposable   Column   (Thermo   Fisher)   for   gravity-flow   affinity

chromatography. The column was washed with 15 mL of ice-cold lysis/wash buffer,

and the protein was eluted with ice-cold elution buffer (100 mM Tris-HCl, 500 mM

NaCl, 1 mM DTT, 5% glycerol, 250 mM imidazole, pH 8.0). Pooled fractions were

concentrated (Amicon Centrifugal concentrator with Ultracel membrane, 10 kDa

MWCO), and the buffer exchanged to running buffer (100 mM Tris-HCl, 150 mM

NaCl, 1 mM DTT, 5% glycerol, pH 8.0). The resulting protein sample was subjected

to size exclusion chromatography using a Superdex 200 increase 10/300 GL (Cytiva)

column equilibrated with running buffer with 0.5 mL/min flow rate using running

buffer as a mobile phase. Fractions of interest were analysed by SDS-PAGE, snap

frozen in liquid nitrogen and stored at -80 °C until further use.

KwaA expression and purification

The genes encoding KwaA (UniProt: P0DW45) and KwaB (UniProt: P0DW46) from

Escherichia coli O55 were codon-optimized and synthesized by IDT. They were then

individually cloned into the pET21a vector (Novagen) with ampicillin resistance and

the pYB100 vector (NovoPro) with kanamycin resistance. KwaA was fused to a C-

terminal Flag tag, while KwaB was fused to a C-terminal His6 tag.

The O55 KwaA membrane protein was overexpressed in C43 cells, a derivative

of *Escherichia coli* BL21(DE3), and induction was carried out using 0.5 mM IPTG

(GoldBio) at 16 °C for 20 hours. The cells were harvested by centrifugation and

resuspended in a lysis buffer (25 mM Tris, pH 8.0, 500 mM NaCl, 2 mM β-

mercaptoethanol, and cOmplete (EDTA-free protease inhibitor). Cell lysis was

achieved through sonication, after which 1% (w/v) Lauryl Maltose Neopentyl Glycol

(LMNG, Anatrace) and 0.1% (w/v) cholesteryl hemisuccinate (CHS, Anatrace) were

added to solubilize the proteins at 4 °C for 3 hours. Insoluble material was cleared by

centrifuging at 22,000 rpm for 1 hour using a JA-20 fixed-angle rotor (Avanti J-E

series centrifuge, Beckman Coulter). The resulting supernatant was incubated with

anti-FLAG M2 affinity gel (SIGMA, A2220) while rotating at 4 °C for another 3 hours.

The sample was subsequently loaded onto a gravity column and thoroughly washed

with Wash Buffer I (25 mM Tris, pH 8.0, 500 mM NaCl, 2 mM β-mercaptoethanol,

0.1% (w/v) LMNG, and 0.01% (w/v) CHS), followed by Wash Buffer II (25 mM Tris,

pH 8.0, 500 mM NaCl, 2 mM β-mercaptoethanol, 0.001% (w/v) LMNG, 0.0001%

(w/v) CHS, and 0.00033% Glyco-diosgenin (GDN, Anatrace)). The protein was then

eluted with Wash Buffer II containing 0.2 mg/ml 3× DYKDDDDK Peptide (Pierce)

while rotating for 30 minutes at 4 °C. The elution was then concentrated, and further

purified using size exclusion chromatography (SEC) on a Superose 200 Increase

10/300 GL column (GE Healthcare) equilibrated with SEC buffer (25 mM Tris, pH

8.0, 500 mM NaCl, 2 mM β-mercaptoethanol, 0.001% (w/v) LMNG, 0.0001% (w/v)

CHS, and 0.00033% GDN). Fractions eluted at around 15 ml, indicating tetrameric

assembly, were collected, concentrated, and prepared for SDS-PAGE and cryo-EM

analysis.

KwaB expression and purification

KwaB protein purification for crystallization began with its expression in BL21 (DE3)

cells, using a C-terminal His tag. Protein expression was induced with 0.5 mM IPTG,

and the culture was grown at 16 °C with shaking at 220 rpm for about 20 hours. The

cells were then harvested by centrifugation at 4,000 rpm for 15 minutes, and the

pellet was resuspended in a lysis buffer containing 25 mM Tris (pH 8.0), 500 mM

NaCl, 5% glycerol, 2 mM β-mercaptoethanol, and PMSF. After cell disruption using

sonication, the lysate was centrifuged at 22,000 rpm for 1 hour to separate the

soluble fraction. The clarified lysate was then applied to a Ni-NTA gravity column,

where the protein underwent flow-through, washing, and elution steps. The eluted

protein solution contained 25 mM Tris (pH 8.0), 150 mM NaCl, 5% glycerol, 2 mM β-

mercaptoethanol, and 500 mM imidazole. To further purify the sample and remove

any nucleic acid contaminants, it was passed through a heparin column. Finally, the

protein underwent gel filtration using a Superose 200 Increase column to exchange it

into the crystallization buffer, which contained 25 mM HEPES (pH 7.5), 150 mM

NaCl, 5% glycerol, and 2 mM DTT.

KwaAB expression and purification

For cryo-EM analysis of the KwaA and KwaB complex, KwaA_flag and KwaB-His6

were co-expressed and purified using anti-FLAG resin, following the same protocol

as that used for isolating KwaA alone. The fraction peaks from the Superose 6

Increase column revealed the high molecular weight KwaAB complex at 11.8 ml and

the low molecular weight KwaAB complex at 14.5 ml, as indicated by the SDS-PAGE

results. Additionally, the MscK protein co-eluted with the high molecular weight

complex, which was also confirmed by SDS-PAGE analysis.

Cryo-EM sample preparation and data collection

For membrane bound KwaA, the purified protein was concentrated to 4 mg/ml using

a Superose 6 Increase column. A 4 µl aliquot of the sample was applied to glow-

discharged holey gold grids (UltrAuFoil 300 mesh R1.2/1.3). The grids were blotted

for 3 seconds with a force setting of 0 at 6 °C and 100% humidity, then rapidly frozen

in liquid ethane using a Vitrobot Mark IV (FEI). Data collection was performed at the

Memorial Sloan Kettering Cancer Center (MSKCC) using a Titan Krios G2

transmission electron microscope (FEI) operating at 300 kV and equipped with a K3

direct detector, controlled by SerialEM software. Movies were recorded in super-

resolution mode with a total electron dose of 53 e^-/Å², a defocus range of -0.8 to -2.2

µm, and a pixel size of 1.064 Å. The cryo-EM sample preparation for the KwaAB

complex was performed similarly to that of KwaA alone. The peak corresponding to

the high molecular weight complex from the Superose 6 Increase column was

collected and concentrated to 4.5 mg/ml for cryo-EM analysis. Data collection was

carried out using a Titan G2 transmission electron microscope (FEI) operating at 300

kV, equipped with a K3 electron detector, and controlled by Leginon software at the

New York Structural Biology Center (NYSBC). The defocus range was set between -

0.8 to -2.2 µm, with a pixel size of 0.826 Å and a total electron dose of 58.73 e^-/Å² in

super-resolution mode.

Cryo-EM data processing

For KwaA structure determination, a total of 6,054 movies was processed using

cryoSPARC[54], Patch motion correction and patch contrast transfer function (CTF)

estimation were applied to correct for drift and estimate CTF parameters,

respectively. Micrographs with ice contamination, high astigmatism, or poor CTF fit

resolution were excluded by setting appropriate threshold ranges using the 'Manually

Curate Exposures' job. High-quality micrographs were then processed with the Blob

picker and extract jobs, resulting in the selection of 5,479,293 particles. Multiple

rounds of two-dimensional (2D) classification were conducted to eliminate junk

particles. After two rounds of ab-initio reconstruction and heterogeneous refinement

into three classes, two distinct volumes with C2 and C4 symmetries were identified.

These were then refined further through homogeneous and non-uniform refinement.

For the C2 symmetry reconstruction, 312,208 particles were used, achieving a

resolution of 3.74 Å, which enabled precise model building. In the case of C4

symmetry reconstruction, 158,542 particles were utilized, reaching a resolution of

4.29 Å, sufficient for fitting the main chain.

To determine the structure of the KwaAB complex, 5,475 micrographs were collected

using the K3 camera, and the movies were processed using the same procedure.

Following particle selection through Blob picking and template-based methods,

1,731,427 particles were chosen for 2D classification. From these 2D images, the

KwaA-KwaB complex and Msck protein were separated, resulting in 182,763

particles for the KwaAB complex and 90,497 particles for the MscK protein. The

particles from the KwaA-KwaB complex were subjected to ab-initio reconstruction to

generate initial models, followed by heterogeneous refinement to enhance the

density map's quality. The best class from the heterogeneous refinement was

selected and further polished using homogeneous and non-uniform refinement,

achieving a 3.75 Å resolution structure with C4 symmetry. This level of detail allowed

for accurate model building of the KwaA tetramer bound to KwaB dimers. For the

MscK protein group, the same protocol as the KwaAB complex was followed,

achieving a resolution of 4.14 Å, which enabled fitting the MscK heptamer in its

closed state. All the resolution estimations were based on a Fourier shell correction

of 0.143 cutoff.

Crystallisation and data collection

KwaB dimer crystals were obtained at 20 °C using the hanging-drop vapor diffusion technique, where 1 μL of the protein solution was combined with 1 μL of reservoir solution containing 0.1 M Tris (pH 8.5) and 5% (w/v) PEG8000. The crystals were cryoprotected by adding 25% glycerol to the reservoir solution before being flash cooled in liquid nitrogen. Diffraction data were collected at the FMX beamline of NSLS-II at Brookhaven National Laboratory and processed using the HKL suite (HKL Research).

Model building, structure refinement and visualization

The KwaA tetramer and KwaB dimer models were predicted using Alphafold2[55], while the MscK model was based on the published heptamer structure (PDB: 7UW5)[56]. The KwaAB model was constructed by fitting the KwaA tetramer and KwaB dimer separately, followed by structure building in Coot[57]. For KwaB crystals, the structure was solved by molecular replacement using PHASER[58], all of the atomic coordinates were refined against the map in PHENIX[59].Figures were prepared using UCSF Chimera[60] and USCF ChimeraX[61] with the final image layout created in Adobe Photoshop and Illustrator.

KwaB mutants' expression and purification

KwaB mutants targeting the DNA-binding site were designed based on the AF3 model. Two mutants, named KwaB 3-mut (R142A/R167A/K187A) and KwaB 6-mut (R142A/R167A/K187A/R233A/N134A/D169A), were constructed as follows: the genes were codon-optimized and synthesized using IDT gBlock fragments, then cloned into the pYB100 vector using the Gibson cloning kit. Expression and purification of the mutants followed the protocol described above for crystallisation of

the KwaB wild type protein. The purified proteins were concentrated to 2 mg/ml for

subsequent functional assays.

Electrophoretic mobility shift assay

For the O55 KwaB binding assays with ssDNA or dsDNA, we utilized a 72 nt

sequence

(5'TGGTTTTTATATGTTTTGTTATGTATTGTTTATTTTCCCTTTAATTTTAGGATAT

GAAAACAAGAATTTATC) as well as its complementary DNA substrates. The DNA

substrates were assembled through self-annealing in an annealing buffer containing

25 mM HEPES (pH 7.5), 150 mM NaCl, and 2 mM $MgCl_2$.For the assembly of DNA

with KwaB and mutant KwaB proteins, 10 nM DNA substrates were incorporated into

a total volume of 20 µl. The ratios of KwaB protein to dsDNA were set at 0, 1, 8, 16,

32, 64, and 128:1, while for ssDNA, the ratios were 0, 1, 4, 8, 16, 32, and 64:1. The

protein and DNA substrates were diluted in an annealing buffer that contained an

additional 5% glycerol for gel loading. After incubating the mixtures on ice for 1 hour,

10 µl of each was loaded onto a 6% DNA Retardation Gel (Invitrogen) and

electrophoresed at 100 V for 35 minutes at 4 °C in 0.5 x TBE buffer. Nucleic acids

were visualized using SYBR Gold nucleic acid gel stain (Invitrogen), incubated in the

dark for 30 minutes, and imaged using a ChemiDoc imaging system.

Additionally, pUC19 plasmid DNA, *E. coli* BL21-AI gDNA, and phage Lambda gDNA

(Thermo Fisher) were used as DNA probes for EMSA assays. 50 ng of dsDNA probe

or 150 ng ssDNA (FN0850) were mixed with purified KwaB in assembly buffer 1 (100

mM Tris-HCl, 150 mM NaCl, 1 mM DTT, 5% glycerol, 10 mM $MgCl_2$, pH 8.0) and

incubated at 37 °C for 30 minutes. For titrations, dilutions ranging from 1400 to 11 ng

of KwaB protein were mixed with the DNA probe prior to complex assembly.

Samples were mixed with blue DNA loading dye (without EDTA) (Thermo Fisher)

and loaded on a 1% agarose gel for electrophoresis. Gels were run at 100 V for 50

minutes in 1X TAE buffer. Nucleic acids were visualised with SYBRSafe or

SYBRGold staining and captured with an Invitrogen iBright 1500 apparatus (Thermo

Fisher).

To assess the impact of mutations in residues involved in the KwaB-DNA binding

surface, purified KwaB wild type (wt) protein and mutants (3-mut and 6-mut) were

concentrated to 2 mg/ml. A 128:1 ratio of KwaB proteins to DNA substrate was used,

with 50 ng of 72-nt single-stranded DNA (ssDNA) or 72-bp double-stranded DNA

(dsDNA) added to a total reaction volume of 10 µl. The mixtures were incubated at

room temperature for 30 minutes. Subsequently, 3 µl of each sample was loaded

onto a 6% DNA retardation gel (12-well), along with 0.2 µl of Quick-Load Purple Low

Molecular Weight DNA Ladder (NEB, N0557S). The gel was run at 100 V for 80

minutes at 4°C in 0.5× TBE running buffer. SYBR Gold was used to stain the gel in

the dark with gentle shaking for 30 minutes, followed by three washes with ddH2O.

The gel was imaged using a ChemiDoc imaging system.

In vitro **nuclease activity**

Purified PCR amplicons, pUC19 plasmid DNA, *E. coli* BL21-AI gDNA, phage

Lambda gDNA (Thermo Fisher) and oligos were used as substrates for cleavage

assays. Target cleavage assays were performed in a 10 µl reaction mixture

containing 50 ng dsDNA or 600 ng ssDNA (FN0850) substrate and 500 ng protein in

cleavage buffer (100 mM Tris-HCl, 150 mM NaCl, 1 mM DTT, 5% glycerol, pH 8.0)

supplemented with ddH$_2$0, MgCl$_2$ (10 mM) or MnCl$_2$ (10 mM), and ATP (1mM).

Assays were allowed to proceed at 37 °C for 2 hours and stopped by addition of

proteinase K (Thermo Fisher) and incubated at 55 °C for 15 minutes. Samples were

mixed with purple DNA loading dye (with EDTA) (NEB) and loaded on a 1% or 1.5%

(oligos only) agarose gel for electrophoresis. Gels were run at 100 V for 50 minutes

in 1X TAE buffer. Nucleic acids were visualised with SYBRSafe or SYBRGold (oligos

only) staining and captured with an Invitrogen iBright 1500 apparatus (Thermo

Fisher).

Fluorescence anisotropy binding assays

To determine DNA binding affinities of KwaB, serial two-fold dilutions of ECOR8

KwaB were incubated with 0.8 nM FAM-labelled 20-nt ssDNA (top oligo, **Table S6**)

or 20-bp dsDNA (top and bottom oligo, **Table S6**) for 3 hours at 37 °C in binding

buffer (150 mM NaCl, 25 mM HEPES pH 7.5, 0.05 Tween-20). Data were recorded

at 37 °C in a CLARIOstar Plus multi-detection plate reader (BMG Labtech) equipped

with a fluorescence polarization optical module (λ_{ex} = 485 nm; λ_{em} = 520 nm). Data

were fit in GraphPad Prism 10 using a one-step binding model, and extrapolated

start and end points were used to normalize the calculation of fraction bound. For

Gam competition binding experiments, 1 µM KwaB was preincubated with 50 nM

ss/dsDNA as described above, and then further supplemented with a final

concentration of 10 µM GamS (New England Biolabs), and then measurements were

performed as described above.

Pull-down assay for GamS and KwaB

The full-length GamS was fused to a C-terminal Flag tag and cloned into the pET21a

vector. It was then co-transfected into *E. coli* with the KwaB-His plasmid at a 4:1

molar ratio, followed by selection based on ampicillin and kanamycin resistance.The

selected colonies were then transferred to 1 L of LB liquid medium, induced, and

harvested using the same method. The bacteria were lysed with Lysis buffer (25 mM

Tris, PH 8.0, 300 mM NaCl, 5% glycerol, 2 mM DTT), followed by low-temperature

sonication and centrifugation to obtain the supernatant. This supernatant was

incubated with Flag beads for 3 hours, after which the beads were washed, and the

target protein was eluted using 3x Flag peptide. The eluate was concentrated and

loaded onto Superose 6 Increase column, and peaks 2 and 3 were collected for gel

analysis. SDS-PAGE results demonstrated the co-elution of KwaB and Gam.

Confocal microscopy

Overnight cultures of control (YFP) or Kiwa-expressing cells were sub-cultured in LB

with antibiotics for 2 hours. Agarose pads were prepared with 1% agarose in LB. For

membrane staining, cells were washed with 1× PBS and resuspended in 0.1 mM

FM4-64 (in PBS). Two µl of bacterial culture were seeded onto the agarose pads

and covered with microscope cover slips prior to imaging. Confocal images and

movies were acquired using a dual point-scanning Nikon A1R-si microscope

equipped with a PInano Piezo stage (MCL), using a 60x PlanApo VC oil objective NA

1.40. Movies and images were acquired in galvanometer scanning mode using

488nm Laser (Coherent, 50mW), and a 561 nm Laser (Coherent, 50mW). Image

processing and quantification was performed using FiJi software. The phage filtrates

were treated with DNase I (1 µg/ml each) for an hour at room temperature. SYTOX

orange was added to a final concentration of 2.5 µM, and incubated overnight at 4

°C. Excess dye was removed by centrifuging the stained lysate on a 100 kDa

Amicon Ultra (UFC5100) filter and rinsing the dye away with fresh LB four times. The

concentrated stained phages were resuspended in 1 ml of fresh LB and stored at 4

°C.

1075 RNAseq

Overnight cultures were diluted with fresh LB supplemented with antibiotics at 1:100

and incubated at 37 °C, 180 rpm for 2 hours. Phages were added to the cells at MOI

10. The cells were harvested by centrifugation after 15 minutes of incubation, and

the RNA immediately extracted with RNeasy Kit (Qiagen). Library construction, rRNA

depletion, and paired-end Illumina sequencing (Novaseq 6,000, 2 × 150 bp

configuration) were performed by Novogene.

1082 Bioinformatic analysis

Information on defence systems in 223,143 prokaryotic genomes was retrieved from

the PADLOC database[13,14]. Only complete Kiwa systems that contained both KwaA

and KwaB gene within the operon, were considered for further analysis (**Table S8**).

For phylogenetic analyses, split KwaB proteins were concatenated into a single

sequence, and KwaA and KwaB proteins were dereplicated using MMseqs2 at 90%

sequence identity threshold[62]. Contigs encoding representative KwaA and KwaB

proteins shorter than 150 and 200 amino acids, respectively, were re-examined.

Partial sequences located at the ends of contigs were discarded, and any sequences

with updated annotations in the latest RefSeq release (v227) were replaced with the

new annotations. The retained sequences (**Tables S9, S10**) were aligned using

Muscle5[63], and sites with more than 99% of gaps were trimmed using ClipKIT[64].

Phylogenetic trees were reconstructed using IQ-Tree2[65], and the optimal model was

picked using ModelFinder from the set of WAG, LG, Q.Pfam, and NQ.Pfam[66].

Support values were estimated using 1,000 ultrafast bootstrap replicates with a hill-

climbing nearest neighbour interchange optimization (UFBoot2)[67], and trees were

visualized using iTOL v6[68]. Contigs encoding KwaA and KwaB from the dereplicated

set were searched for plasmids and prophages using geNomad v1.7[69], and genomic

islands were identified using TreasureIsland with probability cutoff at 0.95[70].

Additionally, 10 genes upstream and downstream of KwaA and KwaB were

annotated using COG profiles[17]. Kiwa genes that are colocalized with at least 2

hallmark genes from MGEs (X category) were labelled as associated with 'other

mobilome' (**Tables S9, S10**).

Protein domains of KwaA and KwaB were identified using HHpred[71].

Transmembrane domains were identified using DeepTMHMM[72]. KwaA proteins with

more or less than 4 transmembrane domains were manually checked for

misannotation of the start codon. Sequence alignments of KwaA C-terminus and

KwaB were constructed using Clustal Omega[73] and visualised using Geneious Prime

v2023.0.1. For KwaB mutants, a total of nine homologs present in the KwaB ECOR8

and ECOR12 cluster were selected and aligned to determine amino acid

conservation. KwaA schematic was visualised and adapted from Protter[74]. All

structural models were built using Alphafold[75] (AF2) software run on the external

COSMIC server[76] using either the multimer or monomer model with complete

database settings. Structural similarity was estimated using DALI[77] for monomers,

and MM-align[78] for multimers. Structures were visualised using ChimeraX v1.6.1[79].

KwaB functionality was predicted using the Biomedical AI Platform webtool SPROF-

GO[29].

Genomes of *E. coli* BL-21-AI and T2 phage used in this study were annotated using

PGAP pipeline[80]. The quality of RNA-seq data was assessed using FastQC, and

low-quality tiles were removed using BBTools[81]. Reads were mapped to the genome

using HISAT2[82], and the ones corresponding to tRNA and rRNA were discarded.

TPM values were calculated using TPMCalculator[83], and for mathematical purposes,

phage genes and bacterial genes were treated as separate datasets.

QUANTIFICATION AND STATISTICAL ANALYSIS

Unless stated otherwise, experiments were performed in biological triplicates and are

represented as the mean and standard deviation. Statistical significance was

calculated by ratio paired t-test or by two-way ANOVA with sidak's multiple

comparison test, with a significance level of 0.05.

Supplemental information titles and legends

Figure S1 Similarity of Kiwa homologues and their anti-phage activity.

Figure S2 Structure analysis of KwaA.

Figure S3 Interactions between KwaB monomers in the symmetrical dimer.

Figure S4 Structure analysis of KwaAB.

Figure S5 Validation of phage staining with SYTOX Orange.

Figure S6 KwaB does not degrade DNA but binding to phage DNA prevents replication.

Figure S7 Inhibition of KwaB by Lambda Gam protein.

Table S1 Cryo-EM data collection and refinement statistics.

Table S2 Crystal data collection and refinement statistics (molecular replacement).

Table S3 Mutation analysis of phage escape mutants. Mutations marked by Breseq under "marginal evidence" are reported if the mutation was supported by more than 20% of the aligned reads.

Table S4 RNAseq data of Kiwa and control (YFP) cells infected with phage T2 for 15 min.

Table S5 Strains and phages used in this study.

Table S6 Primers and gblocks used in this study.

Table S7 Plasmids used in this study.

Table S8 Annotation of 9,797 Kiwa operons in 9,582 prokaryotic genomes based on PADLOC database.

Table S9 Annotation of mobile genetic elements in the dereplicated dataset of KwaA genes.

Table S10 Annotation of mobile genetic elements in the dereplicated dataset of KwaB genes.

References

1. Georjon, H., and Bernheim, A. (2023). The highly diverse antiphage defence systems of bacteria. Nat Rev Microbiol. 10.1038/s41579-023-00934-x.

2. Egido, J.E., Costa, A.R., Aparicio-Maldonado, C., Haas, P.-J., and Brouns, S.J.J. (2022). Mechanisms and clinical importance of bacteriophage resistance. FEMS Microbiol Rev *46*, fuab048. 10.1093/femsre/fuab048.

3. Agapov, A., Baker, K.S., Bedekar, P., Bhatia, R.P., Blower, T.R., Brockhurst, M.A., Brown, C., Chong, C.E., Fothergill, J.L., Graham, S., et al. (2024). Multi-layered genome defences in bacteria. Curr Opin Microbiol *78*, 102436. 10.1016/j.mib.2024.102436.

4. Anzalone, A.V., Koblan, L.W., and Liu, D.R. (2020). Genome editing with CRISPR–Cas nucleases, base editors, transposases and prime editors. Nat Biotechnol *38*, 824-844. 10.1038/s41587-020-0561-9.

5. Millman, A., Melamed, S., Leavitt, A., Doron, S., Bernheim, A., Hör, J., Garb, J., Bechon, N., Brandis, A., Lopatina, A., Ofir, G., Hochhauser, D., Stokar-Avihail, A., et al. (2022). An expanded arsenal of immune systems that protect bacteria from phages. Cell Host Microbe *30*, 1556-1567.e5. 10.1016/j.chom.2022.09.017

6. Doron, S., Melamed, S., Ofir, G., Leavitt, A., Lopatina, A., Keren, M., Amitai, G., and Sorek, R. (2018). Systematic discovery of antiphage defense systems in the microbial pangenome. Science *359*, eaar4120. 10.1126/science.aar4120.

7. Ofir, G., Melamed, S., Sberro, H., Mukamel, Z., Silverman, S., Yaakov, G., Doron, S., and Sorek, R. (2018). DISARM is a widespread bacterial defence

system with broad anti-phage activities. Nat Microbiol *3*, 90-98.

10.1038/s41564-017-0051-0.

8. Goldfarb, T., Sberro, H., Weinstock, E., Cohen, O., Doron, S., Charpak-

Amikam, Y., Afik, S., Ofir, G., and Sorek, R. (2015). BREX is a novel phage

resistance system widespread in microbial genomes. EMBO J *34*, 169-183.

10.15252/embj.201489455.

9. Gao, L., Altae-Tran, H., Böhning, F., Makarova, K.S., Segel, M., Schmid-

Burgk, J.L., Koob, J., Wolf, Y.I., Koonin, E.V., and Zhang, F. (2020). Diverse

enzymatic activities mediate antiviral immunity in prokaryotes. Science *369*,

1077-1084. 10.1126/science.aba0372.

10. Owen, S.V., Wenner, N., Dulberger, C.L., Rodwell, E.V., Bowers-Barnard, A.,

Quinones-Olvera, N., Rigden, D.J., Rubin, E.J., Garner, E.C., Baym, M., and

Hinton, J.C.D. (2021). Prophages encode phage-defense systems with

cognate self-immunity. Cell Host Microbe *29*, 1620-1633.e1628.

10.1016/j.chom.2021.09.002.

11. Rousset, F., Depardieu, F., Miele, S., Dowding, J., Laval, A.-L., Lieberman,

E., Garry, D., Rocha, E.P.C., Bernheim, A., and Bikard, D. (2022). Phages

and their satellites encode hotspots of antiviral systems. Cell Host Microbe *30*,

740-753.e745. 10.1016/j.chom.2022.02.018.

12. Vassallo, C.N., Doering, C.R., Littlehale, M.L., Teodoro, G.I.C., and Laub,

M.T. (2022). A functional selection reveals previously undetected anti-phage

defence systems in the E. coli pangenome. Nat Microbiol *7*, 1568-1579.

10.1038/s41564-022-01219-4.

13. Payne, L.J., Todeschini, T.C., Wu, Y., Perry, B.J., Ronson, Clive W., Fineran,

Peter C., Nobrega, Franklin L., and Jackson, Simon A. (2021). Identification

and classification of antiviral defence systems in bacteria and archaea with

PADLOC reveals new system types. Nucleic Acids Res *49*, 10868-10878.

10.1093/nar/gkab883.

14.   Payne, L.J., Meaden, S., Mestre, M.R., Palmer, C., Toro, N., Fineran,

Peter C., and Jackson, Simon A. (2022). PADLOC: a web server for the

identification of antiviral defence systems in microbial genomes. Nucleic Acids

Res *50*, W541-W550. 10.1093/nar/gkac400.

15.   Ofir, G., Herbst, E., Baroz, M., Cohen, D., Millman, A., Doron, S., Tal, N.,

Malheiro, D.B.A., Malitsky, S., Amitai, G., and Sorek, R. (2021). Antiviral

activity of bacterial TIR domains via immune signaling molecules. Nature

600,116-120. 10.1038/s41586-021-04098-7

16.   Duncan-Lowey, B., McNamara-Bordewick, N.K., Tal, N., Sorek, R., and

Kranzusch, P.J. (2021). Effector-mediated membrane disruption controls cell

death in CBASS antiphage defense. Mol Cell *81*, 5039-5051.e5035.

10.1016/j.molcel.2021.10.020.

17.   Galperin, M.Y., Vera Alvarez, R., Karamycheva, S., Makarova, K.S., Wolf,

Yuri I., Landsman, D., and Koonin, E.V. (2024). COG database update 2024.

Nucleic Acids Res. 10.1093/nar/gkae983.

18.   Parks, D.H., Chuvochina, M., Rinke, C., Mussig, A.J., Chaumeil, P.-A., and

Hugenholtz, P. (2021). GTDB: an ongoing census of bacterial and archaeal

diversity through a phylogenetically consistent, rank normalized and complete

genome-based   taxonomy.   Nucleic   Acids   Res   *50*,   D785-D794.

10.1093/nar/gkab776.

19.   Paulino, C., Kalienkova, V., Lam, A.K.M., Neldner, Y., and Dutzler, R. (2017).

Activation mechanism of the calcium-activated chloride channel TMEM16A

revealed by cryo-EM. Nature *552*, 421-425. 10.1038/nature24652.

20.   Rossmann, M.G., Moras, D., and Olsen, K.W. (1974). Chemical and biological

evolution of a nucleotide-binding protein. Nature *250*, 194-199.

10.1038/250194a0.

21.   Medvedev, K.E., Kinch, L.N., Schaeffer, R.D., and Grishin, N.V. (2019).

Functional analysis of Rossmann-like domains reveals convergent evolution

of topology and reaction pathways. PLoS Comput Biol *15*, e1007569.

10.1371/journal.pcbi.1007569.

22.   Mount, J., Maksaev, G., Summers, B.T., Fitzpatrick, J.A.J., and Yuan, P.

(2022). Structural basis for mechanotransduction in a potassium-dependent

mechanosensitive ion channel. Nat Commun *13*, 6904. 10.1038/s41467-022-

34737-0.

23.   Lopatina, A., Tal, N., and Sorek, R. (2020). Abortive Infection: Bacterial

Suicide as an Antiviral Immune Strategy. Ann Rev Virol *7*, 371-384.

10.1146/annurev-virology-011620-040628.

24.   Aframian, N., and Eldar, A. (2023). Abortive infection antiphage defense

systems: separating mechanism and phenotype. Trends Microbiol *31*, 1003-

1012. 10.1016/j.tim.2023.05.002.

25.   Tabib-Salazar, A., Liu, B., Barker, D., Burchell, L., Qimron, U., Matthews, S.J.,

and Wigneshweraraj, S. (2018). T7 phage factor required for managing RpoS

in Escherichia coli. Proc Natl Acad Sci *115*, E5353-E5362.

10.1073/pnas.1800429115.

26. Tabib-Salazar, A., Liu, B., Shadrin, A., Burchell, L., Wang, Z., Wang, Z.,
Goren, M.G., Yosef, I., Qimron, U., Severinov, K., et al. (2017). Full shut-off of
Escherichia coli RNA-polymerase by T7 phage requires a small phage-
encoded DNA-binding protein. Nucleic Acids Res *45*, 7697-7707.
10.1093/nar/gkx370.

27. Hu, B., Margolin, W., Molineux, I.J., and Liu, J. (2015). Structural remodeling
of bacteriophage T4 and host membranes during infection initiation. Proc Natl
Acad Sci *112*, E4919-E4928. doi:10.1073/pnas.1501064112.

28. Medvedev, K.E., Kinch, L.N., Dustin Schaeffer, R., Pei, J., and Grishin, N.V.
(2021). A Fifth of the Protein World: Rossmann-like Proteins as an
Evolutionarily Successful Structural unit. J Mol Biol *433*, 166788.
10.1016/j.jmb.2020.166788.

29. Yuan, Q., Xie, J., Xie, J., Zhao, H., and Yang, Y. (2023). Fast and accurate
protein function prediction from sequence through pretrained language model
and homology-based label diffusion. Brief Bioinform *24*. 10.1093/bib/bbad117.

30. Kutter, E., Bryan, D., Ray, G., Brewster, E., Blasdel, B., and Guttman, B.
(2018). From Host to Phage Metabolism: Hot Tales of Phage T4's Takeover
of E. coli. Viruses *10*, 387.

31. Michel, B., Boubakri, H., Baharoglu, Z., LeMasson, M., and Lestini, R. (2007).
Recombination proteins and rescue of arrested replication forks. DNA Repair
6, 967-980. 10.1016/j.dnarep.2007.02.016.

32. Mayo-Muñoz, D., Pinilla-Redondo, R., Camara-Wilpert, S., Birkholz, N., and
Fineran, P.C. (2024). Inhibitors of bacterial immune systems: discovery,
mechanisms and applications. Nat Rev Gen. 10.1038/s41576-023-00676-9.

33.   Yirmiya, E., Leavitt, A., Lu, A., Ragucci, A.E., Avraham, C., Osterman, I.,

Garb, J., Antine, S.P., Mooney, S.E., Hobbs, S.J., et al. (2024). Phages

overcome bacterial immunity via diverse anti-defence proteins. Nature *625*,

352-359. 10.1038/s41586-023-06869-w.

34.   Shin, J., Jiang, F., Liu, J.-J., Bray, N.L., Rauch, B.J., Baik, S.H., Nogales, E.,

Bondy-Denomy, J., Corn, J.E., and Doudna, J.A. (2017). Disabling Cas9 by

an anti-CRISPR DNA mimic. Sci Adv *3*, e1701620. 10.1126/sciadv.1701620.

35.   Isaev, A., Drobiazko, A., Sierro, N., Gordeeva, J., Yosef, I., Qimron, U.,

Ivanov, N.V., and Severinov, K. (2020). Phage T7 DNA mimic protein Ocr is a

potent inhibitor of BREX defence. Nucleic Acids Res *48*, 5397-5406.

10.1093/nar/gkaa290.

36.   Krüger, D.H., and Schroeder, C. (1981). Bacteriophage T3 and bacteriophage

T7 virus-host cell interactions. Microbiol Rev *45*, 9-51. 10.1128/mr.45.1.9-

51.1981.

37.   Sakaki, Y., Karu, A.E., Linn, S., and Echols, H. (1973). Purification and

Properties of the γ-Protein Specified by Bacteriophage λ: An Inhibitor of the

Host RecBC Recombination Enzyme. Proc Natl Acad Sci *70*, 2215-2219.

10.1073/pnas.70.8.2215.

38.   Millman, A., Bernheim, A., Stokar-Avihail, A., Fedorenko, T., Voichek, M.,

Leavitt, A., Oppenheimer-Shaanan, Y., and Sorek, R. (2020). Bacterial

Retrons Function In Anti-Phage Defense. Cell *183*, 1551-1561.e1512.

10.1016/j.cell.2020.09.065.

39.   Marsić, N., Roje, S., Stojiljković, I., Salaj-Smic, E., and Trgovcević, Z. (1993).

In vivo studies on the interaction of RecBCD enzyme and lambda Gam

protein. J Bacteriol *175*, 4738-4743. 10.1128/jb.175.15.4738-4743.1993.

40.   Dillingham, M.S., and Kowalczykowski, S.C. (2008). RecBCD Enzyme and
1301   the Repair of Double-Stranded DNA Breaks. Microbiol Mol Biol Rev *72*, 642-
1302   671. 10.1128/MMBR.00020-08.

1303   41.   Cheng, K., Wilkinson, M., Chaban, Y., and Wigley, D.B. (2020). A
1304   conformational switch in response to Chi converts RecBCD from phage
1305   destruction to DNA repair. Nat Struct Mol Biol *27*, 71-77. 10.1038/s41594-
1306   019-0355-2.

1307   42.   Levy, A., Goren, M.G., Yosef, I., Auster, O., Manor, M., Amitai, G., Edgar, R.,
1308   Qimron, U., and Sorek, R. (2015). CRISPR adaptation biases explain
1309   preference for acquisition of foreign DNA. Nature *520*, 505-510.
1310   10.1038/nature14302.

1311   43.   Kuzmenko, A., Oguienko, A., Esyunina, D., Yudin, D., Petrova, M., Kudinova,
1312   A., Maslova, O., Ninova, M., Ryazansky, S., Leach, D., et al. (2020). pAgo-
1313   induced DNA interference protects bacteria from invader DNA. bioRxiv,
1314   2020.2003.2001.971358. 10.1101/2020.03.01.971358.

1315   44.   Lipinska, B., Rao, A., Bolten, B., Balakrishnan, R., and Goldberg, E. (1989).
1316   Cloning and identification of bacteriophage T4 gene 2 product gp2 and action
1317   of gp2 on infecting DNA in vivo. J Bacteriol *171*.

1318   45.   Xie, P., Zhang, H., Qin, Y., Xiong, H., Shi, C., and Zhou, Z. (2023). Membrane
1319   Proteins and Membrane Curvature: Mutual Interactions and a Perspective on
1320   Disease Treatments. Biomolecules *13*, 1772.

1321   46.   Wilkinson, M., Troman, L., Wan Nur Ismah, W.A.K., Chaban, Y., Avison, M.B.,
1322   Dillingham, M.S., and Wigley, D.B. (2016). Structural basis for the inhibition of
1323   RecBCD by Gam and its synergistic antibacterial effect with quinolones. Elife
1324   *5*, e22963. 10.7554/eLife.22963.

47. Kudryavtseva, A.A., Cséfalvay, E., Gnuchikh, E.Y., Yanovskaya, D.D., Skutel,

M.A., Isaev, A.B., Bazhenov, S.V., Utkina, A.A., and Manukhov, I.V. (2023).

Broadness and specificity: ArdB, ArdA, and Ocr against various restriction-

modification systems. Front Microbiol *14*. 10.3389/fmicb.2023.1133144.

48. Osterman, I., Samra, H., Rousset, F., Loseva, E., Itkin, M., Malitsky, S.,

Yirmiya, E., Millman, A., and Sorek, R. (2024). Phages reconstitute NAD+ to

counter bacterial immunity. Nature *634*, 1160-1167. 10.1038/s41586-024-

07986-w.

49. Jiang, S., Chen, C., Huang, W., He, Y., Du, X., Wang, Y., Ou, H., Deng, Z.,

Xu, C., Jiang, L., et al. (2024). A widespread phage-encoded kinase enables

evasion of multiple host antiphage defence systems. Nat Microbiol.

10.1038/s41564-024-01851-2.

50. Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko,

K.A., Tomita, M., Wanner, B.L., and Mori, H. (2006). Construction of

Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio

collection. Mol Sys Biol *2*, 2006.0008. 10.1038/msb4100050.

51. Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences.

Bioinformatics *34*, 3094-3100. 10.1093/bioinformatics/bty191.

52. Luong, T., Salabarria, A.-C., Edwards, R.A., and Roach, D.R. (2020).

Standardized bacteriophage purification for personalized phage therapy. Nat

Prot *15*, 2867-2890. 10.1038/s41596-020-0346-0.

53. Deatherage, D.E., and Barrick, J.E. (2014). Identification of Mutations in

Laboratory-Evolved Microbes from Next-Generation Sequencing Data Using

breseq. In Engineering and Analyzing Multicellular Systems: Methods and

Protocols, L. Sun, and W. Shou, eds. (Springer New York), pp. 165-188.

10.1007/978-1-4939-0554-6_12.

54. Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017).

cryoSPARC: algorithms for rapid unsupervised cryo-EM structure

determination. Nat Methods *14*, 290-296. 10.1038/nmeth.4169.

55. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,

Tunyasuvunakool, K., Bates, R., Zidek, A., Potapenko, A., et al. (2021). Highly

accurate protein structure prediction with AlphaFold. Nature *596*, 583-589.

10.1038/s41586-021-03819-2.

56. Mount, J., Maksaev, G., Summers, B.T., Fitzpatrick, J.A.J., and Yuan, P.

(2022). Structural basis for mechanotransduction in a potassium-dependent

mechanosensitive ion channel. Nat Commun *13*, 6904. 10.1038/s41467-022-

34737-0.

57. Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and

development of Coot. Acta Crystallogr D Biol Crystallogr *66*, 486-501.

10.1107/S0907444910007493.

58. McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni,

L.C., and Read, R.J. (2007). Phaser crystallographic software. J Appl

Crystallogr *40*, 658-674. 10.1107/S0021889807021206.

59. Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N.,

Headd, J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W., et al. (2010).

PHENIX: a comprehensive Python-based system for macromolecular

structure solution. Acta Crystallogr D Biol Crystallogr *66*, 213-221.

10.1107/S0907444909052925.

60.   Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M.,

Meng, E.C., and Ferrin, T.E. (2004). UCSF Chimera--a visualization system

for exploratory research and analysis. J Comput Chem *25*, 1605-1612.

10.1002/jcc.20084.

61.   Pettersen, E.F., Goddard, T.D., Huang, C.C., Meng, E.C., Couch, G.S., Croll,

T.I., Morris, J.H., and Ferrin, T.E. (2021). UCSF ChimeraX: Structure

visualization for researchers, educators, and developers. Protein Sci *30*, 70-

82. 10.1002/pro.3943.

62.   Steinegger, M., and Söding, J. (2017). MMseqs2 enables sensitive protein

sequence searching for the analysis of massive data sets. Nat Biotechnol *35*,

1026-1028. 10.1038/nbt.3988.

63.   Edgar, R.C. (2022). Muscle5: High-accuracy alignment ensembles enable

unbiased assessments of sequence homology and phylogeny. Nat Commun

13, 6968. 10.1038/s41467-022-34630-w.

64.   Steenwyk, J.L., Buida, T.J., III, Li, Y., Shen, X.-X., and Rokas, A. (2020).

ClipKIT: A multiple sequence alignment trimming software for accurate

phylogenomic inference. PLOS Biol *18*, e3001007.

10.1371/journal.pbio.3001007.

65.   Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D.,

von Haeseler, A., and Lanfear, R. (2020). IQ-TREE 2: New Models and

Efficient Methods for Phylogenetic Inference in the Genomic Era. Mol Biol

Evol *37*, 1530-1534. 10.1093/molbev/msaa015.

66.   Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., and

Jermiin, L.S. (2017). ModelFinder: fast model selection for accurate

phylogenetic estimates. Nat Meth *14*, 587-589. 10.1038/nmeth.4285.

67.   Hoang, D.T., Chernomor, O., von Haeseler, A., Minh, B.Q., and Vinh, L.S.

(2017). UFBoot2: Improving the Ultrafast Bootstrap Approximation. Mol Biol

Evol *35*, 518-522. 10.1093/molbev/msx281.

68.   Letunic, I., and Bork, P. (2019). Interactive Tree Of Life (iTOL) v4: recent

updates and new developments. Nucleic Acids Res *47*, W256-W259.

10.1093/nar/gkz239.

69.   Camargo, A.P., Roux, S., Schulz, F., Babinski, M., Xu, Y., Hu, B., Chain,

P.S.G., Nayfach, S., and Kyrpides, N.C. (2024). Identification of mobile

genetic elements with geNomad. Nat Biotechnol *42*, 1303-1312.

10.1038/s41587-023-01953-y.

70.   Banerjee, P., Eulenstein, O., and Friedberg, I. (2024). Discovering genomic

islands in unannotated bacterial genomes using sequence embedding. Bioinf

Adv *4*. 10.1093/bioadv/vbae089.

71.   Söding, J., Biegert, A., and Lupas, A.N. (2005). The HHpred interactive server

for protein homology detection and structure prediction. Nucleic Acids Res *33*,

W244-W248. 10.1093/nar/gki408.

72.   Hallgren, J., Tsirigos, K.D., Pedersen, M.D., Almagro Armenteros, J.J.,

Marcatili, P., Nielsen, H., Krogh, A., and Winther, O. (2022). DeepTMHMM

predicts alpha and beta transmembrane proteins using deep neural networks.

bioRxiv, 2022.2004.2008.487609. 10.1101/2022.04.08.487609.

73.   Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R.,

McWilliam, H., Remmert, M., Söding, J., et al. (2011). Fast, scalable

generation of high-quality protein multiple sequence alignments using Clustal

Omega. Mol Sys Biol *7*, 539. 10.1038/msb.2011.75.

74.   Omasits, U., Ahrens, C.H., Müller, S., and Wollscheid, B. (2013). Protter:

1423   interactive protein feature visualization and integration with experimental

1424   proteomic data. Bioinformatics *30*, 884-886. 10.1093/bioinformatics/btt607.

1425   75.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,

1426   Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., et al. (2021).

1427   Highly accurate protein structure prediction with AlphaFold. Nature *596*, 583-

1428   589. 10.1038/s41586-021-03819-2.

1429   76.   Cianfrocco, M.A., Wong-Barnum, M., Youn, C., Wagner, R., and Leschziner,

1430   A. (2017). COSMIC2: A Science Gateway for Cryo-Electron Microscopy

1431   Structure Determination. Proceedings of the Practice and Experience in

1432   Advanced Research Computing 2017 on Sustainability, Success and Impact.

1433   Association for Computing Machinery.

1434   77.   Holm, L., and Laakso, L.M. (2016). Dali server update. Nucleic Acids Res *44*,

1435   W351-W355. 10.1093/nar/gkw357.

1436   78.   Mukherjee, S., and Zhang, Y. (2009). MM-align: a quick algorithm for aligning

1437   multiple-chain protein complex structures using iterative dynamic

1438   programming. Nucleic Acids Res *37*, e83-e83. 10.1093/nar/gkp318.

1439   79.   Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J.,

1440   Morris, J.H., and Ferrin, T.E. (2023). UCSF ChimeraX: Tools for structure

1441   building and analysis. Prot Sci *32*, e4792. 10.1002/pro.4792.

1442   80.   Li, W., O'Neill, K.R., Haft, D.H., DiCuccio, M., Chetvernin, V., Badretdin, A.,

1443   Coulouris, G., Chitsaz, F., Derbyshire, Myra K., Durkin, A.S., et al. (2020).

1444   RefSeq: expanding the Prokaryotic Genome Annotation Pipeline reach with

1445   protein family model curation. Nucleic Acids Res *49*, D1020-D1028.

1446   10.1093/nar/gkaa1105.

81. Bushnell, B., Rood, J., and Singer, E. (2017). BBMerge – Accurate paired

shotgun read merging via overlap. PLoS One *12*, e0185056.

10.1371/journal.pone.0185056.

82. Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-

based genome alignment and genotyping with HISAT2 and HISAT-genotype.

Nat Biotechnol *37*, 907-915. 10.1038/s41587-019-0201-4.

83. Vera Alvarez, R., Pongor, L.S., Mariño-Ramírez, L., and Landsman, D.

(2018). TPMCalculator: one-step software to quantify mRNA abundance of

genomic features. Bioinformatics *35*, 1960-1962.

10.1093/bioinformatics/bty896.

1457

A

ECOR8
ECOR12
2
O55:H7
D9
ECOR49
1
3
SN82F48
1.0

ECOR8
ECOR12
2
D9
ECOR49
O55:H7
1
3
SN82F48
1.0

Prophages
Plasmids
Genomic islands
Other mobilome

Previously confirmed anti-phage activity

KwaA n = 1,259

KwaB n = 1,262

B

KwaA | KwaB

Campylobacterales
Micrococcales
Mycobacteriales
Bacillales
Lactobacillales
Lachnospirales
Eubacteriales
Bacteroidales
Flavobacteriales
Hyphomicrobiales
Sphingomonadales
Burkholderiales
Xanthomonadales
Moraxellales
Oceanospirillales
Pseudomonadales
Enterobacterales
Vibrionales
Alteromonadales
Aeromonadales

1.0

Count | Count

C

KwaA   KwaB

ECOR49

TM   195aa   DUF4868   319aa

D

	Myo			Sipho			Podo	
	T2	T4	T6	T1	T5	λ_{vir}	T3	T7$_s$
ECOR8							m	m
ECOR12								
ECOR49							m	m
D9								
SN82F48							m	m

Fold defence

<10^1
10^1
10^2
10^3
10^4
10^5
>10^6

m = plaque morphology change

E

% pSEVA637 conjugants

Control   ECOR8   ECOR12   ECOR49   D9   SN82F48

A

KwaA⁴ KwaA³

KwaA¹ KwaA²

LMNG

C2 symmetry

B

KwaA⁴ KwaA³

KwaA¹ KwaA²

C4 symmetry

C

Extra

Intra

ECL1   ECL2

1   2   3   4

NH₂   ICL   COOH

beta-barrel

D

Interface I

Interface II

C2 symmetry

E

Interface I

C4 symmetry

F

C2 symmetry
C4 symmetry

G

KwaB dimer2

KwaB dimer1

CTD   α2   NTD

H

CTD

KwaB¹   KwaB²

NTD

KwaB dimer
(Crystal)

A

B
KwaB-72 nt ssDNA   KwaB-72 bp dsDNA
0 1 4 8 16 32 64   0 1 8 16 32 64 128

C

	circular plasmid	*E. coli* gDNA +plasmid	Phage gDNA
KwaB	− +	− +	− +

kb
10
2.5
1.5
1.0

D
CTD
KwaB¹   KwaB²
NTD
16-mer dsDNA

AF3 model
R233
N134   K187
D169   R167
R142

E
KwaB-72 nt ssDNA   KwaB-72 bp dsDNA
− wt m3 m6   − wt m3 m6

m3: R142A/R167A/K187A

m6: R142A/R167A/K187A/ R233A/N134A/D169A

F

Uninfected   Infected
FM4-64
KwaB

30 min

60 min

1 μm

30 min

| | KwaB foci/cell |
Uninf. Infect.

60 min

| | KwaB foci/cell |
Uninf. Infect.

H

Fraction bound vs KwaB (nM)
dsDNA
ssDNA

G

log fold change vs Genome position (kb)

glutaredoxin GroES dCTP pyrophosphatase sliding clamp
RegA clamp loader RpbA
RNaseH RecA
UvsY td glutaredoxin
DNA ligase Gp59 exonuclease endonuclease
MTase td GT

terminase tail sheath tail fiber anti-sigma factor A
neck protein tail fiber
fibritin capsid protein MotA
neck protein baseplate subunit long tail fiber holin
baseplate subunit tail fiber adhesin