Kiwa rescues RecBCD for anti-phage activity

Thomas C. Todeschini[1], Yi Wu[1], Ameena Naji[1,2], Rupavidhya Mondi[1,3], Franklin L. Nobrega[1]*

[1]School of Biological Sciences, University of Southampton, SO17 1BJ Southampton, UK

*Corresponding author: F.Nobrega@soton.ac.uk

[2] Current address: School of Molecular and Cellular Biology, University of Leeds, Leeds LS2 9JT, UK

[3] Current address: The William Harvey Research Institute, Barts and The London School of Medicine, Queen Mary University of London, Charterhouse Square, London EC1M 6BQ, UK

Abstract

Bacteria have evolved numerous defence systems to combat phage infection, but our understanding of the underlying mechanisms is limited beyond their antiviral function. Here, we provide mechanistic insights into the Kiwa defence system, which is composed of two proteins, the transmembrane protein KwaA and the DUF4868 domain-containing protein KwaB. Using Kiwa from *Escherichia coli* ECOR8 and ECOR12, we found that KwaA detects phage infection by sensing inhibition of the host RNA polymerase. Consequently, the inhibitory effect of KwaA on the effector protein KwaB is relieved, which results in decreased phage DNA replication efficiency through a RecBCD-dependent pathway. Notably, Kiwa restores RecBCD activity even against phages that encode anti-RecBCD proteins. Our findings also reveal that KwaB has antiphage activity in the absence of both KwaA and phage trigger, but the conserved two-protein Kiwa operon implies that strict regulation of KwaB activity by KwaA is necessary to prevent potential auto-immunity.

Keywords: Kiwa, RNAP inhibitor, RecBCD, ATP depletion, phage DNA replication

Bacteria have evolved a set of mechanisms to defend against phage predation.

Some of these defence mechanisms, such as CRISPR-Cas and restriction

modification systems, have been widely studied and used in biotechnology, including

Cas9-based gene editing[1]. However, most of the defence systems were discovered

recently[2-10] and their molecular mechanisms are not well understood beyond their

anti-phage functions. Research on some of these novel defence systems, such as

BREX[5,11,12], DISARM[4,13,14], CBASS[15,16], Thoeris[17,18], Pycsar[19], PARIS[8], Gabija[3,20,21],

Retrons[22], RADAR[6,23,24], DSR[25], Avast[6,26], Gasdermins[27], dCTP deaminase,

dGTPase[28,29], and pVips[30], has made significant progress in elucidating their

molecular activities. Further research is needed to fully characterise the dozens of

defence systems that remain to be explored.

The Kiwa defence system is among the recent discoveries in anti-phage defence

and is composed of two genes, *kwaA* and *kwaB*. *kwaA* encodes a protein with

transmembrane-spanning abilities, while *kwaB* encodes a protein containing the

DUF4868 domain of unknown function. The Kiwa system from *Escherichia coli*

O55:H7 RM12579 was shown to protect against phages Lambda and SECphi18[3],

but there is currently no further understanding of the precise mechanisms underlying

the protective effects of Kiwa.

In this study, we demonstrate that Kiwa defence systems derived from *E. coli*

ECOR8 and ECOR12 are activated by the inhibition of the host RNA polymerase

(RNAP) by phage-encoded proteins. Upon sensing of RNAP inhibition, KwaA

relieves its inhibitory activity on KwaB. KwaB then initiates a RecBCD-mediated

process that reduces the efficiency of phage DNA replication, even in the presence

of phage-encoded anti-RecBCD proteins. Notably, KwaB exhibits anti-phage activity

without the presence of KwaA and the phage trigger. However, the fact that KwaA

and KwaB are present together in the conserved Kiwa operon across various

bacterial species underscores the importance of tightly regulating KwaB activity

through KwaA to avoid potential auto-immunity.

Results

Kiwa orthologues protect against different phages

To explore the diversity of Kiwa within prokaryotic genomes, we collected the Kiwa

operons from all prokaryotic genomes accessible in the NCBI database. Out of the

198,640 genomes analysed, 1.6% were found to contain Kiwa operons, with the

largest proportion (45%) being present in *Escherichia coli* (Supplementary Fig. 1a).

This outcome aligns with the findings of earlier research[3].

By means of hierarchical cluster analysis of the KwaA and KwaB proteins within

Kiwa operons, we observed their distribution into four clusters (Fig. 1a). For KwaA,

we noted substantial diversity in protein size (ranging from 66 to 389 amino acids,

Supplementary Table 1) and the number of transmembrane domains (varying from

zero to five, with four being the most prevalent, Supplementary Fig. 1b). Previous

research has demonstrated that diversity within defence systems of the same

subtype may result in differing specificities of phage protection[2,15,17]. To determine

whether diversity in Kiwa plays a role in defence specificity, we cloned five Kiwa

systems from different clusters into *E. coli* BL21-AI (Fig. 1b, Supplementary Fig. 1c),

which lacks these systems naturally. Infection assays with a panel of coliphages

showed that four of the five Kiwa systems tested exhibited anti-phage activity with

phage specificity that was relatively consistent with the hierarchical clusters (Fig.

1a,c).

Moreover, the analysis of infection dynamics in liquid cultures confirmed that cells

expressing Kiwa could protect against phage infection at both low (0.03) and high (3)

multiplicities of infection (MOI). However, at higher MOI, we observed a delayed

collapse of the culture, suggesting possible cell death (Fig. 1d, Supplementary Fig.

1d). Time post-infection assays demonstrated a decline in phage concentration

within one hour of infection, revealing that the phage was capable of entering the cell

but not replicating (Fig. 1e, Supplementary Fig. 1e).

To sum up, our findings reveal that Kiwa systems from different clusters exhibit

distinct phage specificity patterns in terms of their protective efficacy.

KwaB is the effector of Kiwa and is regulated by an interaction with KwaA

Next, we investigated the role of individual genes for Kiwa's protective function by

exposing single-gene deletions of Kiwa ECOR8 to phages. Our results indicate that

the deletion of KwaB resulted in the loss of protection, whereas the deletion of KwaA

had no effect (Fig. 2a, Supplementary Fig. 2a). Interestingly, the presence of KwaA

in ECOR8 seems to induce cell toxicity, as evidenced by reduced cell growth in the

absence of KwaB, but not in the absence of KwaA (Supplementary Fig. 2b). This

toxic effect is also observed in Kiwa from ECOR49 and D9, but not in Kiwa from

ECOR12 or SN82F48 (Supplementary Fig. 2b). KwaA contains transmembrane (TM)

domains, which are also present in proteins of other characterised defence systems

such as CBASS[15], Pycsar[19], and Retrons[22]. In CBASS, the deletion of the TM

domains of the effector proteins abolished cell toxicity and revealed these to be

essential for functionality[15]. To determine the role of the TM domains in KwaA-

induced cell toxicity, we truncated TM 1 (Δ5-25) and TM 1-2 (Δ5-74), resulting in

reduced cell toxicity upon deletion of TM 1-2 (Supplementary Fig. 2b). Curiously,

both truncations caused a loss of phage protection (Fig. 2a, Supplementary Fig. 2a).

To further investigate the importance of the TM domains in Kiwa protection, we

examined Kiwa SN82F48, which contains only two TM domains and does not

provide phage protection (Fig. 1b). A closer inspection of the coding region of Kiwa

SN82F48 revealed that the start of *kwaA* was located 336 nucleotides upstream of

the annotated start, resulting in a longer open reading frame (ORF) that included four

TM domains. The reconstitution of the longer *kwaA* restored the activity of Kiwa

SN82F48 against two coliphages, providing evidence for the critical role of the four

TM regions in KwaA for anti-phage defence (Fig. 2b, Supplementary Fig. 2c,d).

Curiously, as with Kiwa ECOR12, no toxic effect on cell growth was observed

(Supplementary Fig. 2b).

Based on the observation that protection is abolished only when a TM-truncated

version of KwaA is present, and not when KwaA is completely absent, we

hypothesized that KwaA and KwaB interact with each other, and that KwaB is in a

free form in the cytoplasm when KwaA is absent, but bound to an inactive form of

KwaA when this protein is present in a truncated version. To test our hypothesis, we

conducted pull down assays with these proteins in *E. coli*. Since KwaA is a

transmembrane protein and expression of such proteins is notoriously challenging,

we opted to express only the cytoplasmic C-terminal region of KwaA (Supplementary

Fig. 2e) that we suspect is responsible for the interaction with KwaB, and which is

also essential for phage protection by Kiwa (Fig. 2a). By using the tagged C-terminal

of KwaA we were able to pull down untagged KwaB, as confirmed by size exclusion

chromatography and mass spectrometry (Fig. 2c, Supplementary Fig. 3a), indicating

that these two proteins interact.

Moreover, KwaB is characterised by the domain of unknown function DUF4868,

which is present in a wide range of bacteria. A member of the family of DUF4868-

containing proteins is involved in the synthesis of a bacteriocin toxin as part of a

gene cluster, but the function of the domain itself remains unclear[31]. We show that

point mutations on specific conserved amino acids within this domain (L145A/F147A

and S272A, Supplementary Fig. 3b) result in decreased phage protection (Fig. 2a,

Supplementary Fig. 2a). This supports the importance of the DUF4868 domain for

phage defence by Kiwa. Alphafold[32] predictions suggest that KwaB can form dimers,

and an interesting finding is that amino acid S272 may facilitate this dimerization by

establishing several hydrogen bounds with the facing S272 amino acid in the second

protein. Furthermore, amino acids L145 and F147 are contained within a beta strand

that is situated in a pocket, suggesting a potential for these residues to interact with

a target molecule (Supplementary Fig. 3c).

In summary, our results suggest that KwaB serves as the effector molecule for the

Kiwa defence system, exhibiting anti-phage activity that is not reliant on KwaA.

However, the activity of KwaB is modulated by its interaction with KwaA within the

Kiwa operon.

KwaA senses phage host hijacking proteins to activate defence

We posited that the toxicity associated with some of the Kiwa systems tested was

not a natural effect of the KwaA protein, but rather a consequence of overexpression

of the membrane protein, which can be detrimental to the cell. To investigate this, we

replaced the inducible T7 promoter in Kiwa ECOR8 with a constitutive Anderson

promoter that may aid in proper protein folding[33]. The results showed that Kiwa

ECOR8 expressed under a constitutive promoter did not exhibit any toxicity (Fig. 3a).

Sequencing of the plasmid and bacterial strain confirmed the absence of any genetic

alterations (Supplementary Table 1), providing additional evidence that the toxicity

observed previously in the absence of phage was likely caused by either

overproduction or misfolding of KwaA.

Under the constitutive promoter, phage infection of Kiwa ECOR8 resulted in a

delayed collapse of the culture, especially at high phage MOI, which may suggest an

abortive infection phenotype[34] (Fig. 3a, Supplementary Fig. 4a). To further

investigate this, we set out to identify the trigger of the defence system by evolving

escape phages[35]. We observed that the Kiwa ECOR8 system expressed

constitutively displayed a broader and stronger anti-phage activity compared to the

same system expressed under the T7 inducible promoter (Fig. 3b, Supplementary

Fig. 4b). Notably, the constitutively-expressed Kiwa exhibited particularly high activity

against phage T3, with 10^4-fold protection (Fig. 3b) and decrease in plaque size (Fig.

3c, Supplementary Fig. 4c). Thus, we used phage T3 in our attempt to generate

escape phages. Given the genomic similarities between T7Select (T7s) and T3, and

the distinct patterns of protection provided against these phages by Kiwa ECOR8

(Fig. 3b), we included T7s in our efforts to generate escape mutants. We tested five

T3 and three T7s mutants and observed that Kiwa from ECOR8 and ECOR12 lost

protection against these mutant phages, suggesting that these systems from the

same cluster (light blue, Fig 1a) use identical phage sensing mechanisms (Fig. 3d,

Supplementary Fig. 5a). However, the phage escape mutants were unable to evade

defence by Kiwa D9, from a different cluster (dark green, Fig. 1a), indicating that

Kiwa systems from separate clusters employ different phage sensing mechanisms,

which likely account for the unique protective patterns we previously observed (Fig

1c).

The escape mutants were sequenced, revealing mutations (early stop codon or gene

deletion) in the inhibitors of the stationary phase RpoS/σ^S sigma subunit of the host

RNA polymerase (RNAP) (T7s gp5.7 and T3 T3p30)[36,37] (Fig. 3e, Supplementary

Table 1). To confirm that the RNAP inhibitors trigger defence by Kiwa ECOR8, we

expressed them in *trans* and observed the restoration of protective activity against

the escape phages (Fig. 3f,g, Supplementary Fig. 5b,c). Despite the presence of

mutations in the tail fibre gene of T7s (gp17) and T3 (T3p48), we observed that

expression of the tail fibre in Kiwa cells did not have any impact on phage protection

(Supplementary Fig. 5d).

The expression of the RNAP inhibitors caused a decrease in the cell growth of both

Kiwa and control cells (Fig. 3g, Supplementary Fig. 5e), likely due to their effect on

the bacterial RNAP. This limited our ability to test abortive infection using the trigger

protein from the phage. In addition, we investigated whether Kiwa detected the

RNAP inhibitor or the integrity of RNAP. Inspired by earlier work[29], we conducted

experiments using rifampicin, a known inhibitor of bacterial RNAP, to assess its

impact on the ability of the mutant phages to infect control and Kiwa cells. The

findings demonstrated that in the presence of rifampicin, mutant phage propagation

decreased in Kiwa cells compared to control cells, indicating that inhibition of the

bacterial RNAP triggers Kiwa defence (Supplementary Fig. 5f). This suggests that

the Kiwa defence system monitors the integrity of RNAP instead of directly sensing

gp5.7/T3p30.

To determine which protein in Kiwa is responsible for sensing phage infection, we

assessed the protective activity of Kiwa ECOR8 and its single gene mutants against

the escape phages (Fig. 3h, Supplementary Fig. 5g). As expected, the escape

phage was not inhibited by the complete Kiwa ECOR8 or ΔKwaB, but unexpectedly,

ΔKwaA provided approximately 100-fold protection. These results indicate that the

anti-phage activity of KwaB is independent of a phage trigger and suggest that KwaA

is the protein responsible for sensing phage infection and modulating the effector

KwaB.

To summarize, our findings indicate that KwaA senses RNAP inhibition by phage

proteins. This recognition is likely responsible for releasing the effector protein

KwaB, which is the key player in the Kiwa defence system.

RecBCD is required for Kiwa defence

To gain insight into the effector mechanism of Kiwa, we investigated different

possible outcomes of the defence. We first tested whether Kiwa affects phage

adsorption (Fig. 4a). Our results show that Kiwa from ECOR8 and ECOR12 does not

affect the adsorption of T3 and T4 phages, suggesting that protection is provided at

later stages of infection. As we observed a reduction in both extracellular and

intracellular levels of viable phage (Fig. 4b), we next sought to determine if the

decrease in intracellular phage particles was caused by reduced phage DNA

replication. We extracted and sequenced DNA from control and Kiwa cells infected

with phage at high MOI, both at the start and 30 minutes after phage infection,

revealing a ≈25% decrease in phage DNA replication efficiency in Kiwa cells

compared to control cells (Fig. 4c). This suggests that the defence system either

halts infection prior to the expression of phage proteins responsible for DNA
replication, or acts directly on the phage DNA.

To test the former, we explored the possible abortive infection mechanism of Kiwa.
As KwaA is a transmembrane protein, which can sometimes be associated with ATP
secretion leading to cellular ATP depletion[38-41], we measured the ATP levels in
control and Kiwa cultures infected with phage. Our results indicate that total ATP is
depleted in phage-infected Kiwa cells (Fig. 4d), suggesting that both extracellular
and intracellular ATP levels are reduced. This implies that ATP secretion is not
involved and another intracellular process must be responsible for ATP consumption.
Importantly, the depletion of ATP is not observed when either of the Kiwa proteins is
absent (Fig. 4e). Given that KwaB alone is sufficient to provide anti-phage protection
(Fig. 2a), our results indicate that ATP depletion is not the effector mechanism of
Kiwa.

The phage RNAP inhibitors (gp5.7/T3p30) that trigger Kiwa defence are middle
genes and are expressed during the same stage as the phage proteins involved in
DNA replication[36,37,42]. This suggests that phage infection proceeds until this stage in
Kiwa cells, leading us to hypothesise that the observed reduction in phage DNA
replication is a direct consequence of a targeted action on phage DNA replication. As
none of the Kiwa proteins are predicted to have nuclease activity, we investigated
the possible involvement of host RecBCD in the Kiwa defence mechanism. RecBCD
is known to be a crucial element of host immunity, with rapid nuclease activity on
phage DNA[43,44]. This complex has been implicated in the activity of other defence
systems, such as CRISPR-Cas[45] and prokaryotic Argonaute[46]. Thus, we
hypothesized that Kiwa may mediate its defence through the RecBCD complex. To

test this hypothesis, we expressed the Kiwa system in a set of *E. coli* strains that had

genomic deletions of the individual RecBCD genes and infected the cells with phage.

Our findings show that RecBCD plays a critical role in Kiwa ECOR8 and ECOR12

defence, as the system becomes inactive against phage infection in its absence (Fig.

4f, Supplementary Fig. 6a). Curiously, RecD seems to play a more relevant role than

RecB or RecC in Kiwa ECOR12, especially against phage T4.

Although RecD is known to possess ATP hydrolysis activity[47], its absence (or the

absence of RecB or RecC) had no impact on ATP depletion in Kiwa cells during

phage infection (Fig. 4g, Supplementary Fig. 6b). The fact that RecBCD is necessary

for Kiwa protection but not for ATP depletion confirms that ATP depletion is not the

effector mechanism of Kiwa and that it is more likely related to the interaction

between KwaA and KwaB.

We hypothesized that KwaB may interact directly with RecD or RecBCD to activate

the defence. However, when we expressed tagged KwaB in *E. coli* and tested for

interaction with RecBCD proteins, none were pulled down with KwaB. During the

SEC experiments aimed at identifying a potential interaction between KwaB and

RecBCD, we observed that KwaB formed oligomers. This was evidenced by a

shorter column retention time of the protein compared to when KwaB was co-

expressed with the C-terminal of KwaA (Supplementary Fig. 4c). Mass spectrometry

confirmed the presence of KwaB in all fractions. These findings suggest that the

activation of KwaB may require oligomerization.

To further explore the interplay between KwaB and RecBCD in Kiwa defence, we

assessed the level of protection provided by KwaB individually in cells with complete

RecBCD or deleted individual genes. Our results indicate that KwaB provides full

protection only in the presence of complete RecBCD (Fig. 4h, Supplementary Fig.

6d), suggesting that KwaB and RecBCD work in concert to control phage infection.

Overall, our findings show that Kiwa defends against phage infection through a

mechanism that involves activation of RecBCD by KwaB.

Discussion

In this study, we propose a mechanism for Kiwa defence that involves the activation

of Kiwa through the inhibition of host RNAP by phage-encoded proteins, ultimately

resulting in a RecBCD-dependent control of phage DNA replication (Fig. 4i).

We used phage mutants that escaped defence to identify the activators of Kiwa

ECOR8 and ECOR12 defence, which were found to be the inhibitors of *E. coli* RNAP

σ^s of phages T7s (gp5.7) and T3 (T3p30). Additional experiments using a

transcription inhibitor, rifampicin, established that Kiwa detects the inhibition of host

RNAP instead of the phage proteins. This is consistent with the observation that

other defence system, such as dCTP deaminase, dGTPase, AVAST type III, and

toxIN[29,35,48], also sense phage infection via binding to RNAP inhibitor or guarding the

integrity of the host RNAP function, indicating that this is a common strategy

employed by multiple defence systems. Interestingly, the phage-encoded RNAP

inhibitor that activates Kiwa from ECOR8 and ECOR12 does not activate Kiwa from

E. coli D9, suggesting that distantly related Kiwa operons are activated by different

triggers, resulting in their distinct anti-phage specificities.

Complementation assays in *trans* showed that the RNAP inhibitor proteins activate

Kiwa defence only when the full system (KwaA+KwaB) is present, while KwaB alone

is capable of providing anti-phage defence in the presence or absence of the phage-

encoded RNAP inhibitor. Because of this we propose that KwaA is the protein

responsible for sensing phage infection in Kiwa.

Sensing of host RNAP inhibition by KwaA likely controls the activity of KwaB, which

we found to bind to the C-terminal of KwaA. However, when we deleted the C-

terminal of KwaA, protection was abolished, even though KwaB should be active in

the cytosol. We hypothesise that some other part of KwaA interacts with KwaB and

limits its activity, mostly likely the 20 amino acids of KwaA predicted to be cytosolic

(G75-F94). Moreover, analysis of the ATP levels in Kiwa cells revealed that the

interaction between KwaA and KwaB results in a reduction of ATP. However, this

ATP depletion is not linked to the protective function of KwaB and is not considered

the effector mechanism of Kiwa. Instead, we suggest that the interaction between

the Kiwa proteins or between these proteins and the trigger may cause the observed

ATP depletion.

KwaB contains the DUF4868 domain, which is commonly found in the human gut

microbiome. A homologue of this domain from *Lactobacillus casei* CRL 705 is

involved in the biosynthesis of a bacteriocin toxin[31]. The specific function of this

domain remains unknown, but its conservation across different KwaB homologues

and the loss of protection upon key amino acid substitutions suggests its significance

for anti-phage defence. Our study demonstrates that KwaB, with or without KwaA,

facilitates phage defence through a RecBCD-dependent mechanisms that hinders

the efficiency of phage DNA replication. Although pull-down assays with tagged

KwaB expressed in *E. coli* did not reveal any direct stable interaction between KwaB

and RecBCD proteins, this interaction may be transient or weak, or the effect may be

indirect. Notably, the protective effect of RecBCD is only observed in the presence of

KwaB, regardless of whether KwaA is present or not. Other defence systems have

also been shown to work alongside RecBCD, such as the CRISPR-Cas systems that

uses DNA fragments created by RecBCD to acquire new spacers targeting

phages[45,49], and the Retron Ec48 system that guards RecBCD activity and prompts

abortive infection upon inhibition of this complex by phage-encoded proteins[22]. The

mechanism of RecBCD dependence in Kiwa appears to be distinct from other

defence systems and uses the inherent anti-phage properties of RecBCD. It is

relevant to note that phage-encoded inhibitors of RecBCD do not seem to affect

Kiwa's defence mechanism, as protection against phages T4, T3, and T7s is

observed regardless of their encoded inhibitors (gp2, T3p31, and gp5.9,

respectively). This suggests that Kiwa can restore the RecBCD activity towards

these phages, either by directly modulating its activity or by preventing the RecBCD

inhibitors from affecting RecBCD.

While we could not definitively confirm that Kiwa results in an abortive infection

phenotype, we propose that this is the case at least against phages that cause host

DNA degradation. Phages T3, T7, and T4 use nucleases that rapidly degrade the

bacterial host chromosome at the early stages of phage infection[50,51]. As Kiwa

senses phage infection by detecting phage-encoded RNAP inhibitors expressed at

the middle stage of the phage infection cycle[36], it is likely that the host chromosome

is damaged beyond repair by the time the defence mechanism kicks in. Therefore,

RecBCD activity may result in degradation of phage DNA, but will not be adequate to

save the cell infected with phages that degrade the host chromosome. Additionally,

the fact that KwaB can provide protection in the absence of KwaA, but is still present

with KwaA in operons, implies that KwaB might be toxic and that its activity needs to

be regulated for cell survival. While our assays showed no evidence of a toxic effect

of KwaB on the cells, it is possible that the RecBCD-mediated action of KwaB could

have harmful effects on the cells in the long term, such as increased mutation rates.

Altogether, our findings reveal a new mechanism for defence against phages in

which Kiwa proteins reinstate RecBCD activity to regulate phage DNA replication,

independent of the presence of phage-encoded RecBCD inhibitors.

Methods

Methods

Bacterial strains and phages

E. coli Dh5α, and BL21-AI, BL21-AIΔRecB, BL21-AIΔRecC, and BL21-AIΔRecD[45]

were grown at 37 °C in Lysogeny Broth (LB) for liquid cultures or LB agar (LBA, 1.5

% (w/v) agar) for solid cultures. Whenever applicable, LB was supplemented with

chloramphenicol (25 µg/mL), spectinomycin (50 µg/mL), Isopropyl ß-D-1-

thiogalactopyranoside (IPTG, 1 mM) or L-Arabinose (0.2 %), to ensure maintenance

of plasmids or induce protein expression. Phage infection was performed in LB at 37

°C. Phages were propagated by infecting *E. coli* BL21-AI grown to an optical density

at 600 nm (OD$_{600}$) of 0.2-0.4 with 5-10 µL of a phage lysate. Infected cultures were

grown for at least 4 hours or until culture collapse. The culture was then centrifuged

at 11,000 × *g*, 4 °C for 15 min to remove cell debris, and the supernatant was filter

sterilized through a 0.22 µm filter. All strains and phages used in this study are listed

in Supplementary Table 2.

Plasmid and strain construction

Kiwa systems from *E. coli* ECOR8, ECOR12, ECOR49, and D9 were amplified by

Q5 polymerase (New England Biolabs) using primers indicated in Supplementary

Table 3. Kiwa from *Ralstonia mannitolilytica* SN82F48 was ordered synthetically as a

gBlock (Integrated DNA Technologies) (Supplementary Table 3). Kiwa systems were

cloned into pACYCDuet-1 (Novagen EMD Millipore) by restriction enzyme cloning or

Gibson assembly (NEBuilder Hifi DNA assembly Master Mix). When appropriate,

deletions, mutations and truncations of Kiwa operons were engineered by around-

the-horn PCR with primers listed in Supplementary Table 3. A plasmid with Kiwa

ECOR8 under a constitutive promoter (pUOS023) was constructed by substitution of

the T7 promoter and lac operator of pUOS017 with the constitutive Anderson

promoter J23104 (http://parts.igem.org/Promoters/Catalog/Anderson)[33]. For protein

purification, KwaA and KwaB genes of ECOR8 were cloned into the 13S-S plasmid

(Berkeley MacroLab) under an inducible T7 promoter, with a 6X His SUMO TEV N-

terminal tag. 13S-S ECOR8 KwaA C-terminus was constructed by around-the-horn

PCR on pUOS025. GFP was amplified from pARACDF (Addgene 47516) and phage

proteins gp5.7/T3p30 and gp17/T3p48 were amplified from the genome of T7s or T3

respectively using the primers indicated in Supplementary Table 3. These proteins

were cloned into pCDF1-b (Novagen) under the control of a T7 promoter. All

plasmids were transformed into *E. coli* Dh5α, extracted using the GeneJET Plasmid

Miniprep kit (Thermo Fisher), verified by Sanger sequencing (Eurofins Genomics)

and transformed into electrocompetent *E. coli* BL21-AI. Plasmids built for this study

are listed in Supplementary Table 4.

Plaque assays

Phages were ten-fold serially diluted in LB and spotted on soft agar plates (0.7 %

(w/v) agar) of *E. coli* BL21-AI containing a control plasmid or individual Kiwa

systems. For inducible systems, overnight cultures were diluted 1:50 in LB with 0.2

% L-arabinose and 1 mM IPTG, and grown for 4 hours at 37 °C with agitation before

preparing bacterial lawns for spot assays. For spot assays in strains expressing

phage proteins *in trans*, induction was performed on plate by supplementing soft

agar with L-arabinose and IPTG. Plates were left to dry at room temperature and

incubated overnight at 37 °C. Plaque forming units (PFUs) were determined by

counting the plaques after overnight incubation. Whenever plaques were too small to

be counted, a clear lysis area was set to be equal to 100 PFUs. Fold defence was

determined by calculating the ratio of phage concentration of Kiwa expressing strain

in relation to control strain. Difference in phage plaque size produced in a Kiwa strain

relative to a control was determined by measuring the plaque area of 100 individual

plaques of phages T3, T4, T7, NBG2 and S1-BL21L.

Turbidity assays

Overnight bacterial cultures of *E. coli* BL21-AI with a control plasmid or individual

Kiwa systems were diluted 1:100 in LB, supplemented with L-arabinose and IPTG if

required, and grown at 37 °C with agitation until cultures reached an OD_{600} of ≈0.3.

Cultures were normalized to an OD_{600} of 0.1 and dispensed in a 96-well plate.

Growth was monitored every 10 min in a ClarioStar plus plate reader at 37 °C with

agitation. When OD_{600} reached ≈0.3, cells were either kept uninfected or infected

with phage at the desired MOI. Growth was then monitored for 500 min post

infection. For cells expressing phage proteins *in trans*, L-arabinose (0.2%) and ITPG

(1 mM) were added simultaneously with the phage.

Time post infection assays

Overnight bacterial cultures of *E. coli* BL21-AI with a control plasmid or individual

Kiwa systems were diluted 1:100 in LB medium, supplemented with L-arabinose (0.2

%) and IPTG (1 mM), and grown at 37 °C with agitation until an OD_{600} of ≈0.3.

Cultures were normalized and infected with phage T3 or T4 at an MOI of 3. At 0, 60

and 90 min post infection, a sample was taken, serially diluted, and phages were

quantified by plaque assay on a phage-sensitive bacterial lawn (*E. coli* BL21-AI +

pUOS016). PFUs were counted after overnight incubation at 37 °C.

Adsorption assays

Overnight bacterial cultures of *E. coli* BL21-AI with a control plasmid or individual

Kiwa systems were diluted 1:100 in LB medium, supplemented with L-arabinose (0.2

%) and IPTG (1 mM), and grown at 37 °C with agitation until an OD of ≈0.3. Then,

500 µL of bacterial culture were infected with phage T3 or T4 at an MOI of 0.005,

and incubated for 15 min at 37 °C with agitation. A sample was taken and serially

diluted to obtain the total phage concentration. Subsequently, cells were centrifuged

(8,000 × *g*, 1 min) and a sample from the supernatant was taken and serially diluted

to obtain the non-adsorbed phage concentration. Dilutions were spotted on a phage-

sensitive bacterial lawn (*E. coli* BL21-AI + pUoS016). The concentration of adsorbed

phages was determined by subtracting non-adsorbed phage concentration from the

total phage concentration. The percentage of adsorbed phages was calculated as

the ratio between adsorbed and total phages.

Intra- and extracellular phage concentrations

Overnight bacterial cultures of *E. coli* BL21-AI with a control plasmid or Kiwa ECOR8

were diluted 1:100 in LB medium, supplemented with L-arabinose (0.2 %) and IPTG

(1 mM), and grown at 37 °C with agitation until cultures reached an OD_{600} of ≈0.3.

Cultures were normalized and infected with phage T4 at an MOI of 10^{-5}. At 90

minutes post infection a sample was taken and centrifuged (8,000 × *g*, 1 min, 4 °C).

The supernatant was serially diluted to obtain the extracellular phage concentration.

The remaining supernatant was discarded and the cell pellet was washed once with

ice cold LB and harvested by centrifugation (8,000 × g, 1 min, 4 °C). The

supernatant was discarded and the cell pellet was re-suspended in ice cold LB, after

which a 1:1 volume of chloroform was added. The sample was mixed by vortexing,

and the aqueous layer was extracted after centrifugation (8,000 × g, 5 min, 4 °C) and

serially diluted to obtain the intracellular phage concentration. Dilutions were spotted

on a phage-sensitive bacterial lawn (*E. coli* BL21-AI + pUOS016), and the PFUs

were counted after overnight incubation at 37 °C.

DNA replication assay

Overnight bacterial cultures of *E. coli* BL21-AI with a control plasmid or individual

Kiwa systems were diluted 1:100 in LB medium, supplemented with L-arabinose (0.2

%) and IPTG (1 mM), and grown at 37 °C with agitation until cultures reached an

OD_{600} of ≈0.3. Cultures were normalized and infected with phage T3 or T4 at an MOI

of 3. At 0 and 30 min post infection, a 5 mL sample was taken and centrifuged

(15,000 × g, 5 min, 4 °C). The cell pellets were snap frozen using liquid nitrogen and

stored at -80 °C. Total DNA was extracted using the GeneJET Genomic DNA

Isolation kit (Thermo Fisher), using the Gram-negative protocol. Samples were

sequenced at SeqCenter (Pittsburgh, PA, USA), where sample libraries were

prepared using the Illumina DNA Prep kit and IDT 10bp UDI indices, and sequenced

on an Illumina NextSeq 2000, producing 2x151bp reads. Demultiplexing, quality

control and adapter trimming was performed with bcl-convert (v3.9.3). Reads were

aligned to *E. coli* BL21-AI and phage T3 or T4 reference genomes (CP047231,

NC_003298 and NC_000866) using MiniMap2[52] in Geneious prime 2023.0.1.

Replication efficiency was determined by calculating the increase of phage genomes

to the number of cells at time 0. The percentage was calculated by comparison of

Kiwa infected cells to control cells over time.

RecBCD mutant assays

463   Strains with individual deleted RecBCD subunits were transformed by

464   electroporation with a control plasmid or individual Kiwa systems and recovered on

465   LBA supplemented with kanamycin (50 µg/mL). Colonies were picked and grown

466   overnight at 37 °C. Spot assays were performed with Phage T3 and T4 as described

467   above for inducible systems.

ATP assays

Overnight bacterial cultures of *E. coli* BL21-AI, or *E.coli* BL21-AI ΔRecB , ΔRecC or

ΔRecD with a control plasmid or individual Kiwa systems were diluted 1:100 in LB

medium, supplemented with L-arabinose (0.2 %) and IPTG (1 mM), and grown at 37

°C with agitation until an OD_{600} of ≈0.3. Cultures were normalized and either kept

uninfected or infected with phage T3 or T4 at an MOI of 3 or 0.03. At 0 and 30 min

post infection a sample was taken and serially diluted, and ATP concentration was

determined by following the BacTiterGlo cell viability kit protocol (Promega).

Luminescence was measured in ClarioStar plus plate reader at room temperature.

ATP levels of Kiwa expressing cells were compared to control cells at 0 and 30 min

post infection.

Isolation and sequencing of escape phages

Escape phages were isolated by picking single plaques at the lowest dilution of a

spot assay on a lawn of *E. coli* cells expression Kiwa ECOR8 under a constitutive

promoter. The resistant phenotype of each escape phage was determining by

comparing the EOP of the escape phage to the EOP of the wild type phage in the

presence of Kiwa ECOR8. Isolated phages were further propagated by infecting a

liquid culture of Kiwa ECOR8 as described above. Phage lysates were pelleted

following the NaCl/PEG8000 precipitation protocol as previously described[53]. The

phage pellets were re-suspended in LB, and treated with DNase I and RNase A (1

µg/mL each) for 30 min at room temperature. EDTA (20 mM), proteinase K (50

µg/mL) and SDS (0.5%) were added to the suspension and incubated overnight at

56 °C. DNA was extracted using the Zymo Clean & Concentrator-5 kit (Zymo

Research). Samples were sequenced at SeqCenter (Pittsburgh, PA, USA), where

sample libraries were prepared using the Illumina DNA Prep kit and IDT 10bp UDI

indices, and sequenced on an Illumina NextSeq 2000, producing 2x151bp reads.

Demultiplexing, quality control and adapter trimming was performed with bcl-convert

(v3.9.3). Mutations were identified by mapping sequencing reads to the respective

reference phage genomes (NC_003298 for T3 and Supplementary Table 1 for T7s)

using Breseq[54] v0.37.0 with default parameters. Only mutations that occurred in the

isolated mutant but not in the wild type phage were considered. Silent mutations

within protein-coding regions were also neglected.

Rifampicin assay

Overnight bacterial cultures of *E. coli* BL21-AI or BL21-AI ECOR8c were diluted

1:100 in LB medium and grown at 37 °C with agitation until an OD600 of ≈0.3.

Cultures were infected with phage T3 mutant n.2 at an MOI of 0.6 and treated with

Rifampicin (100 µg/mL) or kept untreated. At 0, 30, 60, and 90 min post infection a

sample was taken and serially diluted. Phages were quantified by plaque assay on a

phage-sensitive bacterial lawn (*E. coli* BL21-AI + pUOS022), and the PFUs were

counted after overnight incubation at 37 °C.

Bacterial growth upon induction of phage trigger

509 Overnight cultures of *E. coli* BL21-AI ECOR8c or a control with a plasmid encoding

510 phage T3 gp30 were normalised to an OD_{600} of 1.0. Normalised cultures were

511 serially diluted in LB and spotted onto LBA plates with or without L-arabinose (0.2 %)

512 and IPTG (1 mM)). Plates were incubated overnight at 37 °C for colony forming units

513 (CFU) counting.

Protein expression and purification

Overnight cultures of *E. coli* BL21-AI with His-SUMO-TEV::ECOR8 KwaA C-terminus

and ECOR8 KwaB, or His-SUMO-TEV::ECOR8 KwaB with mVenus YFP as a

control were diluted in LB and grown at 37 °C with agitation until an OD_{600} of ≈0.4.

Grown cultures were incubated on ice for 1 hour and protein expression was induced

with L-arabinose (0.2 %) and IPTG (1 mM), followed by overnight incubation at 20 °C

and 150 rpm. The overnight cultures were harvested by centrifugation (8,000 × g, 30

min, 4 °C). The supernatant was discarded and the cell pellets were re-suspended in

lysis/wash buffer (50 mL per 1 L of initial culture, 100 mM Tris-HCl, 150 mM NaCl, 1

mM DTT, 5% glycerol, 25 mM imidazole, pH 7.5) supplemented with Pierce™

Protease Inhibitor Tablets EDTA-free (Thermo Fisher). Cell lysis was performed

three times in a cooled French press (15Gpa). The cell lysate was centrifuged

(17,000 × g, 30 min, 4 °C) and debris were removed by filtration (0.45 μm PES). The

filtered supernatant was incubated with HisPur™ Ni-NTA Resin (Thermo Fisher)

(500 μL/50 mL lysate) pre-washed with 20 mL of lysis/wash buffer for 30 min at 4 °C.

The sample was loaded onto a 5 mL Pierce™ Disposable Column (Thermo Fisher)

for gravity-flow affinity chromatography. The column was washed with 15 mL of ice-

cold lysis/wash buffer, and the protein was eluted with ice-cold elution buffer (100

mM Tris-HCl, 150 mM NaCl, 1 mM DTT, 5% glycerol, 250 mM imidazole, pH 7.5).

Pooled fractions were concentrated (Amicon Centrifugal concentrator with Ultracel

membrane, 3 kDa MWCO), and the buffer exchanged to running buffer (100 mM

Tris-HCl, 150 mM NaCl, 1 mM DTT, 5% glycerol, pH 7.5). The resulting protein

sample was subjected to size exclusion chromatography using a Superdex 200

increase 10/300 GL (Cytiva) column equilibrated with running buffer with 0.5 mL/min

flow rate using running buffer as a mobile phase. Fractions of interest were analysed

by SDS-PAGE, snap frozen in liquid nitrogen and stored at -80 °C until further use.

Fractions containing protein brands corresponding to the size of tagged KwaA C-

terminus, tagged KwaB or KwaB were analysed by mass spectrometry as detailed

below.

Protein identification by mass spectrometry

The proteins in the purified fractions were analysed by mass spectrometry at the

Mass Spectrometry Facility of the Department of Biotechnology at Delft University of

Technology (Netherlands), as previously described[55]. Briefly, proteins were identified

following in-solution digestion, or after SDS-PAGE analysis followed by in-gel

digestion. In solution digestion was performed by mixing 100 µL of each protein

solution with 440 µL of 200 mM ammonium bicarbonate buffer (ABC). To this

solution, 30 µL of 10 mM dithiothreitol (DTT) solution was added, and incubated at

37 ˚C, 300 rpm for 1 hour. Next, 30 µL of 20 mM Iodoacetamide (IAA) solution was

added and incubated in the dark, at room temperature, for 30 minutes. 5 µL of

Trypsin solution (0.1 µg/µL) were added to the solution, and the solution was

incubated at 37°C, 300 rpm overnight. Samples were desalted and cleaned using

solid phage extraction and dried afterwards. The dried samples were resuspended in

15 μL 3% acetonitrile in water plus 0.1 % formic acid under careful vortexing. An

aliquot corresponding to approximately 100 ng protein digest was analysed using a

nano-LC EASY 1200 (Thermo Scientific, Germany) coupled online to a QE plus

Orbitrap mass spectrometer (Thermo Scientific, Germany), which was operated in

data-dependent acquisition (DDA) mode, acquiring MS2 spectra of the top 10

peptide signals. Mass spectrometric raw data were analysed using PEAKS Studio X

(Bioinformatics Solutions Inc., Canada) by comparing fragmentation spectra to a

database consisting of the target protein sequences and the *E. coli* BL21 AI

proteome (NCBI). Database searching was performed allowing 20 ppm parent mass

error, 0.2 Da fragment mass error, 2 missed cleavages, carbamidomethylation as

fixed, and oxidation and deamidation as variable peptide modifications. Peptide

matches and protein identifications were filtered for 1% false discovery rate. Protein

identifications with >1 unique peptide sequences were considered as confident

matches.

Bioinformatic analysis

All prokaryotic genomes were retrieved from NCBI, and Kiwa containing genomes

were identified using the Prokaryotic Antiviral Defense LOCator (PADLOC)[10]. Only

complete Kiwa systems that contained at least one KwaA and one KwaB gene within

the operon, were considered for further analysis (Supplementary Table 5). KwaA and

KwaB protein sequences were aligned using Clustal Omega[56] with default

parameters and the noise was removed with CIAlign[57]. The multiple sequence

alignment was used on IQ-tree[58] to generate a tree with -alrt 1500, -bb 1500

parameters, followed by masking of regions with non-homologous adjacent

characters using prequal. iTOL[59] was used for tree visualisation. Protein domains of

KwaA and KwaB were identified using HHpred[60]. Transmembrane domains were

identified using DeepTMHMM[61], and protein size was predicted using the ProtParam

webtool[62]. Sequence alignments of KwaA C-terminus and KwaB were computed by

Clustal Omega[56] and visualised using Geneious Prime v2023.0.1. For KwaB

mutants, a total of nine homologs present in the KwaB ECOR8 and ECOR12 cluster

were selected and aligned to determine amino acid conservation.

Statistical analysis

Unless stated otherwise, experiments were performed in biological triplicates and are

represented as the mean and standard deviation. Statistical significance was

calculated by ration paired t-test or by two-way ANOVA with sidak's multiple

comparison test, with a significance level of 0.05.

Data availability

All data supporting the findings of this study are available within the paper and in the

Supplementary Data. Raw data is available from Figshare via the following DOIs:

Sequencing reads of phage escape mutants

https://doi.org/10.6084/m9.figshare.21652433.v1, sequencing reads of phage

infection of Kiwa cells https://doi.org/10.6084/m9.figshare.21648587.v1, sequencing

reads of ECOR8c https://doi.org/10.6084/m9.figshare.22093256.v1, and mass

spectrometry data https://doi.org/10.6084/m9.figshare.22147931.v1. All unique

bacterial strains, phages, and plasmids generated in this study are available from the

corresponding author upon request.

Acknowledgements

We thank Morgen Hedges for bacteriophage illustrations, Cristian Aparicio-
Maldonado for helping with protein purification, and Dita Heikens and Martin Pabst
from TU Delft for the mass spectroscopy analysis. This work was supported by grant
RGS\R2\222312 from the Royal Society to F.L.N.

Author contributions

Conceptualization, F.L.N.; Methodology; T.C.T. and F.L.N.; Formal Analysis, T.C.T.,
F.L.N.; Investigation, T.C.T., Y.W., A.N., and R.M.; Visualization, T.C.T.; Data
Curation, T.C.T., Y.W., F.L.N.; Writing – Original Draft, T.C.T., F.L.N.; Writing –
Review & Editing, F.L.N.; Resources, F.L.N.; Funding Acquisition, F.L.N..

Declaration of interests

The authors declare no competing interests.

Figure legends

Fig. 1: Kiwa orthologues provide defence against multiple coliphages. a Clustering of KwaA (left) and KwaB (right) homologues in prokaryotic genomes. The tree was built using only non-redundant sequences. Each cluster is identified by a specific colour shading. The tree includes annotations for Kiwa operons that were tested in this study, and any Kiwa operons that have previously been confirmed to provide anti-phage activity are indicated with a shield symbol. **b** Schematic representation of the Kiwa operons tested in this study, obtained from *Escherichia coli* (ECOR8, ECOR12, ECOR49, D9) and *Ralstonia mannitolilytica* (SN82F48) strains. TM, transmembrane domain. DUF, domain of unknown function. **c** Anti-phage activity of five Kiwa operons shown as fold protection of efficiency of platting assays. m, plaque morphology change relative to a control strain expressing mVenusYFP. **d** Growth curves of Kiwa ECOR12 or control cells infected by phage T4 at an MOI of 3 or 0.03. Data shows the mean of three biological replicates. **e** Effect of Kiwa ECOR8 (left) and ECOR12 (right) on phage propagation. Data represents plaque forming units (PFU) per mL (PFU/mL) of phage T4 at different time points post infection of Kiwa or control cells. Asterisks mark statistically significant PFU/mL decreases relative to a control (Two-way ANOVA, $p < 0.05$). Bar graphs represent the average of 3 biological replicates with individual points overlaid.

Fig. 2: Transmembrane domains of KwaA and DUF4868 domain of KwaB are essential for the anti-phage activity by Kiwa. a Impact of deletions, truncations and point mutations on the ability of Kiwa ECOR8 to defend against phages T3 and T4. The domain organisation of Kiwa ECOR8 is shown on top, with transmembrane (TM) domains of KwaA in dark green, C-terminal domain of KwaA, in light green, and

point mutations in the DUF4868 domain of KwaB in orange. The fold defence was

calculated by dividing the number of PFU (plaque-forming units) per mL of control

and Kiwa strains. Asterisk indicate a statistically significant decrease in defence

compared to Kiwa ECOR8 (Two-way ANOVA, p <0.05). The bars represent the

average of two biological replicates with individual data points overlaid. **b** Effect of

reconstituing two TM domains of KwaA on the defensive activity of SN82F48 against

phages T3 and T7s. A schematic of the SN82F48 operon with added TM domains in

green is shown on top. The fold defence was calculated as the ratio of PFU/mL on

control and Kiwa strains. Asterisks indicate a statistically significant decrease in

defence compared to truncated SN82F48 (Two-way ANOVA, p <0.05). The bars

represent the average of two biological replicates with individual data points overlaid.

c Pulldown assay of ECOR8 KwaA C-terminus and KwaB. The KwaA C-terminus

was N-terminally tagged with a HIS-Sumo-TEV tag and co-expressed with KwaB. A

schematic of the heterologous expression system used for protein overexpression is

shown on top. The size exclusion chromatogram shows absorbance at 280 nm on

the left, with elution volume corresponding to the peak run on SDS-PAGE on the

right.

Figure 3. Kiwa is activated upon detecting inhibition of host RNA polymerase.

a Growth curves of *E. coli* with Kiwa ECOR8 under expression of a constitutive

promoter (ECOR8c) or control cells infected with phage at MOI of 3 or 0.03. Lines

show the mean of three biological replicates. **b** Kiwa ECOR8c exhibits increased

defence against three phages in comparison to ECOR8 (T7 inducible promoter). The

data represents fold defence in relation to control cells, and it is the average of two

biological replicates. Plaque morphology change compared to control is indicated by

"m". **c** Effect of ECOR8c on plaque size of infecting coliphages. The data represents

plaque sizes of phages infecting ECOR8c cells relative to the average size of

plaques on controls cells. Asterisks indicate a statistically significant decrease in

plaque size in Kiwa cells compared to the control strain (Two-way ANOVA, $p < 0.05$).

The mean of 100 replicates is shown, with individual points overlaid. **d** Reduction of

defence by Kiwa ECOR8, ECOR12 or D9 when infected by phage escape mutants

(n.1-n.5). Fold defence is calculated as the ratio of PFU/mL on Kiwa and control

strains. The average of two biological replicates is shown, with individual data points

overlaid. Asterisks indicate a statistically significant decrease in defence compared

to control (Two-way ANOVA, $p < 0.05$). **e** Mutations in gp5.7 of phage T7s and T3p30

of phage T3 allow evasion of Kiwa defence. Phages with complete deletions of gp5.7

or T3p30 are annotated separately. The full list of mutations and deletions in the

escape mutants is detailed in Supplementary Table 2. **f** Reconstitution of gp5.7 and

T3p30 recovers ECOR8c defence activity against phage escape mutants. Fold

defence was calculated as the ratio of PFU/mL in Kiwa and control. Asterisks

indicate a statistically significant increase of defence in relation to a control of

ECOR8c-GFP (Ratio paired t-test, $p < 0.05$). The bars represent the mean of two

biological replicates, with individual data points overlaid. **g** Growth curves of *E. coli*

Kiwa ECOR8c expressing T3p30, gp5.7 or GFP (control) infected with phage escape

mutants at an MOI of 0.6 or 0.06. The curves represent the mean of three biological

replicates. **h** Effect of deletion of Kiwa genes on the infectivity of phage escape

mutants. Asterisks indicate a statistically significant decrease of phage concentration

(PFU/mL) in relation to control cells (Two-way ANOVA, $p < 0.05$). The bars represent

the average of two biological replicates, with individual data points overlaid. Images

are representatives of phage mutant T3 n.2 infecting cells at a decreasing phage

concentration.

Figure 4. Kiwa reduces phage DNA replication via a RecBCD-dependent

mechanism. a Effect of Kiwa on phage absorption to *E. coli* cells. The percentage of

phages adsorbed to cells after 15 min was calculated relative to the total amount of

phage. The bar graphs show the mean of three biological replicates with individual

points overlaid. **b** Quantification of intra- and extracellular phages on cultures of *E.*

coli expressing Kiwa ECOR8 or a control. Data shows plaque forming units

(PFU/mL) of phage T4 at 90 minutes post infection. Asterisks indicate a statistically

significant decrease of PFU compared to control cells (Two-way ANOVA, $p < 0.05$).

The bars shows the mean of three biological replicates, with individual data points

overlaid. **c** Efficiency of phage DNA replication during infection of control and Kiwa

cells. Total DNA was extracted from cells infected with phage at an MOI of 3. The

data represents the replication efficiency of phage DNA per minute over 30 minutes

of infection, compared to control cells, and is shown as the mean of two biological

replicates, with individual data points overlaid. **d** Effect of Kiwa on total ATP levels

during phage infection. Data represents ratio of total ATP levels of cells expressing

Kiwa relative to the control. Cells where either kept uninfected or infected with phage

T3 or T4 at an MOI of 3 or 0.03. Asterisks indicate a statistically significant decrease

of ATP in relation to uninfected cells at the same time point (Two-way ANOVA, p

<0.05). The bars show the average of two biological replicates with individual data

points overlaid. **e** Effect of deleting Kiwa genes on total ATP levels during phage

infection. **f** Effect of deleting RecBCD subunits on Kiwa defence. Data represents

PFU/mL of phage T4 infecting control or Kiwa cells. Asterisks indicate a statistically

significant decrease of PFU in relation to control cells with complete RecBCD (Two-

way ANOVA, p <0.01). The bars show the average of two biological replicates, with

individual data points overlaid. **g** Effect of the deletion of RecD on total ATP levels

during phage infection. Data represents ratio of total ATP levels of Kiwa cells with or

without RecD relative to the control (no defence). Cells where either kept uninfected

or infected with phage T3 or T4 at an MOI of 3. Asterisks indicate a statistically

significant decrease of ATP compared to uninfected cells at the same time point

(Two-way ANOVA, p <0.01). The bars show the average of two replicates, with

individual data points overlaid. **h** Effect of deleting RecBCD subunits on KwaB

defence activity. Data represents PFU/mL of phage T4 infecting control or ECOR8

KwaB (ΔKwaA) cells with a complete RecBCD complex or single subunit deletions.

Asterisks indicate a statistically significant decrease of PFU compared to complete

RecBCD expressing cells (Two-way ANOVA, p <0.01). The bars show the average

of two biological replicates, with individual data points overlaid. **i** A model for the

mechanism of the Kiwa defence system. Phage infection is sensed by the inhibition

of the host RNA polymerase by phage proteins, leading to an ATP depletion event

that causes the release and activation of KwaB, likely via oligomerisation. The active

KwaB re-activates RecBCD, leading to a decrease in phage DNA replication to

control the spread of the infection.

Supplementary data

Supplementary Fig. 1: Distribution of Kiwa homologues in bacterial genomes and
their anti-phage activity.

Supplementary Fig. 2: Anti-phage activity of modified Kiwa systems from *E. coli*
ECOR8 and SN82F48.

Supplementary Fig. 3: KwaB oligomerisation and conservation.

Supplementary Fig. 4: *E. coli* ECOR8c has increased defence against phage
infection.

Supplementary Fig. 5: Mutant phage challenges of Kiwa cells expressing phage
proteins in *trans*.

Supplementary Fig. 6: The effect of RecBCD and KwaB on Kiwa defence.

Supplementary Table 1: Mutation analysis of bacterial strains and phage escape
mutants.

Supplementary Table 2: Strains and phages used in this study.

Supplementary Table 3: Primers and gblocks used in this study.

Supplementary Table 4: Plasmids used in this study.

Supplementary Table 5: Kiwa homologues used for hierarchical clustering.

References

1   Anzalone, A. V., Koblan, L. W. & Liu, D. R. Genome editing with CRISPR–
Cas nucleases, base editors, transposases and prime editors. *Nat Biotechnol*
38, 824-844, doi:10.1038/s41587-020-0561-9 (2020).

2   Millman, A. *et al.* An expanding arsenal of immune systems that protect
bacteria from phages. *bioRxiv*, doi:10.1101/2022.05.11.491447 (2022).

3   Doron, S. *et al.* Systematic discovery of antiphage defense systems in the
microbial pangenome. *Science* **359**, eaar4120, doi:10.1126/science.aar4120
(2018).

4   Ofir, G. *et al.* DISARM is a widespread bacterial defence system with broad
anti-phage activities. *Nat Microbiol* **3**, 90-98, doi:10.1038/s41564-017-0051-0
(2018).

5   Goldfarb, T. *et al.* BREX is a novel phage resistance system widespread in
microbial genomes. *EMBO J* **34**, 169-183, doi:10.15252/embj.201489455
(2015).

6   Gao, L. *et al.* Diverse enzymatic activities mediate antiviral immunity in
prokaryotes. *Science* **369**, 1077-1084, doi:10.1126/science.aba0372 (2020).

7   Owen, S. V. *et al.* Prophages encode phage-defense systems with cognate
self-immunity. *Cell Host Microbe* **29**, 1620-1633.e1628,
doi:10.1016/j.chom.2021.09.002 (2021).

8   Rousset, F. *et al.* Phages and their satellites encode hotspots of antiviral
systems. *Cell Host Microbe* **30**, 740-753.e745,
doi:10.1016/j.chom.2022.02.018 (2022).

9   Vassallo, C. N., Doering, C. R., Littlehale, M. L., Teodoro, G. I. C. & Laub, M.
T. A functional selection reveals previously undetected anti-phage defence
systems in the E. coli pangenome. *Nat Microbiol* **7**, 1568-1579,
doi:10.1038/s41564-022-01219-4 (2022).

10   Payne, L. J. *et al.* Identification and classification of antiviral defence systems
in bacteria and archaea with PADLOC reveals new system types. *Nucleic*
Acids Res **49**, 10868-10878, doi:10.1093/nar/gkab883 (2021).

11   Isaev, A. *et al.* Phage T7 DNA mimic protein Ocr is a potent inhibitor of BREX
defence. *Nucleic Acids Res* **48**, 5397-5406, doi:10.1093/nar/gkaa290 (2020).

12   Gordeeva, J. *et al.* BREX system of Escherichia coli distinguishes self from
non-self by methylation of a specific DNA site. *Nucleic Acids Res* **47**, 253-265,
doi:10.1093/nar/gky1125 (2018).

13   Bravo, J. P. K., Aparicio-Maldonado, C., Nobrega, F. L., Brouns, S. J. J. &
Taylor, D. W. Structural basis for broad anti-phage immunity by DISARM. *Nat*
Commun **13**, 2987, doi:10.1038/s41467-022-30673-1 (2022).

14   Aparicio-Maldonado, C. *et al.* Class I DISARM provides anti-phage and anti-
conjugation   activity   by   unmethylated   DNA   recognition.   *bioRxiv*,
doi:10.1101/2021.12.28.474362 (2021).

15   Duncan-Lowey, B., McNamara-Bordewick, N. K., Tal, N., Sorek, R. &
Kranzusch, P. J. Effector-mediated membrane disruption controls cell death in
CBASS   antiphage   defense.   *Mol   Cell*   **81**,   5039-5051.e5035,
doi:10.1016/j.molcel.2021.10.020 (2021).

16   Cohen, D. *et al.* Cyclic GMP–AMP signalling protects bacteria against viral
infection. *Nature* **574**, 691-695, doi:10.1038/s41586-019-1605-5 (2019).

17   Ofir, G. *et al.* Antiviral activity of bacterial TIR domains via signaling molecules
that trigger cell death. *bioRxiv*, doi:10.1101/2021.01.06.425286 (2021).

18   Ka, D., Oh, H., Park, E., Kim, J.-H. & Bae, E. Structural and functional
evidence of bacterial antiphage protection by Thoeris defense system via
NAD+ degradation. *Nat Commun* **11**, 2816, doi:10.1038/s41467-020-16703-w
(2020).

19   Tal, N. *et al.* Cyclic CMP and cyclic UMP mediate bacterial immunity against
phages. *Cell* **184**, 5728-5739.e5716, doi:10.1016/j.cell.2021.09.031 (2021).

20   Cheng, R. *et al.* The prokaryotic Gabija complex senses both viral
transcription and DNA metabolism for antiviral defense. Research Square,
doi: 10.21203/rs.3.rs-1703025/v1 (2022).

21   Cheng, R. *et al.* A nucleotide-sensing endonuclease from the Gabija bacterial
defense system. *Nucleic Acids Res* **49**, 5216-5229, doi:10.1093/nar/gkab277
(2021).

22   Millman, A. *et al.* Bacterial Retrons Function In Anti-Phage Defense. *Cell* **183**,
1551-1561.e1512, doi:10.1016/j.cell.2020.09.065 (2020).

23   Gao, Y. *et al.* Molecular basis of RADAR anti-phage supramolecular
assemblies. *Cell*, doi:10.1016/j.cell.2023.01.026 (2023).

24 Duncan-Lowey, B. *et al.* Cryo-EM structure of the RADAR supramolecular
anti-phage defense complex. *Cell*, doi:10.1016/j.cell.2023.01.012 (2023).

25 Garb, J. *et al.* Multiple phage resistance systems inhibit infection via SIR2-
dependent NAD+ depletion. *Nat Microbiol* **7**, 1849-1856, doi:10.1038/s41564-
022-01207-8 (2022).

26 Gao, L. A. *et al.* Prokaryotic innate immunity through pattern recognition of
conserved   viral   proteins.   *Science*   **377**,   eabm4096,
doi:10.1126/science.abm4096 (2022).

27 Johnson, A. G. *et al.* Bacterial gasdermins reveal an ancient mechanism of
cell death. *Science* **375**, 221-225, doi:10.1126/science.abj8432 (2022).

28 Hsueh, B. Y. *et al.* Phage defence by deaminase-mediated depletion of
deoxynucleotides   in   bacteria.   *Nat   Microbiol*   **7**,   1210-1220,
doi:10.1038/s41564-022-01162-4 (2022).

29 Tal, N. *et al.* Bacteria deplete deoxynucleotides to defend against
bacteriophage infection. *Nat Microbiol* **7**, 1200-1209, doi:10.1038/s41564-022-
01158-0 (2022).

30 Bernheim, A. *et al.* Prokaryotic viperins produce diverse antiviral molecules.
Nature **589**, 120-124, doi:10.1038/s41586-020-2762-2 (2021).

31 Cuozzo, S. A., Sesma, F., Palacios, J. M., de Ruíz Holgado, A. d. P. & Raya,
R. R. Identification and nucleotide sequence of genes involved in the
synthesis of lactocin 705, a two-peptide bacteriocin from Lactobacillus casei
CRL 705. *FEMS Microbiol Let* **185**, 157-161, doi:10.1016/S0378-
1097(00)00093-8 (2000).

32 Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold.
Nature **596**, 583-589, doi:10.1038/s41586-021-03819-2 (2021).

33 Florea, M. *et al.* Engineering control of bacterial cellulose production using a
genetic toolkit and a new cellulose-producing strain. *Proc Natl Acad Sci U S A*
113, E3431-E3440, doi:10.1073/pnas.1522985113 (2016).

34 Lopatina, A., Tal, N. & Sorek, R. Abortive Infection: Bacterial Suicide as an
Antiviral Immune Strategy. *Ann Rev Virol* **7**, 371-384, doi:10.1146/annurev-
virology-011620-040628 (2020).

35 Stokar-Avihail, A. *et al.* Discovery of phage determinants that confer
sensitivity   to   bacterial   immune   systems.   *bioRxiv*,
doi:10.1101/2022.08.27.505566 (2022).

36　Tabib-Salazar, A. *et al.* T7 phage factor required for managing RpoS in Escherichia coli. *Proc Natl Acad Sci U S A* **115**, E5353-E5362, doi:10.1073/pnas.1800429115 (2018).

37　Tabib-Salazar, A. *et al.* Full shut-off of Escherichia coli RNA-polymerase by T7 phage requires a small phage-encoded DNA-binding protein. *Nucleic Acids Res* **45**, 7697-7707, doi:10.1093/nar/gkx370 (2017).

38　Schmitt, C. K., Kemp, P. & Molineux, I. J. Genes 1.2 and 10 of bacteriophages T3 and T7 determine the permeability lesions observed in infected cells of Escherichia coli expressing the F plasmid gene pifA. *J Bacteriol* **173**, 6507-6514, doi:10.1128/jb.173.20.6507-6514.1991 (1991).

39　Parma, D. H. *et al.* The Rex system of bacteriophage lambda: tolerance and altruistic cell death. *Genes Dev* **6**, 497-510, doi:10.1101/gad.6.3.497 (1992).

40　Cheng, X., Wang, W. & Molineux, I. J. F exclusion of bacteriophage T7 occurs at the cell membrane. *Virology* **326**, 340-352, doi:10.1016/j.virol.2004.06.001 (2004).

41　Millman, A., Melamed, S., Amitai, G. & Sorek, R. Diversity and classification of cyclic-oligonucleotide-based anti-phage signalling systems. *Nat Microbiol* **5**, 1608-1615, doi:10.1038/s41564-020-0777-y (2020).

42　Cecchini, N., Schmerer, M., Molineux, I. J., Springman, R. & Bull, J. J. Evolutionarily Stable Attenuation by Genome Rearrangement in a Virus. *G3 Gene Genome Genet* **3**, 1389-1397, doi:10.1534/g3.113.006403 (2013).

43　Dillingham, M. S. & Kowalczykowski, S. C. RecBCD Enzyme and the Repair of Double-Stranded DNA Breaks. *Microbiol Mol Biol Rev* **72**, 642-671, doi:10.1128/MMBR.00020-08 (2008).

44　Cheng, K., Wilkinson, M., Chaban, Y. & Wigley, D. B. A conformational switch in response to Chi converts RecBCD from phage destruction to DNA repair. *Nat Struct Mol Biol* **27**, 71-77, doi:10.1038/s41594-019-0355-2 (2020).

45　Levy, A. *et al.* CRISPR adaptation biases explain preference for acquisition of foreign DNA. *Nature* **520**, 505-510, doi:10.1038/nature14302 (2015).

46　Kuzmenko, A. *et al.* pAgo-induced DNA interference protects bacteria from invader DNA. *bioRxiv*, doi:10.1101/2020.03.01.971358 (2020).

47　Chen, H.-W., Randle, D. E., Gabbidon, M. & Julin, D. A. Functions of the ATP hydrolysis subunits (RecB and RecD) in the nuclease reactions catalyzed by

the RecBCD enzyme fromEscherichia coli11Edited by G. Smith. *J Mol Biol*
278, 89-104, doi:10.1006/jmbi.1998.1694 (1998).

48   Guegler, C. K. & Laub, M. T. Shutoff of host transcription triggers a toxin-
antitoxin system to cleave phage RNA and abort infection. *Mol Cell* **81**, 2361-
2373.e2369, doi:10.1016/j.molcel.2021.03.027 (2021).

49   Zabrady, K., Zabrady, M., Kolesar, P., Li, A. W. H. & Doherty, A. J. CRISPR-
Associated Primase-Polymerases are implicated in prokaryotic CRISPR-Cas
adaptation. *Nat Commun* **12**, 3690, doi:10.1038/s41467-021-23535-9 (2021).

50   Kutter, E. *et al.* From Host to Phage Metabolism: Hot Tales of Phage T4's
Takeover of E. coli. *Viruses* **10**, 387 (2018).

51   Krüger, D. H. & Schroeder, C. Bacteriophage T3 and bacteriophage T7 virus-
host cell interactions. *Microbiol Rev* **45**, 9-51, doi:10.1128/mr.45.1.9-51.1981
(1981).

52   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*
34, 3094-3100, doi:10.1093/bioinformatics/bty191 (2018).

53   Luong, T., Salabarria, A.-C., Edwards, R. A. & Roach, D. R. Standardized
bacteriophage purification for personalized phage therapy. *Nat Prot* **15**, 2867-
2890, doi:10.1038/s41596-020-0346-0 (2020).

54   Deatherage, D. E. & Barrick, J. E. in *Engineering and Analyzing Multicellular*
Systems: Methods and Protocols (eds Lianhong Sun & Wenying Shou) 165-
188 (Springer New York, 2014).

55   Pabst, M. *et al.* A general approach to explore prokaryotic protein
glycosylation reveals the unique surface layer modulation of an anammox
bacterium. *ISME J* **16**, 346-357, doi:10.1038/s41396-021-01073-y (2022).

56   Sievers, F. *et al.* Fast, scalable generation of high-quality protein multiple
sequence alignments using Clustal Omega. *Mol Sys Biol* **7**, 539,
doi:10.1038/msb.2011.75 (2011).

57   Tumescheit, C., Firth, A. & Brown, K. CIAlign: A highly customisable
command line tool to clean, interpret and visualise multiple sequence
alignments. *PeerJ* **10**, e12983 (2022).

58   Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A
Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood
Phylogenies. *Mol Biol Evol* **32**, 268-274, doi:10.1093/molbev/msu300 (2014).

59   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and
new developments. *Nucleic Acids Res* **47**, W256-W259,
doi:10.1093/nar/gkz239 (2019).
60   Söding, J., Biegert, A. & Lupas, A. N. The HHpred interactive server for
protein homology detection and structure prediction. *Nucleic Acids Res* **33**,
W244-W248, doi:10.1093/nar/gki408 (2005).
61   Hallgren, J. *et al.* DeepTMHMM predicts alpha and beta transmembrane
proteins using deep neural networks. *bioRxiv*, doi:10.1101/2022.04.08.487609
(2022).
62   Gasteiger, E. *et al.* in *The Proteomics Protocols Handbook* (ed John M.
Walker) 571-607 (Humana Press, 2005).
936

a

O55:H7 RM12579
ECOR49
D9
ECOR8
O55:H7 RM12579
ECOR49
ECOR8
D9
ECOR12
ECOR12
SN82F48
SN82F48

KwaA n = 3177

0.5

KwaB n = 3177

0.5

🛡 Previously confirmed anti-phage activity

b

KwaA   KwaB   DUF4868
TM

ECOR8   186aa   276aa
ECOR12   172aa   294aa
ECOR49   195aa   319aa
D9   195aa   306aa
SN82F48   110aa   312aa

c

	Myo				Sipho		Podo	
	T4	NBG1	NBG2	S1-BL21L	T1	Lambda$_{vir}$	T3	T7$_s$
ECOR8		m	m				m	m
ECOR12			m				m	m
ECOR49		m	m				m	
D9								
SN82F48								

Fold defence

<10¹ 10¹ 10² 10³ 10⁴ 10⁵ >10⁵

m = plaque morphology change

d

MOI   3.0   0.03   No phage
Control
Kiwa

ECOR12
T4

OD600

1.2
0.9
0.6
0.3

100   300   500
T (min)

e

MOI = 10⁻⁵ T4

Control   ECOR8

Phage (PFU/mL)

10¹¹
10⁹
10⁷
10⁵

ND ND

0   60   90
Time post infection (min)

MOI = 3   T4

Control   ECOR12

0   60   90
Time post infection (min)

a

KwaA   KwaB

ECOR8

residue (aa)   0 25   74   141   L145A/F147A   S272A

T3   T4

b

KwaA   KwaB

SN82F48

ORF   Complete   Truncated

c

His-SUMO-TEV::
ECOR8 KwaA C-term   ECOR8 KwaB

13S-S   +   pACYC-Duet1