Odor modality is transmitted to cortical brain regions from the olfactory bulb

Michelle F. Craft[1], Andrea K. Barreiro[2], Shree Hari Gautam[3], Woodrow L. Shew[3], Cheng Ly[1,*]

1 Department of Statistical Sciences and Operations Research, Virginia Commonwealth University, Richmond,

Virginia, United States

2 Department of Mathematics, Southern Methodist University, Dallas, Texas, United States

3 Department of Physics, University of Arkansas, Fayetteville, Arkansas, United States

∗cly@vcu.edu ; 1015 Floyd Avenue, Richmond, Virginia 23284

Running Title: OB encodes ortho versus retro odor input

Abstract

Odor perception is the impetus for important animal behaviors, most obviously for feeding but also for mating and communication. There are two predominate modes of odor processing: odors pass through the front of the nose (orthonasal) while inhaling and sniffing, or through the rear (retronasal) during exhalation and while eating. Despite the importance of olfaction for an animal's well-being and specifically that ortho and retro naturally occur, it is unknown how the modality (ortho versus retro) is even transmitted to cortical brain regions, which could significantly affect how odors are processed and perceived. Using multi-electrode array recordings in anesthetized rats, we show that mitral cells in rat olfactory bulb reliably transmit ortho versus retronasal modality with ethyl butyrate, a common food odor. Drug manipulations affecting $GABA_A$ that control synaptic inhibition lead to worse decoding of ortho versus retro, independent of whether overall inhibition increases or decreases, suggesting that the olfactory bulb circuit may naturally favor encoding this important aspect of odors. Detailed data analysis paired with a firing rate model that captures population trends in spiking statistics shows how this circuit can encode odor modality. We have not only demonstrated that ortho/retro information is encoded to downstream brain regions, but models and analyses reveal network dynamics that promotes this encoding.

Key words: retronasal olfaction; decoding accuracy; ethyl butyrate; firing rate model; synaptic inhibition

New and Noteworthy: Whether ortho (sniffing odors) versus retro (exhalation and eating) is encoded from the olfactory bulb to other brain areas is not completely known. Using multi-electrode array recordings in anesthetized rats, we show that the olfactory bulb transmits this information downstream via spikes. Altering inhibition degrades ortho/retro information on average. We use theory and computation to explain our results, which should have implications on cortical processing considering that only food odors occur retronasally.

Introduction

Olfaction is driven by odor molecules entering the nasal cavity, inducing a cascade of action potentials in the nervous system that transmit and process odor information. There are two routes by which odor molecules can enter the nasal cavity: through the nostrils during inhalation and sniffing (**ortho**), or from the throat during exhalation and while eating (**retro**). Orthonasal stimulation is by far the most commonly studied modality, despite the fact that retonasal delivery naturally occurs during eating. Prior imaging studies have shown differences in the activation of the regions of the nasal epithelium with ortho and retro stimulation (Sanganahalli et al., 2020; Gautam and Verhagen, 2012b; Furudono et al., 2013). Small et al. (2005) showed that ortho versus retro information is transmitted to cortical brain regions in humans, but none of these studies clarifies how this occurs. Thus, how ortho versus retro information is transmitted to cortical brain regions, and the core network mechanisms that facilitate encoding this information are largely unknown.

These questions are important for several reasons. Humans are able to discriminate whether food odors are delivered ortho or retronasally without being told which modality and without active sniffing (Frasnelli et al., 2008), but it is unknown whether this information is propagated from the 'bottom up' or if detection is solely from higher brain regions. Imaging studies have shown that at least with food odors, ortho/retro are two distinct modalities, rather than 2 different routes to the same modality, independent of odor intensity (Bender et al., 2009). Also, there is evidence that cortical processing depends on whether odors are delivered ortho- versus retro-nasally (Blankenship et al., 2019). Humans also report perceiving smells differently depending on ortho or retro modality (Hannum et al., 2021; Small et al., 2005). This study provides a more complete picture of these phenomena.

Before odor information reaches the brain, it is processed in the olfactory bulb (**OB**) and relayed to higher brain regions via excitatory mitral cells (**MC**s) (and tufted cells). Thus, the OB is critical for determining whether ortho versus retronasal odors are encoded before being conveyed to the brain for processing and perception. More broadly, OB activity is tied to odor

perception (Mandairon and Linster, 2009). While the role of inhibitory cells in influencing OB processing of ortho versus retro odors is unknown, inhibition is known to play a key role in many other aspects of OB processing. OB inhibition is known to alter activity patterns that represent odors (Wilson and Laurent, 2005), granule cells that provide inhibition reflect changes in odor concentration (Tan et al., 2010), and OB inhibition levels alter odor discrimination dynamics (Abraham et al., 2010). These facts motivated our experiments that consist of *in vivo* rat recordings of MCs in the OB with both decreased and increased levels of $GABA_A$ synaptic inhibition via drug manipulations. The rats were anesthetized and a double tracheotomy was performed to allow air flow during breathing to bypass the nasal cavity, thus allowing precise control of ortho versus retro stimulation.

We show that the mode of olfaction (ortho versus retro) is indeed transmitted to cortical brain regions with a common food odor, ethyl butyrate (**EB**). We find that encoding is generally good and well above chance level for the intact circuit and even with altered levels of inhibition. Thus downstream brain regions may readily have access to ortho versus retronasal odor modality encoded via firing rate from the OB. Importantly, the altered circuits with both increases (via **Mus**cimol application, a $GABA_A$ agonist) and decreases in inhibition (via **Bic**uculline, a $GABA_A$ antagonist) result in worse individual encoding on average. This suggests that the intact OB circuit may be advantageous for encoding ortho versus retro modality.

Our data enables investigation of the OB circuit mechanisms that promote this encoding with a computational model. We show how a simple excitatory-inhibitory (**E-I**) reciprocally coupled model captures the modality dependent differences in population firing rate. Our data shows that the $GABA_A$ (ant-)agonists do not significantly alter the odor-evoked population firing rate with slower longer-lasting retronasal stimulation, in contrast to ortho stimulation. Our model captures these aspects of our data, exploiting the temporal differences, i.e., the slow retro stimulation leads to short-term synaptic depression of the excitatory synapse to diminish the effects of the inhibition. Furthermore, we model trial-to-trial spike rate variability to be largely consistent with our data, so

that the model can encode orth-/retro-like stimuli. We find that average encoding is degraded when inhibition is either increased or decreased, as observed in our data. Given the commonality of reciprocally coupled excitatory-inhibitory circuits, these results may apply outside of the OB, specifically to where stimuli have temporal differences.

Materials and methods

Ethics statement

All procedures were carried out in accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health and approved by University of Arkansas Institutional Animal Care and Use Committee (protocol #14049). Isoflurane and urethane anesthesia were used and urethane overdose was used for euthanasia.

Data and code availability

See `https://github.com/michellecraft64/Modality` for MATLAB code implementing the data analysis and firing rate model.

See `https://www.dx.doi.org/10.6084/m9.figshare.14877780` for raw data collected from the Shew Lab.

Electrophysiological recordings

Data were collected from 8 adult male rats (240-427 g; *Rattus Norvegicus*, Sprague-Dawley outbred, Harlan Laboratories, TX, USA) housed in an environment of controlled humidity (60%) and temperature (23°C) with 12h light-dark cycles. The experiments were performed in the light phase.

Surgical preparations. Anesthesia was induced with isoflurane inhalation and maintained with urethane (1.5 g/kg body weight (**bw**) dissolved in saline, intraperitoneal injection (**ip**)). Dex-

amethasone (2 mg/kg bw, ip) and atropine sulphate (0.4 mg/kg bw, ip) were administered before performing surgical procedures. Throughout surgery and electrophysiological recordings, core body temperature was maintained at 37°C with a thermostatically controlled heating pad. To isolate the effects of olfactory stimulation from breath-related effects, we performed a double tracheotomy surgery as described previously (Gautam and Verhagen, 2012b). A Teflon tube (OD 2.1 mm, upper tracheotomy tube) was inserted 10 mm into the nasopharynx through the rostral end of the tracheal cut. Another Teflon tube (OD 2.3 mm, lower tracheotomy tube) was inserted into the caudal end of the tracheal cut to allow breathing, with the breath bypassing the nasal cavity. Both tubes were fixed and sealed to the tissues using surgical thread. Local anesthetic (2% Lidocaine) was applied at all pressure points and incisions. Subsequently, a craniotomy was performed on the dorsal surface of the skull over the right olfactory bulb (2 mm × 2 mm, centered 8.5 mm rostral to bregma and 1.5 mm lateral from midline).

Olfactory stimulation. A Teflon tube was inserted into the right nostril to deliver orthonasal stimuli, and the left nostril was sealed by suturing. The upper tracheotomy tube inserted into the nasopharynx was used to deliver odor stimuli retronasally. Odorized air was delivered for 1 second in duration at 1 minute intervals, with a flow rate of 250 ml/min and 1% of saturated vapor. Two odors were used: ethyl butyrate (**EB**, a food odor) and 1-Hexanol (**Hex**, a non-food odor). Here we limited our analysis to EB trials because food odors are perceived ortho- and retro-nasally (Small et al., 2005); non-food odors do not naturally occur retronasally.

Electrophysiology. A 32-channel microelectrode array (MEA, A4x2tet, NeuroNexus, MI, USA) was inserted 400 μm deep from dorsal surface of OB targeting tufted and mitral cell populations. The MEA probe consisted of 4 shanks (diameter: 15 μm, inter-shank spacing: 200 μm), each with eight iridium recording sites arranged in two tetrode groups near the shank tip (inter-tetrode spacing: 150 μm, within tetrode spacing 25 μm). Voltage was measured with respect to an AgCl ground pellet placed in the saline-soaked gel foams covering the exposed brain surface around the inserted MEAs. Voltages were digitized with 30 kHz sample rate (Cereplex + Cerebus, Blackrock

Microsystems, UT, USA). Recordings were band-pass filtered between 300 and 3000 Hz and semi-

automatic spike sorting was performed using Klustakwik software, which is well suited to the type

of electrode arrays used here (Rossant et al., 2016).

Pharmacology. We topically applied GABA antagonists (bicuculline 20 μM) or agonists (mus-

cimol 20 μM) dissolved in artificial cerebrospinal fluid (**ACSF**). Gel foam pieces soaked in the

ACSF+drug solutions were placed on the surface of OB surrounding the recording electrodes. The

gel foams were placed 10 minutes before beginning a recording.

Data analysis

Data was collected *in vivo* from the mitral cell (**MC**) layer in the olfactory bulb (**OB**) of multiple

anesthetised rats using a multi-electrode array recording. The data consisted of spike recordings

of multiple MC spike responses to EB and Hex presented by the two routes of stimulation, or-

thonasally and retronasally, for a total of 20 trials, 10 for each ortho and retro, with the order

alternating (10 ortho before 10 retro, then 10 retro before 10 ortho trials). The spike counts were

calculated using 100 ms overlapping time windows. Three separate drug preparations were used

in order to analyze inhibitory effects on MC spiking responses: no drug (control), Bicuculline

(GABA$_A$ antagonist, i.e., decreasing inhibition), and Muscimol (GABA$_A$ agonist, i.e., increasing

inhibition). Data was obtained from 8 viable rats in total. After spike sorting, we identified and

eliminated units (cells) that had firing rates below 0.008 Hz or more than 49 Hz, calculated over

the entire recording duration. We took this conservative approach to eliminate potential artifacts

yielding units that had unrealistically high or low firing; see Table 2 for the reported number of

total rats and individual cells for each drug preparation. Note that some cells around the odor onset

($t = 0$) will have much larger (> 49 Hz) or smaller firing rates than our criteria to eliminate units,

but the entire recording duration is much longer with approximately a minute between trials.

Defining single cell decoding

We used a form of linear discriminant analysis (**LDA**) for decoding accuracy. Decoding accuracy is defined as the fraction of trials correctly classified by a threshold that maximizes decoding accuracy. Letting $s_k(t)$ be the spike train of the k^{th} trial ($k = 1, 2, ..., N$) that at each time t is either $= 0$ (no spike) or $= \delta(t - t_{j,k})$, where $t_{j,k}$ denotes the j^{th} spike on the k^{th} trial . The entries of the spiking rate $\vec{x}$ are:

$$x_k = \frac{1}{T_{ev}} \int_0^{T_{ev}} s_k(t)\, dt \tag{1}$$

where again $t = 0$ is the time the odor is presented. There are $N = 20$ trials total (10 ortho, 10 retro) for each individual cell. We consider all possible threshold values θ_k:

$$\begin{cases} \theta_k = x_k - \epsilon, & \text{if } x_k = \max(\vec{x}) \\ \theta_k = x_k + \epsilon, & \text{else} \end{cases}$$

and select an optimal threshold θ^* and classification scheme (decode stimulus as ortho if x_k is less than or greater than θ^*) that results in the best decoding accuracy for each cell. Here $\epsilon \approx 10^{-16}$ is machine precision and is there to avoid trivial threshold (when $x_k = \max(\vec{x})$ without ϵ the decoding accuracy is 50%) and so that $x_k \neq \theta_k$ so each is categorized as ortho or retro. Since decoding accuracy is the percent of correctly classified trials (ortho or retro) and $N = 20$, it varies between 0.5 (chance) and up to 1 in increments of 0.05.

To systematically compare differences in decoding accuracy, we select a fixed time window for all drug preparations and modalities based on two statistical tests (two sample $t-$test assuming unequal variances and Wilcoxon rank sum test) and their $p-$values. We evaluated decoding accuracy from $T_{ev} = 100$ ms to $T_{ev} = 1$ s in 100 ms increments in the evoked state for each drug preparation. We analyzed differences in mean between no drug preparation and Bicuculline, as well as no drug preparation and Muscimol. This resulted in 4 different $p-$values for each time window; we found that $T_{ev} = 900$ ms had the most significant $p-$values for both combinations of no drug/Bicuculline

175 and no drug/Muscimol, see Fig A2A,C. Figure A2B,D shows the population decoding accuracies

176 (mean across the population and one standard deviation in shaded region).

Statistical significance

We use three tests to assess whether a given effect is statistical significant: 1) two-sample $t-$test

with unequal variance, 2) Wilcoxon rank sum test, 3) one-way ANOVA. Each of these tests are

used to rule out the null hypothesis that the population means are the same in two different cat-

egories (Cohen, 2013). Each test is different and has various underlying assumptions: 1) normal

samples, 2) non-parametric test assuming independent groups and equal variance, 3) including

sample variance to assess differences in means assuming equal variance and normally distributed

residuals. We consider three tests to provide a more complete picture of the results and to demon-

strate robustness, or lack thereof, for a given effect.

In addition to the $p-$values that indicate the probability of the null hypothesis holding (no ef-

fect, same population means), we also report the effect sizes. We use the conventions outlined in

Cohen (2013); Tomczak and Tomczak (2014) that provide formulas for effect sizes, and a qualita-

tive 'rule of thumb' for when effect sizes are: **small, medium** or **large**. For 2 groups of length n_1

and n_2 respectively, the effect sizes we use:

$$\text{For } t - \text{test, use:} \quad \frac{|\mu_1 - \mu_2|}{\sigma_p}, \text{ Cohen's } d, \text{ where} \tag{2}$$

$$\sigma_p = \sqrt{\frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2}}. \tag{3}$$

$$\text{For Wilcoxon rank sum test, use:} \quad \frac{|\zeta|}{\sqrt{n_1 + n_2}}, \text{ where } \zeta \text{ is the z-score.} \tag{4}$$

$$\text{For one-way ANOVA, use:} \quad \frac{SS_{group}}{SS_{total}}, \text{ ratio of particular sum of squares.} \tag{5}$$

191 The variables μ_j and S_j^2 denote the sample mean and variances of the j^{th} group. Effect sizes

192 for $t-$test and Wilcoxon rank sum test with values: $(0, 0.2]$ are considered **small**, $(0.2, 0.5]$ are

¹⁹³ **medium**, $(0.5, 0.8]$ and above are **large** (Cohen, 2013; Tomczak and Tomczak, 2014). Effect

¹⁹⁴ sizes for one-way ANOVA with values $(0, 0.01]$ are considered **small**, $(0.01, 0.06]$ are **medium**,

¹⁹⁵ $(0.06, 0.14]$ and above are **large** (Cohen, 2013).

Population decoding

¹⁹⁷ The two methods are applied to each population of simultaneously recorded MCs.

¹⁹⁸ <u>PCA+LDA</u>: For each recording, Principle Components Analysis (**PCA**) was applied to the

¹⁹⁹ concatenated matrix X of size $20 \times M$, where M is the number of MCs (Fig 6A), rows 1–10

²⁰⁰ correspond to the spiking rate with ortho and rows 11–20 correspond to retro. We only use the

²⁰¹ first two principal components and apply Linear Discriminate Analysis (**LDA**) to find a line in the

²⁰² 2-dimensional plane that maximizes classification accuracy (Cunningham and Yu, 2014). We used

²⁰³ built-in MATLAB routines `pca`, `fitcdiscr`, `predict` (see GitHub for code).

²⁰⁴ <u>SVM</u>: We use support vector machine with supervised 'learning' to classify data into ortho or

²⁰⁵ retro. We use a nonlinear classification method with Gaussian kernels, and Bayesian optimization

²⁰⁶ to find the best kernel scale and box constraints for the SVM (we insured that for all recordings and

²⁰⁷ drug preparations, the objective was minimized within the allowable function evaluations). After

²⁰⁸ the SVM is fit to the data, the decoding accuracy is the correct classification rate obtained from

²⁰⁹ the MATLAB function `kfoldLoss` using 10-fold cross-validation. We used built-in MATLAB

²¹⁰ routines (see GitHub for code).

Firing rate model

²¹² We use a space-clamped Wilson-Cowan rate model of coupled E-I cells: E (MC, excitatory) and I

²¹³ (inhibitory from PGC, GC, etc). The models of the respective cell firing rate, $A_j(t)$ $j \in (M, P)$,

²¹⁴ are represented by the following ordinary differential equations:

$$\tau_E \frac{dA_E}{dt} = -A_E(t) + F\Big(w_{orn,E}I_{orn}(t) - w_{EI}S_I(t)\Big) \tag{6}$$

$$\tau_I \frac{dA_I}{dt} = -A_I(t) + F\Big(w_{orn,I}I_{orn}(t) + w_{IE}S_E(t)\Big) \tag{7}$$

where the synaptic term, S_j $j \in \{E, I\}$, is defined by synapses with rise and decay time scales (same τ_{rise}, τ_{decay} for E/I) as follows:

$$\tau_{\text{decay}} \frac{dS_j(t)}{dt} = -S_j(t) + X_j(t) \tag{8}$$

$$\tau_{\text{rise}} \frac{dX_j(t)}{dt} = -X_j(t) + \tau_{\text{rise}} A_j(t). \tag{9}$$

In Eqs 6 and 7, $I_j(t)$ is the sum of external currents (e.g., stimulus input and other currents) that varies over time to account for stimulus input, $w_{orn,j}$ are the coupling strengths of olfactory receptor neuron (**ORN**) input to j, w_{jk} are the coupling strengths from cell k to j, and the transfer function F is a threshold linear function:

$$F(x) = \max(x, 0). \tag{10}$$

Table 1: **Parameter values in firing rate model (Eqs (6)–(12)). Last 2 rows describe the parameter ρ in simulated trial variability using negative binomial distribution.**

Parameter	Value		
τ_E	10 ms		
τ_I	5.5 ms		
τ_{decay}	10 ms		
τ_{rise}	2 ms		
τ_w	100 ms		
$w_{orn,E}$	1		
$w_{orn,I}$	1.1875 (ortho), 0.3 (retro)		
	Drug Preparation:		
	No Drug	**Bic**	**Mus**
w_{EI}	0.2	0.15	0.25
ρ for ortho	0.32	0.32	0.32
ρ for retro	0.32	0.53	0.052

Without synaptic depression, we fix $w_{IE} = 1$. With synaptic depression, $w_{IE}(t)$ is governed by:

$$\tau_w \frac{dw_{IE}}{dt} = -w_{IE} + F_d\Big(A_E(t)\Big) \tag{11}$$

$$F_d(x) = \frac{1}{1 + e^{(x-0.8)}}. \tag{12}$$

with initial value $w_{IE}(0)$ set to its steady-state value at $t = -0.5\,\text{s}$ in Eqs(6)–(12).

The trial-to-trial variability of spiking rate is modeled as a negative binomial random variable:

$$\mu_{NB} = \frac{r(1-\rho)}{\rho} \tag{13}$$

Given output from the Wilson-Cowan model $A_E(t)$, we set:

$$\mu_{NB} = \frac{1}{T_{ev}} \int_0^{T_{ev}} A_E(t) dt \tag{14}$$

by manually determining the ρ parameter in the negative binomial distribution (see Table 1) for each drug preparation, and setting the parameter $r = \mu_{NB}\rho/(1-\rho)$.

Results

We collected spike data with multi-electrode array recordings of urethane anesthetized rats from the OB mitral cell layer, where each cell (MC) was subject to the same odor delivered with ortho and retro stimulation (Fig 1Ai), repeated for 10 trials for each modality. In addition, two drugs were applied to alter the circuit: a GABA$_A$ antagonist (**Bic**uculline) and a GABA$_A$ agonist (**Mus**cimol), see Table 2. For reference, Fig 1Aii shows the population-averaged firing rate (trial-average and population-averaged firing rate) by modality for a given drug preparation, with $t = 0$ denoting time of odor presentation, held for 1 s. This is commonly referred to as the peri-stimulus time histogram

 (**PSTH**).

Table 2: **Number of rats and respective individual cells for each drug preparation subject to the food odor ethyl butyrate (EB).**

	No Drug	Bicuculline	Muscimol
Rats	8	4	3
Cells	913	413	419

Individual MCs encode ortho versus retro

To remain agnostic to how higher brain regions decode population activity, we predominately consider individual MCs odor modality encoding, a logical first step (Rolls and Treves, 2011). We use odor-evoked firing rate in a given trial as the source of encoding. Specifically, the firing rate in the k^{th} trial, x_k, is the sum of the evoked spike counts normalized by time window $T_{ev} = 0.9$ s:

$$x_k = \frac{1}{T_{ev}}\Big(\# \text{ Evoked spikes in } T_{ev}\Big) \tag{15}$$

for $k = 1, 2, ..., 20$, same as Eq (1) in **Materials and methods**. The time window T_{ev} is systematically chosen from a range of possible values, see Fig A2A,C in Appendix and **Materials and methods** for details. To assess whether MCs might prefer one modality over the other, we compare the trial-averaged firing rate for ortho and retro as a measure of selectivity (Fig 1Aiii). Most MCs 'prefer' or spike more with retro than ortho ($\approx 63\,\%$), and the proportions do not vary much across drug preparations (but see Appendix and Table A1 for analysis of Off-Response cells).

We define decoding accuracy as the fraction of trials correctly classified by a threshold that maximizes decoding accuracy; this is a measure of information accessible by an ideal observer from each MC individually (Fig 1B). We only consider EB, a common food odor, because typically only food odors are perceived retronasally (Small et al., 2005). It has been shown that non-food odors artificially delivered retronasally can be perceived as well as food odors (Small et al., 2005). However, numerous studies have shown that human perception of non-food odors deliv-

Figure 1: **Individual mitral cells encode modality. Ai)** Experiment setup to test whether OB MCs encode ortho versus retro stimulation, with example raster plot for 1 MC. **Aii)** Population-averaged firing rate (trial- and population-averaged), i.e., **PSTH** for each modality and drug preparation; $t = 0$ denotes when ethyl butyrate (food odor) is presented. **Aiii)** The proportion of population that fire more for ortho (or retro), i.e., ortho selective $\Leftrightarrow$ higher trial averaged firing rate for ortho than retro. **B)** Schematic of how decoding accuracies are calculated for each MC based on trial firing rates: $\mu_{R/O}$ are the trial-averaged firing rates. Shaded regions correspond to incorrect decoding, unshaded regions to correct decoding. **C)** Distribution of decoding accuracies of MC with 3 different preparations: intact no drug (gray), less inhibition via Bic (green) and more inhibition via Mus (purple). Mean decoding accuracy for no drug is 0.724; with Bic and Mus the means are: 0.680, 0.690 (resp.). Differences are all statistically significant ($\alpha = 0.01$ with two-sample $t-$test assuming unequal var, Wilcoxon rank sum test, and one-way ANOVA) with mostly 'medium' effect size, see Table 3. Windows were determine systematically to maximize $p-$values in testing average decoding accuracy, see Fig A2A,C and **Materials and methods**.

256 ered retronasally is degraded compared to food odors (Hannum et al., 2021; Bender et al., 2009;

257 Frasnelli et al., 2008).

258   The OB circuit encodes ortho versus retro very well. Figure 1C shows the spiking activity of

₂₅₉ individual MCs encode (measured by decoding accuracy) ortho versus retro for the vast majority

₂₆₀ of cells. MCs have a wide range of decoding accuracies varying between 0.5 (chance) and 1

₂₆₁ (perfect decoding). With no drugs, the mean decoding accuracy (0.724) is better than both Bic and

₂₆₂ Mus, which have lower average decoding accuracies of 0.680, 0.690, respectively. These averages

₂₆₃ are numerically small in difference, but note that the range of decoding accuracies is only 0.5.

₂₆₄ Furthermore, the differences in the mean decoding accuracy are all statistically significant ($\alpha =$

₂₆₅ 0.01) using three tests: two-sample $t-$test. Table 3 demonstrates that the $p-$values are all small,

₂₆₆ and that the effect size is generally 'medium', with some 'small' (see **Statistical significance**

₂₆₇ section in **Materials and methods**).

Table 3: **Population Decoding Accuracy: significance measured by $p-$values of average decoding accuracies over MC, shown in Figure 1C.** Using 3 tests: two-sample $t-$test assuming unequal variance, Wilcoxon rank sum test, and one-way ANOVA. Bottom table shows effect size (see **Materials and methods**).

Relationship / $p-$value	$t-$test	Rank Sum	ANOVA
Bic < ND	3.6×10^{-10}	4.9×10^{-6}	1.8×10^{-8}
Mus < ND	1.2×10^{-4}	1.8×10^{-6}	7.1×10^{-5}

Relationship / Effect Size	$t-$test	Rank Sum	ANOVA
Bic < ND	0.336 (medium)	0.126 (small)	0.0237 (medium)
Mus < ND	0.235 (medium)	0.131 (small)	0.0118 (medium)

₂₆₈   Figure 2 shows a more complete picture of the entire (pooled) population's spike rate statistics

₂₆₉ across drug preparations and modality, beyond the PSTH in Figure 1Aii. There are very strong

₂₇₀ positive correlations between the mean firing rates and standard deviations (across trials) in all

₂₇₁ drug preparations and both modalities (see Figure A3, and Table A2), consistent with results in

₂₇₂ Ly et al. (2021) reported for MC in OB. Notably, the *opposite* relationship is observed in the vast

₂₇₃ majority of cortical neurons (Churchland et al., 2010).

Figure 2: **Detailed spiking rate statistics of pooled MC populations. A**) Colormap showing trial-averaged spiking rate of individual MC (individual PSTH), each of the 6 panels sorted by time-averaged firing rate (4 s total, 2 s in spontaneous and evoked, but only showing 1 s of spontaneous for brevity). **B**) Colormap showing trial variability of spiking rate (measured by standard deviation across trials) of individual MC, using the same six corresponding indices from sorting in A). To see relative differences across MC, the color scales were effectively thresholded so that the larger values did not obscure the entire population. See Fig A3 for more on firing rate and trial-to-trial variability relationship.

Drug effects on average population spiking

To better understand the dynamics of MC decoding accuracy, we consider how $GABA_A$ (ant-)agonists effect population activity. Although population activity is a coarse measure that neglects

individual cell heterogeneity, it can still be insightful for determining average trends. A charac-
teristic of an individual MC that might be indicative of encoding fidelity is the distance between
the scaled trial-averaged firing rate of ortho and retro stimulation $|\frac{\mu_R}{\sigma_R} - \frac{\mu_O}{\sigma_O}|$ (Fig 1B). We expect
larger distances to coincide with better decoding accuracy than smaller distances, if all other fac-
tors are equal. Fig 3A shows moderate to strong correlations (Pearson's and Spearman's rank)
between $|\frac{\mu_R}{\sigma_R} - \frac{\mu_O}{\sigma_O}|$ and decoding accuracies for all drug preparations. Although other factors af-
fect decoding accuracy since the spread in the horizontal direction is large for a given decoding
accuracy value and the correlations are not 100%. Nevertheless, this motivates considering the
population-averaged spiking activity as a means to better understand the dynamics of decoding
accuracy.

The affects of the GABA$_A$ (ant-)agonists on population firing rate depend on the odor modal-
ity; only ortho evoked firing rates show statistically significant differences in the mean (over MC
population) firing rate (Fig 3B). As expected, Mus (increased inhibition) results in lower firing
rates and Bic (less inhibition) results in higher firing rates compared to no drug, especially in the
spontaneous state. However, the evoked firing rates are **only** significantly different ($\alpha = 0.01$)
across drug preparations with ortho stimulation, and **not** significantly different with retro stimu-
lation. This holds for nearly all time points (see * in $p-$value plots of two sample $t-$test with
unequal variance). This effect is clear in the right panels of Fig 3Bi and Bii. For completeness, the
Appendix shows the PSTH in Figs 1Aii and 3B for a longer timeframe (Fig A1A,B).

If we assume these population trends (Fig 3B) hold individually for MCs (i.e., drugs only effect
spiking rate in ortho but not retro), then we might expect decoding accuracies to change in specific
ways: if ortho firing rate is less than retro with no drug, then Bic should increase firing rate (shift
histograms to the right) causing more overlap and lower decoding accuracy. For exposition pur-
poses, these results are detailed in **Appendix** and Fig A4 where predictions on decoding changes
based on individual MC's relationship of ortho versus retro firing rate are partially verified.

Figure 3: **Population activity: retronasal response is more robust to changes in OB inhibition.**
A) Distance between scaled trial-averaged mean $|\frac{\mu_R}{\sigma_R} - \frac{\mu_O}{\sigma_O}|$ is correlated with decoding accuracy. The 'Correlation' is Pearson's correlation coefficient, and 'Rank Correlation' is Spearmen's. **B**) Drug effects on evoked population PSTH are different with statistically significant effects only with ortho stimulation **(Bi)** but not retro **(Bii)**. The right-panels with 3 subplots show the time series of $p-$values from two-sample $t-$test assuming unequal variances, comparing pairs of drug preparations.

Drug effects on trial-to-trial variability

The trial-to-trial variability plays a key role in decoding accuracy differences. We analyze trial variance for each MC to test whether the population average is different across drug preparations. We find with Mus application, the average trial variability is smallest, followed by Bic, then no drug (i.e., Mus < Bic < ND), for both ortho and retro. The significance measured by $p-$values and effect size are shown in Tables 4 and 5 for ortho and retro trials, respectively, using the same

three statistical tests as before. For ortho stimulation, Mus < ND is significant for two of the three tests ($\alpha = 0.01$), while Bic < ND and Mus < Bic are only significant with Wilcoxon rank sum test. For retro stimulation, again only the Wilcoxon rank sum test shows this relationship is significant (all with $\alpha = 0.01$ except for Bic < ND for retro where $p = 0.038$). Overall, the Bic < ND relationship is not as strong as the others, and the trend Mus < Bic < ND is stronger with ortho stimulation. Note that the effect size is considered 'small' in all cases, see Tables 4 and 5.

Table 4: **Ortho stimulation only: significance ($p-$values) of average (over MC) trial variance differences between drugs with various statistical tests.** Using same 3 tests as before; **ND**='no drug'; bottom table shows effect size.

Relationship / $p-$value	$t-$test	Rank Sum	ANOVA
Mus < ND	1.8×10^{-3}	7.7×10^{-8}	0.021
Mus < Bic	0.055	2.4×10^{-13}	0.053
Bic < ND	0.82	5.0×10^{-4}	0.82

Relationship / Effect Size	$t-$test	Rank Sum	ANOVA
Mus < ND	0.137 (small)	0.147 (small)	0.00402 (small)
Mus < Bic	0.134 (small)	0.254 (medium)	0.00450 (small)
Bic < ND	0.0137 (small)	0.0955 (small)	4.06×10^{-5} (small)

Table 5: **Retro stimulation only: significance ($p-$values) of average (over MC) trial variance differences between drugs with various statistical tests.** Using same 3 tests as before; **ND**='no drug'; bottom table shows effect size.

Relationship / $p-$value	$t-$test	Rank Sum	ANOVA
Mus < ND	0.039	1.9×10^{-4}	0.11
Mus < Bic	0.57	7.6×10^{-7}	0.57
Bic < ND	0.26	0.038	0.30

Relationship / Effect Size	$t-$test	Rank Sum	ANOVA
Mus < ND	0.0953 (small)	0.102 (small)	0.00193 (small)
Mus < Bic	0.0399 (small)	0.171 (small)	3.98×10^{-4} (small)
Bic < ND	0.0610 (small)	0.0569 (small)	7.97×10^{-4} (small)

Within all of the 3 drug preparations, the average trial variances for ortho compared to retro are statistically indistinguishable, consistent with Craft et al. (2021) who did similar analyses but on only 1 rat instead of 8 rats.

Model connects network dynamics and decoding accuracy

To reveal the neural network dynamics that explain our experimental data results, we use a simple firing rate model. Our rich data set with 3 total drug preparations enables a model framework that is highly constrained by data (Xiao et al., 2021; Ly et al., 2021; Barreiro et al., 2017). In addition, we incorporate known differences in temporal dynamics of ORN inputs to MCs with ortho versus retro stimulation (Sanganahalli et al., 2020; Gautam and Verhagen, 2012b; Furudono et al., 2013; Scott et al., 2007; Carey et al., 2009). Ortho stimulation results in fast increase and fast decrease of ORN inputs to MCs while retro input results in slower increase and slower decrease ORN inputs than ortho (Fig 4B). Fig 2c of Furudono et al. (2013) suggests that the amplitude of ORN inputs with ortho are larger than retro (Fig 4B). We develop a principled model that is structurally the same with both odor modalities but effectively have different dynamics stemming from different ORN inputs (Craft et al., 2021); note that paper had a different purpose with data only from intact circuit, i.e., it did not include drug manipulations. We decided not to use the biophysical model in Craft et al. (2021) because: i) individual MC spiking statistics are modulated by net inhibitory inputs, distinguishing interneurons is a complication more relevant for joint statistics of MC populations; ii) resolving trial-average spike statistics is more computationally expensive, making parameter tuning to capture *both* firing rate differences and average decoding accuracies across 3 drug preparations impractical; iii) calculating many decoding accuracies where each parameter set has an optimal threshold that has to be determined exhaustively (see Fig 5B,C below) is a necessary additional component here (not in Craft et al. (2021)) that is also computationally expensive.

We first set out to implement an OB model that accounts for the large changes in population PSTH with $GABA_A$ (ant-)agonists with ortho stimulation only, and small (or no significant) changes in population PSTH with retro (Fig 3B). We use a reciprocally coupled 2 cell E-I network (Fig 4A), consistent with OB networks that have many such E-I pairs (granule and periglomerular cells are reciprocally connected to MCs with fast dendrodendritic synapses (Rall et al., 1966)). We

model the drug effects Bic (Mus) by decreasing (increasing) the coupling strength from I to E cell by 25% of the baseline value ($w_{EI} = 0.2$ with no drugs, arbitrary units, see Eqs (6)–(7) and Table 1).

The data shows that affects of drugs altering inhibition are largely absent with slower longer-lasting retro stimulation (Fig 3B), thus finding a neural network principle that effectively removes the inhibitory cell input is imperative, and might be indicative of the core mechanisms. We hypothesize that short-term plasticity on the excitatory synapse ($E \rightarrow I$) could be a factor in explaining the differences of population firing between ortho versus retro with GABA$_A$ (ant-)agonists because of known temporal differences in ORN inputs. There is indeed evidence of plasticity in the OB circuit (Cang and Isaacson, 2003), specifically for short-term synaptic depression in the main OB (Wang et al., 2012) and from MC to granule cells (Dietz and Murthy, 2005). Since the relative differences in firing rate with ortho are already large without plasticity (1st column in Table 6), honing in on decreasing the firing rate differences in retro (3rd column in Table 6) would make the model more consistent with the data (Fig 3B). We use short-term synaptic depression as a means to nonlinearly change the retro firing rates to diminish the differences because retro has longer lasting stimuli. Although Dietz and Murthy (2005) showed that $E \rightarrow I$ synapses in OB have both facilitation and depression with facilitation appearing more prominent, we found that short-term facilitation made our model much less consistent with our data.

Table 6: **Relative differences of firing rate model between no drug and inhibitory drugs in both modalities.** All percent differences (top table) are relative to no drug: $\left| \int_0^{T_{ev}} A_E^{\text{Drug}}(t)\, dt - \int_0^{T_{ev}} A_E^{\text{No Drug}}(t)\, dt \right| \Big/ \int_0^{T_{ev}} A_E^{\text{No Drug}}(t)\, dt$. Bottom table shows average firing rate $\frac{1}{T_{ev}} \int_0^{T_{ev}} A_E(t)\, dt$ (Hz) , i.e., the components of the relative differences.

Relative Differences (%)	Ortho		Retro	
	Without Depression	With Depression	Without Depression	With Depression
No Drug / Bicuculline	37.87	38.16	16.08	6.04
No Drug / Muscimol	30.31	45.09	12.90	6.25

$\frac{1}{T_{ev}} \int_0^{T_{ev}} A_E(t)\, dt$	Ortho		Retro	
	Without Depression	With Depression	Without Depression	With Depression
No Drug	6.11	8.58	9.64	15.65
Bicuculline	8.43	11.86	11.19	16.59
Muscimol	4.26	4.71	8.40	14.67

Figure 4: **Firing rate model captures effects of drugs, with better alignment with data when including synaptic depression. A)** Schematic of reciprocally coupled E-I pair. **B)** Fixed ORN input profiles for ortho and retro based on experiments (Sanganahalli et al., 2020; Gautam and Verhagen, 2012b; Furudono et al., 2013; Scott et al., 2007; Carey et al., 2009) (see main text). **C)** Wilson-Cowan firing rate model (Eqs (6)–(12)) output for ortho (top row) and retro (bottom row) stimulation, compared when synaptic depression is included (solid lines) or not (dashed lines). Left column: E cell (MC) output; center column: I cell output; right column: synaptic strengths (W_{IE}). Data from experiments also shown for comparison (dot-dashed lines) in left column.

Figure 4C shows comparisons of various model outputs (Eqs (6)–(12)). The model indeed follows the firing rate modulation with $GABA_A$ (ant-)agonists with both odor modalities, whether or

not synaptic depression is included. Specifically, ortho MC firing rates (top left in Fig 4) with different drug preparations are well-separated, while the retro MC firing rates are pretty close together (bottom left in Fig 4). The effects of synaptic depression significantly narrows the differences in retro population PSTH among all 3 drug applications, while with ortho the differences across drug preps remain large (see Table 6 for relative differences). Taken together, synaptic depression is crucial for capturing the data (Fig 3B). With retro stimuli and all 3 drug preparations, the $E \rightarrow I$ synaptic connections are depressed for longer so that the I-cell fires less, giving a boost to E-cell firing. In both modalities, there is overall higher E (MC) firing with short-term synaptic depression because of less inhibition (for all 3 drug preparations). Synaptic depression effectively diminishes the reciprocal loop so that the inhibitory drug effects are weakened. In principle Mus should have the opposite effect as Bic, but: the coupling strength w_{EI} has moderate variation: 0.15 (Bic), 0. 2 (ND), 0.25 (Mus). Also, we modeled short-term depression so that inhibition is almost completely turned off with Retro, resulting in MC being effectively decoupled from the I cell. Our model does not have realistic levels of synaptic depression, Cang and Isaacson (2003) report moderate levels of synaptic depression. And anesthetized rats have less active granule cells (Cang and Isaacson, 2003) than awake, so other interneurons would have to be active for our model to be more consistent.

With the neural dynamics captured in our model, we turn to connecting the model to decoding accuracy. The firing rate model does not explicitly have trial-to-trial variability, an important piece for decoding accuracy. A natural choice would be to assume Poisson statistics for the spike counts, but this single parameter distribution results in trial variance ordered by the firing rate: Mus<ND<Bic, which is inconsistent with our data and has perfect (100%) decoding accuracy with these firing rate models. Capturing the trial variability trends in our data seems to require tunable parameters that are not directly tied to the (mean) firing rate. We use a simple two parameter model that makes simulation easy and transparent: the negative binomial distribution. Note that there may be other models of trial variability that are more realistic relation to MCs, but that is

unlikely to be as pragmatic for capturing firing rate <u>and</u> decoding accuracy trends. Specifically, for each drug (ND/Bic/Mus) and each input (ortho/retro), we fit a negative binomial distribution to the average decoding accuracy. The two distribution parameters were chosen so that: 1) the mean of distribution coincided with the model spiking rate $\frac{1}{T_{ev}} \int_0^{T_{ev}} A_E(t)\,dt$ (solid curves in Fig 4C, left column), and 2) the decoding accuracy matches the averages in the experimental data.

The negative binomial distribution has distribution: $\frac{\Gamma(x+r)}{x!\Gamma(r)}\rho^r(1-\rho)^x$ with parameters: $\rho \in (0,1)$ and $r > 0$ that are not directly related to the physiology. We manually (by trial and error) determine the parameter ρ for each modality and drug preparation, which are all the same ρ except for retro with Bic and Mus (see Table 1), after which the r parameter is determined by:

$$ r = \frac{\rho}{1-\rho}\left(\frac{1}{T_{ev}} \int_0^{T_{ev}} A_E(t)\,dt\right) \tag{16} $$

since the mean of the negative binomial is $\frac{r(1-\rho)}{\rho}$. We chose ρ for each regime so that the resulting decoding accuracy is close to the averages from the experiments: 0.724 (no drug), 0.680 (Bic), and 0.690 (Mus).

The results are summarized in Fig 5A. We simulated 50,000 trials and found the optimal threshold to get decoding accuracies that match the data.

We check whether our model results are consistent with the trial-to-trial variability of our data by calculating the variances of firing rate over trials, which is simply $\frac{r(1-\rho)}{\rho^2}$ for the negative binomial distribution. The values listed in Table 7 show that trial variance is consistent with the data: Mus < ND < Bic, except for retro Mus firing that has the largest variance. Recall in our data that the trends for variance of firing rate was weakest with retro stimulation with larger $p-$values (Table 5).

A test of our model is to determine whether the specified parameter values for ρ results in ND generally having better decoding accuracy than Bic and Mus. We start with the base model of spiking rate: $\frac{1}{T_{ev}} \int_0^{T_{ev}} A_E(t)\,dt$ (solid curves in Fig 4C, left column) and perturb these values to

Figure 5: **Model decoding accuracies with simulated trial variability. A)**: Histograms of simulated firing rate assuming negative binomial trial-to-trial variability with the mean set to: $\frac{1}{T_{ev}} \int_0^{T_{ev}} A_E(t)\, dt$ (see Eqs (6)–(12), and solid curves in left column of Fig 4 for $A_E(t)$). **B)–C)** Systematically varying the mean to show that the specified ρ parameter (Table 1) in negative binomial distribution generally results in no drug having best decoding accuracy (see main text for details). **C)** Same model results in **B)** but showing relationship with distance between (scaled) means; red circles correspond to regimes in **A)**.

Table 7: **Simulated trial variance of the model:** $\frac{r(1-\rho)}{\rho^2} = \frac{\mu}{\rho}$.

	Muscimol	No Drug	Bicuculline
Ortho	14.73	26.83	37.06
Retro	282.12	48.90	31.31

412 get different means: $\mu^*_{O/R} = \frac{1}{T_{ev}} \int_0^{T_{ev}} A_E(t)\, dt \pm M_{O/R}$, where $M_O \in \{-2, \ldots, 4\}$ Hz on 25

413 equally spaced points and $M_R \in \{-6, \ldots, 9\}$ Hz on 31 equally spaced points. Once a mean is

414 specified, we use the same ρ (Table 1) and set r^* to correspond to the mean μ^*:

$$r^* = \frac{\rho}{1-\rho}\mu^*; \tag{17}$$

we simulate 50,000 trials to get a decoding accuracy value for each set of parameters. We get

775 decoding accuracy values corresponding to the different (μ^*_O, μ^*_R) combinations for each drug

preparation (Fig 5B). Importantly, our model does not soundly capture many details of realistic

MC heterogeneities in our data (Fig 1C and Fig 5B have different shapes), but Figure 5B at least

matches the result that ND on balance has better decoding accuracy than Bic and Mus. Finally,

Figure 5C shows how decoding accuracy varies with scaled distance $|\frac{\mu^*_R}{\sigma_R} - \frac{\mu^*_O}{\sigma_O}|$. The decoding

accuracies can vary for a given drug preparation, but the ND and Bic have strong positive correla-

tions (like the data, see Fig 3A), but Mus has strong negative correlations that is inconsistent with

the data.

Population decoding results

Thus far we have focused on decoding odor modality using the firing rate of individual MCs.

While this approach avoids making assumptions about how higher level cortical regions might use

the population as a whole (Rolls and Treves, 2011), population coding is known to be important for

complicated tasks (Saxena and Cunningham, 2019). We apply standard population coding metrics

to assess how well odor modality can be decoded from populations of MCs. We show that modality

is well-encoded, but that the differences with GABA$_A$ maniuplations are minor.

We applied two approaches to population decoding. First, we used principal component analy-

sis (**PCA**) to project the population response onto a two-dimensional subspace of highest variance,

and then applied linear discriminant analysis (**LDA**) to find an optimal decoder in that subspace

Figure 6: **Population decoding of ortho versus retro is reliable and robust to altered inhibition. A)** Each drug preparation has a different number of recordings and each recording has a different number of simultaneously recorded MCs. Showing recording number (sorted) and MC population size. **B)** Decoding accuracies (one for each simultaneous recording) for all drug preparations using a 2D PCA dimension reduction following by LDA. Inset shows variance explained by PCA (sorted) by recording. **C)** Decoding accuracies using full dimensional support vector machine (**SVM**) classification. Average decoding accuracies by both methods (**B**)–**C**)) are statistically indistinguishable (minimum $p-$value for all relationships and all 3 statistical tests is: $p \geq 0.32$ for PCA+LDA, $p \geq 0.13$ for SVM).

(PCA+LDA). Second, we used a support vector machine (**SVM**), a common supervised learning algorithm to (nonlinearly) classify data. With both methods, we find odor modality is still encoded well above chance even when inhibition is altered (Fig 6B,C). Not only is the average decoding accuracy (across recordings) higher than before (Fig 1C) but there are no longer decoding accuracies near chance level (0.5). This is perhaps expected because these metrics (PCA+LDA, SVM) exploit the higher dimensional population activity to perform optimal for a binary classification problem when in reality the MCs encode other aspects of stimuli. Note that nearly 35% of recordings with no drug and Mus have perfect decoding accuracy (Fig 6B,C).

In contrast to the case of a single MC's encoding (Fig 1C), we do not find any statistically significant differences between the mean decoding accuracy across drug preparations. In fact, the minimum $p-$value for all relationships and all 3 statistical tests is $p \geq 0.32$ for PCA+LDA, $p \geq 0.13$ for SVM. We expect population coding to perform better than individual cell coding, especially given the large dimensionality of population activity and that these metrics (PCA+LDA, SVM) were optimized for a binary classification task.

Discussion

We have shown that MCs in the OB encode odor modality with our experimental data, and develop an OB model inspired by many prior studies and constrained by our rich data to analyze plausible network components that support these dynamics. Our study uncovers further details for how ortho versus retro information is encoded to early cortical regions via the OB, which is consistent with reports that humans are able to discriminate whether food odors are delivered ortho or retronasally without being told which modality (Frasnelli et al., 2008). Imaging studies of the human mouth have shown that with food odors, ortho stimuli delivery versus retro are two distinct modalities, rather than 2 different stimulus routes to the same modality, independent of odor intensity (Bender et al., 2009). Based on *in vivo* rat data, we found significant differences in decoding accuracies (our proxy for encoding assuming an ideal observer) for classifying ortho and retronasal odors across different drug preparations. The intact (no drug) circuit had the best (average) decoding accuracy, suggesting that the OB might be structured to encode this important aspect of odor input.

Our model framework used prior results, data constraints (Xiao et al., 2021; Ly et al., 2021; Barreiro et al., 2017), and simplicity to account for the results in our data. Many prior studies (Sanganahalli et al., 2020; Gautam and Verhagen, 2012b; Furudono et al., 2013; Scott et al., 2007; Carey et al., 2009) showed that there are differences in the temporal dynamics ORN response to ortho versus retro stimulation. We used this to investigate the OB circuit components that promote efficient coding of individual mitral cells with drug manipulations of inhibitory synapses. Further data analysis revealed that inhibitory drugs have a stronger effect on the population firing with ortho stimulation than retro. Using these insights, we constructed a standard firing rate model that captures the various drug effects for ortho versus retro stimulus, using only a pair of $E - I$ cells. Including synaptic depression was crucial for capturing the different drug effects of firing rate. We did not explore varying the many parameters of the model to find a 'best fit' to data, rather we wanted a simple yet informative model that demonstrates a plausible mechanism. Lastly, we

simulated trial-to-trial variability by assuming a negative binomial distribution with mean defined by the firing rate model output. The resulting model captured both decoding accuracy trends (ND is better) and some aspects of firing rate trial variability.

There have been prior studies that implemented an OB circuit model with varying levels of realism, but none that we are aware of that uses an OB model to account for coding of odor modality and firing rate dynamics. We previously developed a realistic biophysical OB network model (Craft et al., 2021) and used it to account for different neural network dynamics (trial-averaged) with ortho/retro inputs, but did not focus on decoding accuracy for a given trial (which is more computationally taxing). In our prior work (Craft et al., 2021), the PSTH for ortho versus retro are different than here (Figs 1Aii, 3B) because we previously used only 1 rat versus 8 rats here. There are biophysical models of particular OB cells (Viertel and Borisyuk, 2019; Li and Cleland, 2013) and of the OB network (Bathellier et al., 2006; Li and Cleland, 2017), and other have used models (firing rate or Bayesian) to investigate coding of mixed odors (Grabska-Barwińska et al., 2017) and learning new odors (and their concentration levels) (Hiratani and Latham, 2020), but none that we know of that focus on coding of odor modality.

A limitation of this study is the use of one odor (EB, a food odor) with one concentration. We collected data from a non-food odor (1-Hexanol), but since typically only food odors experienced retronasally (Small et al., 2005), we did not include 1-Hexanol in our study. It is well-known that non-food odors delivered retronasally degrade human perception of odors (Hannum et al., 2021; Bender et al., 2009; Frasnelli et al., 2008). Other food odors have different physicochemical properties that impact differences in ortho versus retro responses (Scott et al., 2007), so the generalization of our results merits further investigation. Concentration of odors alters firing rate (Tan et al., 2010) and is thus a confounding factor in distinguishing ortho vs retro from firing rates alone. Our data was collected from anesthetized rats with forced air to model ortho/retro (Gautam and Verhagen, 2012b,a; Gautam et al., 2014) which has clear advantages: control and a fair decoding 'task' so that ortho and retro are mechanically similar with same stimulus duration.

In addition to the intact circuit, our data also had both increased and decreased $GABA_A$ synaptic inhibition with direct manipulations in the OB. Via pooling data from many rats, we had a large number MCs. Another limitation is that the rats were not awake and were not eating. We are currently unaware of any data with awake rodents that record MC spiking activity for the purposes of comparing ortho and retro (food) odors during feeding. There are freely available MC spike data with awake rodents (mice in Bolding and Franks (2018, 2021)) but not during feeding and not with direct manipulation of OB inhibitory synapses.

Since the temporal dynamics of odors can be crucial in OB processing, we systematically varied the evoked time window and even considered removing the first 300 ms of the evoked state (see Fig A2). We found that the population-average (and standard deviation) decoding accuracy varied slightly (Fig A2B,D), and indeed with the first part of the evoked state removed the average decoding accuracy increased. In the context of our study, the temporal patterning differences between ortho and retro have minor effects on decoding. In awake, intact animals, the timing of the breath cycle determines the timing of ortho and retro stimulation (Shusterman et al., 2011). In principle, brain areas downstream from OB could use a breath-timing-based decoding scheme to entirely distinguish ortho from retro. However, our experiments show that timing is not the only mechanism that impacts ortho versus retro coding. In our experiments, the breath of the anesthetized rats did not pass through the nose at all; it was redirected through a tracheotomy tube. Thus, the timing of ortho and retro signals in our experiments was decoupled from breath timing and we kept the stimulation parameters (concentration and flow rate) the same, changing only the direction of airflow. Yet, we found differences in ortho and retro response in OB. Also, without breathing or sniffing, humans can correctly report ortho versus retro odor modality with food odor without being told the modality (Frasnelli et al., 2008). Thus, we conclude that the OB encodes ortho and retro stimuli not just with the timing of the breath, but also with other mechanisms. Nonetheless, it is likely that intact, awake animals would employ both timing (inhale vs exhale) and the mechanisms we identify here. Disentangling the relative importance of these different

possible contributions to ortho versus retro coding will require further experiments.

Our results predominately center on population averages of individual MC decoding accuracy because it is a logical first step (Rolls and Treves, 2011). Given the importance of population coding for many complicated tasks (Saxena and Cunningham, 2019), we also considered 2 standard population decoding accuracies measures: PCA+LDA and SVM and found in both cases that the intact circuit was not statistically significant better (on average). However, decoding accuracy was again much higher than chance levels and better compared to average individual MC decoding accuracy. Developing a heterogeneous OB model that accounts for the population coding results and dynamics is a potential next step but beyond the scope of this current study.

Author Contributions

MC programmed and implemented models in software, including the data analysis. MC and CL developed the computational models. MC, AKB, WLS, CL conceptualized and developed the project. SHG and WLS designed the experiments and collected the data. MC and CL drafted the original manuscript, including visualizations. MC, AKB, SHG, WLS, CL edited the manuscript. CL supervised the project.

References

Abraham, N. M., V. Egger, D. R. Shimshek, R. Renden, I. Fukunaga, R. Sprengel, P. H. Seeburg, M. Klugmann, T. W. Margrie, A. T. Schaefer, et al. (2010). Synaptic inhibition in the olfactory bulb accelerates odor discrimination in mice. Neuron 65(3), 399–411.

Barreiro, A., S. Gautam, W. Shew, and C. Ly (2017). A theoretical framework for analyzing coupled neuronal networks: Application to the olfactory system. PLoS Computational Biology 13, e1005780.

Bathellier, B., S. Lagier, P. Faure, and P.-M. Lledo (2006). Circuit properties generating gamma oscillations in a network model of the olfactory bulb. Journal of neurophysiology 95(4), 2678–2691.

Bender, G., T. Hummel, S. Negoias, and D. M. Small (2009). Separate signals for orthonasal vs. retronasal perception of food but not nonfood odors. Behavioral neuroscience 123(3), 481.

Blankenship, M. L., M. Grigorova, D. B. Katz, and J. X. Maier (2019). Retronasal odor perception requires taste cortex, but orthonasal does not. Current biology 29(1), 62–69.

Bolding, K. and K. Franks (2021). Simultaneous extracellular recordings from mice olfactory bulb (ob) and piriform cortex (pcx) and respiration data in response to odor stimuli and optogenetic stimulation of ob. CRCNS. org Date accessed: January.

Bolding, K. A. and K. M. Franks (2018). Recurrent cortical circuits implement concentration-invariant odor coding. Science 361(6407).

Cang, J. and J. S. Isaacson (2003). In vivo whole-cell recording of odor-evoked synaptic transmission in the rat olfactory bulb. Journal of Neuroscience 23(10), 4108–4116.

Carey, R. M., J. V. Verhagen, D. W. Wesson, N. Pírez, and M. Wachowiak (2009). Temporal structure of receptor neuron input to the olfactory bulb imaged in behaving rats. Journal of neurophysiology 101(2), 1073–1088.

Churchland, M., B. Yu, et al. (2010). Stimulus onset quenches neural variability: a widespread cortical phenomenon. Nature Neuroscience 13, 369–378.

Cohen, J. (2013). Statistical power analysis for the behavioral sciences. Routledge.

Craft, M., A. Barreiro, S. Gautam, W. Shew, and C. Ly (2021). Differences in olfactory bulb mitral cell spiking with ortho- and retronasal stimulation revealed by data-driven models. PLoS Computational Biology 17(9), e1009169.

Cunningham, J. P. and B. M. Yu (2014). Dimensionality reduction for large-scale neural recordings. Nature neuroscience 17(11), 1500–1509.

Dietz, S. B. and V. N. Murthy (2005). Contrasting short-term plasticity at two sides of the mitral–granule reciprocal synapse in the mammalian olfactory bulb. The Journal of physiology 569(2), 475–488.

Frasnelli, J., M. Ungermann, and T. Hummel (2008). Ortho-and retronasal presentation of olfactory stimuli modulates odor percepts. Chemosensory Perception 1(1), 9–15.

Furudono, Y., G. Cruz, and G. Lowe (2013). Glomerular input patterns in the mouse olfactory bulb evoked by retronasal odor stimuli. BMC neuroscience 14(1), 45.

Gautam, S. H., S. M. Short, and J. V. Verhagen (2014). Retronasal odor concentration coding in glomeruli of the rat olfactory bulb. Frontiers in Integrative Neuroscience 8, 81–81.

Gautam, S. H. and J. V. Verhagen (2012a). Direct behavioral evidence for retronasal olfaction in rats. PloS one 7(9), e44781.

Gautam, S. H. and J. V. Verhagen (2012b). Retronasal odor representations in the dorsal olfactory bulb of rats. The Journal of Neuroscience 32(23), 7949–7959.

Grabska-Barwińska, A., S. Barthelmé, J. Beck, Z. F. Mainen, A. Pouget, and P. E. Latham (2017). A probabilistic approach to demixing odors. Nature Neuroscience 20, 98–106.

Hannum, M. E., J. A. Fryer, and C. T. Simons (2021). Non-food odors and the duality of smell: Impact of odorant delivery pathway and labeling convention on olfactory perception. Physiology & Behavior 238, 113480.

Hiratani, N. and P. E. Latham (2020). Rapid bayesian learning in the mammalian olfactory system. Nature communications 11(1), 3845.

Li, G. and T. A. Cleland (2013). A two-layer biophysical model of cholinergic neuromodulation in olfactory bulb. Journal of Neuroscience 33(7), 3037–3058.

Li, G. and T. A. Cleland (2017). A coupled-oscillator model of olfactory bulb gamma oscillations. PLoS computational biology 13(11), e1005760.

Ly, C., A. K. Barreiro, S. H. Gautam, and W. L. Shew (2021). Odor-evoked increases in olfactory bulb mitral cell spiking variability. iScience 24(9), 102946.

Mandairon, N. and C. Linster (2009). Odor perception and olfactory bulb plasticity in adult mammals. Journal of neurophysiology 101(5), 2204–2209.

Meredith, M. (1992). Neural circuit computation: complex patterns in the olfactory bulb. Brain research bulletin 29(1), 111–117.

Rall, W., G. M. Shepherd, T. S. Reese, and M. W. Brightman (1966). Dendrodendritic synaptic pathway for inhibition in the olfactory bulb. Experimental neurology 14(1), 44–56.

Rolls, E. T. and A. Treves (2011). The neuronal encoding of information in the brain. Progress in neurobiology 95(3), 448–490.

Rossant, C., S. Kadir, D. Goodman, J. Schulman, M. Hunter, A. Saleem, A. Grosmark, M. Belluscio, G. Denfield, A. Ecker, A. Tolias, S. Solomon, G. Buzsáki, M. Carandini, and K. Harris (2016). Spike sorting for large, dense electrode arrays. Nature neuroscience 19(4), 634–641.

Sanganahalli, B. G., K. L. Baker, G. J. Thompson, P. Herman, G. M. Shepherd, J. V. Verhagen, and F. Hyder (2020). Orthonasal versus retronasal glomerular activity in rat olfactory bulb by fmri. NeuroImage 212.

Saxena, S. and J. P. Cunningham (2019). Towards the neural population doctrine. Current opinion in neurobiology 55, 103–111.

Scott, J. W., H. P. Acevedo, L. Sherrill, and M. Phan (2007). Responses of the rat olfactory epithelium to retronasal air flow. Journal of neurophysiology 97(3), 1941–1950.

Shusterman, R., M. C. Smear, A. A. Koulakov, and D. Rinberg (2011). Precise olfactory responses tile the sniff cycle. Nature neuroscience 14(8), 1039–1044.

Small, D. M., J. C. Gerber, Y. E. Mak, and T. Hummel (2005). Differential neural responses evoked by orthonasal versus retronasal odorant perception in humans. Neuron 47(4), 593–605.

Tan, J., A. Savigner, M. Ma, and M. Luo (2010). Odor information processing by the olfactory bulb analyzed in gene-targeted mice. Neuron 65(6), 912–926.

Tomczak, M. and E. Tomczak (2014). The need to report effect size estimates revisited. an overview of some recommended measures of effect size. Trends in sport sciences 21(1).

Viertel, R. and A. Borisyuk (2019). A computational model of the mammalian external tufted cell. Journal of theoretical biology 462, 109–121.

Wang, Z.-J., L. Sun, and T. Heinbockel (2012). Cannabinoid receptor-mediated regulation of neuronal activity and signaling in glomeruli of the main olfactory bulb. Journal of Neuroscience 32(25), 8475–8479.

Wilson, R. I. and G. Laurent (2005). Role of gabaergic inhibition in shaping odor-evoked spatiotemporal patterns in the drosophila antennal lobe. Journal of Neuroscience 25(40), 9069–9079.

Xiao, Z.-C., K. K. Lin, and L.-S. Young (2021). A data-informed mean-field approach to mapping of cortical parameter landscapes. PLoS computational biology 17(12), e1009718.

Appendix

The main text focuses on the 1 s duration of the odor, which is longer than most breath cycles. However, there are dynamics beyond the duration of the odor that may be insightful and of interest, so the population PSTHs extended to 5 s after odor onset are shown in Figures A1A,B. By visual inspection, there appears to be an increase in the firing rate after the odor is off, e.g., ortho with Bic and retro with Mus. Whether this change is statistical significant, and the underlying mechanisms to explain this are fascinating but beyond the scope of this current study. For the interested readers, we report the fraction of 'Off-Response' cells that have higher firing rates when the odor is removed than with odor, a well-known phenomena dating back many decades (Meredith, 1992). We report the fraction of the MC population the have a higher firing rate (time and trial-averaged) in $1 \leq t \leq 2$ s than in $0 \leq t < 1$ s in Table A1.

Table A1: **Fraction of 'Off-Response' cells: cells that have a higher firing rate (time and trial-averaged) in the immediate time (1 s) after odor is removed than during the 1 s odor stimulation.**

Fraction of Off-Response MCs	ND	Bic	Mus
Ortho	30.12%	36.08%	27.45%
Retro	27.05%	18.64%	42.00%

Table A2: **Correlation between trial-mean and trial-standard deviation of spiking rate.** There is a strong positive relationship, see Figure A3. This is consistent with MC in OB reported in Ly et al. (2021), but notably is *inconsistent* with cortical neurons (Churchland et al., 2010).

	Ortho		Retro	
	Pearson's	**Spearman's Rank**	**Pearson's**	**Spearmen's Rank**
No Drug	0.932	0.986	0.935	0.995
Bicculline	0.930	0.989	0.937	0.988
Muscimol	0.907	0.989	0.920	0.996

Figure A1: **Supplemental data analysis: extended time of PSTHs in main text.** Extended timeframe for Figure 1Aii (**A**) and Figure 3B (**B**).

Decoding accuracy dynamics from drug effects on population-averaged firing rates

If we assume the population trends in Fig 3B) hold individually for MCs (drugs only effect spiking rate with ortho but not retro), then we might expect decoding accuracies to change in specific ways, as illustrated in Fig A4**Ai,Aii**. If ortho spiking is less than retro with no drug ($\mu_O < \mu_R$, see Fig A4**Ai**), Bic should increase spiking (shift gray curve to the right, green curve) causing more overlap and lower decoding accuracy, while Mus should decrease spiking (shift gray curve left, purple curve) leading to less overlap and higher decoding accuracy. Similar arguments can be made for when retro spiking is less than ortho spiking ($\mu_R < \mu_O$, Fig A4Aii). Thus, the predictions from observations of the population firing rates (Fig 2**B**) are:

1. When $\mu_O < \mu_R$:

   (a) Bicuculline = less accuracy*

   (b) Muscimol = more accuracy

2. When $\mu_R < \mu_O$:

   (a) Muscimol = less accuracy*

   (b) Bicuculline = more accuracy

The results of these predictions are shown in Fig A4**B,C**. Predictions 1a and 2a (Fig A4**B**) denoted by asterisk are statistically significant in difference ($t-$test ($p_{ND,Bic} = 5.3{\times}10^{-16}$, $p_{ND,Mus} = 1.8{\times}10^{-6}$), Wilcoxon rank sum test ($p_{ND,Bic} = 3.1{\times}10^{-14}$, $p_{ND,Mus} = 6.9{\times}10^{-7}$), and one-way ANOVA ($p_{ND,Bic} = 1.1 \times 10^{-15}$, $p_{ND,Mus} = 5.6 \times 10^{-6}$)). The other two predictions (1b and 2b, Fig A4**C**) do not precisely hold; interestingly, it is because no drug preparation retains a higher average decoding accuracy than with the two drug applications. Two caveats: in Fig A4**Ci**, with Mus the average decoding accuracy is better when $\mu_O < \mu_R$ (purple filled) compared to all cells $\mu_O < \mu_R$ & $\mu_O \geq \mu_R$ (dark purple outline) ($t-$test ($p_{Mus-all,Mus} = 1.2 \times 10^{-8}$), Wilcoxon rank sum test ($p_{Mus-all,Mus} = 9.4 \times 10^{-6}$), and one-way ANOVA ($p_{Mus-all,Mus} = 3.0 \times 10^{-6}$)); in Fig A4**Cii**, the difference between Bic and no drug are no longer statistically significant ($t-$test ($p_{ND,Bic} = 0.22$), Wilcoxon rank sum test ($p_{ND,Bic} = 0.58$), and one-way ANOVA ($p_{ND,Bic} = 0.26$)), so that with $\mu_R < \mu_O$, cells with Bic and no drug have indistinguishable averages. This all shows that the observation that ortho stim is more effected with GABA$_A$ manipulation than retro has consequences for decoding accuracy modulation, even if the predictions are imperfect.

Figure A2: **How we choose the time window; decoding accuracy as time window varies.** Varying time windows on evoked firing rate to select for data analysis (**A, C**) and decoding accuracy. **A)** Vertical axes show $p-$values (log-scale) from comparisons of average decoding accuracies with different drug preparations using two different statistical tests: two sample $t-$test assuming unequal variance (black) and Wilcoxon rank sum test (magenta). Open circles compares no drug to Bic, stars compare no drug to Mus (see legend). **B)** Showing the population average decoding accuracy and $\pm$ one standard deviation in shaded region. Longer time windows generally lead to larger average decoding accuracy. **C), D)** are similar to **A), B)** except the first 300 ms of the evoked state are removed to test whether decoding accuracies increase are significant; this was based on the population PSTH being similar for ortho and retro immediately after stimulus onset (Fig 1Aii). Decoding accuracies are significantly different ($\alpha = 0.01$, denoted by dashed black line) for a large range of time windows, but we chose 0.9 s for evoked because this value resulted in the smallest $p-$values generally, the exception being denoted by the red arrows where other time bins have *slightly* smaller $p-$values. We decided to keep the entire time, not excluding the first 300 ms, for simplicity and because the $p-$values are similar. All data analysis is with EB odor input.

Figure A3: **Trial-averaged firing rate is positively correlated with trial-to-trial variability.** For each MC in a given modality and drug preparation, showing the trial-averaged firing rate versus the standard deviation of firing rate (across trials). **A)** With orthonosal stimulation; **B)** with retronasal stimulation. We plot the time average to resolve the many instances with no spiking so that each MC has a single number for the firing rate and a single number for the standard deviation.

Figure A4: **Decoding accuracy dynamics from drug effects on population-averaged firing rates.**
Left column: cells where $\mu_O < \mu_R$; Right column: cells where $\mu_R < \mu_O$. **Ai–Aii**) predicted
decoding accuracy for different drug preparations. **Bi**) Histogram of decoding accuracies for no
drug and Bic when $\mu_O < \mu_R$, and no drug with muscimol when $\mu_R < \mu_O$ (**Bii**); triangle is
mean decoding accuracy over all such MCs. Differences in middle row (**Bi,Bii**) are statistically
significant. **Ci**) Histogram of decoding accuracies for no drug and muscimol when $\mu_O < \mu_R$, also
showing the histogram for muscimol decoding accuracy for all cells (i.e., not only those where
$\mu_O < \mu_R$) in dark purple outline. Although no drug is better than muscimol, muscimol with cells
$\mu_O < \mu_R$ have better decoding than all cells with muscimol. **Cii**) No drug and Bic when $\mu_R < \mu_O$
do not have statistically significant differences in mean. See text for $p-$values.