1

Widespread epistasis shapes RNA Polymerase II active site function and evolution

Authors: Bingbing Duan[1], Chenxi Qiu[2], Sing-Hoi Sze[3, 4], Craig Kaplan[1*]

Affiliations:

[1] Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260, USA

[2] Department of Genetics, Harvard Medical School, Boston, MA 02215, USA

[3] Department of Computer Science and Engineering, Texas A&M University, College Station, TX 77843, USA

[4] Department of Biochemistry & Biophysics, Texas A&M University, College Station, TX 77843, USA

Contact information: craig.kaplan@pitt.edu

ABSTRACT

Multi-subunit RNA Polymerases (msRNAPs) are responsible for transcription in all kingdoms of life. These enzymes rely on dynamic, highly conserved active site domains such as the so-called "trigger loop" (TL) to accomplish steps in the transcription cycle. Mutations in the RNA polymerase II (Pol II) TL confer a spectrum of biochemical and genetic phenotypes that suggest two main classes, which decrease or increase catalysis or other nucleotide addition cycle (NAC) events. The Pol II active site relies on networks of residue interactions to function, and mutations likely perturb these networks in ways that may alter mechanisms. We have undertaken a structural genetics approach to reveal residue interactions within and surrounding the Pol II TL – determining its "interaction landscape" – by deep mutational scanning in *Saccharomyces cerevisiae* Pol II. This analysis reveals connections between TL residues and surrounding domains, demonstrating that TL function is tightly coupled to its specific enzyme context.

INTRODUCTION

Transcription from cellular genomes is performed by conserved multi-subunit RNA polymerases (msRNAPs)[1-3]. While bacteria and archaea use a single msRNAP to transcribe all RNAs[4-6], eukaryotes have at least three msRNAPs (Pol I, II, and III) for different RNA types[7-10]. RNA synthesis by msRNAPs occurs by iterative nucleotide addition cycles (NAC) of nucleotide selection, catalysis and polymerase translocation[11-15]. The msRNAPs active sites accomplish these NAC steps using two conformationally flexible domains, the bridge helix (BH) and trigger loop (TL)[5,12,16-19]. The multi-functional natures of the BH and TL likely underlie their striking conservation, serving as interesting models for studying the function and evolution of extremely constrained protein domains.

Most catalytic cycle events involve concerted conformational changes in the TL, and potentially the BH[12-14,19-23]. The extent to which these conformations are supported by, or communicate with, the surrounding enzymatic context can be examined by structural or computational approaches, but functional importance needs to be determined experimentally. The BH is straight in most msRNAP structures[7,8,12,13,24], but appears kinked in *Thermus thermophilus* (bacteria) RNAP[5,25]. Simulations suggest these dynamics may promote msRNAP translocation[16,25-28]. The TL adopts various conformations to support its multiple functions. During each NAC, the TL nucleotide interaction region (NIR) discriminates correct NTPs from mismatched ones, initiating a shift from a catalytic-disfavoring "open" state to a catalytic-favoring "closed" state[11,20,29,30]. This closure promotes catalysis[20,31], and subsequent TL reopening is proposed to support polymerase translocation to the next template DNA position, allowing for the next NAC[32-34](**Fig. 1a**). Disrupting the TL with mutations affects transcription at all stages[13,17,23,35-39]. For instance, NIR mutations impair TL-substrate interactions, resulting in reduced catalysis and elongation rate in vitro (Loss of function, LOF)[17,23,36,37,40,41]. Mutations in the TL hinge and C-terminal regions appear to shift the TL toward the active state, enhancing catalysis and elongation rate but impairing transcription fidelity (Gain of function, GOF)[17,42,43]. Mutations may also affect other transcription processes such as proofreading, pausing and termination[12,43-46], as TL confirmations are associated with these activities in addition to catalysis and translocation.

All TL conformational dynamics and functions are likely balanced by residue interactions within and around the TL[41,45,47-49]. Embedded in the conserved active site, the TL interacts with surrounding domains such as the BH and α46-47 helices, forming a five-helix bundle enclosing a hydrophobic pocket that supports these dynamics **(Fig. 1a)**[12,42]. Mutations in TL adjacent domains (e.g., BH GOF T834P and LOF T834A, funnel helix α-21 GOF S713P, and Rpb2 GOF Y769F)[17,23,25,26,35-37] **(Fig. 1a)**, and in a Pol II subunit which does not directly contact Pol II active site, such as Rpb9[50], also alter catalytic activity and fidelity, showing phenotypes seemingly identical to TL mutants, indicating that TL dynamics and function are finely balanced and could be sensitive to allosteric effects[50-53]. Determining the extent and nature of the pathways connecting the TL to the rest of the polymerase will reveal the functional networks that shape its dynamics and the potential pathways through which Pol II has evolved or may be controlled by elongation factors.

Physical and functional intramolecular interactions between amino acids define protein function and evolvability[54-57]. Genetic interactions where one mutation modulates or alters the phenotype of another are commonly referred to as epistasis. Interactions can be of different types such as suppressive, masking, or synthetic interactions. In this study, we use "epistasis" specifically to

describe cases where mutant phenotypes allele-specifically depend on the identities of other
amino acids. Epistasis contributes to protein evolvability by providing a physical context and an
evolutionary window in which some intolerable mutations may be tolerated[58-60]. Recent studies
have shown how mutations can alter protein function, allostery, and evolvability, suggesting that
even conserved residues are subject to unique, context dependent epistatic constraints[61-68].
Consistent with this, distinct phenotypes for the same conserved residues have been observed
in various proteins, including yeast Pol I and Pol II[66,67,69]. For example, the TL domain of yeast
Pol I is incompatible with Pol II, despite sharing 70% of sequence identity, while the similarly
distinct Pol III TL maintains some function[70]. It is not known generally if highly conserved TL
sequences are compatible in a heterologous msRNAP context. Answering this question would
reveal how tightly integrated the TL is by coevolution with its embedded context, or the extent to
which its functions are self-contained due to its very high conservation.

Functional interactions between residues can be revealed by genetic interactions of double
mutants[17,37,71-73]. A systematic analysis is needed to fully describe and understand the
requirements for each mutant's phenotype, and the nature of the interactions that control Pol II
activity. Previous studies from our lab on a small subset of site-directed substitutions identified
distinct types of Pol II double mutant interactions, including suppression, synthetic sickness and
lethality (activity-additive), less-than-additive interactions (sometimes referred to as epistasis),
and sign-epistasis[17,37,72]. Suppression interactions are common between LOF and GOF
mutants, as expected if each mutant is individually acting in the double mutant, balancing
opposing effects on activity. Similarly, synthetic lethality or sickness were common between
mutants of the same class (GOF or LOF), consistent with greater defects arising from combining
partial losses of TL function. However as just noted, we also observed mutant combinations
where the double mutant was not sicker than the single mutants for mutants of the same class,
suggesting these mutants might act at the same step. Additionally, we had observed a potential
case of sign-epistasis, where a mutant's phenotype class appeared dependent on the identity of
another residue. For example, the GOF TL substitution Rpb1 F1084I was unexpectedly lethal
with the LOF TL substitution Rpb1 H1085Y, contrary to the predicted mutual suppression for a
GOF/LOF combination[37,72]. This suggested that F1084I required H1085 for its GOF
characteristics and became a LOF mutant in the presence of H1085Y (**Fig. 1b**).

Deducing large-scale residue interaction networks is challenging. To accomplish this for Pol II,
we developed experimental and analytical schemes to extend the previously established "Pol II
phenotypic landscape"[17,37,72,74] to 15,174 double and multiple mutants within the *S. cerevisiae*
Pol II TL and between the TL and adjacent domains (a "Pol II Interaction landscape"). Our
analyses indicate that TL function and evolution are dominated by widespread epistasis. The TL
faces extensive constraints despite its high conservation, arising from residues within the TL
and its interacting domains, resulting in highly similar TLs being incompatible inside yeast Pol II
to a large degree. Additionally, individual mutants with similar biochemical and genetic
phenotypes have quite distinct genetic interaction networks, revealing differential functional
requirements for their abilities to confer altered Pol II activities. Some mutants minimally change
the genetic behavior of most TL substitutions when combined, suggesting that TL conformations
or interactions remain largely unchanged, with altered TL dynamics hypothesized to be the sole
difference. In contrast, other alleles greatly perturb the genetic behavior of many TL
substitutions, indicating these substitutions change the active site landscape in ways that would
otherwise be undetectable. Together, these results suggest small changes in the Pol II active
site can dramatically change its constraints and therefore, its potential evolutionary paths.

Furthermore, our findings highlight potential communication networks through evolutionary
coupling of Pol II residues within the active site, which may underlie allosteric communication
paths to the active site for elongation factors.

RESULTS

A system for dissection of the Pol II active site genetic interaction landscape. We
developed an experimental and analytical framework, termed the Pol II TL interaction
landscape, to dissect residue interactions that shape Pol II TL function and evolution in *S.*
cerevisiae. We designed and synthesized 15,174 variants representing evolutionary haplotypes,
all possible Pol II TL single mutants and a subset of targeted double mutants in ten libraries
(Supplemental Table 1). This approach follows our prior analysis of the TL phenotypic
landscape[17] with modifications (**Methods** and **Extended Data Fig. 1a**). Libraries were
transformed, screened under selected conditions (**Methods**), and sequenced (**Fig. 1c,**
Extended Data Fig. 1a). Mutant growth phenotypes are calculated based on relative allele
frequency shifts compared to a control condition and normalized to the WT under identical
conditions. Biological replicates indicate high reproducibility (**Extended Data Fig. 1b-c**).
Individual libraries were min-max normalized[75] to account for scaling differences between
libraries (**Fig. S1a**) and the same mutants present among different libraries indicate high
correlation of fitness determinations in each library (**Fig. S1b-c**).

We defined a conceptual framework for evaluating genetic interactions among TL mutations.
First, we assume that independent mutant effects combine in log-additive manner. This means
that predicted double mutant fitness defects should be the combination of both single mutant
defects, as is standardly assumed[60,71,73,76]. Deviations from log-additive expectations represent
genetic interactions. To quantify any potential deviation, we compared the observed fitness of a
double mutant measured in our high-throughput experiment to its expected fitness, calculated
as the addition of constituent single substitutions' fitnesses, which are expressed as a log value.
We term these differences as a "deviation score" (Deviation score = observed fitness –
expected fitness; **Extended Data Fig. 2**). We consider a deviation score between -1 and 1 to be
consistent with additive behavior, meaning independent contributions of single mutants to the
double mutant fitness. A deviation score > 1 is taken as a positive genetic interaction (e.g.,
suppression), while a score < -1 indicates a negative genetic interaction (e.g., synthetic
sickness or lethality).

Importantly, our phenotyping analyses determine both general mutant fitness and predict their
class of in vitro biochemical defect. Our previous studies showed that phenotypic profiles from
conditional growth assays, reflecting Pol II-activity dependent phenotypes in vivo, correlate with
the measured biochemical activity in vitro[17,37,77]. For example, catalytically hyperactive GOF
mutants have a phenotypic profile based on sensitivity to mycophenolic acid (MPA) and weak
suppression of two genetic reporters (*gal10Δ56* and *lys2-128∂*), while catalytically defective LOF
mutants are MPA resistant, show strong suppression of *gal10Δ56* and fail to suppress *lys2-128∂*
(more fully described in Methods). Using the established correlation between growth profiles
(MPA and media used to detect phenotypes of mutants on *gal10Δ56* and *lys2-128∂*
backgrounds) and known/predicted in vitro catalytic effects (GOF or LOF) of 65 TL
alleles[17,37,53,78], we trained two multiple logistic regression models to distinguish Pol II mutant
classes, and applied them to all viable mutants (growth fitness score > -6.5) of designed
variants, classifying them into three groups, GOF (22.96%), LOF (28.11%) and unclassified
(mutants not belonging to either class, 48.93%) (**Extended Data Fig. 3a,** Methods). To visually

inspect the classifications, we applied t-SNE projection and k-means clustering for all measured
mutants across all growth conditions. We observed separated GOF and LOF clusters consistent
with logistic regression classifications (**Extended Data Fig. 3b-c**). Notably, mutants with similar
defects may be further distinguished in different clusters, suggesting their similar phenotypic
defects may be related to different transcription defects (**Extended Data Fig. 3c**). By classifying
Pol II active site mutants as "GOF" or "LOF" based on predicted in vitro biochemical
phenotypes, we establish baseline expectations for their genetic interactions. We previously
observed activity-additive interactions, meaning suppression between mutants of different
classes (GOF+LOF) or synthetic sickness/lethality within the same class (GOF+GOF or
LOF+LOF). Together with the calculated deviation scores, we can distinguish specific epistatic
interactions from activity-dependent effects. We classified an interaction as epistasis when we
observed positive fitness deviation in mutants of the same activity class (GOF+GOF,
LOF+LOF), where we would otherwise expect synthetic sickness or lethality if mutants were
functioning independently. Conversely, we defined sign epistasis for situations where negative
deviations in combinations between the classes (GOF+LOF), where suppression would be
expected for independent mutants (**Fig. 1d, Extended Data Fig. 2**).

We compared interactions observed in sequencing to growth assays for 59 previously examined
double mutants[17,37], with 48 matching our observations. The other 11 were categorized as ultra-
sick/lethal and therefore inaccessible to sequencing analysis (**Supplemental Table 2**). We
constructed 50 additional mutants to assess interactions by patch assay and spot assay, and
these aligned with our high-throughput results (**Extended Data Fig. 4**).

Widespread incompatibility of conserved TL sequences with yeast Pol II suggests the TL
is highly constrained by its coevolved environment. We previously found that identical
mutations in a residue conserved between the Pol I and Pol II TLs yielded different biochemical
phenotypes[70,79]. Additionally, the yeast Pol I TL was incompatible within the yeast Pol II enzyme,
implying that TL function in that case depended on its enzymatic context[70,79]. To determine the
scope of TL-Pol II incompatibility, we designed a library containing evolutionary TL variants from
bacterial, archaeal, and eukaryotic msRNAPs and determined their compatibility in the yeast Pol
II context (**Fig. 2a**). Eukaryotic TL alleles were more compatible than those from archaea or
bacteria, with Pol II alleles being the most compatible (**Fig. 2b, Extended Data Fig. 5a-b**),
consistent with evolutionary distance. Growth fitness in the Pol II background showed a slight
negative correlation with the number of TL substitutions for most TLs but not those from bacteria
(**Extended Data Fig. 5c**), likely because of the bacterial TLs being largely incompatible with Pol
II. Despite this, some archaeal TLs provide yeast Pol II viability, yet several Pol II TLs were
defective or lethal. These results suggest widespread coevolution between TL sequences and
that the enzymatic background, beyond ultra-conserved positions, shapes TL function in
individual enzymes. We have explored the nature of this incompatibility and complex intra-TL
genetic interactions further in a separate study[80].

Allele-specific interactions suggest unique properties of individual mutants with similar
phenotypes. To determine the potential residue interactions surrounding the TL in the yeast Pol
II environment, we analyzed 12 previously studied GOF and LOF mutants (eight within the TL
and four outside), each combined with >600 perturbations in the form of all single substitution
TL mutants (**Fig. 3a**). These 12 mutants serve as probes for the genetic interaction space of the
TL, revealing how it might be altered in allele-specific fashion by each "probe" mutation.
Importantly, TL conformational dynamics and function are balanced by internal residue

interactions within the TL and by TL-external interactions between the TL and TL-proximal
domains. While GOF and LOF mutants in TL-adjacent domains appear similar to GOF and LOF
TL mutants, their underlying mechanisms or specific residue dependencies are not known. We
therefore compared interaction networks among probe mutants with similar biochemical and
phenotypic defects to distinguish if changes to TL function might reflect simple alterations to TL
dynamics, or additional alterations to folding trajectories or conformations.

We observed distinct interaction networks among probe mutants, even for those with similar
apparent biochemical and growth defects (**Fig. 3b-d**), indicating that mutants differentially
impact the Pol II active site, potentially reflective of different constraints and distinct properties of
individual mutants, as discussed below. A simple expectation is that mutants with altered TL
dynamics yet maintenance of conformational states should show additive interactions, because
the effects of substitutions on function would be maintained. In contrast, substitutions altering TL
conformations/folding or propagating changes across the active site, may be subject to allele-
specific interactions *i.e.* prominent epistasis. We observed that most probe mutants were placed
on continuum of effects based on the balance between additive (independent) interactions and
epistatic (dependent or allele-specific) interactions in double mutants, which we interpret as the
degree to which they propagate their effects across the active site (**Fig. 3b-c, Extended Data
Fig. 6, and Fig. S2**). Probe mutants appeared separable based on the extents of their additive
or epistatic interactions, where greater numbers of independent interactions (additive) would
indicate fewer effects on other residues' functions. In contrast, an increase in dependent
interactions (sign epistasis) would indicate greater changes in other mutant's characters and
presumably residue function. For example, Y769F, a GOF TL-proximal mutant in Rpb2 link
domain (**Fig. 1a**), uniquely suppresses some TL lethal substitutions among all GOF probe
mutants (**Extended Data Fig. 7**), indicating specific functional interactions between Y769F and
these substitutions. Moreover, two TL-adjacent GOF mutants, Rpb1 S713P (funnel α-helix 21)
and T834P (BH) displayed largely distinct interaction networks despite similarly increased
biochemical activity and genetic phenotypes as single mutants[17] **(Fig 3c, Extended Data Fig.
8a)**. S713P broadly suppressed LOF TL substitutions (96 instances), consistent with generic
enhancement and preservation of TL dynamics or residue function. In contrast, T834P exhibited
sparser suppression (33 instances) but much greater sign epistasis than S713P (102 versus 38
instances), especially with P1099, where all substitutions showed sign epistasis with T834P but
not S713P (**Extended Data Fig. 8a**). These results suggest that T834P's GOF function requires
most TL residues to be WT, is especially reliant on proline at 1099, and that T834P alteration to
the BH extensively changes the TL's functional space, likely through folding or altered
conformations.

A similar distinction was observed between two internal TL GOF mutants, Rpb1 E1103G and
F1084I **(Fig. 3c, Extended Data Fig. 8b)**. E1103G showed widespread suppression of LOF TL
substitutions (184 instances), consistent with its proposed role in altering TL dynamics to
promote TL closure[23,37] and which allows TL mutants to maintain their effects. In contrast,
F1084I showed more limited suppression (43 instances) while showing widespread synthetic
lethality due to predicted sign epistasis, indicating that F1084I requires many WT residues for
GOF characteristics and itself may switch to a LOF upon many additional TL perturbations.

The most striking example came from two LOF substitutions at the ultra-conserved Rpb1 H1085
(Extended Data Fig. 8c). This histidine, which contacts incoming NTP substrates[5,12], is the
target for the Pol II inhibitor α-amanitin[36,81], and promotes catalysis[12,13]. Initial structural data and

molecular dynamics simulations suggested that H1085 functions as a general acid for Pol II
catalysis[82-85]. Our discovery that H1085L was well-tolerated[17], and subsequent experiments
from the Landick lab[86,87], have led to their proposal that the TL histidine functions as a positional
catalyst where a similarly sized leucine substitution supports catalysis with mild effects on
biochemistry and growth. If H1085Y and L substitutions represent a continuum of positional
catalyst activity, then their interaction networks would be expected to be similar, and differing
only in magnitude, not identity or type of interactions. Contrary to this, distinct interaction
patterns were observed (**Fig. 3c, Extended Data Fig. 8c**). Most GOF mutants suppressed
H1085Y but not H1085L. Instead, H1085L showed synthetic lethality with most GOF mutants
(putative sign epistasis). For example, almost all substitutions at E1103 showed sign epistasis
with H1085L but not H1085Y (**Fig. S2b, Extended Data Fig. 8c**). The distinction between
H1085L and H1085Y is further highlighted in PCA analysis (**Fig. 3d**).

Functional and epistatic interactions within the TL hydrophobic pocket tune Pol II
activity. Several allele-specific epistatic interactions were observed, with some of the most
extensive occurring between A1076 substitutions and L1101S (**Fig. 4a**), suggesting tight
coupling between A1076 and L1101 for Pol II function. These two hydrophobic residues,
together with other hydrophobic residues in TL helices, likely stabilize the open TL conformation
through hydrophobic packing of the TL helices (**Fig. 4c**). Consistent with this, another pair of
adjacent residues, M1079 and G1097, also showed allele-specific interactions (**Fig. 4b-c**).
Additionally, three GOF/GOF combinations between L1101S and A1076 substitutions showed
WT-like behavior for the double mutants based on our regression model for mutants, indicating
mutual suppression due to one of the singles switching to a different class (sign epistasis),
further supporting a tight functional interaction between 1101 and 1076 (**Fig. S3a**).

We also observed allele-specific interactions for predicted lethal mutants. Our lethality threshold
is more conservative than the actual lethal threshold and therefore, some classified lethal but
actually slow-growing mutants have enough growth in phenotypic assays for GOF or LOF class
determination. Among 21 ultra-sick/lethal TL substitutions predicted as GOF, suppression by
LOF mutants was expected and observed (**Fig. S3b**). For example, lethal A1076 substitutions
were suppressed by LOF probe mutants, suggesting their severe defects result from strong
GOF status. However, some A1076 substitutions were also suppressed by the GOF probe
mutant L1101S, indicating allele-specific mutual suppression or sign-epistasis, consistent with
A1076-L1101 interactions described above (**Fig. S3c**). Similarly, ultra-sick/lethal LOF
substitutions were commonly suppressed by GOF mutants, but some were suppressed by
specific LOF mutants (**Fig. S3b**). For example, S1091G was suppressed by most GOF mutants
and specifically by the LOF V1094D (**Fig. S3c**), suggesting allele-specific compensation.
Furthermore, some lethal mutants exhibited interactions with specific GOF or LOF probe
mutants. For example, F1084R, was specifically suppressed by LOF probe Q1078S and GOF
probe Y769F (**Fig. S3c**). Notably, F1084, Q1078S and Y769 are in close proximity in the
substrate bound, catalytic-favoring TL conformation. These types of allele-specific interactions
potentially allow the TL and adjacent residues to evolve and differentiate while maintaining
essential functions.

We note that strong epistasis is much more prevalent in the Pol II system than in other
quantified proteins[57,88-91] (**Fig. S3d**), likely due to the high rate of suppressive interactions
stemming from Pol II mutants' opposing effects on catalysis.

Mapping TL-internal residue interaction networks suggests widespread epistasis within the TL. To determine TL-internal interaction networks, we selected 2-4 substitutions for each TL residue, representing diverse phenotypes (GOF, LOF, lethal, or unclassified), and combined them with the selected substitutions across all other TL positions. This curated set of 3790 double mutants captures potential interactions between any two TL residue positions (**Fig. 5a**). Observed fitness of these double mutants was compared to predictions from the additive model, revealing widespread deviation from the model (i.e. large interactions)(r^2=0.21), much smaller than the r^2 (about 0.65-0.75) reported in other studies[57,73,88,89,91] (**Fig. 5b**), likely reflecting our selection criteria for mutations with diverse phenotypes. In general, mutations of the same class were additive or exacerbating, with some exceptions that reveal functional dependencies between residues (**Fig. 5c, 5e, Extended Data Fig. 9**). Additionally, the strongest deviations from expectations were between mutations of opposing phenotypic classes (GOF+LOF combinations). Here, commonly we observed suppression as predicted for mutations acting independently in 43% of GOF+LOF combinations but we also observed unexpected lethality or exacerbation (sign epistasis) for 41% of GOF+LOF combinations. Distinct patterns of sign epistasis for GOF/LOF combinations provide the ability to distinguish functional requirements for GOF alleles to be able to confer GOF phenotypes (discussed below). Interactions were distributed throughout the TL, covering every residue, supporting connectivity across the TL (**Fig. 5d**). Notably, epistasis clustered within the C-terminal TL helix and adjacent regions (The far-right panel of **Fig. 5d**), supporting functional dependencies of TL-C terminal residues on one another, consistent with their proposed function in collaboratively stabilizing the open TL.

Genetic interactions reveal insights into lethal or unclassified individual mutants. Most lethal mutants could be suppressed by at least one predicted GOF mutant (**Fig. 5f, Extended Data Fig. 9**), suggesting their lethality likely comes from reduced activity (LOF) below a viable threshold. However, this suppressibility also suggests that individual lethal substitution activity defects are mostly close to the viable threshold. This suggests that the contributions of most individual residues represent only a portion of total TL activity, which is expected to provide a ~1000 fold increase in catalytic activity[92-94]. Suppressibility of lethal residues by GOF alleles is also consistent with the greater probability of lethal substitutions being LOF rather than GOF. However, two lethal mutations at A1076 were suppressed by most LOF mutations or specific lethal mutants, but not GOF mutants, implying that their lethality resulted from being GOF. Additionally, unclassified single mutants generally showed limited interactions with GOF, LOF, or lethal classes, consistent with small or mostly independent effects. However, a few unclassified mutants showed suppression in combination with GOF mutants, suggesting they are either potential atypical LOF not detected by phenotypic analysis, or allele-specific interactions (**Fig. 5g, Extended Data Fig. 9**).

We reasoned that evolutionarily observed lethal substitutions would be closer to viability than those not observed and would therefore be more likely to be suppressible by Pol II GOF mutants. To determine this, we compared the suppressibility of substitutions observed in existing msRNAPs that are lethal in *S. cerevisiae* Pol II versus substitutions that are lethal but not present in extant msRNAPs by analyzing the maximum positive deviation scores from all double mutants containing individually lethal substitutions. Singly-lethal substitutions (single substitutions that are lethal on their own in yeast Pol II) identified in our msRNAP TL MSA showed higher maximum deviation scores than substitutions lethal on their own in yeast and not present in the MSA (**Fig. 6a, Extended Data Fig. 5b**). Consistently, most L1081 substitutions cannot be suppressed, whereas a L1081M, (commonly a WT residue in bacteria and archaeal

RNAPs) showed the highest suppressibility (**Fig. 6b**). These results indicate that singly lethal
mutants in *S. cerevisiae* Pol II observed elsewhere msRNAP evolution, on average, maintain
greater functionality and/or are more easily suppressed by single additional mutants in yeast Pol
II. Interestingly, most lethal substitutions in A1087 and all lethal substitutions in G1088 cannot
be suppressed (**Fig. 6b**), suggesting the strict restriction at these positions, consistent with the
structural observation that A1087 and G1088 are in a small pocket between the bridge and
funnel helices[12,17]. The TL has been estimated to enhance catalytic activity by 500-1000
fold[92,93,95], while yeast viability tolerates only ~10-fold effects[36]. We conclude that lethal mutants
observed as functional residues in other species are likely closer to the viability threshold, as
might be predicted to result from a series of small adaptive steps that allow their function.

Co-evolution of Pol II active site residues form distinct sectors within specific domains
and extend to the surfaces of Rpb1 and Rpb2. Our analyses suggest that even a highly
conserved domain like the Pol II TL can be tightly integrated through coevolution of less
conserved residues and that substitution tolerance is shaped by widespread epistasis. Changes
in residue identities alter interaction networks, shaping the Pol II active site across evolution. To
identify the coevolving residue networks in Pol II active site we employed statistical coupling
analysis (SCA) for residues within Rpb1 and Rpb2, the two largest subunits of Pol II, which
include the entire active site. SCA is well suited for identifying coupled residue sectors,
representing putative residue communication networks[96-98]. From 283 linked Rpb1/Rpb2
sequences, we identified 53 coevolving sectors (**Fig. S4**). Residues within the active site form
four major sectors (**Fig. 7a-e**). First, sector 37 includes many Rpb2 link domain residues and
most TL residues, including those at the NIR and some at the TL C-terminal (**Fig. 7b**). This
sector tightly surrounds the substrate binding pocket (**Extended Data Fig. 10a**), befitting critical
roles in catalysis. Second, sector 15 consists of four BH residues, most residues in switch 1
(part of the clamp domain), and many cleft residues, including the α46-47 loop, which
accommodates incoming DNA[8,13] (**Fig. 7c**). Residues in this sector directly interact with Pol II
subunits Rpb5 and Rpb6 (**Extended Data Fig. 10b**), consistent with Rpb5's role in interacting
with the incoming DNA[99]. While the α46-47 loop forms a five-helix bundle with two TL helices
and the BH helix[42] (**Fig. 4c**), coupling analysis indicates it is primarily evolutionarily coupled with
cleft residues instead of the TL. This suggests that the α46-47 loop may influence TL function by
constraining the space for TL movement, rather than directly interacting through specific residue
identities. Third, sector 1 involves active site residues surrounding the Mg^{2+} at the catalytic site
and most F-loop residues (**Fig. 7d**). This sector forms a potential communication path that
wraps around the TL and BH, connects to the BH through two BH residues, and extends to
Rpb9, itself a subunit that forms important inputs to the TL and is required for transcription
fidelity[50,100,101]. This pathway likely mediates communication between Rpb9 and the Pol II active
site (**Extended Data Fig. 10c**). Fourth, sector 2 contains most BH residues are coupled with the
Rpb2 Fork and hybrid binding domain, forming a network linking the BH to Rpb2 residues (**Fig.**
7e). This network connects to Rpb5, which interacts with the downstream DNA, Rpb6, a shared
subunit in three eukaryotic msRNAPs, Rpb9, and Rpb3 and Rpb10, which function in Pol II
assembly[102] (**Extended Data Fig. 10d**), representing a residue communication path potentially
tied to various transcription activities.

Notably, coupling is not restricted to residues that are within or near the active site. Some
sectors extend from the active site to the surface of Rpb1/Rpb2 (**Fig. S5a-f**), suggesting that
active site function, potentially through the TL, could be modulated by allosteric interactions
through the binding of transcription factors. Residues at interfaces with elongation factors can

be coupled with those factors yet some residues at interfaces can be coupled with residues
distal to the interface, suggesting potential pathways for allosteric interactions. Examples of
these are shown for TFIIS (**Fig. S6**), LEO1 (Leo1), ELOF1 (Elf1), and SUPT5H (Spt5) (**Fig. S7**).
Together, our findings highlight potential communication networks through evolutionary coupling
that may underlie regulation paths from Pol II surface to active site.

DISCUSSION

How individual mutants alter a protein's function is not necessarily straightforward at the
mechanistic level. Amino acid substitutions both remove functionality of the WT residue but
replace that functionality with something different. By altering the local environment within a
protein or potentially propagating effects to distant locations, each substitution potentially can be
quite different. These differences may not be apparent as phenotypic outputs, and phenotypic
assays may not have granularity to distinguish different biophysical behaviors if they result in
similar outputs. For Pol II mutants, even high-resolution phenotypic analyses, such as gene
expression profiling or genetic interaction profiling between Pol II mutants and deletions in other
yeast genes[53], suggest that LOF and GOF mutants represent a continuum of defects that match
enzymatic activity in vitro. Therefore, these profiles also appear dependent on the output of Pol
II activity defects and can't distinguish potential differences in underlying mechanism.

Through systematic detection of genetic interactions within the Pol II active site, we have
identified functional relationships between amino acids across the TL and between TL
substitutions and others. In the absence of double mutant epistasis analyses it would not be
possible to differentiate similar alleles from one another. L1101S and E1103G, for example, are
two GOF alleles very close to each other in Pol II structure and confer similar phenotypic
landscapes across various growth conditions. Here, we find that their distinct interactions
support that substitutions at 1101 and 1103 target distinct residue networks (**Fig. 3c, 4a,
Extended Data Fig. 6, Fig. S2**). L1101 functions in the TL hydrophobic pocket while E1103
interacts and co-evolves with a number of TL external residues that together support
interactions that maintain the open TL conformation. We also observed connections between TL
C-terminal residues that suggest a limit to how disruptions to structure there can alter Pol II
activity (**Fig. 5d**). Helix-disrupting LOF proline substitutions in at least two TL positions showed
epistasis with multiple substitutions in the back of the TL (1094-1098), suggesting that their
functions require TL C-terminal helix structure and in the absence of that structure (proline
disruption) effects are no longer additive.

The strongest epistatic interactions were observed between two pairs of hydrophobic residues,
A1076 and L1101, and M1079 and G1097 (**Fig. 4**), suggesting their interdependence in
maintaining TL open conformation stability[42]. Considering the TL's multiple active site roles, the
observed interaction networks may also contribute to other NAC steps or off pathway steps (*eg.*
pausing or backtracking). Structural studies have proposed that these hydrophobic residues
form a pocket stabilizing the TL by bundling two TL helices with the BH and α46-47 loop in a
five-helix bundle[28,42]. However, our coupling analysis suggests that α46-47 loop is evolutionarily
coupled with cleft residues rather than the TL, indicating the α46-47 loop may indirectly impact
TL function by limiting the space available for TL movement, rather than through direct
interactions with TL residues (**Fig. 7c**).

Elongation factors bind Pol II and alter its activity, but the mechanisms by which they do so are
not known[103,104]. We observed a high level of genetic interactions between residues outside the

TL and residues within it, including allele-specific reshaping of TL mutant space upon single substitution outside the TL (**Fig. 3**). The fact that minor mutational changes outside the TL can apparently functionally perturb the TL would be consistent with the idea that minor alterations to Pol II structure upon elongation factor binding could easily propagate into the active site via the TL or the BH. As an example, human Rtf1 has been observed to project a domain into the Pol II structure adjacent to the BH (in yeast, this region is occupied instead by Rpb2[105]). These contacts have been proposed to alter Pol II activity. We would propose that the paths for such alteration to activity may follow the coupling sectors we have observed by SCA (**Fig. 7**).

How different individual substitutions are under the surface is critical for understanding plasticity in protein mechanisms and how they might be altered by evolutionary change. A key open question in nucleic acid polymerase mechanisms is the paths for protons in the reaction (for example, deprotonation of the synthesized strand 3′-OH and protonation of pyrophosphate leaving group, for example[82,83,85,87,106-108]). For msRNAPs, the association with the incoming NTP by a universally conserved histidine led to the proposal that this residue might donate a proton during the reaction [12,84,106]. Some substitutions at this position can provide minimal essential function (e.g. tyrosine, arginine), while others are only moderately defective (glutamine). Surprisingly, we found that H1085L was very-well tolerated for growth [17] and the Landick lab has proposed this substitution supports catalysis through positional but not chemical effects[86,87]. Our studies here were quite surprising in that they indicated that L1085 Pol II has unique behavior when perturbed by all possible TL substitutions and is entirely distinct from H1085Y (where we have direct observations of all possible intra-TL doubles) or H1085A or H1085Q (curated doubles) (**Extended Data Fig. 8c, Fig. S8**). These residue specific behaviors suggest that each substitution may have different properties, and compatibility with function may not necessarily represent similar function under the surface.

Evolutionary change over time can alter protein function but it can also alter protein functional plasticity. Recent work from the Thornton lab elegantly demonstrates that phenotypes of substitutions to residues conserved over hundreds of millions of years can change over evolutionary time and can do so unpredictably and transiently during evolution[62]. msRNAPs have structures and functions conserved over billions of years, and deep within their active sites is a mobile domain, the TL, that has large functional constraints on its sequence. The TL sequence must be able to fold into multiple states and maintain recognition of the same substrates across evolutionary space and is shows high identity even between distantly related species. Here we show that the TL, and likely the entire Pol II active site, exhibits a great amount of plasticity through non-conserved positions that are essential for compatibility of the TL and surrounding domains. A limitation of our study is its reliance primarily on genetic and statistical coupling analysis. Future biochemical and biophysical studies will give mechanistic insight into the activity defects underlying the functional connections we observe, for example connecting kinetic properties or structural states to specific mutants. Nonetheless, our results illustrating widespread epistasis and allele-specific effects of single and double mutants predict that comparative analyses among Pol I, II, and III will reveal shared but also enzyme-specific mechanisms due to higher order epistasis shaping the functions of conserved residues.

ACKNOWLEDGMENTS

We thank Dr. Anne-Ruxandra Carvunis (U. Pittsburgh) and Dr. Steve Lockless (Texas A&M) for discussions and advice. We thank Zhizhen Wang and Muyao Lin from the Pitt Statistical Consulting Center for their advice on checking the reproducibility of our data. We acknowledge

funding from NIH R01GM097260 for initiation of this project and NIH R35GM144116 for this
work. This research was supported in part by the University of Pittsburgh Center for Research
Computing, RRID:SCR_022735, through the resources provided. Specifically, this work used
the HTC cluster, which is supported by NIH award number S10OD028483.

AUTHOR CONTRIBUTIONS

BD performed sequencing library construction, data analysis, and made figures, drafted and
revised the manuscript. CQ designed the mutant libraries, performed screening experiments,
and contributed to revising the manuscript. SHZ provided allele counts from raw sequencing
data. CDK conceived the project, guided analyses and interpretation of data, provided funding,
and revised the manuscript.

COMPETING INTERESTS

The authors declare no conflict of interest.

 FIGURES AND FIGURE LEGENDS

Figure 1. Schematics of the Pol II active site interaction landscape. a. The Pol II active site
is embedded in the center of a 12-subunit complex (left panel). Pol II functions are supported by
distinct TL conformational states. An open TL (PDB: 5C4X)[42] and closed TL (PDB: 2E2H)[12]

conformations are shown in the middle panel. GOF mutations have been identified in the TL and
its proximal domains (right panel), suggesting TL mobility and function may be impacted by
adjacent residues. **b.** Examples of inter-residue genetic interactions. WT residues are shown in
grey circles with number indicating residue position in Rpb1. Mutant substitutions are shown in
colored circles, with color representing mutant class. Colored lines between mutant substitutions
represent types of genetic interactions. **c.** Overview of experimental approach. We synthesized
10 libraries of TL variants represented by colored stars. Libraries were transformed into WT or
mutated yeast strains. A selection assay was subsequently performed by scraping and replating
the transformants onto different media for phenotyping. DNA was extracted from yeast from all
conditions, and went through TL region amplification, and Illumina sequencing. Read counts for
variants on general conditions were used to determine growth fitness, while read counts on
other conditions were used to determine the phenotypic fitness landscape (see **Methods**). **d.**
Overview of analytical approach for determining interaction landscape. Mutant conditional
growth fitnesses were calculated using allele frequencies under selective growth conditions and
subjected to two logistic regression models for classification/prediction of catalytic defects.
Double mutant interactions were computed using growth fitness. Classification allowed epistatic
interactions to be deduced from double mutant growth fitness (see **Methods**).

531
Figure 2. Contextual epistasis shapes TL evolution. a. Schematic for the TL evolutionary
haplotypes library. We selected 662 TL haplotypes representing TL alleles from bacterial,
archaeal and the three conserved eukaryotic msRNAPs. These TL alleles were transformed into
yeast and were phenotyped under selective conditions. **b.** Fitness of evolutionarily observed TL
haplotypes in the yeast Pol II background. The Pol II WT TL fitness (0) is labeled as dotted line.
Kruskal-Wallis test was performed for comparison and significant levels ($P < 0.05$) were labeled.

Figure 3. Pol II TL interaction landscape distinguishes mutants with similar phenotypes.

a. Design of the targeted double mutant libraries. All possible substitutions at each TL residue (represented with a simplified format in the left panel) and twelve "probe" mutations (eight within the TL and four in TL-proximal domains) (middle panel) were combined with to generate 7280 double mutants (right panel). 7276 mutants passed the reproducibility filter and were used for interaction analyses. **b.** The percentage of functional interactions observed for each probe mutant with viable GOF or LOF TL substitutions. Epistasis and sign epistasis are labeled with colored lines. **c.** Pol II-TL functional interaction landscape with interactions represented by deviation scores. The upper panel shows interactions of GOF probe mutants in combination of viable GOF or LOF TL substitutions. The lower panel shows interactions of combinations with LOF probe mutants. **d**. Principal component analysis (PCA) of deviation scores across double mutant interactions for 12 probe mutants (see **Methods**).

Figure 4. Pol II TL interaction landscape reveals functional dependency of proximal residues. a-b. Specific epistatic interactions observed between hydrophobic residues A1076 and L1101 (A), and M1079 and G1097 are shown as heatmaps (B). The x-axis of both heatmaps are 20 substitutions ordered by predicted phenotypic classes, and the color of substitution represents the phenotypic class of the substitution. GOF substitution is in green, LOF is in blue, unclassified is in gray, and lethal (fitness < -6.5) is in black. **c.** The epistatic interactions we identified between A1076 and L1101, together with M1079 and G1097 are shown on the five-helix bundle of Pol II active site (PDB: 5C4X)[42].

Figure 5. Widespread epistasis in the Pol II TL interaction landscape. a. Design of the pairwise double mutant library. We curated 2-4 substitutions for each TL residue (in total 90 substitutions, n(GOF) = 18, n(LOF) = 30, n(Unclassified) = 19, n(Lethal) = 23) and combined

them with each other to generate double mutants. 3910 double mutants representing
combinations between any two TL residues were measured and 3790 of them passed the
reproducibility filter. WT TL residue positions are indicated with magenta arch. Phenotype
classes of single substitutions are shown as colored circles (GOF in green, LOF in blue) while
unclassified mutants are in grey and lethal mutants are in black. **b.** An xy-plot of observed
double mutant growth fitness measured in our experiment (Y-axis) and expected fitness from
the addition of two constituent single mutants' fitnesses (X-axis). N (positive) = 612. N
(Negative) = 1402. N (Additive) = 1776. N (Sum)=3790. Lethal threshold (-6.5) is labeled with
dotted lines on X and Y axis. The additive line where $X \pm 1 = Y$ is indicated with dashed line.
Simple linear regression was performed, and the best fit equation is $Y = 0.52X - 2.55$, $r^2 = 0.21$,
$P < 0.0001$. **c.** Percent of interactions observed from each combination group. N (LOF/LOF) =
412. N (GOF/GOF) = 156. N (GOF/LOF) = 534. Epistasis and sign epistasis are indicated with
colored lines. **d.** Various groups of interactions are displayed in a network format for clarity. In
these networks, the TL structure is simplified to a circle, with the TL-N terminal and C-terminal
of the TL connected. WT TL residues are represented by magenta arches, and selected
substitutions are labeled beside these arches. The color of the label represents the phenotype
of each substitution. A colored line connects two substitutions if there is interaction between
them, with the line color representing the interaction type. For example, in the "Epistasis and
sign epistasis" network, we selected three substitutions at K1102: K1102P, K1102L and
K1102A. They show epistasis with V1094D and R, T1095P, S1096P, V1098R and P1099F and
V. These interactions are labeled with red lines. The observed epistasis, indicated by red lines,
accumulated at the TL-C terminal, indicating a functional dependency in this region. **e-g.** The
intra-TL functional interaction heatmaps of various combinations. Double mutant deviation
scores are shown in the heatmap. Annotations at the top and left indicate the curated single
mutants and their predicted phenotypic classes from multiple logistic regression modeling.
GOF/GOF, LOF/LOF, and GOF/LOF combinations are shown in **e**. Combinations with lethal
single substitutions are in F. Combinations with unclassified mutants are in **g**.

Figure 6. Contextual epistasis shapes TL evolution. a. A comparison of the maximum deviation score of each TL lethal single substitution that is present in any evolutionary TL haplotypes from bacterial, archaeal or eukaryotic Pols versus those that have not been observed in any species. The evolutionary TL haplotypes were from multiple sequence alignments (MSA). 9 substitutions were found in an MSA of 542 archaeal TL sequences that are lethal when present in yeast as a single substitution. 17 were found in an MSA of 1403 bacterial TLs, 5 were found in 749 Pol I TLs, 7 were found in 499 Pol II TLs, and 5 were found in 539 Pol III TLs. Evolutionarily observed lethal substitutions were compared to those unobserved in our TL MSA. The percentage of total suppressible lethal single mutants for each group is labeled at the bottom of the plot. Boxes are: center line, median; box limits, second and third quartiles; whiskers, maximum and minimum points. Statistical comparison was done with the Mann-Whitney test and the significant levels (P < 0.05) are shown in the figure. **b.** A heatmap displays the highest deviation scores of lethal single substitutions. The x-axis represents the wild-type residues and positions of the yeast Pol II TL, while the y-axis shows 20 substitutions. Viable

606 substitutions are marked with gray boxes, and wild-type amino acids are indicated with gray
607 boxes with black edge. The highest deviation scores of lethal substitutions are shown in colored
608 boxes, ranging from white to orange, reflecting the strength of suppression.

609
Figure 7. Ultra-conserved TL co-evolves with Pol II residues through diverse pathways. a.
Structure representations of Pol II active site. Pol II subunits are showed with surface in different
colors**.** The active site domains are in cartoon format. Rpb1 active site residues 346-375 and
436-508. Rpb1 TL, 1076-1106. Rpb1 BH, 810-846. Rpb1 α20-21 loop, 672-737. Rpb1 F loop,
747-810. Rpb1 α46-47 loop, 1339-1378. Rpb1 switch 1, 1385-1407. Rpb1 clamp, 1-95, 235-
346, and 1395-1436. Rpb2 link domain, 757-776. Rpb2 Fork, 465-547. Rpb2 hybrid binding,
750-852. Rpb2 wall, 852-973. Rpb2 protrusion, 44-218. Rpb2 lobe, 218-405. **b-e**. Four major
coevolution sectors of Pol II active site residues were identified through statistical coupling
analysis. The residues within each active site domain and the total number of residues in each
sector are labeled. These sectors are mapped on the yeast Pol II Rpb1/Rpb2 structure (PDB:
5C4X)[42].

621
Extended Data Fig.1 Pol II deep mutational scanning is highly reproducible. a. Single
mutant growth fitness from mutants in libraries constructed from synthesized oligos correlated

well with our previous library constructed by a random building block approach when plating
conditions were the same. Qiu *et al*[17] plated at a lower density (colony plating) that we
speculated added noise to the analysis. When plating densely ("dense" and "lawn" conditions)
our new and old libraries showed highly reproducible fitness determinations for single mutants.
b-c. Biological replicates for each library showed high reproducibility for all conditions. Pearson
correlation of each library was calculated with three replicates for viable mutant fitness on all
selective conditions.

631
Extended Data Fig.2 Detection of functional interactions by deviation score. For a pseudo
double mutant *ab*, the difference between its observed fitness (*ab*) and expected fitness (*ab*)
adding the fitness of two constituent single mutants (*a* and *b*) determines the type of interaction
between the two mutants. Positive or negative interactions are determined if the deviation score
is greater than 1 or smaller than –1. Specific epistatic interactions are further distinguished from
general suppression or synthetic sick or lethal interactions using predicted mutant catalytic
defect classes (GOF or LOF).

639
Extended Data Fig.3 Classification of mutant catalytic defects with machine learning

algorithms. a. ROC curves for two multiple logistic regression models used to determine
mutant catalytic class. Using 65 mutants with validated in vitro determined catalytic defects and
conditional growth fitness measured in our experiment, we trained two models to classify
variants as GOF or LOF. The GOF AUROC is 0.9889 (P ≤ 0.0001), whereas the LOF ROC is
0.9914 (P ≤ 0.0001). The predicted vs. observed graphs display the predicted probability of 65
known mutants would be GOF or LOF. The threshold we used to determine GOF or LOF
mutations is shown by lines at 0.75. Details of the models are in Supplemental Table 6. Among
the 6054 viable mutants, 1390 were classified as GOF (22.96%), 1702 as LOF (28.11%), and
2962 remained unclassified (48.93%). **b.** Left: t-SNE projection of all mutants (n=15174) with
perplexity = 50. Right: k-means cluster of all mutants with 20 clusters. The t-SNE and k-means
projections suggest GOF are in 3 clusters (cluster 2, 14, and 16), LOF are in 2 clusters (cluster
3 and 18), and unclassified mutants are in 2 clusters (11 and 15). Most ultra-sick/lethal mutants
(fitness <= -6.5) are projected together into 13 clusters, likely due to significant noise from low
read counts across conditions. **c.** Feature plot of viable mutations in t-SNE and k-means
projections (n=6054). Ultra-sick/lethal mutations were removed and the viable mutants were
projected with t-SNE (perplexity = 100) and K-means (10 clusters). GOF were grouped into 4
clusters (4, 5, 7 and 10) and LOF were in 4 clusters (1, 3, 6, and 9). Each spot in the projection
represents a mutant and it is colored based on the fitness of the mutant in selective conditions.
GOF and LOF mutants in different clusters are related to various phenotype patterns. GOF
clusters 7 and 10 are defined by strong MPA^S, while clusters 4 and 5 show slight MPA^S, Gal^R,
Mn^S, but strong Lys$^+$. Slight Form^S is a common feature across four GOF clusters. LOF clusters
3 and 6 show slight Mn^R, while clusters 1 and 9 are strongly Mn^R and Gal^R. There are three
common features in all LOF clusters: MPA^R, Form^S, and Lys$^-$. Cluster 8, which mostly contain
unclassified mutants, appear defined by Gal super sensitivity, indicating a potential specific
defect defining this cluster.

Extended Data Fig.4 Genetic interactions detected with spot assay and patch assay are consistent with high-throughput results. To verify the genetic interactions observed in high throughput experiment, we picked 50 mutants to measure the genetic interactions with spot assay (**a**) or patch assay (**b**). They are consistent. The fitness, deviation scores and corresponding genetic interactions measured in high throughput experiment are indicated in **a.** Specifically, the labeled observed fitness indicates observed general growth fitness that was measured on SC-Leu+5FOA.The names of GOF mutants are labeled with green color, LOF mutants are labeled with blue, ultra-sick to lethal mutants are labeled with black color.

Extended Data Fig.5 Contextual epistasis affects fitness of TL haplotypes. a. Distributions of deviation scores of the TL haplotypes in each group. **b.** Comparison of the mean deviation scores of lethal single substitutions that are present in different species and those that are absent in any species. Standard deviation values are also shown in the bar plot. ANOVA

multiple comparison was applied to compare the mean deviation score of the "Absent" group to
each of the other groups and significant levels (P < 0.05) are shown in the figure. **c.** An xy-plot
of evolutionary observed TL haplotypes fitness versus the numbers of substitutions in the
haplotypes. Simple linear regression was performed for each plot. Bacteria fitness vs count: Y =
0.004267*X – 8.660, r^2=2.152e-005. Archaea fitness vs count: Y = -0.3406*X - 4.175, r^2=0.1568.
Pol I fitness vs count: Y = -0.7818*X + 1.235, r^2=0.1521. Pol II fitness vs count: Y = -0.3943*X –
1.132, r^2=0.06535. Pol III fitness vs count: Y = -0.4148*X – 3.468, r^2=0.06984. The P values of
the slopes are labeled.

688
Extended Data Fig.6 The functional interaction landscape of probe mutants. The functional
interaction landscape is shown as a heatmap. The upper part of the heatmap shows all Pol II TL
single mutant growth fitness profiling across several phenotypes and the single mutants were
ordered by hierarchical clustering with Euclidean distance. The lower part of the heatmap shows
double mutant deviation scores where a colored block at the interaction of x and y coordinates
indicates deviation score of the double mutant.

Extended Data Fig.7 Allele-specific interactions. Unique interactions observed between TL
substitutions and probe mutants. For each substitution, we analyzed the interquartile range
(IQR) of their deviation scores with all probe mutants. Any substitution with deviation score(s)
outside of the IQR were extracted and called as unique interaction(s). 127 substitutions with
unique interactions with probe mutants were found out of 620 and are shown in the heatmap.
The heatmap was hierarchical clustered with Euclidean distance for both rows and columns.

All   **Positive**   **Epistatic**

a

S713P

n=304/620   n=96/620   n=38/620

T834P

n=376/620   n=33/620   n=102/620

Kruskal-Wallis test
P < 0.0001

b

E1103G

n=395/600   n=184/600   n=35/600

F1084I

n=334/600   n=43/600   n=58/600

Kruskal-Wallis test
P < 0.0001

c

H1085Y

n=447/600   n=85/600   n=18/600

H1085L

n=315/600   n=32/600   n=56/600

Kruskal-Wallis test
P < 0.0001

● GOF
● LOF
● No obvious phenotype
● Lethal
— TL position
— Positive interactions
— Negative interactions
— Synthetic lethal
— Epistasis
— Sign epistasis

Extended Data Fig.8 Interaction networks of selected probe mutants. The TL is shown in
circle with WT residues and positions labeled. All 20 substitutions of each TL residue are
represented by a magenta arc under each WT residue, with tick marks representing individual
substitutions at that position and are colored by mutant class. Comparison of interaction
networks between S713P and T834P (**a**), E1103G and F1084I (**b**) and H1085Y and H1085L (**c**)
showed the differences are significant (P < 0.0001). The comparisons were performed with
Kruskal-Wallis test with P-value correction with Dunn's multiple comparisons test (Supplemental
Table 7).

711
Extended Data Fig.9 The intra-TL functional interaction landscape. The intra-TL functional
interaction landscape is shown as a heatmap. Annotations at the top and right indicate the 90
curated single mutants and their predicted phenotypic classes from multiple logistic regression
modeling. The upper part of the heatmap shows single mutant growth fitness profiling across
multiple phenotypes ordered by groups predicted with logistic regression models. The lower part
of the heatmap shows double mutant deviation scores where a colored block at the interaction
of x and y coordinates indicates deviation score of the double mutant.

719
Extended Data Fig.10 Coevolutionary sectors of Pol II active site residues extend to the
surface of Pol II, forming connections with other Pol II subunits. a. The coevolutionary
sector of Rpb1 TL and Rpb2 link domain residues surrounds the substrate ATP, indicating its
potential role in catalysis**.** The sector is mapped on the yeast Pol II structure with TL in open
state (PDB: 5C4X) and closed state (PDB: 8U9R). **b**. Residues in sector 15 connect with Rpb5
and Rpb6. **c.** Residues on sector 1 connect with Mg^{2+} and Rpb9. **d.** Residues in sector 2
connect with Rpb5, Rpb6, Rpb3, Rpb9 and Rpb10. These sectors are mapped on the yeast Pol
II structure (PDB: 5C4X)[42].

METHODS

Design and Synthesis of TL mutant libraries. We updated and extended the fitness dataset
of Qiu *et al*[17]. sing a similar methodology, but with adjusted conditions and a second-generation
mutant library strategy to generate a complete Pol II TL mutation-phenotype map and examine
genetic interactions. Mutants were constructed by oligo synthesis (Agilent) and screened for

phenotypes previously established as informative for Pol II mutant biochemical defects.
Programmed oligonucleotide library pools included all 620 single TL residue substitutions and
deletions for Rpb1 amino acids 1076-1106 (Library 1), 3914 pairwise double substitutions
(Library 2), 4800 targeted double substitutions (Library 6), and 3373 multiple substitutions
(Library 3-5), along with the WT *S. cerevisiae* Pol II TL allele at a level of ~15% of the total
variants, enabling precise quantification (see Supplemental Table 1, 3). Each synthesized
region contained a mutated or WT Pol II TL sequence and two flanking regions at the 5′ and 3'
ends of the TL-encoding sequence. These flanking regions also contained designed "PCR
handle" (20bp) sequences, allowing distinct subsets of oligos to be amplified from synthesized
pools using selected primers for PCR, and additional flanking WT Pol II sequences allow for
further extension of homology arms by PCR "sewing" (Details are in Supplemental Method 3).

Introduction of Libraries into yeast. Synthesized mutant pools were transformed into yeast
(CKY283) along with an *RPB1*-encoding plasmid where the TL-encoding sequence was
replaced with an *Mlu*I restriction site for linearization as described in Qiu et al[17]. This strategy
allows construction of *rpb1* mutant libraries by gap repair between library fragments and the
linearized vector. Briefly, the synthesized oligo pools were amplified by limited cycles of
emulsion PCR to limit template switching. Extension of flanking homology arms of ~200 bp were
added by PCR sewing. Amplified TL sequences with extended flanking regions were co-
transformed with linearized pRS315-derived *CEN LEU2* plasmid (pCK892) into CKY283,
allowing gap repair via homologous flanking regions. In our experience phenotypes of integrated
alleles and single copy plasmids are qualitatively very similar but this has not been
systematically compared quantitatively. To detect potential residue-residue interactions between
the TL and TL-proximal domains including the Rpb1 Bridge Helix (BH), Funnel Helix alpha-21
and Rpb2, the Pol II TL single mutant pool (Library 1, 620 mutant alleles and 111 WT alleles)
was co-transformed individually with gapped plasmids encoding an additional *rpb1* allele T834P
(Library 7), T834A (Library 8), or Funnel Helix alpha-21 S713P (Library 10)) into CKY283
respectively, or with the gapped WT *RPB1* plasmid into a strain with the genomic mutation, *rpb2*
Y769F (Library 9). These co-transformations created double mutants between the TL and TL-
proximal mutants. The WT allele in single mutant pool represented the single probe mutant due
to substitutions outside the TL on the plasmid or in the strain background. To distinguish
between a fully WT TL and a WT TL representing the TL of a mutant allele elsewhere, a WT Pol
II TL allele with a silent mutant at T1083 (WT codon ACC was replaced with ACT) was co-
transformed with plasmid containing gapped WT *RPB1* in a WT strain in parallel. 15% of the
transformants with silent mutation were mixed with transformants of double mutants. The silent
mutation allowed us to distinguish the WT and the single mutants.

Phenotyping libraries with different conditions. Each transformation was done in three
biological replicates. After transformation, Leu⁺ colonies were collected from **SC-Leu** plates and
replated on **SC-Leu+5FOA** to select for cells having lost the *RPB1 URA3* plasmid and to
measure growth defects. 5-FOA-resistant colonies were then scraped and replated on different
conditions to assess functional differences between WT, GOF, and LOF mutants[17,37]. **SC-Leu +**
20 µg/L MPA (Fisher Scientific) evaluates transcription start site (TSS) shifting. MPA depletes
GTP levels in yeast. To generate GTP, *IMD2* expression is required, but its promoter has
multiple transcription start sites (TSS)[109,110]. Transcripts from the downstream TSS express
functional *IMD2* upon MPA treatment. WT and LOF mutants are resistant to MPA due to their
ability to use downstream TSSs for *IMD2* expression, while GOF mutants are sensitive as they
can only use upstream TSS sand fail to express functional *IMD2*[53,77]. **SC-Leu + 15 mM Mn**

(Sigma): Mn^{2+} compromises transcription fidelity[111] and likely also increases Pol II catalysis;
while WT and LOF mutants are resistant, GOF mutants are sensitive either due to decreased
fidelity or exacerbated catalytic increase. **YPRaf** and **YPRafGal**: the *gal10Δ56* allele
compromised *GAL10* 3'-end formation and polyadenylation, leading to transcription read-
through that inhibits the downstream *GAL7* gene [112,113]. The transcription readthrough in WT
gal10Δ56 cells leads to accumulation of a toxic galactose metabolite because of defects in
GAL7 expression. Some GOF and most LOF mutants show varying galactose resistance, likely
through increasing termination in the *GAL10-GAL7* intergenic region or otherwise reducing
read-through interference with *GAL7*. YPRaf serves as a control for YPRafGal, to demonstrate
that the defect on YPRafGal is galactose toxicity. **SC-Lys** tests lysine auxotrophy. The *lys2-*
122∂ allele in the strain background has a *Ty1* transposable element insertion, which prevents
normal expression of *LYS2*[114]. WT and LOF mutants are Lys⁻, while GOF mutants utilize a
normally silent promoter within the *Ty1 ∂* element insertion, becoming Lys⁺. **SC-Leu + 3%**
formamide (JT Baker), formamide destabilizes protein hydrogen bonds[115], with most mutants
showing resistance except for slight sensitivity in LOF mutants. Details of cell numbers plated
on each plate and screening time of each plate are in Supplemental Table 5. Details of the high
efficiency transformation protocol are in Supplemental Method 1.

Generation of libraries for quantification by amplicon sequencing. Genomic DNA of each
screened library was extracted using the Yeastar genomic DNA kit according to manufacturer's
instructions (Zymo Research). To ensure adequate DNA for sequencing, the TL regions of all
libraries were amplified with PCR cycles that were verified to be in the linear range by qPCR to
minimize disturbance of allele distributions, and under emulsion PCR conditions (EURx
Micellula DNA Emulsion & Purification (ePCR) PCR kit) to limit template switching. Details are in
Supplemental Method 2 and 3. To multiplex samples, we employed a dual indexing strategy
wherein 10 initial barcodes for differentiating 10 mutant libraries were added during the initial
amplification using 10 pairs of custom primers. In a second amplification, 28 primers containing
28 NEB indices were used to add a second index for distinguishing conditions and replicates
(NEBNext Multiplex Oligos for Illumina) (see Supplemental Table 4). As a result, a sample-
specific barcodes were present for each set of variants. The indexed, pooled samples were
sequenced by single end sequencing on an Illumina Next-Seq (150nt reads). On average, over
11 million reads were obtained for individual samples with high reproducibility from two rounds
of sequencing.

Data cleaning and fitness calculation and normalization. Reads of mutants were sorted into
appropriate libraries and conditions by detecting particular indices after sequencing. Read
counts were estimated by a codon-based alignment algorithm to distinguish reads that exactly
matched designated codons of mutants [116]. To clean the data, mutant reads with coefficients of
variation greater than 0.5 in the control condition (SC-Leu) were excluded from the analysis.
The mutant read count was increased by 1 to calculate the allele frequency under different
conditions. To measure and compare the phenotypes of all mutants, mutant phenotypic score
(fitness) was calculated by allele frequency change of a mutant under selective conditions
relative to the unselective condition comparing to the frequency change of WT. The formula for
calculating fitness is shown below.

Fitness (mut) = log [$f^{mut, sele}$ / $f^{mut, unsele}$ - log [$f^{WT, sele}$ / $f^{WT, unsele}$]

We applied min-max normalization to bring the median growth fitness of mutants measured at
ten libraries to the same level for direct comparison (formula is shown below). In each library,

we divided mutants into several groups based on their allele counts on the control condition.
Mutants with read count differences of less than 10 are present in one group. The WT growth
fitness was set as the maximum value and the minimum fitness in each group was the
minimum. Min-max normalization was used to equalize the growth fitness into the same range
between various groups inside each library. Additionally, we utilized min-max normalization to
level the mutant fitness across all ten libraries with WT fitness as Max and minimal fitness in
each library as the minimum. As a result, mutant growth fitness was scaled to one range and
could be used to determine genetic interactions.

832
$$X' = \frac{X - Xmin}{Xmax - Xmin}$$

Determination of functional interactions. The genetic interactions between single
substitutions were determined by comparing the multiple-substitution mutant normalized median
fitness to the log additive of the single substitution normalized median fitness. The simplified
formula is as follows:

Deviation score $(M_1M_2M_3)$ = Fitness $(M_1M_2M_3)$ – [Fitness (M_1) + Fitness (M_1) + Fitness (M_3)]

(1). -1 < Deviation score < 1, the interaction among the constituent single mutants is additive
and mutants are acting independently.

(2). Deviation score $\geq$ 1, the interaction is non-additive and is positive, including suppression
and epistatic interactions.

(3). Deviation score $\leq$ -1. the interaction is non-additive and is negative, including synthetic sick,
synthetic lethal, and sign epistasis interactions.

Any mutation with fitness smaller than the lethal threshold (-6.50) was classified as an ultra-sick/
lethal mutant and its fitness was normalized to -6.50 for calculation of the deviation score.
Synthetic sickness and synthetic lethality were distinguished by whether a double mutant is
viable or lethal (fitness is greater than or equals to the lethal threshold -6.5) when two
constituent mutations are viable. Synthetic lethality can be further classified into two types.
First, additive-synthetic lethality was determined when the expected double mutant fitness
calculated by additive model was lethal (expected fitness = -6.5) and the observed double
mutant fitness was also lethal (fitness = -6.5) (in this case the deviation score = 0). Second, the
beyond-additive synthetic lethality was determined when the expected double mutant was viable
(expected fitness > -6.5) while the observed double mutant fitness was lethal (fitness = -6.5) (in
this case the deviation score < 0). To separate these two situations in our figures, we labeled
additive synthetic lethality as black and beyond-additive synthetic lethality as purple. Details of
formulas are in Supplemental Method 4.

Mutant classification using two multiple logistic regression models. We trained two
multiple logistic regression models to distinguish GOF and LOF mutants using phenotypic
fitness from SC-Leu+MPA, SC-Lys, and YPRafGal conditions of 65 single mutants. This dataset
included 20 mutants with transcription rates measured by biochemical experiments (13 GOF
and seven LOF)[17,37,53], and 45 additional mutants that were hierarchical clustered with verified
mutants (12 GOF, 26 LOF, 1 WT, and 6 that were not GOF or LOF mutants)[17]. Both models
incorporated intercept, main effects, and two-way interactions were involved in defining both
models, with a cutoff threshold of 0.75 applied to classify mutants in both of the GOF and LOF
models.

Model for predicting the probability of a mutant being a GOF:

y
$= \dfrac{1}{1 + e\textasciicircum(1.816 + 2.542 * fMPA - 1.942 * fLys + 0.06566 * fGal - 0.5297 * fMPA * fLys - 0.08373 * fMPA * fGal + 0.02556 * fLys * fGal)}$

Model for predicting the probability of a mutant being LOF:
y
$= \dfrac{1}{1 + e\textasciicircum(1.916 - 1.392 * fMPA - 1.328 * fLys - 0.8353 * fGal - 0.01112 * fMPA * fLys - 0.2992 * fMPA * fGal + 0.8823 * fLys * fGal)}$

Both models showed accuracy, with the area under ROC close to one **(Extended Data Fig.**
2A). The details are provided in Supplemental Table 6.

Principal component analysis (PCA). Deviation scores of curated and probe double mutants
were analyzed in PCA. The scripts used R language v4.0.3 (https://www.R-project.org/) with R
packages tidyverse v1.3.1 (https://www.tidyverse.org), prompt (stats v3.6.2
(https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/prcomp)), ggplot2 v3.3.3
(https://ggplot2.tidyverse.org), dplyr v1.0.6 (https://dplyr.tidyverse.org), and missMDA v1.18
(https://dplyr.tidyverse.org).

t-SNE projection. Allele frequencies for all mutants in nine conditions with three replicates were
analyzed by t-SNE (Perplexity = 50) or k-means (clusters =20). Thirteen clusters with ultra-sick
to lethal mutants as majority were eliminated. The remaining mutants were analyzed again with
t-SNE (Perplexity = 100) and k-means (cluster =10). The scripts utilized R language v4.0.3
(https://www.R-project.org/), along with R packages Rtsne v0.15
(https://github.com/jkrijthe/Rtsne), ggplot2 v3.3.3 (https://ggplot2.tidyverse.org), k-means (stats
v3.6.2 (https://www.rdocumentation.org/packages/stats/versions/3.6.2/topics/kmeans)).

Statistical coupling analysis. 5787 eukaryotic homologous sequences of yeast Rpb1 from a
published multiple sequence alignment (MSA) and 4000 Rpb2 sequences from NCBI nr
database were cleaned and used in the statistical coupling analysis[87]. Sequence identity was
reducing to 90% with T-coffee package v12.00.7fb08c2[117]. Pol I, II, and III sequences were
separated based on a phylogenetic tree constructed with FastTree 2[118]. Finally, 283 matched
Pol II Rpb1 and Rpb2 sequences, with each matched pair sharing the same species name,
were connected with 10 Ns and aligned with T-coffee. This newly generated MSA was used for
statistical coupling analysis with the python-based package pySCA v6.1[97]. The scripts were
adapted from https://github.com/ranganathanlab/pySCA.

Supplemental Tables

Supplemental Table 1. Library summary

Supplemental Table 2. Double mutant interactions observed in high throughput system and
growth assays in published papers

Supplemental Table 3. Strains and plasmids

Supplemental Table 4. Primers

Supplemental Table 5. Phenotyping details

Supplemental Table 6. MLR_models_summary

Supplemental Table 7. Kruskal-Wallis tests

Supplemental methods

Supplemental Method 1. High efficiency large scale chemical yeast transformation protocol.

Supplemental Method 2. Emulsion PCR set up with EURx Micellula DNA Emulsion &
Purification (ePCR) PCR kit.

Supplemental Method 3. Amplification/transformation/screening of mutant libraries and
sequencing pool preparation.

Supplemental Method 4. Formulas of calculating functional interactions.

DATA AVAILIABILITY

Raw sequencing data has been deposited on the NCBI SRA (Sequence Read Archive)
database under BioProject PRJNA948661. Processed mutant count, fitness and processing
codes are available through GitHub (https://github.com/Kaplan-Lab-Pitt/TLs_Screening.git).
Strains and plasmids will be provided upon request. Source data are provided with this paper.

CODE AVAILIABILITY

The codes for calculating deviation scores, PCA, t-SNE projection, statistical coupling analysis,
and generating figures are available in GitHub (https://github.com/Kaplan-Lab-
Pitt/TLs_Screening.git).

REFERENCES

1. Cramer, P. Multisubunit RNA polymerases. *Curr Opin Struct Biol* **12**, 89-97 (2002).
2. Werner, F. & Grohmann, D. Evolution of multisubunit RNA polymerases in the three
domains of life. *Nat Rev Microbiol* **9**, 85-98 (2011).
3. Allison, L.A., Moyle, M., Shales, M. & Ingles, C.J. Extensive homology among the largest
subunits of eukaryotic and prokaryotic RNA polymerases. *Cell* **42**, 599-610 (1985).
4. Zhang, G. et al. Crystal Structure of Thermus aquaticus Core RNA Polymerase at 3.3 Å
Resolution. *Cell* **98**, 811-824 (1999).
5. Vassylyev, D.G. et al. Crystal structure of a bacterial RNA polymerase holoenzyme at
2.6 A resolution. *Nature* **417**, 712-9 (2002).
6. Hirata, A., Klein, B.J. & Murakami, K.S. The X-ray crystal structure of RNA polymerase
from Archaea. *Nature* **451**, 851-4 (2008).
7. Gnatt, A.L., Cramer, P., Fu, J., Bushnell, D.A. & Kornberg, R.D. Structural basis of
transcription: an RNA polymerase II elongation complex at 3.3 A resolution. *Science*
292, 1876-82 (2001).
8. Cramer, P., Bushnell, D.A. & Kornberg, R.D. Structural basis of transcription: RNA
polymerase II at 2.8 angstrom resolution. *Science* **292**, 1863-76 (2001).
9. Fernandez-Tornero, C. et al. Crystal structure of the 14-subunit RNA polymerase I.
Nature **502**, 644-9 (2013).
10. Hoffmann, N.A. et al. Molecular structures of unbound and transcribing RNA polymerase
III. *Nature* **528**, 231-6 (2015).
11. Malinen, A.M. et al. Active site opening and closure control translocation of multisubunit
RNA polymerase. *Nucleic Acids Res* **40**, 7442-51 (2012).

12. Wang, D., Bushnell, D.A., Westover, K.D., Kaplan, C.D. & Kornberg, R.D. Structural
basis of transcription: role of the trigger loop in substrate specificity and catalysis. *Cell*
127, 941-54 (2006).
13. Kaplan, C.D. Basic mechanisms of RNA polymerase II activity and alteration of gene
expression in Saccharomyces cerevisiae. *Biochimica Et Biophysica Acta-Gene*
Regulatory Mechanisms **1829**, 39-54 (2013).
14. Dangkulwanich, M. et al. Complete dissection of transcription elongation reveals slow
translocation of RNA polymerase II in a linear ratchet mechanism. *Elife* **2**, e00971
(2013).
15. Bar-Nahum, G. et al. A ratchet mechanism of transcription elongation and its control.
Cell **120**, 183-93 (2005).
16. Weinzierl, R.O. The nucleotide addition cycle of RNA polymerase is controlled by two
molecular hinges in the Bridge Helix domain. *BMC Biol* **8**, 134 (2010).
17. Qiu, C. et al. High-Resolution Phenotypic Landscape of the RNA Polymerase II Trigger
Loop. *PLoS Genet* **12**, e1006321 (2016).
18. Da, L.T. et al. Bridge helix bending promotes RNA polymerase II backtracking through a
critical and conserved threonine residue. *Nat Commun* **7**, 11244 (2016).
19. Mazumder, A., Lin, M., Kapanidis, A.N. & Ebright, R.H. Closing and opening of the RNA
polymerase trigger loop. *Proc Natl Acad Sci U S A* **117**, 15642-15649 (2020).
20. Wang, B., Predeus, A.V., Burton, Z.F. & Feig, M. Energetic and structural details of the
trigger-loop closing transition in RNA polymerase II. *Biophys J* **105**, 767-75 (2013).
21. Larson, M.H. et al. Trigger loop dynamics mediate the balance between the
transcriptional fidelity and speed of RNA polymerase II. *Proc Natl Acad Sci U S A* **109**,
6555-60 (2012).
22. Fouqueau, T., Zeller, M.E., Cheung, A.C., Cramer, P. & Thomm, M. The RNA
polymerase trigger loop functions in all three phases of the transcription cycle. *Nucleic*
Acids Res **41**, 7048-59 (2013).
23. Kireeva, M.L. et al. Transient reversal of RNA polymerase II active site closing controls
fidelity of transcription elongation. *Mol Cell* **30**, 557-66 (2008).
24. Liu, X., Bushnell, D.A. & Kornberg, R.D. RNA polymerase II transcription: structure and
mechanism. *Biochim Biophys Acta* **1829**, 2-8 (2013).
25. Kaplan, C.D. & Kornberg, R.D. A bridge to transcription by RNA polymerase. *J Biol* **7**, 39
(2008).
26. Tan, L., Wiesler, S., Trzaska, D., Carney, H.C. & Weinzierl, R.O. Bridge helix and trigger
loop perturbations generate superactive RNA polymerases. *J Biol* **7**, 40 (2008).
27. Silva, D.A. et al. Millisecond dynamics of RNA polymerase II translocation at atomic
resolution. *Proc Natl Acad Sci U S A* **111**, 7665-70 (2014).
28. Lin, G. et al. Structural basis of transcription: RNA polymerase II substrate binding and
metal coordination using a free-electron laser. *Proc Natl Acad Sci U S A* **121**,
e2318527121 (2024).
29. Fong, N. et al. Pre-mRNA splicing is facilitated by an optimal RNA polymerase II
elongation rate. *Genes Dev* **28**, 2663-76 (2014).
30. Xu, L. et al. Dissecting the chemical interactions and substrate structural signatures
governing RNA polymerase II trigger loop closure by synthetic nucleic acid analogues.
Nucleic Acids Res **42**, 5863-70 (2014).
31. Vassylyev, D.G. et al. Structural basis for substrate loading in bacterial RNA
polymerase. *Nature* **448**, 163-8 (2007).
32. Da, L.T., Wang, D. & Huang, X. Dynamics of pyrophosphate ion release and its coupled
trigger loop motion from closed to open state in RNA polymerase II. *J Am Chem Soc*
134, 2399-406 (2012).

33. Liu, B., Zuo, Y. & Steitz, T.A. Structures of E. coli sigmaS-transcription initiation
complexes provide new insights into polymerase mechanism. *Proc Natl Acad Sci U S A*
113, 4051-6 (2016).

34. Seibold, S.A. et al. Conformational coupling, bridge helix dynamics and active site
dehydration in catalysis by RNA polymerase. *Biochim Biophys Acta* **1799**, 575-87
(2010).

35. Malagon, F. et al. Mutations in the Saccharomyces cerevisiae RPB1 gene conferring
hypersensitivity to 6-azauracil. *Genetics* **172**, 2201-9 (2006).

36. Kaplan, C.D., Larsson, K.M. & Kornberg, R.D. The RNA polymerase II trigger loop
functions in substrate selection and is directly targeted by alpha-amanitin. *Mol Cell* **30**,
547-56 (2008).

37. Kaplan, C.D., Jin, H., Zhang, I.L. & Belyanin, A. Dissection of Pol II trigger loop function
and Pol II activity-dependent control of start site selection in vivo. *PLoS Genet* **8**,
e1002627 (2012).

38. Matthew H. Larsona, Jing Zhoub,1, Craig D. Kaplanc,1, Murali Palangatd,2, Roger D.
Kornberge, Robert Landickd, and Steven M. Blocka,b,f,3. Trigger loop dynamics mediate
the balance between the transcriptional fidelity and speed of RNA polymerase II. (2012).

39. Kireeva, M.L. et al. Molecular dynamics and mutational analysis of the catalytic and
translocation cycle of RNA polymerase. *BMC Biophys* **5**, 11 (2012).

40. Windgassen, T.A. et al. Trigger-helix folding pathway and SI3 mediate catalysis and
hairpin-stabilized pausing by Escherichia coli RNA polymerase. *Nucleic Acids Res* **42**,
12707-21 (2014).

41. Nayak, D., Voss, M., Windgassen, T., Mooney, R.A. & Landick, R. Cys-pair reporters
detect a constrained trigger loop in a paused RNA polymerase. *Mol Cell* **50**, 882-93
(2013).

42. Barnes, C.O. et al. Crystal Structure of a Transcribing RNA Polymerase II Complex
Reveals a Complete Transcription Bubble. *Mol Cell* **59**, 258-69 (2015).

43. Cheung, A.C. & Cramer, P. Structural basis of RNA polymerase II backtracking, arrest
and reactivation. *Nature* **471**, 249-53 (2011).

44. Sydow, J.F. et al. Structural basis of transcription: mismatch-specific fidelity mechanisms
and paused RNA polymerase II with frayed RNA. *Mol Cell* **34**, 710-21 (2009).

45. Sekine, S., Murayama, Y., Svetlov, V., Nudler, E. & Yokoyama, S. The ratcheted and
ratchetable structural states of RNA polymerase underlie multiple transcriptional
functions. *Mol Cell* **57**, 408-21 (2015).

46. Kuldell, J.C. & Kaplan, C.D. RNA Polymerase II Activity Control of Gene Expression and
Involvement in Disease. *Journal of Molecular Biology* (2024).

47. Kettenberger, H., Armache, K.J. & Cramer, P. Complete RNA polymerase II elongation
complex structure and its interactions with NTP and TFIIS. *Mol Cell* **16**, 955-65 (2004).

48. Lennon, C.W. et al. Direct interactions between the coiled-coil tip of DksA and the trigger
loop of RNA polymerase mediate transcriptional regulation. *Genes Dev* **26**, 2634-46
(2012).

49. Hein, P.P. et al. RNA polymerase pausing and nascent-RNA structure formation are
linked through clamp-domain movement. *Nat Struct Mol Biol* **21**, 794-802 (2014).

50. Kaster, B.C., Knippa, K.C., Kaplan, C.D. & Peterson, D.O. RNA Polymerase II Trigger
Loop Mobility: INDIRECT EFFECTS OF Rpb9. *J Biol Chem* **291**, 14883-95 (2016).

51. Taatjes, A.C.S.a.D.J. Structure and mechanism of the RNA polymerase II transcription
machinery. (2020).

52. Leng, X.Y. et al. Organismal benefits of transcription speed control at gene boundaries.
Embo Reports **21**(2020).

53. Braberg, H. et al. From structure to systems: high-resolution, quantitative genetic
analysis of RNA polymerase II. *Cell* **154**, 775-88 (2013).

54. Tesileanu, T., Colwell, L.J. & Leibler, S. Protein sectors: statistical coupling analysis
versus conservation. *PLoS Comput Biol* **11**, e1004091 (2015).
55. Phillips, P.C. The language of gene interaction. *Genetics* **149**, 1167-71 (1998).
56. Breen, M.S., Kemena, C., Vlasov, P.K., Notredame, C. & Kondrashov, F.A. Epistasis as
the primary factor in molecular evolution. *Nature* **490**, 535-8 (2012).
57. Starr, T.N. & Thornton, J.W. Epistasis in protein evolution. *Protein Sci* **25**, 1204-18
(2016).
58. Ortlund, E.A., Bridgham, J.T., Redinbo, M.R. & Thornton, J.W. Crystal structure of an
ancient protein: evolution by conformational epistasis. *Science* **317**, 1544-8 (2007).
59. Karageorgi, M. et al. Genome editing retraces the evolution of toxin resistance in the
monarch butterfly. *Nature* **574**, 409-412 (2019).
60. Phillips, P.C. Epistasis--the essential role of gene interactions in the structure and
evolution of genetic systems. *Nat Rev Genet* **9**, 855-67 (2008).
61. Faure, A.J. et al. Mapping the energetic and allosteric landscapes of protein binding
domains. *Nature* **604**, 175-183 (2022).
62. Park, Y., Metzger, B.P.H. & Thornton, J.W. Epistatic drift causes gradual decay of
predictability in protein evolution. *Science* **376**, 823-830 (2022).
63. Ding, D. et al. Co-evolution of interacting proteins through non-contacting and non-
specific mutations. *Nat Ecol Evol* **6**, 590-603 (2022).
64. Kondrashov, A.S., Sunyaev, S. & Kondrashov, F.A. Dobzhansky-Muller incompatibilities
in protein evolution. *Proc Natl Acad Sci U S A* **99**, 14878-83 (2002).
65. Lunzer, M., Golding, G.B. & Dean, A.M. Pervasive cryptic epistasis in molecular
evolution. *PLoS Genet* **6**, e1001162 (2010).
66. Natarajan, C. et al. Epistasis among adaptive mutations in deer mouse hemoglobin.
Science **340**, 1324-7 (2013).
67. Doud, M.B., Ashenberg, O. & Bloom, J.D. Site-Specific Amino Acid Preferences Are
Mostly Conserved in Two Closely Related Protein Homologs. *Mol Biol Evol* **32**, 2944-60
(2015).
68. Starr, T.N. et al. Shifting mutational constraints in the SARS-CoV-2 receptor-binding
domain during viral evolution. *Science* **377**, 420-424 (2022).
69. Haddox, H.K., Dingens, A.S., Hilton, S.K., Overbaugh, J. & Bloom, J.D. Mapping
mutational effects along the evolutionary landscape of HIV envelope. *Elife* **7**(2018).
70. Viktorovskaya, O.V. et al. Divergent contributions of conserved active site residues to
transcription by eukaryotic RNA polymerases I and II. *Cell Rep* **4**, 974-84 (2013).
71. Mani, R., St Onge, R.P., Hartman, J.L.t., Giaever, G. & Roth, F.P. Defining genetic
interaction. *Proc Natl Acad Sci U S A* **105**, 3461-6 (2008).
72. Qiu, C. & Kaplan, C.D. Functional assays for transcription mechanisms in high-
throughput. *Methods* **159-160**, 115-123 (2019).
73. Lin, X. et al. Nested epistasis enhancer networks for robust genome regulation. *Science*
377, 1077-1085 (2022).
74. Fowler, D.M. & Fields, S. Deep mutational scanning: a new style of protein science. *Nat*
Methods **11**, 801-7 (2014).
75. Sergey Ioffe, C.S. Batch Normalization: Accelerating Deep Network Training by
Reducing Internal Covariate Shift. *arXiv:1502.03167v3* (2015).
76. Hill, W.G., Goddard, M.E. & Visscher, P.M. Data and theory point to mainly additive
genetic variance for complex traits. *PLoS Genet* **4**, e1000008 (2008).
77. Malik, I., Qiu, C., Snavely, T. & Kaplan, C.D. Wide-ranging and unexpected
consequences of altered Pol II catalytic activity in vivo. *Nucleic Acids Research* **45**,
4431-4451 (2017).
78. (!!! INVALID CITATION !!! 17,37).

79. Scull, C.E., Ingram, Z.M., Lucius, A.L. & Schneider, D.A. A Novel Assay for RNA
Polymerase I Transcription Elongation Sheds Light on the Evolutionary Divergence of
Eukaryotic RNA Polymerases. *Biochemistry* **58**, 2116-2124 (2019).
80. Duan, B., Qiu, C., Lockless, S.W., Sze, S.H. & Kaplan, C.D. Higher-order epistasis
within Pol II trigger loop haplotypes. *Genetics* **228**(2024).
81. Brueckner, F. & Cramer, P. Structural basis of transcription inhibition by alpha-amanitin
and implications for RNA polymerase II translocation. *Nat Struct Mol Biol* **15**, 811-8
(2008).
82. Carvalho, A.T., Fernandes, P.A. & Ramos, M.J. The Catalytic Mechanism of RNA
Polymerase II. *J Chem Theory Comput* **7**, 1177-88 (2011).
83. Huang, X. et al. RNA polymerase II trigger loop residues stabilize and position the
incoming nucleotide triphosphate in transcription. *Proc Natl Acad Sci U S A* **107**, 15745-
50 (2010).
84. Castro, C. et al. Nucleic acid polymerases use a general acid for nucleotidyl transfer. *Nat
Struct Mol Biol* **16**, 212-8 (2009).
85. Unarta, I.C., Goonetilleke, E.C., Wang, D. & Huang, X. Nucleotide addition and cleavage
by RNA polymerase II: Coordination of two catalytic reactions using a single active site.
J Biol Chem **299**, 102844 (2023).
86. Mishanina, T.V., Palo, M.Z., Nayak, D., Mooney, R.A. & Landick, R. Trigger loop of RNA
polymerase is a positional, not acid-base, catalyst for both transcription and
proofreading. *Proc Natl Acad Sci U S A* **114**, E5103-E5112 (2017).
87. Palo, M.Z., Zhu, J., Mishanina, T.V. & Landick, R. Conserved Trigger Loop Histidine of
RNA Polymerase II Functions as a Positional Catalyst Primarily through Steric Effects.
Biochemistry **60**, 3323-3336 (2021).
88. Araya, C.L. et al. A fundamental protein property, thermodynamic stability, revealed
solely from large-scale measurements of protein function. *Proc Natl Acad Sci U S A* **109**,
16858-63 (2012).
89. Melamed, D., Young, D.L., Gamble, C.E., Miller, C.R. & Fields, S. Deep mutational
scanning of an RRM domain of the Saccharomyces cerevisiae poly(A)-binding protein.
RNA **19**, 1537-51 (2013).
90. Harms, M.J. & Thornton, J.W. Analyzing protein structure and function using ancestral
gene reconstruction. *Curr Opin Struct Biol* **20**, 360-6 (2010).
91. Fowler, D.M. et al. High-resolution mapping of protein sequence-function relationships.
Nat Methods **7**, 741-6 (2010).
92. Toulokhonov, I., Zhang, J., Palangat, M. & Landick, R. A central role of the RNA
polymerase trigger loop in active-site rearrangement during transcriptional pausing. *Mol
Cell* **27**, 406-19 (2007).
93. Yuzenkova, Y. et al. Stepwise mechanism for transcription fidelity. *BMC Biol* **8**, 54
(2010).
94. Diss, G. & Lehner, B. The genetic landscape of a physical interaction. *Elife* **7**(2018).
95. Wang, W., Walmacq, C., Chong, J., Kashlev, M. & Wang, D. Structural basis of
transcriptional stalling and bypass of abasic DNA lesion by RNA polymerase II. *Proc Natl
Acad Sci U S A* **115**, E2538-E2545 (2018).
96. Halabi, N., Rivoire, O., Leibler, S. & Ranganathan, R. Protein sectors: evolutionary units
of three-dimensional structure. *Cell* **138**, 774-86 (2009).
97. Rivoire, O., Reynolds, K.A. & Ranganathan, R. Evolution-Based Functional
Decomposition of Proteins. *PLoS Comput Biol* **12**, e1004817 (2016).
98. Salinas, V.H. & Ranganathan, R. Coevolution-based inference of amino acid interactions
underlying protein function. *Elife* **7**(2018).
99. Cramer, P. Structure and function of RNA polymerase II. *Adv Protein Chem* **67**, 1-42
(2004).

100. Walmacq, C. et al. Rpb9 subunit controls transcription fidelity by delaying NTP
sequestration in RNA polymerase II. *J Biol Chem* **284**, 19601-12 (2009).
101. Ziegler, L.M., Khaperskyy, D.A., Ammerman, M.L. & Ponticelli, A.S. Yeast RNA
polymerase II lacking the Rpb9 subunit is impaired for interaction with transcription factor
IIF. *J Biol Chem* **278**, 48950-6 (2003).
102. Nouraini, S., Archambault, J. & Friesen, J.D. Rpo26p, a subunit common to yeast RNA
polymerases, is essential for the assembly of RNA polymerases I and II and for the
stability of the largest subunits of these enzymes. *Mol Cell Biol* **16**, 5985-96 (1996).
103. Cramer, P. Organization and regulation of gene transcription. *Nature* **573**, 45-54 (2019).
104. Schier, A.C. & Taatjes, D.J. Structure and mechanism of the RNA polymerase II
transcription machinery. *Genes Dev* **34**, 465-488 (2020).
105. Vos, S.M., Farnung, L., Linden, A., Urlaub, H. & Cramer, P. Structure of complete Pol II-
DSIF-PAF-SPT6 transcription complex reveals RTF1 allosteric activation. *Nat Struct Mol*
Biol **27**, 668-677 (2020).
106. Castro, C. et al. Two proton transfers in the transition state for nucleotidyl transfer
catalyzed by RNA- and DNA-dependent RNA and DNA polymerases. *Proc Natl Acad Sci*
U S A **104**, 4267-72 (2007).
107. Gregory, M.T., Gao, Y., Cui, Q. & Yang, W. Multiple deprotonation paths of the
nucleophile 3'-OH in the DNA synthesis reaction. *Proc Natl Acad Sci U S A* **118**(2021).
108. Belogurov, G.A. & Artsimovitch, I. The Mechanisms of Substrate Selection, Catalysis,
and Translocation by the Elongating RNA Polymerase. *J Mol Biol* **431**, 3975-4006
(2019).
109. Jenks, M.H., O'Rourke, T.W. & Reines, D. Properties of an intergenic terminator and
start site switch that regulate IMD2 transcription in yeast. *Mol Cell Biol* **28**, 3883-93
(2008).
110. Kuehner, J.N. & Brow, D.A. Regulation of a eukaryotic gene by GTP-dependent start site
selection and transcription attenuation. *Mol Cell* **31**, 201-11 (2008).
111. Qiu, C. et al. Thiolutin has complex effects in vivo but is a direct inhibitor of RNA
polymerase II in vitro. *Nucleic Acids Res* **52**, 2546-2564 (2024).
112. Greger, I.H. & Proudfoot, N.J. Poly(A) signals control both transcriptional termination and
initiation between the tandem GAL10 and GAL7 genes of Saccharomyces cerevisiae.
Embo J **17**, 4771-9 (1998).
113. Kaplan, C.D., Holland, M.J. & Winston, F. Interaction between transcription elongation
factors and mRNA 3'-end formation at the Saccharomyces cerevisiae GAL10-GAL7
locus. *J Biol Chem* **280**, 913-22 (2005).
114. Simchen, G., Winston, F., Styles, C.A. & Fink, G.R. Ty-mediated gene expression of the
LYS2 and HIS4 genes of Saccharomyces cerevisiae is controlled by the same SPT
genes. *Proc Natl Acad Sci U S A* **81**, 2431-4 (1984).
115. Aguilera, A. Formamide sensitivity: a novel conditional phenotype in yeast. *Genetics*
136, 87-91 (1994).
116. Sing-Hoi Sze, C.D.K. Codon-Based Sequence Alignment for Mutation Analysis by High-
Throughput Sequencing. in *2018 IEEE 8th International Conference on Computational*
Advances in Bio and Medical Sciences (ICCABS) (2018).
117. Notredame, C., Higgins, D.G. & Heringa, J. T-Coffee: A novel method for fast and
accurate multiple sequence alignment. *J Mol Biol* **302**, 205-17 (2000).
118. Price, M.N., Dehal, P.S. & Arkin, A.P. FastTree 2--approximately maximum-likelihood
trees for large alignments. *PLoS One* **5**, e9490 (2010).

1193