Dissociable contributions of the amygdala and ventral hippocampus to stress-induced changes in defensive behavior

Zachary T. Pennington[1], Alexa R. LaBanca[1], Patlapa Sompolpong[1], Zoe Christenson Wick[1],

Yu Feng[1], Zhe Dong[1], Taylor R. Francisco[1], Lingxuan Chen[1], Sasha L. Fulton[1], Ian Maze[1,2,3],

Tristan Shuman[1], Denise J. Cai[1]

1. Nash Family Department of Neuroscience, Icahn School of Medicine at Mount Sinai
2. Department of Pharmacological Sciences, Icahn School of Medicine at Mount Sinai
3. Howard Hughes Medical Institute, Icahn School of Medicine at Mount Sinai

Correspondence: Denise.Cai@mssm.edu and Zachary.Pennington@mssm.edu

Abstract:

Severe stress can produce multiple persistent changes in defensive behavior. While much is known about the circuits supporting stress-induced associative fear responses, how circuit plasticity supports the broader changes in defensive behavior observed after severe stress remains unclear. Here, we find that stress-induced plasticity in the ventral hippocampus (vHC) and basolateral amygdala (BLA) support doubly dissociable defensive behavioral changes. Stress-induced protein synthesis in the BLA was found to support lasting enhancements in stress sensitivity but not enhancements in exploratory anxiety-related behaviors, whereas protein synthesis in the vHC was found to support enhancements in anxiety-related behavior but not enhancements in stress sensitivity. Like protein synthesis, neuronal activity of the BLA and vHC were found to differentially support the expression of these same defensive behaviors. Lastly, blockade of associative fear had no impact on stress-induced changes in anxiety-related behavior. These findings highlight that multiple memory-systems support stress-induced defensive behavior changes.

Keywords:

Basolateral amygdala, ventral hippocampus, stress, fear, anxiety, PTSD, memory, stress-enhanced fear learning, defensive behavior

INTRODUCTION

In immediate response to stressful and life-threatening events, animals display evolutionarily conserved defensive responses, including changes in heart rate and respiration, stress hormone release, as well as the behavioral initiation of fight, flight, and freezing [1-6]. If sufficiently strong, stressful events can also instantiate persistent changes in how animals interact with their environment. Perhaps most extensively studied are associative fear responses, in which animals engage in defensive behaviors such as freezing and/or flight when re-exposed to environmental cues present at the time of the initial stressful experience [1,7-12]. However, after severe stress, animals also display alterations in foraging and exploration in uncertain environments [3,5,13,14] (often referred to as anxiety-related behavior), as well as heightened responses to future stressful events [14-17]. These long-lasting defensive behavioral changes are fundamental to anxiety disorders, which include fear of stress-related cues, heightened stress responses, and reduced environmental engagement; and are frequently predated by the experience of severe stress [18-21].

It is often assumed that many of the defensive behavioral changes observed in the aftermath of stress are fundamentally associative in nature – animals could either be responding to cues that were directly present at the time of stress, or stimuli resembling these cues to some degree (i.e., stimulus generalization) [22-30]. For example, it is well-documented that startle responses are potentiated by the presence of associative fear cues [31,32], suggesting that associative stimuli may drive heightened responses to aversive events after stress. Moreover, it is possible that following stress, alterations in exploration in anxiety-related behavior tests such as the elevated-plus maze could be accounted for by shared features with the environment in which the stressor took place. Lastly, several reports document altered associative fear learning and generalization in humans with anxiety disorders [23,24,33-35]. In light of these findings, broad emphasis has been placed on associative learning processes governing the lasting consequences of stress. However, the explanatory reach of an associative framework has its limits. Pre-weaning rodents incapable of forming associative fear memories have been found to nevertheless display decreased exploration in exploratory anxiety-related behavior tests, as well as heightened responses to subsequent aversive stimuli following stressful experiences [14]. Moreover, extinguishing fear to stress-associated cues does not necessarily mitigate sensitized responses to new stressors [15,36,37]. These findings

highlight the persistence of some stress-induced behavioral phenotypes despite weak

associative fear, indicating a potential dissociation. As such, it could be the case that multiple

memory systems – associative and non-associative – support the enduring consequences of

stress on defensive behavior. However, a direct biological dissociation of such memory

systems has remained elusive. If discovered, this would have broad implications for the

treatment of anxiety disorders, potentially explaining why treatments focused on associative

processes are ineffective in some individuals [38-40].

Here, we explore the contributions of stress-induced plasticity within the ventral

hippocampus (vHC) and basolateral amygdala (BLA) to the enduring impacts of stress on

different defensive behaviors. Neuronal activity within both the BLA and vHC are well known

to regulate defensive behaviors [8,41-48]. However, whether stress-induced plasticity within these

structures act in concert to support a common defensive behavioral process, or whether they

support distinct defensive behavior changes, is unclear. We find that plasticity within each

brain region supports separate defensive behavior changes in response to stress,

demonstrating unique functions of these structures, and supporting the view that multiple

memory systems underly stress-induced defensive behavioral changes.

RESULTS

Acute severe stress produces multiple, lasting, changes in defensive behavior.

We first sought to establish a behavioral protocol in which a single acute stressor produces lasting changes in multiple defensive behaviors, adapting a prior model that has been used extensively in mice and rats [14,15,49-51] (Fig 1A). Animals were placed in a distinct environment where they received 10 foot-shocks during a 10-min period ('trauma', T), or were placed in the same environment but did not receive foot-shocks ('no-trauma', NT). A week later, multiple defensive behaviors were assessed. In the light-dark box, an exploratory test that captures rodents' natural avoidance of well-lit places and is sensitive to anxiolytics [52,53], trauma-exposed animals showed increased anxiety-related behavior, reflected in more time spent in the dark side of the light-dark box (Fig 1C. $t_{47.5}=5.5$, $p<0.001$). To assess associative fear, animals were returned to the trauma environment (trauma recall). As expected, trauma-exposed animals spent large amounts of time freezing (Fig 1D. $t_{40.7}=11.5$, <0.001). Lastly, we assessed the animals' stress sensitivity by placing the animals in a novel environment, in which they showed very little initial freezing at baseline (0.1% baseline freezing in controls vs 1.3% in trauma-exposed; $t_{42.9}=2.8$, $p<0.01$). A loud auditory startle stimulus was then presented (novel stressor). When returned to this environment the next day, trauma-exposed animals showed substantially more freezing (Fig 1E. $t_{47.6}=5.79$, <0.001). Notably, although this phenomenon is often termed stress-enhanced fear learning, learning rate analysis suggests the enhanced learning likely reflects heightened sensitivity to the aversive stimulus rather than an enhanced learning rate (Fig S1).

Next, as a preliminary means of establishing that these defensive behaviors convey information about unique biobehavioral processes, we correlated these phenotypes in a large group of trauma-exposed animals, as high inter-phenotype correlations would suggest shared biological origins. We found minimal to no relationship between behavioral tests (Fig 1F-H. trauma recall and light-dark: $R^2=0.04$, $p=0.19$; trauma recall and novel stressor: $R^2=0.19$, $p<0.01$; light-dark and novel stressor: $R^2=0.13$, $p=0.32$). Though each of these measures is likely subject to imperfect test-retest reliability, these findings nevertheless suggest that these phenotypes may be independently governed.

Figure 1: Acute stress produces multiple, lasting, changes in defensive behavior. A) Animals were exposed to a distinct environment in which they received 10 foot-shocks (trauma, T) or were placed in the sameenvironment and received no foot-shocks (no-trauma, NT). A week later, they were tested in the light-dark test of anxiety-related behavior, associative fear of the trauma environment, and their response to a novel stressor in a new environment. B) Trauma-exposed animals displayed high levels of post-shock freezing during the trauma, C) increased time in the dark in the light-dark test, D) strong associative fear of the trauma environment, and E) increased fear of the novel stressor environment when placed in this environment on the final test day. F-H) Examining a large set of trauma-exposed animals, these post-trauma phenotypes were found to be poorly correlated, suggesting they reflect unique biological processes. NT=23 & T=40 mice. p<.05 (*), p<0.01 (**), p<0.001 (***).

Stress-induced protein synthesis in the BLA and vHC produce distinct changes in defensive behavior.

In order to assess how stress-induced plasticity supports persistent changes in defensive behavior, we utilized post-stress administration of the protein synthesis inhibitor anisomycin, as protein synthesis is known to support the consolidation of many forms of memory and to regulate synaptic plasticity [54-60]. Further, because manipulations of protein synthesis can be done after a learning experience, it provides a means of disrupting the consolidation of an experience without altering its initial encoding. To validate that the emergence of the observed defensive behavioral changes are indeed supported by stress-induced protein synthesis, we first tested the effects of systemically administered anisomycin (Fig 2B-D). Animals underwent trauma and were immediately after given 3 injections of anisomycin or vehicle (at 0, 4, and 8 hours). Alternatively, animals were placed in the same environment but did not receive trauma and were treated with vehicle. A week later, relative to no-trauma controls, trauma-exposed animals treated with vehicle exhibited increases in anxiety-related behavior in the light dark-test (Fig 2B. $t_{12.1}=4.9$, $p<0.001$), associative fear in the trauma recall test (Fig 2C. $t_{8.4}=7.2$, $p<0.001$), and heightened fear of the novel stressor environment (Fig 2D. $t_{14}=2.7$, $p=0.02$). Post-trauma anisomycin administration reduced all of these stress-induced defensive behaviors relative to trauma-exposed animals given vehicle (Fig 2B-D. light-dark test: $t_{16.3}=2.4$, $p=0.031$; trauma recall: $t_{8.4}=6.6$, $p<0.001$; novel stressor: $t_{15.9}=2.9$, $p<.01$).

Next, we assessed the impacts of targeting trauma-induced protein synthesis specifically in the BLA or vHC, regions previously linked to regulating defensive behaviors. Mice had indwelling cannulas implanted above either the BLA or vHC (Fig 2E. See Fig S3 for placement in all animals). After surgical recovery, animals then underwent trauma and immediately thereafter received intracranial infusions of anisomycin or vehicle. Alternatively, they experienced no trauma and were treated immediately with vehicle. Animals treated with vehicle in the BLA or vHC showed no behavioral differences and are collapsed here (Fig S2). Moreover, prior to vehicle/anisomycin treatment, no differences were observed in freezing during the trauma session for animals that underwent trauma (Group: $F_{2,76}=1.1$, $p=0.33$; Group x Time: $F_{20,760}=0.5$, $p=0.96$). As expected, relative to no-trauma controls, trauma-exposed animals treated with vehicle exhibited heightened anxiety-related behavior in the light-dark test

(Fig 2F. $t_{47.5}$=5.5, p<0.001), a strong associative trauma memory (Fig 2G. $t_{40.7}$=11.5, p<0.001),

and heightened fear of the novel stressor (Fig 2H. $t_{47.6}$=5.8, p<0.001). These behaviors were

differentially affected by blocking trauma-induced protein synthesis in the BLA and vHC. In the

light-dark test, anisomycin in the vHC greatly attenuated trauma-induced increases in animals'

preference for the dark (Fig 2F. $t_{28.2}$=3.1, p<0.01). However, anisomycin in the BLA was

without effect ($t_{44.9}$=1.09, p=0.28). In the trauma recall test, anisomycin in either the BLA or

vHC were effective at reducing associative freezing relative to trauma controls (Fig 2G. BLA:

$t_{56.9}$=8.2, p<0.001, vHC: t_{44}=4.2, p<0.001), though the BLA appeared to contribute to a more

sizable degree (Fig 2G. T:BLA-ani vs T:vHC-ani: $t_{30.5}$=2.4, p=0.02). Lastly, anisomycin in the

BLA was able to block the enhanced sensitivity to a novel stressor, whereas anisomycin in the

vHC was without effect (Fig 2H. BLA: $t_{56.4}$=4.1, p<0.001, vHC: $t_{46.2}$=1.6, p=0.11). These

findings highlight that while stress-induced plasticity in the BLA is paramount for associative

fear and heightened stress sensitivity, it is not necessary for alterations in anxiety-related

behavior. Conversely, stress-induced plasticity in the vHC is essential for increased anxiety-

related behavior, and to a lesser extent associative fear recall, but not heightened stress

sensitivity. Importantly, the finding that blockade of protein synthesis in the BLA had a

profound impairment on associative fear for the trauma environment but no detectable effect

on the light-dark test further suggests a dissociation between anxiety-related behavior and

associative fear.

Figure 2: Stress-induced protein synthesis in the BLA and vHC support distinct changes in defensive behavior. A) Immediately after trauma (T) or no-trauma (NT), animals were administered 3 injections of anisomycin (ani) or vehicle (veh). A week later, they were tested in the light-dark test of anxiety-related behavior, associative fear of the trauma environment, and their response to a novel stressor in a new environment. B) Systemic administration of anisomycin after trauma reduced time in the dark side in the light-dark test, C) associative freezing in the trauma environment, D) and reduced freezing in the novel stressor environment when placed in this environment on the final test day. E) Example placement of cannula injectors in the BLA and vHC for intracranial anisomycin infusions. F) Anisomycin in the vHC after trauma reduced subsequent time in the dark side of the light-dark test, whereas anisomycin in the BLA was without effect. G) Anisomycin in the BLA or vHC reduced associative fear of the trauma environment. H) Anisomycin in the BLA reduced freezing in the novel stressor test relative to controls, whereas anisomycin in the vHC was without effect. For systemic injections, NT: veh=7, T:veh=9, and T:ani=10 mice. For intracranial infusions, NT:veh=23, T:veh=40, T:ani-BLA=19 , and T:ani-vHC=20 mice. P<.05 (*), p<0.01 (**), p<0.001 (***).

Distinct stress-induced defensive behaviors require activity of the BLA and vHC

The prior findings indicate that stress-induced plasticity within the BLA and vHC support

the induction of distinct post-stress phenotypes. We next sought to address whether activity

within these structures also differentially contributes to the expression of these behaviors after

trauma. To inhibit neural activity in the BLA or vHC, we utilized the inhibitory ionotropic

designer receptor PSAM4-GlyR (PSAM) in conjunction with its activating ligand uPSEM-817-

tartrate (uPSEM) [61].

To first verify that we could reversibly alter the activity of the BLA and vHC to change

defensive behavior, we used the PSAM-uPSEM system to inactivate the BLA or vHC during

trauma memory recall, as the activity of both regions have been shown to be important for

associative fear memories [62]. A pan-neuronal virus expressing PSAM was infused into either

of these two structures a month prior to behavioral testing (Fig 3A. See Fig S4 for placement).

Then, animals received the trauma previously described and a week later were tested in a

trauma recall test, once after receiving an injection of uPSEM and once after receiving an

injection of vehicle. No differences were observed in freezing behavior during the trauma

experience for animals with the PSAM receptor in the BLA versus the vHC (Region: $F_{1,14}=2$,

p=0.18; Region x Time: $F_{10,140}=0.5$, p=0.91). However, in the recall test, inhibition of either the

BLA or vHC via administration of the agonist uPSEM reduced associative freezing (Fig 3C.

Drug: $F_{1,14=}12.1$, p<0.01; Drug X Region: $F_{1,14}=0.2$, p=0.67).

Next, to assess the contributions of BLA and vHC neural activity to the expression of

enhanced anxiety-related behavior and stress sensitivity after trauma, PSAM-expressing virus

was infused into either of these structures, or animals underwent a control surgery in which

PBS was infused (CTRL) (Fig 3D. See Fig S5 for placement). A month later, all animals

underwent the trauma procedure and were tested for anxiety-related behavior and stress

sensitization. Notably, no behavioral differences were observed during the initial trauma,

suggesting that expression of the receptor alone had no effect on the acquisition or expression

of conditioned freezing (Fig 3E. Group: $F_{2,23}=0.3$, p=0.72; Group x Time: $F_{20,230}=1.6$, p=0.1). A

week later, animals were tested in the light-dark test after administration of uPSEM.

Consistent with our finding that stress-induced protein synthesis within these structures

supports distinct behaviors, inhibition of the vHC produced a dramatic decrease in time spent

on the dark side (Fig 3F. $t_{11.7}=5$, p<0.001), whereas inhibition of the BLA was without effect

(Fig 3F. $t_{16.8}$=0.4, p=0.68). Next, we tested the animals drug-free in a trauma recall session in

order to ensure that prior inactivation during the light-dark test did not influence subsequent

behavior, as well as to further confirm that receptor expression alone did not influence memory

recall. As expected, there was no difference in freezing between groups (Fig 3G. $F_{2,23}$=0.5,

p=0.6). Lastly, we inactivated either the BLA or the vHC via uPSEM administration prior to the

novel stressor. The following day, when animals were returned to the novel stressor

environment, BLA-inactivated animals froze less than controls (Fig 3H. t_{13}=2.4, p=0.04),

whereas vHC inhibition did not alter freezing ($t_{13.8}$=1.3, p=0.23). In summary, inactivation of

the BLA/vHC replicated the effects observed with protein synthesis inhibition, where the BLA

supports heightened stress sensitivity, and the vHC supports enhanced anxiety-related

behavior.

Figure 3. Distinct stress-induced defensive behaviors require activity of the BLA and vHC. A) A pan-neuronal virus expressing the inhibitory ionotropic receptor PSAM4-GlyR (PSAM) was expressed in either the BLA or vHC a month before behavior, or animals underwent a control surgery (Ctrl) in which PBS was infused. B) To assess efficacy of PSAM-mediated neuronal silencing on behavior, a group of animals expressing PSAM in either the BLA or vHC underwent trauma, and week later their associative recall of the trauma environment was tested in the presence or absence of the PSAM ligand uPSEM. C) Inhibition of either the BLA or vHC was able to reduce trauma memory recall. D) In a separate set of animals that underwent trauma, we then tested the effects of inhibiting the BLA or vHC during testing of anxiety-related behavior in the light-dark test and administration of the novel stressor. Animals had PSAM-expressing virus infused into the BLA, vHC, or underwent Ctrl surgery. E) No differences were observed in freezing during trauma, indicating no effect of receptor expression alone. F) Inhibition of the vHC, but not the BLA, was able to attenuate stress-induced increases in time spent in the dark side in the light-dark test. G) In a drug-free test, there were no differences in trauma memory recall. H) Inhibition of the BLA, but not the vHC, during the novel stressor was able to reduce subsequent freezing when placed back in this environment. For 2B/2C, BLA=9 and vHC=7 mice. For 2D-2H, Ctrl=9, PSAM-BLA=10, and PSAM-vHC=7 mice. p<.05 (*), p<0.01 (**), p<0.001 (***).

Activity of the vHC, but not BLA, is necessary for multiple anxiety-related behaviors.

We selected the light-dark test instead of other exploratory measures of anxiety-related

behavior because initial pilot studies suggested behavior in this test was more reliably altered

by prior stress. To assess whether our findings generalize to other measures of exploratory

anxiety-related behavior, PSAM expressing virus was infused into the BLA or vHC a month

before behavioral testing. Animals were then given the trauma and a week later were tested in

the open field, elevated plus maze, and the light-dark tests, once with vehicle and once with

uPSEM for each test (drug and testing order counterbalanced; see Fig S6 for placement).

Again, inhibition of the vHC, but not the BLA, reduced time in the dark side of the light-dark test

(Fig 4B. vHC: $F_{1,7}=14.9$, $p<0.01$, BLA: $F_{1,6}=0.01$, $p=0.92$). Additionally, inhibition of the vHC

reduced time in the closed arms of the elevated plus maze (Fig 4C. $F_{1,7}=55.6$, $p<0.001$). Of

some interest, inhibition of the BLA produced a slight increase in time in the closed arms

($F_{1,6}=10.2$, $p=0.02$), although this effect does not survive multiple comparisons across all

anxiety-related measures. Lastly, neither inhibition of the BLA nor the vHC affected behavior

in the open field test (Fig 4D. Drug: $F_{1,13}=1.7$, $p=0.211$; Drug X Region: $F_{1,13}=0$, $p=0.96$).

These findings confirm that the effects of vHC inhibition generalize to at least one other

commonly used measure of exploratory anxiety-related behavior, and further suggests that

inhibition of the BLA does not alter these measures.

Figure 4. Activity of the vHC, but not BLA, is necessary for multiple anxiety-related behaviors. A) Animals expressing PSAM in either the BLA or vHC were subjected to trauma and then tested in multiple exploratory anxiety-related tests, both with and without uPSEM. B) Inhibition of the vHC reduced time in the dark side in the light-dark test, B) as well as time in the closed arms of the elevated plus maze. C) Neither inhibition of the BLA nor the vHC altered distance from the center in the open field test. BLA=7 and vHC=8 mice. p<.05 (*), p<0.01 (**), p<0.001 (***).

DISCUSSION

A wealth of literature supports the notion that the BLA and vHC regulate defensive

behaviors [8,41-48]. In light of reciprocal connections between these structures [63-65], it would be

reasonable to hypothesize that stress-induced plasticity within the BLA and vHC subserve a

common defensive process (or processes). While the findings presented here do not negate

this possibility, they nevertheless highlight that under certain conditions the BLA and vHC

support dissociable defensive behavioral functions, both at the levels of stress-induced

plasticity and neuronal activity. Stress-induced protein synthesis within the BLA was found to

be critical to stress-evoked enhancements in stress sensitivity, whereas stress-induced protein

synthesis within the vHC had no bearing on this defensive phenotype. Conversely, stress-

induced protein synthesis within the vHC, but not the BLA, was found to support heightened

exploratory anxiety-related behaviors. Suppressing neural activity within the BLA or vHC was

of similar effect to protein synthesis blockade.

Notably, it could be argued that although stress-induced plasticity within each of these

structures supports the induction of distinct defensive phenotypes, connectivity between the

two structures could still be fundamental to the expression of both phenotypes. That is, stress-

induced plasticity within one structure may support behavior changes via its connections with

the other. Our results suggest otherwise. If plasticity within the vHC was supporting increased

anxiety-related behaviors via its connections with the BLA, inhibition of either structure would

be expected to have an effect on anxiety-related behavior. Similarly, if plasticity within the BLA

regulated stress sensitivity through its connections with the vHC, inhibition of the vHC would

be expected to alter this phenotype. However, blockade of activity within each of these

structures produced dissociable impacts on behavior, mirroring the effects of protein synthesis

blockade. As such, these structures appear to contain distinct systems through which stress

produces lasting changes in behavior.

The dissociation of the contributions of the BLA and vHC to the defensive behaviors

studied here is not entirely without precedent, although side by side comparisons of their

functions are limited. For instance, despite the large literature on the role of the BLA in

associative fear learning [9-11,41,42,66-69], several studies have reported that inhibition of the BLA is

without effect on exploratory anxiety-related behaviors [70-73], although discrepancies also exist

[74]. Additionally, a recent report found that optogenetic stimulation of projections from the vHC

to the BLA do not regulate exploratory anxiety-related behavior [75]. This corroborates the

hypothesis that vHC regulates exploratory-anxiety related behavior through its connection with

other down-stream structures such as the hypothalamus [75-77] or medial prefrontal cortex [44].

While stimulation of BLA terminal fibers in the vHC has been found to alter anxiety-related

behavior [71], this may reflect a general effect of exciting the vHC, as opposed to the natural role

served by BLA to vHC efferents. In concert with our findings, these results broadly suggest

that the vHC may regulate exploratory anxiety-related behavior in a manner distinct from its

connections with the BLA.

Relatively few studies have examined how stress sensitivity is altered as a function of

prior experience. However, consistent with our findings, existing work supports a role for the

BLA. Either inactivation of the BLA or blockade of glucocorticoid receptors in the BLA prior to

an acute stressor have been found to reduce subsequent enhancements in responding to an

aversive stimulus [49].   Further, antagonism of ghrelin receptors in the BLA during chronic stress

is able to block subsequently enhanced responses to a new stressor[78]. Our results strengthen

the proposition that plasticity within the BLA supports these changes by utilizing a post-stress

manipulation of anisomycin, which does not interfere with the initial stress experience.   By

contrast, outside of the work presented here, little is known about the contributions of the vHC

to stress-induced changes in stress sensitivity. Thus, the finding that enhancements in stress

sensitivity following aversive experience are independent of plasticity and activity of the vHC is

entirely novel.

Blockade of stress-induced plasticity in the BLA and vHC were both able to impair

associative fear learning for the initial stressor, and blockade of activity within either structure

was similarly able to impair associative memory recall. That said, blockade of stress-induced

plasticity within the BLA had a much more pronounced effect than the same manipulation

performed in the vHC. This discrepancy is perhaps in line with prior reports pointing to a more

equivocal role of the vHC in associative fear. For instance, modulation of vHC granule cells

have been found to modulate anxiety-related behaviors, but not associative fear learning [45];

another report indicates that inactivation of the vHC disrupts the acquisition of auditory, but not

contextual, fear learning [79]; and newer findings suggest vHC to BLA projections are able to

modulate the acquisition of contextual fear, although these effects are relatively small in size

[75,80]. While the precise role of the vHC in associative fear learning is yet to be determined, it is

possible that connections between the vHC and BLA permit modulation of the more heavily
BLA-mediated associative fear learning.

Plasticity within the BLA was necessary for the acquisition and expression of both
heightened associative memories of a stressful event and the heightened sensitivity to novel
stressors after prior stress. Furthermore, these two phenotypes were correlated, albeit weakly.
It may be argued that these defensive phenotypes are one in the same. Prior evidence, as
well as data presented here, stands in opposition to this possibility. First, extinction of the
associative memory for an initial stressor has been found to leave the enhanced response to a
second stressor intact [15,36,37]. Second, early life stress at a time point when rodents are not
able to form associative memories nevertheless leaves animals with heightened responses to
subsequent stressors in adulthood [14]. Third, pharmacological blockade of NMDA receptors
during initial stress that provides near-complete loss of associative memory for that event do
not reduce the heightened responding to subsequent aversive events [15]. Lastly, despite the
finding presented here that inactivation of the BLA and vHC were able to equivalently impair
trauma memory recall, only inactivation of the BLA was able to alter the heightened sensitivity
to a novel stressor. Therefore, despite both phenotypes' dependence upon the BLA,
associative memory for a stressor and the enhanced responding to subsequent stress are
dissociable. It could be that synapse- and ensemble-specific plasticity within the amygdala
supports the associative memory for a specific stressor, whereas a broader form of non-
associative plasticity within the amygdala supports sensitized stress responses. Future
studies need to disentangle how plasticity within the amygdala supports these two forms of
learning.

It is particularly striking that only manipulations of the vHC were able to alter stress
induced-changes in exploratory anxiety-related behaviors despite both the BLA's and vHC's
role in associative fear learning. If exploratory anxiety-related behaviors reflected
generalization of the associative memory for a stressor, as is sometimes assumed, one would
expect that manipulations that affect associative fear memories would also affect anxiety-
related behaviors. Instead, manipulations of the BLA produced a profound impact on
associative fear but no change in exploratory anxiety-related behavior, indicating that these
behaviors are supported by distinct neural processes. Furthermore, we saw no evidence of a
correlation between measures of associative fear for an initial stressor and exploratory anxiety-

related behavior. This is in line with prior reports where near-complete abolition of an

associative fear memory did not reduce stress-induced changes in anxiety-related

behavior[14,15], and indicate that these phenotypes are supported by unique biological

processes.   Here, we extend these results by showing where plasticity underlying stress-

induced changes in exploratory anxiety-related behavior occur, in the vHC.

In closing, these results shed new light on how stress-induced plasticity within the BLA

and vHC support the formation of defensive behavioral phenotypes. Furthermore, they

highlight just how distinct components of memories for stressful events might be. Similar to

the separate memory systems in the brain supporting episodic and procedural learning of the

same event, we find that different defensive behaviors induced by stress are supported by

distinct brain regions. This has important clinical implications for the treatment of anxiety

disorders and other stress-associated mental health conditions. First, it suggests that clinically

targeting one stress-induced defensive behavior, or the circuits that support that behavior, may

leave others wholly unaffected. Perhaps some of the existing gaps in treatment result from a

failure to adequately target the spectrum of defensive processes altered in these conditions.

Second, by understanding the relationship between specific defensive behaviors and their

biology, in combination with knowing the specific defensive behaviors altered in a particular

mental health condition, we may make greater headway in the treatment of these conditions.

For instance, associative learning processes are more likely affected in some mental health

conditions, while anxiety-related behaviors might be more affected in others, and these

differences undoubtedly covary with different neuronal patterns. Indeed, it is known that the

different anxiety disorders are not only symptomatically different, but characterized by unique

brain activity patterns[81]. By understanding these inter-relationships, we may more rapidly find

the appropriate key for each lock as we try to advance treatments for stress-associated mental

health conditions.

ACKNOWLEDGEMENTS

This work was supported by NIMH DP2 MH122399, NIMH R01 MH120162, Brain

Research Foundation Award, Klingenstein-Simons Fellowship, NARSAD Young Investigator

Award, McKnight Memory and Cognitive Disorder Award, One Mind-Otsuka Rising Star

Research Award, Hirschl/Weill-Caulier Award, Mount Sinai Distinguished Scholar Award, and

Friedman Brain Institute Award, to DJC; NIMH K99 MH131792 and BBRF Young Investigator

Award to ZTP; NINDS F32 NS116416 to ZCW; AES Predoctoral Research Fellowship to YF;

and NINDS R01 NS116357 and NIA RF1 AG072497 to TS. The authors would like to thank

Dr. Scott Russo and Dr. Roger Clem for their helpful comments on this work.

AUTHOR CONTRIBUTIONS

ZTP and DJC conceived of the overarching research goals, designed the experiments,

and oversaw the experiments. ZTP analyzed the experimental data and prepared the initial

manuscript. ZTP, ARL, PS, ZCW, YF, ZD, TRF, LC, SLF, IM, TS and DJC contributed to

interpretation of the results and edited the manuscript. ZTP, ALB, PS, ZCP, YF, ZD, TRF, LC,

and SF performed experiments. ZTP and PD designed software for analysis of behavioral

data. DJC, IM, TS, ZTP, ZCW and YF secured funding.

DECLARATION OF INTERESTS

The authors declare no competing interests.

STAR METHODS

Animals*:*

All animals were adult male C57BL/6J mice obtained from Jackson Laboratories, aged

2-6 months. Animals were housed in a temperature- and humidity-controlled vivarium on a

12/12 light-dark cycle (lights on at 7 a.m.). All experimental procedures were approved by the

Icahn School of Medicine at Mount Sinai's IACUC.

Behavioral testing:

For all experiments, animals were singly housed beginning 1 week prior to the start of

behavioral testing and were handled by the experimenters for approximately 1 min/day for 5

days during this time. When systemic injections were to be given, animals were additionally

briefly habituated to restraint 2-3 times. Animals were habituated to being transported from the

vivarium to the laboratory 2-3 times to mitigate transport serving as an associative cue.

Trauma and trauma recall: Animals were transported from the vivarium in their cages on

a cart to the experimental testing room, which was well lit and had a fan providing ambient

sound. Animals were then placed in a brightly lit experimental testing chamber with a grid floor

(Med Associates), scented with 5% Simple Green solution. During trauma, after a 5 min

period of baseline exploration, animals received 10, 1 sec, 1 mA, scrambled foot-shocks, with

an inter-shock interval of 30 sec. Animals were taken out of the testing chamber 30 sec after

the last shock and returned to the vivarium in their home cage. For trauma recall sessions,

animals were transported to the same experimental testing chamber for an 8 min test session.

Novel stressor and novel stressor recall: Animals were transported from the vivarium in

P1000 pipet boxes and carried in a dark box to the experimental testing room, which was dark

except a dim red light. Animals were then placed inside of a dark testing chamber (Med

Associates) with a flat plexiglass floor and a curved back wall. The chamber was scented with

1% acetic acid solution. After a 3 min baseline period, animals were exposed to a single loud

auditory stimulus (3 sec, 130 dB white noise, 0 ms rise time) that was delivered by a speaker

attached to the wall. Animals were removed 10 sec later and returned to the vivarium. For

novel stressor recall sessions, animals were transported to the same experimental testing

chamber for an 8 min test session.

Exploratory anxiety-related tests: The light-dark test was conducted using two
interconnected square compartments with an open top (each compartment measured 7.5 in
width x 11.25 in height), separated by a 1.5 in wide passageway that could be closed with an
opaque sliding divider. One chamber was made of all white acrylic, while the walls of the other
were covered in matte black wallpaper and had a red acrylic floor. Overhead lighting provided
luminance of 50 lux on the light side. After a 1 min baseline period in which animals were
confined to the dark side, the central divider was raised and the animals could freely explore
both sides of the light-dark box. The open field test was conducted in a circular arena (19 in
diameter; 10 in height) made of white acrylic. A circular field was utilized in order to avoid the
need to define arbitrary center/outer areas. Animals were placed along one wall and allowed
to explore for 5 min. Luminance was approximately 50 lux. For the elevated plus maze, each
arm measured 2 3/8 in wide and 13.75 in long. The floor of the maze was made of white
acrylic, and the enclosed arms had black walls (8 in high). Luminance of the open arms was
25 lux. Animals were placed at the end of a closed arm and then allowed to explore freely for
5 min. For all anxiety-related behavior tests, apparatus were cleaned with 70% ethanol
between test sessions and behavior was captured with an overhead webcam. These sessions
were conducted in an otherwise dark room with a fan providing ambient background noise.
Learning Rate Analysis: To examine differences in the acquisition of associative fear
after trauma (Fig S1), animals experienced a 10 foot-shock trauma (1 sec, 1 mA foot-shocks,
distributed pseudo-randomly over the course of an hour), or were placed in the same
environment but received no shocks. 19-20 days later, all animals were then placed in a novel
environment and received one tone-shock pairing per day across 7 days (Tone = 30 sec, 75
dB white noise. Foot-shock = 2 sec, 0.25 mA). A lower amplitude foot-shock was utilized
because it is known to produce lower asymptotic freezing levels [82]. The tone was presented
after 5 min baseline and co-terminated with shock. Data is compared qualitatively to
predictions from the Rescorla-Wager Model [83]. All animals received a 20 minute habituation
session in the novel environment before training began. Additionally, all mice were tested for
trauma recall twice (8 min/test), once the day after trauma and once the day before habituation
to the novel stressor environment. Of note, these mice were initially used for another
experiment designed to assess the impact of trauma on conditioned cocaine place preference.
All mice received 3, 20 mg/kg, intraperitoneal injections of cocaine hydrochloride between

trauma and tone-shock pairings. That said, no differences in preference were found between

groups (data available at github.com/ZachPenn/BLAvHC_Dissoc).

Behavior quantification: For analysis of freezing and motion in conditioning chambers,

Med Associates Video Freeze software was used to analyze videos acquired from a near infra-

red camera located in the chamber [84]. For measuring distance travelled and time spent in

regions of interest in exploratory anxiety-related behavior tests, ezTrack was used [85,86]. With

the exception of freezing during the trauma and novel stressor session, all measures reflect

the average across the entire session. For freezing during the trauma, time was binned into

the 300 sec baseline and then 10 post-shock periods. Each post-shock period was 20 sec in

length and begun 10 sec after shock offset.

Surgery:

For surgery, anesthesia was induced with 5% isoflurane and subsequently maintained

at 1-2%. Body temperature was maintained during surgery and recovery with a heating pad

below the animal, and ophthalmic ointment was applied to lubricate the eyes. All surgeries

followed aseptic surgical technique. For viral surgeries, 100 nL of AAV5-hSyn-PSAM4-GlyR-

IRES-EGFP (2.4 x 10^{13} GC/mL; Addgene 119741) was infused into the BLA (AP: -1.4; ML: 3.3;

DV: -5) or vHC (AP: -3; ML: 3.2; DV: -4.5) at 2 nL/sec via glass pipettes. Alternatively, an

equivalent volume of sterile PBS was infused. 10 min was allowed for diffusion before

removing the injector, irrigating the incision with saline, and suturing the incision site. For

cannulation surgeries, 26 gauge guide cannula (P1 Technologies; 8IC315GMNSPC) were

implanted overlying the BLA (AP: -1.4; ML: 3.2; DV: -3.5) or vHC (AP: -3; ML: 3.2; DV: -3), and

affixed to the skull with dental cement and super glue. A skull screw was also implanted

during surgery to help secure the head cap (P1 Technologies; 00-96X1/16). After surgery,

dummy cannula that extended 1.5 mm below the guide cannula were inserted (P1

Technologies 8IC315DCMNSP). Following surgery, animals were given 20 mg/kg ampicillin

and 5 mg/kg carprofen (s.c.) per day for 7 days and body weight and general disposition were

monitored. All surgeries followed aseptic surgical technique.

Anisomycin experiments:

For experiments in which anisomycin (Sigma A9789) was administered systemically, we

utilized a dose of 150 mg/kg (10 mL/kg, s.c.), consistent with prior literature [87,88]. Because

numerous waves of protein synthesis have been found to support memory consolidation

[59,89,90], we opted to administer anisomycin 3 times, once every 4 hours. In line with prior

reports [91,92], this should maintain approximately 90% blockade of protein synthesis for 12

hours. For experiments in which anisomycin was administered intracranially, 33 gauge

injectors (P1 Technologies; 8IC315IMNSPC) attached via PE-20 tubing (Instech) to a Harvard

syringe pump (Harvard Apparatus, #55-2222) were utilized to infuse anisomycin (10 ng/nL) at

a rate of 150 nL/min. 300 nL of anisomycin solution was administered per hemisphere in the

vHC. 200 nL was administered per hemisphere in the BLA. Control animals were infused with

an equivalent volume of 1X PBS. Following infusions, injectors were left in place for 1 min

before removal. Again, anisomycin was infused 3 times, once every four hours. Prior to

testing, animals were habituated to handling such that infusions could be done while mice

were gently held by the experimenter. Additionally, all animals received a habituation infusion

of 1X PBS 2-3 days prior to the trauma day. Anisomycin was first dissolved in a small volume

of 0.1 N HCL (90% PBS, 10% 1 N HCL), brought near concentration with the addition of 1X

PBS, and the pH was then normalized to 6-7 by the addition of 1 N NaOH.

PSAM experiments:

For PSAM experiments, actuation of PSAM4-GlyR was achieved through intraperitoneal

administration of 1 mg/kg uPSEM-817-tartrate (Tocris), 15-20 minutes prior to behavior, at a

volume of 10 mL/kg (dissolved in saline). For the study in which the effects of inhibiting the

vHC or BLA were assessed on multiple measures of exploratory anxiety-related behavior,

each animal underwent each of these tests twice, once with uPSEM and once with vehicle.

Tests occurred in a fixed order across two weeks, with open-field on Monday, EPM on

Wednesday, and Light-Dark on Friday. However, drug order was counterbalanced, such that

half the animals that received uPSEM on the first open field test received saline on the first

EPM, and so forth.

Histology:

At the end of behavioral testing, animals that underwent surgical manipulation were
deeply anesthetized, and their brains were then extracted and placed in paraformaldehyde
overnight at 4C. For animals with cannula implants, 100 nl of DAPI (0.5 mg/mL) was infused
prior to brain extraction, but after anesthesia, to mark cannula placement. The next day,
brains were transferred to 30% sucrose in 1X PBS and left at 4C to sink before being frozen
and sectioned at 50 um on a cryostat. Tissue was then mounted on slides and either cover-
slipped using mounting media with DAPI (Vector Laboratories, #H-1200-10) for checking viral
placement or cover-slipped with non-fluorescent mounting media (Vector Laboratories, #H-
1000-10) after a green nucleic acid stain. For green nucleic acid staining, slides were
submerged in 50 mM Sytox Green (diluted in 1X PBS from 5 mM. Thermo Fisher #S7020) for
10 min and then washed 3x in 1X PBS. Tissue was then imaged on a Leica DM6
epifluorescent microscope. Viral expression and cannula placement was evaluated using the
mouse brain atlas of Franklin and Paxinos [93].
545
Analysis:
All analyses were performed using RStudio. All data and statistical analysis are
available at github.com/ZachPenn/BLAvHC_Dissoc. Groups sizes are listed in each figure
legend. Briefly, omnibus ANOVA were conducted using the package ezANOVA with type 3
degrees of freedom. The white adjustment was implemented to correct for heterogeneity of
variance using heteroscedasticity corrected standard errors ('hc3'). For repeated measures
ANOVA, the Greenhouse-Geisser correction was implemented when the assumption of
sphericity was not met. Planned comparisons and post-hoc t-tests (Welch unequal variance)
were performed after omnibus significance was detected. Post-hoc tests were evaluated
against a modified criterion calculated using the Dunn-Sidak method in order to keep family-
wise type 1 error at 0.05. F and t values are rounded to the nearest tenth. Where F values
were less than .1, F is listed as 0.

KEY RESOURCE TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and virus strains		
AAV5-hSyn-PSAM4-GlyR-IRES-EGFP (2.4 x 10^{13} GC/mL)	Magnus et al, 2019[61]	Addgene: 119742
Chemicals, peptides, and recombinant proteins		
Anisomycin from *Streptomyces griseolus*	Millipore Sigma	Millipore Sigma: A9789
uPSEM 817 tartrate	Tocris	Tocris: 6866
Experimental models: Organisms/strains		
Mouse: C57BL/6J	The Jackson Laboratory	JAX: 000664
Software and algorithms		
Med Associates Video Freeze	Med Associates[84]	Med Associates: SOF-843
ezTrack	Pennington et al, 2019[86]	www.github.com/denisecailab/eztrack

References and Literature Cited

1. Fendt, M., and Fanselow, M.S. (1999). The neuroanatomical and neurochemical basis of conditioned fear. Neurosci Biobehav Rev *23*, 743-760.

2. Bolles, R.C. (1970). Species-specific defense reations and avoidance learning. Psychological Review *77*, 32-48.

3. Fanselow, M.S., and Lester, L.S. (1988). A functional behavioristic approach to aversively motivated behavior. In Evolution and Learning, R.C. Bolles, and M.C. Beecher, eds. (Erlbaum), pp. 185-211.

4. Cannon, W.B. (1915). Bodily changes in pain, hunger, fear and rage: An account of recent researches into the function of emotional excitement. (D. Appleton and Componay).

5. Blanchard, R.J., Blanchard, D.C., Rodgers, J., and Weiss, S.M. (1990). The characterization and modelling of antipredator defensive behavior. Neurosci Biobehav Rev *14*, 463-472. 10.1016/s0149-7634(05)80069-7.

6. Blanchard, R.J., and Blanchard, D.C. (1969). Passive and active reactions to fear-eliciting stimuli. J Comp Physiol Psychol *68*, 129-135. 10.1037/h0027676.

7. Bienvenu, T.C.M., Dejean, C., Jercog, D., Aouizerate, B., Lemoine, M., and Herry, C. (2021). The advent of fear conditioning as an animal model of post-traumatic stress disorder: Learning from the past to shape the future of PTSD research. Neuron *109*, 2380-2397. 10.1016/j.neuron.2021.05.017.

8. Maren, S. (2005). Synaptic mechanisms of associative memory in the amygdala. Neuron *47*, 783-786. 10.1016/j.neuron.2005.08.009.

9. Maren, S. (2003). The amygdala, synaptic plasticity, and fear memory. Ann N Y Acad Sci *985*, 106-113.

10. Fanselow, M.S., and LeDoux, J.E. (1999). Why we think plasticity underlying Pavlovian fear conditioning occurs in the basolateral amygdala. Neuron *23*, 229-232.

11. Davis, M. (1992). The role of the amygdala in fear and anxiety. Annu Rev Neurosci *15*, 353-375. 10.1146/annurev.ne.15.030192.002033.

12. Fadok, J.P., Krabbe, S., Markovic, M., Courtin, J., Xu, C., Massi, L., Botta, P., Bylund, K., Müller, C., Kovacevic, A., et al. (2017). A competitive inhibitory circuit for selection of active and passive fear responses. Nature *542*, 96-100. 10.1038/nature21047.

13. Adamec, R.E., and Shallow, T. (1993). Lasting effects on rodent anxiety of a single exposure to a cat. Physiol Behav *54*, 101-109. 10.1016/0031-9384(93)90050-p.

14. Poulos, A.M., Reger, M., Mehta, N., Zhuravka, I., Sterlace, S.S., Gannam, C., Hovda, D.A., Giza, C.C., and Fanselow, M.S. (2014). Amnesia for early life stress does not preclude the adult development of posttraumatic stress disorder symptoms in rats. Biol Psychiatry *76*, 306-314. 10.1016/j.biopsych.2013.10.007.

15. Rau, V., DeCola, J.P., and Fanselow, M.S. (2005). Stress-induced enhancement of fear learning: an animal model of posttraumatic stress disorder. Neurosci Biobehav Rev *29*, 1207-1223. 10.1016/j.neubiorev.2005.04.010.

16. Pynoos, R.S., Ritzmann, R.F., Steinberg, A.M., Goenjian, A., and Prisecaru, I. (1996). A behavioral animal model of posttraumatic stress disorder featuring repeated exposure to situational reminders. Biol Psychiatry *39*, 129-134. 10.1016/0006-3223(95)00088-7.

17. Servatius, R.J., Ottenweller, J.E., and Natelson, B.H. (1995). Delayed startle sensitization distinguishes rats exposed to one or three stress sessions: further

evidence toward an animal model of PTSD. Biol Psychiatry *38*, 539-546. 10.1016/0006-
3223(94)00369-E.

18.   American Psychiatric Association. (2013). American Psychiatric Association: Diagnostic
and Statistical Manual of Mental Disorders, Fifth Edition. (American Psychiatric
Publishing).

19.   Moreno-Peral, P., Conejo-Cerón, S., Motrico, E., Rodríguez-Morejón, A., Fernández, A.,
García-Campayo, J., Roca, M., Serrano-Blanco, A., Rubio-Valera, M., and Bellón, J.
(2014). Risk factors for the onset of panic and generalised anxiety disorders in the
general adult population: a systematic review of cohort studies. J Affect Disord *168*,
337-348. 10.1016/j.jad.2014.06.021.

20.   Kessler, R.C., Gruber, M., Hettema, J.M., Hwang, I., Sampson, N., and Yonkers, K.A.
(2008). Co-morbid major depression and generalized anxiety disorders in the National
Comorbidity Survey follow-up. Psychol Med *38*, 365-374.
10.1017/S0033291707002012.

21.   Blanco, C., Rubio, J., Wall, M., Wang, S., Jiu, C.J., and Kendler, K.S. (2014). Risk
factors for anxiety disorders: common and specific effects in a national sample. Depress
Anxiety *31*, 756-764. 10.1002/da.22247.

22.   Keiser, A.A., Turnbull, L.M., Darian, M.A., Feldman, D.E., Song, I., and Tronson, N.C.
(2017). Sex Differences in Context Fear Generalization and Recruitment of
Hippocampus and Amygdala during Retrieval. Neuropsychopharmacology *42*, 397-407.
10.1038/npp.2016.174.

23.   Lissek, S., Bradford, D.E., Alvarez, R.P., Burton, P., Espensen-Sturges, T., Reynolds,
R.C., and Grillon, C. (2014). Neural substrates of classically conditioned fear-
generalization in humans: a parametric fMRI study. Soc Cogn Affect Neurosci *9*, 1134-
1142. 10.1093/scan/nst096.

24.   Lissek, S., Kaczkurkin, A.N., Rabin, S., Geraci, M., Pine, D.S., and Grillon, C. (2014).
Generalized anxiety disorder is associated with overgeneralization of classically
conditioned fear. Biol Psychiatry *75*, 909-915. 10.1016/j.biopsych.2013.07.025.

25.   Zelikowsky, M., Bissiere, S., Hast, T.A., Bennett, R.Z., Abdipranoto, A., Vissel, B., and
Fanselow, M.S. (2013). Prefrontal microcircuit underlies contextual learning after
hippocampal loss. Proc Natl Acad Sci U S A *110*, 9938-9943.
10.1073/pnas.1301691110.

26.   Xu, W., and Südhof, T.C. (2013). A neural circuit for memory specificity and
generalization. Science *339*, 1290-1295. 10.1126/science.1229534.

27.   Wiltgen, B.J., Zhou, M., Cai, Y., Balaji, J., Karlsson, M.G., Parivash, S.N., Li, W., and
Silva, A.J. (2010). The hippocampus plays a selective role in the retrieval of detailed
contextual memories. Curr Biol *20*, 1336-1344. 10.1016/j.cub.2010.06.068.

28.   Starita, F., Kroes, M.C.W., Davachi, L., Phelps, E.A., and Dunsmoor, J.E. (2019).
Threat learning promotes generalization of episodic memory. J Exp Psychol Gen *148*,
1426-1434. 10.1037/xge0000551.

29.   Dunsmoor, J.E., Kroes, M.C.W., Braren, S.H., and Phelps, E.A. (2017). Threat intensity
widens fear generalization gradients. Behav Neurosci *131*, 168-175.
10.1037/bne0000186.

30.   Dunsmoor, J.E., Otto, A.R., and Phelps, E.A. (2017). Stress promotes generalization of
older but not recent threat memories. Proc Natl Acad Sci U S A *114*, 9218-9223.
10.1073/pnas.1704428114.

31.   Davis, M. (1989). Neural systems involved in fear-potentiated startle. Ann N Y Acad Sci
563, 165-183. 10.1111/j.1749-6632.1989.tb42197.x.
32.   Brown, J.S., Kalish, H.I., and Farber, I.E. (1951). Conditioned fear as revealed by
magnitude of startle response to an auditory stimulus. J Exp Psychol *41*, 317-328.
10.1037/h0060166.
33.   Kaczkurkin, A.N., Burton, P.C., Chazin, S.M., Manbeck, A.B., Espensen-Sturges, T.,
Cooper, S.E., Sponheim, S.R., and Lissek, S. (2017). Neural Substrates of
Overgeneralized Conditioned Fear in PTSD. Am J Psychiatry *174*, 125-134.
10.1176/appi.ajp.2016.15121549.
34.   Grillon, C., and Morgan, C.A. (1999). Fear-potentiated startle conditioning to explicit and
contextual cues in Gulf War veterans with posttraumatic stress disorder. J Abnorm
Psychol *108*, 134-142.
35.   Acheson, D.T., Geyer, M.A., Baker, D.G., Nievergelt, C.M., Yurgil, K., Risbrough, V.B.,
and Team, M.-I. (2015). Conditioned fear and extinction learning performance and its
association with psychiatric symptoms in active duty Marines.
Psychoneuroendocrinology *51*, 495-505. 10.1016/j.psyneuen.2014.09.030.
36.   Long, V.A., and Fanselow, M.S. (2012). Stress-enhanced fear learning in rats is
resistant to the effects of immediate massed extinction. Stress *15*, 627-636.
10.3109/10253890.2011.650251.
37.   Hassien, A.M., Shue, F., Bernier, B.E., and Drew, M.R. (2020). A mouse model of
stress-enhanced fear learning demonstrates extinction-sensitive and extinction-resistant
effects of footshock stress. Behav Brain Res *379*, 112391. 10.1016/j.bbr.2019.112391.
38.   Difede, J., Olden, M., and Cukor, J. (2014). Evidence-based treatment of post-traumatic
stress disorder. Annu Rev Med *65*, 319-332. 10.1146/annurev-med-051812-145438.
39.   Bradley, R., Greene, J., Russ, E., Dutra, L., and Westen, D. (2005). A multidimensional
meta-analysis of psychotherapy for PTSD. Am J Psychiatry *162*, 214-227.
10.1176/appi.ajp.162.2.214.
40.   Cukor, J., Olden, M., Lee, F., and Difede, J. (2010). Evidence-based treatments for
PTSD, new directions, and special challenges. Ann N Y Acad Sci *1208*, 82-89.
10.1111/j.1749-6632.2010.05793.x.
41.   Maren, S., Aharonov, G., and Fanselow, M.S. (1996). Retrograde abolition of
conditional fear after excitotoxic lesions in the basolateral amygdala of rats: absence of
a temporal gradient. Behav Neurosci *110*, 718-726.
42.   Blanchard, D.C., and Blanchard, R.J. (1972). Innate and conditioned reactions to threat
in rats with amygdaloid lesions. J Comp Physiol Psychol *81*, 281-290.
43.   Feinstein, J.S., Adolphs, R., Damasio, A., and Tranel, D. (2011). The human amygdala
and the induction and experience of fear. Curr Biol *21*, 34-38.
10.1016/j.cub.2010.11.042.
44.   Padilla-Coreano, N., Bolkan, S.S., Pierce, G.M., Blackman, D.R., Hardin, W.D., Garcia-
Garcia, A.L., Spellman, T.J., and Gordon, J.A. (2016). Direct Ventral Hippocampal-
Prefrontal Input Is Required for Anxiety-Related Neural Activity and Behavior. Neuron
89, 857-866. 10.1016/j.neuron.2016.01.011.
45.   Kheirbek, M.A., Drew, L.J., Burghardt, N.S., Costantini, D.O., Tannenholz, L., Ahmari,
S.E., Zeng, H., Fenton, A.A., and Hen, R. (2013). Differential control of learning and
anxiety along the dorsoventral axis of the dentate gyrus. Neuron *77*, 955-968.
10.1016/j.neuron.2012.12.038.

46.   Adhikari, A., Topiwala, M.A., and Gordon, J.A. (2010). Synchronized activity between
699   the ventral hippocampus and the medial prefrontal cortex during anxiety. Neuron *65*,
700   257-269. 10.1016/j.neuron.2009.12.002.
701   47.   Fanselow, M.S., and Dong, H.W. (2010). Are the dorsal and ventral hippocampus
702   functionally distinct structures? Neuron *65*, 7-19. 10.1016/j.neuron.2009.11.031.
703   48.   Bannerman, D.M., Yee, B.K., Good, M.A., Heupel, M.J., Iversen, S.D., and Rawlins,
704   J.N. (1999). Double dissociation of function within the hippocampus: a comparison of
705   dorsal, ventral, and complete hippocampal cytotoxic lesions. Behav Neurosci *113*,
706   1170-1188. 10.1037//0735-7044.113.6.1170.
707   49.   Perusini, J., Meyer, E., Rau, V., Avershal, J., Rajbhandari, A., Hoffman, A., Nocera, N.,
708   Condro, M., Waschek, J., Spigelman, I., and Fanselow, M. (2014). Mechanisms
709   underlying the induction and expression of fear sensitization following acute traumatic
710   stress. Pavlovian Society.
711   50.   Rau, V., and Fanselow, M.S. (2009). Exposure to a stressor produces a long lasting
712   enhancement of fear learning in rats. Stress *12*, 125-133.
713   10.1080/10253890802137320.
714   51.   Pennington, Z.T., Trott, J.M., Rajbhandari, A.K., Li, K., Walwyn, W.M., Evans, C.J., and
715   Fanselow, M.S. (2020). Chronic opioid pretreatment potentiates the sensitization of fear
716   learning by trauma. Neuropsychopharmacology *45*, 482-490. 10.1038/s41386-019-
717   0559-5.
718   52.   Chaouloff, F., Durand, M., and Mormède, P. (1997). Anxiety- and activity-related effects
719   of diazepam and chlordiazepoxide in the rat light/dark and dark/light tests. Behav Brain
720   Res *85*, 27-35. 10.1016/s0166-4328(96)00160-x.
721   53.   Crawley, J.N. (1981). Neuropharmacologic specificity of a simple animal model for the
722   behavioral actions of benzodiazepines. Pharmacol Biochem Behav *15*, 695-699.
723   10.1016/0091-3057(81)90007-1.
724   54.   Smith, A.C.W., Jonkman, S., Difeliceantonio, A.G., O'Connor, R.M., Ghoshal, S.,
725   Romano, M.F., Everitt, B.J., and Kenny, P.J. (2021). Opposing roles for striatonigral and
726   striatopallidal neurons in dorsolateral striatum in consolidating new instrumental actions.
727   Nat Commun *12*, 5121. 10.1038/s41467-021-25460-3.
728   55.   Hernandez, P.J., Sadeghian, K., and Kelley, A.E. (2002). Early consolidation of
729   instrumental learning requires protein synthesis in the nucleus accumbens. Nat
730   Neurosci *5*, 1327-1331. 10.1038/nn973.
731   56.   Santini, E., Ge, H., Ren, K., Peña de Ortiz, S., and Quirk, G.J. (2004). Consolidation of
732   fear extinction requires protein synthesis in the medial prefrontal cortex. J Neurosci *24*,
733   5704-5710. 10.1523/JNEUROSCI.0786-04.2004.
734   57.   Schafe, G.E., and LeDoux, J.E. (2000). Memory consolidation of auditory pavlovian fear
735   conditioning requires protein synthesis and protein kinase A in the amygdala. J
736   Neurosci *20*, RC96.
737   58.   Nader, K., Schafe, G.E., and Le Doux, J.E. (2000). Fear memories require protein
738   synthesis in the amygdala for reconsolidation after retrieval. Nature *406*, 722-726.
739   10.1038/35021052.
740   59.   Bourtchouladze, R., Abel, T., Berman, N., Gordon, R., Lapidus, K., and Kandel, E.R.
741   (1998). Different training procedures recruit either one or two critical periods for
742   contextual memory consolidation, each of which requires protein synthesis and PKA.
743   Learn Mem *5*, 365-374.

744 60. Kandel, E.R. (2001). The molecular biology of memory storage: a dialogue between
745   genes and synapses. Science *294*, 1030-1038. 10.1126/science.1067020.
746 61. Magnus, C.J., Lee, P.H., Bonaventura, J., Zemla, R., Gomez, J.L., Ramirez, M.H., Hu,
747   X., Galvan, A., Basu, J., Michaelides, M., and Sternson, S.M. (2019). Ultrapotent
748   chemogenetics for research and potential clinical applications. Science *364*.
749   10.1126/science.aav5282.
750 62. Sierra-Mercado, D., Padilla-Coreano, N., and Quirk, G.J. (2011). Dissociable roles of
751   prelimbic and infralimbic cortices, ventral hippocampus, and basolateral amygdala in the
752   expression and extinction of conditioned fear. Neuropsychopharmacology *36*, 529-538.
753   10.1038/npp.2010.184.
754 63. Hintiryan, H., Bowman, I., Johnson, D.L., Korobkova, L., Zhu, M., Khanjani, N., Gou, L.,
755   Gao, L., Yamashita, S., Bienkowski, M.S., et al. (2021). Connectivity characterization of
756   the mouse basolateral amygdalar complex. Nat Commun *12*, 2859. 10.1038/s41467-
757   021-22915-5.
758 64. Canteras, N.S., Simerly, R.B., and Swanson, L.W. (1992). Connections of the posterior
759   nucleus of the amygdala. J Comp Neurol *324*, 143-179. 10.1002/cne.903240203.
760 65. Canteras, N.S., and Swanson, L.W. (1992). Projections of the ventral subiculum to the
761   amygdala, septum, and hypothalamus: a PHAL anterograde tract-tracing study in the
762   rat. J Comp Neurol *324*, 180-194. 10.1002/cne.903240204.
763 66. Sprengelmeyer, R., Young, A.W., Schroeder, U., Grossenbacher, P.G., Federlein, J.,
764   Büttner, T., and Przuntek, H. (1999). Knowing no fear. Proc Biol Sci *266*, 2451-2456.
765   10.1098/rspb.1999.0945.
766 67. Fanselow, M.S., and Kim, J.J. (1994). Acquisition of contextual Pavlovian fear
767   conditioning is blocked by application of an NMDA receptor antagonist D,L-2-amino-5-
768   phosphonovaleric acid to the basolateral amygdala. Behav Neurosci *108*, 210-212.
769 68. Davis, M. (1992). The role of the amygdala in fear-potentiated startle: implications for
770   animal models of anxiety. Trends Pharmacol Sci *13*, 35-41.
771 69. Rumpel, S., LeDoux, J., Zador, A., and Malinow, R. (2005). Postsynaptic receptor
772   trafficking underlying a form of associative learning. Science *308*, 83-88.
773   10.1126/science.1103944.
774 70. Ribeiro, A.M., Barbosa, F.F., Munguba, H., Costa, M.S., Cavalcante, J.S., and Silva,
775   R.H. (2011). Basolateral amygdala inactivation impairs learned (but not innate) fear
776   response in rats. Neurobiol Learn Mem *95*, 433-440. 10.1016/j.nlm.2011.02.004.
777 71. Felix-Ortiz, A.C., Beyeler, A., Seo, C., Leppla, C.A., Wildes, C.P., and Tye, K.M. (2013).
778   BLA to vHPC inputs modulate anxiety-related behaviors. Neuron *79*, 658-664.
779   10.1016/j.neuron.2013.06.016.
780 72. Tye, K.M., Prakash, R., Kim, S.Y., Fenno, L.E., Grosenick, L., Zarabi, H., Thompson,
781   K.R., Gradinaru, V., Ramakrishnan, C., and Deisseroth, K. (2011). Amygdala circuitry
782   mediating reversible and bidirectional control of anxiety. Nature *471*, 358-362.
783   10.1038/nature09820.
784 73. Moreira, C.M., Masson, S., Carvalho, M.C., and Brandão, M.L. (2007). Exploratory
785   behaviour of rats in the elevated plus-maze is differentially sensitive to inactivation of
786   the basolateral and central amygdaloid nuclei. Brain Res Bull *71*, 466-474.
787   10.1016/j.brainresbull.2006.10.004.
788 74. Bueno, C.H., Zangrossi, H., and Viana, M.B. (2005). The inactivation of the basolateral
789   nucleus of the rat amygdala has an anxiolytic effect in the elevated T-maze and

790 light/dark transition tests. Braz J Med Biol Res *38*, 1697-1701. 10.1590/s0100-
791 879x2005001100019.

75. Jimenez, J.C., Su, K., Goldberg, A.R., Luna, V.M., Biane, J.S., Ordek, G., Zhou, P., Ong, S.K., Wright, M.A., Zweifel, L., et al. (2018). Anxiety Cells in a Hippocampal-Hypothalamic Circuit. Neuron *97*, 670-683.e676. 10.1016/j.neuron.2018.01.016.

76. Bang, J.Y., Sunstrum, J.K., Garand, D., Parfitt, G.M., Woodin, M., Inoue, W., and Kim, J. (2022). Hippocampal-hypothalamic circuit controls context-dependent innate defensive responses. Elife *11*. 10.7554/eLife.74736.

77. Yan, J.J., Ding, X.J., He, T., Chen, A.X., Zhang, W., Yu, Z.X., Cheng, X.Y., Wei, C.Y., Hu, Q.D., Liu, X.Y., et al. (2022). A circuit from the ventral subiculum to anterior hypothalamic nucleus GABAergic neurons essential for anxiety-like behavioral avoidance. Nat Commun *13*, 7464. 10.1038/s41467-022-35211-7.

78. Meyer, R.M., Burgos-Robles, A., Liu, E., Correia, S.S., and Goosens, K.A. (2014). A ghrelin-growth hormone axis drives stress-induced vulnerability to enhanced fear. Mol Psychiatry *19*, 1284-1294. 10.1038/mp.2013.135.

79. Maren, S., and Holt, W.G. (2004). Hippocampus and Pavlovian fear conditioning in rats: muscimol infusions into the ventral, but not dorsal, hippocampus impair the acquisition of conditional freezing to an auditory conditional stimulus. Behav Neurosci *118*, 97-110. 10.1037/0735-7044.118.1.97.

80. Jimenez, J.C., Berry, J.E., Lim, S.C., Ong, S.K., Kheirbek, M.A., and Hen, R. (2020). Contextual fear memory retrieval by correlated ensembles of ventral CA1 neurons. Nat Commun *11*, 3492. 10.1038/s41467-020-17270-w.

81. Etkin, A., and Wager, T.D. (2007). Functional neuroimaging of anxiety: a meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. Am J Psychiatry *164*, 1476-1488. 10.1176/appi.ajp.2007.07030504.

82. Ozawa, T., Ycu, E.A., Kumar, A., Yeh, L.F., Ahmed, T., Koivumaa, J., and Johansen, J.P. (2017). A feedback neural circuit for calibrating aversive memory strength. Nat Neurosci *20*, 90-97. 10.1038/nn.4439.

83. Rescorla, R.W., and Wagner, A.R. (1972). A theory of Pavlovian conditioning: Variations in the effectiveness of reinforcement and nonreinforcement. In Classical conditioning: II. Current research and theory, B.A. H, and P.W. F, eds. (Appleton-Century-Crofts), pp. 64-99.

84. Anagnostaras, S.G., Wood, S.C., Shuman, T., Cai, D.J., Leduc, A.D., Zurn, K.R., Zurn, J.B., Sage, J.R., and Herrera, G.M. (2010). Automated assessment of pavlovian conditioned freezing and shock reactivity in mice using the video freeze system. Front Behav Neurosci *4*. 10.3389/fnbeh.2010.00158.

85. Pennington, Z.T., Diego, K.S., Francisco, T.R., LaBanca, A.R., Lamsifer, S.I., Liobimova, O., Shuman, T., and Cai, D.J. (2021). ezTrack-A Step-by-Step Guide to Behavior Tracking. Curr Protoc *1*, e255. 10.1002/cpz1.255.

86. Pennington, Z.T., Dong, Z., Feng, Y., Vetere, L.M., Page-Harley, L., Shuman, T., and Cai, D.J. (2019). ezTrack: An open-source video analysis pipeline for the investigation of animal behavior. Sci Rep *9*, 19979. 10.1038/s41598-019-56408-9.

87. Lattal, K.M., and Abel, T. (2001). Different requirements for protein synthesis in acquisition and extinction of spatial preferences and context-evoked fear. J Neurosci *21*, 5773-5780. 10.1523/JNEUROSCI.21-15-05773.2001.

835   88.   Frankland, P.W., Ding, H.K., Takahashi, E., Suzuki, A., Kida, S., and Silva, A.J. (2006).
836   Stability of recent and remote contextual fear memory. Learn Mem *13*, 451-457.
837   10.1101/lm.183406.
838   89.   Lattal, K.M., and Abel, T. (2004). Behavioral impairments caused by injections of the
839   protein synthesis inhibitor anisomycin after contextual retrieval reverse with time. Proc
840   Natl Acad Sci U S A *101*, 4667-4672. 10.1073/pnas.0306546101.
841   90.   Grecksch, G., and Matthies, H. (1980). Two sensitive periods for the amnesic effect of
842   anisomycin. Pharmacol Biochem Behav *12*, 663-665. 10.1016/0091-3057(80)90145-8.
843   91.   Wanisch, K., and Wotjak, C.T. (2008). Time course and efficiency of protein synthesis
844   inhibition following intracerebral and systemic anisomycin treatment. Neurobiol Learn
845   Mem *90*, 485-494. 10.1016/j.nlm.2008.02.007.
846   92.   Flood, J.F., Rosenzweig, M.R., Bennett, E.L., and Orme, A.E. (1973). The influence of
847   duration of protein synthesis inhibition on memory. Physiol Behav *10*, 555-562.
848   10.1016/0031-9384(73)90221-7.
849   93.   Franklin, K., and Paxinos, G. (2008). The mouse brain in stereotaxic coordinates, 3
850   Edition (Elsevier Inc.).
851