A generic binding pocket for small molecule I_{Ks} activators at the extracellular inter-subunit interface of KCNQ1 and KCNE1 channel complexes

Magnus Chan[1,*], Harutyun Sahakyan[3,4,*], Jodene Eldstrom[1,*], Daniel Sastre[1], Yundi Wang[1], Ying Dou[1], Marc Pourrier[1], Vitya Vardanyan[2,#], and David Fedida[1,#]

[1] Department of Anesthesiology, Pharmacology and Therapeutics, University of British Columbia, Vancouver, British Columbia, Canada, V6T1Z3.

[2] Molecular Neuroscience Group, Institute of Molecular Biology, NAS RA, 7 Hasratyan St. 0014 Yerevan, Armenia.

[3] Laboratory of Computational Modeling of Biological Processes, Institute of Molecular Biology of NAS RA, 0014, Yerevan, Armenia.

[4] Present Address: National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda MD, USA.

[*] Contributed equally to the work in this paper.

[#] Email addresses for correspondence: david.fedida@ubc.ca and v_vardanyan@mb.sci.am

Keywords: ion channels, mefenamic acid, DIDS, I_{Ks}

Running Title: K^+ channel activator binding pocket

Dr. David Fedida
Life Sciences Centre, 2350 Health Sciences Mall, Room 2.301 Vancouver, British Columbia, Canada V6T 1Z3
Telephone: +1 (604) 822-5806, Email: david.fedida@ubc.ca

Abbreviations
I_{Ks}, delayed cardiac rectifier K^+ current; LQTS, long QT syndrome; DIDS, 4,4'-diisothiocyano-2,2'-stilbenedisulfonic acid; SITS, 4-acetamido-4'-isothiocyanatostilbene-2,2'-disulfonic acid; HMR1556, (3R,4S)-(+)-N-[3-hydroxy-2,2-dimethyl-6-(4,4,4-trifluorobutoxy) chroman-4-yl]-N-methylmethanesulfonamide; G-V, conductance-voltage relationship; k, slope factor; $V_{1/2}$, voltage at half-maximal activation; WT, wild type; ps-I_{Ks}, pseudo-KCNE1-KCNQ1; MD, molecular dynamics; MM, molecular mechanics; PBSA, Poisson-Boltzmann surface area; GBSA,

46 generalized Born surface area; RMSD, root mean square deviation; RMSF, root mean square
47 fluctuation; POPC, 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine.
53 **<u>Significance Statement</u>**
54
Combined, KCNQ1 and KCNE1 subunits generate the I_{Ks} current. Activating I_{Ks} has been
identified as a promising therapeutic strategy to treat arrhythmogenesis resulting from delayed
repolarization. In this study, we uncovered a common drug-induced binding site accessed by two
structurally diverse I_{Ks} activators, mefenamic acid and DIDS. Located in the extracellular
interface where KCNE1 interacts with KCNQ1, we propose that binding of drugs to this location
impairs channel closing and leads to enhanced current activation. This is shown to be particularly
beneficial at higher pulse rates and explains how such drugs may make an important contribution
to the electrical repolarization reserve in the heart.
63
64
65
66

<u>Abstract</u>
The cardiac I_{Ks} ion channel comprises KCNQ1, calmodulin, and KCNE1 in a dodecameric
complex which provides a repolarizing current reserve at higher heart rates and protects from
arrhythmia syndromes that cause fainting and sudden death. Pharmacological activators of I_{Ks} are
therefore of interest both scientifically and therapeutically for treatment of I_{Ks} loss-of-function
disorders. One group of chemical activators are only active in the presence of the accessory
KCNE1 subunit and here we investigate this phenomenon using molecular modeling techniques
and mutagenesis scanning in mammalian cells. A generalized activator binding pocket is formed
extracellularly by KCNE1, the domain-swapped S1 helices of one KCNQ1 subunit and the
pore/turret region made up of two other KCNQ1 subunits. A few residues, including K41, A44
and Y46 in KCNE1, W323 in the KCNQ1 pore, and Y148 in the KCNQ1 S1 domain, appear
critical for the binding of structurally diverse molecules, but in addition, molecular modeling
studies suggest that induced fit by structurally different molecules underlies the generalized
nature of the binding pocket. Activation of I_{Ks} is enhanced by stabilization of the KCNQ1-
S1/KCNE1/pore complex, which ultimately slows deactivation of the current, and promotes
outward current summation at higher pulse rates. Our results provide a mechanistic explanation
of enhanced I_{Ks} currents by these activator compounds and provide a map for future design of
more potent therapeutically useful molecules.

Introduction

Potassium ion (K^+) channel activators are important compounds in human health, as partial or complete loss of function of many K^+ channels may lead to inherited or acquired diseases that have significant morbidity and mortality. The delayed cardiac rectifier potassium current, I_{Ks}, plays an important role, especially at high heart rates, in the physiological shortening of the cardiac action potential (Sanguinetti et al., 1996). Unsurprisingly, mutations in both KCNQ1 (α-subunit) and KCNE1 (β-subunit), which when paired together give rise to the I_{Ks} current (Barhanin et al., 1996; Sanguinetti et al., 1996; Bendahhou et al., 2005), have been implicated in cardiac arrhythmia syndromes such as long QT syndrome (LQTS) and atrial fibrillation (Jervell and Lange-Nielsen, 1957; Wang et al., 1996; Chen et al., 2003; Eldstrom and Fedida, 2011; Olesen et al., 2014). With nearly all mutations seen in LQTS patients identified as loss-of-function and 50% of those loss-of-function mutations identified in the KCNQ1 subunit (Hedley et al., 2009; Ackerman et al., 2011), enhancing and activating I_{Ks} currents has long been suggested as a promising therapeutic approach for treating LQTS. It is a curiosity of I_{Ks} channels that the known activators fall into two general groupings: those that work best on the α-subunit, KCNQ1, alone, including zinc pyrithione, L-364,373, and one of the most studied activators, ML277 (Mattmann et al., 2012; Yu et al., 2013; Xu et al., 2015; Eldstrom et al., 2021); and those that work best in the presence of the auxiliary β-subunit, KCNE1, compounds like mefenamic acid and DIDS (4,4'-diisothiocyano-2,2'- stilbene disulfonic acid) (Abitbol et al., 1999; Wang et al., 2020).

Using cryo-EM, we recently visualized the binding of ML277 deep in the central core of KCNQ1 channels in a pocket lined inferiorly by the S4-S5 linker, laterally by the S5 and S6 helices of two separate subunits, and above by pore domain residues (Willegems et al., 2022). The location of the binding pocket and its structural inter-relationships help to explain the underlying mechanism of action of ML277, its specificity for KCNQ1, and the lack of efficacy due to steric hindrance in the presence of a β-subunit. However, it is known neither where activators of I_{Ks} bind that *require* the presence of the β-subunit, such as phenylboronic acid (Mruk and Kobertz, 2009), hexachlorophene, stilbenes such as DIDS and SITS (4-acetamido-4'-isothiocyanatostilbene-2,2'-disulfonic acid), and diclofenac acid derivatives such as mefenamic acid (Abitbol et al., 1999; Zheng et al., 2012; Wang et al., 2020), nor how they mediate their activator action. We have some clues for DIDS and mefenamic acid that their binding sites are not in the central channel core, as is the case for ML277, but depend on KCNE1 β-subunit residues at the extracellular surface of the channel, residues 39-43 in KCNE1 for DIDS (Abitbol et al., 1999) and K41 for mefenamic acid (Wang et al., 2020). Our recent cysteine scanning data revealed that although other extracellular KCNE1 residues in the same region to varying degrees impacted the effect of mefenamic acid, only the K41C mutation completely abolished mefenamic acid effect up to a concentration of 1 mM (Wang et al., 2020). Previous cross-linking studies have identified key interactions between this extracellular region of KCNE1 and the S1 and S6 transmembrane segments of KCNQ1 (Xu et al., 2008; Chung et al., 2009), suggesting that residues in either of these regions could also provide important clues to explain mefenamic acid's mechanism of action.

To further explore the dependence of residues in KCNE1 and those in adjacent KCNQ1 sites on binding of mefenamic acid to I_{Ks}, we first examined the role of K41C in preventing the drug effect. Docking in combination with molecular dynamics (MD) simulations of mefenamic acid binding to I_{Ks} followed by mutagenesis were used to map out critical KCNE1 and KCNQ1

residues. Further, we expanded on the idea that the stilbene, DIDS (Abitbol et al., 1999), which
is structurally quite different from mefenamic acid, shares a common binding site. Our results
showed that both compounds bind in the same general region formed by elements of the pore and
S6 domains of KCNQ1 and the near extracellular region of KCNE1, but depend on different
critical residues for their binding stability. Exposure of the channel complex to either compound
induces subtle structural changes that subsequently stabilize the conformation of the S1/outer
pore/S6 of KCNQ1 and slow the I_{Ks} deactivation gating kinetics. The results suggest the
existence of a common drug-induced binding site and a mechanism of action for small molecule
I_{Ks} activators which is distinct from that of specific compounds that activate KCNQ1 alone.
140
141

Results

The mefenamic acid binding site on the KCNQ1/KCNE1 complex

Exposure of wild type I_{Ks} complexes (4:4 ratio; WT EQ) to 100 µM mefenamic acid transforms the slowly activating I_{Ks} current into one with an almost linear waveform and completely inhibits tail current decay at -40 mV (Figure 1A) (Abitbol et al., 1999; Unsöld et al., 2000; Wang et al., 2020). G-V relations obtained from peak initial tail currents show that 100 µM mefenamic acid hyperpolarizes the G-V ($\Delta V_{1/2}$ = -105.7 mV, Figure 1B, 1C) and decreases the slope (control k = 19.4 mV, Mef k = 41.3 mV, Table 1). We previously showed that introduction of a cysteine mutation at residue K41 in all four KCNE1 subunits (4:4 ratio of mutant K41C-KCNE1 to KCNQ1; K41C-EQ) itself had only a minor effect on the G-V, but prevented changes to currents and the G-V relationship on exposure to 100 µM or 1 mM mefenamic acid (Figure 1B, 1C and Table 1) (Wang et al., 2020). The data suggest that residue K41 in KCNE1 is critical to the action of mefenamic acid, but give no information why K41 is so important, and prompted us to question how this residue and other adjacent residues in KCNE1, and those nearby on KCNQ1 acted together to form a binding pocket for mefenamic acid.

Figure 1 near here

Mefenamic acid binding site predicted by molecular modeling.

Initially, to visualise potential drug binding sites and understand how mutation of KCNE1 and KCNQ1 residues might prevent drug action, *in silico* experiments of drug docking with subsequent MD simulations were performed on a model of I_{Ks} channels. The I_{Ks} model was constructed based on the recent cryo-EM structure of KCNQ1-KCNE3, which is thought to represent the activated-open state of the channel complex (Sun and MacKinnon, 2020). Taking into consideration the sequence similarity of KCNE1 and KCNE3 subunits in their transmembrane segments, it has been suggested that the main interface of these subunits with KCNQ1 is preserved in this region. Our initial data indicated that the extracellular residues of KCNE1 are involved in the action of I_{Ks} activators, so we constructed a model where external KCNE3 residues R53-Y58 were substituted with homologous KCNE1 residues, D39-A44 (Figure 2A). The resulting 4:4 I_{Ks} channel complex was termed pseudo-KCNE1-KCNQ1, ps-I_{Ks}, and Figure 2B shows the essential elements of the ps-I_{Ks} subunits which form extracellular interface of KCNQ1 and KCNE and served as a basis for drug docking and MD simulations. Details of the docking procedure are described in the Materials and Methods section and schematically summarized in Figure 2-figure supplement 1.

Figure 2 near here

Briefly, conformational sampling was performed on ps-I_{Ks} residues D39-A44, and conformations showing the lowest free energy were selected for docking using a four-dimensional (4D) docking approach to find the best binding pose of the ligand. A conformation with the best docking score (Figure 2B) shows mefenamic acid binding to the pocket formed between extracellular KCNE1 residues, the external S6 transmembrane helix of one subunit, and the S1 transmembrane domain of the neighboring subunit and the pore turret of a third subunit. The estimated volume of the pocket in the mefenamic acid bound state is ~307 $\mathring{A}^3$ with a hydrophobicity value of ~0.65 kcal/mol (Figure 2D). The free energy of the mefenamic acid-ps-I_{Ks} interaction estimated by the MM/GBSA method from analysis of 300 ns MD simulations was -37.9± 1.22 kcal/mol (Figure

4A, Table 4), while a similar value of -31.8 ± 1.46 kcal/mol was calculated using the Poisson-
Boltzmann surface area (MM/PBSA) model. The MMGBSA/MMPBSA data were used to
identify ps-I_{Ks} residues contributing to the free energy of interaction with mefenamic acid for
further analysis (Figure 2- figure supplement 2) though it should be noted that root mean square
deviation (RMSD) analysis of trajectories indicated that the binding was dynamic as there were
other binding conformations possible (Figure 2 – figure supplement 3). Mefenamic acid in this
complex was stabilized by two hydrogen bonds formed between the drug and ps-KCNE1
residues Y46 and E43 (Figure 2C, Supplemental movie S1). Analysis of MD simulations showed
that these two H-bonds were the most stable, the Y46 H-bond to mefenamic acid had a mean
frequency of 79.7%, and E43 22.5% of the drug bound time (Figure 2 – figure supplement 4;
Table 2).
199

The energy decomposition per amino acid using MM/GBSA and MM/PBSA methods revealed
several residues in KCNQ1 and psKCNE1 with significant contributions to mefenamic acid
coordination (Figure 2C, Figure 2-figure supplement 2). As expected, these are the KCNE1
residues located at the external region of the auxiliary subunit – amino acids K41 to Y46. In
addition, residues W323 and V324 located on the S6 helix as well as residues L142, Q147 and
Y148 located on the S1 helix exhibited the lowest interaction free energy scores. We focused on
functional validation of the KCNE1 residues K41, L42, E43, A44 and Y46, and KCNQ1 residues
W323, V324, L142, Q147 and Y148 by mutation to cysteine, alanine and/or tryptophan, and
examining the sensitivity of fully saturated EQ channel complexes to 100 µM mefenamic acid.
Although A300, located in the turret region was identified as potentially important (but less so
than KCNE residues and W323), we could not get adequate expression from A300C for
functional analysis.
212

Mutational impact on EQ current changes induced by mefenamic acid.
The effect of mutations on the current waveform and tail current response to 100 µM mefenamic
acid treatment was examined on I_{Ks} channels identified as: x-EQ-y where "x" denotes a KCNE1
mutation and "y" denotes a KCNQ1 mutation. In the absence of mefenamic acid (control), most
mutations, with the exception of EQ-L142C (Figure 3-figure supplement 2), produced slowly
activating currents with rapid tail current decay (Figure 3A).
In the presence of 100 µM mefenamic acid, the waveforms of WT EQ, S6 and pore mutations
EQ-V324A, EQ-V324W, EQ-Q147C and EQ-Y148C were all transformed into ones with
instantaneous current onset and slowed tail current decay (Figure 3A, Figure 3-figure supplement
1). Only the EQ-W323A waveform and tail current were largely unaffected by 100 µM
mefenamic acid (like K41C-EQ, Figure 1). The EQ-W323C and A44C-EQ current waveforms
were also unchanged by 100 µM mefenamic acid, but their tail current decay was slowed (Figure
3A, Figure 3-figure supplement 1). We interpreted this to mean that mefenamic acid binds to
EQ-W323C and A44C-EQ mutant open channels and slows closing, but that the drug-channel
complex is less stable and mefenamic acid unbinds during the interpulse interval, which relieves
drug action between pulses. The summary of the normalized response of the different mutants to
mefenamic acid is shown in Figure 3D. G-V plots were obtained from the tail current amplitudes
in the absence and presence of 100 µM mefenamic acid, and unlike WT EQ (Figure 1), minimal
change in the shape and position of the EQ-W323A G-V plot was seen after exposure to 100 µM
mefenamic acid ($V_{1/2}$ shift of -14.1 mV compared with -105.7 mV seen in WT EQ (Figure 1,
Figure 3B, 3E, and Table 1). For the W323C mutation, a less drastic decrease in size compared

to alanine, the $V_{1/2}$ shift seen with 100 µM mefenamic acid increased (EQ-W323C $\Delta V_{1/2}$ = -26.5
mV), suggesting that the size of the W323 residue is important (Figure 3E and Table 1).
However, when the neighboring V324 KCNQ1 residue was mutated to a smaller (alanine) or
bulkier (tryptophan) residue they both showed the same response to mefenamic acid. Both the
V324W and V324A mutations reduced $V_{1/2}$ shifts caused by 100 µM mefenamic acid to between
-14 to -19 mV (Figure 3E, Table 1, Figure 3-figure supplement 1).
240
241 **Figure 3 near here**
242
Extracellular S1 residues identified in the MD simulations also proved to be important for
mefenamic acid binding, although less so than W323 and K41 residues found in the KCNQ1
extracellular regions of the S6 segment and KCNE1, respectively. Compared to WT, EQ-Y148C
reduced the $V_{1/2}$ shift after exposure to 100 µM mefenamic acid (EQ-Y148C $\Delta V_{1/2}$ = -19.3 mV)
and lessened the slope of the G-V relationship (Figure 3C). The Q147C mutant on the other
hand, only partially prevented the $V_{1/2}$ shift observed after mefenamic acid treatment (EQ-
Q147C $\Delta V_{1/2}$ = -37.6 mV, Figure 3E).
Other KCNE1 residues located at the N-terminal limit of the KCNE1 transmembrane segment
(L42, E43, A44, Y46) were also investigated. Unlike with K41C, but similar to WT EQ, tail
current decay was inhibited in the presence of 100 µM mefenamic acid in all of these mutants,
reflected by the normalized response (Figure 3D, Figure 3-figure supplement 1). In addition, a
reduced but statistically significant shift in $V_{1/2}$ was observed with 100 µM mefenamic acid
compared to control in all mutants tested except K41C. (K41C $\Delta V_{1/2}$ = -3.8 mV; L42C $\Delta V_{1/2}$ = -
37.1 mV; E43C $\Delta V_{1/2}$ = -24.4 mV; A44C $\Delta V_{1/2}$ = -9.7 mV; Figure 3E). Representative current
traces and G-V curves for all mutants are shown in Figure 3- figure supplement 1. Consistent
with previous literature (Gofman et al., 2012; Wang et al., 2012; Kuenze et al., 2020), Y46C-EQ
in control conditions produced a current with faster activation and a complex GV curve which
made analysis and an assignment of slope and $V_{1/2}$ difficult, though a small left shift in the GV
curve was visible (data not shown). In lieu of Y46C, G-V data from the Y46A-EQ mutant
showed a potent effect of mefenamic acid, equivalent to that of WT with a $\Delta V_{1/2}$ = -106.2 mV
(Figure 3-figure supplement 1D). These results suggest that mutation of residues further away
from the N-terminus of KCNE1 than K41 has diminishing effects on the activator action of
mefenamic acid.
266
Augmented activation of EQ-L142C in the absence of mefenamic acid
Unlike other mutations, the control EQ-L142C current waveform displayed an almost
instantaneous current onset, and tail currents showed no decay with our standard protocol (Figure
3-figure supplement 2A). As the control EQ-L142C current waveform qualitatively resembled
WT EQ currents in the presence of 100 µM mefenamic acid, G-V plots at different interpulse
intervals were compared. At the standard interval of 15 s, the EQ-L142C G-V plot (blue) closely
overlapped that of WT EQ in the presence of Mef (solid grey), and at a 30 s interpulse interval,
the position of the WT EQ + Mef (grey open circles) and EQ-L142C plots (green) both
depolarized significantly (Figure 3-figure supplement 2B). It appeared that the L142C mutation
augmented channel activation as much as 100 µM mefenamic acid on WT channels.
In addition, mefenamic acid made the EQ-L142C G-V plot voltage independent, overlapping
with the G-V obtained for EQ-L142C with a 7 s interpulse interval (pink and red respectively,

Figure 3- figure supplement 2B). The data indicate that EQ-L142C is still responsive to mefenamic acid but does so from a heightened state of activation (Table 3).

MD simulations of mutant ps-I_{Ks} channels exposed to mefenamic acid.
To explore the role played by critical residues, *in silico* homologous mutations K41C, Y46C and W323A were introduced into the mefenamic acid-bound ps-I_{Ks} channel (Mef-ps-I_{Ks}) and the stability of the Mef-mutant complexes was assessed in MD simulations. Remarkably, mefenamic acid detached from mutant K41C and W323A channels within 120 ns in all three independent MD simulations with an AMBER force field (Supplemental movies S2, S3, Table 4). In contrast, mefenamic acid remained bound during the entire simulation time to WT and Y46C-ps-I_{Ks} complexes (Supplemental movie S1, S4, Table 4). The last mutant was tested as we could not determine the functional effect of mefenamic acid on this residue in electrophysiological experiments, but in the simulation, Y46C-ps-I_{Ks} behaved like Y46A from electrophysiological data (Figure 3, Figure 3 – figure supplement 1). Mefenamic acid unbound from A44C-ps-I_{Ks} channel complexes within 80, 100 and 110 ns during three different simulations, changing binding pose several times before doing so.
Similar results were obtained in MD simulations with the CHARMM force field where Mef-ps-I_{Ks} was embedded in the lipid membrane, although in two of five 1250 ns long-duration simulations we observed unbinding of Mef from WT ps-I_{Ks} complexes, which suggested that the lipid environment allowed a more dynamic interaction between Mef and IKs. In contrast, Mef left the binding site in all five 500 ns runs for W323A and K41C mutants (Figure 4 – figure supplement 1), and when the Y46C mutant was placed in a lipid environment, mefenamic acid left the binding site or changed its binding conformation in three of five 500 ns MD simulations. A significantly *reduced* free interaction energy (ΔG) of ligand binding for K41C and W323A mutant Mef-ps-I_{Ks} complexes in 300 ns simulations was observed compared to WT Mef-ps-I_{Ks} (Figure 4A, Table 4), and the small but statistically significant change in free energy observed for the Y46C mutant complex confirms that the Y46 residue is not as important as K41 and W323 for mefenamic acid binding. It should be noted that the absolute values of ΔG or corresponding dissociation constants (K_d) calculated from MD simulations do not reflect apparent ΔG and K_d values determined from electrophysiological experiments or biochemical essays.

The flexibility of the external ps-KCNE1 protein residues of the mutants W323A, K41C and Y46C ps-I_{Ks} channels was also analyzed from trajectories with 300 ns duration, by monitoring their average root mean square fluctuation (RMSF) during the last 100 ns of simulations after the detachment of mefenamic acid from the molecular complex or in the case of Y46C without introduction of the ligand. The RMSF values obtained from the mutant channels were then compared to that of WT ps-I_{Ks} channels where MD simulations of the same duration were conducted after removing the mefenamic acid molecule from the complex (Figure 4B). The results indicate that K41C and W323A mutations modestly increased the RMSF of D39-A44 ps-KCNE1 residues when compared to the WT ps-I_{Ks} channel complex without mefenamic acid bound (Figure 4C, 4D). The Y46C mutation seems to impact the mobility of the N-terminal portion of KCNE1 much less (Figure 4E), perhaps owing to its location further into the membrane. These results suggest that the side chains of K41 and W323 residues normally stabilize the conformation of the external region of KCNE1, so that mutation of these residues

324 increases random motion and reduces the ability of drugs like mefenamic acid to remain bound
325 at this site.
326
327 **Figure 4 near here**
328
Mutational impact on I_{Ks} current changes induced by DIDS.
To establish whether the binding pocket for mefenamic acid can be generalized to other I_{Ks}
activators, we also examined the binding of the structurally unrelated I_{Ks} activator, DIDS.
Stilbenes such as DIDS (Figure 5A) also activate I_{Ks} (Abitbol et al., 1999; Bollmann et al.,
2020), and given their molecular differences from fenamates, it is of interest to explore common
structural and dynamic features of their binding to the I_{Ks} channel complex. 100 µM DIDS had
no effect on endogenous currents in GFP-transfected tsA201 cells but treatment of WT EQ
channels with 100 µM DIDS transformed the slowly activating waveform into one with faster
onset (although not instantaneous like mefenamic acid) and inhibited tail current decay (Figure
5B). G-V plots were obtained from the tail amplitudes in the absence and presence of 100 µM
DIDS (Figure 5C, 5D). The overall shape, slope, and $V_{1/2}$ of the WT EQ G-V relationship
changed with 100 µM DIDS in the direction of increased activation ($\Delta V_{1/2}$ = -46.6 mV; Table 5;
control k = 20.3 mV, DIDS k = 25.3 mV). However, the effects on WT EQ were less pronounced
with DIDS than mefenamic acid at the 100 µM concentration.
343
344 **Figure 5 near here**
345
The ps-I_{Ks} construct and docking procedures used previously to explore the binding site of
mefenamic acid were used as a basis for DIDS docking and subsequent MD simulations. We
found that DIDS also bound to a location formed between extracellular KCNE1 residues, the
external S6 transmembrane helix of one subunit, the S1 transmembrane domain of the
neighboring subunit, and the pore turret of the opposite subunit (Figure 6A). In docking
experiments, DIDS is stabilized by its hydrophobic and van der Waals contacts with KCNQ1 and
KCNE1 subunits as well as by two hydrogen bonds formed between the drug and ps-KCNE1
residue L42 and KCNQ1 residue Q147 (Figure 6B). These hydrogen bonds however were not
stable during MD simulations and instead H-bonding occurs between E43, Y46, and I47 in
KCNE1 as well as W323, S298 and Y299 in KCNQ1 Figure 6- figure supplement 1, Table 2).
The energy decomposition per amino acid using MM/GBSA and MM/PBSA methods based on
300 ns simulations revealed several residues in KCNQ1 and psKCNE1 with significant
contributions to DIDS coordination (Figure 6C). Although residues identified as critical to
mefenamic acid binding and action also generally appear to be important for DIDS, especially
W323, K41 was noticeably not as critical to DIDS coordination, since the ΔG values for this
residue were significantly smaller compared to those of the Mef (see Figure 2-figure supplement
2).
363
364 **Figure 6 here**
365
Similar to mefenamic acid, DIDS stayed bound to WT ps-I_{Ks}, but detached from the W323A
mutant during two of three runs of 300 ns duration, and from Y46C channels in one of three runs
after 240, 239 and 122 ns of simulation, respectively (Supplemental movies S5, S7, Table 6).
Interestingly, DIDS did not unbind from K41C mutant ps-I_{Ks} channels, even after 300 ns of

simulation, although the binding pose did change in one run (Supplemental movie S6).
Calculations of free interaction energy shown in Table 6 using both MM/PBSA and MM/GBSA
methods are of mean values before the detachment of the drug if that occurred. If DIDS
detachment did not occur during 300 ns MD runs, the average free energy was calculated over
the duration of the entire MD run for that construct. Only DIDS binding to W323A, and to Y46C
in one simulation run, showed significantly lower free energy than DIDS binding to WT, using
both methodologies. The small change in free energy and more stable binding of DIDS to K41C
suggest that this residue is not as important as W323 or Y46 for DIDS binding. In longer 500 ns
simulations with the channel complexes embedded in lipid, trajectory analysis revealed clusters
of DIDS poses in complex with the WT ps-I_{Ks} channel, which are much more closely related than
those identified for mefenamic acid (Figure 6 – figure supplement 2, note the ordinate scale).
Still, DIDS remained bound to WT ps-I_{Ks}, during all runs throughout 500 ns simulations, and
both W323A and Y46C mutants unbound during 500 ns simulations (all runs for Y46C, Figure 6
– figure supplement 3) as was seen with the with AMBER force field. In this model the major
difference was that the K41C mutant unbound in three runs. These model predictions were tested
experimentally using fully saturated mutant I_{Ks} channel complexes, K41C-EQ, L42C-EQ, A44C-
EQ, Y46A-EQ, EQ-W323A, EQ-L142C and EQ-Y148C (Figure 7A, Figure 7-figure supplement
1). Y46 was identified as an important residue for DIDS binding in MD simulations, but as
before, the complexity of the GV curve and low functional expression of the Y46C construct led
us to substitute Y46A for this mutation in electrophysiology experiments (Figure 7 and Figure 7-
figure supplement 1).
391
392 **Figure 7 near here**
393
K41C-EQ G-V plots were obtained from peak tail current amplitudes in the absence and
presence (Figure 7B) of 100 µM DIDS. Unlike with mefenamic acid, but in agreement with the
modeling, K41C only partially prevented the action of DIDS. Treatment of K41C-EQ with 100
µM DIDS hyperpolarized the $V_{1/2}$ and changed the shape of the G-V relationship compared to
control (Figure 7B, 7D, Table 5). The current waveform of K41C-EQ activated more quickly
with less sigmoidicity after treatment with DIDS, and tail current decay was slowed as indicated
by the normalized response (Figure 7B, 7C). As K41C-EQ remained responsive to DIDS, and
L42C responded similarly to K41C-EQ (for K41C-EQ, $\Delta V_{1/2}$ was -24.7 mV, and for L42C-EQ
$\Delta V_{1/2}$ was -34 mV), we hypothesized that DIDS binding to I_{Ks} channel complexes may involve
KCNE1 residues closer to the transmembrane region. In agreement with this idea, A44C showed
little response to 100 µM DIDS both in terms of current waveform and tail current decay. The
shift in $V_{1/2}$ was also reduced compared to WT-EQ (A44C-EQ $\Delta V_{1/2}$ = -18.5 mV; WT EQ $\Delta V_{1/2}$
= -46.6mV, Figure 7C, D). In Y46A, the effect of DIDS on the current waveform was greatly
reduced as was slowing of tail current decay (Figure 7A, 7C). We could only estimate an
activation $V_{1/2}$ for Y46A as the G-V curve was right shifted to very positive potentials and non-
saturating. Nevertheless, hyperpolarization of the $V_{1/2}$ (Figure 7D, Figure 7-figure supplement 1)
in the presence of DIDS was observed, suggesting that unstable and short-lived binding of the
drug to the channel complex was sufficient to interfere with channel gating.
412
The modeling studies (Figure 6) suggested that W323 remained a key residue for I_{Ks} activator
sensitivity, and in agreement with this, EQ-W323A was only partially responsive to DIDS. In
EQ-W323A, the tail current decay was affected but the slowly-activating current waveform was

preserved in the presence of DIDS, supporting the idea that the drug dissociates from the channel
complex between pulses (Figure 7A, 7C, Table 6). However, a significant shift in the $V_{1/2}$
remained ($\Delta V_{1/2}$ = -27.3 mV; Figure 7D, Table 5). Comparable effects observed in EQ-W323A,
A44C-EQ, and Y46A-EQ are consistent with the idea that the DIDS binding pocket involves the
N-terminal end of S6 and the deeper extracellular surface of the KCNE1 transmembrane
segment. In EQ-Y148C, the $V_{1/2}$ and the shape of the G-V plot were not altered by 100 µM
DIDS (Figure 7D, Table 5), and tail current decay was not markedly slowed (Figure 7A, 7C).
This result suggests that unlike mefenamic acid, DIDS binding determinants extend to the S1
region of KCNQ1 and that Y148 is an important contributor.
Notably, as with mefenamic acid, the L142C-EQ G-V plot became voltage-independent after
treatment with 100 µM DIDS (data not shown) and so it proved hyper-responsive to both of the
I_{Ks} activators, DIDS and mefenamic acid.
428
429

Discussion

Mefenamic acid, a nonsteroidal anti-inflammatory drug (NSAID), and the structurally distinct stilbenes DIDS and SITS, among other compounds, have previously been identified by numerous groups to enhance I_{Ks} currents in various expression systems including canine and guinea-pig ventricular myocytes (Magyar et al., 2006; Toyoda et al., 2006) as well as heterologous expression systems such as *Xenopus laevis* oocytes, CHO, COS-7, tsA201 and LM cells (Busch et al., 1994; Abitbol et al., 1999; Unsöld et al., 2000; Toyoda et al., 2006; Wang et al., 2020). The extracellular region of KCNE1 between residues 39-44 was found important in mediating the effect of DIDS on I_{Ks} (Abitbol et al., 1999) and residue K41, located on the extracellular end of KCNE1, was found to be critical in mediating mefenamic acid's activating effect on the fully saturated I_{Ks} channel complex (Wang et al., 2020). Consistent with this idea, when all four WT KCNE1 subunits are replaced with mutant K41C-KCNE1, mefenamic acid up to a concentration of 1 mM is largely ineffective (Figure 1). Unlike in the WT complex, there is little change to the current waveform, slope or $V_{1/2}$ of the G-V plot during activator exposure, suggesting that all the drug binding site(s) on the channel complex are impaired, or that the mechanism of action is disabled, or that the mutation causes a combination of these actions. In the present study, we analysed these activator compound actions using molecular modeling approaches, and complementary mutational analysis. Our data describe the formation of a drug binding pocket between the immediate extracellular residues of KCNE1, S1 and pore residues of two KCNQ1 subunits, stabilized by the presence of structurally different activator compounds. The effect of mutations that negate the activator compound actions is to destabilize the binding pocket itself, reduce drug binding and limit activator residency time on the channel complex.

Mefenamic acid binding site in I_{Ks}.

Using data that suggested an extracellular binding site for mefenamic acid was at the KCNE1-channel interface, MD simulations were used to explore drug-channel interactions further. The drug binding pocket was defined as the extracellular space formed by KCNE1, the domain-swapped S1 helices of one KCNQ1 subunit and the pore/turret region made up of two other KCNQ1 subunits (Figure 2; Supplemental movie S1). Docking and molecular simulations identified a binding site made up of several residues across all of these elements (Figure 2 and Figure 2-figure supplement 2, Table 7). Mutagenesis revealed that most mutations impacted at least one out of the three mefenamic acid effects: $V_{1/2}$ shift, GV-plot slope change, and current waveform change related to slowed deactivation. However, only three of the mutants impacted all of the mefenamic acid actions (Figure 3). These were W323A in KCNQ1, as well as K41C, and A44C (to a lesser extent) in KCNE1, which define key components of the binding pocket for mefenamic acid. That W323 forms an important medial wall of the hydrophobic pocket to which mefenamic acid binds is suggested by data from mutants in which the size of the side chain at this location is systematically made smaller to cysteine and then alanine, which results in diminished effectiveness of the drug, particularly with alanine. This mutation resulted in the detachment of mefenamic acid as well as lower interaction energy of drug-channel complexes during MD simulations (Figure 4A; Supplemental movie S3, Figure 4-figure supplement 1) particularly loss of interaction strength with KCNE1 (Table 7). Likewise, MD simulations revealed detachment of mefenamic acid and reduced interaction energy values for K41C-Mef-I_{ks} complexes (Supplemental movie S2, Figure 4A) again accompanied by loss of interaction strength with KCNE1 (Table 7). The drug also detached from the A44C complex, another mutant with a reduced response in terms of changes in waveform, $V_{1/2}$ and slope. In contrast, while the

free interaction energy of the mutant Y46C channel was significantly decreased in comparison to the WT channel complex (Figure 4A), detachment of mefenamic acid from the mutant channel during 300 ns MD simulations was not observed (Supplemental movie S4, Table 4), though when placed in a lipid environment mefenamic did completely unbind in two out of five 500 ns simulations with Y46C, suggesting a more dynamic interaction in this model (Figure 4-figure supplement 1C). Comparison of energies at specific residues shows a loss of interaction with Q147 and Y148 in S1 with this mutant and a small shift towards the turret at S298 and A300 (Figure 8A; Table 7). While we were not able to obtain good electrophysiological data from Y46C, data from Y46A suggested that mefenamic acid still had potent actions on this mutant. Overall, the combination of binding studies and electrophysiological data indicate that Y46 was not as important for mefenamic acid binding and action as other KCNE1 N-terminal residues.

Common binding site for I_{Ks} activators, mefenamic acid and DIDS.

Although it was suggested that DIDS and mefenamic acid have the same binding site (Abitbol et al., 1999), the extent of overlap was unknown. Initially, we examined the effect of DIDS on WT EQ. Consistent with previous studies (Abitbol et al., 1999; Bollmann et al., 2020) we confirmed that 100 µM DIDS enhanced WT EQ activity (Figure 5B) with a $V_{1/2}$ shift of -46.6 mV and a decrease in the slope of the G-V curve (Table 5). As most I_{Ks} activators are dependent on the KCNQ1-KCNE1 stoichiometry, this more potent effect of DIDS seen in our study may be explained by the higher dose we used and our KCNE1-KCNQ1 linked channel constructs, which ensured fully KCNE1-saturated complexes. Furthermore, due to the large size and complex folds found on the surface of oocytes, higher drug concentrations than those used for cultured cells are often required in order to facilitate a similar effect in both expression systems (Kvist et al., 2011). All previous studies utilized *Xenopus laevis* oocytes, whereas in this study transiently transfected tsA201 cells were used.

The results of in silico experiments, including some binding properties and stability of some mutations, were only partially validated by electrophysiology data, which might be explained by the limitations of the applied methods. Nevertheless, docking analyses revealed that both mefenamic acid and DIDS bind to the same general extracellular inter-subunit interface with some differences in key residues revealed by the electrophysiology data. Mutation Y148 in KCNQ1 to a cysteine (Y148C) or Y46 in KCNE1 to an alanine (Y46A) was found to reduce effects of 100 µM DIDS, particularly in the case of Y148C (Figure 7). These data suggest that DIDS resides deeper in the binding pocket formed by KCNE1 residues and the KCNQ1 pore/S6, and so was less dependent on the side chain of K41 to retain the activator on the channel complex (compare Movies S1 and S6). DIDS associated more strongly with the pore of the Y148C mutant, particularly with V324 and T327, and less across the ps-KCNE1, S1 and turret regions (Table 7). The Y46C mutation resulted in shifts away from ps-KCNE1 and towards the S1 domain (T144, E146 and Q147). In the case of A44C and DIDS, the changes were more subtle, with stronger interactions moving from central residues in KCNE1 (K41, L42 and E43) and S1 (V141 and L142; coloured grey in Figure 8B) to more peripheral residues (I145, E146 and Q147) and lower in the KCNE TMD (A44 and M45; coloured magenta in Figure 8B).

Figure 8 near here

Proposed mechanism of action for mefenamic acid and DIDS.

Molecular modeling and docking revealed that mefenamic acid and DIDS induce conformational changes in the channel upon binding, to shape a binding pocket formed by residues from the external S1 domain, KCNE1 and the pore domain of I_{Ks} (Figure 2, Supplemental movies S1, S5). This cryptic binding pocket is not detectable in the absence of the drug (Figure 4B), which suggests that it has been induced in a similar manner to previous reports of toxin interactions with KcsA-Kv1.3 that induce conformational changes in both the toxin and the channel structure to generate a high-affinity binding site (Lange et al., 2006; Zachariae et al., 2008). Indeed, analysis of the binding site before and after mefenamic acid unbinds, shows the involved residues in the channel complex moving into the space vacated by mefenamic acid (Figure 4 – figure supplement 2), a collapse of the pocket created by the drug-channel interactions.

Given the high modeled binding energy of the channel-drug activated state complexes, ~-39 kcal/mol for mefenamic acid and ~-35 kcal/mol for DIDS, but the relatively low affinities, given the micromolar concentrations required experimentally, we can imagine a dynamic complex where mefenamic acid and DIDS bind/unbind from I_{Ks} at high frequency. Slowed deactivation may therefore be the result of these rapid binding/unbinding interactions slowing the dissociation of the S1/KCNE1/pore domain/drug complex by either providing steric hindrance to dissociation or by stabilizing the activated complex. MD simulations suggest the latter is most likely the case. Crosslinking studies have previously shown that placing cysteines at key locations in the KCNE1 N-terminus, the top of S1 and in S6 can lead to disulfide bond formation and slowing or elimination of deactivation (Chung et al., 2009) similar to what we observe when I_{Ks} is exposed to mefenamic acid and DIDS. Our data indicate that mutation of residue W323 to an alanine would not only destabilize the external S1/KCNE1/pore domain interface (Figure 4D, Table 1) but also eliminate direct hydrophobic contacts which normally occur between the W323 side chain and mefenamic acid, thus facilitating drug dissociation (Figure 2C, Supplemental movie S3).

A reduced interaction with S1 could also conceivably curtail the ability of I_{Ks} activators to slow dissociation of the activated complex and restore faster deactivation rates such that there is no longer an enhanced step current in the presence of the drug, as seen in mutants K41C and A44C for mefenamic acid (Figure 1A and 3A) and A44C, Y46A, W323A and Y148C for DIDS (Figure 7A). Similarly, electrostatic interactions of residue K41, which acts as a lid for the binding pocket for mefenamic acid (Supplemental movie S2), could help stabilize the S1/KCNE1/pore complex by its contacts with drug, which in its turn links the static pore domain to the dynamic voltage-sensor. Mutation of residue K41 to a cysteine may prevent stable I_{Ks} activator binding to the channel by increasing the fluctuations of the external KCNE1 region (Figure 4C) and by reducing the contacts with the drug (Figure 2C). In simulations, mefenamic acid was seen dissociating from mutant K41C, A44C, and W323A channel complexes much more often compared to wild type I_{Ks}, and this explains why the electrophysiological effects of I_{Ks} activators were prevented by amino acid substitutions at these locations (Supplemental movies S2, S3). Shifts in, and particularly reductions in ps-KCNE1 association for both drugs (Table 7) that lead to loss of efficacy suggest that it is the interactions with KCNE1 and S1 that are key to maintaining the activated state. This is similar to the proposal for tight interactions between S1 and KCNE3 in maintaining constitutive activity in KCNE3-associated channels (Kasuya and Nakajo, 2022).

In close proximity to this drug-binding pocket are known gain-of function mutations in S1,
S140G and V141M, which have previously been reported to slow deactivation in the presence of
KCNE1 (El Harchi et al., 2010; Peng et al., 2017). As their current waveforms are qualitatively
similar to those seen after exposure of WT EQ to I_{Ks} activators, we propose that the same
mechanism may underlie the effects of mefenamic acid, DIDS and these I_{Ks} S1 mutations. In
addition, we find that mutation of the neighboring residue, L142, also produces current
waveforms and G-V plots which mirror those seen when WT EQ is treated with 100 µM
mefenamic acid (Figure 3-figure supplement 2) or DIDS. In the cryo-EM structure of KCNQ1-
KCNE3 (Sun and MacKinnon, 2020), residue L142 interacts with KCNE3, whereas V141
interacts with both KCNE3 and the pore domain. This suggests that both the V141M and L142C
mutations could directly alter the interaction of the S1 domain with the pore and/or the position
or movement of KCNE1. The co-evolved interface between the extracellular end of S1 and the
pore domain is thought to be important for bracing the VSD, to allow efficient force transmission
to the pore (Lee et al., 2009), but can also impact permeation, as the S140G and V141M
mutations also enhance rubidium permeation through I_{Ks} complexes (Peng et al., 2017). The
importance of this S1-pore coupling to channel function is supported by mutational analyses of
S1 residues (Chen et al., 2003; Hong et al., 2005; Wang et al., 2011; Campbell et al., 2013) as
well as the L142C mutation examined in this study (Figure 3-figure supplement 2), which all
display current waveforms with instantaneous onset. Incidentally, A300, a residue in the pore
region of KCNQ1, which also interacts with both mefenamic acid and DIDS in MD simulations
(Figure 2-figure supplement 2, Figure 6), has also been implicated in the same gain-of-function
cleft as S140G and V141M (Smith et al., 2007). The V300T mutant has an ~-20 mV shift in $V_{1/2}$
of activation and faster rates of activation (Bianchi et al., 2000), once again showing how a
mutation can mirror the effects of the two activators studied here.
590
Conclusion
Binding of mefenamic acid and DIDS to the extracellular end of KCNE1 and the KCNQ1 S6 and
S1 helices is facilitated by a number of key residues. Residue K41 acts as a "lid" holding
mefenamic acid in place, while residue W323 impacts the size of the binding pocket. Size
reduction mutations of either residue destabilize mefenamic acid binding and ultimately lead to
drug detachment from the channel complex. This explains why, when all four I_{Ks} subunits are
mutated, as in the case of K41C-EQ, EQ-W323A and A44C-EQ, little to no effect of the drug is
seen. The larger drug, DIDS, interacts with many of the same residues but those deeper in the
pocket appear more important than for mefenamic acid. Furthermore, the qualitative similarities
between the S1 mutant channel, EQ-L142C and WT EQ in the presence of 100 µM mefenamic
acid suggest that I_{Ks} activators most likely cause their effects by modulating interactions between
the S1 helix, pore turret, KCNE1 and the S6 helix. Upon binding, both DIDS and mefenamic
acid induce conformational changes in an occult binding pocket and stabilize the
S1/KCNE1/pore complex, which ultimately slows deactivation. The results indicate that this
extracellular inter-subunit interface forms a generalized binding site which different drugs can
access and through which induce common effects on channel activation and deactivation. The
presence of such a binding site and the variable nature of I_{Ks} complex composition may serve as
starting points for future drug development projects targeted at discovering therapeutically-useful
I_{Ks} agonists.
610
611

Materials and Methods

Drug docking and Molecular Dynamic simulations

A model of the I_{Ks} channel complex – termed pseudo-KCNE1-KCNQ1 (ps-I_{Ks}) – was created based on the cryogenic electron microscopy (cryo-EM) structure of KCNQ1-KCNE3 (PDB ID: 6v01) (Sun and MacKinnon, 2020). In this structure, the extracellular residues of KCNE3, R53-Y58, were substituted with homologous KCNE1 residues D39-A44. Conformational sampling was then performed for substituted residues and the lowest free energy conformations were selected for subsequent docking experiments applying a four-dimensional (4D) docking approach which accounts for the flexibility of the receptor site configuration (Bottegoni et al., 2009). For both docking and MD simulations only one ligand was introduced per channel complex to ensure a stable complex throughout sampling. After docking, a receptor ensemble with multiple conformations of the putative binding site region was created via another round of conformational sampling for the external part of ps-KCNE1 and its KCNQ1 neighborhood (8 Å cut-off distance) and generated conformations were used for a new docking iteration. A ligand-channel conformation with the lowest free energy was chosen from the final docking iterations. The docking and conformational sampling as well as substitution of amino acids were performed with ICM-pro 3.8 software (Neves et al., 2012).

The coordinates of mefenamic acid-bound ps-I_{Ks} channel complexes with the lowest free energy were then used for two sets of MD simulations with AMBER and CHARMM force fields (see below). Considering the complexity of the binding site located on the periphery of the pore, psKCNE1 and VSD, docking and subsequent MD simulations were performed for only one binding site (Figure 2D). To keep the voltage-sensing domain (VSD) in its activated state conformation, we restrained the PIP2 molecules in their cryo-EM positions with a force constant of 1000 kJ/mol/nm^2 during MD simulations.

Three 300 ns duration simulations were performed in a water environment with AMBER20 using a ff14SB force field for protein and GAFF/AM1-BCC scheme for the ligand parameterization and calculation of the atomic point charges (Jakalian et al., 2002; Case et al., 2005; Maier et al., 2015). The complex was solvated with TIP3P water models and Na$^+$/Cl$^-$ at 100 mM concentration. The system was minimized and equilibrated in the NVT and NPT ensembles for 10 ns, gradually releasing spatial restraints from the backbone and sidechains. During the last 5 ns only backbone atoms were restrained with a force constant of 50 kJ/mol/nm^2. The same procedure was used for equilibration of mutant complexes after introduction of mutations using ICM-pro 3.8 software. We used the Langevin thermostat with a collision frequency of 2 ps^{-1}, and the reference temperature 310 K, and Monte Carlo barostat with reference pressure at 1 bar (Oliver et al., 1997; Wu et al., 2016). The long-range electrostatic interactions with a cut-off at 10 Å were treated with the Particle Mesh Ewald (PME) algorithm. Bonds involving only hydrogens were constrained with the SHAKE algorithm and a 2 fs integration time step was used.

Five 500 ns duration simulations were performed using a CHARMM 36m force field and ps-I_{Ks} complex inserted into a POPC membrane. The same MD parameters described in our previous study were used for this set of simulations (https://doi.org/10.1038/s41467-022-31526-7).

Trajectories from MD simulations were clustered with TTclust based on ligand and its binding site RMSD (https://doi.org/10.1038/s41467-022-31526-7). The elbow method was used to find the optimum number of clusters. From each cluster, a centroid with the lowest RMSD to all other ligand conformations in the cluster was selected as a representative structure. GROMACS-2021.4 was used for RMSD, RMSF, and H-bond calculation. RMSD was calculated from the

initial position of the ligand after the least squares fit alignment of the protein backbone. For
RMSF calculations only backbone and C-beta atoms were used.
The 2D diagrams and other molecular visualizations were generated by ICM-pro and VMD
software. The MM/PBSA and MM/GBSA methods were used to calculate the free energy of
binding (Miller et al., 2012). We collected 1000 snapshots with equal intervals (every 0.3 ns) if
ligands did not leave the binding site during 300 ns. A schematic representation of the general
workflow for ps- I_{Ks} model construction and drug docking can be found in Figure 2-figure
supplement 1.

Reagents and solutions
DIDS (Sigma-Aldrich, Mississauga, ON, Canada) was prepared as a 50 mM stock solution
dissolved in 100% dimethyl sulfoxide. Stock DIDS solution was diluted in control whole-cell
bath solution to obtain a final DIDS concentration of 100 µM which was perfused onto
mammalian cells for whole-cell experiments. All other reagents and solutions were prepared as
described (Wang et al., 2020). Mefenamic acid (Tocris Bioscience, Oakville, ON, Canada) was
used to activate I_{Ks} at concentrations of 100 µM and 1 mM.

Molecular biology, cell culture and whole cell patch clamp
tsA201 transformed human embryonic kidney 293 cells were first cultured, then plated for
whole-cell experiments and, finally, transfected with Lipofectamine2000 (Murray et al., 2016;
Westhoff et al., 2019; Wang et al., 2020). All mutations were first generated using site-directed
mutagenesis and Pfu turbo, then sequence confirmed. Whole-cell experiments were conducted
24-48 h post transfection. For wildtype (WT) EQ and mutant x-EQ-Y (where "x" denotes a
KCNE1 mutation and "y" denotes a KCNQ1 mutation; for example, K41C-EQ, EQ-W323A),
cells were transfected with a linked KCNE1 and KCNQ1 cDNA (2-3 µg was used) which
assembles as a fully saturated 4:4 ratio of KCNE1 to KCNQ1. All constructs were also co-
transfected with 0.8 µg of GFP to allow transfected cells to be identified. Data were obtained
using an Axopatch 200B amplifier, Digidata 1440A digitizer and pCLAMP 11 software
(Molecular Devices, LLC, San Jose, CA).

Data analysis
Conductance-voltage (G-V) relationships were obtained from the normalized peak of the initial
tail current (G/Gmax) and plotted against the corresponding voltage. G-V plots were fitted with a
Boltzmann sigmoid equation to obtain the voltage at half-maximal activation ($V_{1/2}$) and slope (k)
values (Tables 1, 3 and 5). For each EQ mutant, the change in activation $V_{1/2}$ ($\Delta V_{1/2}=V_{1/2}$ in the
presence of drug-$V_{1/2}$ control) was also determined (Figures 3E and 7D). In some cases, the foot
of the G-V curve did not reach 0 (essentially in the presence of drug) and as a result, the $V_{1/2}$ was
read directly from the normalized plots at the voltage point where the curve crossed the 0.5 value
on the y coordinate. Slowing of tail current decay was used as a measure of mefenamic acid and
DIDS impact on WT and mutated EQ channel complexes. Specifically, the peak to end
difference currents were calculated by subtracting the minimum amplitude of the deactivating
current from the peak amplitude of the deactivating current. The difference current in mefenamic
acid or DIDS was normalized to the maximum control (in the absence of drug) difference current
and subtracted from 1.0 to obtain the normalized response (Figures 3D and 7C).

GraphPad Prism 9 (GraphPad Software, San Diego, CA) was used to analyze all the data. Where
applicable, unpaired t-test or one-way ANOVA followed by the Fisher's least significant
difference (LSD) test was used to determine statistical significance. In all figures ****, ***, **,
* denotes a significance where $p<0.0001$, $p<0.001$, $p<0.01$ and $p<0.05$, respectively. All data in
the figures and tables are shown as mean $\pm$ SD or SEM. Bar graphs showing mean $\Delta V_{1/2}$
(Figures 3E and 7D) were generated by calculating changes in $V_{1/2}$ induced by drug treatment vs.
control in separate cells. $\Delta V_{1/2}$ values reported in the Results were calculated from the mean $V_{1/2}$
values shown in Tables 1 and 3.

710
711
712
Acknowledgements

715   We thank Fariba Ataei for her assistance in cell culture and for making mutants.

716
717
Authorship Contributions

719
Participated in research design:
Jodene Eldstrom, David Fedida, Vitya Vardanyan, Harutyun Sahakyan, Magnus Chan, Marc Pourrier, Yundi Wang

Conducted experiments:
Magnus Chan, Harutyun Sahakyan, Daniel Sastre, Marc Pourrier, Jodene Eldstrom, Yundi Wang, Ying Dou, David Fedida

Performed data analysis:
Magnus Chan, Jodene Eldstrom, Harutyun Sahakyan, Yundi Wang, Vitya Vardanyan

Wrote or contributed to the writing of the manuscript:
David Fedida, Harutyun Sahakyan, Jodene Eldstrom, Magnus Chan, Marc Pourrier, Yundi Wang, Vitya Vardanyan

734
735
736

References

Abitbol I, Peretz A, Lerche C, Busch AE, and Attali B (1999) Stilbenes and fenamates rescue the loss of I KS channel function induced by an LQT5 mutation and other IsK mutants. *EMBO Journal*. **18**:4137-4148.

Ackerman MJ, Priori SG, Willems S, Berul C, Brugada R, Calkins H, Camm AJ, Ellinor PT, Gollob M, Hamilton R, Hershberger RE, Judge DP, Le Marec H, McKenna WJ, Schulze-Bahr E, Semsarian C, Towbin JA, Watkins H, Wilde A, Wolpert C, and Zipes DP (2011) HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies: this document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Europace*. **13**:1077-1109.

Barhanin J, Lesage F, Guillemare E, Fink M, Lazdunski M, and Romey G (1996) K v LQT1 and IsK (minK) proteins associate to form the I Ks cardiac potassium current. *Nature*. **384**:78-80.

Bendahhou S, Marionneau C, Haurogne K, Larroque MM, Derand R, Szuts V, Escande D, Demolombe S, and Barhanin J (2005) In vitro molecular interactions and distribution of KCNE family with KCNQ1 in the human heart. *Cardiovascular Research*. **67**:529-538.

Bianchi L, Priori SG, Napolitano C, Surewicz KA, Dennis AT, Memmi M, Schwartz PJ, and Brown AM (2000) Mechanisms of I Ks suppression in LQT1 mutants. *American Journal of Physiology: Heart and Circulatory Physiology*. **279**:H3003-H3011.

Bollmann E, Schreiber JA, Ritter N, Peischard S, Ho HT, Wunsch B, Strunker T, Meuth S, Budde T, Strutz-Seebohm N, and Seebohm G (2020) 4,4'-Diisothiocyanato-2,2'-Stilbenedisulfonic Acid (DIDS) Modulates the Activity of KCNQ1/KCNE1 Channels by an Interaction with the Central Pore Region. *Cell Physiol Biochem*. **54**:321-332.

Bottegoni G, Kufareva I, Totrov M, and Abagyan R (2009) Four-dimensional docking: a fast and accurate account of discrete receptor flexibility in ligand docking. *Journal of medicinal chemistry*. **52**:397-406.

Busch AE, Herzer T, Wagner CA, Schmidt F, Raber G, Waldegger S, and Lang F (1994) Positive regulation by chloride channel blockers of I sK channels expressed in Xenopus oocytes. *Mol.Pharmacol*. **46**:750-753.

Campbell CM, Campbell JD, Thompson CH, Galimberti ES, Darbar D, Vanoye CG, and George AL, Jr. (2013) Selective targeting of gain-of-function KCNQ1 mutations predisposing to atrial fibrillation. *Circ Arrhythm Electrophysiol*. **6**:960-966.

Case DA, Cheatham TE, 3rd, Darden T, Gohlke H, Luo R, Merz KM, Jr., Onufriev A, Simmerling C, Wang B, and Woods RJ (2005) The Amber biomolecular simulation programs. *Journal of computational chemistry*. **26**:1668-1688.

Chen YH, Xu SJ, Bendahhou S, Wang XL, Wang Y, Xu WY, Jin HW, Sun H, Su XY, Zhuang QN, Yang YQ, Li YB, Liu Y, Xu HJ, Li XF, Ma N, Mou CP, Chen Z, Barhanin J, and Huang W (2003) KCNQ1 gain-of-function mutation in familial atrial fibrillation. *Science*. **299**:251-254.

Chung DY, Chan PJ, Bankston JR, Yang L, Liu G, Marx SO, Karlin A, and Kass RS (2009) Location of KCNE1 relative to KCNQ1 in the I(KS) potassium channel by disulfide cross-linking of substituted cysteines. *Proc.Natl.Acad.Sci.U.S.A*. **106**:743-748.

El Harchi A, Zhang H, and Hancox JC (2010) The S140G KCNQ1 atrial fibrillation mutation
affects 'I(KS)' profile during both atrial and ventricular action potentials. *J Physiol*
Pharmacol. **61**:759-764.
Eldstrom J, and Fedida D (2011) The voltage-gated channel accessory protein KCNE2: multiple
ion channel partners, multiple ways to long QT syndrome. *Expert.Rev.Mol.Med.* **13**:e38.
Eldstrom J, McAfee DA, Dou Y, Wang Y, and Fedida D (2021) ML277 regulates KCNQ1
single-channel amplitudes and kinetics, modified by voltage sensor state. *J Gen Physiol*.
153.
Gofman Y, Shats S, Attali B, Haliloglu T, and Ben-Tal N (2012) How does KCNE1 regulate the
Kv7.1 potassium channel? Model-structure, mutations, and dynamics of the Kv7.1-
KCNE1 complex. *Structure*. **20**:1343-1352.
Hedley PL, Jorgensen P, Schlamowitz S, Wangari R, Moolman-Smook J, Brink PA, Kanters JK,
Corfield VA, and Christiansen M (2009) The Genetic Basis of Long QT and Short QT
Syndromes: A Mutation Update. *Human Mutation*. **30**:1486-1511.
Hong K, Piper DR, Diaz-Valdecantos A, Brugada J, Oliva A, Burashnikov E, Santos-de-Soto J,
Grueso-Montero J, Diaz-Enfante E, Brugada P, Sachse F, Sanguinetti MC, and Brugada
R (2005) De novo KCNQ1 mutation responsible for atrial fibrillation and short QT
syndrome in utero. *Cardiovascular Research*. **68**:433-440.
Jakalian A, Jack DB, and Bayly CI (2002) Fast, efficient generation of high-quality atomic
charges. AM1-BCC model: II. Parameterization and validation. *Journal of computational*
chemistry. **23**:1623-1641.
Jervell A, and Lange-Nielsen F (1957) Congenital deaf-mutism, functional heart disease with
prolongation of the Q-T interval and sudden death. *Am Heart J*. **54**:59-68.
Kasuya G, and Nakajo K (2022) Optimized tight binding between the S1 segment and KCNE3 is
required for the constitutively open nature of the KCNQ1-KCNE3 channel complex.
Elife. **11**.
Kuenze G, Vanoye CG, Desai RR, Adusumilli S, Brewer KR, Woods H, McDonald EF, Sanders
CR, George AL, Jr., and Meiler J (2020) Allosteric mechanism for KCNE1 modulation
of KCNQ1 potassium channel activation. *Elife*. **9**.
Kvist T, Hansen KB, and Brauner-Osborne H (2011) The use of Xenopus oocytes in drug
screening. *Expert Opin Drug Discov*. **6**:141-153.
Lange A, Giller K, Hornig S, Martin-Eauclaire MF, Pongs O, Becker S, and Baldus M (2006)
Toxin-induced conformational changes in a potassium channel revealed by solid-state
NMR. *Nature*. **440**:959-962.
Lee SY, Banerjee A, and MacKinnon R (2009) Two separate interfaces between the voltage
sensor and pore are required for the function of voltage-dependent K(+) channels.
PLoS.Biol. **7**:e47.
Magyar J, Horvath B, Banyasz T, Szentandrassy N, Birinyi P, Varro A, Szakonyi Z, Fulop F, and
Nanasi PP (2006) L-364,373 fails to activate the slow delayed rectifier K+ current in
canine ventricular cardiomyocytes. *Naunyn Schmiedebergs Arch Pharmacol*. **373**:85-89.
Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, and Simmerling C (2015)
ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from
ff99SB. *Journal of chemical theory and computation*. **11**:3696-3713.
Mattmann ME, Yu H, Lin Z, Xu K, Huang X, Long S, Wu M, McManus OB, Engers DW, Le
UM, Li M, Lindsley CW, and Hopkins CR (2012) Identification of (R)-N-(4-(4-
methoxyphenyl)thiazol-2-yl)-1-tosylpiperidine-2-carboxamide, ML277, as a novel,

potent and selective K(v)7.1 (KCNQ1) potassium channel activator. *Bioorg Med Chem Lett*. **22**:5936-5941.

Miller BR, 3rd, McGee TD, Jr., Swails JM, Homeyer N, Gohlke H, and Roitberg AE (2012) MMPBSA.py: An Efficient Program for End-State Free Energy Calculations. *Journal of chemical theory and computation*. **8**:3314-3321.

Mruk K, and Kobertz WR (2009) Discovery of a novel activator of KCNQ1-KCNE1 K channel complexes. *PLoS One*. **4**:e4236.

Murray CI, Westhoff M, Eldstrom J, Thompson E, Emes R, and Fedida D (2016) Unnatural amino acid photo-crosslinking of the channel complex demonstrates a KCNE1:KCNQ1 stoichiometry of up to 4:4. *Elife*. **5**.

Neves MA, Totrov M, and Abagyan R (2012) Docking and scoring with ICM: the benchmarking results and strategies for improvement. *Journal of computer-aided molecular design*. **26**:675-686.

Olesen MS, Nielsen MW, Haunso S, and Svendsen JH (2014) Atrial fibrillation: the role of common and rare genetic variants. *Eur J Hum Genet*. **22**:297-306.

Oliver DS, Cunha LB, and Reynolds AC (1997) Markov chain Monte Carlo methods for conditioning a permeability field to pressure data. *Mathematical Geology*. **29**:61–91.

Peng G, Barro-Soria R, Sampson KJ, Larsson HP, and Kass RS (2017) Gating mechanisms underlying deactivation slowing by two KCNQ1 atrial fibrillation mutations. *Sci Rep*. **7**:45911.

Sanguinetti MC, Curran ME, Zou A, Shen J, Spector PS, Atkinson DL, and Keating MT (1996) Coassembly of K v LQT1 and minK (IsK) proteins to form cardiac I Ks potassium channel. *Nature*. **384**:80-83.

Smith JA, Vanoye CG, George AL, Jr., Meiler J, and Sanders CR (2007) Structural models for the KCNQ1 voltage-gated potassium channel. *Biochemistry*. **46**:14141-14152.

Sun J, and MacKinnon R (2020) Structural Basis of Human KCNQ1 Modulation and Gating. *Cell*. **180**:340-347 e349.

Toyoda F, Ueyama H, Ding WG, and Matsuura H (2006) Modulation of functional properties of KCNQ1 channel by association of KCNE1 and KCNE2. *Biochemical and Biophysical Research Communications*. **344**:814-820.

Unsöld B, Kerst G, Brousos H, HÅbner M, Schreiber R, Nitschke R, Greger R, and Bleich M (2000) KCNE1 reverses the response of the human K + channel KCNQ1 to cytosolic pH changes and alters its pharmacology and sensitivity to temperature. *Pflugers Archiv European Journal of Physiology*. **441**:368-378.

Wang Q, Curran ME, Splawski I, Burn TC, Millholland JM, VanRaay TJ, Shen J, Timothy KW, Vincent GM, De Jager T, Schwartz PJ, Toubin JA, Moss AJ, Atkinson DL, Landes GM, Connors TD, and Keating MT (1996) Positional cloning of a novel potassium channel gene: KVLQT1 mutations cause cardiac arrhythmias. *Nat.Genet*. **12**:17-23.

Wang Y, Eldstrom J, and Fedida D (2020) The I Ks Ion Channel Activator Mefenamic Acid Requires KCNE1 and Modulates Channel Gating in a Subunit-Dependent Manner. *Mol Pharmacol*. **97**:132-144.

Wang Y, Zhang M, Xu Y, Jiang M, Zankov DP, Cui M, and Tseng GN (2012) Probing the structural basis for differential KCNQ1 modulation by KCNE1 and KCNE2. *J.Gen.Physiol*. **140**:653-669.

Wang YH, Jiang M, Xu XL, Hsu KL, Zhang M, and Tseng GN (2011) Gating-related molecular
motions in the extracellular domain of the IKs channel: implications for IKs
channelopathy. *J Membr Biol*. **239**:137-156.
Westhoff M, Eldstrom J, Murray CI, Thompson E, and Fedida D (2019) I Ks ion-channel pore
conductance can result from individual voltage sensor movements. *Proc Natl Acad Sci U*
S A. **116**:7879-7888.
Willegems K, Eldstrom J, Kyriakis E, Ataei F, Dou Y, Russo S, Van Petegem F, and Fedida D
(2022) Structural and electrophysiological basis for the modulation of KCNQ1 channel
currents by ML277. *Nature Communications*. **13**.
Wu X, Brooks BR, and Vanden-Eijnden E (2016) Self-guided Langevin dynamics via
generalized Langevin equation. *Journal of computational chemistry*. **37**:595-601.
Xu X, Jiang M, Hsu KL, Zhang M, and Tseng GN (2008) KCNQ1 and KCNE1 in the IKs
channel complex make state-dependent contacts in their extracellular domains.
J.Gen.Physiol. **131**:589-603.
Xu Y, Wang Y, Zhang M, Jiang M, Rosenhouse-Dantsker A, Wassenaar T, and Tseng GN
(2015) Probing binding sites and mechanisms of action of an I(Ks) activator by
computations and experiments. *Biophys J*. **108**:62-75.
Yu H, Lin Z, Mattmann ME, Zou B, Terrenoire C, Zhang H, Wu M, McManus OB, Kass RS,
Lindsley CW, Hopkins CR, and Li M (2013) Dynamic subunit stoichiometry confers a
progressive continuum of pharmacological sensitivity by KCNQ potassium channels.
Proc.Natl.Acad.Sci.U.S.A. **110**:8732-8737.
Zachariae U, Schneider R, Velisetty P, Lange A, Seeliger D, Wacker SJ, Karimi-Nejad Y,
Vriend G, Becker S, Pongs O, Baldus M, and de Groot BL (2008) The molecular
mechanism of toxin-induced conformational changes in a potassium channel: relation to
C-type inactivation. *Structure*. **16**:747-754.
Zheng Y, Zhu X, Zhou P, Lan X, Xu H, Li M, and Gao Z (2012) Hexachlorophene is a potent
KCNQ1/KCNE1 potassium channel activator which rescues LQTs mutants. *PLoS One*.
7:e51820.

899
900

Footnotes

This research was funded by Natural Sciences and Engineering Research Council of Canada
(grant
#RGPIN-2022-03021), Canadian Institutes of Health Research (#PJT-175024) and Heart and
Stroke Foundation of Canada (#G-21-0031566) grants to D.F. and grants from the Volkswagen
Foundation (#AZ86659 and AZ 92111) to V.V.. MC holds an NSERC CGS-M scholarship.
Y.W. holds a CIHR– Vanier CGS scholarship.

Table 1. Mean $V_{1/2}$ of activation (mV) and slope values (*k*-factor, mV) in the absence and presence of mefenamic acid for fully saturated I_{Ks} channel complexes. A statistical difference in $V_{1/2}$ compared to control is shown as p-value, determined using an unpaired t-test. NS denotes not significant. SEM is denoted by ±.

	Control			**100 µM or 1 mM Mefenamic Acid ***			**p-value**
	V$_{1/2}$	***k-factor***	**n**	**V$_{1/2}$**	***k-factor***	**n**	
Wildtype EQ	25.4 ± 2.4	19.4 ± 1.2	6	-80.3 ± 4.1	41.3 ± 8.4	3	<0.0001
K41C-EQ	15.2 ± 1.1	18.4 ± 1.7	4	11.4 ± 1.0	19.4 ± 0.8	4	<0.05
				16.7 ± 2.0	19.8 ± 1.4	3	NS
L42C-EQ	68.9 ± 1.5	21.5 ± 3.7	3	31.8 ± 0.4	14.8 ± 4.27	3	<0.01
E43C-EQ	70.8 ± 1.8	27.0 ± 2.8	4	46.4±2.1	29.1 ± 1.7	4	<0.01
A44C-EQ	4.1 ± 1.8	17.6 ±1.4	4	-5.6 ± 2.7	18.3 ± 2.0	4	<0.05
Y46A-EQ**	76.4 ± 1.5	57.3 ± 1.6	3	-29.8 ± 4.1	18.9 ± 3.1	3	<0.05
EQ-W323A	47.8 ± 2.7	23.7 ± 2.2	4	33.7 ± 1.2	28.5 ± 3.1	3	<0.05
EQ-W323C	54.0 ± 2.1	22.2 ± 2.1	4	27.5 ± 3.3	25.6 ± 1.8	4	<0.05
EQ-V324A	34.4 ± 2.3	20.7 ± 1.1	6	15.5 ± 2.3	27.9 ± 1.6	5	<0.05
EQ-V324W	41.0 ± 2.4	18.4 ± 0.3	4	27.3 ±6.0	25.5 ± 1.3	4	NS
EQ-Q147C	63.9 ± 5.8	25.4 ± 2.0	4	26.3 ± 5.7	32.7 ± 1.7	4	<0.05
EQ-Y148C	36.8 ± 0.3	20.3 ± 0.4	4	17.5 ± 4.0	36.7 ± 3.7	4	<0.05

* For K41C-EQ, the concentration of mefenamic acid used was either 100 µM (upper row values) or 1 mM (lower row values). For all other constructs, 100 µM mefenamic acid was used.
** An estimation of the activation $V_{1/2}$ was calculated from a right shifted, non-saturating GV curve.

921 **Table 2.** Mean frequency of H-bonds between MEF or DIDS and residues in ps- I_{Ks} during 500
922 ns MD simulations or until drug unbound.

923 MEF357-N1 indicates the nitrogen of aminobenzoic acid; MEF357:O1 and O2 are the oxygens

	Donor	Hydrogen	Acceptor	% Frames with H-Bonds	SEM
MEF	MEF-N1	MEF-H1	GLU43-O	22.5	10.2
	TYR46-N	TYR46-HN	MEF-O2	79.7	4.2
	ILE47-N	ILE47-HN	MEF-O2	7.9	2.7
DIDS	DIDS-O2	DDS357-H10	TRP323-NE1	12.9	2.1
	DIDS-O1	DDS357-H9	GLU43-OE2	25.1	4.0
	DIDS-O1	DDS357-H9	GLU43-OE1	25.2	4.6
	ILE47N	ILE47-HN	DIDS-O5	22.0	3.2
	TYR46-N	TYR46-HN	DIDS-O6	56.8	10.4
	TYR46-N	TYR46-HN	DIDS-O5	7.1	1.8
	TYR299N	TYR299-HN	DIDS-N2	11.9	0.6
	SER298-OG	SER298-HG1	DIDS-N2	26.0	10.6

924 of the aminobenzoic acid; DDS357:O1-O6 are the oxygens of the sulfonic acids; DDS357:N1
925 and N2 are nitrogens of the isothiocyanates. N =5 for each

Table 3. Mean $V_{1/2}$ of activation (mV) and slope values (k-factor, mV) for WT EQ treated with 100 µM mefenamic acid, and untreated mutant EQ-L142C. Interpulse interval used is as indicated. SEM is denoted by ±. An interpulse interval of 7 s created such a dramatic change in the shape of the EQ-L142C G-V plot (see Figure 3- figure supplement 1) that a Boltzmann curve could not be fit and $V_{1/2}$ and k-values are therefore not available.

	$V_{1/2}$	*k-factor*	n
WT EQ + 100 µM mefenamic acid: Interpulse interval 15 s	-80.3 ± 4.1	41.3 ± 8.4	3
WT EQ + 100 µM mefenamic acid: Interpulse interval 30 s	26.7 ± 10.5	66.6 ± 28	8
EQ-L142C control: Interpulse interval 15 s	-80.3 ± 4.5	30.0 ± 2.9	6
EQ-L142C control: Interpulse interval 30 s	-28.7 ± 19	62.5 ± 9.6	3

Table 4. Average free interaction energies of MEF-bound ps-I_{Ks} complexes calculated according to MM/GBSA methods from three independent MD simulation runs. Mean values are in kcal/mol, ± SD. For W323A and K41C mutations, calculations correspond to interval of simulations before the detachment of ligand from molecular complex. Note that unbinding occurred in K41C and W323A in all 3 simulations with 300 ns duration, e.g. Supplemental movies S2 and S3.

Run	Method	Wild Type	W323A	Y46C	K41C
I	GBSA	-39.3 ± 4.1	-13.0 ± 5.1	-31.2 ± 2.9	-10.5 ± 3.8
		-	Unbinding after ~75 ns	-	Unbinding after ~25 ns
II	GBSA	-39.0 ± 3.3	-23.1 ± 3.1	-32.2 ± 3.8	-17.0 ± 2.6
		-	Unbinding after ~120 ns	-	Unbinding after ~70 ns
III	GBSA	-35.5 ± 2.6	-22.6 ± 4.1	-28.8 ± 5.7	-17.2 ± 2.5
		-	Unbinding after ~85 ns	-	Unbinding after ~20 ns

Table 5. Mean $V_{1/2}$ of activation (mV) and slope values (k-factor, mV) in the absence and presence of 100 μM DIDS for fully saturated I_{Ks} channel complexes. A statistical difference in $V_{1/2}$ compared to control is shown as p-value determined using an unpaired t-test. NS denotes not significant. SEM is denoted by ±

	Control			100 μM DIDS			p-value
	$V_{1/2}$	*k-factor*	n	$V_{1/2}$	*k-factor*	n	
Wildtype EQ	30.5 ± 4.3	20.3 ± 0.9	8	-16.1 ± 2.8	25.3 ± 1.9	5	<0.001
K41C-EQ	23.1 ± 3.0	20.2 ± 1.8	3	-1.6 ± 3.7	24.0 ± 3.6	3	<0.01
L42C-EQ	55.4 ± 3.3	19.9 ± 1.5	3	21.4 ±10.7	28.5 ± 2.6	4	<0.05
A44C-EQ	24.1 ± 1.7	29.6 ± 3.3	4	5.6 ± 2.4	17.6 ± 1.8	3	<0.05
*Y46A-EQ	75.2 ± 2.1	59.1 ± 5.3	5	25.8 ± 0.9	39.2 ± 1.4	5	<0.0001
EQ-Y148C	51.8±4.6	23.5±1.7	4	40.0 ± 3.1	26.3 ± 1.5	5	**NS**
EQ-W323A	50.7 ± 3.6	23.9 ± 1.6	4	23.4 ± 4.9	24.1 ± 1.3	4	<0.05

* An estimation of the activation $V_{1/2}$ was calculated from a right shifted, non-saturating GV curve.

Table 6. Average free interaction energies of DIDS-bound ps-I_{Ks} complexes calculated according to MM/PBSA and MM/GBSA methods from three independent MD simulation runs of 300 ns duration. Mean values are in kcal/mol, ± SD. For W323A and Y46C mutations, calculations correspond to interval of simulations before the detachment of ligand from molecular complex. Note that unbinding occurred in W323A in 2 of 3 simulations, and K41C did not unbind but the binding pose shifted – see Supplemental movie S6.

Run	Method	Wild Type	W323A	Y46C	K41C	A44C	Y148C
I	PBSA	-32.3 ± 6.7	-25.7 ± 4.9	-35.5 ± 6.3	-33.5 ± 4.4	-29.2 ± 4.2	-34.1± 4.6
	GBSA	-35.5 ± 4.6	-30.2 ± 4.5	-42.7 ± 5.4	-41.5 ± 5.8	-36.8 ± 2.9	-40.7 ± 4.0
			unbinding after 240 ns		changed binding pose after ~140-150 ns		
II	PBSA	-32.6 ± 4.8	-25.3 ± 5.5	-25.7 ± 6.5	-36.4 ± 4.2	-29.0 ± 4.0	-29.3 ± 3.7
	GBSA	-36.0 ± 4.8	-30.0± 4.8	-32.4 ± 6.4	-31.6 ± 4.0	-34.0 ± 5.7	-31.8 ± 4.7
				unbinding after ~100-132 ns			
III	PBSA	-29.2 ± 8.1	-27.1± 3.9	-38.3 ± 4.8	-35.1 ± 4.3	-31.2 ± 45.0	-35.7 ± 5.8
	GBSA	-34.0 ± 8.9	-35.1± 4.8	-42.5 ± 4.7	-30.8 ± 3.9	-35.6 ± 5.6	-40.3 ± 5.9
			unbinding after ~240 ns				

Table 7. Average free energy of mefenamic acid and DIDS bound WT and mutant ps-I_{Ks} complexes. Values are calculated according to the MM/GBSA method from three independent MD simulation runs and further broken down by residue and channel region. For K41C and W323C, calculations correspond to the interval of simulations before detachment of the ligand from the complex. Values are in kcal/mol. Mutated residues are in bold italics.

	Residue	WT MEF	K41C MEF	Y46C MEF	W232A MEF	WT DIDS	A44C DIDS	Y46C DIDS	Y148C DIDS	W323A DIDS
Total		-38.02	-16.39	-31.40	-20.31	-34.29	-35.40	-39.16	-37.60	-34.33
Drug		-19.69	-8.79	-15.25	-11.65	-17.42	-17.52	-21.21	-18.18	-19.17
Channel		-17.91	-6.52	-16.61	-7.55	-19.21	-18.40	-16.94	-19.21	-15.10
Ch+Drug		-37.60	-15.31	-31.86	-19.20	-36.63	-35.92	-38.15	-37.39	-34.27
Difference		-0.42	-1.08	0.46	-1.11	2.34	0.52	-1.01	-0.21	-0.06
	K41	-1.548	***-0.349***	-1.411	-0.954	-0.639	-0.439	-1.148	-0.283	0.083
	L42	-0.970	-0.895	-0.943	-1.193	-1.061	-0.798	-0.165	-0.320	-0.587
	E43	-2.681	-0.472	-2.257	-0.309	-1.312	-0.660	-0.014	-2.264	-0.705
	A44	-1.906	-0.886	-2.035	-1.102	-1.778	***-2.062***	-1.027	-1.061	-1.780
E1/E3	**M45**	-1.744	-0.163	-1.913	-0.064	-1.756	-2.154	-1.222	-2.810	-1.633
	Y46	-2.391	-0.121	***-2.702***	-0.083	-1.985	-1.797	***-1.245***	-2.377	-1.603
	I47	-0.771	-0.136	-0.412	-0.203	-2.036	-1.993	-1.004	-0.621	-1.528
	L48	-0.010	-0.003	-0.025	0.005	-0.045	-0.100	-0.233	-0.081	-0.035
Sum		**-12.02**	**-3.03**	**-11.70**	**-3.90**	**-10.61**	**-10.00**	**-6.06**	**-9.82**	**-7.79**
	T322	-0.125	-0.047	-0.091	-0.968	-0.200	-0.213	-0.127	-0.152	-0.410
	W323	-2.159	-0.736	-2.226	***-0.932***	-3.312	-3.395	-2.279	-3.566	***-1.590***
Pore	**V324**	-0.644	-0.037	-0.123	-0.047	-0.172	-0.210	-0.111	-2.016	-0.063
	G325	-0.038	-0.006	-0.006	0.017	-0.004	0.000	0.003	-0.048	0.032
	K326	-0.185	-0.015	-0.024	0.008	0.038	0.058	0.091	0.000	0.118
	T327	-0.130	-0.015	-0.020	0.027	-0.040	-0.022	-0.018	-1.298	-0.014
Sum		**-3.28**	**-0.86**	**-2.49**	**-1.89**	**-3.69**	**-3.78**	**-2.44**	**-7.08**	**-1.93**
	V141	-0.061	-0.014	-0.247	-0.006	-0.569	-0.411	-0.660	-0.076	-0.504
	L142	-0.861	-0.434	-1.052	-0.267	-1.525	-0.822	-1.285	-0.460	-1.036
	S143	-0.028	0.004	-0.014	-0.063	-0.008	-0.006	-0.104	0.026	-0.017
	T144	-0.091	-0.006	-0.242	-0.526	-0.015	-0.122	-0.982	-0.002	-0.001
S1	**I145**	0.004	0.005	0.002	0.009	0.003	-0.329	-0.067	-0.009	-0.001
	E146	0.000	0.048	0.056	0.114	0.028	-0.014	-1.601	0.113	0.013
	Q147	-0.230	-0.365	0.006	-0.244	0.012	-0.143	-1.061	-0.974	-0.054
	Y148	-0.565	-1.556	0.006	-0.128	-0.428	-0.336	0.002	***-0.022***	-0.223
Sum		**-1.83**	**-2.32**	**-1.49**	**-1.11**	**-2.50**	**-2.18**	**-5.76**	**-1.40**	**-1.82**
	G297	-0.024	-0.001	-0.010	-0.027	0.016	-0.027	-0.089	-0.001	-0.076
	S298	-0.320	-0.163	-0.419	-0.427	-0.791	-0.848	-1.107	-0.216	-1.026
Turret	**Y299**	-0.103	-0.046	-0.104	-0.052	-0.718	-0.598	-0.563	-0.167	-0.673
	A300	-0.618	-0.183	-0.636	-0.431	-1.130	-1.112	-0.751	-0.652	-1.093
	D301	0.311	0.075	0.248	0.299	0.231	0.172	-0.181	0.136	-0.683
	A302	-0.024	-0.003	-0.011	-0.007	-0.011	-0.020	0.009	-0.008	-0.015
Sum		**-0.78**	**-0.32**	**-0.93**	**-0.65**	**-2.40**	**-2.43**	**-2.68**	**-0.91**	**-3.57**

Supplemental movies 1-4

Mefenamic Acid 150 ns videos

Supplemental movies 5-7

DIDS 250 ns videos

Supplemental movies 1-4. MD simulations at the molecular level of binding of mefenamic Acid to WT EQ, and K41C-EQ, W323A-EQ, Y46C-EQ mutants. Note that videos are longer than the actual simulations, durations stated below.

Movie S1. 80 ns simulation of mef binding to WT ps-KCNE1-KCNQ1.

Movie S2. 150 ns simulation of mef binding to K41C ps-KCNE1-KCNQ1

Movie S3. 110 ns simulation of mef binding to W323A ps-KCNE1-KCNQ1.

Movie S4. 120 ns simulation of mef binding to Y46C ps-KCNE1-KCNQ1

Supplemental movies 5-7. MD simulations at the molecular level of binding of DIDS to WT EQ, K41C-EQ, and W323A-EQ.

Movie S5. 220 ns simulation of DIDS binding to WT ps-KCNE1-KCNQ1.

Movie S6. 250 ns simulation of DIDS binding to K41C ps-KCNE1-KCNQ1

Movie S7. 250 ns simulation of DIDS binding to W323A ps-KCNE1-KCNQ1.

Figure 1. K41C-KCNE1 mutants prevent the agonist effect of mefenamic acid.
(A) Current traces of WT EQ (top) and K41C-EQ (bottom) in the presence of 100 µM mefenamic acid (Mef). A 4 s protocol was used with pulses from -150 mV or higher to +100 mV, in 10 mV steps, followed by a repolarization step to -40 mV for 1 s. Holding potential and interpulse interval were -80 mV and 15 s, respectively. **(B)** G-V plots obtained from WT EQ tail currents (triangles) and K41C-EQ (circles) in the absence (control: black) and presence of Mef (100 µM: grey; 1 mM: light blue). Boltzmann fits were: WT EQ control (n = 6): $V_{1/2}$ = 25.4 mV, k = 19.4 mV; WT EQ 100 µM Mef (n = 3): $V_{1/2}$ = -80.3 mV, k = 41.3 mV; K41C-EQ control (n = 4): $V_{1/2}$ = 15.2 mV, k = 18.4 mV; K41C-EQ 100 µM Mef (n = 4): $V_{1/2}$ = 11.4 mV, k = 19.4 mV; and K41C-EQ 1 mM Mef (n = 3): $V_{1/2}$ = 16.7 mV, k = 19.8 mV. Error bars shown are SEM. **(C)** Summary plot of $V_{1/2}$ change ($\Delta V_{1/2}$) for WT EQ in the presence of 100 µM mefenamic acid, WT EQ vs K41C-EQ in control and K41C-EQ in the presence of 100 µM and 1 mM mefenamic acid. Data are shown as mean ± SEM and unpaired t-test was used. **** denotes a significant $\Delta V_{1/2}$ compared to control where p<0.0001.

Figure 2. MD prediction of mefenamic acid binding site in the ps-I_{Ks} model. **(A)** Pseudo-KCNE1 (ps-KCNE1) used to predict Mef binding site. Extracellular residues of KCNE1 (top), ps-KCNE1 (middle) and KCNE3 (bottom). Below, cartoon topology of the single transmembrane ps-KCNE1 β-subunit and the six transmembrane KCNQ1 α-subunit. S1-S4 transmembrane segments form the voltage sensor domain and S5-S6 form the pore domain. **(B)** Binding pose of Mef (yellow) in external region of the ps-I_{Ks} channel complex obtained with docking (side view). Pore domain residues are blue, ps-KCNE1 subunit red, and the VSD of a neighbouring subunit is in yellow. **(C)** Ligand interaction map of Mef with ps-I_{Ks} from molecular docking. Size of residue ellipse is proportional to the strength of the contact. The distance between the residue label and ligand represents proximity. Grey parabolas represent accessible surface for large areas. Light grey ellipses indicate residues in van der Waals contacts, light green ellipses are hydrophobic contacts, and light blue are H-bond acceptors. Red borders indicate KCNE1, yellow are KCNQ1 VSD, and blue are pore residues. Dashed lines indicate H-bonds. The 2D diagram was generated by ICM pro software with cut-off values for hydrophobic contacts of 4.5 Å and hydrogen bond strength of 0.8. Further details in Materials and Methods. **(D)** Mef binding pose observed in MD simulations in space-fill to highlight pocket formed by external S1 (yellow), S6 (blue) transmembrane domains of KCNQ1, and extracellular region of the ps-KCNE1 subunit (red). This binding conformation is the most frequent binding pose of Mef observed in ~50 % of frames and corresponds to the blue-framed conformation in Figure 2-figure supplement 3.

Figure 2-figure supplement 1. Drug docking and MD simulation workflow. Schematic representation of the general workflow for construction of ps-I_{Ks}, Mef docking to the model ps-I_{Ks} channel complex, and MD simulations (See Materials and Methods).

psKCNE1/KCNQ1 molecular complex with mefenamic acid (KCNE1 D39-A44)

Figure 2-figure supplement 2. Energy decomposition per amino acid for mefenamic acid binding to ps-I_{Ks}. Poisson-Boltzmann Surface Area (MM/PBSA; blue) and Generalized Born Surface Area (MM/GBSA; orange) methods were used to estimate the interaction free energy contribution of each residue in the Mef-bound ps-I_{Ks} complex. Free energy for residues located in ps-KCNE1, pore domain and voltage-sensor domain of KCNQ1, that stand above background, are shown as enlarged panels. Error bars represent ±SD. 1000 frames for each simulation (n=3) were analyzed.

Figure 2-figure supplement 3. Clustering of MD trajectories based on mefenamic acid conformations. **(A)** The clustering dendrogram based on Mef RMSD illustrates the similarity of Mef conformations in MD trajectories. **(B)** Centroids from each cluster with the lowest RMSD compared to all other conformations in the cluster. The colour frame around the panels shows to which cluster the depicted conformation belongs.

Figure 2-figure supplement 4. H-bonds formed between mefenamic acid and protein residues in its binding site during MD simulations. MEF-N1 indicates the nitrogen of aminobenzoic acid; MEF:O1 and O2 are the oxygens of the aminobenzoic acid. Red lines indicate at which time point and between which atoms H-bond were formed.

Figure 3. Current waveform and G-V changes induced by mefenamic acid in binding site mutants.
(A) Current traces from WT EQ and key residue mutants in control (black) and 100 µM Mef (colors).
(B) EQ-W323A and **(C)** EQ-Y148C current traces in control (top) and presence of 100 µM Mef (below). G-V plots in control (black) and presence of 100 µM Mef (colors). Boltzmann fits were: EQ-W323A control (n = 4): $V_{1/2}$ = 47.8 mV, k = 23.7 mV; EQ-W323A Mef (n = 3): $V_{1/2}$ = 33.7 mV, k = 28.5 mV; EQ-Y148C control (n = 4): $V_{1/2}$ = 36.8 mV, k = 20.3 mV; and EQ-Y148C Mef (n = 4): $V_{1/2}$ = 17.5 mV, k = 36.7 mV). Voltage steps were from -80 mV to +100 mV for 4 s, followed by a 1s repolarization to -40 mV. Interpulse interval was 15 s. Error bars shown are ± SEM. **(D)** Summary plot of the normalized response to 100 µM Mef (see Materials and Methods). Data are shown as mean +SEM and one-way ANOVA statistical test was used. **p<0.01 and *p<0.05 denote a significantly reduced response compared to WT EQ. N.S. denotes not significant. **(E)** Change in $V_{1/2}$ ($\Delta V_{1/2}$) for WT EQ and each I_{Ks} mutant in control versus mefenamic acid. Data are shown as mean -SEM and unpaired t-test was used, where *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 indicate significant change in $V_{1/2}$ comparing control to the presence of drug. n-values for mutants in D and E are stated in Table 1.

Figure 3-figure supplement 1. Current waveform and G-V changes induced by mefenamic acid for all binding site mutants. (A)-(H) current traces in control (above) and presence of 100 μM Mef (below) and G-V plots for binding site mutants, as indicated. Cells were held at -80 mV, then pulsed from -90 mV to +100 mV for 4 s followed by -40 mV for 2 s. Error bars shown are ± SEM. Refer to Table 1 for Boltzmann fit values.

Figure 3-figure supplement 2. Augmented activation of EQ-L142C in the absence of mefenamic acid.

(A) EQ-L142C current traces in control solutions. Voltage steps were from -130 mV or higher in 10 mV steps to +100 mV for 4 s, followed by a repolarization step to -40 mV for 1 s. Holding potential was -80 mV. Interpulse interval was as indicated above, 15 s, 30 s and 7 s. **(B)** G-V plot of WT EQ in control (interpulse interval 15 s: black) and in the presence of 100 µM Mef (interpulse interval 15 s: grey closed circle; 30 s: grey open circle) and EQ-L142C in control solutions (interpulse interval 7 s: red; 15 s: blue; 30 s: green). Boltzmann fits were: WT EQ 15 s control (n = 6): $V_{1/2}$ = 25.4 mV, k = 19.4 mV; WT EQ 15 s Mef (n = 3): $V_{1/2}$ = -80.3 mV, k = 41.3 mV; WT EQ 30 s Mef (n = 8); $V_{1/2}$ = 26.7 mV, k = 66.6 mV; EQ-L142C 15 s control (n = 6) $V_{1/2}$ = -80.3 mV, k = 30.0 mV; EQ-L142C 30 s control (n = 3) $V_{1/2}$ = -28.7 mV, k = 62.5. Due to the dramatic change in G-V plot shape, Boltzmann fits were not possible for EQ-L142C with an interpulse interval of 7 s. All mutations are forced saturated I_{Ks} channel complexes. Error bars shown are ± SEM.

Figure 4. K41C, Y46C, and W323A mutant impact on mefenamic acid binding energy and flexibility of external KCNE1 residues.

(A) Average free interaction energy of Mef-bound ps-I_{Ks} complexes calculated using MM/GBSA methods from three independent MD simulation runs. For K41C and W323A mutations, calculations correspond to interval of simulations before the detachment of ligand from molecular complex. * and *** denote significant differences in average free interaction energy compared to WT. Student`s unpaired t-test was used for comparison of groups. **(B)** Surface representation of WT ps-I_{Ks} after removal of Mef. The pore residues are in blue and the VSD of a neighbouring subunit is yellow.

(C-E) Root mean square fluctuations (RMSF) of ps-KCNE residues (Å) in the ps-I_{ks} complex during the last 100 ns of simulations. Three separate MD simulations shown for WT ps-I_{Ks} channel without Mef (black lines) and three for K41C (C, red), W323A (D, blue) after ligand detachment, and Y46C in absence of ligand (E, purple). Dashed lines show average values of RMSF calculated from three simulations. ***p = 0.001; **p =>0.01 *p =>0.05 using an unpaired t-test. GROMACS software was used for RMSF analysis.

Figure 4-figure supplement 1. Root-mean-square deviations (RMSD) of mefenamic acid from its initial position during MD simulations. (A) K41C-, **(B)** W323A-, and **(C)** Y46C-I_{Ks} mutants in the presence of mefenamic acid. Each color represents an individual simulation run.

Figure 4 – figure supplement 2. Result of loss of mefenamic acid from the binding site.
Figure depicts the binding pocket viewed from above (left) and the side (right) during a 1250
ns MD simulation before drug detachment (grey superimposed structure) and after (red
superimposed structure). Mefenamic acid is represented as licorice and colored green.
Snapshots for superimposition were collected every 10 ns. Visible from the snapshots is that
when the drug leaves the binding site (after 500 ns, structures colored red), the N-terminal
residues of ps-KCNE1, as well as W323 and other residues that form the pocket, shift toward
the binding site that mefenamic acid previously occupied.

Figure 5. Effect of DIDS on I_{Ks}. **(A)** Molecular structure of mefenamic acid and DIDS. **(B)** WT EQ current in control (black) and exposed to 100 µM DIDS over time (grey). Pulses were from -80 to +60 mV every 15 s, and current traces are shown superimposed. Lower panel shows no effect on currents from GFP-only transfected cell exposed to 100 µM DIDS over time (grey). **(C)** Current traces from WT EQ in control and presence of 100 µM DIDS as indicated. Pulses were from -80 to +100 mV for 4s, with a 1 s repolarization to -40 mV. Interpulse interval was 15 s. **(D)** Corresponding G-V plot in control (black) and DIDS (grey) from data as shown in panel C. Boltzmann fits were: WT EQ control (n = 8): $V_{1/2}$ = 30.5 mV, k = 20.3 mV; WT EQ in the presence of DIDS (n = 5): $V_{1/2}$ = -16.1 mV, k = 25.3 mV. Error bars shown are ± SEM.

Figure 6. Ligand interaction and energy decomposition per amino acid for DIDS binding to ps-I_{ks}.
(A) Binding pose of DIDS (yellow) in external region of the ps-I_{Ks} channel complex obtained with molecular docking (side view). The residues of the pore domain are coloured in blue, ps-KCNE1 subunit in red, and the voltage-sensor domain of a neighbouring subunit is presented in yellow. **(B)** Ligand interaction map of DIDS with ps-I_{Ks} from molecular docking. Size of residue ellipse is proportional to the strength of the contact. Light grey indicates residues in van der Waals contacts, light green hydrophobic contacts, and light blue are hydrogen bond acceptors. Red borders indicate KCNE1 residues, yellow are KCNQ1 VSD residues, and blue are pore residues. Dashed lines indicate H-bonds. The distance between the residue label and ligand represents proximity. Grey parabolas represent accessible surface for large areas. The 2D diagram was generated by ICM pro software with a cut-off value for hydrophobic contacts 4.5 Å and hydrogen bond strength 0.8. **(C)** Energy decomposition per amino acid for DIDS binding to ps-I_{Ks}. Generalized Born Surface Area (MM/GBSA; orange) and Poisson-Boltzmann Surface Area (MM/PBSA; blue) methods were used to estimate the interaction free energy contribution of each residue in the DIDS-bound ps-I_{Ks} complex. The lowest interaction free energy for residues in ps-KCNE1 and selected KCNQ1 domains are shown as enlarged panels (n=3 for each point). Error bars indicate ±SD.

Figure 6- figure supplement 1. **H-bonds formed between DIDS and protein residues in its binding site during MD simulations.** DDS:O1-O6 are the oxygens of the sulfonic acids; DDS:N1 and N2 are nitrogens of the isothiocyanates. Red lines indicate at which time point and between which atom H-bonds were formed.

Figure 6 -figure supplement 2. Clustering of MD trajectories based on DIDS conformations.
(A) The clustering dendrogram based on RMSD illustrates the similarity of DIDS conformations in MD trajectories. **(B)** Centroids from each cluster with the lowest RMSD compared to all other conformations in the cluster were selected and overlayed for comparison. The color of the DIDS molecule shows to which cluster the depicted conformation belongs .

Figure 6- figure supplement 3. Root-mean-square deviations (RMSD) of DIDS from its initial position during MD simulations. (A) WT I_{ks}, and mutations **(B)** K41C, **(C)** W323A, and **(D)** Y46C complexes. Individual runs are shown in separate colors, n=5 for each.

Figure 7. Binding site mutants important for the action of DIDS.

(A) Current traces from WT EQ and key mutants in the absence (control; black) and presence (colors) of 100 µM DIDS. **(B)** Data and G-V plots in control (black) and DIDS (teal). Boltzmann fits were: K41C-EQ control (n = 3): $V_{1/2}$ = 23.1 mV, k = 20.2 mV; K41C-EQ DIDS (n = 5): $V_{1/2}$ = -1.6 mV, k = 24.0 mV. Voltage steps from a holding potential of -80 mV to +70 mV for 4 s, followed by repolarization to -40 mV for 1 s. Interpulse interval was 15 s. Error bars shown are ± SEM. All calibration bars denote 0.5 nA/0.5 s. **(C)** Summary plot of the normalized response to 100 µM DIDS. Data are shown as mean ± SEM and *p<0.05, **p<0.01, ***p<0.001 denote significant change in mutant versus WT currents (one-way ANOVA, see Materials and Methods). For calculation, see Materials and Methods. **(D)** Change in $V_{1/2}$ ($\Delta V_{1/2}$) for WT EQ and each I_{Ks} mutant in control versus DIDS. Data are shown as mean ± SEM and unpaired t-test was used, where *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 indicate significant change in $V_{1/2}$ comparing control to the presence of drug. n-values for mutants in C and D are stated in Table 5.

Figure 7 – figure supplement 1. Current waveform and G-V changes induced by DIDS for all binding site mutants. (A)-(E) current traces in control (top) and presence of 100 μM DIDS (bottom). For binding site mutants, cells were held at -80 mV, then pulsed to -90 mV to +100 mV for 4 s followed by -40 mV for 2 s. G-V plots in control (black) and presence of 100 μM Mef (colors). Error bars are ± SEM. Refer to Table 5 for Boltzmann fit values.

Figure 8. Subtle differences in activator binding to Y46C- and A44C-ps-IKs mutant channels.
(A) Mefenamic acid bound to the WT ps-IKs channel. Residues that are part of the binding site are shown in stick format colored green, except those residues that were important in the WT channel that had reduced ΔG in Y46C (in grey). Residues that had slight increases in ΔG values in Y46C are shown in magenta (S298, A300). Mefenamic acid is in cyan.
(B) DIDS bound to the WT ps-IKs channel. Residues that are part of the binding site are shown in stick format colored green except those residues that were important in the WT channel that had reduced ΔG in the A44C mutant (in grey). Residues that had slight increases in ΔG values in A44C are shown in magenta. DIDS is in cyan. Images were made with the PyMOL Molecular Graphics System, Version 2.0 Schrödinger, LLC.

Supplemental movies 1-4

Mefenamic Acid 150 ns videos

Supplemental movies 1-4. MD simulations at the molecular level of binding of mefenamic Acid to WT EQ, and K41C-EQ, W323A-EQ, Y46C-EQ mutants. Note that videos are longer than the actual simulations, durations stated below.

Movie S1. 80 ns simulation of mef binding to WT ps-KCNE1-KCNQ1.

Movie S2. 150 ns simulation of mef binding to K41C ps-KCNE1-KCNQ1

Movie S3. 110 ns simulation of mef binding to W323A ps-KCNE1-KCNQ1.

Movie S4. 120 ns simulation of mef binding to Y46C ps-KCNE1-KCNQ1

DIDS 250 ns videos

S5

WT+ DIDS

S6

K41C

S7

W323A

Supplemental movies 5-7. MD simulations at the molecular level of binding of DIDS to WT EQ, K41C-EQ, and W323A-EQ.

Movie S5. 220 ns simulation of DIDS binding to WT ps-KCNE1-KCNQ1.

Movie S6. 250 ns simulation of DIDS binding to K41C ps-KCNE1-KCNQ1

Movie S7. 250 ns simulation of DIDS binding to W323A ps-KCNE1-KCNQ1.