Invertebrate herbivory damage of lowland plant species decreases after an experimental shift to higher altitudes

Karolína Hrubá [1,2], Dagmar Hucková [1], & Jan Klečka [1,*]

[1] *Institute of Entomology, Biology Centre of the Czech Academy of Sciences, České Budějovice, Czech Republic*

[2] *Department of Zoology, Faculty of Science, University of South Bohemia, České Budějovice, Czech Republic*

Keywords: plant-herbivore interactions, enemy release hypothesis, climate change, range shift, altitudinal gradient, elevational gradient, functional traits

* corresponding author: Jan Klečka, e-mail: jan.klecka@entu.cas.cz

Abstract

Many species of plants and animals shift to higher altitudes in response to the ongoing climate warming. Such shifts of species distributions lead to the co-occurrence of species that have not previously lived in the same environment and allow the emergence of novel plant-animal interactions with potential implications for species diversity and community composition in mountain habitats. According to the enemy release hypothesis, the spread of plants in new geographic regions may be facilitated by the reduction of damage caused by natural enemies, such as herbivores. While the importance of this mechanism for the spread of invasive exotic species has been established, it is unclear whether the movement of plants uphill within their native region in response to increasing temperatures may be also facilitated by the reduction of herbivory at sites above their current upper altitudinal limit. In our study, we experimentally tested this hypothesis. We compared herbivory damage of six species of thermophilous lowland plants grown in pots exposed to herbivores at their native sites in the lowland and at sites above their current upper altitudinal limit. As a control, we also measured herbivory damage of six plants growing naturally across the entire range of altitude. We found that lowland plants had reduced herbivory damage when they were moved to highland sites, while herbivory damage of species naturally growing at both altitudes did not differ. Changes of herbivory damage were modulated by leaf dry matter content and to a lesser degree also by specific leaf area and plant height. Our results provide support for the enemy release hypothesis in the novel context of altitudinal range shifts. We conclude that the reduction of herbivory damage may facilitate the spread of plants above their current upper altitudinal limit in response to increasing temperature.

1. Introduction

Global warming leads to large-scale changes of the geographic distribution of many organisms, such as shifts to higher altitudes or towards the poles (Kerr et al., 2015; Freeman et al., 2018; Fazlioglu et al., 2020). Such range shifts lead to the emergence of communities with novel species combinations with important consequences for local biodiversity (Hobbs et al., 2006; 2009). For example, the spread of herbivores to higher altitudes can affect the composition of mountain plant communities (Descombes et al. 2020). The spread of plants uphill can affect mountain species which have to compete for resources (Nomoto & Alexander, 2021) or for pollinators (Hernández-Castellano et al., 2020) with the incoming plants. However, our knowledge of these processes and their long-term ecological impacts is very limited and there is a number of open questions related to factors affecting the establishment and spread of new species in mountain habitats and their direct and indirect effects on mountain communities mediated by interactions with species at other trophic levels (Tomiolo & Ward, 2018; Nomoto & Alexander, 2021).

Causes and ecological consequences of the spread of plants to new geographic regions were studied mostly in invasive exotic species (Vilà et al., 2009; Simberloff et al., 2013). According to the enemy release hypothesis, their spread to new areas can be facilitated by the reduction of damage caused by herbivores or pathogens because the plants escape specialized natural enemies occurring in their native range (Kambo & Kotanen, 2014; Keane & Crawley, 2002; Costan et al., 2022). Herbivores attacking the exotic plants in their invasive range are usually generalists which may not be adapted to cope with chemical defences of the exotic plants (Cappuccino & Carpenter, 2005; Enge et al., 2012). Lower herbivory damage provides a competitive advantage to exotic plants, which facilitates their spread (Keane & Crawley, 2002). Such reduction of herbivory was detected also at smaller scales, e.g., in isolated populations of invasive species at the margins of their geographic range (Kambo & Kotanen, 2014). We hypothesize that a similar reduction of herbivory may facilitate the spread of plants to higher altitudes within their native range in response to climate change.

Altitudinal range shifts are a striking case when the reorganization of local community structure can be observed at a small spatial scale because of the steep temperature gradient in the mountains. The responses of individual species are highly variable (Parmesan & Yohe, 2003). While the optimum altitude for some plant species recorded by Lenoir et al. (2008) shifted by several hundred meters over a few decades, it changed little or even decreased in other cases. Hence, altitudinal range shifts do not represent the movement of entire communities upwards. Rather, species move at a different pace, which creates novel combinations of species that have not coexisted previously in high-altitude habitats. This trend has been accelerating during the last few decades, so mountain communities are increasingly affected by an influx of novel species from lower altitudes (Walther et al., 2005; Steinbauer et al., 2018). However, we do not know well how such shifts affect the interactions of the plants with herbivores, pollinators, or seed dispersers (Tomiolo & Ward, 2018; Hernández-Castellano et al., 2020).

Native species shifting to higher altitudes move over a relatively short distance, but still represent novel species in the high-altitude communities. The main difference compared to invasive exotic species, which come from distant regions, is that mountain communities often contain species which are phylogenetically closely related to plants coming from the lowland. This raises the question whether the shifts of lowland plants to higher altitudes may be facilitated by reduced herbivory as observed in invasive exotic species (Kambo & Kotanen, 2014; Keane & Crawley, 2002; Costan et al., 2022). In addition, the abundance and activity of generalist herbivorous invertebrates often decreases with increasing elevation because of less favourable climate compared to the lowlands (Scheidel & Bruelheide, 2001; Scheidel et al., 2003; Moreira et al., 2017), which may result in lower herbivory damage at higher elevations (Reynolds & Crossley, 1997; Scheidel et al., 2003; Pellisier et al., 2014; Galmán et al., 2018).

Herbivory damage can be affected not only by phylogenetic relatedness and increasing elevation but also by plant functional traits (Pérez-Harguindeguy et al., 2003). Higher leaf dry matter content (LDMC) typical for plants whose leaves are tougher, have higher longevity, and slow growth decreases leaf palatability for herbivores. On the contrary, leaves with high specific leaf area (SLA), which are thinner and softer, are often preferred by herbivores (Reese et al., 2016). Also, the height of a plant is an important indicator of invertebrate herbivory, because some flightless herbivores, in particular slugs and snails, forage close to the ground (Hahn et al., 2011; Hrubá et al., 2022). In general, plant species with tough leaves with high LDMC, low SLA, and taller size are less attractive for herbivores and generally have lower herbivory damage (Reese et al., 2016; Těšitel et al., 2021; Hrubá et al., 2022).

Despite the ubiquitous nature of altitudinal range shifts and their potential impact on plant-animal interactions, we are not aware of any empirical studies focusing on the impact of novel species on the relationships of plants with herbivores, with the exception of a study by Descombes et al., (2020) who transferred herbivorous locusts to higher altitudes. Therefore, we conducted an experimental study where we transferred six species of lowland thermophilous plants to altitudes above their current upper altitudinal limit and tested whether this led to a reduction of their herbivory damage using plants naturally growing across the entire range of altitude as a control. We also aimed to test the effect of several plant traits known to affect herbivory, such as leaf dry matter content (LDMC), specific leaf area (SLA), and plant height (Descombes et al., 2017; Těšitel et al., 2021; Wang et al., 2022). Specifically, we asked the following questions:

1. Do lowland species suffer lower damage by herbivores when experimentally moved to altitudes above their current upper altitudinal limit, as predicted by the enemy release hypothesis?
2. Are plant species naturally growing along the entire range of altitude more damaged by herbivores in the lowland where we expect higher abundance of herbivores?
3. Which plant traits influence herbivory damage and its differences between the lowland and mountain sites?

2. Methods

2.1 Study sites

The study was carried out in the southern part of the Czech Republic. We conducted our experiment at 14 lowland sites (528 – 609 m a.s.l.) in the surroundings of the city of Český Krumlov and six highland sites (856 – 1020 m a.s.l.) in the nearby Šumava Mountains (Table 1, Figure 1). In both cases, the sites were meadows with a relatively high diversity of plant species. The choice of lowland sites was based on the occurrence of plant species selected for the experiment (see below). We aimed to include at least six sites where each plant species grows. Each lowland site contained one to four target species growing there naturally. The highland sites were all located at an altitude higher than the highest known occurrence of the target lowland species in the region based on the Pladias database (Chytrý et al., 2021). Several sites were located on public land inside protected landscape areas with a low level of protection (Blanský Les and Šumava), so conducting the experiments was possible with the permission of their management. The remaining sites were on private land, and we obtained permission for our work from the owners.

Figure 1. The location of the study area and the scheme of our experimental design. (a) The study sites were located in the southern part of the Czech Republic in the area shown in blue. The map was plotted using the rworldmap package in R (South, 2011). Map lines delineate study areas and do not necessarily depict accepted national boundaries. (b) Thermophilous lowland plants were exposed in pots at lowland sites where they grow naturally and at highland sites above their current upper altitudinal limit.

Table 1. The list of study sites in the lowland and the highland. Altitude and GPS coordinates of the centre of each site are provided.

Site ID	Site name	Altitude (m a.s.l.)	GPS
Highland 1	Arnoštov	856	48.89307N, 13.99915E
Highland 2	Horní Sněžná	1020	48.88743N, 13.93954E
Highland 3	Čtyři domy	956	48.89779N, 13.95513E
Highland 4	Strážný 1	982	48.93432N, 13.71026E
Highland 5	Strážný 2	973	48.92935N, 13.73084E
Highland 6	Strážný 3	910	48.92762N, 13.71185E
Lowland 1	Brloh	588	48.92492N, 14.22733E
Lowland 2	Cvičák	570	48.82559N, 14.31408E
Lowland 3	Kájov	553	48.81425N, 14.27224E
Lowland 4	Kalamandra	543	48.82179N, 14.27911E
Lowland 5	Křenovský dvůr	566	48.82147N, 14.24758E
Lowland 6	Lazec	609	48.83522N, 14.26014E
Lowland 7	Liščí hora	573	48.82858N, 14.32326E
Lowland 8	Český Krumlov 1	543	48.82323N, 14.32137E
Lowland 9	Český Krumlov 2	528	48.81838N, 14.30662E
Lowland 10	Český Krumlov 3	530	48.81854N, 14.30744E
Lowland 11	Vyšný 1	579	48.82637N, 14.30220E
Lowland 12	Vyšný 2	581	48.83102N, 14.29278E
Lowland 13	Vyšný 3	590	48.83088N, 14.30778E
Lowland 14	Vyšný 4	578	48.83046N, 14.29691E

2.2 Study species

We selected two groups of plant species (Table 2) for our study. The first group was experimental, grown in pots. This group consists of six thermophilous species from three families (Asteraceae: *Centaurea jacea, C. scabiosa,* and *Senecio jacobea*; Lamiaceae: *Origanum vulgare and Salvia verticillata*; and Plantaginaceae: *Veronica teucrium*) naturally growing only at the lowland sites. The upper altitudinal limit of these species in the region is lower than the altitude of the highland sites chosen for the experiment (Chytrý et al., 2021). We collected seeds of these species at the lowland sites in August – September 2020. We sowed the seeds into germination trays with a mixture of garden soil, compost, and sand (2:1:1 ratio) with the addition of a small quantity of limestone in March 2021. We transferred the seedlings individually into pots (diameter 11 cm) about two weeks after germination and grew them in a greenhouse to prevent access by herbivores.

The second group of plant species, which served as a control in the experiment, consisted of six plant taxa which grow naturally at the lowland as well as the highland sites. These species from five families (Rosaceae: *Fragaria* spp.; Campanulaceae: *Campanula patula*; Plantaginaceae: *Veronica chamaedrys and Plantago lanceolata*; Dipsacaceae: *Knautia arvensis*;

and Hypericaceae: *Hypericum* spp.) (Table 2) were selected on the basis of their high abundance at our focal lowland and highland sites. The selection included four taxa identified at the species level and two identified at the genus level. Specifically, *Fragaria* spp. and *Hypericum* spp. include the mixture of two very similar species (*Fragaria vesca* and *F. viridis*; *Hypericum perforatum* and *H. maculatum*), which we consider equivalent for the purpose of our analyses. The presence of the same species or genera at both lowland and highland sites allowed us to compare the natural level of herbivory between the two altitudes.

Table 2. The list of plant species used and the summary of functional trait data. Origin distinguishes potted plants growing originally in the lowland (Potted) and plants naturally growing at both altitudes (Natural). Mean value of three functional traits measured in the end of the experiment is also provided for each species; separately for the lowland and highland sites. LDMC is expressed as the proportion of dry matter in plant leaves, i.e., dry weight (g) / fresh weight (g). SLA is expressed as the proportion of leaf area (mm^2) / leaf mass (mg).

Species	Origin	Lowland			Highland		
		Height (cm)	SLA (mm^2/mg)	LDMC	Height (cm)	SLA (mm^2/mg)	LDMC
Fragaria spp.	Natural	17.78	20.14	0.3858	17.18	29.25	0.3284
Campanula patula L.	Natural	29.43	36.85	0.2095	35.69	40.38	0.1778
Hypericum spp.	Natural	39.94	26.26	0.3166	32.63	30.87	0.3118
Plantago lanceolata L.	Natural	23.05	14.12	0.1959	23.52	18.77	0.1843
Veronica chamaedris L.	Natural	18.92	30.57	0.2484	21.04	27.55	0.2837
Knautia arvensis (L.) Coult.	Natural	48.14	18.47	0.2062	60.07	18.35	0.2011
Centaurea jacea L.	Potted	10.14	22.72	0.1472	14.02	23.84	0.1335
Centaurea scabiosa L.	Potted	12.83	22.07	0.1330	15.33	23.10	0.1252
Salvia verticillata L.	Potted	9.40	26.83	0.1375	15.24	33.03	0.1209
Veronica teucrium L.	Potted	9.45	37.07	0.1723	7.55	34.94	0.1786
Origanum vulgare L.	Potted	16.35	31.17	0.2251	16.60	35.64	0.2004
Senecio jacobea L.	Potted	6.51	24.51	0.1201	9.54	29.23	0.1045

2.3 Experimental design and data collections

We conducted the field experiment during June−July 2021. We placed 15 individuals of each potted species at six lowland sites where the species grows naturally and at all six highland sites. The pots were distributed in the meadow in three loose clusters per species set >5 m apart, each containing five pots arranged in a 1x1 m square. Each pot was secured by a wire stuck into the ground, which prevented the fall of the pot, e.g., in the case of strong wind. We left the pots with seedlings exposed to herbivores at experimental sites for exactly three weeks.

After three weeks, we measured the height of all potted individuals. Then, we cut the whole individuals and placed them individually in zip lock plastic bags with a piece of filter paper soaked with water. We also randomly chose 15 individuals of the selected plant species naturally growing at both lowland and highland sites (Table 2), measured their height, cut them at ground level, and also put them in zip lock bags. The bags with the plants were stored in polystyrene boxes with ice preventing the wilting of plants and enabling the leaves to fully saturate by water. In the laboratory, we cut all leaves of each plant individual. The leaves were weighed immediately after releasing from zip lock plastic bags to obtain fresh weight of each individual, followed by scanning of the leaves within one day. Afterwards, all scanned leaves were dried in the oven at 70°C for 72 hours (Pérez-Harguindeguy et al., 2003) and dry weight was then measured. We used ImageJ software to measure the total area of the scanned leaf for each individual. If present, the parts of the scanned leaves that appeared to be eaten by herbivores were manually completed to resemble the original shape of the leaves. The corresponding area was calculated using the ImageJ software. Herbivory damage was expressed as 1 − (actual leaf area / leaf area of the completed leaf), i.e., the proportion of the leaf area missing, apparently due to herbivory (Hrubá et al., 2022). We also measured specific leaf area (SLA), and leaf dry matter content (LDMC) following the protocol of Pérez-Harguindeguy et al. (2003). All data are available in Figshare (Hrubá et al. 2023).

2.4 Data analysis

We conducted all analyses in the R software version 4.0.3 (R Core Team 2020) using linear mixed effect models (LMM) using *lme4* (Bates et al., 2015) and *lmerTest* (Kuznetsova et al., 2017) packages. We tested the effect of plant origin (naturally growing vs. potted plants originating in the lowland), altitude (lowland vs. highland), and plant traits on herbivory damage, including interactions shown in Table 3. Herbivory damage, the response variable, was expressed as the proportion of the leaf area eaten by herbivores and was logit-transformed prior to analysis. The explanatory variables plant height and SLA were log-transformed and LDMC, expressed as the proportion of leaf dry weight / fresh weight, was logit-transformed. Plant species and site were used as random effects.

We also used linear mixed effect models (LMM) to compare the values of individual traits between plants from the lowland and the highland and between potted and naturally growing plants. We fitted LMMs with individual traits as response variables, plant origin, altitude, and their interaction as predictors, and plant species and site identity as random factors.

We evaluated statistical significance of fixed effects using F-tests implemented in the R package *lmerTest* (Kuznetsova et al., 2017). To test differences among plant species (a random effect), we compared LMMs with and without plant species using a likelihood ratio test.

3. Results

Herbivory damage was driven by the interaction between the plant origin (potted lowland plants vs. plants naturally growing at both elevations) and the altitude (LMM, interaction plant origin x. altitude: $F = 35.33$, $DF = 1, 1596.7$, $P < 10^{-15}$; Table 3). While herbivory damage was reduced in the highland compared to the lowland in potted plants originating in the lowland, it was independent of the elevation in plants naturally growing at both elevations (Figs. 2 and 3).

In addition, herbivory damage was also affected by plant height, SLA, and LDMC. We detected statistically significant interactions of plant height and SLA with plant origin and altitude (Table 3). However, herbivory damage mostly decreased with increasing SLA, increasing LDMC, and increasing plant height, although with differences in the slope between potted and naturally growing plants and between plants in the lowland and the highland (Fig. 2, Table 3). Mean herbivory damage differed among plant species (comparison of LMMs with and without species as a random effect: $\chi^2 = 349.23$, $P < 10^{-15}$, Fig. 3).

Table 3. The results of the linear mixed effects model testing the dependence of the herbivory damage on plant origin, altitude, and plant traits. The model included species and site as random effects. Model estimates are plotted in Fig. 2. DF = degrees of freedom.

Predictor	F	DF	P
Altitude	1.87	1, 573.7	0.1715
Plant origin	12.43	1, 73.0	**0.0001**
LDMC	12.36	1, 16.3	**0.0028**
SLA	7.57	1, 22.0	**0.0117**
Height	1.98	1, 153.1	0.1615
Altitude x Plant origin	35.33	1, 1596.7	**< 0.0001**
Altitude x LDMC	8.63	1, 662.4	**0.0034**
Altitude x SLA	10.34	1, 1619.8	**0.0013**
Altitude x Height	1.09	1, 1664.2	0.2969
Plant origin x LDMC	0.72	1, 18.8	0.4075
Plant origin x SLA	2.75	1, 25.2	0.1098
Plant origin x Height	10.82	1, 206.4	**0.0012**
Altitude x Plant origin x LDMC	0.03	1, 1007.9	0.8590
Altitude x Plant origin x SLA	25.67	1, 1708.2	**< 0.0001**
Altitude x Plant origin x Height	5.03	1, 1636.3	**0.0250**

Measured plant traits varied among species, according to comparisons of LMMs with and without species as a random effect: height (χ^2 = 2045.2, P <10^{-15}), SLA (χ^2 = 346.79, P <10^{-15}), and LDMC (χ^2 = 911.64, P <10^{-15}). Naturally growing and potted plants collected in the highland were on average taller by 4.52 cm (F = 5.21, DF = 1, 14.728, P = 0.0377), had SLA higher by 3.85 (F = 7.60, DF = 1,15.17, P = 0.0146), and LDMC lower by 0.022 (F = 7.23, DF = 1,15.027, P = 0.0168). Potted plants were on average shorter by 14.51 cm compared to the naturally growing plants we sampled (F = 20.07, DF = 1, 10.01, P = 0.0012), had similar SLA (F = 0.41, DF = 1, 9.98, P = 0.5346), and LDMC lower by 0.107 (F = 11.74, DF = 1, 10.01, P = 0.0065).

Figure 2. Herbivory damage was reduced in the highland in potted plants originating in the lowland and was affected by functional traits. (a) Herbivory damage (the proportion of leaf area consumed by herbivores) of plants naturally growing at both elevations (in green) did not differ between the lowland and the highland sites, while potted plants originating in the lowland (in pink) had lower herbivory damage in the highland. The effect of (b) LDMC, (c) SLA, and (d) plant height on herbivory damage in the lowland and the highland, affected also by the interaction with the plant origin (potted lowland plants vs. plants naturally growing at both elevations). The plot shows estimates (mean ± standard error) from a generalised liner mixed effects model specified in Table 3. The y-axis is displayed using the logit scale.

Figure 3. Herbivory damage of individual species. Box plots show the values of herbivory damage (the proportion of leaf area consumed by herbivores) of individual species at the lowland and highland sites. Data for potted plants originating in the lowland are shown in the upper row and data for plants naturally growing at both elevations are shown in the lower row. Results of linear mixed effects models comparing the herbivory damage in the lowland and the highland separately for each species are also provided (site used as a random effect). The y-axis is displayed using the logit scale.

4. Discussion

4.1 Lowland plants had reduced herbivory damage when moved to higher altitudes

The results of our field experiment show that several thermophilous plant species originally growing in the lowland were significantly less damaged by invertebrate herbivores in the highland, based on the measurement of herbivory damage in individuals grown in pots. These results hold in comparison to control measurements of herbivory damage in plants naturally growing in both the lowland and the highland which revealed no significant difference in natural herbivory between the two altitudes. Our results thus provide support for the enemy release hypothesis (Costan et al., 2022; Kambo & Kotanen, 2014) in a new context, specifically as a potential mechanism facilitating the uphill shift of plants induced by climate change.

The enemy release hypothesis was previously confirmed in exotic species, such as *Impatiens glandulifera* (Hejda & Pyšek, 2006) and *Heracleum mantegazzianum* (Pyšek, 1991) in Europe, which suffer decreased level of herbivory after they spread to new regions. Two main mechanisms of enemy release have been documented. First, a plant can escape its specialised herbivores which occur in its native range but not in the invasive range (Keane & Crawley, 2002; Costan et al., 2022). It can still be consumed by generalised herbivores, but the overall herbivory damage is reduced because of the absence of specialised herbivores. The second mechanism, also called the novel weapon hypothesis, is that the exotic plant may contain defence compounds not present in plants growing in the invaded community, which means that local herbivores are evolutionarily naive to these defence compounds and cannot consume the exotic plant (Cappuccino & Carpenter, 2005; Enge et al., 2012). This mechanism can reduce damage caused also by generalised herbivores.

Both mechanisms can explain the observations that plant species which are phylogenetically more distant from other native species in the community have a higher chance to become invasive (Omer et al., 2022). For this reason, it was previously unclear whether plants shifting their range over short distances, such as plants shifting to higher altitudes (Walther et al., 2005; Lenoir et al., 2008; Steinbauer et al., 2018), may also benefit from enemy release. The short geographic distance travelled by the plants mean that they are likely to encounter herbivores feeding on related species and thus preadapted to attack the new plants. However, our results clearly demonstrate that plants can benefit from reduced herbivory when they move to higher altitudes in the same region. The exact mechanism is not clear, but we assume that it is driven by differences in the composition of the herbivore communities in the highland compared to the lowland, such as lower abundance of slugs (Scheidel & Bruelheide, 2001), who were apparently the main herbivores of *Centaurea jacea* and *C. scabiosa* in the lowland based on the type of leaf damage and slime trails we observed on the plants.

4.2 No difference in herbivory damage of species naturally growing across the entire range of altitude

In comparison to lowland plants in the pots, the naturally growing plant species did not show significant differences in herbivory damage between lowland and highland sites. These species are all common in grassland habitats and occur across the entire climatic gradient from the warmest lowlands to the highest mountain ranges in the Czech Republic (Chytrý et al., 2021). We can thus be confident that the populations we sampled occur in our study area for a long time and can be attacked by generalised herbivores as well as by coevolved specialists (Futuyma & Agrawal, 2009; Maron et al., 2019). The six plant species/genera come from several different families (*Rosaceae, Campanulaceae, Plantaginaceae, Dipsacaceae, Hypericaceae*), which vary in defence mechanisms, and the comparison of herbivory damage of these plants between the lowland and the highland can thus provide a robust estimate of the difference in the overall abundance and activity of herbivorous invertebrates between the two altitudes (Scheidel et al., 2003).

A number of studies conducted in the temperate region reported the decrease of herbivory damage of naturally growing plants with increasing altitude (Reynolds & Crossley, 1997; Scheidel et al., 2003; Pellisier et al., 2014, Galman et al., 2018), although this pattern is not universal and some studies reported no change of herbivory with altitude, increasing herbivory with altitude, or a hump-shaped relationship with a mid-elevation peak of herbivory damage (Scheidel & Bruelheide, 2001; Scheidel et al., 2003; Moreira et al., 2017). It is assumed that when herbivory decreases at higher altitudes it is because of harsh environmental conditions, in particular lower temperature, which limits the abundance and activity of generalist herbivorous invertebrates (Scheidel & Bruelheide, 2001; Scheidel et al., 2003; Moreira et al., 2017).

In our study, we compared only two elevations separated by ca. 400 m of altitude, because our aim was to experimentally test the effect of a realistic altitudinal shift of plant distribution on herbivory, rather than to provide a detailed description of changes of herbivory damage along an altitudinal gradient. Hence, it is likely that the lack of differences in herbivory damage of naturally growing plants we observed between the lowland and highland sites was caused by the relatively small difference in altitude leading to similar total abundance of herbivores.

4.3 The effect of plant traits on herbivory damage

Overall, plant species differed in herbivory damage and all three measured traits, i.e., height, LDMC, and SLA, which reflects their different evolutionary histories translating into specific anti-herbivore strategies (Carmona et al., 2011; War et al. 2018). Differences in functional traits, particularly LDMC, are known to affect leaf palatability to herbivores (Schädler et al., 2003; Descombes et al., 2017) and variation of herbivory damage across different plant species (Schädler et al., 2003; Těšitel et al., 2021; Hrubá et al., 2022).

In our case, both potted and naturally growing plants collected in highlands were on average taller, had lower LDMC, and naturally growing plants also had higher SLA compared to individuals in the lowland. This indicates higher palatability of the highland plants in accordance with the findings of Descombes et al. (2017), who reported a decrease of LDMC and an increase in experimentally measured leaf palatability with the increasing altitude in the Swiss Alps, which may be related to lower abundance of herbivores in mountain habitats (Scheidel & Bruelheide, 2001; Scheidel et al., 2003; Moreira et al., 2017). Plant growth and functional traits may be affected interactively by water and nutrient availability (Cossani et al., 2012; Akter & Klečka, 2022), which we did not measure. However, potted plants were planted in a standardised soil mixture, which suggests that higher humidity in the highland habitats was likely driving these trait differences. Potted plants were on average shorter and had slightly lower LDMC than naturally growing plants, both of which simply follows from the fact that potted plants were younger, grown from seeds sown in the greenhouse ca. 3 months before the start of the experiment.

Herbivory rate was consistently negatively related to LDMC in both habitats and in both naturally growing and potted plants, which corresponds to previous findings (Schädler et al., 2003; Těšitel et al., 2021; Hrubá et al., 2022). Leaves with lower LDMC are usually more palatable for herbivores (Pálková & Lepš, 2008; Descombes et al., 2017), likely because they are easier to chew. However, some herbivores may prefer tougher leaves with higher LDMC, as reported in locusts by Dostálek et al. (2020). In contrast to LDMC, the effect of plant height and SLA on herbivory damage was less consistent and was pronounced mostly in potted plants in the lowland, where a negative relationship between herbivory damage and both plant height and SLA was apparent. Potted plants were on average shorter than naturally growing plants, which makes them more susceptible to flightless herbivores such as slugs and snails, more abundant in the lowland, which can consume whole seedlings but are less capable of reaching taller plants (Hahn et al., 2011), which are thus exposed to a restricted set of invertebrate herbivores.

Similarly to our results, Poorter et al. (2004) and Dostálek et al. (2020) recorded that herbivory damage decreased with increasing SLA. However, in our case, the negative effect of SLA was driven mostly by potted plants at the lowland sites, while the relationship between herbivore damage and SLA in naturally growing plants was weak or absent. This is consistent with the fact that SLA did not significantly influence herbivory damage in previous studies by Whitfeld et al., (2012) and Hrubá et al. (2022). On the other hand, multiple studies reported a positive relationship between SLA and herbivory damage (Dirzo, 1980; Kozlov et al., 2015; Reese et al., 2016). Plants with higher SLA are also more attractive to slugs (Moshgani et al., 2017). Both SLA and LDMC serve as proxies for leaf toughness, which is functionally important for chewing herbivores. However, nutrient content and specific secondary metabolites also affect herbivory, but their effect is taxon-specific (Cornell & Hawkins, 2003; Kelly & Bowers, 2016). In our case, we measured only the total herbivore damage without identifying the herbivores feeding on the plants, so measuring additional traits, such as leaf chemistry, in more detail would not yield more insight into the interspecific differences in herbivory damage.

5. Conclusion

Our main hypothesis, i.e., that lowland plant species would be less damaged by herbivores when they move to altitudes above their current upper altitudinal limit, was confirmed. Our data also show that plants with lower LDMC were less damaged by herbivores, while the effect of SLA and plant height was less pronounced and variable across altitudes. In conclusion, our results suggest that altitudinal shifts in the distribution of plant species may be facilitated by a decrease of herbivory damage in the newly colonized environment, as predicted by the enemy release hypothesis. While reduced herbivory may facilitate the spread of plant species from lower altitudes to mountain habitats, long-term changes of mountain plant communities will also depend on interactions of plants with herbivores shifting to higher altitudes (Descombes et al., 2020). Improved understanding of these processes will be necessary to evaluate the impact of altitudinal shifts of species distribution on biodiversity of mountain habitats.

Acknowledgments

We would like to thank Barbora Hazuková, Daniela Kameníčková, Lydie Plačková and Daniel Jackwerth for their help with experimental work in the field and with processing of leaf samples in the lab. We are grateful to the management of the Blanský Les Protected Landscape Area, Šumava Protected Landscape Area and all landowners for their cooperation and permission to conduct the study. We would also like to thank Ondřej Mudrák for his comments on the experimental design and on the previous version of the manuscript.

References

Akter, A., & Klečka, J. (2022). Water stress and nitrogen supply affect floral traits and pollination of the white mustard, *Sinapis alba* (Brassicaceae). *PeerJ, 10,* e13009. https://doi.org/10.7717/peerj.13009

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software, 67*(1), 1–48. https://doi.org/10.18637/jss.v067.i01

Carmona, D., Lajeunesse, M. J., & Johnson, M. T. (2011). Plant traits that predict resistance to herbivores. *Functional Ecology, 25*(2), 358–367. https://doi.org/10.1111/j.1365-2435.2010.01794.x

Carpenter, D., & Cappuccino, N. (2005). Herbivory, time since introduction and the invasiveness of exotic plants. *Journal of Ecology, 93*(2), 315–321. https://doi.org/10.1111/j.1365-2745.2005.00973.x

Chytrý, M., Danihelka, J., Kaplan, Z., Wild, J., Holubová, D., Novotný, P., ... & Pyšek, P. (2021). Pladias database of the Czech flora and vegetation. *Preslia, 93*(1), 1–87. https://doi.org/10.23855/preslia.2021.001

Cornell, H. V., & Hawkins, B. A. (2003). Herbivore responses to plant secondary compounds: a test of phytochemical coevolution theory. *American Naturalist, 161*(4), 507–522. https://doi.org/10.1086/368346

Cossani, C. M., & Reynolds, M. P. (2012). Physiological traits for improving heat tolerance in wheat. *Plant Physiology, 160*(4), 1710–1718. https://doi.org/10.1104/pp.112.207753

Costan, C.-A., Godsoe, W. K., Bufford, J. L., Marris, J. W. M., & Hulme, P. E. (2022). Can the enemy release hypothesis explain the success of *Rumex* (Polygonaceae) species in an introduced range? *Biological Invasions, 24*(9), 2665–2677. https://doi.org/10.1007/s10530-022-02810-w

Descombes, P., Marchon, J., Pradervand, J.-N., Bilat, J., Guisan, A., Rasmann, S., & Pellissier, L. (2017). Community-level plant palatability increases with elevation as insect herbivore abundance declines. *Journal of Ecology, 105*(1), 142–151. https://doi.org/10.1111/1365-2745.12664

Descombes, P., Pitteloud, C., Glauser, G., Defossez, E., Kergunteuil, A., Allard, P. M., Rasmann, S., & Pellissier, L. (2020). Novel trophic interactions under climate change promote alpine plant coexistence. *Science, 370*(6523), 1469–1473. https://doi.org/10.1126/science.abd7015

Dostálek, T., Rokaya, M. B., & Münzbergová, Z. (2020). Plant palatability and trait responses to experimental warming. *Scientific Reports, 10*(1), 10526. https://doi.org/10.1038/s41598-020-67437-0

Enge, S., Nylund, G. M., Harder, T., & Pavia, H. (2012). An exotic chemical weapon explains low herbivore damage in an invasive alga. *Ecology, 93*(12), 2736–2745. https://doi.org/10.1890/12-0143.1

Dirzo, R. (1980). Experimental Studies on slug-plant interactions: I. The acceptability of thirty plant species to the slug *Agriolimax caruaneae*. *Journal of Ecology, 68*(3), 980–998. https://doi.org/10.2307/2259470

Fazlioglu, F., Wan, J. S. H., & Chen, L. (2020). Latitudinal shifts in mangrove species worldwide: Evidence from historical occurrence records. *Hydrobiologia, 847*(19), 4111–4123. https://doi.org/10.1007/s10750-020-04403-x

Freeman, B. G., Lee–Yaw, J. A., Sunday, J. M., & Hargreaves, A. L. (2018). Expanding, shifting and shrinking: The impact of global warming on species' elevational distributions. *Global Ecology and Biogeography, 27*(11), 1268–1276. https://doi.org/10.1111/geb.12774

Futuyma, D. J., & Agrawal, A. A. (2009). Macroevolution and the biological diversity of plants and herbivores. *Proceedings of the National Academy of Sciences, 106*(43), 18054–18061. https://doi.org/10.1073/pnas.0904106106

Galmán, A., Abdala–Roberts, L., Zhang, S., Berny–Mier y Teran, J. C., Rasmann, S., & Moreira, X. (2018). A global analysis of elevational gradients in leaf herbivory and its underlying drivers: Effects of plant growth form, leaf habit and climatic correlates. *Journal of Ecology, 106*(1), 413-421. https://doi.org/10.1111/1365-2745.12866

Hahn, P. G., Draney, M. L., & Dornbush, M. E. (2011). Exotic Slugs Pose a Previously Unrecognized Threat to the Herbaceous Layer in a Midwestern Woodland. *Restoration Ecology, 19*(6), 786–794. https://doi.org/10.1111/j.1526-100X.2010.00710.x

Hejda, M., & Pyšek, P. (2006). What is the impact of Impatiens glandulifera on species diversity of invaded riparian vegetation? *Biological Conservation, 132*(2), 143–152. https://doi.org/10.1016/j.biocon.2006.03.025

Hernández–Castellano, C., Rodrigo, A., Gómez, J. M., Stefanescu, C., Calleja, J. A., Reverté, S., & Bosch, J. (2020). A new native plant in the neighborhood: effects on plant–pollinator networks, pollination, and plant reproductive success. *Ecology, 101*(7), e03046. https://doi.org/10.1002/ecy.3046

Hobbs, R. J., Arico, S., Aronson, J., Baron, J. S., Bridgewater, P., Cramer, V. A., Epstein, P. R., Ewel, J. J., Klink, C. A., Lugo, A. E., Norton, D., Ojima, D., Richardson, D. M., Sanderson, E. W., Valladares, F., Vilà, M., Zamora, R., & Zobel, M. (2006). Novel

ecosystems: Theoretical and management aspects of the new ecological world order. *Global Ecology and Biogeography*, *15*(1), 1–7. https://doi.org/10.1111/j.1466-822X.2006.00212.x

Hobbs, R. J., Higgs, E., & Harris, J. A. (2009). Novel ecosystems: Implications for conservation and restoration. *Trends in Ecology & Evolution*, *24*(11), 599–605. https://doi.org/10.1016/j.tree.2009.05.012

Hrubá, K., Lisner, A., & Lepš, J. (2022). Effects of phylogeny, traits, and seasonality on invertebrate herbivory damage in a meadow community. *Acta Oecologica, 117*, 103871. https://doi.org/10.1016/j.actao.2022.103871

Hrubá, K., Hucková, D., & Klečka, J. (2023). Data: Invertebrate herbivory damage of lowland plant species decreases after an experimental shift to higher altitudes. Figshare. https://doi.org/10.6084/m9.figshare.22159244

Kambo, D., & Kotanen, P. M. (2014). Latitudinal trends in herbivory and performance of an invasive species, common burdock (Arctium minus). *Biological Invasions, 16*(1), 101–112. https://doi.org/10.1007/s10530-013-0506-z

Keane, R. M., & Crawley, M. J. (2002). Exotic plant invasions and the enemy release hypothesis. *Trends in Ecology & Evolution*, *17*(4), 164–170. https://doi.org/10.1016/S0169-5347(02)02499-0

Kelly, C. A., & Bowers, M. D. (2016). Preference and performance of generalist and specialist herbivores on chemically defended host plants. *Ecological Entomology*, *41*(3), 308-316. https://doi.org/10.1111/een.12305

Kerr, J. T., Pindar, A., Galpern, P., Packer, L., Potts, S. G., Roberts, S. M., Rasmont, P., Schweiger, O., Colla, S. R., Richardson, L. L., Wagner, D. L., Gall, L. F., Sikes, D. S., & Pantoja, A. (2015). Climate change impacts on bumblebees converge across continents. *Science*, *349*(6244), 177–180. https://doi.org/10.1126/science.aaa7031

Kozlov, M. V., Lanta, V., Zverev, V., & Zvereva, E. L. (2015). Background losses of woody plant foliage to insects show variable relationships with plant functional traits across the globe. *Journal of Ecology*, *103*(6), 1519–1528. https://doi.org/10.1111/1365-2745.12471

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest package: Tests in linear mixed effects models. *Journal of Statistical Software*, *82*(13), 1–26. https://doi.org/10.18637/jss.v082.i13

Lenoir, J., Gégout, J. C., Marquet, P. A., de Ruffray, P., & Brisse, H. (2008). A significant upward shift in plant species optimum elevation during the 20th century. *Science*, *320*(5884), 1768–1771. https://doi.org/10.1126/science.1156831

Maron, J. L., Agrawal, A. A., & Schemske, D. W. (2019). Plant–herbivore coevolution and plant speciation. *Ecology*, *100*(7), e02704. https://doi.org/10.1002/ecy.2704

Moreira, X., Glauser, G., & Abdala-Roberts, L. (2017). Interactive effects of plant neighbourhood and ontogeny on insect herbivory and plant defensive traits. *Scientific Reports*, *7*(1), 1–9. https://doi.org/10.1038/s41598-017-04314-3

Moshgani, M., van Mil, H. G. J., & de Jong, T. J. (2017). Plant characters related to slug feeding: A re-analysis of feeding studies using trait data from botanical databases and literature: Plant traits and slug feeding. *Annals of Applied Biology*, *170*(1), 116–126. https://doi.org/10.1111/aab.12321

Nomoto, H. A., & Alexander, J. M. (2021). Drivers of local extinction risk in alpine plants under warming climate. *Ecology Letters*, *24*(6), 1157–1166. https://doi.org/10.1111/ele.13727

Omer, A., Fristoe, T., Yang, Q., Razanajatovo, M., Weigelt, P., Kreft, H., Dawson, W., Dullinger, S., Essl, F., Pergl, J., Pyšek, P., & van Kleunen, M. (2022). The role of phylogenetic relatedness on alien plant success depends on the stage of invasion. *Nature Plants*, *8*(8), 906–914. https://doi.org/10.1038/s41477-022-01216-9

Pálková, K., & Lepš, J. (2008). Positive relationship between plant palatability and litter decomposition in meadow plants. *Community Ecology*, *9*(1), 17–27. https://doi.org/10.1556/ComEc.9.2008.1.3

Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, *421*(6918), 37–42. https://doi.org/10.1038/nature01286

Pellissier, L., Roger, A., Bilat, J., & Rasmann, S. (2014). High elevation *Plantago lanceolata* plants are less resistant to herbivory than their low elevation conspecifics: is it just temperature? *Ecography*, *37*(10), 950–959. https://doi.org/10.1111/ecog.00833

Pérez-Harguindeguy, N., Díaz, S., Vendramini, F., Cornelissen, J. H. C., Gurvich, D. E., & Cabido, M. (2003). Leaf traits and herbivore selection in the field and in cafeteria experiments. *Austral Ecology*, *28*(6), 642–650. https://doi.org/10.1046/j.1442-9993.2003.01321.x

Poorter, L., Plassche, M., Willems, S., & Boot, R. G. A. (2004). Leaf Traits and Herbivory Rates of Tropical Tree Species Differing in Successional Status. *Plant Biology*, *6*(6), 746–754. https://doi.org/10.1055/s-2004-821269

Pyšek, P. (1991). *Heracleum mantegazzianum* in the Czech Republic: Dynamics of spreading from the historical perspective. *Folia Geobotanica et Phytotaxonomica*, *26*(4), 439–454. https://doi.org/10.1007/BF02912779

Reese, A. T., Ames, G. M., & Wright, J. P. (2016). Variation in plant response to herbivory underscored by functional traits. *PLOS ONE*, *11*(12), e0166714. https://doi.org/10.1371/journal.pone.0166714

Scheidel, U., Röhl, S., & Bruelheide, H. (2003). Altitudinal gradients of generalist and specialist herbivory on three montane Asteraceae. *Acta Oecologica*, *24*(5–6), 275–283. https://doi.org/10.1016/j.actao.2003.09.004

Scheidel, U., & Bruelheide, H. (2001). Altitudinal differences in herbivory on montane Compositae species. *Oecologia*, *129*(1), 75–86. https://doi.org/10.1007/s004420100695

Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D. A., Aronson, J., Courchamp, F., Galil, B., García-Berthou, E., Pascal, M., Pyšek, P., Sousa, R., Tabacchi, E., & Vilà, M. (2013). Impacts of biological invasions: What's what and the way forward. *Trends in Ecology & Evolution*, *28*(1), 58–66. https://doi.org/10.1016/j.tree.2012.07.013

South A. (2011). rworldmap: a new R package for mapping global data. *The R Journal 3*(1), 35–43. https://doi.org/10.32614/RJ-2011-006

Steinbauer, M. J., Grytnes, J. A., Jurasinski, G., Kulonen, A., Lenoir, J., Pauli, H., ... & Wipf, S. (2018). Accelerated increase in plant species richness on mountain summits is linked to warming. *Nature, 556*(7700), 231–234. https://doi.org/10.1038/s41586-018-0005-6

Reynolds, B. C., & Crossley Jr, D. A. (1997). Spatial variation in herbivory by forest canopy arthropods along an elevation gradient. *Environmental Entomology, 26*(6), 1232-1239. https://doi.org/10.1093/ee/26.6.1232

R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: https://www.R-project.org/

Těšitel, J., Tahadlová, M., Lepš, J., & Hölzel, N. (2021). Linking insect herbivory with plant traits: Phylogenetically structured trait syndromes matter. *Journal of Vegetation Science, 32*(4), e13061. https://doi.org/10.1111/jvs.13061

Tomiolo, S., & Ward, D. (2018). Species migrations and range shifts: A synthesis of causes and consequences. *Perspectives in Plant Ecology, Evolution and Systematics, 33*, 62–77. https://doi.org/10.1016/j.ppees.2018.06.001

Vilà, M., Bartomeus, I., Dietzsch, A. C., Petanidou, T., Steffan-Dewenter, I., Stout, J. C., & Tscheulin, T. (2009). Invasive plant integration into native plant–pollinator networks across Europe. *Proceedings of the Royal Society B: Biological Sciences, 276*(1674), 3887–3893. https://doi.org/10.1098/rspb.2009.1076

Walther, G. R., Beißner, S., & Burga, C. A. (2005). Trends in the upward shift of alpine plants. *Journal of Vegetation Science, 16*(5), 541–548. https://doi.org/10.1111/j.1654-1103.2005.tb02394.x

Wang, B., Tian, C., & Liang, Y. (2022). Leaf traits-mediated effects of tree diversity on insect herbivory on *Populus laurifolia* in a riparian forest ecosystem. *Forest Ecology and Management, 504*, 119777. https://doi.org/10.1016/j.foreco.2021.119777

War, A. R., Taggar, G. K., Hussain, B., Taggar, M. S., Nair, R. M., & Sharma, H. C. (2018). Plant defence against herbivory and insect adaptations. *AoB plants, 10*(4), ply037. https://doi.org/10.1093/aobpla/ply037

Whitfeld, T. J. S., Novotny, V., Miller, S. E., Hrcek, J., Klimes, P., & Weiblen, G. D. (2012). Predicting tropical insect herbivore abundance from host plant traits and phylogeny. *Ecology, 93*(sp8), S211–S222. https://doi.org/10.1890/11-0503.1