Unistrand piRNA clusters are an evolutionary conserved mechanism to suppress endogenous retroviruses across the *Drosophila* genus

Jasper van Lopik*, Maria-Anna Trapotsi, Gregory J. Hannon, Susanne Bornelöv*,# and Benjamin Czech Nicholson#

Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Cambridge CB2 0RE, UK

* These authors contributed equally
# To whom correspondence should be addressed: Susanne.Bornelov@cruk.cam.ac.uk or Ben.Nicholson@cruk.cam.ac.uk

Keywords: *flamenco*, unistrand piRNA cluster, *Drosophila*, PIWI-interacting RNA, ERV, *Gypsy*-family transposon

SUMMARY

The PIWI-interacting RNA (piRNA) pathway plays a crucial role in preventing endogenous genomic parasites, transposable elements (TEs), from damaging the genetic material of animal gonadal cells. Specific regions in the genome, called piRNA clusters, define each species' piRNA repertoire and therefore its capacity to recognize and silence transposons. In the somatic cells of the *Drosophila melanogaster* ovary, the *flamenco* (*flam*) unistrand cluster is the main source of piRNAs and primarily regulates *Gypsy* family TEs that are able to form virus-like particles and infect neighbouring germ cells. Disruption of the *flam* locus or failure to process *flam* precursor transcripts into piRNAs results in sterility, yet it remains unknown whether this silencing mechanism is employed widely across Drosophilidae. Here, using both synteny-based analyses and *de novo* TE annotation, we identify candidate loci sharing both their organisation and TE targeting repertoire with *flam* in widely divergent *Drosophila* species groups. Small RNA-sequencing validated these loci as bona-fide unistrand piRNA clusters and revealed their predominant expression in somatic cells of the ovary, likely to counter TE mobilisation in this tissue. This study provides compelling evidence of co-evolution between virus-like *Gypsy* family transposons and a host defence mechanism in the form of soma-expressed, unistrand piRNA clusters.

INTRODUCTION

Transposable elements (TEs) are widespread across all domains of life. TEs are broadly categorized based on their structure and mobilisation strategies into DNA transposons, which move via a "cut-and-paste" mechanism, and retrotransposons [1]. Retrotransposons replicate via RNA intermediates and are further subdivided into non-LTR elements, including short interspersed nucleotide elements (SINEs) and long interspersed nucleotide elements (LINEs), and long terminal repeats (LTR) elements, which share similarity to endogenous retroviruses (ERVs). LTR transposons and ERVs both encode *gag* and *pol* open reading frames (ORFs), with ERVs and specialised retroelements (also known as errantiviruses) such as *gypsy* and *ZAM* also possessing an *envelope* (*env*) gene. The *env* gene allows virus-like particle formation and cell-to-cell transposition in addition to the "copy-and-paste" mobilisation mechanism intrinsic to all LTR TEs.

The ability of transposons to mobilise and thereby move or copy from one genomic location to another forms a threat to the genome integrity of their host. This activity, if present in gonadal cells, typically results in sterility [2,3]. Multiple molecular mechanisms have evolved to suppress TE activity, including the HUSH complex, KRAB-zinc finger proteins and the PIWI-interaction RNA (piRNA) pathway [4–9]. The animal-specific piRNA pathway is predominantly expressed in gonadal cells and relies on 23-30 nt small RNAs, mainly derived from transposons and/or discrete genomic loci dubbed piRNA clusters, that associate with PIWI-clade Argonaute proteins [8–10]. While piRNA clusters are widely found throughout the animal kingdom, including examples in human, mice, zebrafish and mosquitos [11–14], their function and content varies with not all species showing enrichment of transposon remnants.

In *Drosophila melanogaster*, piRNA clusters are enriched in TE fragments reflecting past and current transposon burden. Depending on their ability to generate piRNAs from one or both genomic strands, piRNA clusters are classified as either unistrand or dual-strand [10]. Dual-strand clusters, as well as the factors ensuring their transcription and export to piRNA processing sites, are expressed specifically in germ cells and appear to be both fast evolving and *Drosophila*-specific [15–17]. In the somatic compartment of the ovary, however, piRNAs are derived mainly from unistrand clusters. Transcripts from unistrand clusters are similar to canonical mRNAs in that they derive from discrete promoters, are spliced, likely polyadenylated,

and are exported via the canonical Nxf1-Nxt1 machinery [16,18,19].

The *Drosophila* ovary contains somatic follicle cells that encapsulate and support the germ cells, including the oocyte. Through genetic experiments, the *flamenco* (*flam*) locus was identified as the master regulator of *Gypsy* family transposons in the follicle cells of *Drosophila melanogaster*, well before the piRNA pathway was discovered [20,21]. Several studies initially attempted to link protein-coding genes in the *flam* region with *Gypsy* repression, but *flam* was eventually found to be a non-coding RNA gene [10] containing numerous LTR TE fragments, from elements such as *idefix*, *ZAM* and *gypsy*, that are predominantly inserted in the same genomic orientation [22]. Following transcription and processing, *flam* gives rise to a diversity of abundant piRNAs with sequence complementarity to these transposons. Loss of *flam* expression or the failure to process its transcripts into piRNAs results in a near complete loss of *Gypsy*-targeting small RNAs, permitting mobilisation of *Gypsy* elements and ultimately resulting in sterility. Interestingly, consistent with their similarity to ERVs, some *env*-containing *Gypsy* TEs expressed specifically in follicle cells were shown to form virus-like particles able to infect germ cells [21,23–25].

The indispensable role of *flam* in TE regulation in *D. melanogaster* has sparked questions about its evolutionary conservation within the wider *Drosophila* genus. Although *flam* has been identified in closely related species, it appeared to be absent in *D. ananassae* and two members of the *obscura* group [26,27]. Making use of numerous recently released high-quality *Drosophila* genome assemblies [28,29], here we systematically searched for *flam*-like unistrand piRNA clusters within the *Drosophila* subgenera *Sophophora* and *Drosophila,* revealing their widespread presence. Our results highlight their unique characteristic architecture and specificity in regulating somatically active LTR elements, particularly those carrying an *envelope* protein that facilitates transfer to germ cells. Collectively, our study suggests a conserved and essential role of somatically expressed unistrand piRNA clusters in the suppression of ERVs across the entire *Drosophila* genus.

RESULTS

Flam is evolutionary conserved beyond the *melanogaster* subgroup

The unistrand cluster *flam* is the major source of piRNAs in somatic follicle cells of the *D. melanogaster* ovary. *Flam* is characterised by an array of antisense oriented remnants of *Gypsy* family transposons, which gives rise to piRNAs complementary to active TEs [30]. As genome assembly quality improved, the size of the *flam* locus has steadily increased from an original annotation of ~180 kilobases (kb) up to an approximate 650 kb [20,31,32]. Despite its indispensable role in TE control [10,20,21], there has been no evidence of *flam* conservation outside of the *melanogaster* subgroup [26]. We therefore asked whether *flam* conservation could be extended further by analysing a total of 193 Drosophilid genomes from 119 species, including recently published high-quality, long-read genome assemblies [28,29].

We first performed a synteny analysis to locate *flam*-syntenic regions in other species by mapping 20 genes up- and downstream of the *D. melanogaster* cluster to each target genome assembly (**Fig. 1a**). Extensive accumulation of TE insertions predominantly in one genomic orientation was suggestive of a unistrand cluster at the expected genomic location across nearly all studied species within the *melanogaster* subgroup (**Fig. S1**). Within the *melanogaster* subgroup, species, where a syntenic *flam* cluster was not apparent, generally had more fragmented genome assemblies. Additionally, *flam*-syntenic candidate clusters were identified in several species outside the *melanogaster* subgroup, including species from the *suzukii* subgroup, but not the *elegans/rhopaloa* subgroups (**Fig. 1b-d**, **Fig. S2**). These clusters ranged from 227 kb (*D. biarmipes*) to 1,085 kb (*D. subpulchrella*) in size (**Fig. 1e**), and, like their *D. melanogaster* counterpart, displayed a clear strand bias with most transposon fragments inserted opposite to the inferred direction of cluster transcription.

Fig. 1: Identification of *flam* across *Drosophila* species
a, Cartoon showing synteny analysis pipeline. **b**, Genome browser tracks showing the *D. melanogaster* (dm6 genome) *flam* region with transposon annotation by RepeatMasker (RM), displaying some neighbouring genes used for synteny analysis. The pie chart to the right indicates LTR content per strand in the cluster region. **c**, MCScan plot showing gene and *flam* synteny between *D. melanogaster* and *D. biarmipes*. **d**, As in (b) but showing the *D. biarmipes* (GCF_018148935 genome) *flam* region. **e**, Phylogenetic tree (left) representation of the *melanogaster, suzukii*, *takahashii*, *eugracilis*, *ficusphila*, *rhopaloa* and *elegans* subgroups, indicating the respective size of their *flam*-syntenic loci in kb (right).

In conclusion, the *flam* locus likely appeared between 13.3 and 15.1 million years ago (MYA), following the emergence of the *elegans/rhopaloa* subgroups, and was detected in 12 species (**Fig. 1e**). The absence of *flam* beyond these species, despite largely conserved gene synteny in the

region and the widespread presence of *Gypsy*-family elements in Drosophilids, prompted the question whether analogous *flam*-like unistrand piRNA clusters exist elsewhere in the genomes of these species.

D. ficusphila possesses a *flam*-like piRNA cluster

Interspersed repeat content was recently estimated across Drosophilids [28], including the subgenera *Drosophila* and *Sophophora* (to which *D. melanogaster* belongs). We noticed that species of the *Drosophila* subgenus generally appeared to have less repeat content compared to those belonging to the *Sophophora* subgenus. We hypothesised that database-driven repeat annotation commonly performed by RepeatMasker underestimates the transposon abundance in less well studied species. Therefore, we constructed de novo TE annotations using Extensive de

novo TE Annotator (EDTA) [33], and found that this indeed improved repeat annotations (**Fig. S3a**).

The species most closely related to *D. melanogaster* clearly lacking a unistrand syntenic *flam* cluster is *D. ficusphila*. Interestingly, with our novel TE annotations, we found a region enriched in LTR elements at the *flam*-syntenic location, however, without any orientation bias of the transposon insertions (**Fig. 2a-b**). RNA-seq and small RNA-seq (sRNA-seq) from ovaries revealed that this locus is transcribed and that its transcripts are processed into piRNAs. These piRNAs emanate from both genomic strands and show a 1U bias characteristic to this class of small RNAs (**Fig. 2b-c**). Overall, this pattern of piRNA production and the organisation of the TE insertions strongly resembles the architecture of a dual-strand piRNA cluster. The production of piRNAs from dual-strand clusters in *D. melanogaster* germ cells is amplified by the

Fig. 2: Identification of a non-syntenic *flam*-like locus in *D. ficusphila*
a, Genome-wide detection of *flam*-like loci in *D. ficusphila* using a sliding window approach. Predicted LTR content (blue, plus strand; red, minus strand) and total repeat content (grey) is shown across the whole genome (100 kb bins). Arrows indicate the *flam*-syntenic region, and the *de novo* identified *flamlike1* region. **b,** Genome browser tracks showing *D. ficusphila* (GCF_018152265 genome) *flam*-syntenic region (black bar) with transposon annotations by RepeatMasker (RM) and EDTA, and gene annotation by NCBI. Uniquely mapping piRNA (cpm) and total RNA levels (ln(cpm+1)) are presented (green/orange, unique mappers; grey, multi-mappers). Mappability is displayed at the bottom. The pie chart to the right indicates LTR content per strand in the cluster region. **c,** Relative piRNA size distribution of piRNAs mapping sense (light brown) and antisense (dark brown) to *D. ficusphila flam*-syntenic region. **d**, Ping-pong signature for piRNA pairs mapping onto the *flam*-syntenic region. **e**, Same as in (b), but for *D. ficusphila flamlike1* region (blue bar). **f**, Relative piRNA size distribution of sense (light brown) and antisense (dark brown) piRNAs mapping to *D. ficusphila flamlike1* region. **g**, Phasing signature (3' end to 5' end distance) for piRNAs mapping onto *flamlike1*. **h**, Zoom-in on genic region indicating presence of a piRNA cluster in the *flam*-syntenic region in *D. ficusphila*. **i**, Macrosynteny plot indicating gene synteny between *D. melanogaster* and *D. ficusphila* highlighting *flam* (red) and *flamlike1* (blue). **j**, Zoom-in on genic region indicating the absence of a piRNA cluster in *flamlike1*-syntenic region in *D. melanogaster*.

germline-specific ping-pong cycle [10,34], which is characterised by the presence of complementary piRNA pairs overlapping by precisely ten nucleotides counted from their 5' ends. Ping-pong and phasing analysis on piRNAs uniquely derived from the *flam*-syntenic cluster in *D. ficusphila* revealed phasing (**Fig. S3b**) and a strong ping-pong signature (**Fig. 2d**), indicating that this piRNA cluster is likely expressed and processed in the germline.

As *D. ficusphila* appears to possess a dual-strand cluster in place of the *flam* locus, it either lacks somatically expressed LTR transposons or controls these TEs by other means. The presence of *Gypsy* family elements in all investigated genomes strongly indicates that *D. ficusphila* has somatically expressed transposons (**Fig. S3c**). We therefore set out to identify non-syntenic unistrand piRNA clusters in *D. ficusphila* that resemble *flam* in terms of its size, *Gypsy*-family TE content and strong enrichment for transposon insertions to be oriented on one genomic strand.

We calculated LTR transposon content across 100 kb sliding windows to scan the entire genome for putative unistrand piRNA clusters (**Fig. 2a**). This identified a ~560 kb region enriched in LTR TEs predominantly on the plus strand of chrUn_025064091 (**Fig. 2a, e**), hereafter referred to as *flamlike1*. Whole ovary sRNA-seq revealed that *flamlike1* produces piRNAs complementary to TE transcripts, thus resembling the expression pattern of a unistrand cluster. Accordingly, piRNAs mapping to this locus also show a strong 1U bias, robust phasing pattern and a weaker ping-pong signature (**Fig. 2f-g**, **Fig. S3d**).

The presence of a dual-strand cluster at the *flam*-syntenic region and the identification of a *flam*-like unistrand cluster elsewhere in the genome prompted us to investigate whether *D. melanogaster* also has a piRNA cluster at the region syntenic to *flamlike1*. A macrosynteny analysis between *D. ficusphila* and *D. melanogaster* indicated that the *flamlike1*-syntenic region is located on chromosome 2L in *D. melanogaster* (**Fig. 2i**). Closer investigation revealed this to be a purely genic region bearing no TE enrichment, whereas the *flam*-syntenic region shows a piRNA cluster in *D. ficusphila* (**Fig. 2h-j**).

Flam-like clusters repeatedly emerged throughout evolution

The observation of a *flam*-like cluster in *D. ficusphila* raises the question if other, more distant species also carry *flam*-like clusters and whether this is an evolutionary conserved genome feature. To determine whether *flam*-like loci can be readily identified across Drosophilids, we applied the genome-wide scanning approach to all 119 species, including *D. melanogaster*. This re-identified *flam* and several *flam*-syntenic loci across nine species of the *melanogaster* group, including *D. suzukii*, *D. takahashii* and *D. biarrmipes* (**Fig. S4**). Publicly available and our own genomics data confirmed that these loci are expressed and produce abundant piRNAs (**Fig. S5**).

Interestingly, we also identified five additional *flam*-like loci outside of the *melanogaster* group in *D. oshimai*, *D. persimilis*, *D. pseudoobscura*, *D. innubila*, and *D. bifasciata* (**Fig. S6**). Of note, *D. persimilis* and *D. pseudoobscura* are closely related species that diverged less than 2 MYA, and synteny analysis revealed that their *flam*-like loci were syntenic (**Fig. S7a**). We named these loci *flamlike2*, *flamlike3* (both *D. persimilis* and *D. pseudoobscura*), *flamlike4*, and *flamlike5* (**Fig. 3a-d**). Publicly available sRNA-seq data available for *D. persimilis* and *D. pseudoobscura* revealed that *flamlike3*

Fig. 3: Multiple *flam*-like clusters are identified across diverse *Drosophila* species

a, Genome browser tracks showing *D. oshimai flamlike2* region with transposon annotation by EDTA (blue, plus strand; red, minus strand) and gene annotation by NCBI. The pie chart to the right indicates LTR content per strand in the cluster region. **b**, As in (a) but showing the *D. innubila flamlike4* region. **c**, As in (a) but showing the *D. bifasciata flamlike5* region. **d**, Genome browser tracks showing *D. persimilis flamlike3* region with transposon annotation by EDTA and gene annotation by NCBI. Uniquely mapping piRNA (cpm) and total RNA levels (ln(cpm+1)) are presented (green/orange, unique mappers; grey, multi-mappers). Mappability is displayed at the bottom. **e**, As in (d) but showing the *D. pseudoobscura flamlike5* syntenic dual-strand cluster. **f**, Phylogenetic tree representation highlighting *flamlike1*, *flamlike2* and *flamlike4* presence and *flamlike3* and *flamlike5* conservation across *Drosophila* species, with *D. melanogaster flam* as a reference (left). Cluster size is indicated in kb (right).

produces vast amounts of piRNAs that predominantly originate from one genomic strand and were in antisense orientation to LTR transposon transcripts (**Fig. S7b**), which was also confirmed by our own sRNA-seq data (**Fig. 3d**, **Fig. S7b**).

We noted that several *flam*-syntenic regions escaped detection through the genome-wide scanning approach in highly fragmented genome assemblies (**Fig. S8**). This observation, together with the presence of syntenic *flam*-like clusters in the closely related species *D. persimilis* and *D. pseudoobscura* raised the possibility that additional syntenic loci might also be present for other candidate unistrand clusters. Synteny analysis of *flamlike3* showed a widespread conservation of this locus within the *pseudoobscura* subgroup, being present in at least five species (**Fig. 3f**, **Fig. S7b**, **Fig. S9a-d**). Similarly, for another group of four species within the *obscura* subgroup we identified *flamlike5* (**Fig. 3c**, **Fig. S9d-g**, **Fig. S10**). Interestingly, *flamlike5* is syntenic to a dual-strand cluster in the species carrying *flamlike3* (**Fig. 3e**, **Fig. S9a,d**, **Fig. S11**). Similar to the *flam*-syntenic region in *D. ficusphila*, these *flamlike5* syntenic dual-strand clusters are enriched for LTR transposons, but do not show an orientation bias. Thus, large *flam*-like unistrand clusters appear to have emerged at various branchings of the *Drosophila* genus (**Fig. 3f**, **Table S1**).

Flam-like clusters are expressed in the somatic follicle cells of the ovary

Identification of unistrand piRNA clusters across 22 *Drosophila* species (excluding *D. melanogaster*), all displaying similar genomic characteristics, raised the hypothesis that these loci have a somatic function similar to that of *flam*. Supporting this, several of these loci were confirmed to be transcribed predominantly from one strand (**Fig. 3d**, **Fig. S7b**), consistent with expression in somatic cells that, in *D. melanogaster*, lack the machinery needed to express and export transcripts from dual-strand clusters. However, since sRNA-seq from whole ovaries captures a mixture of both somatic and germline piRNAs, it remained uncertain if these unistrand piRNA clusters actually operate in the soma.

To determine whether *flam*-like loci are somatically expressed, we generated and sequenced sRNA-seq libraries that were enriched for somatic follicle cells from nine species, which contain the *flam*-syntenic, *flamlike1*, and *flamlike3* piRNA clusters (**Fig. 4a**). Remarkably, these loci displayed a strong piRNA strand bias (between 7- and 70-fold) (**Fig. 4b**, **Fig. S12a**) and produced a greater fraction of piRNAs in somatic cells as compared with whole ovaries (**Fig. 4c-d**). Interestingly, we found that *D. yakuba* and *D. erecta* deviated from this pattern, displaying somatic expression at the 5' end and germline expression towards the 3' end of the *flam*-syntenic region (**Fig. S12b-c**). Together, this indicates that all identified *flam*-like loci produce antisense piRNAs capable of targeting transposons and that they are expressed primarily in the somatic follicle cells of the ovary.

Somatic unistrand piRNA clusters may be universal across *Drosophila*

Next, we asked whether the soma-enriched sequencing data could be used to detect somatically expressed unistrand piRNA clusters in species where genomic sequence alone was insufficient to identify *flam*-like loci. Therefore, we generated total and soma enrichment sRNA-seq libraries for *D. ananassae*, *D. mojavensis*, and *D. virilis*, representing three distinct groups across the *Drosophila* genus. These libraries, together with our previously generated sequencing data, were used to identify major piRNA-producing loci (>35kb) using proTRAC [35] across 12 species. Up to 14 clusters were identified per genome assembly (mean 7.4 ± 3.7, standard deviation) and classified as somatic, intermediate, or germline based on their somatic expression ratio (**Table S2**). Somatic expression was strongly associated with strand bias, both across *D. ananassae*, *D. mojavensis*, and *D. virilis* (**Fig. 4e-f**, **Fig. S12d-e**) as well as the species investigated originally (**Fig. S13**), except for *D. erecta* and *D. yakuba* as these have both a somatic and a germline component arising from the same genomic location. Notably, these somatic piRNA clusters were all enriched in LTR transposons, suggesting that they might play a conserved role in the repression of these TEs in the soma (**Fig. 4d**, **f**, **Fig. S12d-e**).

The sRNA-seq-assisted identification of somatic *flam*-like piRNA clusters in three additional species (**Fig. 4e**) suggests that somatic unistrand piRNA clusters may be a universal feature across *Drosophila* (**Fig. 4g**). In support of a *flam*-like role, we observed that these newly identified clusters also displayed pericentromeric localisation (**Fig. S14a-c**) and at least one of these cluster appears to be conserved across a much wider group of species (**Fig. S14d-g**). Together, this indicates a strong selective pressure to maintain production of transposon-targeting piRNAs in somatic follicle cells.

Unistrand *flam*-like clusters control ERVs in somatic follicle cells

The canonical function of the piRNA pathway is arguably the suppression of parasitic elements in gonadal cells. At some point, an ancestral LTR transposon of the *Gypsy* family obtained an *env*-like ORF, likely from an insect baculovirus, thereby gaining properties of an endogenous retrovirus (ERV) [36–38]. This allowed the TE, in addition to its ability to mobilize across the genome, to move from cell to cell and infect the oocyte from the surrounding somatic follicle cells [36,37]. *D. melanogaster* counters these ERVs through a somatic piRNA pathway in conjunction with the specific expression of the *flam* locus in this tissue [23,39,40]. Whether the silencing of ERVs by somatic piRNAs is conserved in other species is currently unknown, though it has been noted that the infectious properties of the *env* ORF are present and appear repressed in the *obscura* group [38].

Presently characterised *Gypsy* family TEs are predominantly from *D. melanogaster* and therefore insufficient to use for analysis of distantly related *Drosophila* species. To enable an unbiased analysis, we constructed curated transposon consensus sequences for all 193 analysed genome assemblies (**Fig. S15**). Briefly, raw EDTA and RepeatModeler libraries were combined, filtered and deduplicated to retain a minimal set of TE consensus sequences. These were carefully annotated with respect to their open reading frames (ORFs), repeat classification, and genomic distribution (**Table S3**). Using our TE libraries, we identified genome repeat content of up to 55% (**Fig. S16**), extending previous estimates based on

Fig. 4: *Flam*-like and unistrand piRNA clusters are somatically expressed

a, Illustration of cell composition in total ovary (germline cells surrounded by somatic follicle cells) or after somatic follicle cell enrichment, respectively. **b**, Soma enrichment library piRNAs mapping to piRNA clusters in sense or antisense orientation across 9 species. The largest cluster representative is shown for species with multiple genome assemblies. For all clusters see Fig. S12a. **c**, Scatterplot of piRNA strand bias against piRNA soma enrichment across the indicated piRNA clusters (9 species). The largest cluster is shown for species with multiple genome assemblies. For all clusters see Fig. S12b. **d**, Genome browser tracks of *flamlike1* in *D. ficusphila* showing transposon annotations (EDTA), soma-enriched piRNAs (cpm), total piRNAs (cpm), total RNA expression (ln(cpm+1) scale), and mappability. Transposon annotations are shown in red (minus strand) or blue (plus strand). Sequencing data is shown in green or orange for uniquely mapped reads, and grey for multi-mapping reads. **e**, Analysis across *de novo* identified large piRNA clusters. Clusters were classified as soma, intermediate or germline based on the follicle cell versus total ovary piRNA ratio. Strand bias (log₁₀ scale) is shown across each category. All species shown in Fig. S13. **f**, As in (d) but showing a somatic piRNA cluster in *D. virilis*. See also Fig. S12e. **g**, Phylogenetic tree summarising all studied somatic piRNA clusters across all analysed species (n=119). Cluster size represents the mean across all assemblies, except in the 'others' category, where a single cluster is shown. Species names in bold have sRNA-seq data to validate their expression.

RepeatMasker annotations [28]. Our analysis further revealed a strong correlation between genome assembly size and interspersed repeat content (**Fig. S17a**). LTR elements dominated the repeat landscape across most species (**Fig. 5a**, **Fig. S17b-d**), with the *Gypsy* family generally most abundant, representing about 25% of all interspersed repeats (**Fig. S17d**). The *Gypsy* family was also the most frequent in terms of the number of subfamilies (**Fig. S18**) and genomic copies (**Fig. S19**). As expected, *env* ORFs were frequently found in *Gypsy* elements (**Fig. 5b**), but not in the other LTR families *Copia* and *Pao*. Notably, *env* ORFs were found in *Gypsy* elements across all species (**Fig. S17e**), suggesting that its acquisition is ancestral to the

Drosophila genus. It is therefore likely that *Gypsy* ERVs mobilise from somatic follicle cells into the germline across all *Drosophila* species.

Next, we compared our curated TE libraries to 180 well-characterised *Drosophila* transposon subfamilies (**Fig. S20**), revealing that many were present in multiple species. Focusing on TEs well-known in *D. melanogaster*, we found convincing evidence of vertical transmission reflecting their phylogenetic relationship within the *melanogaster* subgroup, but also evidence of horizontal transfer to *D. ercepeae*, *D. bocqueti*, *D. pruinosa*, and the *Zaprionus* genus (**Fig. S20**). Importantly, these observations were supported by cross-species alignment of our sRNA-seq data

Fig. 5: Unistrand *flam*-like clusters are selectively enriched for antisense ERVs
a, Boxplot showing fraction of interspersed repeat content for the indicated repeat classes. Each data point represents one species (n=119). Species with multiple genome assemblies are represented by their mean. **b**, Boxplot showing the number of subfamilies detected per LTR family with either *gag* + *pol* (left) or *gag* + *pol* + *env* (right) ORFs. Each data point corresponds to one species (n=119). Species with multiple genome assemblies are represented by their mean. **c**, Bar plot showing LTR content per strand (left, antisense; right, sense) across all annotated *flam*-like clusters. *Gypsy* elements are shown in red (antisense) or blue (sense) and other LTR elements are shown in grey. Clusters are grouped by synteny to the right. Species and genome assembly (alphabetically sorted) are indicated to the left. **d**, Boxplot displaying *Gypsy* versus other LTR coverage against the genomic average across different unistrand clusters. Each point corresponds to one cluster in one genome assembly. **e**, Scatterplot showing *Gypsy* enrichment against *env* enrichment in unistrand clusters from the indicated species (see "Cluster content analyses" in the Methods for details). Only high-quality LTR transposons are included in the analysis (both *gag* and *pol* and at least one good genomic hit).

(**Fig. S21**), suggesting that similarities in piRNA populations between species is indicative of shared transposon burden.

Analysis of repeat content across all *flam*-like loci revealed a strong enrichment for LTR transposons inserted in antisense orientation (**Fig. 5c**). This is expected, since an LTR TE enrichment and strand bias was part of the criteria used to define the loci. However, when analysing the LTR transposon families, we observed that this strand bias was driven exclusively by *Gypsy* elements and absent for other LTR transposons (**Fig. 5c**). These unistrand clusters thus produce piRNAs specifically targeting *Gypsy* elements in follicle cells. To determine whether *Gypsy* family elements are predominantly occupying these loci, we compared the cluster composition against the mean genomic coverage per LTR family, observing a strong enrichment of *Gypsy* transposons while other LTRs were simultaneously depleted (**Fig. 5d**).

Finally, to determine whether the unistrand clusters specifically capture *Gypsy* elements, we analysed our TE libraries at subfamily level (**Fig. 5b**). For this, we calculated a *Gypsy* enrichment ratio as the conditional probability that an LTR transposon captured by the locus was a *Gypsy*-family element, divided by the probability that an LTR element not captured by the cluster was a *Gypsy* family transposon. In total, 19 out of 21 unistrand clusters, including *D. melanogaster flam*, displayed a positive

enrichment, indicative of a selective capture of *Gypsy* TEs over other LTR transposons (**Fig. 5e**, **Fig. S17f**). Notably, some *Gypsy* subfamilies either never gained or subsequently lost their *env* ORF (**Fig. 5b**). These are expected to have a germline expression pattern. One such example is *Burdock* in *D. melanogaster*, which shares origin with other ERVs, but has lost its *env* and is now expressed exclusively in the germline [27,37]. This suggests that transposons possessing an *env* ORF should be controlled by a somatic piRNA cluster. To test this model, we calculated an *env* enrichment ratio, defined as the conditional probability that a *Gypsy* TE captured by the cluster had the *env* ORF present, divided by the probability that a *Gypsy* not captured possessed an *env* domain. This analysis revealed a selective capture of *env*-containing *Gypsy* elements across 17 out of 21 unistrand clusters (**Fig. 5e**, **Fig. S17g**), including *D. melanogaster flam*. We note that this analysis required high-quality transposon consensus sequences and therefore some of the exceptions may be due to annotation artefacts. However, we speculate that the *flam*-syntenic region in *D. takahashii*, showing neither *Gypsy* or *env* enrichment and displaying limited soma enrichment (**Fig. 4c**) may be in the process of losing its *flam*-like function and converting into a germline-expressed dual-strand piRNA cluster.

Together, our analyses reveal that unistrand *flam*-like piRNA clusters selectively acquired *env*-containing *Gypsy*

family transposable elements in antisense orientation, supporting a conserved role for the piRNA pathway in controlling ERVs in follicle cells across the *Drosophila* genus.

DISCUSSION

New TE insertions can only propagate to the next generation when they are established in the germ cells. Acquisition of a retroviral *env* ORF by an ancestral *Gypsy* family retrotransposon is believed to have transformed it into an endogenous retrovirus. Thereby it gained the means of invading the germline from the somatic follicle cells of the ovary, evading silencing by the germline piRNA pathway. In *D. melanogaster*, a somatic piRNA pathway fuelled by *flam*, the main source of piRNAs in follicle cells, protects the germline against these deleterious elements [20,21,41]. Previous observations of *flam* were limited to species within the *melanogaster* subgroup [18,26,27,30]. Our identification of *flam* within the *suzukii* subgroup together with the absence of any piRNA cluster at the *flam*-syntenic region in and beyond the *rhopaloa* subgroup, places the emergence of *flam* between 13.3 and 15.1 MYA, long before the *melanogaster* subgroup separated from the remainder of the *melanogaster* group around 6.8 MYA.

Despite the absence of a *flam* locus in the *rhopaloa* subgroup and beyond, we identified several unistrand piRNA clusters with characteristics similar to *flam*. Across 24 *Drosophila* species, we identified five additional *flam*-like loci, including *flamlike3* and *flamlike5* present in several species across the *obscura* group. Of note, we confirmed that *flamlike3* produces abundant somatic piRNAs despite the lack of *fs(1)Yb*, a key factor for efficient *flam* processing [42,43], across the *obscura* species group (**Fig. S23**). This supports recent observations of mechanistic divergence, including the loss of *fs(1)Yb* or *Ago3*, across some *Drosophila* species [44]. Analysis of the cluster content revealed that all five *flam*-like loci specifically capture remnants of ERVs in the antisense orientation, consistent with strong selective pressure to generate piRNAs against these elements in somatic cells. We speculate that all *Drosophila* species use *flam*-like piRNA clusters in a somatic branch of the pathway that specifically evolved to repress ERVs. In support, we identified somatic unistrand piRNA clusters in all species sequenced in this study and did not observe any *Drosophila* species lacking the presence of *env*-containing *Gypsy* family elements.

Interestingly, all five piRNA generating loci show substantial size, comparable to their *D. melanogaster* counterpart. Simulations also show that a single large piRNA cluster in a region without recombination is the most efficient way to stop transposon invasion [45]. In agreement with this model, individual dual-strand piRNA clusters may be dispensable for transposon control and hence show poor conservation even across closely related *Drosophila* species [17,46]. Of note, however, the machineries responsible for the dual-strand piRNA cluster expression and export are essential for transposon control and fertility [16,31,47–51]. Most *flam*-like unistrand clusters reported here follow the pattern of a single large locus, supporting the above model. We hypothesise that in addition to being the most efficient way of stopping TE invasion, this may enforce rigid natural selection, as disruption of *flam*-like

piRNA clusters likely result in sterility. The recurring presence of unistrand clusters across the *Drosophila* genus strongly argues for an essential role of these loci, perhaps as a means to produce piRNAs in the soma without access to the germline piRNA expression and export machinery.

Surprisingly, we detected several cases of synteny between unistrand and dual-strand clusters. The *flam*-syntenic region in *D. ficusphila* harbours a dual-strand cluster and several *pseudoobscura* subgroup species have a dual-strand cluster at the *flamlike5*-syntenic location. Our data hint towards a conversion over time of unistrand loci into dual-strand clusters, or vice versa, although it could also reflect a propensity to repeatedly form piRNA clusters at specific genomic locations, likely those with low recombination rate and low selective pressure [46]. In support of the conversion model, signatures of germline *flam* expression have been observed in *D. simulans* and *D. mauritiana* [52]. Additionally, we observed germline expression towards the 3' ends of *D. erecta* and *D. yakuba* *flam*-syntenic clusters. Together, this suggests that unistrand piRNA clusters can lose their somatic identity over time. Some dual-strand clusters thus could be vestigial, where the locus was retained, and its extant function was either acquired after or during the transformation. It will be interesting to determine whether the *D. ficusphila* *flam*-syntenic dual-strand cluster was initially unistranded and lost its promoter, or whether it emerged as a dual-strand cluster that gained a promoter in the *suzukii/melanogaster* ancestor.

Although their transcriptional regulation may differ, the recurrent emergence of *flam*-like loci across the *Drosophila* genus and the wider presence of unistrand clusters within in the animal kingdom hints at convergent evolution, where this mechanism is best equipped to antagonise TE mobilisation. Together, our study opens the door to understanding the co-evolution between virus-like *Gypsy*-family transposons and the host defence mechanisms that silence them. Further characterisation of these novel piRNA clusters as well as the piRNA pathway machinery in these species will allow us and others to test several long-standing hypotheses regarding piRNA cluster emergence, transcriptional regulation, and the licensing of their transcripts for piRNA biogenesis.

METHODS

Genome assemblies and nomenclature

We strived to collect as many high-quality genome assemblies as possible, including multiple ones for the same species when available, to ensure that we maximise the chance to detect novel unistrand clusters and to assess consistency across isolates. In total, we used 193 assemblies from 119 species. All downloads and processing were done by custom scripts (see "Genome_assemblies" at https://github.com/susbo/Drosophila_unistrand_clusters) and are summarised below.

We used two resources of mostly long-read assemblies representing 15 genomes for 15 species [29] and 104 genomes for 101 species [28]. The latter 104 assemblies included a re-assembly of the first 15 assemblies. Although the re-assemblies generally had higher BUSCO scores compared to the original ones [28], we kept both versions for the analysis.

Assemblies for 36 species annotated by the NCBI Eukaryotic Genome Annotation Pipeline (listed on https://www.ncbi.nlm.nih.gov/genome/annotation_euk/all) from

any species within the *Drosophila* genus were downloaded on three separate occasions (2020-10-17, 2021-10-20, and 2022-04-26). Only the most recently annotated genome assembly is listed for each species and as a result, 19 species were represented by a single assembly and 15 species were represented by two different assemblies.

Additionally, we downloaded the droEre1, droSec1, droSim1, droYak2, droAna2, droPer1, dp3, droMoj2, droVir2, and droGri1 assemblies from the UCSC Genome Browser (http://hgdownload.soe.ucsc.edu/goldenPath). GCF_000754195.2 from *D. simulans* and GCF_000005975.2 from *D. yakuba* from NCBI RefSeq and another 12 assemblies used in a recent study of *Drosophila* phylogeny [53] from the NCBI GenBank.

Genome assemblies downloaded from NCBI or the UCSC Genome Browser retained their original identifier. For assemblies downloaded from the 15 or 101 genomes resources, we used 'd15genomes' and 'd101g', respectively. For NCBI genomes, the contig/scaffold identifiers were simplified in UCSC Genome Browser-style based on information in the FASTA description lines. Unplaced contigs/scaffolds were referred to as 'chrUn_nnn' where 'nnn' refers to the numerical part of the 'NW_nnn' identifier, and contigs/scaffolds associated with a chromosome were referred to in 'chrN'_rand_nnn' format. The script used for the replacements is available in the repository above. Species abbreviations and all studied assemblies are listed in **Table S7**.

Synteny analysis for *flam* conservation

Since the clusters themselves are not conserved, we used a synteny analysis (see "Synteny_clusters" at https://github.com/susbo/Drosophila_unistrand_clusters). Briefly, we used the *D. melanogaster* genome as a reference and extracted the 20 unique up- and downstream genes, excluding tRNA, miRNA, snoRNA, asRNA, and sisRNA. For each gene, we extracted the coding sequence (protein-coding genes) or the full transcript (all others). Next, we mapped these sequences onto the genome of interest using blat (v36x6, -minIdentity=25) and filtered the results to keep the best hit (pslCDnaFilter, -minCover=0.2 -globalNearBest=0.0). Finally, we constructed a candidate list with all genomic regions that had at least two gene hits within 1 Mb. These candidate regions were then manually inspected for the presence of a transposon-rich area at the expected syntenic location. Clusters running into assembly breakpoints were labelled as either 5' or 3', depending on whether they were located next to up- or downstream genes.

UCSC Genome Browser shots

We prepared a genome browser assembly hub covering all 193 assemblies, largely following the instructions on http://genomewiki.ucsc.edu/index.php/Assembly_Hubs. For species with NCBI gene predictions available, we used gtfToGenePred to convert the annotations from GTF to genePred format, followed by genePredToBigGenePred to convert it to bigGenePred format. Gene identifiers were replaced by gene symbols, if available, and the resulting file was converted to bigBed. Tracks displaying the best *D. melanogaster*, *D. grimshawi*, and *D. pseudoobscura* gene mappings were prepared for all genomes using blat (v36x6, -minIdentity=90) followed by pslCDnaFilter (-minCover=0.5 -globalNearBest=0.0). The resulting psl file was converted to genePred format using mrnaToGene. Gene identifiers were replaced by symbols and identical names were numbered to allow search. The edited GenePred file was converted to bigBed via bigGenePred. To construct repeat tracks (RepeatMasker, EDTA, or final TE libraries), we followed the instructions on http://genomewiki.ucsc.edu/index.php/RepeatMasker, enabling the display of the tracks, coloured by strand, and grouped by repeat type. De novo identified clusters (proTRAC) were converted from GTF to GenePred, followed by BigGenePred and BigBed. RNA-seq and sRNA-seq tracks were displayed as standard bigWig tracks produced by deepTools. All genome browser shots shown

in this study were made by exporting the assembly hub display as a pdf, followed by manual refinement to enhance readability.

RepeatMasker tracks

Repeat annotations were done using RepeatMasker (v4.1.2, -s -species Drosophilidae -xsmall) with the Dfam v3.5 and RepBase (RepeatMaskerEdition-20181026) databases [54].

Mappability tracks

Mappability tracks for the sRNA-seq were constructed by generating all possible 26-mers from each genome (bedtools, v2.26.0), aligning them back to the genome (bowtie, v1.2.3, -S -n 2 -M 1 --best --strata --nomaqround --chunkmbs 1024 --no-unal), and converting the alignments to bigWig using deepTools bamCoverage (v3.3.2, --binSize 1 --normalizeUsing None --scaleFactor 0.038461). This will construct a per-nucleotide signal between 0 and 1, representing the ability to uniquely map reads to each position.

LTR strand bias and transposon classes in repeat tracks

To retain consistency across all genome browser shots, the percentage of LTR transposons per strand (displayed in **Fig. S1** and others) is always quantified based on the EDTA repeat annotations (see "De novo transposon annotations using EDTA"), whereas cluster content (bar graphs in **Fig. S1** and others) is based on the curated transposon libraries (see "Construction of curated de novo transposon libraries" below).

Synteny visualisation

MCScan tool [55] was used for synteny comparisons and visualisations. The input to MCScan (Python version) was gene sequences in FASTA format and coordinates in BED-like format. This tool calls LAST [56] to perform pairwise synteny search and a single linkage clustering is performed on the LAST output to cluster anchors into synteny blocks. Following this calculation, different visualisations can be produced with MCScan and hence we performed "Macrosynteny" and "Microsynteny" visualisation. The former is a karyotype plot and highlights syntenic regions across species' genomes whereas the latter offers the advantage to investigate local synteny, which focuses on gene-level. Based on the MCScan tool, we created a set of Python scripts that can be used to investigate and visualise gene synteny between *Drosophila* species. The code and examples with instructions are available on GitHub (https://github.com/marianna-trapotsi/MCScan_plot).

Phylogenetic trees

To display our species onto a phylogenetic tree, we used a previously reported IQ-TREE maximum-likelihood analysis for 704 Drosophilidae species [57]. The phylogeny was imported using the R module treeio (v.1.10.0) and species not included in our study were dropped. Visualisation of the tree and metadata was done using ggtree (v2.0.4) and ggnewscale (v0.4.6). Time of divergence estimates were taken from another recent study of *Drosophila* phylogeny [53] based on fossil evidence.

De novo transposon annotations using EDTA

An initial de-novo transposon library was built using EDTA (v1.9.3, --sensitive 1 --anno 1 --evaluate 1) [33]. The EDTA pipeline consists of three steps, detection of LTRs, Helitrons and TIRs. Most genomes were successfully processed, with transposons of all types being detected. However, some runs failed when one of the types were missing and we manually resumed EDTA at the next type for these genomes. Three genome assemblies (Dneo-d101g, Dsal-d101g, and Zind-d101g_BS02) that failed to run with EDTA v1.9.3 did run successfully with v1.9.6, whereas four assemblies (Daca-d101g, Dari-GCF_001654025, Dnav-GCF_001654015, Dwas-d101g) that had problems with v1.9.3 still had to be resumed with v1.9.6 due to not detecting any LTR transposons.

De novo detection of unistrand *flam*-like clusters
To search for *flam*-like clusters, we developed a search strategy based on the known enrichment of LTR transposons arranged in the same orientation in *flam* (see "De-novo_clusters" at https://github.com/susbo/Drosophila_unistrand_clusters). Briefly, repeat annotations from the EDTA were used. The repeats were separated based on strand retaining either only LTR transposons, or all transposons with a predicted class (i.e., not unknown). Overlapping annotations were combined (bedtools merge) and strand-specific transposon coverage was computed (bedtools coverage) across the genome using a 100 kb sliding window with a 5 kb step size (bedtools makewindows, -w 100000 -s 5000). Each genome was manually inspected for regions enriched in LTR transposons, with a strong strand bias, and located outside of centromeric or telomeric regions. This analysis was strongly contingent on assembly quality, but we nevertheless identified 15 clusters that fulfilled the outlined criteria, including several corresponding to *flam* across the *D. melanogaster* group. Six of the initial candidates were found outside of the *D. melanogaster* subgroup. Of these, two species had publicly available sRNA-seq data, and both produced large amounts of piRNAs from one strand only. We therefore concluded that the approach was working. Synteny analysis using these five clusters as starting points (described below) revealed that *D. persimilis* and *D. pseudoobscura* were syntenic.

Fly husbandry
All *Drosophila* species were maintained at room temperature. The origin of each species and their food requirements are indicated in **Table S8**.

Small RNA-seq library preparation
Small RNAs were isolated from 15 species (2-3 replicates each) using the TraPR Small RNA Isolation Kit (Lexogen) following the manufacturer's instructions. sRNA libraries were generated using the Small RNA-Seq Library Prep Kit (Lexogen) with minor modifications. Both primers A3 and A5 as well as the primer RTP were used at 0.5x. Libraries were sequenced 50 nt paired-end on an Illumina NovaSeq 6000 or 75 nt single-end on an Illumina MiSeq sequencing platform generating 29 (+/- 15) million reads per library.

Enrichment for somatic cells
The soma-enrichment sRNA-seq libraries were generated for 12 species (2 replicates each) similar to published protocols [26,58], with modifications. In brief, 75-100 ovary pairs were dissected in ice-cold PBS. Ovaries were dissociated for 18 minutes in 0.25% Trypsin (Sigma-Aldrich) at 25°C, shaking at 800 rpm. Dissociated tissue was pushed through a 40 µm nylon mesh (Greiner Bio-one) washed with equal volume Schneider 2 medium (Thermo Fisher Scientific) and then pelleted. Cell count was quantified using the Luna-FL (Logos Biosystems). Pelleted cells were directly used as input for sRNA isolation using the TraPR Small RNA Isolation Kit (Lexogen).

Publicly available sRNA-seq data
We searched the Sequencing Read Archive (SRA) and Gene Expression Omnibus (GEO) for any sRNA-seq data from *Drosophila* species other than *D. melanogaster*. After excluding two SOLiD sequencing samples, we found 67 samples from 12 species [26,59–63], representing embryo (n=16), female body (n=11), female germline (n=2), female soma (n=2), head (n=18), male body (n=9), follicle-cell enriched ovary (n=3), and testis (n=6).

Processing of sRNA-seq data
All sRNA-seq data were processed using the same analysis pipeline. Trim Galore! (v0.6.4, --stringency 30 -e 0.1 -a TGCTTGGACTACATATGGTTGAGGGTTGTA --length 18 -q 0) was first run to remove an abundant rRNA sequence, followed by a second run (--stringency 5 -e 0.1 --length 18 --max_length 35

-q 0) to remove adapter sequences (specified using '-a'), and any flanking random nucleotides ('--clip_R1' and/or '--three_prime_clip_R1' with appropriate arguments). All samples and their adapter sequences are listed in **Table S9**.

The processed reads were mapped to a miRNA hairpin database (miRBase release 22.1) [64] using bowtie (v1.2.3, -S -n 2 -M 1 -p 20 --best --strata --nomaqround --chunkmbs 1024) with '--un' and '--max' to extract unmapped reads. Reads not mapping to miRNAs were aligned to the respective reference genomes using bowtie (-S -n 2 -M 1 -p 20 --best --strata --nomaqround --chunkmbs 1024). Multi-mapping reads were extracted into a separate BAM file using awk (MQ<10). Alignment metrics are available in **Table S9**.

The BAM files were converted to bigWig using bamCoverage from deepTools [65] (v3.3.2, --binSize 1 --ignoreForNormalization chrM --normalizeUsing CPM --exactScaling --skipNonCoveredRegions --minFragmentLength 23 --maxFragmentLength 30) and additionally '--filterRNAstrand' to separate the two strands, '--scaleFactor' to scale counts per million to reflect all mapped reads, and optionally '--minMappingQuality 50' when extracting uniquely mapped reads.

RNA-seq library preparation
Ovaries from 10-20 flies were dissected in ice-cold PBS and total RNA was extracted using TRIzol (Thermo Fisher Scientific), following the manufacturer's instructions. Ribosomal RNA was depleted using RiboPOOL (siTOOLs Biotech) following the manufacturer's protocol. Additionally, we re-sequenced (paired-end) four published *D. melanogaster* samples treated with RiboZero as previously described [66]. RNA-seq libraries were produced using NEBNext Ultra Directional Library Prep Kit for Illumina, following the manufacturer's instructions for rRNA depleted RNA. Library size distribution was analysed on a TapeStation instrument (Agilent Technologies) using a High Sensitivity D1000 ScreenTape. Libraries were pooled in equal molar ratio, quantified with KAPA Library Quantification Kit for Illumina (Kapa Biosystems) and sequenced paired-end 50 nt on an Illumina NovaSeq 6000.

Publicly available RNA-seq data
We downloaded modENCODE RNA-seq data from *Drosophila* species other than *D. melanogaster* and additional *D. innubila* samples [67,68]. We included 58 samples from 8 species, representing embryo (n=16), female body (n=18), female head (n=1), larvae (n=2), male body (n=18), male head (n=1), and pupae (n=2).

Processing of RNA-seq data
Trim Galore! (v0.6.4, --stringency 6 -e 0.1) was used to remove adapters and low-quality bases. Alignment was done using HiSeq2 [69] (v2.2.0, -max-seeds 100 -q -k 1), keeping at most one alignment for each read. Multi-mapping reads were extracted into a separate BAM file using awk (MQ<10). The BAM files were converted to bigWig using bamCoverage from deepTools [65] (v3.3.2, --binSize 1 --ignoreForNormalization chrM --normalizeUsing CPM --exactScaling --skipNonCoveredRegions) and additionally '--filterRNAstrand' to separate the two strands, '--scaleFactor' to scale the counts per million to all mapped reads, and optionally '--minMappingQuality 50' when extracting uniquely mapped reads. Alignment metrics are available in **Table S10**.

Ping-pong and phasing analyses
Ping-pong and phasing analyses were performed for reads of length 24-28 nt mapping to each cluster region. The deepTools module bamCoverage was used to extract the number of 3' and 5' ends mapping to each position and strand. Ping-pong and phasing signature was calculated following the strategy in [70]. In short, we calculated the ping-pong signature using a 5' end overlap score for overlap x nt as

$$s_x = \sum_{i \,\in\, all\ positions} n_i m_{i+x}$$

where n_i is the number of 5' ends mapping at the plus strand position i and m_{i+x} is the number of 5' ends mapping at the minus strand position $i+x$. The fraction of overlapping reads involved in ping-pong was calculated as $s_{10}/(s_1+...+s_{20})$. A z10 score was defined as $(s_{10}-mean(s_1,...,s_9,s_{11},...,s_{20}))/stdev(s_1,...,s_9,s_{11},...,s_{20})$.

For the phasing signature, we calculated a 3' to 5' end score for distance y as

$$h_y = \sum_{i \,\in\, all\ positions} \min(n_i, m_{i+y})$$

where n_i is the number of 3' ends mapping at position i and m_{i+y} is the number of 5' ends mapping at position $i+y$ at the same strand. The fraction of closely mapped reads with phasing signature was calculated as $h_1/(h_1+...+h_{20})$. A z1 score was calculated as $(h_1-mean(h_2,...,h_{20}))/stdev(h_2,...,h_{20})$. Phasing calculations were done for the plus and minus strand separately.

Detection of regions syntenic to *flam*-like clusters
Synteny analysis for *flam*-like clusters was performed using the same strategy as for *flam*, except that Augustus gene predictions (v3.3.2, --species=fly --UTR=off --singlestrand=true) were used instead of FlyBase annotations. The MAKER-masked genome from the EDTA output was used as genome input to Augustus. Full transcript and coding sequences were extracted from the annotations. Sequences with strong hits to the raw transposon libraries were excluded (blat, -q=dna -t=dna -minIdentity=25; pslCDnaFilter, -minCover=0.2 -globalNearBest=0) and gene predictions shorter than 200 nt were excluded. The blat identity threshold was reduced to 20.

Additionally, within the *obscura* group (*flamlike5*), the closest flanking genes displayed good conservation and we used these to search for syntenic regions using our UCSC Genome Browser session.

Prediction of major piRNA clusters
Cluster predictions were performed using proTRAC (v2.4.4, -pdens 0.01 -swincr 100 -swsize 1000 -clsize 5000 -1Tor10A 0.75 -clstrand 0.5 -pimin 23 -pimax 30 -psize 0.75 -distr 1-99 -nomotif -format SAM) [35] using all available sRNA-seq libraries except head, female_soma, male_body and OSC, with '-repeatmasker' set to RepeatMasker annotations generated by EDTA, and with '-geneset' set to NCBI gene predictions, if available.

Clusters within 40 kb from each other were combined for the analyses. Genes flanking the major *D. melanogaster* clusters were mapped onto each genome using BLAT (v36x6, -minIdentity=25), filtered to retain only the best hit (pslCDnaFilter, -minCover=0.2 -globalNearBest=0.0) and the predicted clusters were subsequently annotated by how many of these genes that were within 1Mb.

Cluster predictions used in the soma-enrichment analysis were performed using the same strategy but restricted to libraries generated for this study and using either only soma-enriched or only total sRNA-seq libraries (2-3 replicates per species and library type). Clusters identified using either somatic or total libraries were concatenated and any clusters within 40 kb from each other were merged. Clusters of size <35 kb were discarded to enable analysis of strand biases across major clusters. Total piRNA coverage per cluster was normalised to counts per million and calculated for soma-enriched and total libraries separately. Somatic clusters were defined as clusters with at least 2-fold soma enrichment over total libraries, and germline clusters were defined as being higher expressed in the total ovary libraries.

Construction of curated *de novo* transposon libraries
In addition to the consensus sequences obtained from EDTA, we also used the RepeatModeler (v2.0.1) output within the EDTA folders to improve detection of LINE elements. We reasoned that we could not provide a list of known LINEs to EDTA, since that would mainly reflect *melanogaster* transposons and would bias the comparisons between the *melanogaster* subgroup and other species.

EDTA and RepeatModeler consensus sequences were combined and further processed using a custom pipeline (see "Transposon_libraries" at https://github.com/susbo/Drosophila_unistrand_clusters). Briefly, raw sequences classified as rRNA, snRNA, tRNA, ARTEFACT, or Simple_repeat were removed, and remaining sequences were clustered using cd-hit-est (v4.8.1, -G 0 -g 1 -c 0.90 -aS 0.90 -n 8 -d 0 -b 500) to combine any sequences with ≥90% identity across ≥90% of the length. Custom scripts were used to select one representative sequence from each cluster, maximising both the number of high-quality genomic hits (blastn, filtered to cover at least 50% of the query sequence) and the length of the sequence. Sequences with fewer than 2 high-quality genomic hits were removed from the transposon libraries.

To prioritise sequences and to detect known transposon domains, we mapped all consensus sequences to *env*, *gag* and *pol* ORFs from RepeatPeps.lib in RepeatMasker (v4.1.2) using blastx (v2.10.0, -max_target_seqs 100 -evalue 1e-3). Sequences covering at least 50% of the full peptide domain were considered true hits.

To detect previously described subfamilies, we used the *Drosophila* transposon canonical sequences database (v10.2, https://github.com/bergmanlab/transposons), including 180 consensus sequences from seven species. We used full_blast (https://github.com/rimjhimroy/Transposon80-80-80, -p megablast -i 95 -qc 98 -l 80 -t 10) to identify hits with ≥95% identity, hit length ≥80 nt, and ≥98% query coverage (i.e., "95/80/98 rule" [71,72]) to the canonical sequences, which were considered to belong to the same subfamily. This was repeated using 90/80/90 and 80/80/80 thresholds to detect more distant similarities.

Transposon class and family were predicted using the RepeatClassifier script in RepeatModeler (v2.0.2a), configured to use RepeatMasker (v4.1.2) with the Dfam (v3.5) and RepBase (RepeatMaskerEdition-20181026) databases [54].

The curated and annotated transposon consensus sequences have been made available (https://github.com/susbo/Drosophila_TE_libraries).

Mapping of sRNA-seq and RNA-seq across species
To evaluate whether different *Drosophila* species express similar piRNAs (**Fig. S21**), we extracted 100,000 filtered reads from each sRNA-seq library and aligned them across all 193 assemblies using bowtie (v1.2.3, -S -n 2 -M 1 -p 10 --best --strata --nomaqround --chunkmbs 1024 --no-unal). Alignment rate for multiple assemblies of the same species were averaged. Most libraries displayed near-zero alignment rate across all species except very closely related ones. A small number of libraries showed elevated baseline alignment, likely due to the presence of other small RNAs (rRNA, tRNA) with high conservation. To remove this effect, we subtracted the median alignment rate from each library, which roughly corresponds to removing reads mapping to all species. A similar strategy was used to map RNA-seq libraries across species (**Fig. S22**) using 200,000 reads aligned with hisat2 (v2.2.0, --max-seeds 100 -q -k 1 -p 10 --no-unal --new-summary –summary-file). Similarly, the median alignment rate was subtracted from each library to reduce false hits driven by abundant non-coding RNAs.

Cluster content analyses
Cluster content analyses in **Fig. 5** use the curated transposon libraries either in the form of genomic coverage (**Fig. 5c-d**) or the number of transposon subfamilies (**Fig. 5e**). To obtain these

estimates we ran RepeatMasker (v4.2.1, -s -xsmall) with the curated transposon libraries (-lib) followed by the buildSummary.pl script to quantify both coverage and copy number within the cluster.

To determine whether clusters were more likely to have captured *Gypsy*-family LTR transposons compared with other LTR transposons we defined a Gypsy enrichment ratio as

$$\text{Gypsy enrichment} = \frac{\text{P}(x \in \text{Gypsy} \mid x \in \text{LTR}, x \in \text{Captured})}{\text{P}(x \in \text{Gypsy} \mid x \in \text{LTR}, x \in \text{Not captured})}$$

And similarly, we defined an *env* enrichment ratio as

$$\text{env enrichment} = \frac{\text{P}(x \in \text{env} \mid x \in \text{Gypsy}, x \in \text{Captured})}{\text{P}(x \in \text{env} \mid x \in \text{Gypsy}, x \in \text{Not captured})}$$

Where *x* is a transposon subfamily, Gypsy is the set of all *Gypsy*-family transposon subfamilies, LTR is the set of all LTR transposons, Captured is the set of all transposon subfamilies found inside the cluster region and Not captured is the set of all other transposon subfamilies.

Conservation of piRNA pathway genes

To avoid false hits to conserved protein domains and to increase sensitivity compared with sequence-based searches, we employed a synteny-based search strategy (see "Synteny_biogenesis_genes" at https://github.com/susbo/Drosophila_unistrand_clusters). Briefly, we used the *D. melanogaster* genome as a reference and extracted the 20 closest genes up- and downstream genes, excluding tRNA, miRNA, snoRNA, asRNA, and sisRNA. For each gene, we extracted the coding sequence (protein-coding genes) or the full transcript (all others). Next, we mapped these sequences onto the genome of interest using blat (v36x6, -maxIntron=500000 -minMatch=2 -minScore=30 -oneOf f=1 -minIdentity=10) and filtered the results to keep the best hit (pslCDnaFilter, -minCover=0.1 -globalNearBest=0.0). Finally,

we constructed a hit list with all genomic regions that had at least two hits within 1 Mb from another. Typically, this resulted in a single hit, which was manually inspected for the presence of the gene of interest.

Data availability

Raw data and genome browser tracks from high-throughput sequencing experiments will be available on GEO.

ACKNOWLEDGEMENTS

We thank Hannon group members for fruitful discussions, Holly Goodrick for help with library preparation and Emma Kneuss for providing early access to unpublished sRNA-seq libraries. We thank the Scientific Computing core facility at CRUK Cambridge Institute for HPC resources. GJH is a Royal Society Wolfson Research Professor (RSRP\R\200001). This research was funded in whole, or in part, by Cancer Research UK (A21143) and the Wellcome Trust (110161/Z/15/Z). The following species were kindly provided by the indicated laboratories: *D. mojavensis, D. persimilis* and *D. pseudoobscura* from Ben Longdon, University of Exeter, UK; *D. ficusphila* from Geoffrey Findlay, College of the Holy cross, Worcester, MA, USA; *D. simulans* from David Stern, Janelia Research Campus, VA, USA; *D. erecta, D. yakuba, D. ananassae* and *D. virilis* from Simon Collier at the Department of Genetics, Fly Facility, University of Cambridge, UK.

AUTHOR CONTRIBUTIONS

JvL, GJH, SB and BCN conceived the study. SB, JvL and BCN designed the experiments and interpreted the results. JvL performed all wet-lab work and preliminary computational analysis. SB performed all computational work, except the MCScan analysis that was performed by MAT. BCN, GJH and SB supervised the work. SB and JvL generated the figures. JvL, SB and BCN wrote the manuscript with input from GJH. All authors read and approved the final version.

COMPETING INTERESTS

The authors declare no competing interests.

REFERENCES

1. Malone, C. D. & Hannon, G. J. Small RNAs as guardians of the genome. *Cell* **136**, 656–668 (2009).
2. Mével-Ninio, M., Pelisson, A., Kinder, J., Campos, A. R. & Bucheton, A. The flamenco Locus Controls the gypsy and ZAM Retroviruses and Is Required for Drosophila Oogenesis. *Genetics* **175**, 1615 (2007).
3. Teixeira, F. K. *et al.* piRNA-mediated regulation of transposon alternative splicing in the soma and germ line. *Nature* **552**, 268–272 (2017).
4. Seczynska, M., Bloor, S., Cuesta, S. M. & Lehner, P. J. Genome surveillance by HUSH-mediated silencing of intronless mobile elements. *Nature 2021 601:7893* **601**, 440–445 (2021).
5. Robbez-Masson, L. *et al.* The hush complex cooperates with trim28 to repress young retrotransposons and new genes. *Genome Res* **28**, 836–845 (2018).
6. Ecco, G., Imbeault, M. & Trono, D. KRAB zinc finger proteins. *Development* **144**, 2719–2729 (2017).
7. Yang, P., Wang, Y. & Macfarlan, T. S. The Role of KRAB-ZFPs in Transposable Element Repression and Mammalian Evolution. *Trends Genet* **33**, 871–881 (2017).
8. Özata, D. M., Gainetdinov, I., Zoch, A., O'Carroll, D. & Zamore, P. D. PIWI-interacting RNAs: small RNAs with big functions. *Nat Rev Genet* **20**, 89–108 (2019).
9. Czech, B. *et al.* PiRNA-guided genome defense: From biogenesis to silencing. *Annual Review of Genetics* Preprint at https://doi.org/10.1146/annurev-genet-120417-031441 (2018).
10. Brennecke, J. *et al.* Discrete Small RNA-Generating Loci as Master Regulators of Transposon Activity in Drosophila. *Cell* **128**, 1089–1103 (2007).
11. Özata, D. M. *et al.* Evolutionarily conserved pachytene piRNA loci are highly divergent among modern humans. *Nat Ecol Evol* **4**, 156–168 (2020).
12. Li, X. Z. *et al.* An Ancient Transcription Factor Initiates the Burst of piRNA Production during Early Meiosis in Mouse Testes. *Mol Cell* **50**, 67–81 (2013).
13. Aguiar, E. R. G. R. *et al.* A single unidirectional piRNA cluster similar to the flamenco locus is the major source of EVE-derived transcription and small RNAs in Aedes aegypti mosquitoes. *RNA* **26**, 581–594 (2020).
14. Houwing, S. *et al.* A role for Piwi and piRNAs in germ cell maintenance and transposon silencing in Zebrafish. *Cell* **129**, 69–82 (2007).
15. Parhad, S. S., Tu, S., Weng, Z. & Theurkauf, W. E. Adaptive Evolution Leads to Cross-Species Incompatibility in the piRNA Transposon Silencing Machinery. *Dev Cell* **43**, 60-70.e5 (2017).

16. Mohn, F., Sienski, G., Handler, D. & Brennecke, J. The rhino-deadlock-cutoff complex licenses noncanonical transcription of dual-strand piRNA clusters in Drosophila. *Cell* **157**, 1364–1379 (2014).

17. Wierzbicki, F., Kofler, R. & Signor, S. Evolutionary dynamics of piRNA clusters in Drosophila. *Mol Ecol* (2021) doi:10.1111/MEC.16311.

18. Goriaux, C., Desset, S., Renaud, Y., Vaury, C. & Brasset, E. Transcriptional properties and splicing of the flamenco piRNA cluster. *EMBO Rep* **15**, 411–418 (2014).

19. Dennis, C., Brasset, E., Sarkar, A. & Vaury, C. Export of piRNA precursors by EJC triggers assembly of cytoplasmic Yb-body in Drosophila. *Nat Commun* **7**, (2016).

20. Prud'homme, N., Gans, M., Masson, M., Terzian, C. & Bucheton, A. Flamenco, a gene controlling the gypsy retrovirus of Drosophila melanogaster. *Genetics* **139**, 697–711 (1995).

21. Pelisson, A. *et al.* Gypsy transposition correlates with the production of a retroviral envelope-like protein under the tissue-specific control of the Drosophila flamenco gene. *EMBO J* **13**, 4401–4411 (1994).

22. Vaury, C., Bucheton, A. & Pelisson, A. The beta heterochromatic sequences flanking the I elements are themselves defective transposable elements. *Chromosoma* **98**, 215–224 (1989).

23. Pelisson, A., Mejlumian, L., Robert, V., Terzian, C. & Bucheton, A. Drosophila germline invasion by the endogenous retrovirus gypsy: Involvement of the viral env gene. *Insect Biochem Mol Biol* **32**, 1249–1256 (2002).

24. Song, S. U., Kurkulos, M., Boeke, J. D. & Corces, V. G. Infection of the germ line by retroviral particles produced in the follicle cells: a possible mechanism for the mobilization of the gypsy retroelement of Drosophila. *Development* **124**, 2789–2798 (1997).

25. Leblanc, P. *et al.* Life cycle of an endogenous retrovirus, ZAM, in Drosophila melanogaster. *J Virol* **74**, 10658–10669 (2000).

26. Chirn, G. W. *et al.* Conserved piRNA Expression from a Distinct Set of piRNA Cluster Loci in Eutherian Mammals. *PLoS Genet* **11**, 1–26 (2015).

27. Malone, C. D. *et al.* Specialized piRNA pathways act in germline and somatic tissues of the Drosophila ovary. *Cell* **137**, 522–535 (2009).

28. Kim, B. Y. *et al.* Highly contiguous assemblies of 101 drosophilid genomes. *Elife* **10**, (2021).

29. Miller, D. E., Staber, C., Zeitlinger, J. & Hawley, R. S. Highly Contiguous Genome Assemblies of 15 Drosophila Species Generated Using Nanopore Sequencing. *G3 (Bethesda)* **8**, 3131–3141 (2018).

30. Goriaux, C., Théron, E., Brasset, E. & Vaury, C. History of the discovery of a master locus producing piRNAs: the flamenco/COM locus in Drosophila melanogaster. *Front Genet* **5**, (2014).

31. Andersen, P. R., Tirian, L., Vunjak, M. & Brennecke, J. A heterochromatin-dependent transcription machinery drives piRNA expression. *Nature* **549**, 54–59 (2017).

32. Munafò, M. *et al.* Channel nuclear pore complex subunits are required for transposon silencing in Drosophila. *Elife* **10**, (2021).

33. Ou, S. *et al.* Benchmarking transposable element annotation methods for creation of a streamlined, comprehensive pipeline. *Genome Biol* **20**, (2019).

34. Gunawardane, L. S. *et al.* A slicer-mediated mechanism for repeat-associated siRNA 5′ end formation in Drosophila. *Science (1979)* **315**, 1587–1590 (2007).

35. Rosenkranz, D. & Zischler, H. proTRAC--a software for probabilistic piRNA cluster detection, visualization and analysis. *BMC Bioinformatics* **13**, (2012).

36. Malik, H. S., Henikoff, S. & Eickbush, T. H. Poised for contagion: evolutionary origins of the infectious abilities of invertebrate retroviruses. *Genome Res* **10**, 1307–1318 (2000).

37. Terzian, C., Pélisson, A. & Bucheton, A. Evolution and phylogeny of insect endogenous retroviruses. *BMC Evol Biol* **1**, (2001).

38. Llorens, J. v. *et al.* Gypsy endogenous retrovirus maintains potential infectivity in several species of Drosophilids. *BMC Evol Biol* **8**, (2008).

39. Duc, C. *et al.* Trapping a somatic endogenous retrovirus into a germline piRNA cluster immunizes the germline against further invasion. *Genome Biol* **20**, (2019).

40. Yoth, M., Gueguen, N., Jensen, S. & Brasset, E. Germline piRNAs counteract endogenous retrovirus invasion from somatic cells. *bioRxiv* 2022.08.29.505639 (2022) doi:10.1101/2022.08.29.505639.

41. Zanni, V. *et al.* Distribution, evolution, and diversity of retrotransposons at the flamenco locus reflect the regulatory properties of piRNA clusters. *Proc Natl Acad Sci U S A* **110**, 19842–19847 (2013).

42. Hirakata, S., Ishizu, H., Fujita, A., Tomoe, Y. & Siomi, M. C. Requirements for multivalent Yb body assembly in transposon silencing in Drosophila. *EMBO Rep* **20**, (2019).

43. Qi, H. *et al.* The Yb body, a major site for Piwi-associated RNA biogenesis and a gateway for Piwi expression and transport to the nucleus in somatic cells. *J Biol Chem* **286**, 3789–3797 (2011).

44. Chary, S. & Hayashi, R. Mechanistic divergence of piRNA biogenesis in Drosophila. *bioRxiv* 2022.11.14.516378 (2022) doi:10.1101/2022.11.14.516378.

45. Kofler, R. Dynamics of Transposable Element Invasions with piRNA Clusters. *Mol Biol Evol* **36**, 1457–1472 (2019).

46. Gebert, D. *et al.* Large Drosophila germline piRNA clusters are evolutionarily labile and dispensable for transposon regulation. *Mol Cell* **81**, 3965-3978.e5 (2021).

47. Klattenhoff, C. *et al.* The Drosophila HP1 homolog Rhino is required for transposon silencing and piRNA production by dual-strand clusters. *Cell* **138**, 1137–1149 (2009).

48. Pane, A., Jiang, P., Zhao, D. Y., Singh, M. & Schüpbach, T. The Cutoff protein regulates piRNA cluster expression and piRNA production in the Drosophila germline. *EMBO J* **30**, 4601–4615 (2011).

49. Czech, B., Preall, J. B., McGinn, J. & Hannon, G. J. A transcriptome-wide RNAi screen in the Drosophila ovary reveals factors of the germline piRNA pathway. *Mol Cell* **50**, 749–761 (2013).

50. Munafò, M. *et al.* Daedalus and gasz recruit armitage to mitochondria, bringing piRNA precursors to the biogenesis machinery. *Genes Dev* **33**, 844–856 (2019).

51. ElMaghraby, M. F. *et al.* A Heterochromatin-Specific RNA Export Pathway Facilitates piRNA Production. *Cell* **178**, 964-979.e20 (2019).

52. Signor, S. *et al.* Rapid evolutionary diversification of the flamenco locus across simulans clade Drosophila species. *bioRxiv* 2022.09.29.510127 (2022) doi:10.1101/2022.09.29.510127.
53. Suvorov, A. *et al.* Widespread introgression across a phylogeny of 155 Drosophila genomes. *Curr Biol* **32**, 111-123.e5 (2022).
54. Bao, W., Kojima, K. K. & Kohany, O. Repbase Update, a database of repetitive elements in eukaryotic genomes. *Mob DNA* **6**, (2015).
55. Tang, H. *et al.* Synteny and collinearity in plant genomes. *Science* **320**, 486–488 (2008).
56. Kiełbasa, S. M., Wan, R., Sato, K., Horton, P. & Frith, M. C. Adaptive seeds tame genomic sequence comparison. *Genome Res* **21**, 487–493 (2011).
57. Finet, C. *et al.* DrosoPhyla: Resources for Drosophilid Phylogeny and Systematics. *Genome Biol Evol* **13**, (2021).
58. Akkouche, A. *et al.* Maternally deposited germline piRNAs silence the tirant retrotransposon in somatic cells. *EMBO Rep* **14**, 458–464 (2013).
59. Vedanayagam, J., Lin, C. J. & Lai, E. C. Rapid evolutionary dynamics of an expanding family of meiotic drive factors and their hpRNA suppressors. *Nat Ecol Evol* **5**, 1613–1623 (2021).
60. Mohammed, J. *et al.* Deep experimental profiling of microRNA diversity, deployment, and evolution across the Drosophila genus. *Genome Res* **28**, 52–65 (2018).
61. Lewis, S. H. *et al.* Pan-arthropod analysis reveals somatic piRNAs as an ancestral defence against transposable elements. *Nat Ecol Evol* **2**, 174–181 (2018).
62. Zhao, Y. *et al.* Run or Die in the Evolution of New MicroRNAs-Testing the Red Queen Hypothesis on De Novo New Genes. *Mol Biol Evol* **38**, 1544–1553 (2021).
63. Kotov, A. A. *et al.* piRNA silencing contributes to interspecies hybrid sterility and reproductive isolation in Drosophila melanogaster. *Nucleic Acids Res* **47**, 4255–4271 (2019).
64. Kozomara, A., Birgaoanu, M. & Griffiths-Jones, S. miRBase: from microRNA sequences to function. *Nucleic Acids Res* **47**, D155–D162 (2019).
65. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res* **44**, W160–W165 (2016).
66. Kneuss, E. *et al.* Specialization of the Drosophila nuclear export family protein Nxf3 for piRNA precursor export. *Genes Dev* **33**, 1208–1220 (2019).
67. Chen, Z. X. *et al.* Comparative validation of the D. melanogaster modENCODE transcriptome annotation. *Genome Res* **24**, 1209–1223 (2014).
68. Hill, T., Koseva, B. S. & Unckless, R. L. The Genome of Drosophila innubila Reveals Lineage-Specific Patterns of Selection in Immune Genes. *Mol Biol Evol* **36**, 1405–1417 (2019).
69. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915 (2019).
70. Han, B. W., Wang, W., Li, C., Weng, Z. & Zamore, P. D. PiRNA-guided transposon cleavage initiates Zucchini-dependent, phased piRNA production. *Science (1979)* **348**, 817–821 (2015).
71. Flutre, T., Duprat, E., Feuillet, C. & Quesneville, H. Considering transposable element diversification in de novo annotation approaches. *PLoS One* **6**, (2011).
72. Wicker, T. *et al.* A unified classification system for eukaryotic transposable elements. *Nat Rev Genet* **8**, 973–982 (2007).