1

Authors: Ori Erlichman[1]**, Shahar Weiss[1], Maria Abu-Arkia[1], Moria Ankary Khaner[1],
Yoram Soroka[1], Weronika Jasinska[2], Leah Rosental[2], Yariv Brotman[2] and Tamar Avin-
Wittenberg[1]*

Author Affiliations

[1] Department of Plant and Environmental Sciences, Alexander Silberman Institute of Life
Sciences, The Hebrew University of Jerusalem, Givat Ram, 9190401, Jerusalem, Israel

[2] Department of Life Sciences, Ben-Gurion University of the Negev, 8410501, Beer-Sheva,
Israel

* Corresponding author: tamar.avin-wittenberg@mail.huji.ac.il

** Current affiliation: Faculty of Biology, Technion-Israel Institute of Technology,
3200003, Haifa, Israel

Title: Autophagy in maternal tissues contributes to *Arabidopsis thaliana* seed
development.

Keywords (5-8, in alphabetical order): *Arabidopsis thaliana,* Autophagy, Maternal
tissue, Seed, Storage compounds.

Word counts for the main sections (Total word count – 5318, Introduction - 756,
Materials and Methods - 1807, Results - 1674, Discussion - 1081)

Number of figures – 7, all in color, **supporting information –** 6 supplemental figures, 2
supplemental tables.

Summary

Seeds are a significant source of food and feed, containing storage compounds that provide nutrients for germination and early seedling growth. Degradation events in both the seed and the mother plant accompany seed development. Autophagy is a conserved degradation mechanism for the breakdown of cellular constituents in the lytic organelle. It participates in various aspects of plant physiology, specifically nutrient availability and yield determination. Autophagy affects nutrient remobilization from mother plants to seeds and functions in the embryo. Though previous studies suggest autophagy is essential for seed development, they were conducted on autophagy-knockout (*atg* mutant) plants, making it impossible to distinguish between the contribution of autophagy in the mother plant and in the embryo to seed development.

Here, we asked how autophagy in the maternal tissue affects seed development. To that end, we performed reciprocal crosses between WT and *atg* mutant *Arabidopsis thaliana* plants. Etiolated F1 plants from maternal *atg* mutants displayed reduced growth, possibly stemming from lower storage protein amounts. Surprisingly, F1 seeds of maternal *atg* mutants exhibited faster germination and altered seed-coat structure. Our results indicate that the observed phenotypes of *atg* mutant seeds are compound phenotypes influenced by autophagy in the mother plant and the developing embryo.

Introduction

Seeds are a major source of food and feed, providing approximately 70% of the world's human caloric intake(Sreenivasulu & Wobus, 2013). The nutritional value of seeds is attributed to storage compounds accumulating in the seed during development (Cernac & Benning, 2004). These include starch, storage lipids (mostly triacylglycerols - TAGs), and storage proteins, the relative amount of which varies between plant species. Storage compounds are degraded during germination and early seedling development to facilitate germination and seedling growth until photosynthesis establishment (Baud *et al.*, 2008). In annual plants, seed development is accompanied by mother plant senescence, resulting in nutrient remobilization to the developing seeds, used to produce storage compounds (Distelfeld *et al.*, 2014). Another vital seed trait is longevity, defined as seed viability after seed dry storage (Nguyen *et al.*, 2012). Seed longevity can directly affect germination and, as a result, crop productivity and food security (Zhou *et al.*, 2019). With the growing world population(Ort *et al.*, 2015), specifically in light of current climate change(Lobell *et al.*, 2011), an increased understanding of nutrient remobilization to seeds and seed longevity is vital to ensure food security.

Macroautophagy (hereafter termed "Autophagy") is a conserved eukaryotic mechanism for the degradation of cytoplasmic constituents in the lytic organelle (vacuoles in yeast and plants and lysosomes in animals)(Bassham *et al.*, 2006; Marshall & Vierstra, 2018). The targets of autophagy are diverse and include long-lived proteins, protein complexes, and whole organelles(Reumann *et al.*, 2010). The autophagy machinery relies on highly-conserved AuTophaGy-related (*ATG*) genes(Bassham, 2009; Avin-Wittenberg *et al.*, 2012; Liu & Bassham, 2012; Ryter *et al.*, 2013), which have been characterized in many plant species (Chung *et al.*, 2009; Kurusu *et al.*, 2014; Zhou *et al.*, 2014). In recent years, autophagy was shown to be involved in many aspects of plant life, from plant development to biotic and abiotic stress response(Avin-Wittenberg *et al.*, 2018).

Autophagy-deficient plants (*atg* mutants) are hypersensitive to carbon and nitrogen starvation and display early senescence and reduced yield(Doelling *et al.*, 2002; Li *et al.*, 2015; Barros *et al.*, 2017), as well as slightly slower germination rates and impaired seedling establishment(Yoshimoto *et al.*, 2014; Avin-Wittenberg *et al.*, 2015). Autophagy was shown to promote nutrient remobilization in plants, functioning in the supply of lipids

and primary metabolites (Masclaux-Daubresse *et al.*, 2014; Avin-Wittenberg *et al.*, 2015; Wada *et al.*, 2015; Barros *et al.*, 2017; Hirota *et al.*, 2018; McLoughlin *et al.*, 2018; McLoughlin *et al.*, 2020; Barros *et al.*, 2021). Moreover, overexpression of *ATG* genes in *Arabidopsis thaliana* (Arabidopsis) was reported to increase seed yield and fatty acid content (Minina *et al.*, 2018).

Autophagy was shown to play a dual role in seed development, affecting both the mother plant and the developing embryo. Studies in Arabidopsis and maize (*Zea mays)* demonstrated that impaired autophagy causes altered nitrogen mobilization from the mother plant to the seed, resulting in changes in seed C/N ratio(Guiboileau *et al.*, 2012; Li *et al.*, 2015). Alternatively, microscopic evidence from wheat suggested the possible involvement of autophagy in the transport of storage proteins to protein storage vacuoles within the developing seeds (Levanony *et al.*, 1992). Furthermore, an additional publication demonstrated the presence of an active autophagy mechanism within developing seeds as well as altered storage protein content and processing in Arabidopsis *atg* mutants (Di Berardino *et al.*, 2018).

The studies described above suggest an essential role for autophagy in seed development. However, as the analysis was performed on knockout mutants, it is impossible to distinguish between the impact of autophagy in the mother plant and in the seed itself on seed development. Moreover, the seed is comprised of several tissues with differing parental contributions. The seed coat is of maternal origin, while the embryo and endosperm are zygotic tissues, and the genomic contribution of the triploid single-layer endosperm is unevenly divided between the female parent (2n) and the male parent (1n) (Bentsink & Koornneef, 2008). In this work, we asked how autophagy activity in the maternal tissue affects seed development. To that end, we performed reciprocal crosses between Arabidopsis wild type (WT) and *atg* mutants to receive *ATG*-heterozygous F1 seeds with different maternal tissues. We discovered F1 seeds differentially accumulated storage compounds according to the genotype of the mother plant. Surprisingly, we observed that F1 seeds from maternal *atg* mutant origin had a faster germination rate, in contrast to homozygous *atg* mutants. This possibly stems from differences in seed coat structure, which also influences seed longevity. Our work suggests that *atg* mutant seeds

display a compound phenotype, merging the impact of autophagy in the mother plant and

the embryo.

Materials and Methods

<u>Plant material and growth conditions</u>

Arabidopsis thaliana ecotype Columbia (Col-0) was used in this study. The lines used were

atg5-1 (SAIL_129B079)(Yoshimoto *et al.*, 2009), *atg7-2* (GK-655B06) (Hofius *et al.*,

2009), NahG over-expression (NahG) (Yoshimoto *et al.*, 2009) and atg5.NahG (Yoshimoto

et al., 2009). Seeds were surface sterilized with Cl_2 for 2h and sown on Nitsch medium

(Duchefa) plates pH 5.8, with 1% sucrose. The seeds were imbibed at 4°C for 72h in the

dark. Plants were grown in a growth chamber (Fitoclima S600PLH, Aralab) at 22°C and

50% humidity in constant light (125 µmol photons m^{-2} s^{-1}) for 2 weeks. They were then

transferred to pots containing Perlite: Vermiculite: Garden mix (1:1:2 v/v) and grown at

22°C under long-day conditions (8h dark / 16h light) with 150 µmol photons m^{-2} s^{-1}.

<u>Plant crosses and genotype validation</u>

We conducted reciprocal crosses between Col-0 and *atg5-1* or *atg7-2*. Seeds were collected

20 days after the cross. PCR analyses were performed for genotype validation. DNA from

10 plants grown from crossed seeds was extracted for each cross (Dellaporta *et al.*, 1983).

The   *atg5-1*   crosses   were   analyzed   with   *ATG5*   forward   primer

(ATTCACTTCCTCCTGGTGAAG), and either *ATG5* reverse primer for the WT allele

(TTGTGCCTGCAGGATAAGCG)   or   T-DNA   left   border   for   the   mutant   allele

(TAGCATCTGAATTTCATAACCAATCTCGATACAC).   The   *atg7-2*   crosses   were

analyzed with *ATG7* forward primer (GATTCAATCAACTCGCTAAGGCGT), and either

ATG7 reverse primer for the WT allele (TGCTAATTCCATGGATCCAAC) or T-DNA

left border for the mutant allele (CCCATTTGGACGTGAATGTAGACAC).

Reciprocal crosses between NahG and atg5.NahG were performed as well.

<u>Embryo development</u>

Siliques (5 from each genotype) were collected 8 and 12 days after flowering (DAF) and

opened. The seeds were discolored for 30min in Hoyer's Solution (7.5g gum arabic, 100g

chloral hydrate, 5ml glycerol, and 30ml water). Discolored seeds (40±5) were then placed

between a slide and coverslip and observed using a Nomarski objective (20x/0.50 DIC M/N2) on an Eclipse 80i microscope (Nikon) coupled to a digital camera.

Carbon starvation and recovery

Reciprocally crossed seeds and seeds of the parental lines were sown on Nitsch plates without exogenous sugars (24 plants per line per plate). After imbibition, the plants were placed in a growth chamber for 2 weeks, covered by aluminum foil for 1 week, and exposed to light for an additional week. The plates were photographed before and after carbon starvation and following a week of recovery.

Hypocotyl length measurements

Reciprocally crossed seeds were sown on Nitsch plates without exogenous sugars. Following imbibition, covered plates were transferred to a growth chamber, and the seeds were grown vertically in dark conditions for 7 days. The plates were photographed, and hypocotyl length was measured by ImageJ (Schneider *et al.*, 2012)(15-40 seedlings per line).

Metabolite extraction for lipidomics and GC-MS analysis

10 seeds from reciprocally crossed lines were weighed using the MX5 microbalance (Mattler Toledo) as previously described (Bromke *et al.*, 2015). Lipids and polar metabolites were extracted from 4-8 biological replicates using the MTBE method (Salem *et al.*, 2016). The organic phase was dried and used for lipid analysis, and the polar phase was dried and used for polar metabolite analysis.

Lipid analysis by liquid chromatography-mass spectrometry (LC-MS)

Vacuum-dried organic phases were processed using ultra-performance liquid chromatography (on a C8 reverse-phase column) coupled with Fourier transform mass spectrometry (Q exactive mass spectrometer; Thermo Fisher Scientific) in positive and negative ionization modes. Processing of chromatograms, peak detection, and integration were performed using REFINER MSH 10 (GeneData). Mass spectrometry data processing included removing fragmentation information, isotopic peaks, and chemical noise. Selected features were annotated using an in-house lipid database (Lapidot-Cohen *et al.*, 2020). More information can be found in Supplemental Table S1.

 Polar metabolite analysis by gas-chromatography-mass spectrometry (GC-MS)

Vacuum-dried polar metabolites were measured by the Agilent 7200B GC/Q-TOF. The injection and separation procedures were performed according to Dahtt *et al.* using the DB-35MS column (Dhatt *et al.*, 2019). Metabolite detection and annotation were performed by Quant software (Agilent), according to an accurate mass library of known markers generated by our group. More information can be found in Supplemental Table S2. Following blank subtraction, the peak area of each metabolite was normalized to the internal standard (i.e., ribitol) and the weight of the sample.

Total protein extraction, quantification, and immunoblotting

Seeds (~2mg, three replicates for each line) were ground. Total proteins were extracted using a protein extraction buffer [50mM pH 7.5 Tris-HCl, 3% SDS, 10% glycerol, 150mM NaCl, $1 \times$ protease inhibitor (Roche), 20mM DTT], based on a previously described extraction buffer (Di Berardino *et al.*, 2018). After the addition of buffer and centrifugation for 15min at 13,000g, supernatants were collected. Samples were incubated in SDS-PAGE sample buffer (Laemmli, 1970) for 10min at 100°C.

Total protein content was measured based on Minamide and Bamburg method(Minamide & Bamburg, 1990)with slight modifications. Bovine serum albumin (BSA) powder dissolved in distilled water to a concentration of 10 mg/ml was used to generate a standard curve, and a series of dilutions were made (0, 0.156, 0.312, 0.625, 1.25 and 2.5mg/ml). A grid of 1.5x1.5 cm squares was drawn on a sheet of Whatman number 3 filter paper and divided into 24 squares. 10µl of the standard curve dilutions and the total protein samples were applied within the squares and stained in Coomassie brilliant blue (0.25% w/v in destain solution: 50% methanol, 10% acetic acid, 40% DDW, Raymond A. Lamb) for 10 minutes.

The Whatman paper was destined with destain solution (50% methanol, 10% acetic acid, 40% distilled water) for 10 minutes and washed in water for 2 minutes. Samples were transferred to a 24-well plate with 200µl DMSO in each well. The plate was mixed gently for 10 minutes. 100µl of each sample was taken to a new 96-well plate in two technical replicates, and the absorbance of the standard curve dilutions and samples were obtained at 600 and 400 nm.

For immunoblotting, total proteins were separated by SDS-PAGE using Tris-Glycine running buffer (Fling & Gregerson, 1986) and transferred to a polyvinylidene difluoride membrane (GE Healthcare Life science), as previously described (Towbin *et al.*, 1979). Equal loading was validated using Coomassie Brilliant Blue staining of the membrane. For immunodetection, antibodies raised against 2S3 Albumin (Agrisera; AS204404) and 12S Globulin (Agrisera; AS204403) were used in combination with horseradish peroxidase-conjugated goat anti-rabbit antibody (Sigma; ab920).

Seed germination assay

Seeds were sown as described above on Nitsch plates without exogenous sugars (five plates per line, 10-60 seeds from different siliques per plate). After imbibition, the plates were transferred to a growth chamber, and the number of germinated seeds was counted every day for 4 days. Germination was scored by radicle protrusion and seedling establishment by the appearance of two green cotyledons. Germination percentage was calculated accordingly.

Seed permeability staining

Dry seeds (five replicates of 20 seeds) of each genotype were placed in a 1.5ml microcentrifuge tube. 1ml of 1% (w/v) aqueous solution of 2,3,5-triphenyltetrazolium chloride (Sigma) was added to each tube as previously described (Vishwanath *et al.*, 2014). The tubes were incubated in an air incubator at 30°C in the dark for a period of 24 and 48 hours. After incubation, seeds (10 seeds of each genotype) were observed for changes in seed color and imaged using a stereomicroscope.

The extraction of formazans was performed as previously described (Vishwanath *et al.*, 2014). After incubation, as mentioned above, seeds were washed twice with distilled water. Then 1ml of 95% ethanol was added to the seeds, and the mix was ground using a mortar and pestle. The ground seed material in ethanol solution was transferred to a microcentrifuge tube, and the final volume was adjusted to 1ml with 95% ethanol. The tubes were immediately centrifuged at 15,000xg for 3min. The supernatant was collected into a spectrophotometer cuvette, and the absorbance of the formazan extracts was measured at 485nm using a spectrophotometer. As the blank, 95% ethanol solution was used.

<u>Seed coat visualization</u>

For seed coat visualization, mature seeds were pre-hydrated with water for 90min and then stained as follows. Ruthenium Red staining for pectins was prepared as previously described (Willats *et al.*, 2001), and hydrated seeds were incubated in 0.01% (w/v) Ruthenium Red (Sigma) for 90min at room temperature. Seeds were washed twice in water and then photographed. Image analysis was performed by ImageJ (Schneider *et al.*, 2012).

Calcofluor staining was conducted as previously described (Willats *et al.*, 2001) using $25\mu g\ ml^{-1}$ calcofluor (Fluorescent Brightener 28, Sigma) for 20min at room temperature (5 seeds for each genotype, cesa5 (SALK_118491) as a negative control). Seeds were washed for 2h in water and then photographed. Observations were conducted using the FV-1200 confocal microscope (Olympus), with a 10x/0.4 or 20x/0.75 objective microscope equipped with 405nm (Excitation) and 430-470nm (Emission) laser diode. Image analysis was performed by ImageJ (Schneider *et al.*, 2012).

For seed coat autofluorescence, mature seeds of WT, *atg5-1*, and *atg7-2* were illuminated and observed with an IX81 microscope (Olympus) using DAPI interference and absorption filter (excitation filter, 350/50nm; emission filter, 460/50nm) coupled to a digital camera (Li *et al.*, 2007).

For phenol dyeing, green seeds (12-16 DAF) of WT, *atg5-1*, and *atg7-2* were incubated in $FeCl_3$ solution (2% in 95% EtOH (Mace, 1963)) for 5min. 95% EtOH was used as A negative control. For cuticle permeability, green seeds (12-16 DAF) of Col-0, *atg5-1*, and *atg7-2* were hydrated for 10min. Hydrated seeds were incubated in an aqueous solution of 0.05% (w/v) Toluidine-Blue (Sigma) for 2 min and then washed with DDW twice (Mace, 1963). Hydration in 0.5% EDTA solution with three DDW washes was used as a positive control.

<u>Scanning electron microscopy (SEM)</u>

Dry seeds were mounted on aluminum stubs (Electron Microscopy Sciences), sputter-coated with 10nm of Au/Pd (5% Pd), and viewed using "Quanta 200", field emission scanning electron microscope (Field Emission Instruments). 6 seeds were examined for each genotype. Measurement of 15 epidermis cells from each seed was performed by ImageJ (Schneider *et al.*, 2012).

Artificial seed aging

Artificial aging was performed as previously described (Nguyen *et al.*, 2012). 50mg of seeds were stored above a saturated NaCl solution in a closed desiccator (relative humidity of 75%) for 70 days. Germination assay was performed on aged and non-aged seeds as described above. For developmental stage scoring, seeds were artificially aged for 35 days and sown on Nitsch plates without exogenous sugars. The plants were grown as described above for 14 days after imbibition. Seedling development was scored daily as previously described (Boyes *et al.*, 2001).

Statistical analysis

The experiments were conducted in a random blocks design with three to six biological replicates of each genotype. Data were statistically examined using analysis of variance and tested for significant (* = $p<0.05$, ** = $p<0.01$) differences using Student's *t-test* from an algorithm embedded in Microsoft Excel. Heatmaps were generated using the MultiExperiment Viewer (MeV) freely available software application(Howe *et al.*, 2010). Principle Component Analysis was performed using Metaboanalyst 5.0(Chong *et al.*, 2019). Mixed model analysis was performed using Prism (GraphPad Software).

Results

<u>F1 progeny of WT and *atg* mutants possess a functional autophagy mechanism</u>

To differentiate between autophagy in the mother plant and in the embryo, we performed

reciprocal crosses between WT and *atg* mutant (*atg5-1* or *atg7-2*) Arabidopsis plants. The

heterozygous genotype of F1 plants was validated by PCR (Fig. S1a,b). We did not observe

any developmental difference between the F1 seeds (Fig. 1a). In addition, we did not detect

any aborted seeds, as previously reported in *atg5-1* plants (Di Berardino *et al.*, 2018).

Homozygous *atg* mutant seeds were shown to display reduced weight (Barros *et al.*, 2017;

Minina *et al.*, 2018). We thus compared the weight of reciprocally crossed F1 seeds. We

did not see a significant difference between the lines, suggesting this phenotype might be

governed by autophagy in the developing embryo (Fig. 1b).

Fig. 1: F1 progeny of WT and *atg* mutants display normal development. (a) F1 seeds were collected at 8DAF and 12DAF, uncolored, and photographed. No aborted embryos were observed. Representative image of 40±5 seeds examined. Scale bar=100μm. (b) Weight of 10 F1 seeds. Data are presented as a box&whiskers plot. No significant differences were observed (n=4-8).

We next wished to determine whether the F1 plants display a functional autophagy

mechanism. One hallmark phenotype of *atg* mutants is hypersensitivity to carbon

starvation (Marshall & Vierstra, 2018). We thus exposed two-week-old F1 seedlings and

their respective paternal lines to carbon starvation and recovery. As can be seen in Fig. 2a-

c and Fig. S1c-d, the *atg* mutant parental lines were more susceptible to carbon starvation

than WT plants and did not recover, as expected. F1 seedlings displayed a WT-like

phenotype, irrespective of the identity of the maternal or paternal lines. *atg* mutant etiolated

seedlings were also shown to exhibit shorter hypocotyls when grown without exogenous sugars (Avin-Wittenberg *et al.*, 2015). Interestingly, when we performed this assay on the F1 progeny of our reciprocal crosses, seedlings stemming from *atg* mutant mother plants had significantly shorter hypocotyls than seedlings from WT mother plants (Fig. 2d,e). We hypothesized this phenotype resulted from altered storage compound amounts deposited in seeds from different mother plants.

As *atg* mutants display an early senescence phenotype (Marshall & Vierstra, 2018), we wished to ascertain whether the observed phenotype did not stem from the premature senescence of the mother plant, resulting in diminished reserve accumulation. We thus performed reciprocal crosses between NahG and *atg5*.NahG plants, in which the overexpression of the bacterial enzyme NahG results in salicylic acid degradation, inhibiting the early senescence phenotype(Yoshimoto *et al.*, 2009). F1 etiolated seedlings from maternal *atg5*.NahG source displayed shorter hypocotyls than seedlings from maternal NahG plants (Fig. 2f). This result suggests that the phenotype observed for F1 seedlings from maternal *atg* mutants is not a result of early senescence.

Fig. 2: Autophagy-related phenotypes of F1 seedlings. a-c: Representative images of F1 progeny of reciprocal crosses between WT and *atg5-1* and their respective parental lines. Seeds were germinated and grown on Nitsch plated without exogenous sucrose for 14 days after imbibition (a). The plants were incubated in the dark for additional 7 days (b). The plants were then returned to light for another 7 days to examine their recovery (c). The experiment was performed in triplicate. (d) Representative image of F1 progeny of reciprocal crosses between WT and *atg5-1* or *atg7-2*. Seeds were sown on Nitsch plates without exogenous sugars and grown in the dark for 7 days after imbibition. (e) Hypocotyl length of the lines described in (d), as quantified by ImageJ. Data are presented as a box&whiskers plot. One asterisk (p<0.05) or two asterisks (p<0.01) denote a significant difference following Student's *t-test* (n=15-30). (f) Hypocotyl length of F1 progeny of reciprocal crosses between NahG and *atg5*.NahG. Seeds were sown on Nitsch plates without exogenous sugars, grown in the dark for 7 days after imbibition, and their hypocotyl length was measured by ImageJ. Two asterisks (p<0.01) denote a significant difference following Student's *t-test* (n=20-30).

1   <u>F1 seeds of maternal *atg* mutant plants display reduced storage protein but not lipid content</u>

2   We next wished to test our hypothesis of reduced storage compounds in F1 seeds of

3   maternal *atg* mutant origin. As Arabidopsis is an oil seed, we performed lipidomics of dry

4   seeds of F1 progeny of reciprocal crosses between WT, *atg5-1,* or *atg7-2*. Though we

5   annotated more than 120 lipids, we did not observe any separation between the groups in

1 the principal component analysis (PCA – Fig. 3a). A closer look at TAG species revealed

2 no significant differences or apparent trends between F1 seeds of WT and *atg* mutant

3 maternal origin (Fig. 3b). These results are in line with previous works, demonstrating no

4 reduction in the total lipid content of *atg* mutant seeds compared to WT seeds (Minina *et*

5 *al.*, 2018 1218).

Fig. 3: Autophagy deficiency in the mother plant does not affect seed lipid content. Dry seeds of F1 progeny of reciprocal crosses between WT and *atg5-1* or *atg7*-2 were collected, and their lipid content was analyzed by UPLC-MS (n=4-8). Detailed results are presented in Supplemental table S1. (a) PCA of lipid levels. (b). Heat map of $\log_2$ of TAG relative levels in comparison to seeds from maternal WT lines. No significant differences between reciprocally crossed lines were identified by student's *t-test*.

We then examined the total protein content of F1 seeds. Surprisingly, dry seeds of F1

progeny of maternal *atg* mutant origin contained lower total protein levels than the

reciprocally-crossed seeds. The difference was significant for the *atg7-2* crosses, and a

similar trend was observed for the *atg5-1* crosses (Fig. 4a). The results indicate that the

protein content, but not the lipid content of the seed, is affected by the mother plan,

suggesting a possible differential regulation of carbon and nitrogen allocation by autophagy

in the mother plant. To confirm our hypothesis, we analyzed the polar metabolites of dry

F1 seeds by GC-MS. Unfortunately, we could not find significant differences between the

1 reciprocally crossed lines for the 17 metabolites annotated and could not detect free amino

2 acids (Fig. S2). Previous work indicated that *atg* mutant seeds display altered storage

Fig. 4: Lack of autophagy in the mother plant alters total seed protein content but not storage-protein processing. Total proteins were extracted from dry seeds of F1 progeny of reciprocal crosses between WT and *atg5-1* or *atg7*-2. (a) Total protein quantification. Data are presented as average ± SE. An asterisk denotes a significant difference in Student's *t-test* (p<0.05, n=3). (b) Total proteins were separated by SDS-PAGE. Top panel – Western blot using anti-12S antibody. Middle panel – Western blot using anti-2S antibody. Bottom panel – Coomassie blue stain. Three biological replicates were performed. P12S – precursors of the 12S protein, α and β – 12S globulin subunits.

protein processing (Di Berardino *et al.*, 2018). We, therefore, performed Western blot

analysis against 12S globulin and 2S albumin. Though we observed a general reduction in

protein amounts, we did not observe a change in the storage protein pattern, indicating

there was no difference in protein processing (Fig. 4b). Our results suggest that the altered

storage-protein processing in *atg* mutant seeds is probably due to a lack of autophagic

activity in the seed, rather than the mother plant.

F1 seeds of maternal *atg* mutant plants display faster germination rates stemming from

altered seed coat morphology

A typical phenotype of *atg* mutants is slightly delayed germination and seedling

establishment (Avin-Wittenberg *et al.*, 2015). We, therefore, examined whether this

phenotype stems from autophagy in the mother plant. We generated reciprocally-crossed

F1 seeds and tested their germination and seedling establishment rates. Surprisingly, we

observed increased germination rates for F1 progeny of *atg* mutant mother plants than WT

mother plants (Fig. 5a). We did not observe significant differences in seedling

establishment (Fig. 5b). We repeated the experiment for F1 seeds of reciprocally-crossed

Fig. 5: F1 seeds of maternal *atg* mutants exhibit faster germination and increased permeability. (a,b) F1 seeds of reciprocal crosses between WT and *atg* mutants were sown on Nitsch plates without exogenous sucrose, imbibed for 72h, and transferred to continuous light conditions. Germination (defined by radicle protrusion) and seedling establishment (defined by the appearance of two green cotyledons) were scored each day for 4 days. (a) Average percent germination is presented ± SE. (b) Average percent of seedling establishment is presented ± SE. An asterisk denotes a significant difference in Student's *t-test* (p<0.05, n=5). (c) Representative image of F1 seeds from reciprocal crosses after incubation for 24h (top panel) or 48h (bottom panel) in tetrazolium chloride solution. Scale bar=0.5mm. (d, e) Quantification of tetrazolium chloride staining, Stained seeds were ground and dissolved in 95% EtOH. absorption in 485nm was measured. Average absorption is presented ± SE for F1 seeds (d) or homozygous parental lines. . One asterisk (p<0.05) or two asterisks (p<0.01) denote a significant difference following Student's *t-test* (n=5).

NahG and *atg5*.NahG plants and received similar results, suggesting the phenotype is

unrelated to early senescence (Fig. S3).

We postulated the higher germination rate of F1 progeny of *atg* mutant mother plants resulted from faster water penetration into the seed. We thus examined seed permeability by tetrazolium chloride staining. Tetrazolium salts are amphipathic cations. After penetrating the dead cells of the seed coat, they are reduced by active dehydrogenases (NADH-dependent reductases) in the embryo to red-colored insoluble precipitates composed of formazans (Berridge, 1996). The intensity of red coloration is directly proportional to the permeability of the seeds. We stained dry F1 seeds for 24 and 48 hours. As shown in Fig. 5c,d F1 seeds of maternal *atg5* plants displayed significantly higher water permeability than F1 seeds of maternal WT origin. A similar, albeit insignificant trend, was observed for the *atg7* reciprocal crosses. Interestingly, we also observed increased water permeability in homozygous *atg* mutant seeds (Fig. 5e).

As water permeability is influenced by the seed coat, which is of maternal origin (Marbach & Mayer, 1974), we took a closer look at the seed coat of F1 seeds and their parental lines. We initially focused on the hilum, the scar left on the seed coat after detachment from the funiculus. In mature Arabidopsis seeds, the hilum is adjacent to the micropyle (where the radicle will emerge) and faces the chalazal pole (Beisson *et al.*, 2007). The hilum contains high levels of suberin, inhibiting the entrance of water and pathogens. We compared the suberin levels of homozygous WT and *atg* mutant seeds by autofluorescence. We did not observe any differences between the lines (Fig. S4a). We next examined the integrity of the cuticle layer that wraps the embryo (Steinbrecher & Leubner-Metzger, 2018). We stained WT and *atg* mutant seeds with toluidine blue (TB), which stains the layers underneath the cuticle when it is damaged. As a positive control, seeds were pre-incubated with EDTA to perforate the coticule. We did not detect any differences between the lines; none were stained, suggesting their cuticle was intact (Fig. S4b). The seed coat contains phenols in many species, inhibiting water entrance (Werker *et al.*, 1979). We stained WT and *atg* mutant seeds with $FeCl_3$ to observe phenols. No genotype displayed dark spots as expected in the presence of phenols (Fig. S4c).

Hydrated Arabidopsis seeds are coated by a gelatinous layer called mucilage, which is mainly composed of cell wall polysaccharides (Voiniciuc *et al.*, 2016). It is deposited in the seed coat epidermal cells during seed development in a structure at the center of the

1   cell termed "columella" and released during imbibition(Griffiths & North, 2017). Mucilage

2   is rich in pectin, and its architecture can be visualized by ruthenium red (RR) dye. When

3   we dyed F1 seeds with RR staining, we found that seeds from maternal *atg5-1* plants

4   displayed lower mucilage content than their reciprocal F1 seeds. We observed a similar

5   trend for F1 seeds of WT and *atg7-2*, yet not statistically significant (Fig. 6a,b). Adherent

Fig. 6: Altered mucilage structure of maternal *atg* mutant F1 seeds. (a,b) F1 seeds of reciprocal crosses between WT and *atg* mutants were stained with ruthenium red solution. (a) Representative images of F1 seeds from reciprocal crosses. Scale bar = 200μm. (b) Quantification of mucilage area. Data are presented as a box&whiskers plot. One asterisk denotes a significant difference following Student's *t-test* (p<0.05, n=10). (c,d) Calcofluor staining for β-glucans of F1 seeds. (c) Representative images of whole-seed views (top panel) and close-ups (bottom panel). Scale bars = 250μm (top panel) and 100μm (bottom panel). (d) Quantification of ray density. Data are presented as a box&whiskers plot. One asterisk denotes a significant difference following Student's *t-test* (p<0.05, n=3-4).

mucilage is partially anchored by cellulosic rays (Sullivan *et al.*, 2011). We performed

calcofluor staining of F1 seeds to visualize mucilage rays. Seeds of maternal *atg5-1* plants

displayed higher ray density than their reciprocal F1 seeds. A similar trend was observed

for the *atg7-2* cross (Fig. 6c,d).

Finally, we assessed the general structure of F1 seeds and seed coat cells by scanning

electron microscopy (SEM). F1 progeny of maternal *atg* mutants displayed altered seed

shape compared to F1 seeds originating from WT mother plants, stemming from reduced

seed width (Fig. 7a-top panel,b). We also observed that the seed coat cells of F1 seeds from

1 *atg* mutant mother plants were misshapen (Fig. 7a-bottom panel). The columella of these

2 cells occupied more of the cell area than in seeds from WT mother plants. When we

Fig. 7: Autophagy in the mother plants has a negative effect on seed shape and aging. (a) Scanning electron microscopy (SEM) of F1 seeds produced by reciprocal crosses between WT and *atg* mutant plants. Top panel, scale bar = 100μm, bottom panel, scale bar = 50μm. The columella tip is marked in purple, and the cell area is marked in yellow (bottom panel). (b) Seed dimensions were measured by ImageJ. Data are presented as a box&whiskers plot. One asterisk denotes a significant difference following Student's *t-test* (p<0.05, n=6). (c) The ratio between the columella and total cell area in individual F1 seeds of WT and *atg7-2* reciprocal crosses. Data are presented as a box&whiskers plot, p-value denotes the statistical difference between the seeds by mixed model analysis (n=12-15 cells per seed). (d) WT and *atg* mutant seeds without artificial aging and following 10 weeks of artificial aging were sown on Nitsch plates without exogenous sucrose, imbibed for 72h, and transferred to continuous light conditions. Germination (defined by radicle protrusion) was scored each day for 4 days. Average percent germination is presented ± SE. One asterisk (p<0.05) or two asterisks (p<0.01) denote a significant difference following Student's *t-test* (n=5, dark gray - *atg5-1*, light gray – *atg7-2*).

3 quantified the ratio between the columella and total cell area, we observed a significantly

4 higher ratio for F1 seeds of maternal *atg7-2* plants than WT plants (Fig. 7c). A similar trend

5 was seen in F1 seeds of *atg5-1* and WT reciprocal crosses (Fig. S5).

<u>*atg* mutants exhibit more rapid seed aging compared to WT plants</u>

The seed coat structure plays an important role in seed longevity (Zhou *et al.*, 2019). We, therefore, performed artificial aging of WT and *atg* mutant seeds and tested their germination ability following aging. Without aging, *atg* mutants displayed slightly delayed germination, as previously described (Avin-Wittenberg *et al.*, 2015). Indeed, all lines were affected by seed aging, showing reduced germination after 10 weeks of artificial aging. As expected, the germination of *atg* mutants was more strongly affected by artificial aging than that of WT seeds, strengthening the link between the altered seed coat of *atg* mutants and seed longevity (Fig. 7d). We also scored seedling development following 5 weeks of artificial aging (Boyes *et al.*, 2001). Artificial aging resulted in delayed development of all the lines. Yet, *atg* mutants developed slower than WT seedlings (Fig. S6).

Discussion

Autophagy is gaining attention as a fundamental mechanism in plants, affecting differentiation (Rodriguez *et al.*, 2020), stress response (Tang & Bassham, 2021), and nutrient availability and remobilization (Magen *et al.*, 2022). Yet, the standard approach to studying autophagy in plants is using ubiquitous knockout or knockdown mutants. Though applicable in many cases, it may obscure our understanding of the function of autophagy in specific tissues or time points, specifically when examining source-sink relationships. *atg* mutant seeds display a plethora of phenotypes, from reduced storage protein processing (Di Berardino *et al.*, 2018) to altered C/N ratio (Guiboileau *et al.*, 2012). Yet, it is impossible to determine the tissue responsible for the phenotype using the current system. We performed reciprocal crosses to separate autophagy in the mother plant and the embryo to address this issue.

The main challenge of our approach is the relatively small number of seeds we could achieve by reciprocal crosses. We were thus unable to perform certain analyses requiring larger material quantities, such as seed coat composition and metabolic flux analysis. Generating transgenic plants in which autophagy is selectively downregulated either in the embryo or the mother plant will allow conducting experiments on a larger scale as well as investigate both sides of the "source-sink coin" concerning autophagy activity. In addition,

a transgenic system could be used to investigate the seed set phenotype of *atg* mutants, which reciprocal crosses can not address. A recent example is a study examining autophagy in root cap cells that used tissue-specific CRISPR to down-regulate autophagy specifically in this tissue (Feng *et al.*, 2022). Nevertheless, we believe our work lays the ground for the in-depth investigation of autophagy in discrete plant tissues and the relationship between tissues.

We determined that F1 seeds displayed WT a carbon starvation phenotype when exposed to dark treatment at the seedling stage, suggesting the autophagy mechanism is active in F1 seedlings (Fig. 2a-c, S1c-e). In contrast, when we examined etiolated seedlings grown without exogenous sugars, F1 seeds from maternal *atg* mutants displayed reduced growth (Fig. 2d,e). These results suggest alerted storage compound accumulation, depending on the presence of autophagy in the mother plant. We observed no significant difference in seed lipid content between F1 seeds (Fig. 3). This aligns with previously reported data, in which homozygous *atg* mutant seeds did not display altered total fatty acid content. Interestingly, the same paper showed that overexpression of *ATG5* or *ATG7* yielded higher fatty acid content (Minina *et al.*, 2018). This could stem from higher sink strength in the mother plant, pushing carbon to the seeds, or increased demand by the embryo.

Unlike our lipid analysis, we observed reduced protein levels in F1 seeds of maternal *atg* mutant origin (Fig. 4a). Previous studies showed that *atg* mutant plants remobilize less nitrogen from the mother plant to developing seeds (Guiboileau *et al.*, 2012; Li *et al.*, 2015). Together with our observations, this strengthens our hypothesis that the altered protein amounts in F1 seeds stem from differential remobilization from the mother plant. Interestingly, we did not observe any difference in storage protein processing, as described for *atg5* mutant seeds (Di Berardino *et al.*, 2018), suggesting this phenotype results from autophagy deficiency in the embryo. The involvement of autophagy in storage protein delivery in seeds has been implied yet hardly investigated, and mostly in monocot, rather than dicot seeds(Levanony *et al.*, 1992; Reyes *et al.*, 2011). More work is needed to elucidate the role of autophagy in storage protein processing.

An important seed tissue we did not address in the scope of this research is the endosperm. This is a triploid tissue that is partly maternal (2n) and partly paternal (1n) (Bentsink &

Koornneef, 2008). This uneven distribution creates a situation in which F1 seeds of maternal WT plants possess more copies of *ATG* genes in their endosperm than F1 seeds of maternal *atg* mutant origin, which we could not directly investigate due to the seed amount issue described above. We are unaware of studies examining the effect of *ATG* gene copy number on plant autophagy activation. However, We did observe that F1 seeds containing a lower copy number of *ATG* genes behaved similarly to WT plants under carbon starvation (Fig. 2a-c, S1c-e), suggesting that one copy is sufficient for autophagy activation. Nonetheless, we cannot disregard the contribution of the endosperm to proper embryo growth and development(Song *et al.*, 2021). Future studies are necessary to investigate the function of autophagy in the seed endosperm, perhaps utilizing monocot seeds that have a larger endosperm.

We were most surprised by the observation that F1 seeds of maternal *atg* mutant plants exhibited faster germination compared to F1 seeds from maternal WT plants (Fig. 5a). We attributed this phenotype to increased water permeability of F1 seeds of maternal *atg* mutant origin (Fig. 5c,d). Interestingly, homozygous *atg* mutant seeds also exhibited increased water permeability, though they are known to germinate slightly slower than WT seeds (Avin-Wittenberg *et al.*, 2015). We postulate that the slower germination of homozygous *atg* mutants is, in fact, a compound phenotype affected by the lack of autophagy in the seed coat and the embryo. Examining the germination of seeds in which autophagy is selectively downregulated in the embryo might help settle this dispute. The reason for the increased water permeability of atg mutants is still unclear. Yet, we could observe several differences in the seed coat of F1 seeds of maternal *atg* mutant origin (Fig. 6, Fig. 7a-c). More extensive analysis of the seed coat of atg mutants is required to fully elucidate the causes of this phenotype. One possibility is the involvement of autophagy in regulating the programmed cell death of the seed coat cells (Sueldo & van der Hoorn, 2017), resulting in deformed seed coat in *atg* mutants. Of note, the increased water permeability is not a result of the early senescence phenotype of the *atg* mutant mother plant, as F1 seeds of stay-green atg mutants displayed a similar phenotype (Fig. S3).

To conclude, we have shown that the seed phenotype of *atg* mutants is a compound phenotype resulting from the lack of autophagy in both the other plant and the embryo.

This is very important, especially in a tissue affected by nutrient remobilization. Our work highlights the importance of studying the role of autophagy in higher resolution while keeping in mind the whole-plant context. We believe that the generation of plant lines in which autophagy is down-regulated explicitly in space or time will prove to be a valuable tool to achieve this goal.

Acknowledgments

We thank Dr. Smadar Harpaz-Saad of the Hebrew University of Jerusalem for her assistance with Arabidopsis seed coat analysis. We also thank Dr. Naomi Melamed-Book from the microscopy core facility at The Hebrew University's Life Science core facility for her technical support with confocal analysis and Dr. Evgenia Blayvas from the Hebrew University's Unit for Nanoscience, for technical support with SEM.

Author Contributions (move after references)

O.E. designed the research, performed research, analyzed data, and wrote the manuscript. S.W., M.A-A, M.A., Y.S., W.J, L.R, and Y.B performed research and analyzed data. T.A-W designed the research and wrote the manuscript. The project was funded by the Israeli Science Foundation, grant number 1942/19.

Competing interests

The authors declare no competing interests.

Supporting information

Fig. S1: F1 seedlings of reciprocal crosses are heterozygous and display a WT phenotype. a,b: PCR analysis for DNA extracted from F1 seedlings (n=5 from different siliques) and parental lines, with primers for the WT allele and T-DNA insertion. (a) atg5-1, (b) atg7-2. c-e: Representative images of F1 progeny of reciprocal crosses between WT and atg7-2 and their respective parental lines. Seeds were germinated and grown on Nitsch plated without exogenous sucrose for 14 days after imbibition (c). The plants were

1 incubated in the dark for additional 7 days (d). The plants were then returned to light for

2 another 7 days to examine their recovery (e). The experiment was performed in triplicate.

Fig. S2: No difference between F1 seeds in primary metabolites. Dry seeds of F1

progeny of reciprocal crosses between WT and *atg5-1* or *atg7-2* were collected, and their

polar metabolite content was analyzed by GC-MS (n=4-8). Detailed results are presented

in Supplemental Data Set S2. (a) PCA of metabolite levels. (b). Heat map of log2 of

metabolite relative levels in comparison to seeds from maternal WT lines. An asterisk

denotes a significant difference between reciprocally crossed lines by Student's *t-test*

(p<0.05).

Fig. S3: The early germination of maternal *atg* mutant seedlings is not salicylic acid-

dependent. (a,b) F1 seeds of reciprocal crosses between NahG and *atg5*.NahG mutants

were sown on Nitsch plates without exogenous sucrose, imbibed for 72h, and transferred

to continuous light conditions. Germination (defined by radicle protrusion) and seedling

establishment (defined by the appearance of two green cotyledons) were scored each day

for 4 days. (a) Average percent germination is presented ± SE. (b) Average percent of

seedling establishment is presented ± SE. An asterisk denotes a significant difference in

Student's *t-test* (p<0.05, n=5).

Fig. S4: No difference between WT and *atg* mutant seeds in several biochemical

factors affecting water permeability. (a) Seed coat autofluorescence was visualized using

DAPI filter. All genotypes show saturation in the hilum region (designated by yellow

arrowheads) because of high Suberin content. 30-50 seeds per genotype were examined.

Scale bar=200µm. (b) TB staining for cuticle permeability. No genotype shows blue

staining as the positive control (pre-incubation in 0.5% EDTA solution). 5 siliques were

examined per genotype. Scale bar=0.5mm. (c) FeCl3 staining for phenols. No genotype

shows dark spots predicted to appear in the presence of phenols. Negative control with

DDW. 5 siliques were examined per genotype. Scale bar=0.5mm.

Fig. S5: Autophagy in the mother plant has a negative effect columella/cell area ratio.

The ratio between the columella and total cell area in individual F1 seeds of WT and *atg5-*

1 reciprocal crosses. Data are presented as a box&whiskers plot, p-value denotes the

statistical difference between the seeds by mixed model analysis (n=12-15 cells per seed).

Fig. S6: Artificial aging affects seedling development of *atg* mutants. WT and *atg*

mutant seeds without artificial aging and following 5 weeks of artificial aging were sown

on Nitsch plates without exogenous sucrose, imbibed for 72h, and transferred to continuous

light conditions. Growth stage progression was scored for seedlings grown vertically. Data

represent average ± SE (n35). Asterisk indicates significant differences from WT in each

aging time (Student's t test; p<0.05).

Supplemental Table S1: Relative lipid content of reciprocally crossed Arabidopsis

thaliana F1 seeds.

Supplemental Table S2: Relative polar metabolite content of reciprocally crossed

Arabidopsis thaliana F1 seeds.

References

Avin-Wittenberg T, Bajdzienko K, Wittenberg G, Alseekh S, Tohge T, Bock R,
Giavalisco P, Fernie AR. 2015. Global analysis of the role of autophagy in cellular
metabolism and energy homeostasis in Arabidopsis seedlings under carbon starvation.
Plant Cell **2**.306-322 :(2)7
Avin-Wittenberg T, Baluška F, Bozhkov PV, Elander PH, Fernie AR, Galili G,
Hassan A, Hofius D, Isono E, Le Bars R, et al. 2018. Autophagy-related approaches for
improving nutrient use efficiency and crop yield protection. *Journal of Experimental*
Botany **69**(6): 1335-1353.
Avin-Wittenberg T, Honig A, Galili G. 2012. Variations on a theme: plant autophagy in
comparison to yeast and mammals. *Protoplasma* **249**(2): 285-299.
Barros J, Magen S, Lapidot-Cohen T, Rosental L, Brotman Y, Araújo W, Avin-
Wittenberg T. 2021. Autophagy is required for lipid homeostasis during dark-induced
senescence. *Plant Physiology* **185**(4): 1542-1558.
Barros JA, Cavalcanti JHF, Medeiros DB, Nunes-Nesi A, Avin-Wittenberg T, Fernie
AR, Araujo W. 2017. Autophagy deficiency compromises alternative pathways of
respiration following energy deprivation. *Plant Physiol* **175** (1):62-76.
Bassham DC. 2009. Function and regulation of macroautophagy in plants. *Biochim*
Biophys Acta **1793**(9): 1397-1403.
Bassham DC, Laporte M, Marty F, Moriyasu Y, Ohsumi Y ,Olsen LJ, Yoshimoto K.
2006. Autophagy in development and stress responses of plants. *Autophagy* **2**(1): 2-11.

Baud S, Dubreucq B, Miquel M, Rochat C, Lepiniec L. 2008. Storage reserve accumulation in Arabidopsis: metabolic and developmental control of seed filling. *Arabidopsis Book* **6**: e0113.

Beisson F, Li Y, Bonaventure G, Pollard M, Ohlrogge JB. 2007. The Acyltransferase GPAT5 Is Required for the Synthesis of Suberin in Seed Coat and Root of Arabidopsis. *The Plant Cell* **19**(1): 351-368.

Bentsink L, Koornneef M. 2008. Seed Dormancy and Germination. *The Arabidopsis Book* **6**.

Berridge MV, Tan, A. S., McCoy, K. D. , Wang, R. . 1996. The Biochemical and Cellular Basis of Cell Proliferation Assays That Use Tetrazolium Salts. *Biochemica* **4**(1): 15-19.

Boyes DC ,Zayed AM, Ascenzi R, McCaskill AJ, Hoffman NE, Davis KR, Görlach J. 2001. Growth Stage–Based Phenotypic Analysis of Arabidopsis. *The Plant Cell* **13**(7): 1499-1510.

Bromke MA, Hochmuth A, Tohge T, Fernie AR, Giavalisco P, Burgos A, Willmitzer L, Brotman Y .2015 .Liquid chromatography high-resolution mass spectrometry for fatty acid profiling. *Plant J* **81**(3): 529-536.

Cernac A, Benning C. 2004. WRINKLED1 encodes an AP2/EREB domain protein involved in the control of storage compound biosynthesis in Arabidopsis .*The Plant Journal* **40**(4): 575-585.

Chong J, Wishart DS, Xia J. 2019. Using MetaboAnalyst 4.0 for Comprehensive and Integrative Metabolomics Data Analysis. *Current Protocols in Bioinformatics* **68**.(1)

Chung T, Suttangkakul A, Vierstra RD. 2009. The ATG autophagic conjugation system in maize: ATG transcripts and abundance of the ATG8-lipid adduct are regulated by development and nutrient availability. *Plant Physiol* **149**(1): 220-234.

Dellaporta SL, Wood J, Hicks JB. 1983. A plant DNA minipreparation: Version II. *Plant Molecular Biology Reporter* **1**(4): 19-21.

Dhatt BK, Abshire N, Paul P, Hasanthika K, Sandhu J, Zhang Q, Obata T, Walia H. 2019. Metabolic Dynamics of Developing Rice Seeds Under High Night-Time Temperature Stress. *Front Plant Sci* **10**: 1443.

Di Berardino J, Marmagne A, Berger A, Yoshimoto K, Cueff G, Chardon F, Masclaux-Daubresse C, Reisdorf-Cren M, Bozhkov P. 2018. Autophagy controls resource allocation and protein storage accumulation in Arabidopsis seeds. *Journal of Experimental Botany* **69**(6): 1403-1414.

Distelfeld A, Avni R, Fischer AM. 2014. Senescence, nutrient remobilization, and yield in wheat and barley. *Journal of Experimental Botany* **65**(14): 3783-3798.

Doelling JH, Walker JM, Friedman EM, Thompson AR, Vierstra RD. 2002. The APG8/12-activating enzyme APG7 is required for proper nutrient recycling and senescence in Arabidopsis thaliana. *J Biol Chem* **277**(36): 33105-33114.

Feng Q, De Rycke R, Dagdas Y, Nowack MK. 2022. Autophagy promotes programmed cell death and corpse clearance in specific cell types of the Arabidopsis root cap. *Current Biology* **32**(9): 2110-2119.e2113.

Fling SP, Gregerson DS. 1986. Peptide and protein molecular weight determination by electrophoresis using a high-molarity tris buffer system without urea. *Anal Biochem* **155**(1): 83-88.

Griffiths JS, North HM. 2017. Sticking to cellulose: exploiting Arabidopsis seed coat mucilage to understand cellulose biosynthesis and cell wall polysaccharide interactions. *New Phytologist* **214**(3): 959-966.

Guiboileau A, Yoshimoto K, Soulay F, Bataille MP, Avice JC, Masclaux-Daubresse C. 2012. Autophagy machinery controls nitrogen remobilization at the whole-plant level under both limiting and ample nitrate conditions in Arabidopsis. *New Phytol* **194**(3): 732-740.

Hirota T, Izumi M, Wada S, Makino A, Ishida H. 2018. Vacuolar Protein Degradation via Autophagy Provides Substrates to Amino Acid Catabolic Pathways as an Adaptive Response to Sugar Starvation in Arabidopsis thaliana. *Plant and Cell Physiology*.

Hofius D, Schultz-Larsen T, Joensen J, Tsitsigiannis DI, Petersen NH, Mattsson O, Jorgensen LB, Jones JD, Mundy J, Petersen M. 2009. Autophagic components contribute to hypersensitive cell death in Arabidopsis. *Cell* **137**(4): 773-783.

Howe E, Holton K, Nair S, Schlauch D, Sinha R, Quackenbush J 2010. MeV : MultiExperiment Viewer. *Biomedical Informatics for Cancer Research*, 267-277.

Kurusu T, Koyano T, Hanamata S, Kubo T, Noguchi Y, Yagi C, Nagata N, Yamamoto T, Ohnishi T, Okazaki Y, et al. 2014. OsATG7 is required for autophagy-dependent lipid metabolism in rice postmeiotic anther development. *Autophagy* **10**(5): 878-888.

Laemmli UK. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**(5259): 680-685.

Lapidot-Cohen T, Rosental L, Brotman Y. 2020. Liquid Chromatography-Mass Spectrometry (LC-MS)-Based Analysis for Lipophilic Compound Profiling in Plants. *Curr Protoc Plant Biol* **5**(2): e20109.

Levanony H, Rubin R, Altschuler Y, Galili G. 1992. Evidence for a novel route of wheat storage proteins to vacuoles. *J Cell Biol* **119**(5): 1117-1128.

Li F, Chung T, Pennington JG, Federico ML, Kaeppler HF, Kaeppler SM, Otegui MS, Vierstra RD. 2015. Autophagic recycling plays a central role in maize nitrogen remobilization. *Plant Cell* **27**(5): 1389-1408.

Li Y, Beisson F, Koo AJ, Molina I, Pollard M, Ohlrogge J. 2007. Identification of acyltransferases required for cutin biosynthesis and production of cutin with suberin-like monomers. *Proc Natl Acad Sci U S A* **104**(46): 18339-18344.

Liu Y, Bassham DC. 2012. Autophagy: pathways for self-eating in plant cells. *Annu Rev Plant Biol* **63**: 215-237.

Lobell DB, Schlenker W, Costa-Roberts J. 2011. Climate Trends and Global Crop Production Since 1980. *Science* **333**(6042): 616-620.

Mace ME. 1963. Histochemical Localization of Phenols in Healthy and Diseased Banana Roots. *Physiologia Plantarum* **16**(4): 915-925.

Magen S, Seybold H, Laloum D, Avin-Wittenberg T. 2022. Metabolism and autophagy in plants—a perfect match. *FEBS Letters*.

Marbach I, Mayer AM. 1974. Permeability of Seed Coats to Water as Related to Drying Conditions and Metabolism of Phenolics. *Plant Physiology* **54**(6): 817-820.

Marshall RS, Vierstra RD. 2018. Autophagy: The Master of Bulk and Selective Recycling. *Annual Review of Plant Biology* **69**.(1)

Masclaux-Daubresse C, Clement G, Anne P, Rouataboul JM, Guiboileau A, Soulay F, Shirasu K, Yoshimoto K. 2014. Stitching together the Multiple Dimensions of Autophagy

Using Metabolomics and Transcriptomics Reveals Impacts on Metabolism, Development, and Plant Responses to the Environment in Arabidopsis .*Plant Cell* **26**(5): 1857-1877.

McLoughlin F, Augustine RC, Marshall RS, Li F, Kirkpatrick LD, Otegui MS, Vierstra RD. 2018. Maize multi-omics reveal roles for autophagic recycling in proteome remodelling and lipid turnover. *Nat Plants* **4**(12): 1056-1070.

McLoughlin F, Marshall RS, Ding X, Chatt EC, Kirkpatrick LD, Augustine RC, Li F, Otegui MS, Vierstra RD. 2020. Autophagy Plays Prominent Roles in Amino Acid, Nucleotide, and Carbohydrate Metabolism during Fixed-Carbon Starvation in Maize. *The Plant Cell* **32**.2699-2724 :(9)

Minamide LS, Bamburg JR. 1990. A filter paper dye-binding assay for quantitative determination of protein without interference from reducing agents or detergents. *Anal Biochem* **190**(1): 66-70.

Minina EA, Moschou PN, Vetukuri RR, Sanchez-Vera V, Cardoso C, Liu Q, Elander PH, Dalman K, Beganovic M, Yilmaz JL, et al. 2018. Transcriptional stimulation of rate-limiting components of the autophagic pathway improves plant fitness. *J Exp Bot*.

Nguyen T-P, Keizer P, van Eeuwijk F, Smeekens S, Bentsink L. 2012. Natural Variation for Seed Longevity and Seed Dormancy Are Negatively Correlated in Arabidopsis *Plant Physiology* **160**(4): 2083-2092.

Ort DR, Merchant SS, Alric J, Barkan A, Blankenship RE, Bock R, Croce R, Hanson MR, Hibberd JM, Long SP, et al .2015 .Redesigning photosynthesis to sustainably meet global food and bioenergy demand. *Proceedings of the National Academy of Sciences* **112**(28): 8529-8536.

Reumann S, Voitsekhovskaja O, Lillo C. 2010. From signal transduction to autophagy of plant cell organelles: lessons from yeast and mammals and plant-specific features. *Protoplasma* **247**(3-4): 233-256.

Reyes FC, Chung T, Holding D, Jung R, Vierstra R, Otegui MS. 2011. Delivery of prolamins to the protein storage vacuole in maize aleurone cells. *Plant Cell* **23**(2): 769-784.

Rodriguez E, Chevalier J, Olsen J, Ansbøl J, Kapousidou V, Zuo Z, Svenning S, Loefke C, Koemeda S, Drozdowskyj PS, et al. 2020. Autophagy mediates temporary reprogramming and dedifferentiation in plant somatic cells. *The EMBO Journal* **39**.(4)

Ryter SW, Cloonan SM, Choi AM. 2013. Autophagy: a critical regulator of cellular metabolism and homeostasis. *Mol Cells* **36**(1): 7-16.

Salem MA, Juppner J, Bajdzienko K, Giavalisco P. 2016. Protocol: a fast, comprehensive and reproducible one-step extraction method for the rapid preparation of polar and semi-polar metabolites, lipids, proteins, starch and cell wall polymers from a single sample. *Plant Methods* **12**: 45.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* **9**(7): 671-675.

Song J, Xie X, Cui Y, Zou J. 2021. Endosperm-Embryo Communications: Recent Advances and Perspectives. *Plants (Basel)* **10**.(11)

Sreenivasulu N, Wobus U. 2013. Seed-development programs: a systems biology-based comparison between dicots and monocots. *Annu Rev Plant Biol* **64**: 189-217.

Steinbrecher T, Leubner-Metzger G. 2018. Tissue and cellular mechanics of seeds. *Current Opinion in Genetics & Development* **51**: 1-10.

Sueldo DJ, van der Hoorn RAL. 2017. Plant life needs cell death ,but does plant cell death need Cys proteases? *FEBS J* **284**(10): 1577-1585.

Sullivan S, Ralet M-C, Berger A, Diatloff E, Bischoff V, Gonneau M, Marion-Poll A, North HM. 2011. CESA5 Is Required for the Synthesis of Cellulose with a Role in Structuring the Adherent Mucilage of Arabidopsis Seeds *Plant Physiology* **156**(4): 1725-1739.

Tang J, Bassham DC. 2021. Autophagy during drought: function, regulation, and potential application. *The Plant Journal* **109**(2): 390-401.

Towbin H, Staehelin T, Gordon J. 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci U S A* **76**(9): 4350-4354.

Vishwanath SJ, Domergue F, Rowland O. 2014. Seed coat permeability test: tetrazolium penetration assay. *Bio-Protocol* **4**(13): e1173-e1173.

Voiniciuc C, Zimmermann E, Schmidt MH-W, Günl M, Fu L, North HM, Usadel B. 2016. Extensive Natural Variation in Arabidopsis Seed Mucilage Structure. *Frontiers in Plant Science* **7**.

Wada S, Hayashida Y, Izumi M, Kurusu T, Hanamata S, Kanno K, Kojima S, Yamaya T, Kuchitsu K, Makino A, et al. 2015. Autophagy supports biomass production and nitrogen use efficiency at the vegetative stage in rice. *Plant Physiol* **168**(1): 60-73.

Werker E, Marbach I, Mayer AM. 1979. Relation Between the Anatomy of the Testa, Water Permeability and the Presence of Phenolics in the Genus Pisum. *Annals of Botany* **43**(6): 765-771.

Willats WG, McCartney L, Knox JP. 2001. In-situ analysis of pectic polysaccharides in seed mucilage and at the root surface of Arabidopsis thaliana. *Planta* **213**(1): 37-44.

Yoshimoto K, Jikumaru Y, Kamiya Y, Kusano M, Consonni C, Panstruga R, Ohsumi Y, Shirasu K. 2009. Autophagy negatively regulates cell death by controlling NPR1-dependent salicylic acid signaling during senescence and the innate immune response in Arabidopsis. *Plant Cell* **21**(9): 2914-2927.

Yoshimoto K, Shibata M, Kondo M, Oikawa K, Sato M, Toyooka K, Shirasu K, Nishimura M, Ohsumi Y. 2014. Organ-specific quality control of plant peroxisomes is mediated by autophagy. *J Cell Sci* **127**(Pt 6): 1161-1168.

Zhou J, Wang J, Yu JQ, Chen Z. 2014. Role and regulation of autophagy in heat stress responses of tomato plants. *Front Plant Sci* **5**: 174.

Zhou W, Chen F, Luo X, Dai Y, Yang Y, Zheng C, Yang W, Shu K. 2019. A matter of life and death: Molecular, physiological, and environmental regulation of seed longevity. *Plant, Cell & Environment* **43**(2): 293-302.