Title: Autophagy in maternal tissues contributes to *Arabidopsis thaliana* seed development.

Authors: Ori Erlichman[1][**], Shahar Weiss[1], Maria Abu-Arkia[1], Moria Ankary Khaner[1], Yoram Soroka[1], Weronika Jasinska[2], Leah Rosental[2], Yariv Brotman[2] and Tamar Avin-Wittenberg[1][*]

Author Affiliations

[1] Department of Plant and Environmental Sciences, Alexander Silberman Institute of Life Sciences, The Hebrew University of Jerusalem, Givat Ram, 9190401, Jerusalem, Israel

[2] Department of Life Sciences, Ben-Gurion University of the Negev, 8410501, Beer-Sheva, Israel

[*] Corresponding author: tamar.avin-wittenberg@mail.huji.ac.il

[**] Current affiliation: Faculty of Biology, Technion-Israel Institute of Technology, 3200003, Haifa, Israel

Running Head: Autophagy in the mother plant and seed development.

Keywords (5-8, in alphabetical order): *Arabidopsis thaliana,* Autophagy, Maternal tissue, Seed, Source-sink relationship, Storage compounds.

Abstract

Seeds are an essential food source, providing nutrients for germination and early seedling growth. Degradation events in the seed and the mother plant accompany seed development. One degradation mechanism is autophagy, facilitating cellular component breakdown in the lytic organelle. Autophagy influences various aspects of plant physiology, specifically nutrient availability and remobilization, suggesting its involvement in source-sink interactions. During seed development, autophagy was shown to affect nutrient remobilization from mother plants and function in the embryo. Yet, these studies examined autophagy-knockout (*atg* mutant) plants, making it impossible to distinguish between the contribution of autophagy in the source (i.e., the mother plant) and the sink tissue (i.e., the embryo).

To address this, we employed a novel approach to differentiate between autophagy in source and sink tissues. We investigated how autophagy in the maternal tissue affects seed development by performing reciprocal crosses between WT and *atg* mutant *Arabidopsis thaliana* plants. Although F1 seedlings possessed a functional autophagy mechanism, etiolated F1 plants from maternal *atg* mutants displayed reduced growth. This was attributed to altered protein but not lipid accumulation in the seeds, suggesting autophagy differentially regulates carbon and nitrogen remobilization. Surprisingly, F1 seeds of maternal *atg* mutants exhibited faster germination, resulting from different seed coat development.

Our study emphasizes the significance of examining autophagy in a tissue-specific manner, revealing valuable insights into the interplay between different tissues during seed development. It sheds light on the tissue-specific functions of autophagy, offering potential for new research into the underlying mechanisms governing seed development and crop yield.

Introduction

Seeds are a major source of food and feed, providing approximately 70% of the world's human caloric intake (Sreenivasulu and Wobus, 2013). The nutritional value of seeds is attributed to storage compounds accumulating in the seed during development (Cernac and Benning, 2004). These include starch, storage lipids (mostly triacylglycerols - TAGs), and storage proteins, the relative amount of which varies between plant species. Storage compounds are degraded during germination and early seedling development to facilitate germination and seedling growth until photosynthesis establishment (Baud et al., 2008). In annual plants, seed development is accompanied by mother plant senescence, resulting in nutrient remobilization to the developing seeds, used to produce storage compounds (Distelfeld et al., 2014). Another vital seed trait is longevity, defined as seed viability after seed dry storage (Nguyen et al., 2012). Seed longevity can directly affect germination and, as a result, crop productivity and food security (Zhou et al., 2019). With a growing world population (Ort et al., 2015), specifically in light of current climate change (Lobell et al., 2011), an increased understanding of nutrient remobilization to seeds and seed longevity is vital to ensure food security.

Nutrient trafficking and partitioning to various tissues in the plant, i.e. source-sink relationship, are vital for seed development (Sonnewald and Fernie, 2018; Tegeder and Masclaux-Daubresse, 2018). Nutrients from source tissues can be used for organ growth and later exported to sink tissues following catabolism and nutrient recycling (Osorio et al., 2014; Li et al., 2015). Nutrient demand of the sink tissue can also affect export from the source (Lemoine et al., 2013). Plant source-sink relationship has been studied for many years. However, there is still a debate as to which process controls plant growth and yield the most (Sonnewald and Fernie, 2018). During seed development, both carbon, in the form of sucrose, and nitrogen, in the form of the amino acids glutamine and asparagine, are transported from the mother plant (source tissue) to the developing seeds (sink tissue) (Dourmap et al., 2023).

Macroautophagy (hereafter termed "Autophagy") is a conserved eukaryotic mechanism for the degradation of cytoplasmic constituents in the lytic organelle (vacuoles in yeast and plants and lysosomes in animals) (Bassham et al., 2006; Marshall and Vierstra, 2018). The targets of autophagy are diverse and include long-lived proteins, protein complexes, and

whole organelles (Reumann et al., 2010). The autophagy machinery relies on highly-conserved AuTophaGy-related (*ATG*) genes (Bassham, 2009; Avin-Wittenberg et al., 2012; Liu and Bassham, 2012; Ryter et al., 2013), which have been characterized in many plant species (Chung et al., 2009; Kurusu et al., 2014; Zhou et al., 2014). In recent years, autophagy was shown to be involved in many aspects of plant life, from plant development to biotic and abiotic stress response (Avin-Wittenberg et al., 2018).

Autophagy-deficient plants (*atg* mutants) are hypersensitive to carbon and nitrogen starvation and display early senescence and reduced yield (Doelling et al., 2002; Li et al., 2015; Barros et al., 2017), as well as slightly slower germination rates and impaired seedling establishment (Yoshimoto et al., 2014; Avin-Wittenberg et al., 2015). Autophagy was shown to promote nutrient remobilization in plants, functioning in the supply of lipids and primary metabolites (Masclaux-Daubresse et al., 2014; Avin-Wittenberg et al., 2015; Wada et al., 2015; Barros et al., 2017; Hirota et al., 2018; McLoughlin et al., 2018; McLoughlin et al., 2020; Barros et al., 2021). Moreover, overexpression of *ATG* genes in *Arabidopsis thaliana* (Arabidopsis) was reported to increase seed yield and fatty acid content (Minina et al., 2018).

Autophagy was shown to play a dual role in seed development, affecting both the mother plant and the developing embryo. Studies in Arabidopsis and maize (*Zea mays)* demonstrated that impaired autophagy causes altered nitrogen mobilization from the mother plant to the seed, resulting in changes in seed C/N ratio (Guiboileau et al., 2012; Li et al., 2015). Alternatively, microscopic evidence from wheat (*Triticum aestivum*) suggested the possible involvement of autophagy in the transport of storage proteins to protein storage vacuoles within the developing seeds (Levanony et al., 1992). Furthermore, an additional publication demonstrated the presence of an active autophagy mechanism within developing seeds as well as altered storage protein content and processing in Arabidopsis *atg* mutants (Di Berardino et al., 2018).

Though autophagy has been established as a major player in plant nutrient remobilization, the work thus far analyzed knockout plants, in which autophagy is inactive in both source and sink tissues. Indeed, although the studies above suggest autophagy is essential for seed development, it was impossible to distinguish between the impact of autophagy in the mother plant and in the seed itself. Moreover, the seed is comprised of several tissues with

differing parental contributions. The seed coat is of maternal origin, while the embryo and endosperm are zygotic tissues, and the genomic contribution of the triploid single-layer endosperm is unevenly divided between the female parent (2n) and the male parent (1n) (Bentsink and Koornneef, 2008). In this work, we devised an experimental system that allowed us to separate the effect of autophagy in source and sink tissues and elucidate how autophagy in the maternal tissue affects seed development. To that end, we performed reciprocal crosses between Arabidopsis wild type (WT) and *atg* mutants to receive *ATG*-heterozygous F1 seeds with different maternal tissues. We discovered F1 seeds differentially accumulated storage compounds according to the genotype of the mother plant. Surprisingly, we observed that F1 seeds from maternal *atg* mutant origin had a faster germination rate, in striking contrast to homozygous *atg* mutants, which germinate slower than WT seeds. This possibly stems from differences in seed coat structure, which also influences seed longevity. Our work suggests that *atg* mutant seeds display a compound phenotype, merging the impact of autophagy in the mother plant and the embryo.

Results

<u>F1 progeny of WT and *atg* mutants possess a functional autophagy mechanism but reduced growth when germinated in the dark</u>

To differentiate between autophagy in the mother plant and in the embryo, we performed reciprocal crosses between WT and *atg* mutant (*atg5-1* or *atg7-2*) Arabidopsis plants. This novel approach allowed us to produce F1 seeds in which the embryos are heterozygous for either *atg5* or *atg7* mutation while maintaining maternal tissues that are homozygous to either WT, *atg5-1,* or *atg7-2* (Fig. 1) mutants. The heterozygous genotype of F1 plants was validated by PCR (Fig. S1a,b).

We did not observe any developmental difference between the F1 seeds (Fig. 2a). In addition, we did not detect any aborted seeds, as previously reported in *atg5-1* plants (Di Berardino et al., 2018). Homozygous *atg* mutant seeds are smaller than WT seeds (Barros et al., 2017; Minina et al., 2018). We thus compared the weight of reciprocally crossed F1 seeds to examine whether we see a similar effect due to different maternal tissues. We did not see a significant difference between the lines, suggesting this phenotype might be governed by autophagy in the developing embryo (Fig. 2b).

We next wished to determine whether the F1 plants display a functional autophagy mechanism. One hallmark phenotype of *atg* mutants is hypersensitivity to carbon starvation (Marshall and Vierstra, 2018). We thus exposed two-week-old F1 seedlings and their respective paternal lines to carbon starvation and recovery. As can be seen in Fig. 3a-c and Fig. S1c-d, the *atg* mutant parental lines were more susceptible to carbon starvation than WT plants and did not recover, as expected. F1 seedlings displayed a WT-like phenotype, irrespective of the identity of the maternal or paternal lines. *atg* mutant etiolated seedlings were also shown to exhibit shorter hypocotyls when grown without exogenous sugars (Avin-Wittenberg et al., 2015). In light of the results of the carbon starvation experiments, we were expecting not to see a difference between the F1 progeny of our reciprocal crosses. Interestingly, when we performed this assay on the F1 seeds, seedlings stemming from *atg* mutant mother plants had significantly shorter hypocotyls than seedlings from WT mother plants (Fig. 3d,e). As etiolated seedlings are reliant on nutrients from the seeds alone, we hypothesized this phenotype resulted from altered storage compound amounts deposited in the seeds from different mother plants.

atg mutants display an early senescence phenotype (Marshall and Vierstra, 2018), which was shown to stem from an accumulation of salicylic acid during plant aging (Yoshimoto et al., 2009). We thus wished to ascertain whether the observed phenotype did not originate from the premature senescence of the mother plant, resulting in diminished reserve accumulation. To that end, we performed reciprocal crosses between NahG and *atg5*.NahG plants. NahG is a bacterial enzyme that degrades salicylic acid, and its overexpression in an *atg* mutant background inhibits the early senescence phenotype (Yoshimoto et al., 2009). F1 etiolated seedlings from maternal *atg5*.NahG plants displayed shorter hypocotyls than seedlings from maternal NahG plants, similar to F1 seedlings from maternal *atg* mutant origin (Fig. 3f). This result suggests the phenotype observed for F1 seedlings from maternal *atg* mutants is not a result of early senescence.

F1 seeds of maternal *atg* mutant plants display reduced storage protein but no difference in lipid content

We next wished to test our hypothesis of reduced storage compounds in F1 seeds of maternal *atg* mutant origin. As Arabidopsis is an oil seed, we performed lipidomics of dry F1 seeds of reciprocal crosses between WT, *atg5-1,* or *atg7-2*. Though we annotated more

than 120 lipids, we did not observe any separation between the groups in the principal component analysis (PCA – Fig. 4a). A closer look at TAG species revealed no significant differences or apparent trends between F1 seeds of WT and *atg* mutant maternal origin (Fig. 4b). These results are in line with previous works, demonstrating no reduction in the total lipid content of *atg* mutant seeds compared to WT seeds (Minina et al., 2018 1218).

We then examined the total protein content of F1 seeds, as they are mostly composed of storage proteins. We extracted total proteins of similar amounts (2mg) of F1 dry seeds. Surprisingly, though no difference in the lipid content was observed, F1 seeds of maternal autophagy-deficient origin contained lower total protein levels than the reciprocally-crossed seeds. The difference was significant for the *atg7-2* crosses, and a similar trend was observed for the *atg5-1* crosses (Fig. 5a). The results indicate that the protein content, but not the lipid content of the seed, is affected by the mother plant, suggesting a possible differential regulation of carbon and nitrogen allocation by autophagy in the mother plant. To confirm our hypothesis, we analyzed the polar metabolites of dry F1 seeds by GC-MS. We could detect 17 metabolites, yet we could not observe significant differences between the reciprocally-crossed lines, including sucrose (Fig. S2). Unfortunately, we did not detect free amino acids and thus could not determine the amounts of glutamine and asparagine in the F1 seeds.

Previous work indicated that homozygous *atg* mutant seeds display altered storage protein processing (Di Berardino et al., 2018). We, therefore, wanted to examine whether the altered protein amounts in F1 seeds correlated with altered storage protein processing. To that end, we performed Western blot analysis against 12S globulin using total proteins extracted from equal seed quantity. Though we observed a general reduction in protein amounts, we did not observe a change in the storage protein pattern (Fig. 5b). For example, no accumulation of globulin precursor (P12S) was observed in F1 seeds, as previously described for *atg5* mutants (Di Berardino et al., 2018). Our results indicate no difference in protein processing in F1 seeds, suggesting that the altered storage-protein processing in *atg* mutant seeds is due to a lack of autophagic activity in the embryo. This is in line with the normal autophagy phenotype of F1 plants (Fi. 3a-c, Fig. S1c-e). We thus show that the reduced protein quality of *atg* mutant seeds stems from a lack of autophagy in the source tissues, while storage protein processing results from autophagy in the sink tissue.

<u>F1 seeds of maternal *atg* mutant plants display faster germination rates stemming from</u>

<u>altered seed coat morphology</u>

A typical phenotype of *atg* mutants is slightly delayed germination and seedling

establishment (Avin-Wittenberg et al., 2015). We wanted to determine whether this

phenotype stems from autophagy in the mother plant and thus tested germination and

seedling establishment rates. We expected no difference in germination or slightly slower

germination in the F1 seeds from maternal *atg* mutant origin. Surprisingly, increased

germination rates were observed for F1 progeny of *atg* mutant mother plants than WT

mother plants (Fig. 6a). No significant differences were seen in seedling establishment of

F1 seeds (Fig. 6b). We repeated the experiment for F1 seeds of reciprocally-crossed NahG

and *atg5*.NahG plants and received similar results, suggesting the phenotype is unrelated

to early senescence (Fig. S3).

We postulated the higher germination rate of F1 progeny of *atg* mutant mother plants

resulted from faster water penetration into the seed. We thus examined seed permeability

by tetrazolium chloride staining. Tetrazolium salts are amphipathic cations. After

penetrating the dead cells of the seed coat, they are reduced by active dehydrogenases

(NADH-dependent reductases) in the embryo to red-colored insoluble precipitates

composed of formazans (Berridge, 1996). The intensity of red coloration is directly

proportional to the permeability of the seeds. We stained dry F1 seeds for 24 and 48 hours.

As shown in Fig. 6c and d, F1 seeds of maternal *atg5-1* plants displayed significantly higher

water permeability than F1 seeds of maternal WT origin. A similar, albeit insignificant

trend, was observed for the *atg7-2* reciprocal crosses. Interestingly, we also observed

increased water permeability in homozygous *atg* mutant seeds, suggesting this phenotype

is ubiquitous in *atg* mutants (Fig. 6e).

As water permeability is influenced by the seed coat, which is of maternal origin (Marbach

and Mayer, 1974), we took a closer look at the seed coat of F1 seeds and their parental

lines. We first verified that *ATG5* and *ATG7* were expressed in the Arabidopsis seed coat.

A previous study performed detailed gene expression analysis in the seed coast at different

developmental stages (Dean et al., 2011); the data is available at the eFP browser

(https://bar.utoronto.ca/efp_seedcoat/cgi-bin/efpWeb.cgi)(Baxter et al., 2007). Both *ATG5*

and *ATG7* were expressed in the seed coat, starting from 3 days post-anthesis (Fig. S4),

strengthening our hypothesis that autophagy is functional in the seed coat during seed
development.

We then wished to characterize how seed-coat traits differed between WT and *atg* mutant
seeds. We first focused on the hilum, the scar left on the seed coat after detachment from
the funiculus. In mature Arabidopsis seeds, the hilum is adjacent to the micropyle (where
the radicle will emerge) and faces the chalazal pole (Beisson et al., 2007). The hilum
contains high levels of suberin, inhibiting the entrance of water and pathogens. We
compared the suberin levels of homozygous WT and *atg* mutant seeds by autofluorescence.
We did not observe any differences between the lines (Fig. S5a). We next examined the
integrity of the cuticle layer that wraps the embryo (Steinbrecher and Leubner-Metzger,
2018). We stained WT and *atg* mutant seeds with toluidine blue (TB), which stains the
layers underneath the cuticle when it is damaged. As a positive control, seeds were pre-
incubated with EDTA to perforate the coticule. We did not detect any differences between
the lines; none were stained, suggesting their cuticle was intact (Fig. S5b). The seed coat
contains phenols in many species, inhibiting water entrance (Werker et al., 1979). We
stained WT and *atg* mutant seeds with $FeCl_3$ to observe phenols. No genotype displayed
dark spots as expected in the presence of phenols (Fig. S5c).

Hydrated Arabidopsis seeds are coated by a gelatinous layer called mucilage, which is
mainly composed of cell wall polysaccharides (Voiniciuc et al., 2016). It is deposited in
the seed coat epidermal cells during seed development in a structure at the center of the
cell termed "columella". The mucilage is maintained in the epidermal cells following cell
death and is released from the seed coat during imbibition (Griffiths and North, 2017).
Mucilage is rich in pectin, and its architecture can be visualized by ruthenium red (RR)
dye. When F1 seeds were dyed with RR staining, we found that seeds from maternal *atg5-
1* plants displayed lower mucilage content than their reciprocal F1 seeds. We observed a
similar trend for F1 seeds of WT and *atg7-2*, yet not statistically significant (Fig. 7a,b).
Adherent mucilage is partially anchored by cellulosic rays (Sullivan et al., 2011). We
performed calcofluor staining of F1 seeds to visualize mucilage rays. Seeds of maternal
atg5-1 plants displayed higher ray density than their reciprocal F1 seeds. A similar trend
was observed for the *atg7-2* cross (Fig. 7c,d).

Finally, we assessed the general structure of F1 seeds and seed coat cells by scanning electron microscopy (SEM). F1 progeny of maternal *atg* mutants displayed altered seed shape compared to F1 seeds originating from WT mother plants, stemming from reduced seed width (Fig. 8a-top panel,b). We also observed that the seed coat cells of F1 seeds from *atg* mutant mother plants were misshapen (Fig. 8a-bottom panel). The columella of these cells occupied more of the cell area than in seeds from WT mother plants. When we quantified the ratio between the columella and total cell area, we observed a significantly higher ratio for F1 seeds of maternal *atg7-2* plants than WT plants (Fig. 8c). A similar trend was seen in F1 seeds of *atg5-1* and WT reciprocal crosses (Fig. S6). We thus postulate the increased water permeability of maternal *atg* mutants stems from altered columella development, leading to differential mucilage accumulation.

atg mutants exhibit more rapid seed aging compared to WT plants

The seed coat structure, as well as seed permeability, play a vital role in seed longevity and aging (Zhou et al., 2019). We thus hypothesized that *atg* mutant seeds would be more sensitive to seed aging. We performed artificial aging of WT and *atg* mutant seeds and tested their germination ability following aging. Without aging, *atg* mutants displayed slightly delayed germination, as previously described (Avin-Wittenberg et al., 2015). Indeed, all lines were affected by seed aging, showing reduced germination after 10 weeks of artificial aging. As expected, the germination of *atg* mutants was more strongly affected by artificial aging than that of WT seeds. This observation strengthens the link between the altered seed coat of *atg* mutants and seed longevity (Fig. 8d). We also scored seedling development following 5 weeks of artificial aging (Boyes et al., 2001) to assess the longer-term effects of seed aging. Artificial aging resulted in delayed development of all the lines. Yet, *atg* mutants developed slower than WT seedlings, further strengthening our assumption that their increased water permeability results in reduced seed longevity (Fig. S7).

Discussion

Autophagy is gaining attention as a fundamental mechanism in plants, affecting differentiation (Rodriguez et al., 2020), stress response (Tang and Bassham, 2021), and nutrient availability and remobilization (Magen et al., 2022). However, the standard

approach to studying plant autophagy involves using ubiquitous knockout or knockdown mutants. Though applicable in many cases, it may obscure our understanding of the function of autophagy in specific tissues or time points, specifically when examining source-sink relationships. *atg* mutant seeds display a plethora of phenotypes, from reduced storage protein processing (Di Berardino et al., 2018) to altered C/N ratio (Guiboileau et al., 2012). Yet, it was impossible to determine the tissue responsible for the phenotype using the current system.

We implemented a novel approach to address this issue described above, performing reciprocal crosses to separate autophagy in the mother plant and the embryo (Fig. 1). This allowed us to determine the role of autophagy exclusively in the mother plant during seed development. We used two well-studied *atg* mutants, *atg5-1* (Yoshimoto et al., 2009) and *atg7-2* (Hofius et al., 2009), to come to general conclusions regarding autophagy function. Although some phenotypes were significant only for one *atg* mutant line, the other line used always showed a similar trend. This aligns with previous studies, in which the phenotypes of different *atg* mutants were not always at the same degree of severity yet displayed a similar trend (Barros et al., 2017; Minina et al., 2018).

The main challenge of our approach is the relatively small number of seeds we could achieve by reciprocal crosses. We were thus unable to perform certain analyses requiring larger material quantities, such as seed coat composition and metabolic flux analysis. Generating transgenic plants in which autophagy is selectively downregulated either in the embryo or the mother plant will allow conducting experiments on a larger scale as well as investigate both sides of the "source-sink coin" concerning autophagy activity. In addition, a transgenic system could be used to investigate the seed set phenotype of *atg* mutants, which reciprocal crosses can not address. A recent example is a study examining autophagy in root cap cells that used tissue-specific CRISPR to down-regulate autophagy specifically in this tissue (Feng et al., 2022). Nevertheless, we believe our work lays the foundation for the in-depth investigation of autophagy in discrete plant tissues and the relationship between tissues.

We determined that F1 seeds displayed a WT carbon starvation phenotype when exposed to dark treatment at the seedling stage, suggesting the autophagy mechanism is active in F1 seedlings (Fig. 2a-c, S1c-e). In contrast, when we examined etiolated seedlings grown

without exogenous sugars, F1 seeds from maternal *atg* mutants displayed reduced growth

(Fig. 2d,e). These results suggest alerted storage compound accumulation, depending on

the presence of autophagy in the mother plant. We observed no significant difference in

seed lipid content between F1 seeds (Fig. 3). This aligns with previously reported data, in

which homozygous *atg* mutant seeds did not display altered total fatty acid content.

Interestingly, the same paper showed that overexpression of *ATG5* or *ATG7* yielded higher

fatty acid content (Minina et al., 2018). This could stem from higher sink strength in the

mother plant, pushing carbon to the seeds, or increased demand by the embryo.

Unlike lipids, we observed reduced protein levels in F1 seeds of maternal *atg* mutant origin

(Fig. 4a). Previous studies showed that *atg* mutant plants remobilize less nitrogen from the

mother plant to developing seeds (Guiboileau et al., 2012; Li et al., 2015). Together with

our observations, this strengthens our hypothesis that the altered protein amounts in F1

seeds stem from differential remobilization from the mother plant. Surprisingly, carbon

remobilization (depicted by the lipid quantity) is less influenced by autophagy than

nitrogen remobilization (depicted by the protein quantity). Interestingly, we did not observe

any difference in storage protein processing, as described for *atg5* mutant seeds (Di

Berardino et al., 2018), suggesting that autophagy in the embryo functions in storage

protein processing. The involvement of autophagy in storage protein delivery in seeds has

been implied yet hardly investigated, and mostly in monocot, rather than dicot seeds

(Levanony et al., 1992; Reyes et al., 2011). More work is needed to elucidate the role of

autophagy in storage protein processing, and we believe downregulating autophagy

specifically in the embryo will shed light on this fascinating topic.

An important seed tissue we did not address in the scope of this research is the endosperm.

This is a triploid tissue that is partly maternal (2n) and partly paternal (1n) (Bentsink and

Koornneef, 2008). This uneven distribution creates a situation in which F1 seeds of

maternal WT plants possess more copies of *ATG* genes in their endosperm than F1 seeds

of maternal *atg* mutant origin (Fig. 1). We could not directly investigate this issue due to

the seed amount issue described above. We are unaware of studies examining the effect of

ATG gene copy number on plant autophagy activation. However, We did observe that F1

seedlings, containing a lower copy number of *ATG* genes, behaved similarly to WT plants

under carbon starvation (Fig. 2a-c, S1c-e), suggesting that one copy is sufficient for

autophagy activation. Nonetheless, we cannot disregard the contribution of the endosperm to proper embryo growth and development (Song et al., 2021). Future studies are necessary to investigate the function of autophagy in the seed endosperm, perhaps utilizing monocot seeds that have a larger endosperm.

We were most surprised by the observation that F1 seeds of maternal *atg* mutant plants exhibited faster germination compared to F1 seeds from maternal WT plants (Fig. 5a). This was unexpected, as homozygous *atg* mutant seeds display delayed germination compared to WT seeds (Avin-Wittenberg et al., 2015). Of note, the early germination phenotype does not result from the early senescence phenotype of the *atg* mutant mother plant, as F1 seeds of stay-green *atg* mutants displayed a similar phenotype (Fig. S3). We attributed this phenotype to increased water permeability of F1 seeds of maternal *atg* mutant origin (Fig. 5c,d). Interestingly, homozygous *atg* mutant seeds also exhibited increased water permeability, though they are known to germinate slightly slower than WT seeds (Avin-Wittenberg et al., 2015). We postulate that the slower germination of homozygous *atg* mutants is, in fact, a compound phenotype affected by the lack of autophagy in the seed coat and the embryo. Examining the germination of seeds in which autophagy is selectively downregulated in the embryo might help differentiate between the two options.

The reason for the increased water permeability of *atg* mutants is still unclear. Yet, we could observe several differences in the seed coat of F1 seeds of maternal *atg* mutant origin, specifically mucilage accumulation and columella shape (Fig. 6, Fig. 7a-c). More extensive analysis of the seed coat of *atg* mutants is required to fully elucidate the causes of this phenotype. One possibility is the involvement of autophagy in regulating the programmed cell death of the seed coat cells (Sueldo and van der Hoorn, 2017), resulting in deformed seed coat in *atg* mutants and improper deposition of mucilage. Our hypothesis is strengthened by the fact both *ATG5* and *ATG7* are expressed in the Arabidopsis seed coat from early stages of seed development (Fig. S4).

To conclude, our work presents a novel approach to dissecting the function of autophagy in source and sink tissues and examining their effect at the whole plant level. We have shown, for the first time, that the seed phenotype of *atg* mutants is a compound phenotype resulting from the lack of autophagy in both the mother plant and the embryo. Moreover, we implicated autophagy in the mother plant in nitrogen, but not carbon remobilization.

Our work highlights the importance of studying the role of autophagy in higher resolution
while keeping in mind the whole-plant context. We believe that the generation of plant
lines in which autophagy is down-regulated explicitly in space or time will prove to be a
valuable tool to achieve this goal.

Materials and Methods

<u>Plant material and growth conditions</u>
Arabidopsis thaliana ecotype Columbia (Col-0) was used in this study. The lines used were
atg5-1 (SAIL_129B079)(Yoshimoto et al., 2009), *atg7-2* (GK-655B06) (Hofius et al.,
2009), NahG over-expression (NahG) (Yoshimoto et al., 2009) and atg5.NahG (Yoshimoto
et al., 2009). Seeds were surface sterilized with Cl_2 for 2h and sown on Nitsch medium
(Duchefa) plates pH 5.8, with 1% sucrose. The seeds were imbibed at 4°C for 72h in the
dark. Plants were grown in a growth chamber (Fitoclima S600PLH, Aralab) at 22°C and
50% humidity in constant light (125 µmol photons m^{-2} s^{-1}) for 2 weeks. They were then
transferred to pots containing Perlite: Vermiculite: Garden mix (1:1:2 v/v) and grown at
22°C under long-day conditions (8h dark / 16h light) with 150 µmol photons m^{-2} s^{-1}.

<u>Plant crosses and genotype validation</u>
We conducted reciprocal crosses between Col-0 and *atg5-1* or *atg7-2*. Seeds were collected
20 days after the cross. PCR analyses were performed for genotype validation. DNA from
10 plants grown from crossed seeds was extracted for each cross (Dellaporta et al., 1983).
The *atg5-1* crosses were analyzed with *ATG5* forward primer
(ATTCACTTCCTCCTGGTGAAG), and either *ATG5* reverse primer for the WT allele
(TTGTGCCTGCAGGATAAGCG) or T-DNA left border for the mutant allele
(TAGCATCTGAATTTCATAACCAATCTCGATACAC). The *atg7-2* crosses were
analyzed with *ATG7* forward primer (GATTCAATCAACTCGCTAAGGCGT), and either
ATG7 reverse primer for the WT allele (TGCTAATTCCATGGATCCAAC) or T-DNA
left border for the mutant allele (CCCATTTGGACGTGAATGTAGACAC).
Reciprocal crosses between NahG and atg5.NahG were performed as well.

 Embryo development

 Siliques (5 from each genotype) were collected 8 and 12 days after flowering (DAF) and

 opened. The seeds were discolored for 30min in Hoyer's Solution (7.5g gum arabic, 100g

 chloral hydrate, 5ml glycerol, and 30ml water). Discolored seeds (40±5) were then placed

 between a slide and coverslip and observed using a Nomarski objective (20x/0.50 DIC

 M/N2) on an Eclipse 80i microscope (Nikon) coupled to a digital camera.

 Carbon starvation and recovery

 Reciprocally crossed seeds and seeds of the parental lines were sown on Nitsch plates

 without exogenous sugars (24 plants per line per plate). After imbibition, the plants were

 placed in a growth chamber for 2 weeks, covered by aluminum foil for 1 week, and exposed

 to light for an additional week. The plates were photographed before and after carbon

 starvation and following a week of recovery.

 Hypocotyl length measurements

 Reciprocally crossed seeds were sown on Nitsch plates without exogenous sugars.

 Following imbibition, covered plates were transferred to a growth chamber, and the seeds

 were grown vertically in dark conditions for 7 days. The plates were photographed, and

 hypocotyl length was measured by ImageJ (Schneider et al., 2012)(15-40 seedlings per

 line).

 Metabolite extraction for lipidomics and GC-MS analysis

 10 seeds from reciprocally crossed lines were weighed using the MX5 microbalance

 (Mattler Toledo) as previously described (Bromke et al., 2015). Lipids and polar

 metabolites were extracted from 4-8 biological replicates using the MTBE method (Salem

 et al., 2016). The organic phase was dried and used for lipid analysis, and the polar phase

 was dried and used for polar metabolite analysis.

 Lipid analysis by liquid chromatography-mass spectrometry (LC-MS)

 Vacuum-dried organic phases were processed using ultra-performance liquid

 chromatography (on a C8 reverse-phase column) coupled with Fourier transform mass

 spectrometry (Q exactive mass spectrometer; Thermo Fisher Scientific) in positive and

 negative ionization modes. Processing of chromatograms, peak detection, and integration

 were performed using REFINER MSH 10 (GeneData). Mass spectrometry data processing

included removing fragmentation information, isotopic peaks, and chemical noise.

Selected features were annotated using an in-house lipid database (Lapidot-Cohen et al.,

2020). More information can be found in Supplemental Table S1.

<u>Polar metabolite analysis by gas-chromatography-mass spectrometry (GC-MS)</u>

Vacuum-dried polar metabolites were measured by the Agilent 7200B GC/Q-TOF. The

injection and separation procedures were performed according to Dahtt *et al.* using the DB-

35MS column (Dhatt et al., 2019). Metabolite detection and annotation were performed by

Quant software (Agilent), according to an accurate mass library of known markers

generated by our group. More information can be found in Supplemental Table S2.

Following blank subtraction, the peak area of each metabolite was normalized to the

internal standard (i.e., ribitol) and the weight of the sample.

<u>Total protein extraction, quantification, and immunoblotting</u>

Seeds (2mg, three replicates for each line) were ground. Total proteins were extracted using

a protein extraction buffer [50mM pH 7.5 Tris-HCl, 3% SDS, 10% glycerol, 150mM NaCl,

$1 \times$ protease inhibitor (Roche), 20mM DTT], based on a previously described extraction

buffer (Di Berardino et al., 2018). After the addition of buffer and centrifugation for 15min

at 13,000g, supernatants were collected. Samples were incubated in SDS-PAGE sample

buffer (Laemmli, 1970) for 10min at 100°C.

Total protein content was measured based on Minamide and Bamburg method(Minamide

and Bamburg, 1990)with slight modifications. Bovine serum albumin (BSA) powder

dissolved in distilled water to a concentration of 10 mg/ml was used to generate a standard

curve, and a series of dilutions were made (0, 0.156, 0.312, 0.625, 1.25 and 2.5mg/ml). A

grid of 1.5x1.5 cm squares was drawn on a sheet of Whatman number 3 filter paper and

divided into 24 squares. 10µl of the standard curve dilutions and the total protein samples

were applied within the squares and stained in Coomassie brilliant blue (0.25% w/v in

destain solution: 50% methanol, 10% acetic acid, 40% DDW, Raymond A. Lamb) for 10

minutes.

The Whatman paper was destined with destain solution (50% methanol, 10% acetic acid,

40% distilled water) for 10 minutes and washed in water for 2 minutes. Samples were

transferred to a 24-well plate with 200µl DMSO in each well. The plate was mixed gently

for 10 minutes. 100µl of each sample was taken to a new 96-well plate in two technical

replicates, and the absorbance of the standard curve dilutions and samples were obtained

at 600 and 400 nm.

For immunoblotting, total proteins were separated by SDS-PAGE using Tris-Glycine

running buffer (Fling and Gregerson, 1986) and transferred to a polyvinylidene difluoride

membrane (GE Healthcare Life science), as previously described (Towbin et al., 1979).

Equal loading was validated using Coomassie Brilliant Blue staining of the membrane. For

immunodetection, 12S Globulin antibody (Agrisera; AS204403) was used in combination

with horseradish peroxidase-conjugated goat anti-rabbit antibody (Sigma; ab920).

<u>Seed germination assay</u>

Seeds were sown as described above on Nitsch plates without exogenous sugars (five plates

per line, 10-60 seeds from different siliques per plate). After imbibition, the plates were

transferred to a growth chamber, and the number of germinated seeds was counted every

day for 4 days. Germination was scored by radicle protrusion and seedling establishment

by the appearance of two green cotyledons. Germination percentage was calculated

accordingly.

<u>Seed permeability staining</u>

<u>D</u>ry seeds (five replicates of 20 seeds) of each genotype were placed in a 1.5ml

microcentrifuge tube. 1ml of 1% (w/v) aqueous solution of 2,3,5-triphenyltetrazolium

chloride (Sigma) was added to each tube as previously described (Vishwanath et al., 2014).

The tubes were incubated in an air incubator at 30°C in the dark for a period of 24 and 48

hours. After incubation, seeds (10 seeds of each genotype) were observed for changes in

seed color and imaged using a stereomicroscope.

The extraction of formazans was performed as previously described (Vishwanath et al.,

2014). After incubation, as mentioned above, seeds were washed twice with distilled water.

Then 1ml of 95% ethanol was added to the seeds, and the mix was ground using a mortar

and pestle. The ground seed material in ethanol solution was transferred to a

microcentrifuge tube, and the final volume was adjusted to 1ml with 95% ethanol. The

tubes were immediately centrifuged at 15,000xg for 3min. The supernatant was collected

into a spectrophotometer cuvette, and the absorbance of the formazan extracts was

measured at 485nm using a spectrophotometer. As the blank, 95% ethanol solution was

used.

<u>Seed coat visualization</u>

For seed coat visualization, mature seeds were pre-hydrated with water for 90min and then

stained as follows. Ruthenium Red staining for pectins was prepared as previously

described (Willats et al., 2001), and hydrated seeds were incubated in 0.01% (w/v)

Ruthenium Red (Sigma) for 90min at room temperature. Seeds were washed twice in water

and then photographed. Image analysis was performed by ImageJ (Schneider et al., 2012).

Calcofluor staining was conducted as previously described (Willats et al., 2001) using 25µg

ml^{-1} calcofluor (Fluorescent Brightener 28, Sigma) for 20min at room temperature (5 seeds

for each genotype, cesa5 (SALK_118491) as a negative control). Seeds were washed for

2h in water and then photographed. Observations were conducted using the FV-1200

confocal microscope (Olympus), with a 10x/0.4 or 20x/0.75 objective microscope

equipped with 405nm (Excitation) and 430-470nm (Emission) laser diode. Image analysis

was performed by ImageJ (Schneider et al., 2012).

For seed coat autofluorescence, mature seeds of WT, *atg5-1*, and *atg7-2* were illuminated

and observed with an IX81 microscope (Olympus) using DAPI interference and absorption

filter (excitation filter, 350/50nm; emission filter, 460/50nm) coupled to a digital camera

(Li et al., 2007).

For phenol dyeing, green seeds (12-16 DAF) of WT, *atg5-1*, and *atg7-2* were incubated in

$FeCl_3$ solution (2% in 95% EtOH (Mace, 1963)) for 5min. 95% EtOH was used as A

negative control. For cuticle permeability, green seeds (12-16 DAF) of Col-0, *atg5-1*, and

atg7-2 were hydrated for 10min. Hydrated seeds were incubated in an aqueous solution of

0.05% (w/v) Toluidine-Blue (Sigma) for 2 min and then washed with DDW twice (Mace,

1963). Hydration in 0.5% EDTA solution with three DDW washes was used as a positive

control.

<u>Scanning electron microscopy (SEM)</u>

Dry seeds were mounted on aluminum stubs (Electron Microscopy Sciences), sputter-

coated with 10nm of Au/Pd (5% Pd), and viewed using "Quanta 200", field emission

scanning electron microscope (Field Emission Instruments). 6 seeds were examined for

each genotype. Measurement of 15 epidermis cells from each seed was performed by ImageJ (Schneider et al., 2012).

Artificial seed aging

Artificial aging was performed as previously described (Nguyen et al., 2012). 50mg of seeds were stored above a saturated NaCl solution in a closed desiccator (relative humidity of 75%) for 70 days. Germination assay was performed on aged and non-aged seeds as described above. For developmental stage scoring, seeds were artificially aged for 35 days and sown on Nitsch plates without exogenous sugars. The plants were grown as described above for 14 days after imbibition. Seedling development was scored daily as previously described (Boyes et al., 2001).

Statistical analysis

The experiments were conducted in a random blocks design with three to six biological replicates of each genotype. Data were statistically examined using analysis of variance and tested for significant (* = $p<0.05$, ** = $p<0.01$) differences using Student's *t-test* from an algorithm embedded in Microsoft Excel. Heatmaps were generated using the MultiExperiment Viewer (MeV) freely available software application (Howe et al., 2010). Principle Component Analysis was performed using Metaboanalyst 5.0 (Chong et al., 2019). Mixed model analysis was performed using Prism (GraphPad Software).

Acknowledgments

We thank Dr. Smadar Harpaz-Saad of the Hebrew University of Jerusalem for her assistance with Arabidopsis seed coat analysis. We also thank Dr. Naomi Melamed-Book from the microscopy core facility at The Hebrew University's Life Science core facility for her technical support with confocal analysis and Dr. Evgenia Blayvas from the Hebrew University's Unit for Nanoscience, for technical support with SEM.

Author Contributions

O.E. designed the research, performed research, analyzed data, and wrote the manuscript. S.W., M.A-A, M.A., Y.S., W.J, L.R, and Y.B performed research and analyzed data. T.A-

W designed the research and wrote the manuscript. The project was funded by the Israeli

Science Foundation, grant number 1942/19.

Competing interests

The authors declare no competing interests.

Data availability

All data presented in the manuscript is available in the figures or supplementary material.

Figure legends

Fig 1. Schematic representation of the experimental system. Reciprocal crosses

between WT and *atg5-1* or *atg7-2* mutants were performed. The illustration denoted the

genetic makeup of the resulting F1 seeds as well as the mother plants.

Fig. 2: F1 progeny of WT and *atg* mutants display normal development. (a) F1 seeds

were collected at 8DAF and 12DAF, uncolored, and photographed. No aborted embryos

were observed. Representative image of 40±5 seeds examined. Scale bar=100µm. (b)

Weight of 10 F1 seeds. Data are presented as a box&whiskers plot. No significant

differences were observed (n=4-8).

Fig. 3: F1 seedlings display a WT under carbon starvation and recovery, yet

differential growth as etiolated seedlings. (a-c) Representative images of F1 progeny of

reciprocal crosses between WT and *atg5-1* and their respective parental lines. Seeds were

germinated and grown on Nitsch plated without exogenous sucrose for 14 days after

imbibition (a). The plants were incubated in the dark for additional 7 days (b). The plants

were then returned to light for another 7 days to examine their recovery (c). The experiment

was performed in triplicate. (d) Representative image of F1 progeny of reciprocal crosses

between WT and *atg5-1* or *atg7-2*. Seeds were sown on Nitsch plates without exogenous

sugars and grown in the dark for 7 days after imbibition. (e) Hypocotyl length of the lines

described in (d), as quantified by ImageJ. Data are presented as a box&whiskers plot. One

asterisk ($p<0.05$) or two asterisks ($p<0.01$) denote a significant difference following

Student's *t-test* (n=15-30). (f) Hypocotyl length of F1 progeny of reciprocal crosses between NahG and *atg5*.NahG. Seeds were sown on Nitsch plates without exogenous sugars, grown in the dark for 7 days after imbibition, and their hypocotyl length was measured by ImageJ. Two asterisks (p<0.01) denote a significant difference following Student's *t-test* (n=20-30).

Fig. 4: Autophagy deficiency in the mother plant does not affect seed lipid content. Dry seeds of F1 progeny of reciprocal crosses between WT and *atg5-1* or *atg7-2* were collected, and their lipid content was analyzed by UPLC-MS (n=4-8). Detailed results are presented in Supplemental table S1. (a) PCA of lipid levels. (b). Heat map of $\log_2$ of TAG relative levels in comparison to seeds from maternal WT lines. No significant differences between reciprocally crossed lines were identified by student's *t-test*.

Fig. 5: Lack of autophagy in the mother plant alters total seed protein content but not storage-protein processing. Total proteins were extracted from similar amounts of dry seeds of F1 progeny of reciprocal crosses between WT and *atg5-1* or *atg7*-2. (a) Total protein quantification. Data are presented as average ± SE. An asterisk denotes a significant difference between reciprocally crossed lines in Student's *t-test* (p<0.05, n=3). (b) Total proteins were separated by SDS-PAGE. Top panel – Western blot using anti-12S antibody. Bottom panel – Coomassie blue stain. Three biological replicates were performed. P12S – precursors of the 12S protein, α and β – 12S globulin subunits.

Fig. 6: F1 seeds of maternal *atg* mutants exhibit faster germination and increased permeability. (a,b) F1 seeds of reciprocal crosses between WT and *atg* mutants were sown on Nitsch plates without exogenous sucrose, imbibed for 72h, and transferred to continuous light conditions. Germination (defined by radicle protrusion) and seedling establishment (defined by the appearance of two green cotyledons) were scored each day for 4 days. (a) Average percent germination is presented ± SE. (b) Average percent of seedling establishment is presented ± SE. An asterisk denotes a significant difference between reciprocally-crossed lines in Student's *t-test* (p<0.05, n=5). (c) Representative image of F1 seeds from reciprocal crosses after incubation for 24h (top panel) or 48h (bottom panel) in tetrazolium chloride solution. Scale bar=0.5mm. (d, e) Quantification of tetrazolium

chloride staining, Stained seeds were ground and dissolved in 95% EtOH. absorption in

485nm was measured. Average absorption is presented $\pm$ SE for F1 seeds (d) or

homozygous parental lines. One asterisk (p<0.05) or two asterisks (p<0.01) denote a

significant difference between reciprocally-crossed lines following Student's *t-test* (n=5).

Fig. 7: Altered mucilage structure of maternal *atg* mutant F1 seeds. (a,b) F1 seeds of

reciprocal crosses between WT and *atg* mutants were stained with ruthenium red solution.

(a) Representative images of F1 seeds from reciprocal crosses. Scale bar = 200µm. (b)

Quantification of mucilage area. Data are presented as a box&whiskers plot. One asterisk

denotes a significant difference following Student's *t-test* (p<0.05, n=10). (c,d) Calcofluor

staining for β-glucans of F1 seeds. (c) Representative images of whole-seed views (top

panel) and close-ups (bottom panel). Scale bars = 250µm (top panel) and 100µm (bottom

panel). (d) Quantification of ray density. Data are presented as a box&whiskers plot. One

asterisk denotes a significant difference between reciprocally-crossed lines following

Student's *t-test* (p<0.05, n=3-4).

Fig. 8: Autophagy in the mother plants has a negative effect on seed shape and aging.

(a) Scanning electron microscopy (SEM) of F1 seeds produced by reciprocal crosses

between WT and *atg* mutant plants. Top panel, scale bar = 100µm, bottom panel, scale bar

= 50µm. The columella tip is marked in purple, and the cell area is marked in yellow

(bottom panel). (b) Seed dimensions were measured by ImageJ. Data are presented as a

box&whiskers plot. One asterisk denotes a significant difference between reciprocally-

crossed lines following Student's *t-test* (p<0.05, n=6). (c) The ratio between the columella

and total cell area in individual F1 seeds of WT and *atg7-2* reciprocal crosses. Data are

presented as a box&whiskers plot, p-value denotes the statistical difference between the

seeds by mixed model analysis (n=12-15 cells per seed). (d) WT and *atg* mutant seeds

without artificial aging and following 10 weeks of artificial aging were sown on Nitsch

plates without exogenous sucrose, imbibed for 72h, and transferred to continuous light

conditions. Germination (defined by radicle protrusion) was scored each day for 4 days.

Average percent germination is presented $\pm$ SE. One asterisk (p<0.05) or two asterisks

(p<0.01) denote a significant difference from WT at each time point following Student's

t-test (n=5, dark gray - *atg5-1*, light gray – *atg7-2*).

Supporting information

Fig. S1: F1 seedlings of reciprocal crosses are heterozygous and display a WT
phenotype. a,b: PCR analysis for DNA extracted from F1 seedlings (n=5 from different
siliques) and parental lines, with primers for the WT allele and T-DNA insertion. (a) atg5-
1, (b) atg7-2. c-e: Representative images of F1 progeny of reciprocal crosses between WT
and atg7-2 and their respective parental lines. Seeds were germinated and grown on Nitsch
plated without exogenous sucrose for 14 days after imbibition (c). The plants were
incubated in the dark for additional 7 days (d). The plants were then returned to light for
another 7 days to examine their recovery (e). The experiment was performed in triplicate.

Fig. S2: No difference between F1 seeds in primary metabolites. Dry seeds of F1
progeny of reciprocal crosses between WT and *atg5-1* or *atg7-2* were collected, and their
polar metabolite content was analyzed by GC-MS (n=4-8). Detailed results are presented
in Supplemental Data Set S2. (a) PCA of metabolite levels. (b). Heat map of log2 of
metabolite relative levels in comparison to seeds from maternal WT lines. An asterisk
denotes a significant difference between reciprocally crossed lines by Student's *t-test*
($p<0.05$).

Fig. S3: The early germination of maternal *atg* mutant seedlings is not salicylic acid-
dependent. (a,b) F1 seeds of reciprocal crosses between NahG and *atg5*.NahG mutants
were sown on Nitsch plates without exogenous sucrose, imbibed for 72h, and transferred
to continuous light conditions. Germination (defined by radicle protrusion) and seedling
establishment (defined by the appearance of two green cotyledons) were scored each day
for 4 days. (a) Average percent germination is presented ± SE. (b) Average percent of
seedling establishment is presented ± SE. An asterisk denotes a significant difference in
Student's *t-test* ($p<0.05$, n=5).

Fig. S4: *ATG5* and *ATG7* are expressed in the seed coat during seed development.
ATG5 (At5g17290 – left panel) and *ATG7* (At5g45900 – right panel) gene expression in
Arabidopsis Col-2 seed coat as measured by Dean *et al*. (2011). Seed coat was sampled at

3,7 and 11 days post-anthesis (DPA), and RNA was analyzed by microarray analysis. Data

were extracted from the Arabidopsis seed coat eFP browser.

Fig. S5: No difference between WT and *atg* mutant seeds in several biochemical

factors affecting water permeability. (a) Seed coat autofluorescence was visualized using

DAPI filter. All genotypes show saturation in the hilum region (designated by yellow

arrowheads) because of high Suberin content. 30-50 seeds per genotype were examined.

Scale bar=200µm. (b) TB staining for cuticle permeability. No genotype shows blue

staining as the positive control (pre-incubation in 0.5% EDTA solution). 5 siliques were

examined per genotype. Scale bar=0.5mm. (c) $FeCl_3$ staining for phenols. No genotype

shows dark spots predicted to appear in the presence of phenols. Negative control with

DDW. 5 siliques were examined per genotype. Scale bar=0.5mm.

Fig. S6: Autophagy in the mother plant has a negative effect columella/cell area ratio.

The ratio between the columella and total cell area in individual F1 seeds of WT and *atg5-*

1 reciprocal crosses. Data are presented as a box&whiskers plot, p-value denotes the

statistical difference between the seeds by mixed model analysis (n=12-15 cells per seed).

Fig. S7: Artificial aging affects seedling development of *atg* mutants. WT and *atg*

mutant seeds without artificial aging and following 5 weeks of artificial aging were sown

on Nitsch plates without exogenous sucrose, imbibed for 72h, and transferred to continuous

light conditions. Growth stage progression was scored for seedlings grown vertically. Data

represent average ± SE (n35). Asterisk indicates significant differences from WT in each

aging time (Student's t test; $p<0.05$).

Supplemental Table S1: Relative lipid content of reciprocally crossed Arabidopsis

thaliana F1 seeds.

Supplemental Table S2: Relative polar metabolite content of reciprocally crossed

Arabidopsis thaliana F1 seeds.

References

Avin-Wittenberg T, Bajdzienko K, Wittenberg G, Alseekh S, Tohge T, Bock R, Giavalisco P, Fernie AR (2015) Global analysis of the role of autophagy in cellular metabolism and energy homeostasis in Arabidopsis seedlings under carbon starvation. Plant Cell **2**7: 306-322

Avin-Wittenberg T, Baluška F, Bozhkov PV, Elander PH, Fernie AR, Galili G, Hassan A, Hofius D, Isono E, Le Bars R, Masclaux-Daubresse C, Minina EA, Peled-Zehavi H, Coll NS, Sandalio LM, Satiat-Jeunemaitre B, Sirko A, Testillano PS, Batoko H, Hawes C (2018) Autophagy-related approaches for improving nutrient use efficiency and crop yield protection. Journal of Experimental Botany **69:** 1335-1353

Avin-Wittenberg T, Honig A, Galili G (2012) Variations on a theme: plant autophagy in comparison to yeast and mammals. Protoplasma **249:** 285-299

Barros J, Magen S, Lapidot-Cohen T, Rosental L, Brotman Y, Araújo W, Avin-Wittenberg T (2021) Autophagy is required for lipid homeostasis during dark-induced senescence. Plant Physiology **185:** 1542-1558

Barros JAS, Cavalcanti JHF, Medeiros DB, Nunes-Nesi A, Avin-Wittenberg T, Fernie AR, Araújo WL (2017) Autophagy Deficiency Compromises Alternative Pathways of Respiration following Energy Deprivation in Arabidopsis thaliana. Plant Physiology **175:** 62-76

Bassham DC (2009) Function and regulation of macroautophagy in plants. Biochim Biophys Acta **1793:** 1397-1403

Bassham DC, Laporte M, Marty F, Moriyasu Y, Ohsumi Y, Olsen LJ, Yoshimoto K (2006) Autophagy in development and stress responses of plants. Autophagy **2:** 2-11

Baud S, Dubreucq B, Miquel M, Rochat C, Lepiniec L (2008) Storage reserve accumulation in Arabidopsis: metabolic and developmental control of seed filling. Arabidopsis Book **6:** e0113

Baxter I, Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ (2007) An "Electronic Fluorescent Pictograph" Browser for Exploring and Analyzing Large-Scale Biological Data Sets. PLoS ONE **2:** e718

Beisson F, Li Y, Bonaventure G, Pollard M, Ohlrogge JB (2007) The Acyltransferase GPAT5 Is Required for the Synthesis of Suberin in Seed Coat and Root of Arabidopsis. The Plant Cell **19:** 351-368

Bentsink L, Koornneef M (2008) Seed Dormancy and Germination. The Arabidopsis Book **6:** e0119

Berridge MV, Tan, A. S., McCoy, K. D., Wang, R. (1996) The Biochemical and Cellular Basis of Cell Proliferation Assays That Use Tetrazolium Salts. Biochemica **4:** 15-19

Boyes DC, Zayed AM, Ascenzi R, McCaskill AJ, Hoffman NE, Davis KR, Görlach J (2001) Growth Stage–Based Phenotypic Analysis of Arabidopsis. The Plant Cell **13:** 1499-1510

Bromke MA, Hochmuth A, Tohge T, Fernie AR, Giavalisco P, Burgos A, Willmitzer L, Brotman Y (2015) Liquid chromatography high-resolution mass spectrometry for fatty acid profiling. Plant J **81:** 529-536

Cernac A, Benning C (2004) WRINKLED1 encodes an AP2/EREB domain protein
involved in the control of storage compound biosynthesis in Arabidopsis. The Plant
Journal **40:** 575-585
Chong J, Wishart DS, Xia J (2019) Using MetaboAnalyst 4.0 for Comprehensive and
Integrative Metabolomics Data Analysis. Current Protocols in Bioinformatics **68**
Chung T, Suttangkakul A, Vierstra RD (2009) The ATG autophagic conjugation system
in maize: ATG transcripts and abundance of the ATG8-lipid adduct are regulated
by development and nutrient availability. Plant Physiology **149:** 220-234
Dean G, Cao Y, Xiang D, Provart NJ, Ramsay L, Ahad A, White R, Selvaraj G, Datla
R, Haughn G (2011) Analysis of Gene Expression Patterns during Seed Coat
Development in Arabidopsis. Molecular Plant **4:** 1074-1091
Dellaporta SL, Wood J, Hicks JB (1983) A plant DNA minipreparation: Version II. Plant
Molecular Biology Reporter **1:** 19-21
Dhatt BK, Abshire N, Paul P, Hasanthika K, Sandhu J, Zhang Q, Obata T, Walia H
(2019) Metabolic Dynamics of Developing Rice Seeds Under High Night-Time
Temperature Stress. Front Plant Sci **10:** 1443
Di Berardino J, Marmagne A, Berger A, Yoshimoto K, Cueff G, Chardon F,
Masclaux-Daubresse C, Reisdorf-Cren M, Bozhkov P (2018) Autophagy
controls resource allocation and protein storage accumulation in Arabidopsis seeds.
Journal of Experimental Botany **69:** 1403-1414
Distelfeld A, Avni R, Fischer AM (2014) Senescence, nutrient remobilization, and yield
in wheat and barley. Journal of Experimental Botany **65:** 3783-3798
Doelling JH, Walker JM, Friedman EM, Thompson AR, Vierstra RD (2002) The
APG8/12-activating enzyme APG7 is required for proper nutrient recycling and
senescence in Arabidopsis thaliana. J Biol Chem **277:** 33105-33114
Dourmap C, Marmagne A, Lebreton S, Clément G, Guivarc'h A, Savouré A,
Masclaux-Daubresse C, Xu G (2023) Carbon and nitrogen remobilization during
seed filling in Arabidopsis is strongly impaired in thepyrroline-5-carboxylate
dehydrogenasemutant. Journal of Experimental Botany **74:** 1489-1500
Feng Q, De Rycke R, Dagdas Y, Nowack MK (2022) Autophagy promotes programmed
cell death and corpse clearance in specific cell types of the Arabidopsis root cap.
Current Biology **32:** 2110-2119.e2113
Fling SP, Gregerson DS (1986) Peptide and protein molecular weight determination by
electrophoresis using a high-molarity tris buffer system without urea. Anal
Biochem **155:** 83-88
Griffiths JS ,North HM (2017) Sticking to cellulose: exploiting Arabidopsis seed coat
mucilage to understand cellulose biosynthesis and cell wall polysaccharide
interactions. New Phytologist **214:** 959-966
Guiboileau A, Yoshimoto K, Soulay F, Bataille MP, Avice JC, Masclaux-Daubresse
C (2012) Autophagy machinery controls nitrogen remobilization at the whole-plant
level under both limiting and ample nitrate conditions in Arabidopsis. New Phytol
194: 732-740
Hirota T, Izumi M, Wada S, Makino A, Ishida H (2018) Vacuolar Protein Degradation
via Autophagy Provides Substrates to Amino Acid Catabolic Pathways as an
Adaptive Response to Sugar Starvation in Arabidopsis thaliana. Plant and Cell
Physiology **59:** 1363-1376

Hofius D, Schultz-Larsen T, Joensen J, Tsitsigiannis DI, Petersen NH, Mattsson O, Jorgensen LB, Jones JD, Mundy J, Petersen M (2009) Autophagic components contribute to hypersensitive cell death in Arabidopsis. Cell **137**: 773-783

Howe E, Holton K, Nair S, Schlauch D, Sinha R, Quackenbush J (2010) MeV: MultiExperiment Viewer. *In* Biomedical Informatics for Cancer Research, pp 267-277

Kurusu T, Koyano T, Hanamata S, Kubo T, Noguchi Y, Yagi C, Nagata N, Yamamoto T, Ohnishi T, Okazaki Y, Kitahata N, Ando D, Ishikawa M, Wada S, Miyao A, Hirochika H, Shimada H, Makino A, Saito K, Ishida H, Kinoshita T, Kurata N, Kuchitsu K (2014) OsATG7 is required for autophagy-dependent lipid metabolism in rice postmeiotic anther development. Autophagy **10:** 878-888

Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature **227:** 680-685

Lapidot-Cohen T, Rosental L, Brotman Y (2020) Liquid Chromatography-Mass Spectrometry (LC-MS)-Based Analysis for Lipophilic Compound Profiling in Plants. Curr Protoc Plant Biol **5:** e20109

Lemoine R, Camera SL, Atanassova R, Dédaldéchamp F, Allario T, Pourtau N, Bonnemain J-L, Laloi M, Coutos-Thévenot P, Maurousset L, Faucher M, Girousse C, Lemonnier P, Parrilla J, Durand M (2013) Source-to-sink transport of sugar and regulation by environmental factors. Frontiers in Plant Science **4**

Levanony H, Rubin R, Altschuler Y, Galili G (1992) Evidence for a novel route of wheat storage proteins to vacuoles. J Cell Biol **119:** 1117-1128

Li F, Chung T, Pennington JG, Federico ML, Kaeppler HF, Kaeppler SM, Otegui MS, Vierstra RD (2015) Autophagic recycling plays a central role in maize nitrogen remobilization. Plant Cell **27:** 1389-1408

Li Y, Beisson F, Koo AJ, Molina I, Pollard M, Ohlrogge J (2007) Identification of acyltransferases required for cutin biosynthesis and production of cutin with suberin-like monomers. Proc Natl Acad Sci U S A **104:** 18339-18344

Liu Y, Bassham DC (2012) Autophagy: pathways for self-eating in plant cells. Annu Rev Plant Biol **63:** 215-237

Lobell DB, Schlenker W, Costa-Roberts J (2011) Climate Trends and Global Crop Production Since 1980. Science **333:** 616-620

Mace ME (1963) Histochemical Localization of Phenols in Healthy and Diseased Banana Roots. Physiologia Plantarum **16:** 915-925

Magen S, Seybold H, Laloum D, Avin-Wittenberg T (2022) Metabolism and autophagy in plants—a perfect match. FEBS Letters **596:** 2133-2151

Marbach I, Mayer AM (1974) Permeability of Seed Coats to Water as Related to Drying Conditions and Metabolism of Phenolics. Plant Physiology **54:** 817-820

Marshall RS, Vierstra RD (2018) Autophagy: The Master of Bulk and Selective Recycling. Annual Review of Plant Biology **69:** 173-208

Masclaux-Daubresse C, Clement G, Anne P, Routaboul JM, Guiboileau A, Soulay F, Shirasu K, Yoshimoto K (2014) Stitching together the Multiple Dimensions of Autophagy Using Metabolomics and Transcriptomics Reveals Impacts on Metabolism, Development, and Plant Responses to the Environment in Arabidopsis. Plant Cell **26:** 1857-1877

McLoughlin F, Augustine RC, Marshall RS, Li F, Kirkpatrick LD, Otegui MS,
Vierstra RD (2018) Maize multi-omics reveal roles for autophagic recycling in
proteome remodelling and lipid turnover. Nat Plants **4:** 1056-1070
McLoughlin F, Marshall RS, Ding X, Chatt EC, Kirkpatrick LD, Augustine RC, Li
F, Otegui MS, Vierstra RD (2020) Autophagy Plays Prominent Roles in Amino
Acid, Nucleotide, and Carbohydrate Metabolism during Fixed-Carbon Starvation
in Maize. The Plant Cell **32:** 2699-2724
Minamide LS, Bamburg JR (1990) A filter paper dye-binding assay for quantitative
determination of protein without interference from reducing agents or detergents.
Anal Biochem **190:** 66-70
Minina EA, Moschou PN, Vetukuri RR, Sanchez-Vera V, Cardoso C, Liu Q, Elander
PH, Dalman K, Beganovic M, Yilmaz JL, Marmon S, Shabala L, Suarez MF,
Ljung K, Novak O, Shabala S, Stymne S, Hofius D, Bozhkov PV (2018)
Transcriptional stimulation of rate-limiting components of the autophagic pathway
improves plant fitness. J Exp Bot **69:**1415-1432
Nguyen T-P, Keizer P, van Eeuwijk F, Smeekens S, Bentsink L (2012) Natural
Variation for Seed Longevity and Seed Dormancy Are Negatively Correlated in
Arabidopsis   Plant Physiology **160:** 2083-2092
Ort DR, Merchant SS, Alric J, Barkan A, Blankenship RE, Bock R, Croce R, Hanson
MR, Hibberd JM, Long SP, Moore TA, Moroney J, Niyogi KK, Parry MAJ,
Peralta-Yahya PP, Prince RC, Redding KE, Spalding MH, van Wijk KJ,
Vermaas WFJ, von Caemmerer S, Weber APM, Yeates TO, Yuan JS, Zhu XG
(2015) Redesigning photosynthesis to sustainably meet global food and bioenergy
demand. Proceedings of the National Academy of Sciences **112:** 8529-8536
Osorio S, Ruan Y-L ,Fernie AR (2014) An update on source-to-sink carbon partitioning
in tomato. Frontiers in Plant Science **5**
Reumann S, Voitsekhovskaja O, Lillo C (2010) From signal transduction to autophagy
of plant cell organelles: lessons from yeast and mammals and plant-specific
features. Protoplasma **247:** 233-256
Reyes FC, Chung T, Holding D, Jung R, Vierstra R, Otegui MS (2011) Delivery of
prolamins to the protein storage vacuole in maize aleurone cells. Plant Cell **23:** 769-
784
Rodriguez E, Chevalier J, Olsen J, Ansbøl J, Kapousidou V, Zuo Z, Svenning S,
Loefke C, Koemeda S, Drozdowskyj PS, Jez J, Durnberger G, Kuenzl F,
Schutzbier M, Mechtler K, Ebstrup EN, Lolle S, Dagdas Y, Petersen M (2020)
Autophagy mediates temporary reprogramming and dedifferentiation in plant
somatic cells. The EMBO Journal **39**
Ryter SW, Cloonan SM, Choi AM (2013) Autophagy: a critical regulator of cellular
metabolism and homeostasis. Mol Cells **36:** 7-16
Salem MA, Juppner J, Bajdzienko K, Giavalisco P (2016) Protocol: a fast,
comprehensive and reproducible one-step extraction method for the rapid
preparation of polar and semi-polar metabolites, lipids, proteins, starch and cell
wall polymers from a single sample. Plant Methods **12:** 45
Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ 25 :years of
image analysis. Nat Methods **9:** 671-675

Song J, Xie X, Cui Y, Zou J (2021) Endosperm-Embryo Communications: Recent
Advances and Perspectives. Plants (Basel) **10**
Sonnewald U, Fernie AR (2018) Next-generation strategies for understanding and
influencing source–sink relations in crop plants. Current Opinion in Plant Biology
43: 63-70
Sreenivasulu N, Wobus U (2013) Seed-development programs: a systems biology-based
comparison between dicots and monocots. Annu Rev Plant Biol **64:** 189-217
Steinbrecher T, Leubner-Metzger G (2018) Tissue and cellular mechanics of seeds.
Current Opinion in Genetics & Development **51:** 1-10
Sueldo DJ, van der Hoorn RAL (2017) Plant life needs cell death, but does plant cell
death need Cys proteases? FEBS J **284:** 1577-1585
Sullivan S, Ralet M-C, Berger A, Diatloff E, Bischoff V, Gonneau M, Marion-Poll A,
North HM (2011) CESA5 Is Required for the Synthesis of Cellulose with a Role
in Structuring the Adherent Mucilage of Arabidopsis Seeds Plant Physiology **156:**
1725-1739
Tang J, Bassham DC (2021) Autophagy during drought: function, regulation, and
potential application. The Plant Journal **109:** 390-401
Tegeder M, Masclaux-Daubresse C (2018) Source and sink mechanisms of nitrogen
transport and use. New Phytologist **217:** 35-53
Towbin H, Staehelin T, Gordon J (1979) Electrophoretic transfer of proteins from
polyacrylamide gels to nitrocellulose sheets: procedure and some applications. Proc
Natl Acad Sci U S A **76:** 4350-4354
Vishwanath SJ, Domergue F, Rowland O (2014) Seed coat permeability test: tetrazolium
penetration assay. Bio-protocol **4:** e1173-e1173
Voiniciuc C, Zimmermann E, Schmidt MH-W, Günl M, Fu L, North HM, Usadel B
(2016) Extensive Natural Variation in Arabidopsis Seed Mucilage Structure.
Frontiers in Plant Science **7**
Wada S, Hayashida Y, Izumi M, Kurusu T, Hanamata S, Kanno K, Kojima S,
Yamaya T, Kuchitsu K, Makino A, Ishida H (2015) Autophagy supports biomass
production and nitrogen use efficiency at the vegetative stage in rice. Plant Physiol
168: 60-73
Werker E, Marbach I, Mayer AM (1979) Relation Between the Anatomy of the Testa,
Water Permeability and the Presence of Phenolics in the Genus Pisum. Annals of
Botany **43:** 765-771
Willats WG, McCartney L, Knox JP (2001) In-situ analysis of pectic polysaccharides in
seed mucilage and at the root surface of Arabidopsis thaliana. Planta **213:** 37-44
Yoshimoto K, Jikumaru Y, Kamiya Y, Kusano M, Consonni C, Panstruga R, Ohsumi
Y, Shirasu K (2009) Autophagy negatively regulates cell death by controlling
NPR1-dependent salicylic acid signaling during senescence and the innate immune
response in Arabidopsis. Plant Cell **21:** 2914-2927
Yoshimoto K, Shibata M, Kondo M, Oikawa K, Sato M, Toyooka K, Shirasu K,
Nishimura M, Ohsumi Y (2014) Organ-specific quality control of plant
peroxisomes is mediated by autophagy. J Cell Sci **127:** 1161-1168
Zhou J, Wang J, Yu JQ, Chen Z (2014) Role and regulation of autophagy in heat stress
responses of tomato plants. Front Plant Sci **5:** 174

Zhou W, Chen F, Luo X, Dai Y, Yang Y, Zheng C, Yang W, Shu K (20 (19A matter of
life and death: Molecular, physiological, and environmental regulation of seed
longevity. Plant, Cell & Environment **43:** 293-302

Fig 1. Schematic representation of the experimental system. Reciprocal crosses between WT and *atg5-1* or *atg7-2* mutants were performed. The illustration denoted the genetic makeup of the resulting F1 seeds as well as the mother plants.

Fig. 2: F1 progeny of WT and *atg* mutants display normal development. (a) F1 seeds were collected at 8DAF and 12DAF, uncolored, and photographed. No aborted embryos were observed. Representative image of 40±5 seeds examined. Scale bar=100µm. (b) Weight of 10 F1 seeds. Data are presented as a box&whiskers plot. No significant differences were observed (n=4-8).

Fig. 3: F1 seedlings display a WT under carbon starvation and recovery, yet differential growth as etiolated seedlings. (a-c) Representative images of F1 progeny of reciprocal crosses between WT and *atg5-1* and their respective parental lines. Seeds were germinated and grown on Nitsch plated without exogenous sucrose for 14 days after imbibition (a). The plants were incubated in the dark for additional 7 days (b). The plants were then returned to light for another 7 days to examine their recovery (c). The experiment was performed in triplicate. (d) Representative image of F1 progeny of reciprocal crosses between WT and *atg5-1* or *atg7-2*. Seeds were sown on Nitsch plates without exogenous sugars and grown in the dark for 7 days after imbibition. (e) Hypocotyl length of the lines described in (d), as quantified by ImageJ. Data are presented as a box&whiskers plot. One asterisk (p<0.05) or two asterisks (p<0.01) denote a significant difference following Student's *t-test* (n=15-30). (f) Hypocotyl length of F1 progeny of reciprocal crosses between NahG and *atg5*.NahG. Seeds were sown on Nitsch plates without exogenous sugars, grown in the dark for 7 days after imbibition, and their hypocotyl length was measured by ImageJ. Two asterisks (p<0.01) denote a significant difference following Student's *t-test* (n=20-30).

Fig. 4: Autophagy deficiency in the mother plant does not affect seed lipid content. Dry seeds of F1 progeny of reciprocal crosses between WT and *atg5-1* or *atg7*-2 were collected, and their lipid content was analyzed by UPLC-MS (n=4-8). Detailed results are presented in Supplemental table S1. (a) PCA of lipid levels. (b). Heat map of log$_2$ of TAG relative levels in comparison to seeds from maternal WT lines. No significant differences between reciprocally crossed lines were identified by student's *t-test*.

Fig. 5: Lack of autophagy in the mother plant alters total seed protein content but not storage-protein processing. Total proteins were extracted from similar amounts of dry seeds of F1 progeny of reciprocal crosses between WT and *atg5-1* or *atg7*-2. (a) Total protein quantification. Data are presented as average ± SE. An asterisk denotes a significant difference between reciprocally crossed lines in Student's *t-test* (p<0.05, n=3). (b) Total proteins were separated by SDS-PAGE. Top panel – Western blot using anti-12S antibody. Bottom panel – Coomassie blue stain. Three biological replicates were performed. P12S – precursors of the 12S protein, α and β – 12S globulin subunits.

Fig. 6: F1 seeds of maternal *atg* mutants exhibit faster germination and increased permeability. (a,b) F1 seeds of reciprocal crosses between WT and *atg* mutants were sown on Nitsch plates without exogenous sucrose, imbibed for 72h, and transferred to continuous light conditions. Germination (defined by radicle protrusion) and seedling establishment (defined by the appearance of two green cotyledons) were scored each day for 4 days. (a) Average percent germination is presented ± SE. (b) Average percent of seedling establishment is presented ± SE. An asterisk denotes a significant difference between reciprocally-crossed lines in Student's *t-test* (p<0.05, n=5). (c) Representative image of F1 seeds from reciprocal crosses after incubation for 24h (top panel) or 48h (bottom panel) in tetrazolium chloride solution. Scale bar=0.5mm. (d, e) Quantification of tetrazolium chloride staining. Stained seeds were ground and dissolved in 95% EtOH. absorption in 485nm was measured. Average absorption is presented ± SE for F1 seeds (d) or homozygous parental lines. One asterisk (p<0.05) or two asterisks (p<0.01) denote a significant difference between reciprocally-crossed lines following Student's *t-test* (n=5).

Fig. 7: **Altered mucilage structure of maternal *atg* mutant F1 seeds.** (a,b) F1 seeds of reciprocal crosses between WT and *atg* mutants were stained with ruthenium red solution. (a) Representative images of F1 seeds from reciprocal crosses. Scale bar = 200μm. (b) Quantification of mucilage area. Data are presented as a box&whiskers plot. One asterisk denotes a significant difference following Student's *t-test* (p<0.05, n=10). (c,d) Calcofluor staining for β-glucans of F1 seeds. (c) Representative images of whole-seed views (top panel) and close-ups (bottom panel). Scale bars = 250μm (top panel) and 100μm (bottom panel). (d) Quantification of ray density. Data are presented as a box&whiskers plot. One asterisk denotes a significant difference between reciprocally-crossed lines following Student's *t-test* (p<0.05, n=3-4).

Fig. 8: Autophagy in the mother plants has a negative effect on seed shape and aging. (a) Scanning electron microscopy (SEM) of F1 seeds produced by reciprocal crosses between WT and *atg* mutant plants. Top panel, scale bar = 100μm, bottom panel, scale bar = 50μm. The columella tip is marked in purple, and the cell area is marked in yellow (bottom panel). (b) Seed dimensions were measured by ImageJ. Data are presented as a box&whiskers plot. One asterisk denotes a significant difference between reciprocally-crossed lines following Student's *t-test* (p<0.05, n=6). (c) The ratio between the columella and total cell area in individual F1 seeds of WT and *atg7-2* reciprocal crosses. Data are presented as a box&whiskers plot, p-value denotes the statistical difference between the seeds by mixed model analysis (n=12-15 cells per seed). (d) WT and *atg* mutant seeds without artificial aging and following 10 weeks of artificial aging were sown on Nitsch plates without exogenous sucrose, imbibed for 72h, and transferred to continuous light conditions. Germination (defined by radicle protrusion) was scored each day for 4 days. Average percent germination is presented ± SE. One asterisk (p<0.05) or two asterisks (p<0.01) denote a significant difference from WT at each time point following Student's *t-test* (n=5, dark gray - *atg5-1*, light gray − *atg7-2*).