# Transcriptomic responses of sponge holobionts to in situ, seasonal anoxia and hypoxia

Brian W. Strehlow[a*], Astrid Schuster[a*], Warren R. Francis[a], Lisa Eckford-Soper[a], Beate Kraft[a], Rob McAllen[b], Ronni Nielsen[c], Susanne Mandrup[c], Donald E. Canfield[a]

[a] Nordcee, Department of Biology, University of Southern Denmark, Odense M, Denmark.
[b] School of Biological, Earth and Environmental Sciences, University College Cork, Cork, Ireland.
[c] Functional Genomics and Metabolism Research Unit, Department of Biochemistry and Molecular Biology, University of Southern Denmark, Odense, Denmark

*Denotes co-first author. Author order was decided based on reverse alphabetical order, since the inverse was used on a previous paper. Both authors have the right to list their name first on their CVs.

## Abstract

Deoxygenation can be fatal for many marine animals; however, some sponge species are tolerant of hypoxia and anoxia. Indeed, two sponge species, *Eurypon* sp. 2 and *Hymeraphia stellifera,* survive seasonal anoxia for months at a time. To understand their tolerance mechanisms, we performed differential gene expression analyses on the sponges, their mitochondria and their microbial symbionts under *in situ* conditions of normoxia, hypoxia and anoxia. Each species possessed a unique microbiome, but the microbiomes of each species were dominated by a species-specific *Thaumarchaeon* and a *Gammaproteobacterium*. Holobiont gene expression was species and oxygen level dependent, though there were some shared interspecific responses to deoxygenation. In general, few changes occurred in the expression of sponge metabolic genes as a function of oxygenation level, indicating that they may remain metabolically active under anoxia. However, ATP synthesis genes were significantly upregulated under hypoxia when compared to normoxia, and genes for DNA replication were downregulated. Mitochondrial gene expression was effectively unchanged under both hypoxia and anoxia. Nevertheless, both anoxia and hypoxia caused upregulation of heat shock proteins (HSPs), indicating cellular level adaptations to deoxygenation stress. A meta-analysis demonstrated that sponge transcriptional responses to anoxia were distinct from those displayed by other invertebrates while dormant, rejecting the hypothesis of sponge dormancy under anoxia. *Thaumarchaeota* symbionts also upregulated stress response genes in hypoxia, while maintaining expression of oxygen-dependent metabolic pathways under hypoxia and anoxia. *Gammaproteobacteria* symbionts showed relatively few noteworthy changes in gene expression in response to anoxia but decreased metabolic gene expression in hypoxia. There was no clear evidence of upregulated anaerobic respiration in the transcriptomes of the sponge holobionts under anoxia or hypoxia. The tolerance of some sponges to prolonged anoxia warrants further investigation and could give them an advantage in future oceans following climate change as well as in ancient oceans when oxygen concentrations were lower than at present.

**Introduction**

Ocean oxygen levels are predicted to decline by 1–7% over the next century (Keeling, Körtzinger, and Gruber 2010). Climate change is expanding marine oxygen-minimum zones, while enhanced nutrient runoff is causing coastal eutrophication and subsequent deoxygenation (Kirchman 2021). Deoxygenation is expected to negatively affect the physiology, metabolism and survival of many marine organisms (Diaz and Rosenberg 2008). As a result of deoxygenation and other anthropogenic stressors, we are facing large scale habitat and biodiversity losses, but the full effects of these stressors are still unknown.

Sessile organisms that cannot move or migrate in response to deoxygenation, are particularly vulnerable. Sponges (Phylum: Porifera Grant, 1836), sessile, ecologically important filter-feeding animals, are hypothesized to suffer from increasing hypoxic/anoxic areas worldwide (Breitburg et al. 2018; Levin 2018; Stramma et al. 2008), and some *in situ* mortality of sponges (along with other taxa) following deoxygenation has been reported (Altieri et al. 2017; Stachowitsch 1984). However, species specific responses to deoxygenation in Porifera are variable, and some species are tolerant of hypoxia. Lab experiments have shown that the demosponge *Halichondria (Halichondria) panicea* Pallas, 1766 can survive low oxygen levels down to 0.5–4% of air saturation (a.s.) (Mills et al. 2014), and *Tethya wilhelma* can survive down to 0.25–2% a.s. for 4 days and <0.05% a.s. for at least one hour (Mills et al. 2018). Additional demosponge species that tolerate similar levels of hypoxia include *Geodia barretti* Bowerbank, 1858 with 7% a.s. (20 μM $O_2$, (Leys and Kahn 2018)) and *Haliclona pigmentifera* Dendy, 1905 (0.50-0.68% a.s., (Gunda and Janapala 2009)). In a recent experimental study, four taxonomically distinct sponge species (*Cliona celata* Grant, 1826, *Suberites carnosus* Johnston, 1842, *Polymastia crocea* Kelly-Borges and Bergquist, 1997 and *Suberites australiensis* Bergquist, 1968) were exposed to various levels of hypoxia (1.5–40% a.s., 0.13–3.3 mg $O_2$ $L^{-1}$) in aquaria. These sponges exhibited species-specific physiological, morphological and behavioral responses to deoxygenation. Sub-lethal oxygen thresholds for these sponges were in the range of 6–20% a.s. (0.48–1.56 mg $O_2$ $L^{-1}$) while mortality was only observed in one of the two species (*P. crocea*) exposed to severe hypoxia lower than 5% a.s. (0.13 mg $O_2$ $L^{-1}$) (Micaroni, Strano, et al. 2021). The sponge species most sensitive to deoxygenation in controlled experiments, *P. crocea*, had a median lethal time (time until death, LT50) of 286 h at ~1.5% a.s. (0.13 mg $O_2$ $L^{-1}$, (Micaroni, Strano, et al. 2021), which was still longer than most marine organisms (reviewed in Micaroni, Strano, et al. 2021 and Hughes et al. 2020).

Few studies, however, have investigated the deoxygenation tolerance of sponges *in situ*. Bayer et al. (2020) found that the glass sponge *Vazella pourtalesii* Schmidt, 1870 lives in low-oxygen conditions (< 4 mL/L) on the Scotian Shelf, Canada. Schuster et al. (2021) found at least nine encrusting demosponge species of the orders Axinellida Lévi, 1953 and Poecilosclerida Topsent, 1928 that tolerate seasonal anoxia in a Marine Lake in Lough Hyne, Ireland for ~three months at a time, and there are more sponge species at this seasonally anoxic site (Bell and Barnes 2000). The anoxia tolerant sponges of Lough Hyne therefore have LT50s that exceed their ca. three month exposure to anoxia, which are rivaled only by the anemone *Bunodosoma cavernatum* Bosc, 1802 (LT50 in anoxia >1.5 months, (Ellington 1982). Moreover, the abundance (percent cover) of encrusting sponges in the seasonally anoxic site of Lough Hyne

90  only slightly decreased over 30 years despite widespread sponge losses in the lough (Micaroni,
91  McAllen, et al. 2021), indicating that these sponges are uniquely suited to their seasonally
92  anoxic niche.
93
Many invertebrates exposed to harsh conditions like anoxia or hypoxia have evolved molecular
response pathways to survive in these environments. For instance, under anoxia, marine
mollusks shift from lipid, carbohydrate and amino acid metabolism to carbohydrate catabolism
as the primary fuel for ATP generation (Larade and Storey 2009). Some mollusks also maximize
anoxic survival time (mollusc LT50s at 0 mg L$^{-1}$: 360–672 h, (Micaroni, Strano, et al. 2021)) by
building up a fixed reserve of internal fuels (primarily glycogen) to increase ATP output and to
reduce acidity. A small suite of upregulated genes also plays a critical role in cell preservation,
facilitating long-term viability under hypoxia (Larade and Storey 2009). Some anemones may
have elements of a facultative anaerobic metabolism (Sassaman and Mangum 1972); however,
even the anoxia-tolerant *B. cavernatum* requires more oxygen than normal to recover after
exposure to anoxia (Ellington 1982).
105
Only one study has investigated the transcriptomic response of a sponge to hypoxia. After four
days of exposure to hypoxia (0.25–2% a.s.), 128 genes within the sponge *T. wilhelma* were
differentially expressed compared to controls, but none of these genes were clearly related to
metabolic or stress responses (Mills et al. 2018). However, one hour exposure to near anoxia
(<0.05% a.s., 105 nM O$_2$) elicited stress responses in *T. wilhelma*'s transcriptome (Mills et al.
2018), demonstrating that transcriptomic responses were mostly observed under very low
oxygen levels. Nevertheless, the underlying mechanisms of sponge deoxygenation tolerance
and long-term anoxia tolerance are largely unknown.
114
Since parts of the hypoxia-inducible factor (HIF) pathway, which mediates the hypoxic
responses in other animals, are lacking in sponges (Mills et al. 2018), we hypothesized that the
sponge microbiome may play a role in their deoxygenation tolerance. Sponges can host diverse
microbiomes that perform vital functions and can constitute large portions of their biomass
(Wilkinson 1978). The host and microbiome together are referred to as a 'holobiont' (N. S.
Webster and Taylor 2012). Some sponge symbionts have anaerobic capabilities, including
sulfate reduction (Hoffmann et al. 2005), denitrification, and anaerobic ammonium oxidation
(anammox) (Hoffmann et al. 2009). In *V. pourtalesii*, the microbial community is thought to be
specialized for the low-oxygen conditions in which it lives (Bayer et al. 2020).
124
The microbiomes of anoxia-tolerant sponges from Lough Hyne were unchanged under anoxia,
hypoxia and normoxia despite drastic changes in the microbial communities of the water and
sediment, indicating that the holobiont as a whole is tolerant to deoxygenation. Furthermore,
Schuster et al. (2021) hypothesized potential metabolic pathways within the microbiome that
might function under deoxygenation. The microbiomes of the anoxia tolerant sponge species
*Eurypon* sp. 2 and *Hymeraphia stellifera* Bowerbank, 1864 were both dominated by two
microbes – one *Thaumarchaeon* and one *Gammaproteobacterium*, though each sponge hosted
a different species from each clade. *Eurypon* sp. 2 harbors the *Thaumarchean* OTU1 (closest
related isolate: *Nitrosopumilus adriaticus* strain NF5), and *H. stellifera* harbors OTU3 (closest
related isolate: *Nitrosopumilus oxyclinae* strain HCE1). For the *Gammaproteobacteria*, *Eurypon*

sp. 2 contains OTU2 (closest related isolate: *Arenicella chitinivorans* strain ALEN 1); whereas *H.*
*stellifera* contains OTU7 (closest related isolate: *A. chitinivorans* strain ALEN 1,  (Schuster et al.
2021).
138
To understand the mechanisms of sponge holobiont tolerance and survival under extended
anoxia, we performed differential gene expression analysis of the sponges *Eurypon* sp. 2 and *H.*
*stellefera* as well as their respective mitochondria and microbial symbionts including
*Thaumarchaeota* and *Gammaproteobacteria*. Sponges were collected from Lough Hyne under
*in situ* normoxia, hypoxia and anoxia. Since neither of these holobionts are established model
species, we generated reference transcriptomes for the sponges and assembled the genomes
of their symbionts and mitochondria from metagenomic sequences. Additionally, we performed
a meta-analysis of the sponge differential expression data to determine if the responses of
Lough Hyne sponges to deoxygenation are similar to those of *T. wilhelma* under deoxygenation,
other invertebrates under heat stress (Dixon et al. 2015), in a dauer state (Sinha, Sommer, and
Dieterich 2012), and during diapause (Gong et al. 2013). Finally, we considered the implications
of sponge deoxygenation tolerance for future oceans as well as early animal evolution under
low atmospheric oxygen concentrations.
152
153 **Materials and Methods**
154
**Sampling**
Sponges were sampled between July 2018 and August 2019 at Labhra Cliff (Lough Hyne
Nature Reserve, Ireland) as described in Schuster et al. (2021) under permit no. R23-27/2018
issued by the Irish Department of Environment, Heritage, and Local Governments. Samples
were taken under various *in situ* oxygen conditions that were categorized as follows: normoxic
(5.3–12 mg L$^{-1}$, 49.3–111% a.s.), hypoxic (1.30–3.56 mg L$^{-1}$, 12.1–33.1% a.s.), and anoxic
(0.00–0.01 mg L$^{-1}$, 0.00–0.93% a.s., i.e. instrument detection limit). All samples were frozen in
liquid nitrogen within ~15 minutes of collection and kept at liquid nitrogen temperatures in a dry
shipper until returned to the laboratory where samples were stored in a -80°C freezer until
extraction. The sponges were identified using molecular barcodes (see Schuster et al. (2021))
as *Eurypon* sp. 2 and *H. stellifera*.
166
**DNA/RNA extraction**
Total RNA was isolated using TRIzol™ Reagent from Invitrogen (ThermoFisher Scientific,
Denmark). For RNA sequencing (RNAseq), eight samples of *H. stellifera* (anoxic = 2, hypoxic =
5, normoxic = 1) were taken, and sixteen samples of *Eurypon* sp. 2 (anoxic = 2, hypoxic = 7,
normoxic = 7) were collected (n = 24 in total). Since these two species are indistinguishable *in*
*situ*, the samples taken for each species were unequal in number. Remaining DNA in the RNA
extracts were removed using the DNase MAX kit (Qiagen).
174
Additionally, high molecular weight (HMW) DNA was extracted from one individual of each
species using the protocol from Vargas et al. (2021) for metagenomic sequencing. The *Eurypon*
sp. 2 sample used for metagenomic sequencing was taken under anoxia (DC24), and RNA was
also sequenced from this sample. Due to limited availability of samples and DNA, a sample of
*H. stellifera* that was not included in the RNAseq (DC297, sampled under normoxia) was used

for metagenome sequencing. DNA and RNA yields were determined using a Qubit™ (Thermo
Fisher Scientific, USA) and a NanoDrop™ spectrophotometer (Thermo Fisher Scientific, US).
RNA quality was evaluated on an Agilent TapeStation using an RNA ScreenTape (Agilent
Technologies).
184
**Reference transcriptomes and metatranscriptomes**
All RNA samples were sequenced twice with two different library preps - one for polyadenylated
(poly A) messenger RNA (mRNA, from eukaryotes) and one for total RNA (to also capture
prokaryotic RNA). Library preparation for the eukaryotic (sponge) RNA was done using
NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina (New England Biolabs, USA).
For the prokaryotic RNAseq, the supernatant from the poly A enrichment was treated with
NEBNext® rRNA Depletion Kit (Bacteria) and then NEBNext® rRNA Depletion Kit
(Human/Mouse/Rat, New England Biolabs, USA) to eliminate ribosomal RNA (rRNA) that
causes sequence biases. This combination of poly A exclusion and rRNA depletion eliminated
the most rRNA compared to other combinations of methods (or single methods in isolation) in
preliminary tests, yielding sequences with only ~25% rRNA. For the eukaryotic (n = 24) and
prokaryotic (n = 24) RNA samples, library preparation was completed using the NEBNext Ultra II
Directional RNA library preparation kit (New England Biolabs, Denmark). Libraries pooled in
equimolar concentrations were sequenced (5M reads per sample, 2 x 50 bp) at the Center for
Functional Genomics and Tissue Plasticity (ATLAS) at SDU, using a NovaSeq 6000 (Illumina,
USA).
201
To ensure good coverage for the final reference transcriptomes, two samples from each species
were sequenced deeper in additional runs. For *H. stellifera*, the sample for deeper sequencing
(DC129, sampled under normoxia) was also present in the sample set above. In contrast, for
*Eurypon* sp. 2 another sample (DC128, sampled under normoxia) was used due to collection
limitations. As in the sequencing above, two libraries were created and sequenced for each
sample - one with eukaryotic RNA and one with prokaryotic RNA. For this deeper sequencing,
mRNA in the sponge transcriptomes was purified using the NucleoTrap mRNA kit (Macherey-
Nagel, Germany). Prokaryotic RNA was rRNA depleted using the Ribo-zero Magnetic kit
(Illumina, USA). An additional purification step was performed for all RNA samples using the
standard protocol for the CleanNGS kit (CleanNA, Netherlands) before sequencing preparation
using the NEBNext Ultra II Directional RNA library preparation kit (New England Biolabs,
Denmark). Samples were diluted in equimolar concentrations, pooled and sequenced (2 x 150
bp) on a HiSeq X platform (Illumina, USA) by DNASense ApS (Aalborg, Denmark).
215
For the eukaryotic RNA, shallow (2 x 50 bp) sequencing generated $9.20 \times 10^8$ and $4.38 \times 10^8$
reads for *Eurypon* sp. 2 and *H. stellifera*, respectively, across all sampled sponges, and deeper
sequences (2 x 150 bp) generated $2.05 \times 10^8$ and reads $3.45 \times 10^8$ for each sponge of the two
species, respectively. For the prokaryotic RNA, shallow sequencing yielded $1.27 \times 10^9$ and 6.68
$\times 10^8$ reads for *Eurypon* sp. 2 and *H. stellifera*, respectively. The deeper prokaryotic sequencing
generated $1.52 \times 10^8$ and $1.20 \times 10^8$ for *Eurypon* sp. 2 and *H. stellifera*, respectively (totals are
shown in Supplemental Table 1).
223

To obtain reference transcriptomes for each sponge species without prokaryotic symbionts, all
eukaryotic RNA sequences were combined into a single file. These sequences were trimmed for
quality control using public scripts (Matz 2015; Meyer 2016) and assembled with Trinity v 2.8.5
(Grabherr et al. 2011). This assembly was filtered for contaminants and annotated using egg-
NOG mapper (Huerta-Cepas et al. 2019). The final sponge transcriptomes included only genes
with metazoan annotations or no annotations. For a detailed description of all filtering and
assembly steps, see Strehlow et al. (2020) and (https://zenodo.org/badge/latestdoi/573040118)
for all scripts used in the current paper. Sponge reference transcriptomes fit within the expected
range of size, gene number, and N50 (Supplemental Table 1) of published sponge
transcriptomes (Strehlow et al. 2020; Riesgo et al. 2014). Additionally, sponge reference
transcriptomes were effectively complete based on Benchmarking Universal Single-Copy
Orthologs (BUSCO) analysis (Supplemental Table 1, (Simão et al. 2015)) and were therefore
suitable for use for subsequent differential expression analysis (see below). Sponge reference
transcriptome assemblies are available through NCBI under BioProject number PRJNA89319.
238

**Reference Metagenome**

239
240     Two reference metagenomes were sequenced for *Eurypon* sp. 2 and *H. stellifera* at DNASense
241     ApS (Aalborg, Denmark). DNA was fragmented to ~550 bp (M220 Focused-ultrasonicator,
242     Covaris, USA) and used for library preparation following the NEBNext Ultra II DNA library
243     preparation kit. The DNA library was paired-end sequenced (2 x 150 bp) on a NovaSeq platform
244     (Ilumina, USA).
245
Sequences were assembled using Megahit v.1.2.9 (D. Li et al. 2016). The final contig fasta files
were used to plot GC (%) vs. mean coverage of mapped reads to identify individual bins. For
this, we used the custom python scripts spadescontigstocovgc.py (Francis 2020a) as well as the
R script contig_gc_coverage.R (Francis 2020b) to extract bins for the *Thaumarchaeota* and
*Gammaproteobacteria* symbiont of each sponge species. Final statistics for each genome are
shown in Supplemental Table 1. Raw sequences are available from NCBI under BioProject
number PRJNA893197. Metagenome assemblies are available on
https://github.com/bstrehlow/Sponge-transcriptomic-responses-to-seasonal-anoxia.
254

**Mitogenomes**

255
256     Using only the deep sequences of total ('prokaryotic') RNA, the mitochondrial genomes of single
257     individuals of *Eurypon* sp. 2 (DC128) and *H. stellifera* (DC129) were assembled following the
258     pipeline in Plese et al. (2019).
259
For future utilization, single contigs that likely represent the mitochondrial genomes of *Eurypon*
sp. 2 and *Hymeraphia stellifera* were retrieved from the reference metagenome in addition to
the coding regions used for differential expression below. The assembled contigs for each
metagenome were blasted against the mitochondrial genome of DC128 (*Eurypon* sp. 2) using
blastn, with an e-value cutoff of $1 \times 10^{-50}$ for *Eurypon* sp. 2 (DC97) and $1 \times 10^{-30}$ for *Hymeraphia*
*stellifera* (DC297). In general, there was only a single hit for each protein coding gene,
confirming a single contig as the mitochondrial genome for each sample. Based on the blast,
these two contigs contained all protein coding genes retrieved from the transcriptome that were

the focus of this analysis. These 'mitogenomes' can be accessed here:
https://zenodo.org/badge/latestdoi/573040118.
270
**Differential Expression Analysis**
Trimmed reads from the shallow sequencing for each individual were mapped back to their
respective reference transcriptome or genome using bowtie2 (version 2.3.5.1, (Langmead and
Salzberg 2012) and mapped reads were counted for each contig using samcount.pl (see Matz
2015). Eukaryotic reads were mapped to the sponge reference transcriptome for each species.
Prokaryotic reads were mapped to their respective metagenomic bins - metagenome assembled
genomes (MAGs), i.e. *Thaumarchaeota* or *Gammaproteobacteria*, within each sponge species.
The mitochondrial reference was broken into its 14 genes and prokaryotic (total RNA) reads
from both species were mapped to their matching mitochondrial reference and counted as
above.
281
Analyses were performed to compare the gene expression of each target (sponge species,
symbiont species, and mitochondria; n=8) under the three different oxygen conditions (anoxia,
hypoxia and normoxia). Differential gene expression analysis was performed with the package
DESeq2 v. 1.22.2 (Love, Huber, and Anders 2016) in R (R Core Team 2016) with (RStudio
Team 2020). Three pairwise comparisons of oxygen condition (normoxia vs. hypoxia, normoxia
vs. anoxia, and anoxia vs. hypoxia) were computed from DESeq2 results. Significantly
differentially expressed genes between each of these pairwise comparisons were identified
using a false discovery rate (FDR) adjusted p-value threshold of 0.1. The ReportingTools
software package (Huntley et al. 2013) was used to create HTML pages displaying4plots of
filtered expressed genes and their annotations in various oxygen conditions
(https://zenodo.org/badge/latestdoi/573040118). We note that comparisons to *H. stellifera* under
normoxia were made with only one normoxic sample due to sampling restrictions. Despite this
sampling limitation, the normoxic data for *H. stellifera* are still shown as a baseline and
reference. For each sponge species, log fold changes (LFC, from DESeq2) in all genes were
calculated so that positive LFCs were upregulated and negative values were downregulated in
hypoxia or anoxia relative to normoxia. For the comparisons between hypoxia and anoxia,
positive LFCs were upregulated in anoxia and vice versa.
299
300
**KOG analysis**
EuKaryotic Orthologous Groups (KOG class) assignments were used to compare large scale
trends and potential correlations in differential gene expression between Lough Hyne sponges
under varied oxygen conditions and external gene expression data.
Delta-ranks, a measure of each KOG class enrichment with up or downregulated genes, were
calculated from LFC data using the KOGMWU package in R (Dixon et al. 2015), and delta-ranks
were plotted using the package pheatmap (Kolde 2013). While it is also possible to calculate
delta-ranks with signed p-values (see KOGMWU supporting documentation), LFC was used
here to include changes between normoxia and other conditions for *H. stellifera*. Since there
was only one normoxic replicant for this species, LFC was a more appropriate metric than
signed p values. Within each pairwise comparison (i.e. hypoxic, anoxic, and anoxic vs hypoxic),
significantly enriched KOGs were also calculated using KOGMWU (FDR-adjusted p < 0.05).

These KOG enrichments were also compared to those calculated from the sponge *Tethya wilhelma* after four days of exposure to hypoxia and one hour of exposure to anoxia (Mills et al. 2018). To test whether KOG enrichment in sponges exposed to variable oxygen conditions was correlated with gene expression in other invertebrate stress responses, delta rank data was included from adult *Acropora millepora* coral exposed to heat stress (31.5 °C for three days, (Dixon et al. 2015). Similarly, to test the similarity of the sponge data to noted decreases in metabolic activity in invertebrates, KOGs enriched in the dauer (dormant) state in *Caenorhabditis elegans* (Sinha, Sommer, and Dieterich 2012) and the diapause state of the midge *Sitodiplosis mosellana* (Gong et al. 2013) were added using delta rank tables calculated by Strader et al. (2016). Correlations between the KOG delta ranks in different datasets were performed using Pearson's Correlation in KOGMWU. Within the significantly enriched KOGs 'replication, recombination and repair' and 'energy production and conversion,' genes that were significantly differentially expressed genes were plotted in heatmaps to further investigate sponge transcriptomic responses to deoxygenation at the gene level in Lough Hyne sponges.

KOG enrichment analysis was performed as above for each of the two major symbionts within the two sponge species; however, Clusters of Orthologous groups (COGs) were used since they are designed for prokaryotes rather than eukaryotic KOGs and are fewer in number.

**Gene ontology (GO) term enrichment analysis**
Log fold changes calculated above were used for rank-based gene ontology (GO) analysis with adaptive clustering (GO_MWU, (Wright et al. 2015)) to determine what GO terms were significantly enriched under various oxygen conditions. These analyses were run individually for each pairwise comparison of oxygen condition in both sponge species. These analyses were not completed for the two symbiont types because GO terms were not available for these microbes.

**Phylogenetic positions of nitroreductase proteins**
Nitroreductase proteins from the *Thaumarchaeota* symbionts of *H. stellifera* and *E.* sp. 2 were extracted using annotations assigned by egg-NOG. Phylogenetic position of nitroreductase proteins from *H. stellifera* and *E.* sp. 2 were investigated by aligning them to other known nitroreductase proteins from Genbank using Mafft v7 (Katoh and Standley 2013). The ModelTest-NG v.0.1.7 function (Edler et al. 2020) implemented within RaxmlGUI 2.0 was used to calculate the evolutionary model. According to AIC, LG+I+G was chosen as the best-fitting model. Maximum Likelihood (ML) phylogeny was then reconstructed using raxmlGUI v.2.0.6 (Edler et al. 2020) with 100 (rapid) bootstraps. The resulting protein tree was rendered by FigTree v.1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

**Results**

**Sponge differential expression**
There were three potential pairwise comparisons for significantly differentially expressed genes - hypoxia vs. normoxia, anoxia vs. normoxia, and anoxia vs. hypoxia. The number of significantly differentially expressed genes in these comparisons for sponges, their mitochondria

and their symbionts are summarized in Table 1. In the sponge *Eurypon* sp. 2, 737, 1,376 and
1,090 genes were significantly differentially expressed under hypoxia (vs. normoxia), anoxia (vs.
normoxia), and anoxia vs. hypoxia, respectively ($p < 0.1$, Table 1). The differentially expressed
genes under each oxygen condition ranged between 1.5–2.8% of the genes of the *Eurypon* sp.
2 reference transcriptome, and 41-48% of these genes are annotated, which reflects the overall
annotation percentage of 39.7% for the whole transcriptome (Supplemental Table 1). In
hypoxia, the majority of these differentially expressed genes were upregulated (68%). Under
anoxia, on the other hand, the majority of genes were downregulated (60%). In anoxia vs.
hypoxia, 57% of genes were upregulated in anoxia compared to hypoxia in *Eurypon* sp. 2
(Table 1).

**Table 1.** Number of significantly differentially expressed genes (DEGs) under deoxygenation in
the sponge holobionts summarized per reference genome or transcriptome. The direction (up or
down) within each paired oxygen condition comparison is relative to the lower oxygen condition.
The number of annotated genes is shown in parentheses.

| | | | Hypoxia | | Anoxia | | Anoxia vs hypoxia | |
|---|---|---|---|---|---|---|---|---|
| **Sponge** | **Reference** | **Total genes in Reference** | Up | Down | Up | Down | Up | Down |
| *Eurypon* sp. 2 | sponge | 49,911 | 499 (239) | 238 (112) | 551 (228) | 825 (340) | 622 (273) | 468 (205) |
| *H. stellifera* | sponge | 62,494 | 465 (125) | 58 (9) | 40 (11) | 244 (87) | 1531 (438) | 623 (286) |
| *Eurypon* sp. 2 | mitochondria | 14 | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 1 (1) | 0 (0) |
| *H. stellifera* | mitochondria | 14 | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| *Eurypon* sp. 2 | *Thaumarchaeota* | 1,556 | 19 (14) | 1 (1) | 3 (3) | 0 (0) | 0 (0) | 0 (0) |
| *H. stellifera* | *Thaumarchaeota* | 1,551 | 17 (12) | 0 (0) | 0 (0) | 0 (0) | 1 (0) | 27 (20) |
| *Eurypon* sp. 2 | *Gammaproteobacteria* | 2,751 | 3 (3) | 29 (29) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| *H. stellifera* | *Gammaproteobacteria* | 3,057 | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 20 (20) | 416 (401) |

In the sponge *H. stellifera*, 523, 284, and 2,154 genes were significantly differentially expressed
in hypoxia, anoxia, and in anoxia vs. hypoxia, respectively ($p < 0.1$, Table 1). Differentially
expressed genes in the different oxygen conditions corresponded to 0.42 - 3.4% of the total

number of genes in the *H. stellifera* transcriptome. Approximately 25-37% of the differentially
expressed genes were annotated, which, as in the other sponge species, roughly corresponded
to the global percentage (37.0%) of genes annotated within the *H. stellifera* reference
transcriptome. Under hypoxia in this species, the vast majority of differentially expressed genes
(89%) were upregulated. As in *Eurypon* sp. 2, the majority of differentially expressed genes
(84%) were downregulated in anoxia in *H. stellifera*. The majority of differentially expressed
genes between anoxia and hypoxia were upregulated in anoxia (71%, Table 1).
386
387  <u>KOG enrichments in sponges</u>
388
The global gene expression patterns of Lough Hyne sponges in response to deoxygenation
were compared across the two species by assessing up and down regulation of KOG classes
(broad functional annotations assigned to genes). Within these sponges, KOG expression in
response to hypoxia and anoxia (versus normoxia) were largely distinct in both species (Figure
1A). However, KOG expression in *H. stellifera* in hypoxia vs. anoxia was significantly positively
correlated, i.e. similar, to that of *Eurypon* sp. 2 under hypoxia ($r = 0.43$, $p < 0.05$, Figure 1J), but
global expression patterns were only similar between the two species in this single case.
396
397
398
399

**Figure 1.** KOG expression profiles of sponges under varied oxygen concentrations compared to each other and external datasets. A. Clustered heatmap (based on correlation) of KOG class (rows) enrichments for upregulated or downregulated genes. Highlighted columns correspond to data generated in this study (Purple: *Hymeraphia stellifera* [Hs], Cyan: *Eurypon* sp. 2 [Es]). Up or down regulation with respect to normoxia is indicated in 'Hypox' (hypoxia) and 'Anox' (anoxia) columns, and differential regulation with respect to hypoxia is shown in 'Hypox vs. Anox' columns, i.e. upregulated KOGs (red delta ranks) are upregulated in anoxia with respect to hypoxia. Within each column on the heatmap, KOGs that are significantly enriched (FDR-adjusted $p < .05$) are marked with asterisks. KOG enrichments from the sponge *Tethya wilhelma* after four days exposure to hypoxia (Tw Hypox Long) and one hour exposure to anoxia (Tw Anox Short), the coral *Acropora millepora* after three days of exposure to heat stress (Am Coral Heat), *Caenorhabditis elegans* in a dauer state (Ce Dauer), and the midge *Sitodiplosis mosellana* in diapause (Sm Diapause) from external studies were included for comparative purposes. B-J. Significant ($p < 0.05$) correlations between the KOG enrichments of different treatments or datasets (columns in A). All other possible comparisons between the various datasets were not significant.

Within *Eurypon* sp. 2, KOG class expression patterns in hypoxia vs. anoxia were significantly positively correlated to those of hypoxia ($r = 0.71$, $p < 0.01$, Figure 1D). Similarly, in *H. stellifera*, KOG enrichment in hypoxia vs. anoxia was positively correlated to that of hypoxia ($r = 0.72$, $p < 0.001$, Figure 1H) and anoxia ($r = 0.46$, $p < 0.05$, Figure 1I)*. All significant correlations between datasets are shown in (Figure 1B-J), and all relationships that were not significantly correlated are not shown.

At the level of individual KOGs, the KOG class 'replication recombination and repair' (RRR) was significantly enriched in all pairwise comparisons of oxygen conditions for both sponge species ($P_{adj} < 0.05$), though in different directions depending on oxygen. It was significantly upregulated in anoxia and anoxia vs. hypoxia but downregulated in hypoxia in both species (Figure 1A).

In *Eurypon* sp. 2,five genes with RRR KOG annotations were significantly upregulated in hypoxia, e.g. HELZ and HELZ2 transcription associated helicases, and 11 genes were significantly downregulated ($p < 0.1$, Supplemental Figure 4A), including five genes from the minichromosome maintenance (MCM) protein complex. In anoxia, two genes were significantly upregulated within RRR in this species - 'siz regulation of ARF protein signal transduction' and DNA polymerase epsilon catalytic subunit ($p < 0.1$, Supplemental Figure 4B). The majority of significantly differentially expressed RRR genes (five) were downregulated in anoxia ($p < 0.1$, Supplemental Figure 4B). The majority of RRR genes (nine) were also downregulated in anoxia vs. hypoxia, with only two significantly upregulated in anoxia (Supplemental Figure 4B). The genes NBN and RFC1 were downregulated in anoxia compared to both normoxia and hypoxia (Supplemental Figure 4B,C). Similarly, the gene 'siz regulation of ARF protein signal transduction' was upregulated in anoxia compared to both normoxia and hypoxia (Supplemental Figure 4B,C).

In *H. stellifera*, three genes were significantly differentially expressed in the RRR KOG class in hypoxia, and all were upregulated ($p < 0.1$, Supplemental Figure 4D). No genes within this class

were significantly differentially expressed in anoxia (vs. normoxia). Eleven genes were
upregulated and eight genes were downregulated in anoxia vs. hypoxia (Supplemental Figure
4E). Similar to *Eurypon* sp. 2, HELZ2 was downregulated in anoxia vs. hypoxia in *H. stellifera*.
In contrast to *Eurypon* sp. 2 a distinct MCM gene (MCM6) was downregulated in anoxia vs.
hypoxia. Additionally, four genes from the structural maintenance of chromosomes protein 6
(SMC6) complex were downregulated in anoxia vs. hypoxia.
452
453 Like RRR, the KOG class 'energy production and conversion' (EPC) was significantly enriched
454 ($P_{adj}$ < 0.05) between all conditions except for anoxia vs. hypoxia in *H. stellifera* ($P_{adj}$ = 0.07).
455 EPC was significantly upregulated in hypoxia in *H. stellifera* but downregulated in all remaining
456 conditions in both sponge species (Figure 1A).
457
Within *Eurypon* sp. 2 specifically, four genes were significantly downregulated and twelve genes
were upregulated within the EPC class in hypoxia (p < 0.1, Supplemental Figure 5A). Various
nuclear genes for processes in the mitochondria were differentially expressed in hypoxia.
SLC25A27 (a mitochondrial membrane uncoupling protein), ndufs8 (a NADH dehydrogenase
core subunit in the mitochondrial membrane - complex I), and Surf1 (part of the complex that
regulates cytochrome c oxidase assembly) were downregulated in hypoxia (Supplemental
Figure 5A). However, other nuclear mitochondrial genes were upregulated, including four
subunits of ATP synthase and two additional genes with annotations corresponding to ATP
synthesis (Supplemental Figure 5A). In anoxia, fourteen genes were upregulated and five genes
were downregulated in the EPC class (Supplemental Figure 5B).
468
As in hypoxia, ndufs8 was downregulated, but no other significant DEGs were shared across
the two oxygen levels. The following genes were also downregulated in anoxia: mitochondrial
sulfite oxidase (SUOX), malate dehydrogenase (MDH1), and two genes belonging to the ATP
gaunido phosphatase family (Supplemental Figure 5B). Conversely, four genes involved in
phosphoenolpyruvate carboxykinase activity were significantly upregulated in anoxia (p < 0.1,
Supplemental Figure 5B). Additionally, three mitochondrial membrane transporter proteins from
the SLC family, two CYB5R4 cytochrome reductase genes, a mitochondrial carrier protein, and
fdx1 cytochrome reductase were upregulated in anoxia. In anoxia vs. hypoxia in *Eurypon* sp. 2,
ten genes were upregulated in anoxia (Supplemental Figure 5C). Upregulated genes included
all of the same genes that were significantly upregulated in anoxia, except for the
phosphoenolpyruvate carboxykinase associated genes. Six genes were significantly
downregulated in anoxia vs. hypoxia. Of these, three were also downregulated in anoxia (vs.
normoxia) and two genes were significantly upregulated in anoxia (Supplemental Figure 5A-C).
It is noteworthy, however, that all of the upregulated genes in anoxia vs. normoxia and hypoxia
were only strongly upregulated in one of the two anoxia replicates (DC56, Supplemental Figure
5 B,C).
485
In *H. stellifera*, eight genes were upregulated and two genes were downregulated in hypoxia. As
in *Eurypon* sp. 2, the ATP5A1 ATP synthase subunit was upregulated in hypoxia. An additional
ATP synthase subunit (ATP6V0C) that was not significantly upregulated in *Eurypon* sp. 2 was
upregulated in *H. stellifera* in hypoxia (p < 0.1, Supplemental Figure 5D). Mitochondrial
sulfide:quinone oxidoreductase (SQRDL), three genes associated with phosphoenolpyruvate

carboxykinase, and a histone demethylase (KDM6A) were also significantly upregulated in
hypoxia in *H. stellifera*. Conversely, PCK1 (also associated with phosphoenolpyruvate
carboxykinase) and MDH1 were significantly downregulated in hypoxia ($p < 0.1$, Supplemental
Figure 5D). In anoxia in *H. stellifera*, ATP5A1 was significantly upregulated, and ATP6V0C,
PCK1, and MDH1 were significantly downregulated (Supplemental Figure 5E). Comparing
anoxia vs. hypoxia, eight genes were upregulated and five genes were downregulated
(Supplemental Figure 5F).

As in hypoxia, SQRDL, four genes associated with phosphoenolpyruvate carboxykinase activity,
and KDM6A were upregulated in anoxia vs. hypoxia. Additionally, two unshared genes,
including the mitochondrial Succinyl-CoA:3-ketoacid coenzyme A transferase 1 (OXCT1) and a
mitochondrial carrier protein, were upregulated in anoxia vs. hypoxia (Supplemental Figure 5F).
Five genes were downregulated in anoxia vs. hypoxia in *H. stellifera*. Two of these four only
exhibited high expression levels in one hypoxic sample (DC105) that drove the observed
difference (Supplemental Figure 5F). The other three downregulated genes in anoxia vs.
hypoxia were mitochondrial creatine kinase (CKMT2),  NAD P transhydrogenase (NNT), and
proto-oncogene tyrosine-protein kinase (Src).

The 'cytoskeleton' KOG class was also significantly enriched ($P_{adj} < 0.01$) in all but one oxygen
condition – anoxia in *Eurypon* sp. 2 ($P_{adj} = 0.1$). Interestingly, enrichment patterns in this KOG
were species specific, being significantly upregulated in hypoxia and anoxia vs. hypoxia in
*Eurypon* sp. 2 and downregulated in all conditions for *H. stellifera*.

The other twenty KOG classes were only significantly enriched in Lough Hyne sponges under a
few (or no) different oxygen conditions and patterns of significant enrichment were generally
condition and species-specific (Figure 1A). With the exception of EPC, KOGs pertaining to
metabolic processes were rarely significantly enriched. Expression of the KOG classes
'carbohydrate metabolism' and  'amino acid transport and metabolism' were not significantly
changed under any oxygen condition ($P_{adj} > 0.05$). There was some significant enrichment of
metabolic KOGs, but only in *Eurypon* sp. 2. 'Coenzyme transport and metabolism' and
'inorganic ion transport and metabolism' were both downregulated in anoxia, and 'nucleic acid
transport metabolism' was downregulated in hypoxia vs. anoxia. For both these conditions
(anoxia and hypoxia vs. anoxia), 'Lipid transport and metabolism' and 'Secondary metabolites
biosynthesis, transport and catabolism' were significantly downregulated. However, the
transcription of the majority of genes labeled with metabolic KOGs remained unchanged under
anoxia and hypoxia.

Comparing the Lough Hyne dataset with external datasets of invertebrate KOG expression in
response to stress, only two positive correlations were found. In *Eurypon* sp. 2, expression
patterns under Anoxia vs. Hypoxia were comparable to those of *T. wilhelma* under short-term
anoxia ($r = 0.44$, $p < 0.05$, Figure 1E). The hypoxic response of *Eurypon* sp. 2 resembled that of
diapause in *S. mosellana* ($r = 0.68$, $p < 0.001$, Figure 1B), but this was the only correlation found
between Lough Hyne sponge data and diapause or dauer states. Importantly, KOG enrichments
under anoxia (versus normoxia) were not positively correlated to any other external datasets for
either sponge species.

Three significant negative correlations, which corresponded to opposing KOG expression
patterns, were observed between Lough Hyne sponges and external datasets. Hypoxia
responses in *H. stellifera* were negatively correlated with those of *A. millepora* under heat stress
($r$ = -0.56, $p$ < 0.01, Figure 1C). The long exposure of *T. wilhelma* to hypoxia was negatively
correlated to both anoxia ($r$ = -0.64, $p$ < 0.01, Figure 1G) and hypoxia vs. anoxia ($r$ = -0.61, $p$ <
0.01, Figure 1F) in *Eurypon* sp. 2.

<u>GO term enrichments in sponges</u>

Generally, significantly enriched GO terms between the different treatments were not shared
between the two sponge species (Supplemental Figures 1-3). The strongest upregulated GO
enrichments ($p$ < 0.01) in hypoxia in *Eurypon* sp. 2 were cellular signaling and 'lipoprotein
particle receptor binding' (Supplemental Figure 1A). For *H. stellifera*, on the other hand, the
strongest upregulated enrichments ($p$ < 0.01) in hypoxia included the terms 'protein tyrosine
kinase binding,' 'molecular adaptor,' 'structural molecule,' and 'structural constituent of
ribosome'; whereas the downregulation of the term 'motor' was strong ($p$ < 0.01, Supplemental
Figure 1B).

Under anoxia, strong upregulation of the 'proteasome binding' term was observed in *Eurypon*
sp. 2 ($p$ < 0.01), and the terms 'phosphoenolpyruvate carboxykinase', 'carbohydrate kinase,'
monovalent inorganic cation transmembrane transporter,' and 'sodium ion transmembrane
transporter' were highly downregulated ($p$ < 0.01, Supplemental Figure 2A). In *H. stellifera*
under anoxia, the terms 'damaged DNA binding' and 'calcium-dependent phospholipid binding'
were highly upregulated, and four terms associated with various aspects of the cytoskeleton
were downregulated ($p$ < 0.01, Supplemental Figure 2B).

In *Eurypon* sp. 2 under anoxia vs. hypoxia, the terms 'lipoprotein receptor binding' and
'proteasome binding' were strongly upregulated in anoxia, but no terms were strongly
downregulated in anoxia versus hypoxia ($p$ < 0.01, Supplemental Figure 3A). The former term
was also upregulated under anoxia vs. normoxia. Conversely, the strongest enrichments in *H.
stellifera* in anoxia vs. hypoxia were downregulated in anoxia and were composed of three of
the four cytoskeletal binding GO terms that were also downregulated in anoxia vs. normoxia
(see above, $p$ < 0.01, Supplemental Figure 3B), but no strong upregulation of GO terms was
observed.

Considering GO enrichments at lower significance levels (0.01 < $p$ < 0.1), within *Eurypon* sp. 2,
the GO terms 'proteasome binding', 'lipoprotein particle receptor binding,' and 'Hsp70 binding'
(HSP: Heat shock protein) were significantly upregulated in all cases of decreased oxygen albeit
at varied significance levels ($p$ < 0.05, Supplemental Figures 1–3A). Between only two levels,
the terms 'transmembrane signaling receptor,' 'secondary active transmembrane transporter,'
and 'molecular transducer' were upregulated in hypoxia but downregulated in anoxia. However,
the term lipoprotein particle receptor binding was downregulated in anoxia and upregulated in
hypoxia ($p$ < 0.1, Supplemental Figures 1,2A). Additionally, the term 'structural constituent of
ribosome' was upregulated in anoxia but downregulated in anoxia vs. hypoxia ($p$ < 0.1,

Supplemental Figures 1,3A). No additional GO terms other than the three present in all oxygen
levels for *Eurypon* sp. 2 were enriched in both the pairwise comparison of anoxia and anoxia vs.
hypoxia.
584
No GO terms were significantly up or downregulated in all three levels of oxygen conditions for
*H. stellifera*. However, some GO terms were significantly enriched in two levels. The term
'ATPase, coupled to transmembrane movement of ions, rotational mechanism' was
downregulated in hypoxia but upregulated in anoxia ($p < 0.1$, Supplemental Figures 1B, 2B
respectively). The GO terms 'I-SMAD binding', thioesterase binding,' and 'ubiquitin-like protein
ligase binding' were all upregulated in hypoxia (vs. normoxia) and anoxia vs. hypoxia, while the
'motor' term was downregulated  ($p < 0.1$, Supplemental Figures 1B, 3B).
592
593 There were only two shared GO enrichments between both sponge species for any given
594 oxygen condition. The terms 'protein tyrosine kinase binding' and 'structural constituent of the
595 ribosome' were upregulated in both species in hypoxia ($p < 0.1$, Supplemental Figure 1),
596 although they were enriched to different degrees of significance. No significantly enriched GO
597 terms were shared between species in normoxia vs. anoxia or hypoxia vs. anoxia.
598
At the gene level, significantly differentially expressed heat shock proteins (Hsps) in both
sponges are shown in Supplemental Figure 8. In *Eurypon* sp. 2, one Hsp gene
(DNAJC8/Hsp40) was downregulated in hypoxia ($p < 0.01$), but four Hsp genes (HSPA8, two
genes for HSP90B1, and Hsp70) were significantly upregulated in hypoxia ($p < 0.1$,
Supplemental Figure 8A). Additionally, HSPA8 and C11orf73 (Hsp70 binding protein) were
upregulated in anoxia vs normoxia ($p < 0.05$, Supplemental Figure 8A). Another Hsp70 gene
was also significantly downregulated in anoxia vs. hypoxia ($p < 0.05$, Supplemental Figure 8A).
In *H. stellifera*, HSP90A1 was upregulated in hypoxia and anoxia ($p < 0.1$, Supplemental Figure
8B). Two additional Hsp genes (both for HSP90B1) were also upregulated in anoxia. Finally, the
two HSP90B1 genes were upregulated in anoxia vs. hypoxia, but HSPA5 was downregulated in
anoxia vs. hypoxia ($p < 0.05$, Supplemental Figure 8B).
610
**Mitochondrial gene expression**
All 14 coding genes from the sponge mitochondria were transcribed in both species and in all
oxygen conditions. There was only one mitochondrial gene (nad2) that was significantly
differentially expressed but only in *Eurypon* sp. 2. This gene was significantly upregulated ($p <
0.1$) in anoxia vs hypoxia (Supplemental Figure 6A, Table 1). Similar counts for each individual
mitochondrial gene across both sponge species and all oxygen conditions were found. For
instance, counts for atp6 were consistently twice as high as counts for atp8 (Supplemental
Figure 6).
619
**Prokaryotic differential expression**
621
*Thaumarchaeota* gene expression
For the *Thaumarchaeota* associated with *Eurypon* sp. 2, 20, 3 and 0 genes were differentially
expressed in hypoxia, anoxia, and hypoxia vs. anoxia, respectively ($p < 0.1$, Table 1). These
genes accounted for 0-1.3% of this *Thaumarchaeota* genome, and 75-100% of the differentially

expressed genes were annotated (Table 1), compared to 69.2% annotated in the entire
reference genome. All but one of the differentially expressed genes in hypoxia were upregulated
(Table 1, Figure 2A). The downregulated gene belonged to the sodium solute symporter family
(SSF, TC 2.A.21, Figure 2A). The upregulated genes in hypoxia included three stress response
genes (COG0589 Universal stress protein [UspA], small heat shock protein [HSP20], and DnaJ
molecular chaperone homology domain), two CBS domains, a member of the lactate/malate
dehydrogenase (LDH MDH) superfamily, a tetratricopeptide (TPR) repeat, two transcription
regulators, and a PII signal transduction protein.

635
**Figure 2.** A-D. All significantly differentially expressed genes for the *Thaumarchaeota* in both sponge
species (left: *E.* sp 2, right: *H. stellifera*. For *Thaumarchaeota* associated with *E.* sp. 2, only two pairwise
comparisons of oxygen conditions had significant differences: A. hypoxia vs. normoxia and C. anoxia vs.
normoxia. In symbionts associated with H. stellifera, there were significantly differentially expressed
genes in hypoxia vs. normoxia (B) and anoxia vs. hypoxia (D). There were no significant differences in the
three genes involved in ammonia oxidation (AmoA, B and C) in either symbiont, but their heatmaps are
shown in E and F for *E.* sp 2 and *H. stellifera*, respectively.

Only three genes were significantly differentially expressed in anoxia in *Eurypon* sp. 2's
*Thaumarchaeota*, and all were upregulated (Table 1, Figure 2C). The upregulated genes in
anoxia were a UbiD family decarboxylase, CAAX protease self-immunity, and the same TPR
repeat that was also upregulated in hypoxia. There were no significantly differentially expressed
genes between anoxia and hypoxia in this *Thaumarchaeota* (Table 1). The genes necessary for
ammonia oxidation, i.e. subunits A, B and C of ammonia monooxygenase (Amo), were present

in this *Thaumarchaeota*, but they were not differentially expressed under deoxygenation (Figure
2E).
652
In the *Thaumarchaeota* symbiont of *H. stellifera*, 17, 0 and 28 genes were significantly
differentially expressed in hypoxia, anoxia and hypoxia vs. anoxia, respectively. Similar to the
symbiont of *Eurypon* sp. 2, all 17 differentially expressed genes in the *H. stellifera* symbiont
were upregulated in hypoxia. These genes included the stress response genes 'universal stress
protein, an HSP20 family protein, and two copies of UspA. HSP20 and UspA were also
upregulated in the *Eurypon* sp. 2 holobiont. Similarly, a PII signal transduction protein, CBS
domain, and two transcription regulators (one identical: TBP stabilizer, and one distinct to *H.*
*stellifera* symbionts) were upregulated in both *Thaumarchaeota* species in hypoxia. Unlike in
*Eurypon* sp. 2, the *Thaumarchaeota* in *H. stellifera* upregulated a helix turn helix ASNC type
domain and a Lrp/AsnC ligand binding domain (Figure 2B). No genes were significantly
differentially expressed in anoxia in this *Thaumarchaeon*.
664
665  Twenty-seven genes were upregulated and one gene (K141_339108_19, unannotated) was
666  downregulated in anoxia vs. hypoxia in the *Thaumarchaeota* within *H. stellifera* (Figure 2D).
667  Most of the upregulated genes in anoxia vs. hypoxia were also upregulated in hypoxia. Some
668  unshared genes included additional transcriptional regulators, an integrase domain, and a
669  putative nitroreductase.
670
671   The phylogenies of this nitroreductase and all nitroreductases from the studied holobionts as
672  well as publicly available nitroreductase sequences from other *Thaumarchaeota*  are shown in
673  Supplemental Figure 9. As in the other holobiont species, *Thaumarchaeota* genes for ammonia
674  oxidation were not significantly differentially expressed with *H. stellifera* symbionts (Figure 2F).
675
676  For both *Thaumarchaeota*, only two COGs ('signal transduction mechanisms' and
677  'transcription') were significantly enriched ($P_{adj}$ < 0.05, Supplemental Figure 7A). These two
678  COGs were both downregulated in hypoxia and anoxia vs. hypoxia within *H. stellifera*
679  symbionts. All other COG classes were comparable in their expression levels regardless of
680  holobiont species or oxygen concentration (Supplemental Figure 7A).
681
*Gammaproteobacteria* gene expression
The *Gammaproteobacteria* associated with *Eurypon* sp. 2 only exhibited significant differential
expression under hypoxia but not in any other oxygen condition (Table 1). In hypoxia, three
genes were upregulated: type 1 restriction-modification system, flavin adenine dinucleotide
binding, and signal sequence binding (Figure 3A). The 29 downregulated genes in hypoxia
included two genes involved in sulfur processing (k212078_2, k237243_12), cytochrome c
(class I), a gene associated with the assembly of the periplasmic c-type cytochromes
(k561840_2), a NQR complex gene, various genes involved in pore formation, a Rhodanese-
like gene, glutamine synthetase (type I), and YbiS YcfS YnhG family protein.
691
692
693

**Figure 3.** All significantly differentially expressed genes for the *Gammaproteobacteria* in both sponge species (A: *Eurypon* sp 2, B: *H. stellifera*. For *Gammaproteobacteria* associated with *Eurypon* sp. 2, there were only significant differences between hypoxia vs. normoxia (A). In symbionts associated with *H. stellifera*, there were only significantly differentially expressed genes in anoxia vs. hypoxia (B,C). All downregulated genes in anoxia vs. hypoxia are shown in B. There were 402 genes that were downregulated in anoxia vs. hypoxia in *Gammaproteobacteria* from *H. stellifera*, so only the top 10 based on p value are shown (C).

In the *Gammaproteobacteria* associated with *H. stellifera*, genes were only significantly differentially expressed between anoxia and hypoxia (436 genes, p <0.1, Table 1, Figure 3B,C). Twenty genes were upregulated in anoxia vs. hypoxia and all were annotated (Figure 3B). Upregulated genes in anoxia vs. hypoxia included five genes associated with the ribosome (k141_198365_15, k141_198365_8, k141_198365_17, and k141_245017_3), various genes for molecular transport, a transmembrane proton pump (k141_226857_1), succinyl-CoA synthetase, and formate dehydrogenase (alpha subunit). There were also 416 genes that were significantly downregulated in anoxia vs. hypoxia (Table 1). The top ten upregulated genes (lowest p values) are exhibited in Figure 3C.

*Gammaproteobacterial* COG expression is shown in Supplemental Figure 7B. Only one COG class, 'translation, ribosomal structure and biogenesis,' was significantly enriched in *Gammaproteobacteria* under the different oxygen conditions ($P_{adj}$ < 0.05). This COG class was downregulated in *Gammaproteobacteria* associated with *Eurypon* sp. 2 under hypoxia as well as those of *H. stellifera* in anoxia vs. hypoxia (Figure 7B).

## Discussion

Sponge holobiont gene expression was significantly changed by exposure to deoxygenation in situ, though responses were generally species-specific, dependent on oxygen level, and varied across holobiont members. The potential functions of this differential expression are discussed

724 below for each member of the two holobionts: sponge, sponge mitochondria, *Thaumarchaeota*
725 and *Gammaproteobacteria*. The implications of this study for past and future oceans are
726 considered.
727
**Differential expression in sponges during deoxygenation**

There were clear differences in gene expression of the sponges in response to normoxia,
hypoxia, and anoxia. In both sponge species, the majority of significantly differentially expressed
genes were upregulated in hypoxia but downregulated in anoxia, when compared to normoxia.
Despite this general similarity in differential expression in response to oxygen conditions
between species, there were also noteworthy, species-specific responses to deoxygenation.
Thus, while both species significantly upregulated genes for responding to cellular stress in
hypoxia and anoxia, the specific genes depended on the species. Therefore, gene expression
depended on both oxygen condition and sponge species. Significantly enriched GO terms were
largely species specific and may indicate general physiological differences between sponge
species. KOG expression in response to deoxygenation was also largely species specific,
except for two key KOGs: 'energy production and conversion' (EPC) and 'replication,
recombination and repair' (RRR). Significant enrichment patterns of these KOGs were nearly
identical for both species, suggesting that differential expression of genes with EPC and RRR
functions is likely crucial to deoxygenation tolerance regardless of species. Interestingly, KOG
expression associated with metabolism, other than in the EPC class, was largely unchanged in
both studied species across all oxygen conditions, suggesting consistency in most metabolic
processes under deoxygenation.

Cellular stress responses were observed in the expression of HSPs, a group of stress response
genes that have been widely studied in marine invertebrates. These proteins are molecular
chaperones involved in protein folding, unfolding, transport and the assembly of protein
complexes, and they are present in nearly all organisms (Z. Li and Srivastava 2004). Both
Lough Hyne sponges upregulated HSP genes in response to hypoxia and anoxia (vs normoxia),
though there was downregulation of some HSP genes in deoxygenation as well. *Eurypon* sp. 2
and *H. stellifera* both upregulated HSP90 genes in hypoxia, but they upregulated different HSPs
in anoxia: HSPA8 and Hsp90 for *Eurypon* sp. 2 and *H. stellifera*, respectively. *Eurypon* sp. 2
also upregulated HSPA8 and Hsp70 in hypoxia but downregulated Hsp40. In long term hypoxia,
*T. wilhelma* exhibited a similar but inverse HSP expression pattern, upregulating Hsp90 but
downregulating Hsp70 (Mills et al. 2018). Sponges under heat stress also upregulate Hsp70
(López-Legentil et al. 2008; Guzman and Conaco 2016; N. Webster et al. 2013) and HSP90
(Guzman and Conaco 2016; N. Webster et al. 2013). Pollutants such as lead and zinc can also
stimulate the upregulation of HSPs in sponges (Efremova et al. 2002; Schröder et al. 2006).
These proteins are likely crucial to sponge stress response, and more work is necessary to fully
understand sponge stress response in general.

Patterns of species and oxygen-condition-dependent expression of GO terms were observed in
sponges in response to deoxygenation. In hypoxia, only one GO term was significantly
upregulated in both species - 'protein tyrosine kinase binding'. Tyrosine kinases are involved in
regulating a wide range of cellular functions, so extrapolating functionality from this GO term is
difficult. However, GO terms with more explicit functions were differentially enriched in both

species. GO terms such as 'proteasome binding' and 'Hsp70 binding' were upregulated in all
cases of decreased oxygen in *Eurypon* sp. 2, suggesting that the species-specific response to
deoxygenation in *Eurypon* sp. 2 may center around repairing or degrading damaged proteins. In
*H. stellifera*, on the other hand, upregulated GO terms related to DNA repair in anoxia, including
'damaged DNA binding', 'DNA binding,' and 'catalysis of DNA and RNA,' so DNA repair may be
more central to the deoxygenation tolerance of this species.

The EPC KOG class was significantly downregulated in all oxygen conditions in both sponges
except for hypoxia and hypoxia vs. anoxia in *H. stellifera*, when it was significantly upregulated
and unchanged, respectively (Figure 1). Within the EPC class, however, there were species-
specific responses to anoxia and hypoxia at the level of individual differentially expressed
genes. Most nuclear ATP synthase genes were upregulated in hypoxia in *Eurypon* sp. 2 but
were unchanged under anoxia. Similarly, two ATP synthase subunits (ATP5A1 and ATP6V0C)
were upregulated in hypoxia in *H. stellifera*. In anoxia, ATP5A1 was also upregulated, but
ATP6V0C was downregulated. Additionally, downregulation of SLC25A27 in *Eurypon* sp. 2 in
hypoxia may increase the coupling across the mitochondrial membrane to improve ATP
synthesis. Interestingly, the electron transport chain genes Ndufs8 and Surf1 were also
downregulated in hypoxia *Eurypon* sp. 2. The NNT gene, also involved in electron transport,
was downregulated in anoxia vs. hypoxia in *H. stellifera*. Nevertheless, most of the genes
associated with the respiration electron transport chain were not differentially expressed, and
electron transport likely continued under deoxygenation. Sponges may therefore have increased
ATP production under hypoxia but not under anoxia. Similarly, mollusks may upregulate the
mitochondrial electron transport system activity and ATP synthesis capacity to improve
respiration efficiency under hypoxia (Hochachka and Lutz 2001).

Glycolysis, which can produce ATP and NADH without oxygen (Stryer 1995), was expected to
increase under anoxia. However, relatively few genes directly or indirectly involved in glycolysis
were differentially expressed in our sponges under deoxygenation. In *Eurypon* sp. 2,
pantothenate kinase 1 (PANK1), ATP-dependent 6-phosphofructokinase (PFKL) and
Phosphoenolpyruvate carboxykinase 2 (PCK2) were significantly upregulated in anoxia, and
only aldehyde dehydrogenase 3 family, member a2 (ALDH3A2) was significantly upregulated in
hypoxia (see Supplemental Reports, Supplemental Figure 5). In *H. stellifera*, acetyl-coenzyme A
synthetase 1 (ACSS1) was significantly downregulated in hypoxia, but PCK1 and pyruvate
kinase (PKM) were upregulated in hypoxia. In anoxia for *H. stellifera*, beta-enolase (ENO3) and
PCK1 were upregulated. These results might indicate that glycolysis rates increased under
anoxia and hypoxia in both species. Due to the consistency in gene expression of the other ~70
genes associated with glycolysis; however, it is more likely that glycolysis continued under
anoxia and hypoxia at normal rates in both species. Either way, the sponges would have at
least some access to anaerobically generated ATP through glycolysis during environmental
deoxygenation. Similarly, gluconeogenesis, the production of glucose for glycolysis during
periods of fasting(Chandramouli et al. 1997), may have increased with deoxygenation. PCK
upregulation in anoxia in both sponge species and hypoxia in *H. stellifera* may indicate
increased gluconeogenesis. It is unclear if sponges are pumping water or feeding during anoxia,
since assays with fluorescein dye did not show obvious pumping in anoxia, but sponges were
definitively pumping under hypoxia (Schuster et al. 2021). However, increased gluconeogenesis
under anoxia might serve to boost energy levels regardless of pumping activity.

For the KOG class 'replication, recombination and repair' (RRR), there was significant
upregulation in anoxia and anoxia vs. hypoxia but down regulation in hypoxia for both sponge
species. In *Eurypon* sp. 2, many MCM genes involved in the initiation of DNA replication were
downregulated in hypoxia but remained unchanged in anoxia. Certain aspects of DNA
replication, e.g. the DNA polymerase epsilon catalytic subunit, may even be upregulated in
anoxia. The pattern was less clear in *H. stellifera*, but MCM genes were downregulated in
anoxia compared to hypoxia. These data indicated that DNA replication may decrease under
hypoxia but remain stable during anoxia. This pattern of a 'stronger' response to hypoxia than
anoxia mirrored that of nuclear ATP synthesis genes.

Between the two study species, there was only one significant, positive correlation of KOG class
expression between hypoxia (and normoxia) in *Eurypon* sp. 2 and hypoxia vs. anoxia in *H.*
*stellifera*, demonstrating the general species-specific responses of most KOGs to
deoxygenation. Species-specific responses between *Eurypon* sp. 2 and *H. stellifera* may reflect
the distinct  microbial communities harbored by each species (Schuster et al. 2021). In contrast,
different coral taxa generally exhibit more correlations in KOG expression in response to
different stressors than sponge taxa do in response to a single stressor, although stress
responses in corals can still differ widely between species and stressor (Kenkel, Mocellin, and
Bay 2020). Within a single sponge species, no correlations were seen between KOG
enrichments in anoxia or hypoxia (vs. normoxia), highlighting the transcriptomic differences
between responses to hypoxia and anoxia. KOG enrichments in hypoxia *versus* anoxia, on the
other hand, were positively correlated with those of anoxia in *Eurypon* sp. 2 and both anoxia
and hypoxia in *H. stellifera*.

When quantitatively compared to transcriptomic datasets from the sponge *T. wilhelma* using
KOGMWU, KOG enrichments in Lough Hyne sponges were only positively correlated between
the sponge *T. wilhelma* under short-term anoxia and *Eurypon* sp. 2 in hypoxia vs. anoxia
(Figure 1E). This positive correlation could indicate similar transcription patterns demonstrated
by these two sponge species transitioning from hypoxia to anoxia. However, no further
similarities between the responses of *T. wilhelma* and Lough Hyne sponges to deoxygenation
were observed. In fact, the long-term exposure of *T. wilhelma* to hypoxia was negatively
correlated to KOG enrichments for both anoxia vs. hypoxia and normoxia vs. anoxia in *Eurypon*
sp. 2 (Figure 1F, G), meaning that KOG expression was generally opposite between these two
species for these conditions in terms of up and down regulation. The physiological responses of
these two species to deoxygenation are presumably very different. As indicated by analyses in
Mills et al.(2018), there was also no correlation between the KOG expression for the exposure
of *T. wilhelma* to anoxia and hypoxia (Figure 1A). These results demonstrate that the large-
scale transcriptomic responses to deoxygenation depend on both the studied species and
oxygen concentration.

Physiological and behavioral responses of sponges to various levels of hypoxia also depend on
sponge species, oxygen level, and exposure duration (Micaroni, Strano, et al. 2021). In their
study (Micaroni, Strano, et al. 2021) found that only one of four tested species, *S. carnosus,*
exhibited a significant change in respiration rate in response to hypoxic (0.5 mg L$^{-1}$---) exposure,
halving its respiration rate. This decrease occurred only after seven days of exposure,
demonstrating the effects of exposure duration on metabolism. *S. australiensis* expanded its

body and aquiferous system and temporarily increased its pumping rate following hypoxic (0.4
mg L$^{-1---}$) exposure, but these behavioral changes were not observed under more severe
hypoxia (0.13 mg L$^{-1---}$). A decrease in behavioral response to more severe hypoxia was
mirrored in gene expression of ATP synthase genes in the current study. There may be an
oxygen threshold, below which increased production of ATP is no longer advantageous. We
hypothesize that these different physiological and behavioral responses underpin tolerance to
hypoxia and may be controlled by differential transcriptomic responses, which warrant further
investigation.

Based on KOG correlation analyses, we can reject the hypothesis that sponges are dormant,
i.e. temporarily ceasing growth, development, and physical activity, under anoxia. Sponge gene
expression under anoxia did not correlate to either diapause or a dauer state in *C. elegans*
*(Sinha, Sommer, and Dieterich 2012)* or *S. mosellana (Gong et al. 2013)*, respectively. Although
these dormancy datasets come from distantly related invertebrates, Strader et al. (2016)
demonstrated a correlation between the KOG expression patterns of a specific phenotype of
coral larvae and diapause in *S. mosellana*, hypothesizing that this potential functional analog of
dormancy could serve to increase larval dispersal. Therefore, the sponges of Lough Hyne could
only be considered dormant under anoxia if their transcriptional method of dormancy was
distinct from that of an insect, nematode and coral, which is possible but not parsimonious given
that animals are monophyletic (Feuda et al. 2017).

Paradoxically, *Eurypon* sp. 2 responses to hypoxia were positively correlated to diapause in *S.*
*mosellana* (Figure 1B). Therefore, the hypoxic response in *Eurypon* sp. 2 species, rather than
the anoxic response, was similar to diapause in terms of global KOG expression patterns. The
RRR kog was also downregulated in *Eurypon* sp. 2 under hypoxia as well as in *S. mosellana*
under diapause, which may partially explain the correlation between these datasets. Moreover,
diapause can be associated with the reduction of cell division (MacRae 2010), which may occur
in sponges under hypoxia but not anoxia based on MCM gene expression. In comparison, KOG
enrichments under anoxia were not positively correlated to any external datasets for either
sponge species (Figure 1), indicating that sponge transcriptomic responses to anoxia are
relatively novel compared to those of other taxa in response to various factors in the existing
literature (Dixon et al. 2015; Strader, Aglyamova, and Matz 2016).
884
885    **Sponge mitochondrial gene expression during deoxygenation**
886
A high degree of transcriptional stability under deoxygenation was observed in sponge
mitochondria. Only one of the 14 mitochondrial, protein coding genes, nad2, was significantly
differentially expressed in the different oxygen conditions, where it was significantly upregulated
in anoxia vs. hypoxia in *Eurypon* sp 2. Both of the sponge species had higher expressions of
nad2 in anoxia than in normoxia or hypoxia, but this difference was only significant in *Eurypon*
sp. 2. Both species also demonstrated remarkably similar and stable patterns of expression for
each mitochondrial gene, including nad2.
894
Surprisingly little is known about differential gene expression in mitochondria, even within
humans. In the limited studies available, mitochondrial gene expression has significantly differed
between human cancer cells (Reznik et al. 2017) and normal cells as well as between different

populations of healthy pancreas tissue (Medini, Cohen, and Mishmar 2021). In these studies
nad2 was significantly downregulated in most studied cancers, though it was still significantly
upregulated in a few cancer types, and nad2 was also significantly differentially expressed
between pancreas cell populations. Based on general upregulation of nuclear genes associated
with respiration in hypoxia, mitochondrial genes were expected to be upregulated in hypoxia
when compared to normoxia and anoxia, but the opposite was true for nad2 in *Eurypon* sp. 2.
The other six nad genes, which code for parts of the same membrane respiratory chain complex
(complex 1, NADH dehydrogenase), however, were not significantly differentially expressed in
either species. In both human studies, when one mitochondrial gene was significantly
differentially expressed, most mitochondrial genes were significantly differentially expressed in
the same direction (Reznik et al. 2017; Medini, Cohen, and Mishmar 2021). The physiological
meaning (if any) of this single, significantly differentially expressed mitochondrial gene in
sponges remains to be determined.
911  
Compared to that of humans, even less is known about mitochondrial gene expression in
marine invertebrates. Mitochondrial gene expression is usually excluded from differential
expression analyses due to the differences in library preparation potentially needed for
sequencing mitochondrial RNA and/or the exclusion of mitochondrial RNA from reference
transcriptomes (*sensu* Meyer (2016). This dearth is partly due to the nature of mitochondrial
transcription, which generates a single continuous transcript. Following transcription, however,
mitochondrial transcripts are broken up into individual coding mRNA, tRNA or rRNA (D'Souza
and Minczuk 2018). After the breakup, the levels of mRNAs from specific genes are controlled
post-transcriptionally (D'Souza and Minczuk 2018). It is therefore possible to have different
counts between the different mitochondrial genes despite their single transcript origin
(Supplemental Figure 6). These consistent 'expression' levels of transcripts across oxygen
conditions for each mitochondrial protein-coding gene following continuous transcription are
likely to be necessary for the maintenance of mitochondrial function.
925  
It remains unclear how sponge mitochondria maintain this transcription from an energetic
standpoint under anoxia. Within the mitochondrial electron transport chain, oxygen is used as
the terminal electron acceptor and is crucial for the generation of ATP through oxidative
phosphorylation. Theoretically, anoxia should cause electron transport and oxidative
phosphorylation in the mitochondria to cease (Chandel and Schumacker 2000). If the
mitochondria are transcribing solely to maintain oxidative phosphorylation machinery until
oxygen returns, then they would need to use at least enough energy (~2 ATP per base) to
transcribe for months at a time without oxygen (Lynch and Marinov 2015), and this energetic
requirement would apply to the transcription of sponge nuclear genes as well. Glycolysis may
provide some anaerobic ATP production, even if it is not upregulated in anoxia or hypoxia.
Alternatively, to maintain ATP production through aerobic respiration, the sponges of Lough
Hyne might have access to oxygen produced by their *Thaumarchaeota* symbionts, a topic
explored in more detail below.
939  

**Microbiome gene expression during deoxygenation**

*Thaumarchaeota* gene expression

*Thaumarchaeota* symbionts from both sponge species exhibited significantly differentially expressed genes under hypoxia. Only the *Thaumarchaeota* from *Eurypon* sp. 2 (OTU1) exhibited differential gene expression under anoxia, and only those of *H. stellifera* (OTU3) differentially expressed genes under anoxia vs. hypoxia. These species-specific responses within the symbionts could result from phylogenetic differences between the two symbionts. OTU3 is more closely related to the free-living *Nitrosopumilus maritimus* than OTU1 (Schuster et al. 2021). Though there were some species-specific responses, overall gene expression in both *Thaumarchaeota* was similar, supporting the hypothesis from Schuster et al. (2021) that these taxonomically similar symbionts between two sponge species might have similar functional roles.

In hypoxia, both *Thaumarchaeota* upregulated the stress response genes USpA and a protein from the HSP20 family. It was surprising that *Thaumarchaeota* may be stressed by hypoxia. free-living thaumarchaeal Nitrosopumilales can be found in high abundances in oxygen-minimum zones (OMZs) where oxygen is either at low concentrations or below detection (Löscher et al. 2012), though some *Thaumarchaeota* OTUs decrease in abundance in suboxic and anoxic zones (Sollai et al. 2019). Members of the genus *Nitrosopumilus*, however, increase their absolute abundance in suboxic regions, but then decrease in abundance with depth in the anoxic zone (Sollai et al. 2019), indicating a preference for suboxic environments. If suboxia is also the preferred niche of sponge *Thaumarchaeota* symbionts, the upregulation of stress response genes in response to hypoxia may correspond to a rise in intracellular oxidative stress following an increase in metabolic activity in hypoxia.

Based on the genomes of their close relatives, the *Thaumarchaeota* in these sponges were hypothesized to have components of a facultatively anaerobic metabolism, including fumarate, nitrite, and sulfite respiration and the ability to ferment (Bayer et al. 2020), which could aid in deoxygenation tolerance. Lactate/malate dehydrogenase, which could be involved in fermentation (Bayer et al. 2020), was upregulated in *Eurypon* sp. 2 *Thaumarchaeota*, but only in hypoxia. Genes for the other anaerobic pathways suggested by Bayer et al. 2020, i.e. succinate dehydrogenase/ fumarate reductase, nitrite reductase and sulfite reductase, were present in both *Thaumarchaeota* in the present study but were not significantly differentially expressed under different oxygen conditions. The genes for nitrite reductase exhibited low expression across all samples (0–20 counts) in both *Thaumarcheaota,* and so did the dehydrogenase/ fumarate reductase gene in the *Thaumarcheaota* of *H. stellifera*. Therefore, these genes with low expression may not be used under the conditions of this study. Only three genes in the *Thaumarchaeota* of *Eurypon* sp. 2 under anoxia were significantly differentially expressed, and none of these were involved in anaerobic metabolism. No genes were differentially expressed in *H. stellifera Thaumarchaeota* under anoxia. Therefore, there was no transcriptomic evidence for the upregulation of anaerobic metabolism within *Thaumarchaeota* in anoxia.

Transcription of ammonium monooxygenase (amo) genes did not significantly change under anoxia or hypoxia, even though the ammonia oxidation pathway is oxygen dependent. Like the

sponge mitochondria, the *Thaumarchaeota* are therefore either transcribing amo genes to be
ready when oxygen returns, or they are obtaining trace oxygen for this pathway from another
source. A recent study demonstrated that *N. maritimus* produces trace levels of oxygen (100–
200 nM) under environmental anoxia, and it uses this oxygen to oxidize ammonia (Kraft et al.
2022). If the closely related sponge *Thaumarchaeota* symbionts are also able to produce
oxygen under anoxic conditions, then they should express amo genes under anoxia as well as
normoxia.

In comparing the proteome during oxygen production under oxygen depleted conditions versus
periods when no oxygen was produced under 100% a.s. by *N. maritimus*, Kraft et al. (2021)
found that proteins upregulated during oxygen production included some AsnC family proteins
and nitroreductases. In *Archaea*, AsnC proteins can regulate metabolic processes, including the
switch between aerobic and anaerobic metabolisms (Lemmens et al. 2019). Similarly,
nitroreductases can have a wide range of metabolic functions (Akiva et al. 2017).
*Thaumarchaeota* within *H. stellifera* upregulated a Lrp/AsnC ligand binding domain under
hypoxia and in anoxia vs. hypoxia. Another AsnC type protein was also upregulated under
hypoxia, and a nitroreductase under anoxia vs. hypoxia. However, these proteins did not directly
correspond to those upregulated in *N. maritimus* during oxygen production (Supplemental
Figure 9), and the latter proteins are not definitively known to be involved in oxygen production,
since the pathway remains unknown.
1000
*Gammaproteobacteria* gene expression

Gene expression in the *Gammaproteobacteria* was species and oxygen-condition dependent. In
*Eurypon* sp. 2, *Gammaproteobacteria* downregulated genes in hypoxia that were associated
with respiration and energy production, including cytochrome c and associated genes (Welchen
et al. 2016). The *Gammaproteobacteria* also downregulated key metabolic genes in hypoxia
such as glutamine synthetase and structural genes like YbiS YcfS YnhG family protein, which is
involved in cell wall production (Magnet et al. 2008). So unlike sponges or *Thaumarchaeota*,
*Gammaproteobacteria* in *Eurypon* sp. 2 (OTU2) may decrease metabolic activity under hypoxia,
though this is likely not the case in *H. stellifera* symbionts (OTU7).

*Gammaproteobacteria* in *H. stellifera* downregulated hundreds of genes in anoxia vs. hypoxia,
including USpA, so, like other members of the holobiont, hypoxia requires a stronger
transcriptomic response and, therefore, may be more stressful than anoxia for these
*Gammaproteobacterial symbionts*. The few upregulated genes in anoxia vs. hypoxia included
genes for energy production, e.g. a transmembrane proton pump, succinyl-CoA synthetase (part
of the citric acid cycle), and formate dehydrogenase (alpha subunit), though other genes
associated with energy production were downregulated. Formate dehydrogenase can function
in both aerobic respiration or in anaerobic respiration(Jormakka, Byrne, and Iwata 2003), but
this enzyme was not significantly upregulated in anoxia compared to normoxia or in the
*Gammaproteobacteria* of *Eurypon* sp. 2 under any condition. So, it is unlikely that this anaerobic
pathway is active in sponge-associated *Gammaproteobacteria* under anoxia.

*Gammaproteobacteria* symbionts were previously hypothesized to use sulfide or thiosulfate as
an electron donor anaerobically coupled to nitrate reduction, removing exogenous, toxic sulfide

from the holobiont under anoxia (Schuster et al. 2021). Although they are not present in OTU2,
OTU7 from *H. stellifera* has the sulfur oxidation genes soxA, soxY, and soxZ, which could
oxidize sulfide and the nitrate reductase genes NAP. However, the sox genes were not
expressed in any samples in the current study, and NAP was expressed at very low levels
across all samples (2-28 counts). Moreover, neither symbiont exhibited significant differential
gene expression in anoxia versus normoxia. Consequently, there was very little transcriptomic
evidence that the anaerobic pathways in these *Gammaproteobacteria* were activated during
environmental anoxia.

**Adaptations of sponge holobionts to low oxygen**

No clear anaerobic respiration responses were observed in the transcriptomes of sponges, their
mitochondria or their two microbial symbionts under environmental anoxia. In the sponges and
their microbes, glycolysis could potentially provide ATP without oxygen. If so, the limited
significant differential expression of glycolysis genes could indicate that this metabolic process
continues to comparable degrees under normoxia, hypoxia and anoxia. Metabolically, sponge
transcription under anoxia was surprisingly similar to normoxia throughout the holobiont
members, and there was no evidence of dormancy within the holobionts. Instead, in the
sponges, hypoxia might elicit increases in energy production machinery such as ATP synthase.
These potential metabolic increases may indicate that within the hypoxic limits studied here
(1.30-3.56 mg $L^{-1}$), sponges perform optimally. Indeed, these limits are above those at which
mortality or substantial physiological changes have been observed for sponges (Micaroni,
Strano, et al. 2021; Kumala et al. 2021; Mills et al. 2014). This increased energy production
could be stressful, however, and the diversion of resources could limit cell division in hypoxia.
Somewhere between the lower limit of hypoxia (1.37 mg $L^{-1}$) and the full anoxia studied here,
sponges may no longer attempt to increase energy efficiency, and therefore elevated
transcription of ATP synthase genes was not observed in anoxia. For both the sponges and
their *Thaumarcheaota* symbionts, cellular stress responses were evident under deoxygenation,
indicating that the holobionts respond to these conditions at a cellular level. The gene
expression associated with anoxia could either be a direct response, via currently unknown
mechanisms, to deoxygenation itself or to the presence of hydrogen sulfide in anoxia.

Given the deoxygenation tolerance of Lough Hyne sponges and their largely unchanged
metabolic gene expression patterns under deoxygenation, the production of oxygen by
*Thaumarchaeota* could also be crucial to holobiont survival. Although any oxygen produced
would be at extremely low concentrations, it would have a short path of diffusion to the sponge
cells, as *Thaumarchaeota* symbionts maintain high cell densities within sponges (Moeller et al.
2019). In addition, mitochondria in general can utilize very low concentrations of oxygen (500
nM) without changing their oxygen consumption rate (Chance 1965; Longmuir 1957; Wilson et
al. 1979). It is plausible that, on a small spatial scale, oxygen generated by *Thaumarchaeota*
could reach these concentrations under anoxia, allowing mitochondrial function to continue in
sponges under anoxia. However, given that the genes responsible are still unknown, this
pathway is still speculative in sponge *Thaumarchaeota*.

1063

1064 **Relevance to future ocean deoxygenation**

1065
Sponges are predicted to be the 'winners' over other benthic organisms in the future due to their
tolerance of ocean warming and acidification associated with climate change (Bell et al. 2018).
Indeed, sponges have already increased in abundance in some locations while corals have
decreased due to altered carbon and nutrient cycling (Pawlik, Burkepile, and Thurber 2016).
The additional stressor of deoxygenation that comes with climate change and coastal
eutrophication has been relatively overlooked compared to ocean warming and acidification. In
other marine taxa, a meta-analysis demonstrated that the incorporation of deoxygenation as a
factor along with warming could substantially decrease survival times of marine organisms
compared to warming alone (Vaquer-Sunyer and Duarte 2011). Since sponges may be more
tolerant than other taxa to changes in all three stressors of temperature, pH and oxygen, they
could indeed become more abundant in a changing ocean. However, even sponges can be
lethally affected by warming, acidification (Bennett et al. 2017) and deoxygenation (Micaroni,
Strano, et al. 2021) in scenarios that might occur with climate change, since tolerance to these
stressors is species-specific. So, surviving sponge species in a warmed, acidified, and
deoxygenated ocean might be abundant, but their abundance, and diversity would likely be
lower than at present. The effects of all three of these stressors together need to be understood
on a much wider range of organisms to accurately predict the composition of future oceans.
More importantly, the causes and effects of anthropogenic climate change need to be
ameliorated as soon as possible to avoid such diversity loss.

**Relevance for animal evolution**

Assuming that the first metazoans were more like an adult sponge than a ctenophore, based on
morphology (Nielsen 2008) and phylogeny (Feuda et al. 2017), then some species of early
sponges or proto-animals could have tolerated low environmental oxygen levels and at least
some periods of benthic anoxia and euxinia in the Neoproterozoic Era (~538.8–1,000 million
years ago) when early animals, including sponges, first evolved (Dohrmann and Wörheide
2017). Lough Hyne sponges are a valuable model system for understanding early animal
evolution. Early sponges could have survived under low and variable levels using mechanisms
like those of Lough Hyne sponges. Therefore, contrary to longstanding hypotheses, the
evolution of animals probably did not require high and stable environmental oxygen levels.

**Acknowledgements**

Field work at Lough Hyne was carried out with the permission of the National Parks and Wildlife
Service of Ireland. We thank Luke Harman and Allen Whittaker of University College Cork. We
also thank the Nordcee lab technicians of the Department of Biology at University of Southern
Denmark for constant support in the laboratory procedures.

## Funding

This project was funded by Villum Fonden grant no. 16518. B.K. was funded by the Villum Young Investigator Grant 25491.

## Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

## Data, scripts, code, and supplementary information availability

Raw sequence data as well as genome and transcriptome assemblies are available at NCBI under BioProject number PRJNA893197 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA893197). Scripts, code, and additional data are available here: https://zenodo.org/badge/latestdoi/573040118. Supplemental figures and tables are available on BioRxiv under 'Supplementary material.'

## References

Akiva, Eyal, Janine N. Copp, Nobuhiko Tokuriki, and Patricia C. Babbitt. 2017. "Evolutionary and Molecular Foundations of Multiple Contemporary Functions of the Nitroreductase Superfamily." *Proceedings of the National Academy of Sciences of the United States of America* 114 (45): E9549–58.

Altieri, Andrew H., Seamus B. Harrison, Janina Seemann, Rachel Collin, Robert J. Diaz, and Nancy Knowlton. 2017. "Tropical Dead Zones and Mass Mortalities on Coral Reefs." *Proceedings of the National Academy of Sciences of the United States of America* 114 (14): 3660–65.

Bayer, Kristina, Kathrin Busch, Ellen Kenchington, Lindsay Beazley, Sören Franzenburg, Jan Michels, Ute Hentschel, and Beate M. Slaby. 2020. "Microbial Strategies for Survival in the Glass Sponge Vazella Pourtalesii." *mSystems*. https://doi.org/10.1128/msystems.00473-20.

Bell, James J., and David K. A. Barnes. 2000. "The Distribution and Prevalence of Sponges in Relation to Environmental Gradients within a Temperate Sea Lough: Vertical Cliff Surfaces." *Diversity Distributions*. https://doi.org/10.1046/j.1472-4642.2000.00091.x.

Bell, James J., Holly M. Bennett, Alberto Rovellini, and Nicole S. Webster. 2018. "Sponges to Be Winners under Near-Future Climate Scenarios." *Bioscience* 68 (12): 955–68.

Bennett, Holly M., Christine Altenrath, Lisa Woods, Simon K. Davy, Nicole S. Webster, and James J. Bell. 2017. "Interactive Effects of Temperature and pCO2 on Sponges: From the Cradle to the Grave." *Global Change Biology* 23 (5): 2031–46.

Breitburg, Denise, Lisa A. Levin, Andreas Oschlies, Marilaure Grégoire, Francisco P. Chavez, Daniel J. Conley, Véronique Garçon, et al. 2018. "Declining Oxygen in the Global Ocean and Coastal Waters." *Science* 359 (6371). https://doi.org/10.1126/science.aam7240.

Chance, B. 1965. "Reaction of Oxygen with the Respiratory Chain in Cells and Tissues." *The Journal of General Physiology* 49 (1): Suppl:163–95.

Chandel, N. S., and P. T. Schumacker. 2000. "Cellular Oxygen Sensing by Mitochondria: Old Questions, New Insight." *Journal of Applied Physiology* 88 (5): 1880–89.

Chandramouli, V., K. Ekberg, W. C. Schumann, S. C. Kalhan, J. Wahren, and B. R. Landau. 1997. "Quantifying Gluconeogenesis during Fasting." *The American Journal of Physiology* 273 (6): E1209–15.

Diaz, Robert J., and Rutger Rosenberg. 2008. "Spreading Dead Zones and Consequences for Marine Ecosystems." *Science* 321 (5891): 926–29.

Dixon, Groves B., Sarah W. Davies, Galina A. Aglyamova, Eli Meyer, Line K. Bay, and Mikhail V. Matz. 2015. "CORAL REEFS. Genomic Determinants of Coral Heat Tolerance across Latitudes." *Science* 348 (6242): 1460–62.

Dohrmann, Martin, and Gert Wörheide. 2017. "Dating Early Animal Evolution Using Phylogenomic Data." *Scientific Reports* 7 (1): 1–6.

D'Souza, Aaron R., and Michal Minczuk. 2018. "Mitochondrial Transcription and Translation: Overview." *Essays in Biochemistry.* https://doi.org/10.1042/ebc20170102.

Edler, Daniel, Johannes Klein, Alexandre Antonelli, and Daniele Silvestro. 2020. "raxmlGUI 2.0: A Graphical Interface and Toolkit for Phylogenetic Analyses Using RAxML." *Methods in Ecology and Evolution,* 373–77.

Efremova, Sofia M., Boris A. Margulis, Irina V. Guzhova, Valeria B. Itskovich, Stephanie Lauenroth, Werner E. G. Müller, and Heinz C. Schröder. 2002. "Heat Shock Protein Hsp70 Expression and DNA Damage in Baikalian Sponges Exposed to Model Pollutants and Wastewater from Baikalsk Pulp and Paper Plant." *Aquatic Toxicology* 57 (4): 267–80.

Ellington, W. Ross. 1982. "Metabolic Responses of the Sea Anemone Bunodosoma Cavernata (Bosc) to Declining Oxygen Tensions and Anoxia." *Physiological Zoology* 55 (3): 240–49.

Feuda, Roberto, Martin Dohrmann, Walker Pett, Hervé Philippe, Omar Rota-Stabelli, Nicolas Lartillot, Gert Wörheide, and Davide Pisani. 2017. "Improved Modeling of Compositional Heterogeneity Supports Sponges as Sister to All Other Animals." *Current Biology: CB* 27 (24): 3864–70.e4.

Francis, Warren. 2020a. "General Sequence Processing Scripts." Bitbucket. 2020. bitbucket.org/wrf/sequences.

———. 2020b. "Make Interactive Plot of Contigs in Genomes or Metagenomes." Gihub. 2020. https://github.com/wrf/lavaLampPlot/blob/master/Rshiny/app.R.

Gong, Zhong-Jun, Yu-Qing Wu, Jin Miao, Yun Duan, Yue-Li Jiang, and Tong Li. 2013. "Global Transcriptome Analysis of Orange Wheat Blossom Midge, Sitodiplosis Mosellana (Gehin) (Diptera: Cecidomyiidae) to Identify Candidate Transcripts Regulating Diapause." *PloS One* 8 (8): e71564.

Grabherr, Manfred G., Brian J. Haas, Moran Yassour, Joshua Z. Levin, Dawn A. Thompson, Ido Amit, Xian Adiconis, et al. 2011. "Full-Length Transcriptome Assembly from RNA-Seq Data without a Reference Genome." *Nature Biotechnology.* https://doi.org/10.1038/nbt.1883.

Gunda, Venu Gopal, and Venkateswara Rao Janapala. 2009. "Effects of Dissolved Oxygen Levels on Survival and Growth in Vitro of Haliclona Pigmentifera (Demospongiae)." *Cell and Tissue Research.* https://doi.org/10.1007/s00441-009-0843-5.

Guzman, Christine, and Cecilia Conaco. 2016. "Gene Expression Dynamics Accompanying the Sponge Thermal Stress Response." *PloS One* 11 (10): e0165368.

Hochachka, P. W., and P. L. Lutz. 2001. "Mechanism, Origin, and Evolution of Anoxia Tolerance in Animals." *Comparative Biochemistry and Physiology. Part B, Biochemistry & Molecular Biology* 130 (4): 435–59.

Hoffmann, Friederike, Ole Larsen, Volker Thiel, Hans Tore Rapp, Thomas Pape, Walter Michaelis, and Joachim Reitner. 2005. "An Anaerobic World in Sponges." *Geomicrobiology Journal* 22 (1-2): 1–10.

Hoffmann, Friederike, Regina Radax, Dagmar Woebken, Moritz Holtappels, Gaute Lavik, Hans Tore Rapp, Marie-Lise Schläppy, Christa Schleper, and Marcel M. M. Kuypers. 2009. "Complex Nitrogen Cycling in the Sponge Geodia Barretti." *Environmental Microbiology* 11

(9): 2228–43.

Huerta-Cepas, Jaime, Damian Szklarczyk, Davide Heller, Ana Hernández-Plaza, Sofia K. Forslund, Helen Cook, Daniel R. Mende, et al. 2019. "eggNOG 5.0: A Hierarchical, Functionally and Phylogenetically Annotated Orthology Resource Based on 5090 Organisms and 2502 Viruses." *Nucleic Acids Research*. https://doi.org/10.1093/nar/gky1085.

Hughes, David J., Rachel Alderdice, Christopher Cooney, Michael Kühl, Mathieu Pernice, Christian R. Voolstra, and David J. Suggett. 2020. "Coral Reef Survival under Accelerating Ocean Deoxygenation." *Nature Climate Change* 10 (4): 296–307.

Huntley, Melanie A., Jessica L. Larson, Christina Chaivorapol, Gabriel Becker, Michael Lawrence, Jason A. Hackney, and Joshua S. Kaminker. 2013. "ReportingTools: An Automated Result Processing and Presentation Toolkit for High-Throughput Genomic Analyses." *Bioinformatics* 29 (24): 3220–21.

Jormakka, Mika, Bernadette Byrne, and So Iwata. 2003. "Formate Dehydrogenase--a Versatile Enzyme in Changing Environments." *Current Opinion in Structural Biology* 13 (4): 418–23.

Katoh, Kazutaka, and Daron M. Standley. 2013. "MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability." *Molecular Biology and Evolution* 30 (4): 772–80.

Keeling, Ralph E., Arne Körtzinger, and Nicolas Gruber. 2010. "Ocean Deoxygenation in a Warming World." *Annual Review of Marine Science* 2: 199–229.

Kenkel, Carly D., Veronique J. L. Mocellin, and Line K. Bay. 2020. "Global Gene Expression Patterns in Porites White Patch Syndrome: Disentangling Symbiont Loss from the Thermal Stress Response in Reef-Building Coral." *Molecular Ecology* 29 (20): 3907–20.

Kirchman, David L. 2021. *Dead Zones: The Loss of Oxygen from Rivers, Lakes, Seas, and the Ocean*. Oxford University Press, USA.

Kolde, R. 2013. *Pheatmap: Pretty Heatmaps. R-Package* (version 0.7.4). http://www2.uaem.mx/r-mirror/web/packages/pheatmap/pheatmap.pdf.

Kraft, Beate, Nico Jehmlich, Morten Larsen, Laura A. Bristow, Martin Könneke, Bo Thamdrup, and Donald E. Canfield. 2022. "Oxygen and Nitrogen Production by an Ammonia-Oxidizing Archaeon." *Science* 375 (6576): 97–100.

Kraft, Beate, Nico Jehmlich, Morten Larsen, Laura Bristow, Martin Könneke, Bo Thamdrup, and Donald E. Canfield. 2021. "Oxygen Production by an Ammonia-Oxidizing Archaeon." *bioRxiv*. https://doi.org/10.1101/2021.04.01.436977.

Kumala, Lars, Morten Larsen, Ronnie N. Glud, and Donald E. Canfield. 2021. "Spatial and Temporal Anoxia in Single-Osculum Halichondria Panicea Demosponge Explants Studied with Planar Optodes." *Marine Biology* 168 (12). https://doi.org/10.1007/s00227-021-03980-2.

Langmead, Ben, and Steven L. Salzberg. 2012. "Fast Gapped-Read Alignment with Bowtie 2." *Nature Methods* 9 (4): 357–59.

Larade, Kevin, and Kenneth B. Storey. 2009. "Living without Oxygen: Anoxia-Responsive Gene Expression and Regulation." *Current Genomics* 10 (2): 76–85.

Lemmens, Liesbeth, Hassan Ramadan Maklad, Indra Bervoets, and Eveline Peeters. 2019. "Transcription Regulators in Archaea: Homologies and Differences with Bacterial Regulators." *Journal of Molecular Biology* 431 (20): 4132–46.

Levin, Lisa A. 2018. "Manifestation, Drivers, and Emergence of Open Ocean Deoxygenation." *Annual Review of Marine Science* 10 (January): 229–60.

Leys, Sally P., and Amanda S. Kahn. 2018. "Oxygen and the Energetic Requirements of the First Multicellular Animals." *Integrative and Comparative Biology*. https://doi.org/10.1093/icb/icy051.

Li, Dinghua, Ruibang Luo, Chi-Man Liu, Chi-Ming Leung, Hing-Fung Ting, Kunihiko Sadakane, Hiroshi Yamashita, and Tak-Wah Lam. 2016. "MEGAHIT v1.0: A Fast and Scalable Metagenome Assembler Driven by Advanced Methodologies and Community Practices."

*Methods*. https://doi.org/10.1016/j.ymeth.2016.02.020.

Li, Zihai, and Pramod Srivastava. 2004. "Heat-Shock Proteins." *Current Protocols in Immunology / Edited by John E. Coligan ... [et Al.]* Appendix 1 (February): Appendix 1T.

Longmuir, I. S. 1957. "Respiration Rate of Rat-Liver Cells at Low Oxygen Concentrations." *Biochemical Journal* 65 (2): 378–82.

López-Legentil, Susanna, Bongkeun Song, Steven E. McMurray, and Joseph R. Pawlik. 2008. "Bleaching and Stress in Coral Reef Ecosystems: hsp70 Expression by the Giant Barrel Sponge Xestospongia Muta." *Molecular Ecology* 17 (7): 1840–49.

Löscher, C. R., A. Kock, M. Könneke, J. LaRoche, H. W. Bange, and R. A. Schmitz. 2012. "Production of Oceanic Nitrous Oxide by Ammonia-Oxidizing Archaea." *Biogeosciences* 9 (July): 2419–29.

Love, Michael I., Wolfgang Huber, and Simon Anders. 2016. "Moderated Estimation of Fold Change and Dispersion for RNA-Seq Data with DESeq2." *BMC Genome Biology* 15 (January): 1–22.

Lynch, Michael, and Georgi K. Marinov. 2015. "The Bioenergetic Costs of a Gene." *Proceedings of the National Academy of Sciences of the United States of America* 112 (51): 15690–95.

MacRae, Thomas H. 2010. "Gene Expression, Metabolic Regulation and Stress Tolerance during Diapause." *Cellular and Molecular Life Sciences: CMLS* 67 (14): 2405–24.

Matz, M. V. 2015. "Assembly and Annotation of Transcriptomes de Novo." Github. 2015. https://github.com/z0on/annotatingTranscriptomes.

Medini, H., T. Cohen, and D. Mishmar. 2021. "Mitochondrial Gene Expression in Single Cells Shape Pancreatic Beta Cells' Sub-Populations and Explain Variation in Insulin Pathway." *Scientific Reports* 11 (1): 466.

Meyer, E. 2016. "A Collection of Scripts for Analyzing Transcriptome Assemblies." Github. 2016. https://github.com/Eli-Meyer/transcriptome_utilities.

Micaroni, Valerio, Rob McAllen, John Turner, Francesca Strano, Christine Morrow, Bernard Picton, Luke Harman, and James J. Bell. 2021. "Vulnerability of Temperate Mesophotic Ecosystems (TMEs) to Environmental Impacts: Rapid Ecosystem Changes at Lough Hyne Marine Nature Reserve, Ireland." *Science of The Total Environment*. https://doi.org/10.1016/j.scitotenv.2021.147708.

Micaroni, Valerio, Francesca Strano, Rob McAllen, Lisa Woods, John Turner, Luke Harman, and James J. Bell. 2021. "Adaptive Strategies of Sponges to Deoxygenated Oceans." *Global Change Biology*, December. https://doi.org/10.1111/gcb.16013.

Mills, Daniel B., Warren R. Francis, Sergio Vargas, Morten Larsen, Coen Ph Elemans, Donald E. Canfield, and Gert Wörheide. 2018. "The Last Common Ancestor of Animals Lacked the HIF Pathway and Respired in Low-Oxygen Environments." *eLife* 7 (February). https://doi.org/10.7554/eLife.31176.

Mills, Daniel B., Lewis M. Ward, Carriayne Jones, Brittany Sweeten, Michael Forth, Alexander H. Treusch, and Donald E. Canfield. 2014. "Oxygen Requirements of the Earliest Animals." *Proceedings of the National Academy of Sciences of the United States of America* 111 (11): 4168–72.

Moeller, Florian U., Nicole S. Webster, Craig W. Herbold, Faris Behnam, Daryl Domman, Mads Albertsen, Maria Mooshammer, et al. 2019. "Characterization of a Thaumarchaeal Symbiont That Drives Incomplete Nitrification in the Tropical Sponge Ianthella Basta." *Environmental Microbiology* 21 (10): 3831–54.

Nielsen, Claus. 2008. "Six Major Steps in Animal Evolution: Are We Derived Sponge Larvae?" *Evolution & Development* 10 (2): 241–57.

Pawlik, Joseph R., Deron E. Burkepile, and Rebecca Vega Thurber. 2016. "A Vicious Circle? Altered Carbon and Nutrient Cycling May Explain the Low Resilience of Caribbean Coral Reefs." *BioScience*. https://doi.org/10.1093/biosci/biw047.

Plese, Bruna, Maria Eleonora Rossi, Nathan James Kenny, Sergi Taboada, Vasiliki Koutsouveli, and Ana Riesgo. 2019. "Trimitomics: An Efficient Pipeline for Mitochondrial Assembly from

Transcriptomic Reads in Nonmodel Species." *Molecular Ecology Resources* 19 (5): 1230–39.

R Core Team. 2016. *R: A Language and Environment for Statistical Computing. Vienna, Austria*. https://www.R-project.org/.

Reznik, Ed, Qingguo Wang, Konnor La, Nikolaus Schultz, and Chris Sander. 2017. "Mitochondrial Respiratory Gene Expression Is Suppressed in Many Cancers." *eLife* 6 (January). https://doi.org/10.7554/eLife.21592.

Riesgo, Ana, Nathan Farrar, Pamela J. Windsor, Gonzalo Giribet, and Sally P. Leys. 2014. "The Analysis of Eight Transcriptomes from All Poriferan Classes Reveals Surprising Genetic Complexity in Sponges." *Molecular Biology and Evolution* 31 (5): 1102–20.

RStudio Team. 2020. *RStudio: Integrated Development for R* (version 4.0.3). X86_64, darwin17.0 . https://www.rstudio.com/products/rstudio/.

Sassaman, Clay, and Charlotte P. Mangum. 1972. "ADAPTATIONS TO ENVIRONMENTAL OXYGEN LEVELS IN INFAUNAL AND EPIFAUNAL SEA ANEMONES." *The Biological Bulletin* 143 (3): 657–78.

Schröder, H. C., S. M. Efremova, B. A. Margulis, I. V. Guzhova, V. B. Itskovich, and W. E. G. Müller. 2006. "Stress Response in Baikalian Sponges Exposed to Pollutants." *Hydrobiologia* 568 (1): 277–87.

Schuster, Astrid, Brian William Strehlow, Lisa Eckford-Soper, Rob McAllen, and Donald Eugene Canfield. 2021. "The Effects of Seasonal Anoxia on the Microbial Community Structure in Demosponges in a Marine Lake (Lough Hyne, Ireland)." *mSphere* 6: e00991–20.

Simão, Felipe A., Robert M. Waterhouse, Panagiotis Ioannidis, Evgenia V. Kriventseva, and Evgeny M. Zdobnov. 2015. "BUSCO: Assessing Genome Assembly and Annotation Completeness with Single-Copy Orthologs." *Bioinformatics* 31 (19): 3210–12.

Sinha, Amit, Ralf J. Sommer, and Christoph Dieterich. 2012. "Divergent Gene Expression in the Conserved Dauer Stage of the Nematodes Pristionchus Pacificus and Caenorhabditis Elegans." *BMC Genomics*. https://doi.org/10.1186/1471-2164-13-254.

Sollai, Martina, Laura Villanueva, Ellen C. Hopmans, Gert-Jan Reichart, and Jaap S. Sinninghe Damsté. 2019. "A Combined Lipidomic and 16S rRNA Gene Amplicon Sequencing Approach Reveals Archaeal Sources of Intact Polar Lipids in the Stratified Black Sea Water Column." *Geobiology* 17 (1): 91–109.

Stachowitsch, Michael. 1984. "Mass Mortality in the Gulf of Trieste: The Course of Community Destruction." *Marine Ecology* 5 (3): 243–64.

Strader, Marie E., Galina V. Aglyamova, and Mikhail V. Matz. 2016. "Red Fluorescence in Coral Larvae Is Associated with a Diapause-like State." *Molecular Ecology* 25 (2): 559–69.

Stramma, Lothar, Gregory C. Johnson, Janet Sprintall, and Volker Mohrholz. 2008. "Expanding Oxygen-Minimum Zones in the Tropical Oceans." *Science* 320 (5876): 655–58.

Strehlow, Brian W., Mari-Carmen Pineda, Carly D. Kenkel, Patrick Laffy, Alan Duckworth, Michael Renton, Peta L. Clode, and Nicole S. Webster. 2020. "Novel Reference Transcriptomes for the Sponges Carteriospongia Foliascens and Cliona Orientalis and Associated Algal Symbiont Gerakladium Endoclionum." *bioRxiv*. bioRxiv. https://doi.org/10.1101/2020.06.26.156463.

Vaquer-Sunyer, Raquel, and Carlos M. Duarte. 2011. "Temperature Effects on Oxygen Thresholds for Hypoxia in Marine Benthic Organisms." *Global Change Biology*. https://doi.org/10.1111/j.1365-2486.2010.02343.x.

Vargas, Sergio, Cüneyt Caglar, Büttner Gabi, Schätzle Simone, Deistler Fabian, and Gert Wörheide. 2021. "Slime Away: A Simple CTAB-Based High Molecular Weight DNA and RNA Extraction Protocol for 'Difficult' Invertebrates." Protocols.io. 2021. https://www.protocols.io/view/slime-away-a-simple-ctab-based-high-molecular-weig-bwcwpaxe.

Webster, Nicole S., and Michael W. Taylor. 2012. "Marine Sponges and Their Microbial Symbionts: Love and Other Relationships." *Environmental Microbiology* 14 (2): 335–46.

Webster, N., R. Pantile, E. Botté, D. Abdo, N. Andreakis, and S. Whalan. 2013. "A Complex Life Cycle in a Warming Planet: Gene Expression in Thermally Stressed Sponges." *Molecular Ecology* 22 (7): 1854–68.

Wilkinson, C. R. 1978. "Microbial Associations in Sponges. II. Numerical Analysis of Sponge and Water Bacterial Populations." *Marine Biology* 49 (2): 169–76.

Wilson, D. F., M. Erecińska, C. Drown, and I. A. Silver. 1979. "The Oxygen Dependence of Cellular Energy Metabolism." *Archives of Biochemistry and Biophysics* 195 (2): 485–93.

Wright, Rachel M., Galina V. Aglyamova, Eli Meyer, and Mikhail V. Matz. 2015. "Gene Expression Associated with White Syndromes in a Reef Building Coral, Acropora Hyacinthus." *BMC Genomics*. https://doi.org/10.1186/s12864-015-1540-2.

Bergquist, Patricia Rose. 1968. "The Marine Fauna of New Zealand: Porifera, Demospongiae, Part 1 (Tetractinomorpha and Lithistida)". *New Zealand Oceanographic Institute Memoir.* 37:27–28.

Bosc, Louis A. G. 1802. "Histoire Naturelle des Crustacés Contenant leur Description et leurs Moeurs; avec Figures Dessinées, d'Après Nature". Paris: Chez Deterville. 2: 1-296, pls. 6-18., available online at https://www.biodiversitylibrary.org/page/27817320. page(s): 221–222.

Bowerbank, J.S. 1864. "A Monograph of the British Spongiadae".Volume 1. Ray Society: London: i-xx, 1-290, pls I-XXXVII., available online at https://biodiversitylibrary.org/page/1883085.

Dendy, Arthur. 1905. "Report on the sponges collected by Professor Herdman, at Ceylon, in 1902". In: Herdman, W.A. (Ed.), Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar. 3 (Supplement 18). Royal Society: London. Pp. 57-246, pls I-XVI. page(s): 143.

Grant, R.E. 1836. Animal Kingdom. Pp. 107-118. In: Todd, R.B. (Ed.), "The Cyclopaedia of Anatomy and Physiology". Volume 1. Sherwood,Gilbert, and Piper: London: 1-813.

Grant, R.E. 1826. "Notice of a New Zoophyte (Cliona celata Gr.) from the Firth of Forth". Edinburgh *New Philosophical Journal* 1: 78-81., available online at https://www.biodiversitylibrary.org/page/2471191. page(s): 81.

Johnston, Georg. 1842. "A History of British Sponges and Lithophytes". (W.H. Lizars: Edinburgh). i-xii, 1-264, pls I-XXV., available online at https://biodiversitylibrary.org/page/35290582. page(s): 146-147.

Kelly-Borges, Michelle., Patricia Rose Bergquist. 1997. "Revision of Southwest Pacific Polymastiidae (Porifera: Demospongiae: Hadromerida) with descriptions of new species of Polymastia Bowerbank, Tylexocladus Topsent, and Acanthopolymastia gen. nov. from New Zealand and the Norfolk Ridge, New Caledonia". *Zealand Journal of Marine and Freshwater Research.* 31 (3): 378–380.

Lévi, Caude. 1953. "Sur une nouvelle classification des Démosponges". *Compte rendu hebdomadaire des séances de l'Académie des sciences*, Paris. 236 (8): 853–855.

Linnaeus, Claude. 1758. "Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata [10th revised edition], vol. 1: 824 pp. Laurentius Salvius: Holmiae. , available online at https://biodiversitylibrary.org/page/726886. page(s): 761.

Magnet S, Dubost L, Marie A, Arthur M, Gutmann L. 2008. "Identification of the L,D-transpeptidases for peptidoglycan cross-linking in Escherichia coli". *J Bacteriol.* 190(13):4782-5. doi: 10.1128/JB.00025-08.

Pallas, Peter S. 1766. "Elenchus zoophytorum sistens generum adumbrationes generaliores et specierum cognitarum succintas descriptiones, cum selectis auctorum synonymis". Fransiscum Varrentrapp, Hagae. pp. 451., available online at https://doi.org/10.5962, page(s): 388.

Schmidt, Oscar. 1870. "Grundzüge einer Spongien-Fauna des atlantischen Gebietes". Wilhelm

Engelmann: Leipzig: iii-iv, 1-88, pls I-VI. page(s): 14-15; pl I fig 1-6.

Stryer, Lubert. 1995. "Biochemistry (Fourth ed.)". New York: W.H. Freeman and Company. pp.
559–565, 574–576, 614–623. ISBN 0-7167-2009-4.

Topsent, Emil. 1928. "Spongiaires de l'Atlantique et de la Méditerranée provenant des croisières
du Prince Albert ler de Monaco". *Résultats des campagnes scientifiques accomplies par le*
*Prince Albert I. Monaco.* 74:1-376, pls I-XI.

Welchen E, Gonzalez DH. 2016.  "Cytochrome c, a hub linking energy, redox, stress and
signaling pathways in mitochondria and other cell compartments". *Physiol Plant.*157(3):310-
21. doi: 10.1111.

1405