1
2
3

Transcriptomic responses of sponge holobionts to in situ, seasonal anoxia and hypoxia

Brian W. Strehlow[ab*], Astrid Schuster[a*], Warren R. Francis[a], Lisa Eckford-Soper[a], Beate Kraft[a],
Rob McAllen[c], Ronni Nielsen[d], Susanne Mandrup[d], Donald E. Canfield[a]

[a] Nordcee, Department of Biology, University of Southern Denmark, Odense M, Denmark.
[b] Littoral Environnement et Sociétés (LIENSs), UMR 7266 CNRS - La Rochelle Université, 2
Rue Olympe de Gouges, 17000, La Rochelle, France.
[c] School of Biological, Earth and Environmental Sciences, University College Cork, Cork,
Ireland.
[d] Functional Genomics and Metabolism Research Unit, Department of Biochemistry and
Molecular Biology, University of Southern Denmark, Odense, Denmark

18
19 ***Denotes co-first author. Author order was decided based on reverse alphabetical order, since the**
20 **inverse was used on a previous paper. Both authors have the right to list their name first on their**
21 **CVs.**

Abstract

Deoxygenation can be fatal for many marine animals; however, some sponge species are tolerant of hypoxia and anoxia. Indeed, two sponge species, *Eurypon* sp. 2 and *Hymeraphia stellifera,* survive seasonal anoxia for months at a time. To understand their tolerance mechanisms, we performed differential gene expression analyses on the sponges, their mitochondria and their microbial symbionts under in situ conditions of normoxia, hypoxia and anoxia. Each species possessed a unique microbiome, but the microbiomes of each species were dominated by a species-specific *Thaumarchaeon* and a *Gammaproteobacterium*. Holobiont gene expression was species- and oxygen-level dependent, though there were some shared interspecific responses to deoxygenation. In general, few changes occurred in the expression of sponge metabolic genes as a function of oxygenation level, indicating that they may remain metabolically active under anoxia. However, ATP synthesis genes were significantly upregulated under hypoxia when compared to normoxia, and genes for DNA replication were downregulated. Mitochondrial gene expression was effectively unchanged under both hypoxia and anoxia. Nevertheless, both anoxia and hypoxia caused upregulation of heat shock proteins (HSPs), indicating cellular level adaptations to deoxygenation stress. A meta-analysis demonstrated that sponge transcriptional responses to anoxia were distinct from those displayed by other invertebrates while dormant, and the hypothesis of sponge dormancy under anoxia was not supported. *Thaumarchaeota* symbionts also upregulated stress response genes in hypoxia, while maintaining expression of oxygen-dependent metabolic pathways under hypoxia and anoxia. *Gammaproteobacteria* symbionts showed relatively few noteworthy changes in gene expression in response to anoxia but decreased metabolic gene expression in hypoxia. There was no clear evidence of upregulated anaerobic respiration in the transcriptomes of the sponge holobionts under anoxia or hypoxia. The tolerance of some sponges to prolonged anoxia warrants further investigation and could give them an advantage in future oceans following climate change as well as in ancient oceans when oxygen concentrations were lower than at present.

Introduction

Ocean oxygen levels are predicted to decline by 1–7% over the next century (Keeling, Körtzinger, and Gruber 2010). Climate change is expanding marine oxygen-minimum zones, while enhanced nutrient runoff is causing coastal eutrophication and subsequent deoxygenation (Kirchman 2021). Deoxygenation is expected to negatively affect the physiology, metabolism and survival of many marine organisms (Diaz and Rosenberg 2008). As a result of deoxygenation and other anthropogenic stressors, we are facing large scale habitat and biodiversity losses, but the full effects of these stressors are still unknown.

Sessile organisms that cannot move or migrate in response to deoxygenation are particularly vulnerable. Sponges (Phylum: Porifera[1]), sessile, ecologically important filter-feeding animals, are hypothesized to suffer from increasing hypoxic/anoxic areas worldwide (Breitburg et al. 2018; Levin 2018; Stramma et al. 2008), and some in situ mortality of sponges (along with other taxa) following deoxygenation has been reported (Altieri et al. 2017; Stachowitsch 1984). However, species-specific responses to deoxygenation in Porifera are variable, and some species are tolerant of hypoxia. Lab experiments have shown that the demosponge *Halichondria (Halichondria) panicea*[2] can survive low oxygen levels down to 0.5–4% of air saturation (a.s.) (Mills et al. 2014), and *Tethya wilhelma* can survive down to 0.25–2% a.s. for 4 days and <0.05% a.s. for at least one hour (Mills et al. 2018). Additional demosponge species that tolerate similar levels of hypoxia include *Geodia barretti*[3] with 7% a.s. (20 µM O2, Leys and Kahn 2018) and *Haliclona pigmentifera*[4] (0.50-0.68% a.s., Gunda and Janapala 2009). In a recent experimental study, four taxonomically distinct sponge species (*Cliona celata*[5], *Suberites carnosus*[6], *Polymastia crocea*[7] and *Suberites australiensis*[8]) were exposed to various levels of hypoxia (1.5–40% a.s., 0.13–3.3 mg O_2 L^{-1}) in aquaria. These sponges exhibited species-specific physiological, morphological and behavioral responses to deoxygenation. Sub-lethal oxygen thresholds for these sponges were in the range of 6–20% a.s. (0.48–1.56 mg O_2 L^{-1}) while mortality was only observed in one of the two species (*P. crocea*) exposed to severe hypoxia lower than 5% a.s. (0.13 mg O_2 L^{-1}) (Micaroni et al. 2021a). The sponge species most sensitive to deoxygenation in controlled experiments, *P. crocea*, had a median lethal time (time until death, LT50) of 286 h at ~1.5% a.s. (0.13 mg O_2 L^{-1}, (Micaroni et al. 2021a), which was still longer than most marine organisms under comparable conditions (reviewed in Micaroni et al. 2021a and Hughes et al. 2020).

Few studies have investigated the deoxygenation tolerance of sponges in situ. Bayer et al. (2020) found that the glass sponge *Vazella pourtalesii*[9] lives in reduced oxygen conditions (< 4 mL/L) on the Scotian Shelf, Canada. Schuster et al. (2021) reported at least nine encrusting

demosponge species of the orders Axinellida[10] and Poecilosclerida[11] that tolerate seasonal
anoxia in a Marine Lake in Lough Hyne, Ireland for ~three months at a time in the summer, and
there are more sponge species at this site (Bell and Barnes 2000). The anoxia-tolerant sponges
of Lough Hyne therefore have LT50s that exceed their ca. three month exposure to anoxia, and,
compared to other animals, their tolerance is rivaled only by the anemone *Bunodosoma*
cavernatum[12] (LT50 in anoxia >1.5 months, Ellington 1982). Moreover, the abundance (percent
cover) of encrusting sponges in the seasonally anoxic site of Lough Hyne only slightly
decreased over 30 years despite widespread sponge losses in the well oxygenated parts of the
lough (Micaroni et al. 2021b), indicating that these sponges are uniquely suited to their
seasonally anoxic niche.
96
Many invertebrates exposed to harsh conditions like anoxia or hypoxia have evolved molecular
response pathways to survive in these environments. For instance, under anoxia, marine
mollusks shift from lipid, carbohydrate and amino acid metabolism to carbohydrate catabolism
as the primary fuel for ATP generation (Larade and Storey 2009). Some mollusks also maximize
anoxic survival time (mollusk LT50s at 0 mg L^{-1}: 360–672 h, see Micaroni, et al. 2021b) by
building up a fixed reserve of internal fuels (primarily glycogen) to increase ATP output and to
reduce acidity. A small suite of upregulated genes also plays a critical role in cell preservation,
facilitating long-term viability under hypoxia (Larade and Storey 2009). Some anemones may
even have elements of a facultative anaerobic metabolism (Sassaman and Mangum 1972).
106
Only one study has investigated the transcriptomic response of a sponge to hypoxia. After four
days of exposure to hypoxia (0.25–2% a.s.), 128 genes within the sponge *T. wilhelma* were
differentially expressed compared to controls, but none of these genes were clearly related to
metabolic or stress responses (Mills et al. 2018). However, one hour exposure to near anoxia
(<0.05% a.s., 105 nM O_2) elicited stress responses in *T. wilhelma's* transcriptome (Mills et al.
2018), demonstrating that transcriptomic responses were mostly observed under very low
oxygen levels.
114
In most metazoans, the hypoxia-inducible factor (HIF) pathway is essential for maintaining
oxygen homeostasis in response to hypoxia (Samenza 2007). However, in sponges and
ctenophores, the likely sister-lineages to other metazoans (Feuda et al. 2017), key components
of the HIF pathway, including orthologs for HIFa and HIFb, are absent (Mills et al. 2018). HIFa is
a transcription factor responsible for triggering metabolic adaptation to deoxygenation in most
metazoans (Samenza 2007). Due to the absence of HIFa in sponges, the mechanisms of
tolerance to hypoxia and anoxia are largely unknown (Mills et al. 2018).
122
123 The sponge microbiome may play a role in sponge deoxygenation tolerance. Sponges can host
124 diverse microbiomes that perform vital functions and can constitute large portions of their
125 biomass (Wilkinson 1978). The host and microbiome together are referred to as a 'holobiont'
126 (Webster and Taylor 2012). Some sponge symbionts have anaerobic capabilities, including
127 sulfate reduction (Hoffmann et al. 2005), denitrification, and anaerobic ammonium oxidation

---

Taxonomic authority citations
[10] Lévi, 1953
[11] Topsent, 1928
[12] Bosc, 1802

128 (anammox) (Hoffmann et al. 2009). In *V. pourtalesii*, the microbial community is thought to be
129 specialized for the low-oxygen conditions in which it lives (Bayer et al. 2020).
130
A previous study has shown that the microbiomes of anoxia-tolerant sponges from Lough Hyne
were unchanged under anoxia, hypoxia and normoxia despite drastic changes in the microbial
communities of the water and sediment, indicating that the holobiont as a whole is tolerant to
deoxygenation (Schuster et al. 2021). Specifically, the microbiomes of the anoxia-tolerant
sponge species *Eurypon* sp. 2 and *Hymeraphia stellifera*[13] were both dominated by two
microbes – one *Thaumarchaeon* and one *Gammaproteobacterium*, though each sponge hosted
a different species from each clade. *Eurypon* sp. 2 harbors the *Thaumarchean* OTU1 (closest
related isolate: *Nitrosopumilus adriaticus* strain NF5), and *H. stellifera* harbors OTU3 (closest
related isolate: *Nitrosopumilus oxyclinae* strain HCE1). For the *Gammaproteobacteria*, *Eurypon*
sp. 2 contains OTU2 (closest related isolate: *Arenicella chitinivorans* strain ALEN 1); whereas *H.*
stellifera contains OTU7 (closest related isolate: *A. chitinivorans* strain ALEN 1, (Schuster et al.
2021). Although their location within Lough Hyne sponges has not yet been determined,
Gammaproteobacteria and *Thaumarchaeota* symbionts are abundant and widely distributed
throughout the sponge tissue in other species (Moeller et al. 2019). These symbionts also
participate in integrated metabolic processes with their host species (Moitinho-Silva et al. 2017).
146
Schuster et al. (2021) hypothesized potential metabolic pathways within these two symbiont
taxa that might function under deoxygenation. Based on the genomes of their close relatives,
the *Thaumarchaeota* in these sponges were hypothesized to have components of a facultatively
anaerobic metabolism, including fumarate, nitrite, and sulfite respiration and the ability to
ferment (Bayer et al. 2020), which could aid in deoxygenation tolerance. A recent study
demonstrated that another close *Thaumarchaeota* relative, *Nitrosopumilus maritimus*, can
produce trace levels of oxygen (100–200 nM) under environmental anoxia (Kraft et al. 2022).
The pathway of oxygen production and the genes involved are not currently known. However,
comparative proteomics suggested that proteins upregulated during oxygen production included
some AsnC family proteins and nitroreductases (Kraft et al. 2021). We therefore hypothesized
that nitroreductases and AsnC family proteins would be upregulated under hypoxia and anoxia
in sponge-associated *Thaumarchaeota. Gammaproteobacteria* symbionts were hypothesized to
use sulfide or thiosulfate as an electron donor anaerobically coupled to nitrate reduction,
removing exogenous, toxic sulfide from the holobiont under anoxia (Schuster et al. 2021).
161
To understand the mechanisms of sponge holobiont tolerance and survival under extended
anoxia, we performed differential gene expression analysis of the sponges *Eurypon* sp. 2 and *H.*
stellifera as well as their respective mitochondria and microbial symbionts including
Thaumarchaeota and *Gammaproteobacteria*. Sponges were collected from Lough Hyne under
in situ normoxia, hypoxia and anoxia. Since neither of these holobionts are established model
species, we generated reference transcriptomes for the sponges and assembled the genomes
of their symbionts and mitochondria from metagenomic sequences. Additionally, we performed
a meta-analysis of the sponge differential expression data to determine if the responses of
Lough Hyne sponges to deoxygenation are similar to those of *T. wilhelma* under deoxygenation,

---

Taxonomic authority citations
[13] Bowerbank, 1864

171   other invertebrates under heat stress (Dixon et al. 2015), in a dauer state (Sinha et al. 2012),
172   and during diapause (Gong et al. 2013). Finally, we considered the implications of sponge
173   deoxygenation tolerance for future oceans as well as early animal evolution under low
174   environmental oxygen concentrations.
175
176   **Materials and Methods**
177
Sampling
Sponges were sampled between July 2018 and August 2019 at Labhra Cliff (Lough Hyne
Nature Reserve, Ireland; N51°30.05309 W9°18.17679) as described in Schuster et al. (2021)
under permit no. R23-27/2018 issued by the Irish Department of Environment, Heritage, and
Local Governments. Dissolved oxygen profiles with depth were determined using a Pro20
dissolved oxygen meter (YSI, USA). Profiles for different sampling times are shown in Schuster
et al. (2021). Sponge samples were taken using SCUBA (depth range: 15-29 m) under various
in situ oxygen conditions that were categorized as follows: normoxic (5.3–12 mg L^{-1}, 49.3–111%
a.s.), hypoxic (1.30–3.56 mg L^{-1}, 12.1–33.1% a.s.), and anoxic (0.00–0.01 mg L^{-1}, 0.00–0.93%
a.s., i.e. instrument detection limit). Oxygen concentration at each sampling point was verified
using a HOBO dissolved oxygen logger (U26-001; Onset, USA). All samples were frozen in
liquid nitrogen within ~15 minutes of collection and kept in a dry shipper until returned to the
laboratory where samples were stored in a -80°C freezer until RNA and DNA extraction.
191
In total 70 sponge samples were taken under anoxia (n = 6), hypoxia (n = 25), and normoxia (n
= 39). Although all these red-orange encrusting sponges were indistinguishable in situ, species
identification using molecular barcodes revealed that there were nine species (see Schuster et
al. 2021). This level of cryptic diversity is common in sponges, and necessitates care in post-
sampling species identification. Two species, *Eurypon* sp. 2 and *H. stellifera,* comprised the
majority of the samples and were represented across the oxygen conditions; therefore, these
two species were selected for the transcriptomic analysis of sponge, symbionts, and
mitochondria in this study. For RNA sequencing (RNAseq), eight samples of *H. stellifera* (anoxic
= 2, hypoxic = 5, normoxic = 1) were taken, and sixteen samples of *Eurypon* sp. 2 (anoxic = 2,
hypoxic = 7, normoxic = 7) were sequenced (n = 24 in total). However, certain oxygen levels
within each species, i.e. anoxia in both species and normoxia in *H. stellifera*, had limited
replicates (n < 3). The implications of this limited replication are considered in the analysis
methods and discussion. Metadata for individual samples, including oxygen concentration,
season, depth, collection date, and individual sample code (e.g. DC##) are included in
Supplemental Table 1.
207
208   **DNA/RNA extraction**
209   Total RNA was isolated using TRIzol^{TM} Reagent from Invitrogen (ThermoFisher Scientific,
210   Denmark), and remaining DNA in the RNA extracts were removed using the DNase MAX kit
211   (Qiagen), following the manufacturers' protocols.
212
Additionally, high molecular weight (HMW) DNA was extracted from one individual of each
species using the protocol from Vargas et al. (2021) for metagenomic sequencing. The *Eurypon*
sp. 2 sample used for metagenomic sequencing was taken under anoxia (sample DC24), and

RNA was also sequenced from this sample. Due to limited availability of samples and DNA, a
sample of *H. stellifera* that was not included in the RNAseq (DC297, sampled under normoxia)
was used for metagenome sequencing. DNA and RNA yields were determined using a Qubit^{TM}
(Thermo Fisher Scientific) and a NanoDrop^{TM} spectrophotometer (Thermo Fisher Scientific).
RNA quality was evaluated on an Agilent TapeStation using an RNA ScreenTape (Agilent
Technologies).
222
Reference transcriptomes and metatranscriptomes
All RNA samples were sequenced twice with two different library preps - one for polyadenylated
(poly A) messenger RNA (mRNA, from eukaryotes) and one for total RNA (to also capture
prokaryotic RNA). Library preparation for the eukaryotic (sponge) RNA was done using
NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina (New England Biolabs, USA).
For the prokaryotic RNAseq, the supernatant from the poly A enrichment was treated with
NEBNext® rRNA Depletion Kit (Bacteria) and then NEBNext® rRNA Depletion Kit
(Human/Mouse/Rat, New England Biolabs, USA) to eliminate ribosomal RNA (rRNA) that
causes sequence biases. This combination of poly A exclusion and rRNA depletion eliminated
the most rRNA compared to other combinations of methods (or single methods in isolation) in
preliminary tests, yielding sequences with only ~25% rRNA. For the eukaryotic (n = 24) and
prokaryotic (n = 24) RNA samples, library preparation was completed using the NEBNext Ultra II
Directional RNA library preparation kit (New England Biolabs, Denmark). Libraries pooled in
equimolar concentrations were sequenced (5M reads per sample, 2 x 50 bp) at the Center for
Functional Genomics and Tissue Plasticity (ATLAS) at the University of Southern Denmark
(SDU), using a NovaSeq 6000 (Illumina, USA).
239
To ensure good coverage for the final reference transcriptomes, two samples from each species
were sequenced deeper in additional runs. For *H. stellifera*, the sample for deeper sequencing
(DC129, sampled under normoxia) was also present in the sample set above. In contrast, for
Eurypon sp. 2 another sample (DC128, sampled under normoxia) was used due to the limited
availability of sample tissue. As in the sequencing above, two libraries were created and
sequenced for each sample - one with eukaryotic RNA and one with prokaryotic RNA. For this
deeper sequencing, mRNA in the sponge transcriptomes was purified using the NucleoTrap
mRNA kit (Macherey-Nagel, Germany). Prokaryotic RNA was rRNA depleted using the Ribo-
zero Magnetic kit (Illumina, USA). An additional purification step was performed for all RNA
samples using the standard protocol for the CleanNGS kit (CleanNA, Netherlands) before
sequencing preparation using the NEBNext Ultra II Directional RNA library preparation kit (New
England Biolabs, Denmark). Samples were diluted in equimolar concentrations, pooled and
sequenced (2 x 150 bp) on a HiSeq X platform (Illumina, USA) by DNASense ApS (Aalborg,
Denmark).
254
For the eukaryotic RNA, shallow (2 x 50 bp) sequencing generated 9.20×10^8 and 4.38×10^8
reads for *Eurypon* sp. 2 and *H. stellifera*, respectively, across all sampled sponges, and deeper
sequences (2 x 150 bp) generated 2.05×10^8 and 3.45×10^8 reads for each sponge of the two
species, respectively. For the prokaryotic RNA, shallow sequencing yielded 1.27×10^9 and 6.68
$\times 10^8$ reads for *Eurypon* sp. 2 and *H. stellifera*, respectively. The deeper prokaryotic sequencing

generated 1.52 x 10^8 and 1.20 x 10^8 for *Eurypon* sp. 2 and *H. stellifera*, respectively (totals are
shown in Supplemental Table 2).
262
To obtain reference transcriptomes for each sponge species without prokaryotic symbionts, all
eukaryotic RNA sequences were combined into a single file. These sequences were trimmed for
quality control using public scripts (Matz 2015; Meyer 2016) and assembled with Trinity v 2.8.5
(Grabherr et al. 2011). This assembly was filtered for contaminants and annotated using egg-
NOG mapper (Huerta-Cepas et al. 2019). The final sponge transcriptomes included only genes
with metazoan annotations or no annotations. For a detailed description of all filtering and
assembly steps, see Strehlow et al. (2020) and (https://zenodo.org/badge/latestdoi/573040118)
for all scripts used in the current paper. Sponge reference transcriptomes fit within the expected
range of size, gene number, and N50 (Supplemental Table 2) of published sponge
transcriptomes (Strehlow et al. 2020; Riesgo et al. 2014). Additionally, sponge reference
transcriptomes were effectively complete based on Benchmarking Universal Single-Copy
Orthologs (BUSCO) analysis (Supplemental Table 2, Simão et al. 2015) and were therefore
suitable for use for subsequent differential expression analysis (see below). Sponge reference
transcriptome assemblies are available through NCBI under BioProject number PRJNA89319.
277
Reference Metagenomes
279 Two reference metagenomes were sequenced for *Eurypon* sp. 2 and *H. stellifera* at DNASense
280 ApS (Aalborg, Denmark). DNA was fragmented to ~550 bp (M220 Focused-ultrasonicator,
281 Covaris, USA) and used for library preparation following the NEBNext Ultra II DNA library
282 preparation kit. The DNA library was paired-end sequenced (2 x 150 bp) on a NovaSeq platform
283 (Ilumina, USA).
284
Sequences were assembled using Megahit v.1.2.9 (Li et al. 2016). The final contig fasta files
were used to plot GC (%) vs. mean coverage of mapped reads to identify individual bins. For
this, we used the custom python scripts spadescontigstocovgc.py (Francis 2020a) as well as the
R script contig_gc_coverage.R (Francis 2020b) to extract bins for the *Thaumarchaeota* and
Gammaproteobacteria symbiont of each sponge species. Final statistics for each genome are
shown in Supplemental Table 2. Raw sequences are available from NCBI under BioProject
number PRJNA893197. Metagenome assemblies are available on
https://github.com/bstrehlow/Sponge-transcriptomic-responses-to-seasonal-anoxia.
293
Mitogenomes
295 Using only the deep sequences of total ('prokaryotic') RNA, the mitochondrial genomes of single
296 individuals of *Eurypon* sp. 2 (DC128) and *H. stellifera* (DC129) were assembled following the
297 pipeline in Plese et al. (2019).
298
For future utilization, single contigs that likely represent the mitochondrial genomes of *Eurypon*
sp. 2 and *Hymeraphia stellifera* were retrieved from the reference metagenome in addition to
the coding regions used for differential expression below. The assembled contigs for each
metagenome were blasted against the mitochondrial genome of DC128 (*Eurypon* sp. 2) using
blastn, with an e-value cutoff of 1x10^{-50} for *Eurypon* sp. 2 (DC97) and 1x10^{-30} for *Hymeraphia*
stellifera (DC297). In general, there was only a single hit for each protein coding gene,

confirming a single contig as the mitochondrial genome for each sample. Based on the blast,
these two contigs contained all protein coding genes retrieved from the transcriptome that were
the focus of this analysis. These 'mitogenomes' can be accessed here:
https://zenodo.org/badge/latestdoi/573040118.
309
Differential Expression Analysis
Trimmed reads from the shallow sequencing for each individual were mapped back to their
respective reference transcriptome or genome using bowtie2 (version 2.3.5.1, Langmead and
Salzberg 2012) and mapped reads were counted for each contig using samcount.pl (see Matz
2015). Eukaryotic reads were mapped to the sponge reference transcriptome for each species.
Prokaryotic reads were mapped to their respective metagenomic bins - metagenome assembled
genomes (MAGs), i.e. *Thaumarchaeota* or *Gammaproteobacteria*, within each sponge species.
The mitochondrial reference was broken into its 14 genes and prokaryotic (total RNA) reads
from both species were mapped to their matching mitochondrial reference and counted as
above.
320
Analyses were performed to compare the gene expression of each target (sponge species,
symbiont species, and mitochondria; n=8) under the three different oxygen conditions (anoxia,
hypoxia and normoxia). Differential gene expression analysis was performed with the package
DESeq2 v. 1.22.2 (Love et al. 2016) in R (R Core Team 2016) with (RStudio Team 2020). Three
pairwise comparisons of oxygen condition (normoxia vs. hypoxia, normoxia vs. anoxia, and
anoxia vs. hypoxia) were computed from DESeq2 results. Significantly differentially expressed
genes between each of these pairwise comparisons were identified using a false discovery rate
(FDR) adjusted p-value threshold of 0.1. The ReportingTools software package (Huntley et al.
2013) was used to create PDF pages displaying 4 plots of significantly differentially expressed
genes and their annotations in various oxygen conditions, as well as box plots, p-values, and
LFCs (https://zenodo.org/badge/latestdoi/573040118). These files are searchable in case
particular genes are of interest to future studies. We note that comparisons to *H. stellifera* under
normoxia were made with only one normoxic sample due to sampling restrictions. Despite this
sampling limitation, the normoxic data for *H. stellifera* are still shown as a baseline and
reference. For each sponge species, log fold changes (LFC, from DESeq2) in all genes were
calculated so that positive LFCs were upregulated and negative values were downregulated in
hypoxia or anoxia relative to normoxia. For the comparisons between hypoxia and anoxia,
positive LFCs were upregulated in anoxia and vice versa.
339
340
KOG analysis
EuKaryotic Orthologous Groups (KOG class) assignments were used to compare large scale
trends and potential correlations in differential gene expression between Lough Hyne sponges
under varied oxygen conditions and external gene expression data from other species and
conditions. Delta-ranks, a measure of each KOG class enrichment with up or downregulated
genes, were calculated from LFC data using the KOGMWU package in R (Dixon et al. 2015),
and delta-ranks were plotted using the package pheatmap (Kolde 2013). While it is also
possible to calculate delta-ranks with signed p-values (see KOGMWU supporting
documentation), LFC was used here to include changes between normoxia and other conditions

for *H. stellifera*. Since there was only one normoxic replicant for this species, LFC was a more
appropriate metric than signed p values. Within each pairwise comparison (i.e. hypoxic, anoxic,
and anoxic vs hypoxic), significantly enriched KOGs were also calculated using KOGMWU
(FDR-adjusted $p < 0.05$).

These KOG enrichments were also compared to those calculated from the sponge *Tethya*
wilhelma after four days of exposure to hypoxia and one hour of exposure to anoxia (Mills et al.
2018). To test whether KOG enrichment in sponges exposed to variable oxygen conditions was
correlated with gene expression in other invertebrate stress responses, delta rank data was
included from adult *Acropora millepora* coral exposed to heat stress (31.5 °C for three days,
(Dixon et al. 2015). Similarly, to test the similarity of the sponge data to noted decreases in
metabolic activity in invertebrates, KOGs enriched in the dauer (dormant) state in
Caenorhabditis elegans (Sinha et al. 2012) and the diapause state of the midge *Sitodiplosis*
mosellana (Gong et al. 2013) were added using delta rank tables calculated by Strader et al.
(2016). Correlations between the KOG delta ranks in different datasets were performed using
Pearson's Correlation in KOGMWU. Within the significantly enriched KOGs 'replication,
recombination and repair' and 'energy production and conversion,' genes that were significantly
differentially expressed were plotted in heatmaps to further investigate sponge transcriptomic
responses to deoxygenation at the gene level in Lough Hyne sponges.

KOG enrichment analysis was performed as above for each of the two major symbionts within
the two sponge species; however, Clusters of Orthologous groups (COGs) were used since
they are designed for prokaryotes rather than eukaryotic KOGs and are fewer in number.

Gene ontology (GO) term enrichment analysis
Log fold changes calculated above were used for rank-based gene ontology (GO) analysis with
adaptive clustering (GO_MWU, Wright et al. 2015) to determine what GO terms were
significantly enriched under various oxygen conditions. These analyses were run individually for
each pairwise comparison of oxygen condition in both sponge species. These analyses were
not completed for the two symbiont types because GO terms were not available for these
microbes.

Phylogenetic positions of nitroreductase proteins
To determine if nitroreductases in sponge symbionts are related to genes that may be involved
in oxygen production in *N. maritimus,* nitroreductase proteins from the *Thaumarchaeota*
symbionts of *H. stellifera* and *Eurypon* sp. 2 were extracted using annotations assigned by egg-
NOG. Phylogenetic position of nitroreductase proteins from *H. stellifera* and *Eurypon* sp. 2 were
investigated by aligning them to other known nitroreductase proteins from Genbank using Mafft
v7 (Katoh and Standley 2013). The ModelTest-NG v.0.1.7 function (Edler et al. 2020)
implemented within RaxmlGUI 2.0 was used to calculate the evolutionary model. According to
AIC, LG+I+G was chosen as the best-fitting model. Maximum Likelihood (ML) phylogeny was
then reconstructed using raxmlGUI v.2.0.6 (Edler et al. 2020) with 100 (rapid) bootstraps. The
resulting protein tree was rendered by FigTree v.1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

Results

Sponge differential expression

There were three pairwise comparisons for significantly differentially expressed genes - hypoxia vs. normoxia, anoxia vs. normoxia, and anoxia vs. hypoxia. The number of significantly differentially expressed genes in these comparisons for sponges, their mitochondria and their symbionts are summarized in Table 1. In the sponge *Eurypon* sp. 2, 737, 1,376 and 1,090 genes were significantly differentially expressed under hypoxia (vs. normoxia), anoxia (vs. normoxia), and anoxia vs. hypoxia, respectively ($p < 0.1$, Table 1). The differentially expressed genes under each oxygen condition ranged between 1.5–2.8% of the genes of the *Eurypon* sp. 2 reference transcriptome, and 41-48% of these genes are annotated, which reflects the overall annotation percentage of 39.7% for the whole transcriptome (Supplemental Table 2). In hypoxia, the majority of these differentially expressed genes were upregulated (68%). Under anoxia, on the other hand, the majority of genes were downregulated (60%). In anoxia vs. hypoxia, 57% of genes were upregulated in anoxia compared to hypoxia in *Eurypon* sp. 2 (Table 1).

412 **Table 1.** Number of significantly differentially expressed genes under deoxygenation in the sponge

413 holobionts summarized per reference genome or transcriptome. The direction (up or down) within each

414 paired oxygen condition comparison is relative to normoxia or in the case of anoxia vs. hypoxia, hypoxia.

415 The number of annotated genes is shown in parentheses.

Sponge	Reference	Total genes in Reference	Hypoxia		Anoxia		Anoxia vs hypoxia	
			Up	Down	Up	Down	Up	Down
Eurypon sp. 2	sponge	49,911	499 (239)	238 (112)	551 (228)	825 (340)	622 (273)	468 (205)
H. stellifera	sponge	62,494	465 (125)	58 (9)	40 (11)	244 (87)	1531 (438)	623 (286)
Eurypon sp. 2	mitochondria	14	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)	0 (0)
H. stellifera	mitochondria	14	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Eurypon sp. 2	*Thaumarchaeota*	1,556	19 (14)	1 (1)	3 (3)	0 (0)	0 (0)	0 (0)
H. stellifera	*Thaumarchaeota*	1,551	17 (12)	0 (0)	0 (0)	0 (0)	1 (0)	27 (20)
Eurypon sp. 2	*Gammaproteobacteria*	2,751	3 (3)	29 (29)	0 (0)	0 (0)	0 (0)	0 (0)
H. stellifera	*Gammaproteobacteria*	3,057	0 (0)	0 (0)	0 (0)	0 (0)	20 (20)	416 (401)

In the sponge *H. stellifera*, 523, 284, and 2,154 genes were significantly differentially expressed

in hypoxia, anoxia, and in anoxia vs. hypoxia, respectively (p < 0.1, Table 1). Differentially

expressed genes in the different oxygen conditions corresponded to 0.42-3.4% of the total

number of genes in the *H. stellifera* transcriptome. Approximately 25-37% of the differentially

expressed genes were annotated, which, as in the other sponge species, roughly corresponded

to the global percentage (37.0%) of genes annotated within the *H. stellifera* reference

transcriptome. Under hypoxia in this species, the vast majority of differentially expressed genes

(89%) were upregulated. As in *Eurypon* sp. 2, the majority of differentially expressed genes

(84%) were downregulated in anoxia in *H. stellifera*. The majority of differentially expressed

genes between anoxia and hypoxia were upregulated in anoxia (71%, Table 1).

KOG enrichments in sponges

The global gene expression patterns of Lough Hyne sponges in response to deoxygenation

were compared across the two species by assessing up and down regulation of KOG classes

(broad functional annotations assigned to genes). Within these sponges, KOG expression in

response to hypoxia and anoxia (versus normoxia) were largely distinct in both species (Figure

1A). Expression patterns of both species were only similar in one case between datasets in this

study. KOG expression in *H. stellifera* under hypoxia versus anoxia significantly positively

correlated with that of *Eurypon* sp. 2 under the same conditions ($r = 0.43$, $p < 0.05$, Figure 1E).

Figure 1. KOG expression profiles of sponges under varied oxygen concentrations compared to each other and external datasets. A. Clustered heatmap (based on correlation) of KOG class (rows) enrichments for upregulated or downregulated genes. Highlighted columns correspond to data generated in this study (Purple: *Hymeraphia stellifera* [Hs], Cyan: *Eurypon* sp. 2 [Es]). Unhighlighted columns represent external datasets. Up or down regulation with respect to normoxia is indicated in 'Hypox' (hypoxia) and 'Anox' (anoxia) columns, and differential regulation with respect to hypoxia is shown in 'Hypox vs. Anox' columns, i.e. upregulated KOGs (red delta ranks) are upregulated in anoxia with respect to hypoxia. Within each column on the heatmap, KOGs that are significantly enriched (FDR-adjusted $p < .05$) are marked with asterisks. KOG enrichments from the sponge *Tethya wilhelma* after four days exposure to hypoxia (Tw Hypox Long) and one hour exposure to anoxia (Tw Anox Short), the coral *Acropora millepora* after three days of exposure to heat stress (Am Coral Heat), *Caenorhabditis elegans* in a dauer state (Ce Dauer), and the midge *Sitodiplosis mosellana* in diapause (Sm Diapause) from external studies were included for comparative purposes. B-E. Significant ($p < 0.05$) correlations between the KOG enrichments for different species and oxygen levels from this study. F-J. Significant correlations between KOG enrichments in this study and those of external datasets. All other possible comparisons between the various datasets were not significant.

Within *Eurypon* sp. 2, KOG class expression patterns in hypoxia vs. anoxia were significantly positively correlated to those of hypoxia ($r = 0.71$, $p < 0.01$, Figure 1B). Similarly, in *H. stellifera*, KOG enrichment in hypoxia vs. anoxia was positively correlated to that of hypoxia ($r = 0.72$, $p < 0.001$, Figure 1C) and anoxia ($r = 0.46$, $p < 0.05$, Figure 1D).

At the level of individual KOGs, the KOG class 'replication recombination and repair' was significantly enriched in all pairwise comparisons of oxygen conditions for both sponge species ($P_{adj} < 0.05$), though in different directions depending on oxygen level. It was significantly upregulated in anoxia and anoxia vs. hypoxia but downregulated in hypoxia in both species (Figure 1A).

In *Eurypon* sp. 2, five genes with replication recombination and repair KOG annotations were significantly upregulated in hypoxia, e.g. HELZ and HELZ2 transcription associated helicases, and 11 genes were significantly downregulated ($p < 0.1$, Supplemental Figure 4A), including five genes from the minichromosome maintenance (MCM) protein complex. In anoxia, two genes were significantly upregulated within the replication recombination and repair KOG in this species - 'siz regulation of ARF protein signal transduction' and DNA polymerase epsilon catalytic subunit ($p < 0.1$, Supplemental Figure 4B). The majority of significantly differentially expressed replication recombination and repair genes (five) were downregulated in anoxia ($p < 0.1$, Supplemental Figure 4B). The majority of replication recombination and repair genes (nine) were also downregulated in anoxia vs. hypoxia, with only two significantly upregulated in anoxia (Supplemental Figure 4B). The genes NBN and RFC1 were downregulated in anoxia compared to both normoxia and hypoxia (Supplemental Figure 4B,C). Similarly, the gene 'siz regulation of ARF protein signal transduction' was upregulated in anoxia compared to both normoxia and hypoxia (Supplemental Figure 4B,C).

In *H. stellifera*, three genes were significantly differentially expressed in the replication
recombination and repair KOG class in hypoxia, and all were upregulated (p < 0.1,
Supplemental Figure 4D). No genes within this class were significantly differentially expressed
in anoxia (vs. normoxia). Eleven genes were upregulated and eight genes were downregulated
in anoxia vs. hypoxia (Supplemental Figure 4E). Similar to *Eurypon* sp. 2, HELZ2 was
downregulated in anoxia vs. hypoxia in *H. stellifera*. In contrast to *Eurypon* sp. 2 a distinct MCM
gene (MCM6) was downregulated in anoxia vs. hypoxia. Additionally, four genes from the
structural maintenance of chromosomes protein 6 (SMC6) complex were downregulated in
anoxia vs. hypoxia.
496
497 Like replication recombination and repair, the KOG class 'energy production and conversion'
498 was significantly enriched (P_{adj} < 0.05) between all conditions except for anoxia vs. hypoxia in
499 *H. stellifera* (P_{adj} = 0.07). The energy production and conversion KOG was significantly
500 upregulated in hypoxia in *H. stellifera* but downregulated in all remaining conditions in both
501 sponge species (Figure 1A).
502
Within *Eurypon* sp. 2 specifically, four genes were significantly downregulated and twelve genes
were upregulated within the energy production and conversion KOG class in hypoxia (p < 0.1,
Supplemental Figure 5A). Various nuclear genes for processes in the mitochondria were
differentially expressed in hypoxia. SLC25A27 (a mitochondrial membrane uncoupling protein),
ndufs8 (a NADH dehydrogenase core subunit in the mitochondrial membrane - complex I), and
Surf1 (part of the complex that regulates cytochrome c oxidase assembly) were downregulated
in hypoxia (Supplemental Figure 5A). However, other nuclear mitochondrial genes were
upregulated, including four subunits of ATP synthase and two additional genes with annotations
corresponding to ATP synthesis (Supplemental Figure 5A). In anoxia, fourteen genes were
upregulated and five genes were downregulated in the energy production and conversion KOG
class (Supplemental Figure 5B).
514
As in hypoxia, ndufs8 was downregulated, but no other significant differentially expressed genes
were shared across the two oxygen levels. The following genes were also downregulated in
anoxia: mitochondrial sulfite oxidase (SUOX), malate dehydrogenase (MDH1), and two genes
belonging to the ATP gaunido phosphatase family (Supplemental Figure 5B). Conversely, four
genes involved in phosphoenolpyruvate carboxykinase activity were significantly upregulated in
anoxia (p < 0.1, Supplemental Figure 5B). Additionally, three mitochondrial membrane
transporter proteins from the SLC family, two CYB5R4 cytochrome reductase genes, a
mitochondrial carrier protein, and fdx1 cytochrome reductase were upregulated in anoxia. In
anoxia vs. hypoxia in *Eurypon* sp. 2, ten genes were upregulated in anoxia (Supplemental
Figure 5C). Upregulated genes included all genes that were significantly upregulated in anoxia,
except for the phosphoenolpyruvate carboxykinase associated genes. Six genes were
significantly downregulated in anoxia vs. hypoxia. Of these, three were also downregulated in
anoxia (vs. normoxia) and two genes were significantly upregulated in anoxia (Supplemental
Figure 5A-C). It is noteworthy, however, that all genes upregulated genes in anoxia and hypoxia
were only strongly upregulated in one of the two anoxia replicates (DC56, Supplemental Figure
5 B,C).
531

In *H. stellifera*, eight genes were upregulated and two genes were downregulated in hypoxia. As
in *Eurypon* sp. 2, the ATP5A1 ATP synthase subunit was upregulated in hypoxia. An additional
ATP synthase subunit (ATP6V0C) that was not significantly upregulated in *Eurypon* sp. 2 was
upregulated in *H. stellifera* in hypoxia (p < 0.1, Supplemental Figure 5D). Mitochondrial
sulfide:quinone oxidoreductase (SQRDL), three genes associated with phosphoenolpyruvate
carboxykinase, and a histone demethylase (KDM6A) were also significantly upregulated in
hypoxia in *H. stellifera*. Conversely, PCK1 (also associated with phosphoenolpyruvate
carboxykinase) and MDH1 were significantly downregulated in hypoxia (p < 0.1, Supplemental
Figure 5D). In anoxia in *H. stellifera*, ATP5A1 was significantly upregulated, and ATP6V0C,
PCK1, and MDH1 were downregulated (Supplemental Figure 5E). Comparing
anoxia vs. hypoxia, eight genes were upregulated and five genes were downregulated
(Supplemental Figure 5F).
544
As in hypoxia, SQRDL, four genes associated with phosphoenolpyruvate carboxykinase activity,
and KDM6A were upregulated in anoxia vs. hypoxia. Additionally, two unshared genes,
including the mitochondrial Succinyl-CoA:3-ketoacid coenzyme A transferase 1 (OXCT1) and a
mitochondrial carrier protein, were upregulated in anoxia vs. hypoxia (Supplemental Figure 5F).
Five genes were downregulated in anoxia vs. hypoxia in *H. stellifera*. Two of these four only
exhibited high expression levels in one hypoxic sample (DC105) that drove the observed
difference (Supplemental Figure 5F). The other three downregulated genes in anoxia vs.
hypoxia were mitochondrial creatine kinase (CKMT2), NAD P transhydrogenase (NNT), and
proto-oncogene tyrosine-protein kinase (Src).
554
555 The 'cytoskeleton' KOG class was also significantly enriched (P_{adj} < 0.01) in all but one oxygen
556 condition – anoxia in *Eurypon* sp. 2 (P_{adj} = 0.1). Interestingly, enrichment patterns in this KOG
557 were species specific, being significantly upregulated in hypoxia and anoxia vs. hypoxia in
558 *Eurypon* sp. 2 and downregulated in all conditions for *H. stellifera*.
559
The other twenty KOG classes were only significantly enriched in Lough Hyne sponges under a
few (or no) different oxygen conditions and patterns of significant enrichment were generally
condition and species-specific (Figure 1A). With the exception of energy production and
conversion, KOGs pertaining to metabolic processes were rarely significantly enriched.
Expression of the KOG classes 'carbohydrate metabolism' and 'amino acid transport and
metabolism' were not significantly changed under any oxygen condition (P_{adj} > 0.05). There was
some significant enrichment of metabolic KOGs, but only in *Eurypon* sp. 2. 'Coenzyme transport
and metabolism' and 'inorganic ion transport and metabolism' were both downregulated in
anoxia, and 'nucleic acid transport metabolism' was downregulated in hypoxia vs. anoxia. For
both these conditions (anoxia and hypoxia vs. anoxia), 'Lipid transport and metabolism' and
'Secondary metabolites biosynthesis, transport and catabolism' were significantly
downregulated. However, the transcription of the majority of genes labeled with metabolic KOGs
remained unchanged under anoxia and hypoxia.
573
Comparing the Lough Hyne dataset with external datasets of invertebrate KOG expression in
response to stress, only two positive correlations were found. In *Eurypon* sp. 2, expression
patterns under anoxia vs. hypoxia were comparable to those of *T. wilhelma* under short-term

anoxia ($r = 0.44$, $p < 0.05$, Figure 1H). The hypoxic response of *Eurypon* sp. 2 resembled that of
diapause in *S. mosellana* ($r = 0.68$, $p < 0.001$, Figure 1F), but this was the only correlation found
between Lough Hyne sponge data and diapause or dauer states. Importantly, KOG enrichments
under anoxia (versus normoxia) were not positively correlated to any other external datasets for
either sponge species.
582
Three significant negative correlations, which corresponded to opposing KOG expression
patterns, were observed between Lough Hyne sponges and external datasets. Hypoxia
responses in *H. stellifera* were negatively correlated with those of *A. millepora* under heat stress
($r = -0.56$, $p < 0.01$, Figure 1G). The long exposure of *T. wilhelma* to hypoxia was negatively
correlated to both anoxia ($r = -0.64$, $p < 0.01$, Figure 1J) and hypoxia vs. anoxia ($r = -0.61$, $p <$
0.01, Figure 1I) in *Eurypon* sp. 2.
589
590   <u>GO term enrichments in sponges</u>
591
Generally, significantly enriched GO terms between the different treatments were not shared
between the two sponge species (Supplemental Figures 1-3). The strongest upregulated GO
enrichments ($p < 0.01$) in hypoxia in *Eurypon* sp. 2 were cellular signaling and 'lipoprotein
particle receptor binding' (Supplemental Figure 1A). For *H. stellifera*, on the other hand, the
strongest upregulated enrichments ($p < 0.01$) in hypoxia included the terms 'protein tyrosine
kinase binding,' 'molecular adaptor,' 'structural molecule,' and 'structural constituent of
ribosome'; whereas the downregulation of the term 'motor' was strong ($p < 0.01$, Supplemental
Figure 1B).
600
Under anoxia, strong upregulation of the 'proteasome binding' term was observed in *Eurypon*
sp. 2 ($p < 0.01$), and the terms 'phosphoenolpyruvate carboxykinase', 'carbohydrate kinase,'
monovalent inorganic cation transmembrane transporter,' and 'sodium ion transmembrane
transporter' were highly downregulated ($p < 0.01$, Supplemental Figure 2A). In *H. stellifera*
under anoxia, the terms 'damaged DNA binding' and 'calcium-dependent phospholipid binding'
were highly upregulated, and four terms associated with various aspects of the cytoskeleton
were downregulated ($p < 0.01$, Supplemental Figure 2B).
608
In *Eurypon* sp. 2 under anoxia vs. hypoxia, the terms 'lipoprotein receptor binding' and
'proteasome binding' were strongly upregulated in anoxia, but no terms were strongly
downregulated in anoxia versus hypoxia ($p < 0.01$, Supplemental Figure 3A). The former term
was also upregulated under anoxia vs. normoxia. Conversely, the strongest enrichments in *H.
stellifera* in anoxia vs. hypoxia were downregulated in anoxia and were composed of three of
the four cytoskeletal binding GO terms that were also downregulated in anoxia vs. normoxia
(see above, $p < 0.01$, Supplemental Figure 3B), but no strong upregulation of GO terms was
observed.
617
Considering GO enrichments at lower significance levels ($0.01 < p < 0.1$), within *Eurypon* sp. 2,
the GO terms 'proteasome binding', 'lipoprotein particle receptor binding,' and 'Hsp70 binding'
(HSP: Heat shock protein) were significantly upregulated in all cases of decreased oxygen albeit
at varied significance levels ($p < 0.05$, Supplemental Figures 1–3A). Between only two levels,

the terms 'transmembrane signaling receptor,' 'secondary active transmembrane transporter,'
and 'molecular transducer' were upregulated in hypoxia but downregulated in anoxia. However,
the term lipoprotein particle receptor binding was downregulated in anoxia and upregulated in
hypoxia ($p < 0.1$, Supplemental Figures 1,2A). Additionally, the term 'structural constituent of
ribosome' was upregulated in anoxia but downregulated in anoxia vs. hypoxia ($p < 0.1$,
Supplemental Figures 1,3A). No additional GO terms other than the three present in all oxygen
levels for *Eurypon* sp. 2 were enriched in both the pairwise comparison of anoxia and anoxia vs.
hypoxia.
630
No GO terms were significantly up or downregulated in all three levels of oxygen conditions for
H. stellifera. However, some GO terms were significantly enriched in two levels. The term
'ATPase, coupled to transmembrane movement of ions, rotational mechanism' was
downregulated in hypoxia but upregulated in anoxia ($p < 0.1$, Supplemental Figures 1B, 2B
respectively). The GO terms 'I-SMAD binding', thioesterase binding,' and 'ubiquitin-like protein
ligase binding' were all upregulated in hypoxia (vs. normoxia) and anoxia vs. hypoxia, while the
'motor' term was downregulated ($p < 0.1$, Supplemental Figures 1B, 3B).
638
639   There were only two shared GO enrichments between both sponge species for any given
640   oxygen condition. The terms 'protein tyrosine kinase binding' and 'structural constituent of the
641   ribosome' were upregulated in both species in hypoxia ($p < 0.1$, Supplemental Figure 1),
642   although they were enriched to different degrees of significance. No significantly enriched GO
643   terms were shared between species in normoxia vs. anoxia or hypoxia vs. anoxia.
644
At the gene level, significantly differentially expressed heat shock proteins (Hsps) in both
sponges are shown in Supplemental Figure 8. In *Eurypon* sp. 2, one Hsp gene
(DNAJC8/Hsp40) was downregulated in hypoxia ($p < 0.01$), but four Hsp genes (HSPA8, two
genes for HSP90B1, and Hsp70) were significantly upregulated in hypoxia ($p < 0.1$,
Supplemental Figure 8A). Additionally, HSPA8 and C11orf73 (Hsp70 binding protein) were
upregulated in anoxia vs normoxia ($p < 0.05$, Supplemental Figure 8A). Another Hsp70 gene
was also significantly downregulated in anoxia vs. hypoxia ($p < 0.05$, Supplemental Figure 8A).
In *H. stellifera*, HSP90A1 was upregulated in hypoxia and anoxia ($p < 0.1$, Supplemental Figure
8B). Two additional Hsp genes (both for HSP90B1) were also upregulated in anoxia. Finally, the
two HSP90B1 genes were upregulated in anoxia vs. hypoxia, but HSPA5 was downregulated in
anoxia vs. hypoxia ($p < 0.05$, Supplemental Figure 8B).
656
Mitochondrial gene expression
All 14 coding genes from the sponge mitochondria were transcribed in both species and in all
oxygen conditions. There was only one mitochondrial gene (nad2) that was significantly
differentially expressed but only in *Eurypon* sp. 2. This gene was significantly upregulated ($p <$
0.1) in anoxia vs hypoxia (Supplemental Figure 6A, Table 1). Similar counts for each individual
mitochondrial gene across both sponge species and all oxygen conditions were found. For
instance, counts for atp6 were consistently twice as high as counts for atp8 (Supplemental
Figure 6).
665
666

Prokaryotic differential expression

Thaumarchaeota gene expression

For the _Thaumarchaeota_ associated with _Eurypon_ sp. 2, 20, 3 and 0 genes were differentially expressed in hypoxia, anoxia, and hypoxia vs. anoxia, respectively ($p < 0.1$, Table 1). These genes accounted for 0-1.3% of this _Thaumarchaeota_ genome, and 75-100% of the differentially expressed genes were annotated (Table 1), compared to 69.2% annotated in the entire reference genome. All but one of the differentially expressed genes in hypoxia were upregulated (Table 1, Figure 2A). The downregulated gene belonged to the sodium solute symporter family (SSF, TC 2.A.21, Figure 2A). The upregulated genes in hypoxia included three stress response genes (COG0589 Universal stress protein [UspA], small heat shock protein [HSP20], and DnaJ molecular chaperone homology domain), two CBS domains, a member of the lactate/malate dehydrogenase (LDH MDH) superfamily, a tetratricopeptide (TPR) repeat, two transcription regulators, and a PII signal transduction protein.

Figure 2. A-D. All significantly differentially expressed genes for the _Thaumarchaeota_ in both sponge species (left: _Eurypon_ sp. 2, right: _H. stellifera_. For _Thaumarchaeota_ associated with _Eurypon_ sp. 2, only two pairwise comparisons of oxygen conditions had significant differences: A. hypoxia vs. normoxia and C. anoxia vs. normoxia. In symbionts associated with _H. stellifera_, there were significantly differentially expressed genes in hypoxia vs. normoxia (B) and anoxia vs. hypoxia (D). There were no significant

690   differences in the three genes involved in ammonia oxidation (AmoA, B and C) in either symbiont, but
691   their heatmaps are shown in E and F for *E.* sp 2 and *H. stellifera*, respectively.
692
Only three genes were significantly differentially expressed in anoxia in *Eurypon* sp. 2's
Thaumarchaeota, and all were upregulated (Table 1, Figure 2C). The upregulated genes in
anoxia were a UbiD family decarboxylase, CAAX protease self-immunity, and the same TPR
repeat that was also upregulated in hypoxia. There were no significantly differentially expressed
genes between anoxia and hypoxia in this *Thaumarchaeota* (Table 1). The genes necessary for
ammonia oxidation, i.e. subunits A, B and C of ammonia monooxygenase (Amo), were present
in this *Thaumarchaeota*, but they were not differentially expressed under deoxygenation (Figure
2E).
701
In the *Thaumarchaeota* symbiont of *H. stellifera*, 17, 0 and 28 genes were significantly
differentially expressed in hypoxia, anoxia and hypoxia vs. anoxia, respectively (Table 1).
Similar to the symbiont of *Eurypon* sp. 2, all 17 differentially expressed genes in the *H. stellifera*
symbiont were upregulated in hypoxia. These genes included the stress response genes
'universal stress protein, an HSP20 family protein, and two copies of UspA. HSP20 and UspA
were also upregulated in the *Eurypon* sp. 2 holobiont. Similarly, a PII signal transduction protein,
CBS domain, and two transcription regulators (one identical: TBP stabilizer, and one distinct to
H. stellifera symbionts) were upregulated in both *Thaumarchaeota* species in hypoxia. Unlike in
Eurypon sp. 2, the *Thaumarchaeota* in *H. stellifera* upregulated a helix turn helix ASNC type
domain and a Lrp/AsnC ligand binding domain (Figure 2B). No genes were significantly
differentially expressed in anoxia in this *Thaumarchaeon*.
713
As in the other holobiont species, *Thaumarchaeota* genes for ammonia oxidation were not
significantly differentially expressed with *H. stellifera* symbionts (Figure 2F). Twenty-seven
genes were upregulated and one gene (K141_339108_19, unannotated) was downregulated in
anoxia vs. hypoxia in the *Thaumarchaeota* within *H. stellifera* (Figure 2D). Most of the
upregulated genes in anoxia vs. hypoxia were also upregulated in hypoxia. Upregulated genes
in anoxia vs. hypoxia unique to the *H. stellifera* holobiont included additional transcriptional
regulators, an integrase domain, and a putative nitroreductase.The phylogenies of this
upregulated nitroreductase and all nitroreductases from the studied holobionts as well as
publicly available nitroreductase sequences from other *Thaumarchaeota* are shown in
Supplemental Figure 9.
724
725   For both *Thaumarchaeota*, only two COGs ('signal transduction mechanisms' and
726   'transcription') were significantly enriched ($P_{adj} < 0.05$, Supplemental Figure 7A). These two
727   COGs were both downregulated in hypoxia and anoxia vs. hypoxia within *H. stellifera*
728   symbionts. All other COG classes were comparable in their expression levels regardless of
729   holobiont species or oxygen concentration (Supplemental Figure 7A).
730
731   *Gammaproteobacteria* gene expression
732   The *Gammaproteobacteria* associated with *Eurypon* sp. 2 only exhibited significant differential
733   expression under hypoxia but not in any other oxygen condition (Table 1). In hypoxia, three
734   genes were upregulated: type 1 restriction-modification system, flavin adenine dinucleotide

binding, and signal sequence binding (Figure 3A). The 29 downregulated genes in hypoxia
included two genes involved in sulfur processing (k212078_2, k237243_12), cytochrome c
(class I), a gene associated with the assembly of the periplasmic c-type cytochromes
(k561840_2), a NQR complex gene, various genes involved in pore formation, a Rhodanese-
like gene, glutamine synthetase (type I), and YbiS YcfS YnhG family protein.

Figure 3. All significantly differentially expressed genes for the *Gammaproteobacteria* in both
sponge species (A: *Eurypon* sp 2, B: *H. stellifera*. For *Gammaproteobacteria* associated with
Eurypon sp. 2, there were only significant differences between hypoxia vs. normoxia (A). In
symbionts associated with *H. stellifera*, there were only significantly differentially expressed
genes in anoxia vs. hypoxia (B,C). All downregulated genes in anoxia vs. hypoxia are shown in B.
There were 402 genes that were downregulated in anoxia vs. hypoxia in *Gammaproteobacteria*
from *H. stellifera*, so only the top 10 based on p value are shown (C).

In the *Gammaproteobacteria* associated with *H. stellifera*, genes were only significantly
differentially expressed between anoxia and hypoxia (436 genes, p <0.1, Table 1, Figure 3B,C).
Twenty genes were upregulated in anoxia vs. hypoxia and all were annotated (Figure 3B).
Upregulated genes in anoxia vs. hypoxia included five genes associated with the ribosome
(k141_198365_15, k141_198365_8, k141_198365_17, and k141_245017_3), various genes for
molecular transport, a transmembrane proton pump (k141_226857_1), succinyl-CoA
synthetase, and formate dehydrogenase (alpha subunit). There were also 416 genes that were
significantly downregulated in anoxia vs. hypoxia (Table 1). The top ten upregulated genes
(lowest p values) are exhibited in Figure 3C.

763 *Gammaproteobacterial* COG expression is shown in Supplemental Figure 7B. Only one COG
764 class, 'translation, ribosomal structure and biogenesis,' was significantly enriched in
765 *Gammaproteobacteria* under the different oxygen conditions ($P_{adj} < 0.05$). This COG class was
766 downregulated in *Gammaproteobacteria* associated with *Eurypon* sp. 2 under hypoxia as well
767 as those of *H. stellifera* in anoxia vs. hypoxia (Figure 7B).
768
Discussion
770
Sponge holobiont gene expression was significantly changed by exposure to deoxygenation in
situ, though responses were generally species-specific, dependent on oxygen level, and varied
across holobiont members. The potential functions of this differential expression are discussed
below for each member of the two holobionts: sponge, sponge mitochondria, *Thaumarchaeota*
and *Gammaproteobacteria*. Changes in gene expression within each member of the holobiont is
discussed in a separate section below, and then the potential functional adaptations to
deoxygenation at the level of the holobiont are summarized. The implications of this study for
past and future oceans are considered.
779
Differential expression in sponges during deoxygenation
Cellular stress responses were observed in the expression of HSPs, a group of stress response
genes that have been widely studied in marine invertebrates. These proteins are molecular
chaperones involved in protein folding, unfolding, transport, assembly of protein complexes, and
the protection of restoration of proteins exposed to stressors, and they are present in nearly all
organisms (Li and Srivastava 2004). Both Lough Hyne sponges upregulated HSP genes in
response to hypoxia and anoxia (vs normoxia), though there was downregulation of some HSP
genes in deoxygenation as well. *Eurypon* sp. 2 and *H. stellifera* both upregulated Hsp90 genes
in hypoxia, but they upregulated different HSPs in anoxia: HSPA8 and Hsp90 for *Eurypon* sp. 2
and *H. stellifera*, respectively. It should be noted that Hsp90 upregulation in *H. stellifera* in
anoxia was driven by only one replicate. However, in the pairwise comparison with the highest
replication in this study, hypoxia vs. normoxia in *Eurypon* sp. 2, Hsp90 genes were significantly
upregulated in hypoxia. Hsp90 expression was also higher in anoxia than normoxia for *Eurypon*
sp. 2, but this was not significant. Based on these trends, it is possible that increased replication
would result in a clearer upregulated signal from Hsp90 genes, since false negatives in gene
expression analyses decrease with more replication (Schurch et al. 2016). *Eurypon* sp. 2 also
upregulated HSPA8 and Hsp70 in hypoxia but downregulated Hsp40. In long term hypoxia, *T.*
wilhelma exhibited a similar HSP expression pattern, upregulating Hsp90 but downregulating
Hsp70 (Mills et al. 2018). Sponges under heat stress also upregulate Hsp70 (López-Legentil et
al. 2008; Guzman and Conaco 2016; Webster et al. 2013) and HSP90 (Guzman and Conaco
2016), as observed under deoxygenation stress in Lough Hyne. Pollutants such as lead and
zinc can also stimulate the upregulation of HSPs in sponges (Efremova et al. 2002; Schröder et
al. 2006). HSPs are therefore crucial to sponge stress response, protecting and restoring
proteins that could be denatured by stressors. HSP expression patterns and other key
transcriptomic responses of the sponge holobionts to deoxygenation are outlined in Figure 4.

Figure 4. Summary of sponge holobiont responses to deoxygenation presented in a tabular format with species on the left (*Es: Eurypon* sp. 2, *Hs: H. stellifera*) and oxygen condition on the top (hypoxia, anoxia, and hypoxia vs. anoxia). The major components of the holobiont are illustrated as follows: sponge cell (tan), mitochondria (red), *Gammaproteobacteria* (green), and *Thaumarchaeota* (purple). Upregulated and downregulated genes and KOGs are indicated with red and blue arrows, respectively. Unchanged gene expression (GE) is noted by the = sign. Correlations (corr.) with external dormancy datasets are indicated along with the direction of the correlation (+ or -). The 'energy production and conversion' and 'replication recombination and repair' KOG classes are abreviated as EPC and RRR, respectively.

In metazoans with a complete HIF pathway, Hsp90 also serves to stabilize HIFa and therefore
plays an important role in hypoxia tolerance (Semenza 2007). In the absence of HIFa in
sponges, Hsp90 may play a greater role in oxygen homeostasis given that it is upregulated
under hypoxia in all sponges that have been examined transcriptomically to date, i.e. *Eurypon*
sp. 2, *H. stellifera* and *T. wilhelma,* and that Hsp90 is upregulated in *H. stellifera* under anoxia.
Hsp90 could stabilize currently unknown proteins that are involved in sponge oxygen regulation
or modulate transcription factors other than HIFa. Further research is needed to determine the
exact role of Hsp90 in sponge oxygen homeostasis, but it presents a good candidate for a gene
underlying sponge tolerance to deoxygenation.

Patterns of species and oxygen-condition-dependent expression of GO terms were observed in
sponges in response to deoxygenation. GO terms such as 'proteasome binding' and 'Hsp70
binding' were upregulated in all cases of decreased oxygen in *Eurypon* sp. 2, suggesting that
the species-specific response to deoxygenation in *Eurypon* sp. 2 may center around repairing or
degrading damaged proteins. In *H. stellifera*, on the other hand, upregulated GO terms related
to DNA repair in anoxia, including 'damaged DNA binding', 'DNA binding,' and 'catalysis of DNA
and RNA,' so DNA repair may be more central to the deoxygenation tolerance of this species.
Both protein and DNA repair upregulation might be a response to reactive oxygen species
(ROS), which can become more abundant under hypoxia due to imbalances in the
mitochondrial electron transport chain (Görlach et al. 2015). If there is any oxygen present in the
sponges under environmental anoxia, e.g. oxygen produced by *Thaumarchaeota* (see Kraft et
al. 2022, _Thaumarchaeota gene expression_ below), then ROSs could cause cellular stress in
anoxia. Otherwise, the presence of sulfide in the water under anoxia (Schuster et al. 2021) or
ROSs other than superoxide (O_2^-) could damage proteins and DNA in anoxic conditions. HSP
genes could then serve to protect and/or restore proteins damaged by sulfide or reactive oxygen
species under hypoxia or anoxia.

The energy production and conversion KOG class was significantly downregulated in all oxygen
conditions in both sponges, indicating potential decreases in energy production, except for
hypoxia and hypoxia vs. anoxia in *H. stellifera*, when it was significantly upregulated and
unchanged, respectively (Figure 1, 4). Within the energy production and conversion KOG class,
however, there were species-specific responses to anoxia and hypoxia at the level of individual
differentially expressed genes. Most nuclear ATP synthase genes were upregulated in hypoxia
in *Eurypon* sp. 2 but were unchanged under anoxia. Similarly, two ATP synthase subunits
(ATP5A1 and ATP6V0C) were upregulated in hypoxia in *H. stellifera*. In anoxia, ATP5A1 was
also upregulated, but ATP6V0C was downregulated. Additionally, downregulation of SLC25A27
in *Eurypon* sp. 2 in hypoxia may increase the coupling across the mitochondrial membrane to
improve ATP synthesis. We note that the electron transport chain genes Ndufs8 and Surf1 were
also downregulated in hypoxia *Eurypon* sp. 2. The NNT gene, also involved in electron
transport, was downregulated in anoxia vs. hypoxia in *H. stellifera*. Nevertheless, most of the
genes associated with the respiration electron transport chain were not differentially expressed,
and electron transport likely continued under deoxygenation. Sponges may therefore have
increased ATP production under hypoxia but not under anoxia. This increase in ATP production
could improve metabolic efficiency under hypoxia. In a similar way, mollusks may upregulate
mitochondrial electron transport system activity and ATP synthesis capacity to improve
respiration efficiency under hypoxia (Hochachka and Lutz 2001).

Glycolysis, which can produce ATP and NADH without oxygen (Stryer 1995), was expected to
increase under anoxia to provide energy. However, relatively few genes directly or indirectly
involved in glycolysis were differentially expressed in our sponges under deoxygenation. In
Eurypon sp. 2, pantothenate kinase 1 (PANK1), ATP-dependent 6-phosphofructokinase (PFKL)
and Phosphoenolpyruvate carboxykinase 2 (PCK2) were significantly upregulated in anoxia,
and only aldehyde dehydrogenase 3 family, member a2 (ALDH3A2) was significantly
upregulated in hypoxia (see Supplemental Reports, Supplemental Figure 5). In *H. stellifera*,
acetyl-coenzyme A synthetase 1 (ACSS1) was significantly downregulated in hypoxia, but
PCK1 and pyruvate kinase (PKM) were upregulated in hypoxia. In anoxia for *H. stellifera*, beta-
enolase (ENO3) and PCK1 were upregulated. Based on these results, it is unlikely that
glycolysis rates increased under anoxia and hypoxia in either species. It is more probable that
glycolysis continued under anoxia and hypoxia at normal rates in both species, since gene
expression in the other ~70 genes associated with glycolysis was not significantly changed
under deoxygenation (Figure 4). Either way, the sponges would have at least some access to
anaerobically generated ATP through glycolysis during environmental deoxygenation.

In the KOG class 'replication, recombination and repair', there was significant upregulation in
anoxia and anoxia vs. hypoxia, but downregulation in hypoxia (vs. normoxia) for both sponge
species. In *Eurypon* sp. 2, many MCM genes involved in the initiation of DNA replication were
downregulated in hypoxia but remained unchanged in anoxia. Certain aspects of DNA
replication, e.g. the DNA polymerase epsilon catalytic subunit, may even be upregulated in
anoxia. The pattern was less clear in *H. stellifera*, but MCM genes were downregulated in
anoxia compared to hypoxia (Figure 4). These data indicated that DNA replication may
decrease under hypoxia but remain stable during anoxia. In hypoxia, decreased DNA replication
and, by extension, growth may be limited by increased oxidative stress and/ or increased
metabolic rates under hypoxia compared to normoxia (see above).

Between the two study species, there was only one significant, positive correlation of KOG class
expression between hypoxia in *Eurypon* sp. 2 and hypoxia vs. anoxia in *H. stellifera*,
demonstrating the general species-specific responses of most KOGs to deoxygenation. Within a
single sponge species, no correlations were seen between KOG enrichments in anoxia or
hypoxia (vs. normoxia), highlighting the transcriptomic differences between responses to
hypoxia and anoxia. KOG enrichments in hypoxia *versus* anoxia, on the other hand, were
positively correlated with those of anoxia in *Eurypon* sp. 2 and both anoxia and hypoxia in *H.*
stellifera.

When quantitatively compared to transcriptomic datasets from the sponge *T. wilhelma* using
KOGMWU, transcriptomic responses in Lough Hyne sponges were only positively correlated
between *T. wilhelma* under short-term anoxia and *Eurypon* sp. 2 in hypoxia vs. anoxia (Figure
1H). This positive correlation could indicate similar transcription patterns demonstrated by *T.*
wilhelma and *Eurypon* sp. 2 transitioning from hypoxia to anoxia. However, no further
similarities between the responses of *T. wilhelma* and Lough Hyne sponges to deoxygenation
were observed. In fact, the long-term exposure of *T. wilhelma* to hypoxia was negatively
correlated to KOG enrichments for both anoxia vs. hypoxia and normoxia vs. anoxia in *Eurypon*
sp. 2 (Figure 1I, J), meaning that KOG expression was generally opposite between these two
species for these conditions in terms of up and down regulation. The physiological responses of

T. wilhelma and *Eurypon* sp. 2 to deoxygenation are presumably very different. These results
demonstrate that large-scale transcriptomic responses to deoxygenation depend on both the
studied species and oxygen concentration.

Based on KOG correlation analyses, the hypothesis that sponges are functionally dormant, i.e.
temporarily ceasing growth, development, and physical activity, under anoxia was not supported
(Figure 4). Sponge gene expression under anoxia did not correlate to either diapause or a dauer
state in *C. elegans* (Sinha et al. 2012) or *S. mosellana* (Gong et al. 2013), respectively.
Although these dormancy datasets come from distantly related invertebrates, Strader et al.
(2016) demonstrated a correlation between the KOG expression patterns of a specific
phenotype of coral larvae and diapause in *S. mosellana*, hypothesizing that this potential
functional analog of dormancy could serve to increase larval dispersal. Therefore, the sponges
of Lough Hyne could only be considered dormant under anoxia if their transcriptional method of
dormancy was distinct from that of an insect, nematode and coral, which is possible but not
parsimonious given that animals are monophyletic (Feuda et al. 2017). Nevertheless, it was
inconclusive whether or not sponges are pumping water or feeding during anoxia, but sponges
were definitively pumping under hypoxia (Schuster et al. 2021). Sponge pumping and filtering
activity remain to be quantified under anoxia and hypoxia in situ.

Paradoxically, *Eurypon* sp. 2 responses to hypoxia were positively correlated to diapause in *S.*
mosellana (Figure 1F, 5). Therefore, the hypoxic response in *Eurypon* sp. 2 species, rather than
the anoxic response, was similar to diapause in terms of global KOG expression patterns. The
replication recombination and repair KOG was also downregulated in *Eurypon* sp. 2 under
hypoxia as well as in *S. mosellana* under diapause, which may partially explain the correlation
between these datasets. Despite the correlation, energy production and conversion KOG
expression patterns between the two species were opposing. The energy production and
conversion KOG was significantly downregulated in hypoxia in *Eurypon* sp. 2 and upregulated in
diapause in *S. mosselana*. These KOG enrichments are calculated from thousands of genes, so
any individual gene within a KOG class might still be upregulated even if the KOG class is
significantly downregulated. Therefore, it is possible that ATP production increased in hypoxia
based on the upregulation of ATP synthase genes, in spite of the downregulation of the energy
production and conversion KOG. Diapause can be associated with the reduction of cell division
(MacRae 2010), which may occur in sponges under hypoxia but not anoxia based on MCM
gene expression. In comparison, KOG enrichments under anoxia were not positively correlated
to any external datasets for either sponge species (Figure 1), indicating that sponge
transcriptomic responses to anoxia are relatively novel compared to those of other taxa in
response to various factors in the existing literature (Dixon et al. 2015; Strader et al. 2016).
938
939   **Sponge mitochondrial gene expression during deoxygenation**
940
A high degree of transcriptional stability under deoxygenation was observed in sponge
mitochondria. Only one of the 14 mitochondrial protein-coding genes, nad2, was significantly
differentially expressed in the different oxygen conditions, where it was upregulated in anoxia
vs. hypoxia in *Eurypon* sp 2 (Figure 4). Both of the sponge species had higher expression of
nad2 in anoxia than in normoxia or hypoxia, but this difference was only significant in *Eurypon*
sp. 2.

947
Surprisingly little is known about differential gene expression in mitochondria, even within
humans. In the limited studies available, mitochondrial gene expression was significantly
different between human cancer cells and normal cells (Reznik et al. 2017) as well as between
different populations of healthy pancreas tissue (Medini et al. 2021). In these studies, nad2 was
significantly downregulated in most studied cancers, though it was still significantly upregulated
in a few cancer types, and nad2 was also significantly differentially expressed between
pancreas cell populations. Based on the general upregulation of nuclear genes associated with
respiration in hypoxia, mitochondrial gene upregulation was expected in hypoxia when
compared to normoxia and anoxia, but the opposite was true for nad2 in *Eurypon* sp. 2 in the
current study. The other six nad genes, which code for parts of the same membrane respiratory
chain complex (complex 1, NADH dehydrogenase), however, were not significantly differentially
expressed in either species. In both human studies, when one mitochondrial gene was
significantly differentially expressed, most other mitochondrial genes were significantly
differentially expressed in the same direction (Reznik et al. 2017; Medini et al. 2021). The
physiological meaning (if any) of this single, significantly differentially expressed mitochondrial
gene in sponges remains to be determined. However, it is most likely that the functionality of the
mitochondrial was unchanged by either anoxia or hypoxia based on gene expression results.
965
Compared to that of humans, even less is known about mitochondrial gene expression in
marine invertebrates. Mitochondrial gene expression is usually excluded from differential
expression analyses due to the differences in library preparation potentially needed for
sequencing mitochondrial RNA and/or the exclusion of mitochondrial RNA from reference
transcriptomes (*sensu* Meyer 2016). This dearth is partly due to the nature of mitochondrial
transcription, which generates a single continuous transcript. Following transcription, however,
mitochondrial transcripts are broken up into individual coding mRNA, tRNA or rRNA (D'Souza
and Minczuk 2018). After the breakup, the levels of mRNAs from specific genes are controlled
post-transcriptionally (D'Souza and Minczuk 2018). It is therefore possible to have different
counts between the different mitochondrial genes despite their single transcript origin
(Supplemental Figure 6). The consistent 'expression' of transcripts across oxygen conditions for
each mitochondrial protein-coding gene following continuous transcription are likely to be
necessary for the maintenance of mitochondrial function.
979
It remains unclear how sponge mitochondria maintain this transcription from an energetic
standpoint under anoxia. Within the mitochondrial electron transport chain, oxygen is used as
the terminal electron acceptor and is crucial for the generation of ATP through oxidative
phosphorylation. Theoretically, anoxia should cause electron transport and oxidative
phosphorylation in the mitochondria to cease (Chandel and Schumacker 2000). If the
mitochondria are transcribing solely to maintain oxidative phosphorylation machinery until
oxygen returns, then they would need to use at least enough energy (~2 ATP per base) to
transcribe for months at a time without oxygen (Lynch and Marinov 2015), and this energetic
requirement would apply to the transcription of sponge nuclear genes as well. Glycolysis may
provide some anaerobic ATP production, even if it is not upregulated in anoxia or hypoxia.
Alternatively, to maintain ATP production through aerobic respiration, the sponges of Lough

Hyne might have access to oxygen produced by their *Thaumarchaeota* symbionts, a topic
explored in more detail below.
993

Microbiome gene expression during deoxygenation

<u>*Thaumarchaeota* gene expression</u>

Thaumarchaeota symbionts from both sponge species exhibited significant differential gene
expression under hypoxia. Only the *Thaumarchaeota* from *Eurypon* sp. 2 (OTU1) exhibited
differential gene expression under anoxia, and only those of *H. stellifera* (OTU3) differentially
expressed genes under anoxia vs. hypoxia. Though there were some species-specific
responses, overall gene expression in both *Thaumarchaeota* was similar, supporting the
hypothesis from Schuster et al. (2021) that these taxonomically similar symbionts between two
sponge species might have similar functional roles.

Relatively few genes involved in anaerobic metabolism were expressed in *Thaumarchaeota*
symbionts under any oxygen condition. Lactate/malate dehydrogenase, which could be involved
in fermentation (Bayer et al. 2020), was upregulated in *Eurypon* sp. 2 *Thaumarchaeota*, but only
in hypoxia. Genes for the other anaerobic pathways suggested by Bayer et al. 2020, i.e.
succinate dehydrogenase/ fumarate reductase, nitrite reductase and sulfite reductase, were
present in both *Thaumarchaeota* genomes in the present study but were not significantly
differentially expressed under different oxygen conditions. The genes for nitrite reductase
exhibited low expression across all samples (0–20 counts) in both *Thaumarcheaota,* and so did
the dehydrogenase/ fumarate reductase gene in the *Thaumarcheaota* of *H. stellifera*. Therefore,
these genes with low expression may not be used under the conditions of this study. Only three
genes in the *Thaumarchaeota* of *Eurypon* sp. 2 under anoxia were significantly differentially
expressed, and none of these were involved in anaerobic metabolism. No genes were
differentially expressed in *H. stellifera Thaumarchaeota* under anoxia. Therefore, there was no
transcriptomic evidence for the upregulation of anaerobic metabolism within *Thaumarchaeota* in
anoxia.

Transcription of ammonium monooxygenase (AMO) genes did not significantly change under
anoxia or hypoxia, even though the ammonia oxidation pathway is oxygen dependent. Like the
sponge mitochondria, the *Thaumarchaeota* are therefore either transcribing AMO genes to be
ready when oxygen returns, or they are obtaining trace oxygen for this pathway by synthesizing
oxygen themselves (Kraft et al. 2022). If sponge *Thaumarchaeota* symbionts are also able to
produce oxygen under anoxic conditions like their relative *N. maritimus*, then they should
express AMO genes under anoxia as well as normoxia. The proposed pathway for oxygen
production in *N. maritimus* involves the reduction of nitrite to nitric oxide that is dismutated into
oxygen and nitrous oxide (Kraft et al. 2022). Water sampled for nitrite concentrations in Lough
Hyne using established methods (Grasshoff et al. 1999) revealed that nitrite concentrations
under anoxia ranged from 0.2–1.2 µM (Supplemental Table 3), increasing with depth from
where seasonal anoxia and hypoxia occur. Therefore, the conditions in Lough Hyne would
theoretically allow for *Thaumarchaeota* to produce oxygen under anoxia.

AsnC family proteins and nitroreductase expression was upregulated during oxygen production
in *N. maritimus* (Kraft et al. 2021). In *Archaea*, AsnC proteins can regulate metabolic processes,
including the switch between aerobic and anaerobic metabolisms (Lemmens et al. 2019).
Similarly, nitroreductases can have a wide range of metabolic functions (Akiva et al. 2017).
Thaumarchaeota within *H. stellifera* upregulated a Lrp/AsnC ligand binding domain under
hypoxia and in anoxia vs. hypoxia. Another AsnC type protein was also upregulated under
hypoxia, and a nitroreductase was upregulated under anoxia vs. hypoxia. However, these
proteins did not directly correspond to those upregulated in *N. maritimus* during oxygen
production (Supplemental Figure 9), and the latter proteins are not definitively known to be
involved in oxygen production, since the pathway remains unknown.

In hypoxia, both *Thaumarchaeota* upregulated the stress response genes USpA and a protein
from the HSP20 family (Figure 4). It was surprising that *Thaumarchaeota* may be stressed by
hypoxia since free-living thaumarchaeal Nitrosopumilales can be found in high abundances in
oxygen-minimum zones (OMZs) where oxygen is either at low concentrations or below
detection (Löscher et al. 2012), though some *Thaumarchaeota* OTUs decrease in abundance in
suboxic and anoxic zones (Sollai et al. 2019). Members of the genus *Nitrosopumilus*, however,
increase their absolute abundance in suboxic regions, but then decrease in abundance with
depth in the anoxic zone (Sollai et al. 2019), indicating a preference for suboxic environments. If
suboxia is also the preferred niche of sponge *Thaumarchaeota* symbionts, the upregulation of
stress response genes in response to hypoxia may correspond to a rise in intracellular oxidative
stress following an increase in metabolic activity in hypoxia, like their sponge hosts. In this case,
these stress response proteins serve a protective and/or restorative function for other crucial
proteins and structures damaged by oxidative stress.
1054
Gammaproteobacteria gene expression

Gene expression in the *Gammaproteobacteria* was species and oxygen-condition dependent. In
Eurypon sp. 2, *Gammaproteobacteria* downregulated genes in hypoxia that were associated
with respiration and energy production, including cytochrome c and associated genes (Welchen
et al. 2016). The *Gammaproteobacteria* also downregulated key metabolic genes in hypoxia
such as glutamine synthetase and structural genes like YbiS YcfS YnhG family protein, which is
involved in cell wall production (Magnet et al. 2008). So unlike sponges or *Thaumarchaeota*,
Gammaproteobacteria in *Eurypon* sp. 2 (OTU2) may decrease metabolic activity under hypoxia,
though this is likely not the case in *H. stellifera* symbionts (OTU7).

Gammaproteobacteria in *H. stellifera* downregulated hundreds of genes in anoxia vs. hypoxia,
including USpA, so, like other members of the holobiont, hypoxia requires a stronger
transcriptomic response and, therefore, may be more stressful than anoxia for these
Gammaproteobacterial symbionts. The few upregulated genes in anoxia vs. hypoxia included
genes for energy production, e.g. a transmembrane proton pump, succinyl-CoA synthetase (part
of the citric acid cycle), and formate dehydrogenase (alpha subunit), although other genes
associated with energy production were downregulated. Formate dehydrogenase can function
in both aerobic respiration and in anaerobic respiration (Jormakka, et al. 2003), but this enzyme
was not significantly upregulated in anoxia compared to normoxia or in the

Gammaproteobacteria of *Eurypon* sp. 2 under any condition. So, it is unlikely that this anaerobic
pathway is active in sponge-associated *Gammaproteobacteria* under anoxia.

Gammaproteobacteria symbionts were previously hypothesized to oxidize sulfide under anoxia
(Schuster et al. 2021). Although they are not present in OTU2, OTU7 from *H. stellifera* has the
sulfur oxidation genes soxA, soxY, and soxZ, which could oxidize sulfide, as well as the nitrate
reductase gene NAP. However, the sox genes were not expressed in any samples in the
current study, and NAP was expressed at very low levels across all samples (2-28 counts).
Moreover, neither symbiont exhibited significant differential gene expression in anoxia versus
normoxia. Consequently, there was very little transcriptomic evidence that the anaerobic
pathways in these *Gammaproteobacteria* were activated during environmental anoxia.

Summary of potential adaptations of sponge holobionts to low oxygen

Metabolically, sponge transcription under anoxia was surprisingly similar to normoxia throughout
the holobiont members, and there was no evidence of dormancy within the holobionts in anoxia
(Figure 4). No clear anaerobic respiration responses were observed in the transcriptomes of
sponges, their mitochondria or their two microbial symbionts under environmental anoxia. The
limited significant differential expression of glycolysis genes could indicate that this metabolic
process continues to comparable degrees under normoxia, hypoxia, and anoxia. Therefore, in
the sponges and their microbes, glycolysis could potentially provide ATP without oxygen (Figure
4).

Hypoxia, rather than anoxia, elicited upregulation in energy production machinery such as ATP
synthase (Figure 4). These potential metabolic increases suggest that within the hypoxic limits
studied here (1.30-3.56 mg L^{-1}), sponges perform optimally. Indeed, these limits are above
those at which mortality or substantial physiological changes have been observed for sponges
(Micaroni et al. 2021; Kumala et al. 2021; Mills et al. 2014). This increased energy production
could be stressful, however, and the diversion of resources could limit DNA replication cell
division in hypoxia (Figure 4). Somewhere between the lower limit of hypoxia (1.37 mg L^{-1}) and
the full anoxia studied here, sponges may no longer attempt to increase energy efficiency, and
therefore elevated transcription of ATP synthase genes was not observed in anoxia.

For both the sponges and their *Thaumarcheaota* symbionts, cellular stress responses were
evident under deoxygenation. Both sponge species upregulated HSP genes under anoxia and
hypoxia. These genes may be crucial for sponge deoxygenation tolerance in these species
(Figure 4). We hypothesize that Hsp90 plays a role in sponge oxygen homeostasis, but this
needs to be tested. The *Thaumarchaeota* symbionts upregulated HSP20 and other stress
response genes in anoxia and hypoxia for *H. stellifera* and hypoxia within *Eurypon* sp. 2 (Figure
4). In hypoxia, archaeal and host upregulation of stress response genes may have been a
response to oxidative stress following increased metabolic activities. In anoxia, cellular stress
response might result from the presence of sulfide. Additionally, post-transcriptional, metabolic
changes may play a role in deoxygenation tolerance, and proteomic and metabolomic changes
should be examined in future studies.

Given the largely unchanged metabolic gene expression patterns under hypoxia or anoxia in
these sponges, the production of oxygen by *Thaumarchaeota* could also be crucial to holobiont
survival (noted with a question mark in Figure 4). Although any oxygen produced would be at
extremely low concentrations, it would have a short path of diffusion to the sponge cells, as
Thaumarchaeota symbionts maintain high cell densities within sponges (Moeller et al. 2019). In
addition, mitochondria in general can utilize very low concentrations of oxygen (500 nM) without
changing their oxygen consumption rate, and cells can still respire below that threshold albeit at
a lower rate (Chance 1965; Longmuir 1957; Wilson et al. 1979). In culture, oxygen generated by
Thaumarchaeota reached 100-200 nM (Kraft et al. 2022). If sponge *Thaumarchaeota* produce
oxygen at similar levels under anoxia, cellular respiration could continue in sponges under
anoxia, though likely at lower respiration rates when compared to normoxia. However, given
that the genes responsible are still unknown, the production of oxygen by sponge
Thaumarchaeota needs to be verified experimentally.

Relevance to future ocean deoxygenation

Sponges are predicted to be the 'winners' over other benthic organisms in the future due to their
tolerance of ocean warming and acidification associated with climate change (Bell et al. 2018).
Indeed, sponges have already increased in abundance in some locations while corals have
decreased due to altered carbon and nutrient cycling (Pawlik et al. 2016). The additional
stressor of deoxygenation that comes with climate change and coastal eutrophication has been
relatively overlooked compared to ocean warming and acidification. In other marine taxa, a
meta-analysis demonstrated that the incorporation of deoxygenation as a factor along with
warming could substantially decrease survival times of marine organisms compared to warming
alone (Vaquer-Sunyer and Duarte 2011). Since sponges may be more tolerant than other taxa
to changes in all three stressors of temperature, pH and oxygen, they could indeed become
more abundant in a changing ocean. However, even sponges can be lethally affected by
warming, acidification (Bennett et al. 2017) and deoxygenation (Micaroni et al. 2021) in
scenarios that might occur with climate change, since tolerance to these stressors is species-
specific. So, surviving sponge species in a warmed, acidified, and deoxygenated ocean might
be abundant, but their abundance, and diversity would likely be lower than at present. The
effects of all three of these stressors together need to be understood on a much wider range of
organisms to accurately predict the composition of future oceans. More importantly, the causes
and effects of anthropogenic climate change need to be ameliorated as soon as possible to
avoid such diversity loss.

Relevance for animal evolution

Assuming that the first metazoans were more like an adult sponge than a ctenophore, based on
morphology (Nielsen 2008) and phylogeny (Feuda et al. 2017), then some species of early
sponges or proto-animals could have tolerated low environmental oxygen levels and at least
some periods of benthic anoxia and euxinia in the Neoproterozoic Era (~538.8–1,000 million
years ago) when early animals, including sponges, first evolved (Dohrmann and Wörheide
2017). Lough Hyne sponges are a valuable model system for understanding early animal
evolution. Early sponges could have survived under low and variable oxygen levels using
mechanisms like those of Lough Hyne sponges. Therefore, contrary to longstanding

hypotheses, the evolution of animals probably did not require high and stable levels of environmental oxygen.

Acknowledgements

Field work at Lough Hyne was carried out with the permission of the National Parks and Wildlife Service of Ireland. We thank Luke Harman and Allen Whittaker of University College Cork. We also thank the Nordcee lab technicians of the Department of Biology at University of Southern Denmark for constant support in the laboratory.

Funding

This project was funded by Villum Fonden grant no. 16518. B.K. was funded by the Villum Young Investigator Grant 25491.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

Raw sequence data as well as genome and transcriptome assemblies are available at NCBI under BioProject number PRJNA893197 (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA893197). Scripts, code, and additional data are available here: https://zenodo.org/badge/latestdoi/573040118. Supplemental figures and tables are available on BioRxiv under 'Supplementary material.'

References

Akiva E., Copp J.N., Tokuriki N., Babbitt P.C. 2017. "Evolutionary and Molecular Foundations of Multiple Contemporary Functions of the Nitroreductase Superfamily." *Proceedings of the National Academy of Sciences of the United States of America* 114(45):E9549–58. https://doi.org/10.1073/pnas.1706849114.

Altieri A.H., S.B. Harrison S.B., Seemann J., Collin R., Diaz R.J., Knowlton N. 2017. "Tropical Dead Zones and Mass Mortalities on Coral Reefs." *Proceedings of the National Academy of Sciences of the United States of America* 114 (14): 3660–65. https://doi.org/10.1073/pnas.1621517114.

Bayer K., Busch K., Kenchington E., Beazley L., Franzenburg S., Michels J., Hentschel U., Slaby B.M. 2020. "Microbial Strategies for Survival in the Glass Sponge Vazella Pourtalesii." *mSystems* 5(4):e00473-20. https://doi.org/10.1128/msystems.00473-20.

Bell J.J., Barnes D.K.A. 2000. "The Distribution and Prevalence of Sponges in Relation to Environmental

Gradients within a Temperate Sea Lough: Vertical Cliff Surfaces." *Diversity Distributions* 6:283–303. https://doi.org/10.1046/j.1472-4642.2000.00091.x.

Bell J.J., Bennett H.M., Rovellini A., Webster N.S. 2018. "Sponges to Be Winners under Near-Future Climate Scenarios." *Bioscience* 68(12):955–68. https://doi.org/10.1093/biosci/biy142.

Bennett H.M., Altenrath C., Woods L., Davy S.K, Webster N.S., Bell J.J. 2017. "Interactive Effects of Temperature and pCO2 on Sponges: From the Cradle to the Grave." *Global Change Biology* 23(5): 2031–46. https://doi.org/10.1111/gcb.13474.

Bergquist P. R. 1968. "The Marine Fauna of New Zealand: Porifera, Demospongiae, Part 1 (Tetractinomorpha and Lithistida)". *New Zealand Oceanographic Institute Memoir* 37:27–28.

Bosc L.A.G. 1802. "Histoire Naturelle des Crustacés Contenant leur Description et leurs Moeurs; avec Figures Dessinées, d'Après Nature". Paris: Chez Deterville. 2: 1–296, pls. 6-18., available online at https://www.biodiversitylibrary.org/page/27817320. page(s): 221–222.

Bowerbank J.S. 1864. "A Monograph of the British Spongiadae".Volume 1. Ray Society: London: i-xx, 1-290, pls I-XXXVII., available online at https://biodiversitylibrary.org/page/1883085.

Breitburg D., Levin L.A., Oschlies A., Grégoire M., Chavez F.P., Conley D.J., Véronique Garçon, et al. 2018. "Declining Oxygen in the Global Ocean and Coastal Waters." *Science* 359 (6371): 1–11. https://doi.org/10.1126/science.aam7240.

Chance, B. 1965. "Reaction of Oxygen with the Respiratory Chain in Cells and Tissues." *The Journal of General Physiology* 49 (1): Suppl:163–95. https://doi.org/10.1085/jgp.49.1.163.

Chandel N.S., Schumacker P.T. 2000. "Cellular Oxygen Sensing by Mitochondria: Old Questions, New Insight." *Journal of Applied Physiology* 88(5):1880–89. doi:10.1152/jappl.2000.88.5.1880.

Dendy, Arthur. 1905. "Report on the sponges collected by Professor Herdman, at Ceylon, in 1902". In: Herdman, W.A. (Ed.), Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar. 3 (Supplement 18). Royal Society: London. Pp. 57-246, pls I-XVI. page(s): 143.

Diaz RJ., Rutger R. 2008. "Spreading Dead Zones and Consequences for Marine Ecosystems." *Science* 321 (5891): 926–29. DOI:10.1126/science.1156401.

Dixon G.B., Sarah W. Davies S.W., Aglyamova, G.A. Meyer E., Bay L.K., Matz M.V. 2015. "Coral Reefs. Genomic Determinants of Coral Heat Tolerance across Latitudes." *Science* 348 (6242): 1460–62. DOI: 10.1126/science.1261224.

Dohrmann M., Wörheide G. 2017. "Dating Early Animal Evolution Using Phylogenomic Data." *Scientific Reports* 7(1):1–6. https://doi.org/10.1038/s41598-017-03791-w.

D'Souza A.R., Minczuk M. 2018. "Mitochondrial Transcription and Translation: Overview." *Essays in Biochemistry* 62(3):309–320. https://doi.org/10.1042/ebc20170102.

Edler D., Klein J., Antonelli A., Silvestro D. 2020. "raxmlGUI 2.0: A Graphical Interface and Toolkit for Phylogenetic Analyses Using RAxML." *Methods in Ecology and Evolution* 12:373–77. https://doi.org/10.1111/2041-210X.13512.

Efremova S.M., Margulis B.A., Guzhova I.V., Itskovich V.B., Lauenroth S., Müller W.E.G., Schröder H.C. 2002. "Heat Shock Protein Hsp70 Expression and DNA Damage in Baikalian Sponges Exposed to Model Pollutants and Wastewater from Baikalsk Pulp and Paper Plant." *Aquatic Toxicology* 57(4):267–80. DOI:10.1016/s0166-445x(01)00209-0. PMID: 11932006.

Ellington W. R. 1982. "Metabolic Responses of the Sea Anemone Bunodosoma Cavernata (Bosc) to Declining Oxygen Tensions and Anoxia." *Physiological Zoology* 55 (3): 240–49. https://www.jstor.org/stable/30157888.

Feuda R., Dohrmann M., Pett W., Philippe H., Rota-Stabelli O., Lartillot N., Wörheide G., Pisani D. 2017. "Improved Modeling of Compositional Heterogeneity Supports Sponges as Sister to All Other Animals." *Current Biology: CB* 27 (24): 3864–70.e4. https://doi.org/10.1016/j.cub.2017.11.008.

Francis, W. 2020a. "General Sequence Processing Scripts." *Bitbucket.* 2020. bitbucket.org/wrf/sequences.

Francis, W. 2020b. "Make Interactive Plot of Contigs in Genomes or Metagenomes." *Gihub.* 2020. https://github.com/wrf/lavaLampPlot/blob/master/Rshiny/app.R.

Gong Z-J, Wu Y-Q., Miao J., Duan Y., Jiang Y-L., Li T. 2013. "Global Transcriptome Analysis of Orange Wheat Blossom Midge, Sitodiplosis Mosellana (Gehin) (Diptera: Cecidomyiidae) to Identify Candidate Transcripts Regulating Diapause." *PloS One* 8 (8): e71564. https://doi.org/10.1371/journal.pone.0071564.

Görlach A., Dimova E.V., Petry A., Martínes-Ruiz A., Hernansan-Agustín P., Rolo A.P., Palmeira C.M., Kietzmann T. 2015. "Reactive oxygen species, nutrition, hypoxia and diseases: Problems solved?" Redox Biology 6:372–385. https://doi.org/10.1016/j.redox.2015.08.016.

Grabherr M.G., Haas B.J., Yassour M., Levin J.Z., Thompson D.A., Amit I., Adiconis X., et al. 2011. "Full-Length Transcriptome Assembly from RNA-Seq Data without a Reference Genome." *Nature Biotechnology* 29:644–652. https://doi.org/10.1038/nbt.1883.

Grant, R.E. 1836. Animal Kingdom. Pp. 107-118. In: Todd, R.B. (Ed.), "The Cyclopaedia of Anatomy and Physiology". Volume 1. Sherwood,Gilbert, and Piper: London: 1-813.

Grant R.E. 1826. "Notice of a New Zoophyte (*Cliona celata* Gr.) from the Firth of Forth". Edinburgh *New Philosophical Journal* 1: 78-81., available online at https://www.biodiversitylibrary.org/page/2471191. page(s): 81.

Grasshoff K., Kremling K., Ehrhardt M. 1999. "Methods of Seawater Analysis", 3rd ed. Wiley-VCH, New York 600p.

Gunda V.G., Janapala V.R. 2009. "Effects of Dissolved Oxygen Levels on Survival and Growth in Vitro of *Haliclona pigmentifera* (Demospongiae)." *Cell and Tissue Research* 337(3):527–35. https://doi.org/10.1007/s00441-009-0843-5.

Guzman C., Conaco C. 2016. "Gene Expression Dynamics Accompanying the Sponge Thermal Stress Response." *PloS One* 11(10):e0165368. https://doi.org/10.1371/journal.pone.0165368.

Hochachka P.W., Lutz P.L. 2001. "Mechanism, Origin, and Evolution of Anoxia Tolerance in Animals." *Comparative Biochemistry and Physiology. Part B, Biochemistry & Molecular Biology* 130 (4): 435–59. https://doi.org/10.1016/S1096-4959(01)00408-0.

Hoffmann F., Larsen O., Thiel V., Rapp H.T., Pape T., Michaelis W., Reitner J. 2005. "An Anaerobic World in Sponges." *Geomicrobiology Journal* 22 (1-2): 1–10. https://doi.org/10.1080/01490450590922505.

Hoffmann F., Radax R., Woebken D., Holtappels M., Lavik G., Rapp H.T., Schläppy M-L., Schleper C., Kuypers M.M.M. 2009. "Complex Nitrogen Cycling in the Sponge *Geodia barretti*." *Environmental Microbiology* 11 (9): 2228–43. https://doi.org/10.1111/j.1462-2920.2009.01944.x.

Huerta-Cepas J., Szklarczyk D., Heller D., Hernández-Plaza A, Forslund S.K., Cook H., Mende D.R., et al. 2019. "eggNOG 5.0: A Hierarchical, Functionally and Phylogenetically Annotated Orthology Resource Based on 5090 Organisms and 2502 Viruses." *Nucleic Acids Research* 47(D1):D309–D314. https://doi.org/10.1093/nar/gky1085.

Hughes D.J., Alderdice R., Cooney C., Kühl M., Pernice M., Voolstra C.R., Suggett D.J. 2020. "Coral Reef Survival under Accelerating Ocean Deoxygenation." *Nature Climate Change* 10 (4): 296–307. https://doi.org/10.1038/s41558-020-0737-9.

Huntley, M.A., Larson J.L., Chaivorapol C., Becker G., Lawrence M., Hackney J.A., Kaminker J.S. 2013. "ReportingTools: An Automated Result Processing and Presentation Toolkit for High-Throughput Genomic Analyses." *Bioinformatics* 29(24):3220–21. https://doi.org/10.1093/bioinformatics/btt551.

Johnston, Georg. 1842. "A History of British Sponges and Lithophytes". (W.H. Lizars: Edinburgh). i-xii, 1-264, pls I-XXV., available online at https://biodiversitylibrary.org/page/35290582. page(s): 146-147.

Jormakka Mika., Byrne B., Iwata S. 2003. "Formate Dehydrogenase – a Versatile Enzyme in Changing Environments." *Current Opinion in Structural Biology* 13(4):418–23. https://doi.org/10.1016/S0959-440X(03)00098-8.

Katoh K., Standley D.M. 2013. "MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability." *Molecular Biology and Evolution* 30(4):772–80. https://doi.org/10.1093/molbev/mst010

Keeling R.E., Körtzinger A., Gruber N. 2010. "Ocean Deoxygenation in a Warming World." *Annual Review of Marine Science* 2: 199–229. https://doi.org/10.1146/annurev.marine.010908.163855.

Kelly-Borges M., Bergquist P.R. 1997. "Revision of Southwest Pacific Polymastiidae (Porifera: Demospongiae: Hadromerida) with descriptions of new species of Polymastia Bowerbank, Tylexocladus Topsent, and Acanthopolymastia gen. nov. from New Zealand and the Norfolk Ridge, New Caledonia". *Zealand Journal of Marine and Freshwater Research.* 31(3): 378–380. https://doi.org/10.1080/00288330.1997.9516772.

Kirchman D.L. 2021. "Dead Zones: The Loss of Oxygen from Rivers, Lakes, Seas, and the Ocean". *Oxford University Press,* USA. 1–224.

Kolde R. 2013. *Pheatmap: Pretty Heatmaps. R-Package* (version 0.7.4). http://www2.uaem.mx/r-mirror/web/packages/pheatmap/pheatmap.pdf.

Kraft B., Jehmlich N., Larsen M., Bristow L.A., Könneke M., Thamdrup B., Canfield D.E. 2022. "Oxygen and Nitrogen Production by an Ammonia-Oxidizing Archaeon." *Science* 375 (6576): 97–100. DOI: 10.1126/science.abe673.

Kraft B., Jehmlich N., Larsen M., Bristow L.A., Könneke M., Thamdrup B., Canfield D.E. 2021. "Oxygen

Production by an Ammonia-Oxidizing Archaeon." *bioRxiv*.
https://doi.org/10.1101/2021.04.01.436977.
Kumala L., Larsen M., Glud R.N., Canfield D.E. 2021. "Spatial and temporal anoxia in single-osculum
Halichondria panicea demosponge explants studied with planar optodes." *Marine Biology* 168:173.
https://doi.org/10.1007/s00227-021-03980-2.
Langmead B., Salzberg S.L. 2012. "Fast Gapped-Read Alignment with Bowtie 2." *Nature Methods*
9(4):357–59. https://doi.org/10.1038/nmeth.1923.
Larade K., Kenneth B.S. 2009. "Living without Oxygen: Anoxia-Responsive Gene Expression and
Regulation." *Current Genomics* 10(2):76–85. DOI: 10.2174/138920209787847032.
Lemmens L., Maklad H.R., Bervoets I., Peeters E. 2019. "Transcription Regulators in Archaea:
Homologies and Differences with Bacterial Regulators." *Journal of Molecular Biology* 431(20): 4132–
46. https://doi.org/10.1016/j.jmb.2019.05.045.
Lévi C. 1953. "Sur une nouvelle classification des Démosponges". *Compte rendu hebdomadaire des*
séances de l'Académie des sciences, Paris. 236 (8): 853–855.
Levin L.A. 2018. "Manifestation, Drivers, and Emergence of Open Ocean Deoxygenation." *Annual Review*
of Marine Science 10:229–60. https://doi.org/10.1146/annurev-marine-121916-063359.
Leys S.P., Kahn A.S. 2018. "Oxygen and the Energetic Requirements of the First Multicellular Animals."
Integrative and Comparative Biology 58(4): 666–676. https://doi.org/10.1093/icb/icy051
Li D., Luo R., Liu C-M., Leung C-M, Ting H-F, Sadakane K., Yamashita H., Lam T-W.. 2016. "MEGAHIT
v1.0: A Fast and Scalable Metagenome Assembler Driven by Advanced Methodologies and
Community Practices." *Methods* 102:3–11. https://doi.org/10.1016/j.ymeth.2016.02.020.
Li Z., Srivastava P. 2004. "Heat-Shock Proteins." *Current Protocols in Immunology* 58(1):A.1T.1–A.1T.6
https://doi.org/10.1002/0471142735.ima01ts58.
Longmuir I.S. 1957. "Respiration Rate of Rat-Liver Cells at Low Oxygen Concentrations." *Biochemical*
Journal 65 (2):378–82. https://doi.org/10.1042/bj0650378.
López-Legentil S., Song B., McMurray S.E., Pawlik J.R. 2008. "Bleaching and Stress in Coral Reef
Ecosystems: hsp70 Expression by the Giant Barrel Sponge Xestospongia Muta." *Molecular Ecology*
17(7):1840–49. https://doi.org/10.1111/j.1365-294X.2008.03667.x.
Löscher C.R., Kock A., Könneke M., LaRoche J., Bange H.W., Schmitz R.A. 2012. "Production of
Oceanic Nitrous Oxide by Ammonia-Oxidizing Archaea." *Biogeosciences* 9(July):2419–29.
https://doi.org/10.5194/bg-9-2419-2012
Love M.I., Huber W., Anders S. 2016. "Moderated Estimation of Fold Change and Dispersion for RNA-
Seq Data with DESeq2." *BMC Genome Biology* 15:1–22. https://doi.org/10.1186/s13059-014-0550-
8.
Lynch M., Marinov G.K. 2015. "The Bioenergetic Costs of a Gene." *Proceedings of the National Academy*
of Sciences of the United States of America 112(51):15690–95.
https://doi.org/10.1073/pnas.151497411.
MacRae T.H. 2010. "Gene Expression, Metabolic Regulation and Stress Tolerance during Diapause."
Cellular and Molecular Life Sciences: CMLS 67(14):2405–24. https://doi.org/10.1007/s00018-010-
0311-0.
Magnet S., Dubost L., Marie A,. Arthur M., Gutmann L. 2008. "Identification of the L,D-transpeptidases for
peptidoglycan cross-linking in Escherichia coli". *J Bacteriol.* 190(13):4782-5.
https://doi.org/10.1128/jb.00025-08
Matz M.V. 2015. "Assembly and Annotation of Transcriptomes de Novo." Github. 2015.
https://github.com/z0on/annotatingTranscriptomes.
Medini H., Cohen T., Mishmar D. 2021. "Mitochondrial gene expression in single cells shape pancreatic
beta cells´ subpopulations and explain variation in insulin pathway. *Scientific Reports* 11:466.
https://doi.org/10.1038/s41598-020-80334-w.
Meyer E. 2016. "A Collection of Scripts for Analyzing Transcriptome Assemblies." Github. 2016.
https://github.com/Eli-Meyer/transcriptome_utilities.
Micaroni V., McAllen R., Turner J., Strano F., Morrow C., Picton B., Harman L., Bell J.J. 2021a.
"Vulnerability of Temperate Mesophotic Ecosystems (TMEs) to Environmental Impacts: Rapid
Ecosystem Changes at Lough Hyne Marine Nature Reserve, Ireland." *Science of The Total*
Environment 789:1–15. https://doi.org/10.1016/j.scitotenv.2021.147708.
Micaroni V., Strano F., McAllen R., Woods L., Turner J., Harman L., Bell J.J. 2021b. "Adaptive Strategies
of Sponges to Deoxygenated Oceans." *Global Change Biology* 28(6): 1972–1989.
https://doi.org/10.1111/gcb.16013.

Mills D.B., Francis W.R., Vargas, S., Larsen M., Elemans C.P., Canfield D.E., Wörheide G. 2018. "The Last Common Ancestor of Animals Lacked the HIF Pathway and Respired in Low-Oxygen Environments." *eLife* 7:e31176. https://doi.org/10.7554/eLife.31176.

Mills D.B., Ward L.M., Jones C., Sweeten B., Forth M., Treusch A.H., Canfield D.E. 2014. "Oxygen Requirements of the Earliest Animals." *Proceedings of the National Academy of Sciences of the United States of America* 111 (11): 4168–72. https://doi.org/10.1073/pnas.1400547111.

Moeller F.U., Webster N.S., Herbold C.W., Behnam F., Domman D., Albertsen M., Maria Mooshammer M., et al. 2019. "Characterization of a Thaumarchaeal Symbiont That Drives Incomplete Nitrification in the Tropical Sponge *Ianthella basta*." *Environmental Microbiology* 21 (10): 3831–54. https://doi.org/10.1111/1462-2920.14732.

Moitinho-Silva L., Díez-Vives C., Batani G., Esteves, A.I.S., Jahn, M.T., Thomas T. 2017. "Integrated metabolism in sponge-microbe symbiosis revealed by genome-centered metatranscriptomics." *ISME Journal* 11(7), 1651–1666. https://doi.org/10.1038/ismej.2017.25

Nielsen C. 2008. "Six Major Steps in Animal Evolution: Are We Derived Sponge Larvae?" *Evolution & Development* 10(2):241–57. https://doi.org/10.1111/j.1525-142X.2008.00231.x.

Pallas P.S. 1766. "Elenchus zoophytorum sistens generum adumbrationes generaliores et specierum cognitarum succinatas descriptiones, cum selectis auctorum synonymis". Fransiscum Varrentrapp, Hagae. pp. 451., available online at https://doi.org/10.5962, page(s): 388.

Pawlik J.R., Burkepile D.E., Thurber R.V. 2016. "A Vicious Circle? Altered Carbon and Nutrient Cycling May Explain the Low Resilience of Caribbean Coral Reefs." *BioScience*. https://doi.org/10.1093/biosci/biw047.

Plese B., Rossi M.E., Kenny N.J., Taboada S., Koutsouveli V., Riesgo A. 2019. "Trimitomics: An Efficient Pipeline for Mitochondrial Assembly from Transcriptomic Reads in Nonmodel Species." *Molecular Ecology Resources* 19(5):1230–39. https://doi.org/10.1111/1755-0998.13033.

R Core Team. 2016. *R: A Language and Environment for Statistical Computing. Vienna, Austria*. https://www.R-project.org/.

Reznik E., Wang Q., La K., Schultz N., Sander C. "Mitochondrial respiratory gene expression is suppressed in many cancers". eLife. 2017;6:e21592. doi:10.7554/eLife.21592.

Riesgo A., Farrar N., Windsor P.J., Giribet G., Leys S.P. 2014. "The Analysis of Eight Transcriptomes from All Poriferan Classes Reveals Surprising Genetic Complexity in Sponges." *Molecular Biology and Evolution* 31 (5): 1102–20. https://doi.org/10.1093/molbev/msu057.

RStudio Team. 2020. *RStudio: Integrated Development for R* (version 4.0.3). X86_64, darwin17.0 . https://www.rstudio.com/products/rstudio/.

Samenza G.L. 2007. "Life with Oxygen". *Science* 318(5847): 62–64. https://www.science.org/doi/10.1126/science.1147949

Sassaman C., Mangum C.P. 1972. "Adaptations To Environmental Oxygen Levels In Infaunal And Epifaunal Sea Anemones." *The Biological Bulletin* 143 (3): 657–78. https://doi.org/10.2307/1540189.

Schmidt O. 1870. "Grundzüge einer Spongien-Fauna des atlantischen Gebietes". Wilhelm Engelmann: Leipzig: iii-iv, 1-88, pls I-VI. page(s): 14-15; pl I fig 1–6.

Schröder H.C., Efremova S.M., Margulis B.A., Guzhova I.V., Itskovich V.B., Müller W.E.G. 2006. "Stress Response in Baikalian Sponges Exposed to Pollutants." *Hydrobiologia* 568 (1): 277–87. https://doi.org/10.1007/s10750-006-0302-1.

Schurch N.J., Schofield P., Gierliński M., Cole C., Sherstnev A., Singh V., Wrobel N., Gharbi K., Simpson G.G., Owen-Hughes T., Blaxter M., Barton G.J. 2016. "How many biological replicates are needed in an RNA-seq experiment and which differential expression tool should you use?" *RNA* 22(6):839-51. doi: 10.1261/rna.053959.115.

Schuster A., Strehlow B.W., Eckford-Soper L., McAllen R., Canfield D.E. 2021. "Effects of Seasonal Anoxia on the Microbial Community Structure in Demosponges in a Marine Lake in Lough Hyne, Ireland." *mSphere* 6(1):e00991–20. https://doi.org/10.1128/msphere.00991-20.

Simão F. A., Waterhouse R.M., Ioannidis P., Kriventseva E.V., Zdobnov E.M. 2015. "BUSCO: Assessing Genome Assembly and Annotation Completeness with Single-Copy Orthologs." *Bioinformatics* 31(19):3210–12. https://doi.org/10.1093/bioinformatics/btv351.

Sinha A., Sommer R.J., Dieterich C. 2012. "Divergent Gene Expression in the Conserved Dauer Stage of the Nematodes Pristionchus Pacificus and Caenorhabditis Elegans." *BMC Genomics* 13, 254. https://doi.org/10.1186/1471-2164-13-254.

Sollai M., Villanueva L., Hopmans E.C., Reichart G-J., Damsté J.S.S. 2019. "A combined lipidomic and 16S rRNA gene amplicon sequencing approach reveals archaeal sources of intact polar lipids in the

stratified Black Sea water column." *Geobiology* 17(1):91–109. https://doi.org/10.1111/gbi.12316.

Stachowitsch M. 1984. "Mass Mortality in the Gulf of Trieste: The Course of Community Destruction." *Marine Ecology* 5 (3):243–64. https://doi.org/10.1111/j.1439-0485.1984.tb00124.x.

Strader M.E., Aglyamova G.V., Matz M.V. 2016. "Red Fluorescence in Coral Larvae Is Associated with a Diapause-like State." *Molecular Ecology* 25(2):559–69. https://doi.org/10.1111/mec.13488.

Stramma L., Johnson G.C., Sprintall J., Mohrholz F. 2008. "Expanding Oxygen-Minimum Zones in the Tropical Oceans." *Science* 320 (5876): 655–58. DOI: 10.1126/science.115384.

Strehlow B.W., Pineda M-C., Kenkel C.D., Laffy P., Duckworth A., Renton M., Clode P.L., Webster N.S. 2020. "Novel Reference Transcriptomes for the Sponges Carteriospongia Foliascens and Cliona Orientalis and Associated Algal Symbiont Gerakladium Endoclionum." *Coral Reefs* 40:9–13. https://doi.org/10.1007/s00338-020-02028-z

Stryer L. 1995. "Biochemistry (Fourth ed.)". New York: W.H. Freeman and Company. pp. 559–565, 574–576, 614–623. ISBN 0-7167-2009-4.

Topsent E. 1928. "Spongiaires de l'Atlantique et de la Méditerranée provenant des croisières du Prince Albert ler de Monaco". *Résultats des campagnes scientifiques accomplis par le Prince Albert I. Monaco.* 74:1-376, pls I-XI.

Vaquer-Sunyer R., Duarte C.M. 2011. "Temperature Effects on Oxygen Thresholds for Hypoxia in Marine Benthic Organisms." *Global Change Biology*. https://doi.org/10.1111/j.1365-2486.2010.02343.x.

Vargas S., Caglar C., Büttner G., Schätzle S., Deistler F., Wörheide G. 2021. "Slime Away: A Simple CTAB-Based High Molecular Weight DNA and RNA Extraction Protocol for 'Difficult' Invertebrates." Protocols.io. 2021. https://www.protocols.io/view/slime-away-a-simple-ctab-based-high-molecular-weig-bwcwpaxe.

Webster N.S., Taylor M.W. 2012. "Marine Sponges and Their Microbial Symbionts: Love and Other Relationships." *Environmental Microbiology* 14(2):335–46. https://doi.org/10.1111/j.1462-2920.2011.02460.x.

Webster N., Pantile R., Botté E., Abdo D., Andreakis N., Whalan S. 2013. "A Complex Life Cycle in a Warming Planet: Gene Expression in Thermally Stressed Sponges." *Molecular Ecology* 22(7):1854–68. https://doi.org/10.1111/mec.12213.

Welchen E., Gonzalez D.H. 2016. "Cytochrome c, a hub linking energy, redox, stress and signaling pathways in mitochondria and other cell compartments". *Physiol Plant.*157(3):310-21. https://doi.org/10.1111/ppl.12449.

Wilkinson C.R. 1978. "Microbial Associations in Sponges. II. Numerical Analysis of Sponge and Water Bacterial Populations." *Marine Biology* 49 (2): 169–76. https://doi.org/10.1007/BF00387116.

Wilson D.F., Erecińska M., Drown C., Silver I.A. 1979. "The Oxygen Dependence of Cellular Energy Metabolism." *Archives of Biochemistry and Biophysics* 195(2):485–93. https://doi.org/10.1016/0003-9861(79)90375-8.

Wright, R.M., Aglyamova G.V., Meyer E., Matz M.V. 2015. "Gene Expression Associated with White Syndromes in a Reef Building Coral, *Acropora hyacinthus*." *BMC Genomics* 16, 371 https://doi.org/10.1186/s12864-015-1540-2.