SARS-CoV-2 ORF3c suppresses immune activation by inhibiting innate sensing

Martin Müller[1], Alexandra Herrmann[2], Shigeru Fujita[3,4], Keiya Uriu[3,4], Carolin Kruth[1], Adam Strange[3], Jan Eric Kolberg[1], Markus Schneider[1], Jumpei Ito[3], Armin Ensser[2], The Genotype to Phenotype Japan (G2P-Japan) Consortium, Kei Sato[3,4,5,6,7,8], Daniel Sauter[1,3,9,*]

[1]Institute for Medical Virology and Epidemiology of Viral Diseases, University Hospital Tübingen, 72076 Tübingen, Germany

[2]Institute for Clinical and Molecular Virology, University Hospital, Friedrich-Alexander-Universität Erlangen-Nürnberg, 91054 Erlangen, Germany

[3]Division of Systems Virology, Department of Microbiology and Immunology, The Institute of Medical Science, The University of Tokyo, Minato-ku, Tokyo, Japan

[4]Graduate School of Medicine, The University of Tokyo, Minato-ku, Tokyo, Japan

[5]International Research Center for Infectious Diseases, The Institute of Medical Science, The University of Tokyo, Minato-ku, Tokyo, Japan

[6]International Vaccine Design Center, The Institute of Medical Science, The University of Tokyo, Minato-ku, Tokyo, Japan

[7]Graduate School of Frontier Sciences, The University of Tokyo, Kashiwa, Chiba, Japan

[8]CREST, Japan Science and Technology Agency, Kawaguchi, Saitama, Japan

[9]Lead contact

Correspondence: daniel.sauter@med.uni-tuebingen.de

ABSTRACT

SARS-CoV-2 proteins are translated from subgenomic RNAs (sgRNAs). While most of these

sgRNAs are monocistronic, some viral mRNAs encode more than one protein. For example,

the *ORF3a* sgRNA also encodes ORF3c, an enigmatic 41-amino acid peptide. Here, we show

that ORF3c suppresses RIG-I- and MDA5-mediated immune activation and interacts with the

signaling adaptor MAVS. In line with this, ORF3c inhibits IFN-β induction. This

immunosuppressive activity of ORF3c is conserved among members of the subgenus

sarbecovirus, including SARS-CoV and coronaviruses isolated from bats. Notably, however,

the SARS-CoV-2 delta and kappa variants harbor premature stop codons in ORF3c

demonstrating that this reading frame is not essential for efficient viral replication *in vivo*. In

agreement with this, disruption of ORF3c did not significantly affect SARS-CoV-2 replication

in CaCo-2 or CaLu-3 cells. In summary, we here identify ORF3c as an immune evasion factor

that suppresses IFN-β induction, but is dispensable for efficient replication of SARS-CoV-2.

INTRODUCTION

Since the emergence of the COVID-19 pandemic, all canonical proteins of SARS-CoV-2 have been extensively characterized for their expression, structure and function. In addition to its prototypical genes, however, SARS-CoV-2 harbors several smaller open reading frames (ORFs) that frequently overlap with other ORFs and may also contribute to efficient viral replication. For example, the ORF3b peptide encoded by *ORF3a* subgenomic RNA (sgRNA) was shown to suppress the induction of type I interferon (IFN) (Konno et al., 2020). Intriguingly, naturally occurring variants of ORF3b differ in their immunosuppressive activity and may be responsible for phenotypic differences between SARS-CoV and SARS-CoV-2 (Konno *et al.*, 2020). Moreover, several short upstream ORFs (uORFs) have been suggested to regulate translation of downstream genes such as *ORF7b* (Finkel et al., 2021). Thus, non-canonical ORFs of SARS-CoV-2 may also be important determinants of viral immune evasion, spread and/or pathogenicity.

Nevertheless, most of the cryptic ORFs of SARS-Cov-2 remain poorly characterized, and several open questions remain: Do they encode proteins or are they merely a result of selection pressures acting on overlapping reading frames? Do these ORFs exert any regulatory activity, e.g. by modulating translation of downstream ORFs via leaky scanning or ribosomal re-initiation? Do they code for functional proteins that contribute to efficient immune evasion and/or replication of SARS-CoV-2? Are the respective peptides or proteins immunogenic?

One interesting cryptic open reading frame is *ORF3c*, located at nt 25457-25579 of the Wuhan-Hu-1 reference genome. This ORF was independently described by different groups and has received several alternative names: ORF3c (Firth, 2020; Jungreis et al., 2021b), ORF3h (for hypothetical) (Cagliani et al., 2020), 3a.iORF1 (Finkel *et al.*, 2021), and ORF3b (Pavesi, 2020). Following the homology-based nomenclature proposed by Jungreis and colleagues (Jungreis et al., 2021a), we will refer to this open reading frame as *ORF3c* hereafter. Like *ORF3b*, *ORF3c* is one of several open reading frames overlapping with *ORF3a*. *In silico* analyses suggested

that the respective ORF3c protein may harbor a transmembrane domain (Firth, 2020) and act

as a viroporin (Cagliani *et al.*, 2020). Nevertheless, its exact function and relevance for viral

replication have remained unclear.

Here, we show that SARS-CoV-2 ORF3c encodes a stable 41-amino acid peptide that

suppresses the induction of IFN-β expression. Mechanistic analyses revealed that it inhibits

innate sensing induced by RIG-I and MDA5, and interacts with the downstream adaptor protein

MAVS. In line with a relevant function *in vivo*, ORF3c orthologs from different sarbecoviruses

share this immunosuppressive activity. However, we also identify SARS-CoV-2 lineages that

spread efficiently in the human population despite premature stop codons in their *ORF3c* genes.

Furthermore, disruption of *ORF3c* did not affect SARS-CoV-2 replication in CaCo-2 and

CaLu-3 cells. Thus, our findings identify ORF3c as an immune evasion factor that inhibits

innate sensing cascades, but is not essential for efficient viral replication.

RESULTS

SARS-CoV-2 *ORF3c* encodes a peptide suppressing IFN-β promoter activation

The *ORF3a* gene of the SARS-CoV-2 reference genome Wuhan-Hu-1 overlaps with several

shorter open reading frames that have the potential to encode for peptides of at least 10 amino

acids in length (Fig. 1A). Translation initiation downstream of the start codon of *ORF3a* may

be enabled via non-canonical translation mechanisms, such as leaky scanning, ribosomal

shunting and/or re-initiation (Firth and Brierley, 2012). In line with this, the start codon of

ORF3c is part of a strong Kozak sequence, and *in silico* analyses predict ORF3c expression

from the ORF3a sgRNA via leaky scanning (Gleason et al., 2022) (Fig. 1B). Indeed, ribosome

profiling studies confirmed that *ORF3c* is translated in SARS-CoV-2-infected cells (Finkel *et*

al., 2021). The same study also found evidence for translation of *ORF3d-2* (Finkel *et al.*, 2021),

which is in agreement with the detection of ORF3d-specific antibodies in sera from previously

SARS-CoV-2-infected individuals (Hachim et al., 2022). In contrast, we found no evidence for

antibodies against ORF3c in SARS-CoV-2 convalescent sera (Fig. S1)

To characterize the stability and potential activity of cryptic ORF3 peptides, we generated

expression vectors for the individual peptides harboring a C-terminal HA-tag (without codon-

optimization). Apart from the *ORF3a* construct, *ORF3c* and *ORF3d/ORF3d-2* code for stable

proteins that are readily detectable in transfected cells (Fig. 1C). The remaining peptides were

not (ORF3b-2, ORF3b-3, ORF3b-4) or only poorly (ORF3b, ORF3e) detectable. Together with

the ribosome profiling study and detection of antibodies in convalescent sera, this demonstrates

that the *ORF3a* sgRNA encodes at least three additional stable peptides, ORF3c and ORFd-2

in SARS-CoV-2 infected cells.

Since several accessory proteins of SARS-CoV-2 (e.g., ORF3b, ORF6) have been shown to

suppress the induction of interferons (Kimura et al., 2021; Konno *et al.*, 2020; Miorin et al.,

2020), we hypothesized that some of the cryptic ORF3 peptides may exert similar immune-

modulatory activities. Indeed, a luciferase reporter assay revealed that ORF3c significantly

suppresses the activation of the IFN-β promoter in response to a constitutively active mutant of

the pattern recognition receptor RIG-I (Fig. 1D). Notably, ORF3c was also more active than

the previously described IFN antagonist ORF3b, which suppressed IFN-β promoter activity

only at higher concentrations or upon codon-optimization (Konno *et al.*, 2020) (data not

shown). To test whether ORF3c is able to suppress immune activation upon viral infection, we

monitored endogenous *IFNB1* expression upon infection with Sendai virus (SeV), a potent

inducer of RIG-I-mediated type I IFN expression (Strahle et al., 2006). As expected, SeV

induced *IFNB1* expression in a dose-dependent manner (Fig. 1E). However, *IFNB1* mRNA

levels were reduced by about 70% in the presence of SARS-CoV-2 ORF3c. Together, these

findings demonstrate that SARS-CoV-2 *ORF3c* encodes a stable peptide that suppresses the

production of IFN-β upon viral sensing.

ORF3c interacts with MAVS and suppresses both RIG-I- and MDA5-mediated immune activation

To elucidate the mechanisms underlying the inhibitory activity of ORF3, we analyzed different steps of the innate RNA sensing cascade culminating in the induction of IFN-β expression (Fig. 2A). The *IFNB1* promoter harbors binding sites for both IRF3 and NF-κB. As expected, disruption of the NF-κB binding site reduced responsiveness to RIG-I-mediated activation (Fig. 2B). However, ORF3c still dose-dependently reduced promoter activation (Fig. 2B), demonstrating that ORF3c does not selectively target NF-κB activation. Next, we activated the sensing cascade at different steps via over-expressing MDA5, MAVS or a constitutively active mutant of IRF3. While ORF3c suppressed MDA5-mediated immune activation (Fig. 2C, left panel), it failed to efficiently suppress *IFNB1* promoter activation in response to MAVS or IRF3 over-expression (Fig. 2C, middle and right panels). Together, these findings suggest that ORF3c targets immune activation upstream or at the level of the signaling adaptor MAVS. In line with this, SARS-CoV-2 ORF3c weakly co-immunoprecipitated with MAVS, while we found no evidence for an interaction with RIG-I, MDA5 or TBK1 (Fig. 2D). The modest co-immunoprecipitation of ORF3c was also observed when a MAVS mutant lacking its CARD domain was used for pull-down (Fig. 2E), demonstrating that this domain is dispensable for the interaction. To map residues in ORF3c involved in its immunosuppressive activity, we performed an alanine scan (Fig. 2F). While all ORF3c mutants tested still reduced IFN-β promoter activity, the double mutants L2A/L3A and I6A/L7A significantly reduced the immunosuppressive effect of SARS-Cov-2 ORF3c. In summary, these results identify residues in the N-terminus of ORF3c that contribute to its inhibitory activity and demonstrate that ORF3c suppresses IFN-β expression independently of the pattern recognition receptor, i.e. downstream of RIG-I and MDA5.

The immunosuppressive activity of ORF3c is conserved among sarbecoviruses

In silico analyses of the ORF3 locus revealed that essentially all sarbecoviruses harbor an

ORF3c gene encoding a 40- or 41-amino-acid peptide (Firth, 2020). In contrast, the remaining

ORF3 genes are only poorly conserved or vary substantially in their length (Fig. 3A). The

absence of *ORF3c* from other subgenera of betacoronaviruses suggests that this open reading

frame emerged after the divergence of sarbeco- and hibecoviruses. To test whether the

immunosuppressive activity of ORF3c is also conserved, we characterized several orthologs

representing human and bat isolates of the SARS-CoV- and SARS-CoV-2-like clusters (Fig.

3B). Titration experiments revealed that all ORF3c peptides tested significantly suppress IFN-β

promoter activation (Fig. 3C). However, ORF3c of SARS-CoV-2 Wuhan Hu-1 and a closely

related bat coronavirus (ZXC21) suppressed IFN-β promoter activation more efficiently than

ORF3c of SARS-CoV Tor2 and SARS-CoV-2 BANAL-20-50. In summary, these findings

demonstrate that not only the *ORF3c* gene itself, but also the immunosuppressive activity of

the respective protein is conserved among the sarbecovirus subgenus of betacoronaviruses.

A natural R36I polymorphism does not affect the immunosuppressive activity of ORF3c

Since the emergence of SARS-CoV-2 in 2019, several mutations have occurred throughout the

viral genome. One notable mutation is G25563T (Fig. 4A), which is found in the beta, eta, iota

and mu variants and has been associated with increased transmission fitness (Oulas et al., 2021).

Although G25563T simultaneously introduces non-synonymous mutations in ORF3a (Q57H),

ORF3c (R36I) and ORF3d (E31*) (Fig. 4A), previous studies focused only on possible

phenotypic consequences of the Q57H change in ORF3a. Notably, however, the R36I mutation

in ORF3c is predicted to result in a conformational change (Fig. 4B) and a transmembrane

domain in the C-terminal half of ORF3c (Fig. 4C). We therefore analyzed whether the R36I

change may affect the subcellular localization and/or immune-suppressive activity of SARS-

CoV-2 ORF3c. Immunofluorescence microscopy revealed that ORF3c of the Wuhan-Hu-1

reference strain and a R36I mutant thereof are similarly distributed throughout the cytoplasm

(Fig. 4D). Moreover, both ORF3c variants dose-dependently suppressed RIG-I-mediated IFN-β

promoter activation to a similar extent (Fig. 4E). Thus, the G25563T polymorphism of the

SARS-CoV-2 beta, eta, iota and mu variants does not seem to alter the IFN-suppressive activity

of ORF3c.

Some SARS-CoV-2 variants harbor premature stop codons in *ORF3c*

To better understand the relevance of an intact *ORF3c* gene for viral spread, we screened the

GISAID SARS-CoV-2 sequence repository for PANGO (sub)lineages harboring a premature

ORF3c stop codon in at least 20% of the isolates. We identified two mutations fulfilling these

criteria: the first one (C25469T) introduces a premature stop codon (Q5*) in ORF3c and an

S26L change in ORF3a (Fig. 4F). It is present in about 44% of B.1.617 isolates and in almost

all sequences of the B.1.617.1 (delta) and B.1.617.2 (kappa) sublineages (Fig. 4G). The second

mutation (del25498-25530) represents an in-frame deletion in ORF3a and can be found in about

80% of all B.1.630 isolates. This deletion results in the loss of the initiation codon of ORF3d

and a premature stop codon (Y14*) in ORF3c. The presence of premature ORF3c stop codons

in a substantial fraction of B.1.617 and B.1.630 lineages suggests that ORF3c is dispensable for

efficient viral replication *in vivo* and/or may be compensated by changes elsewhere in the

genome.

ORF3c is dispensable for efficient SARS-CoV-2 replication

To assess the importance of *ORF3c* for efficient viral replication, we generated SARS-CoV-2

Wuhan Hu-1 variants harboring inactivating mutations in this gene. Introducing premature stop

codons into *ORF3c* is not possible without simultaneously introducing non-synonymous

mutations in ORF3a and/or ORF3d. Using circular polymerase extension reaction (CPER)

(Torii et al., 2021), we therefore mutated the start codon of ORF3c to threonine (M1T), which

resulted in a silent mutation in the overlapping *ORF3a* gene (D22D) (Fig. 5A, left panel). After

rescue and validating successful introduction of the mutation, we infected CaCo-2 and CaLu-3

cells with an MOI of 0.1. Quantification of SARS-CoV-2 RNA copies in the culture

supernatants over a period of three days revealed that ORF3c M1T replicated as efficiently as

wild type SARS-CoV-2 (Fig. 5A, right panels). In a parallel experiment, we introduced a

premature stop codon (Q5*) in ORF3c, mimicking the mutation that can naturally be found in

the SARS-CoV-2 delta and kappa variants (Fig. 4F, 5B, left panel). Since the respective

nucleotide change introduces a S26L mutation in ORF3a, we simultaneously disrupted *ORF3a*

by introducing a premature stop codon (R6*) downstream of a methionine and potential

alternative start codon at position 5. While loss of ORF3a markedly reduced replicative fitness

in CaCo-2 cells, the virus still replicated efficiently in CaLu-3 cells (Fig. 5B, right panels). As

observed for the ORF3c M1T mutant, introduction of ORF3c Q5* did not significantly affect

the replicative fitness of the virus.

Since several (accessory) proteins of SARS-CoV-2 have been shown to suppress the induction

of IFN-β and/or IFN-β-mediated immune activation (Min et al., 2021; Xia et al., 2020), we

hypothesized that the immunosuppressive effect of ORF3c may be masked by other viral

factors. One potent suppressor of IRF3-mediated IFN-β induction is ORF6 (Kimura *et al.*, 2021;

Li et al., 2020). We therefore generated a BAC clone of SARS-CoV-2, in which *ORF6* was

replaced by a *YFP* reporter gene, and introduced the ORF3c M1T mutation described above

(Fig. 5C, left panel). This clone is based on the B.1 variant, and therefore additionally harbors

the ORF3c R36I mutation described above. Viral spread was monitored by live-cell imaging

and quantification of YFP-expressing (i.e. infected) cells. Again, disruption of *ORF3c* did not

result in impaired viral spread, and the ORF3c M1T virus replicated as efficiently as its ORF3c

wild type counterpart (Fig. 5C, right panels). Thus, while ORF3c and its immunosuppressive

activity are conserved among sarbecoviruses, the respective gene is dispensable for viral

replication in CaCo-2 and CaLu-3 cells.

DISCUSSION

Several lines of evidence suggested that the *ORF3c* gene of SARS-CoV-2 codes for a peptide that plays a role in viral replication: (1) the respective open reading frame is conserved in different sarbecoviruses and shows synonymous site conservation (Firth, 2020; Jungreis *et al.*, 2021b; Nelson et al., 2020) (Fig. 3), (2) upstream ATGs do not show a strong initiation context and may allow ORF3c translation via leaky scanning (Firth, 2020) (Fig. 1), (3) ribosomal profiling demonstrated that ORF3c is translated in SARS-CoV-2 infected cells (Finkel *et al.*, 2021), (4) ORF3c shows a high density of CD8+ T cell epitopes (Nelson *et al.*, 2020), (5) *in silico* analyses predict a conserved transmembrane domain in ORF3c and a potential role as viroporin (Cagliani *et al.*, 2020; Firth, 2020). Still, the exact function of ORF3c, and its contribution to efficient viral replication have remained unclear.

Here, we identify ORF3c as an immune evasion factor of SARS-CoV-2 and other sarbecoviruses that inhibits the induction of IFN-β upon activation of innate sensing cascades. Using luciferase reporter assays, we demonstrate that ORF3c suppresses activation of the *IFNB1* promoter by the pattern recognition receptors (PRRs) and RNA sensors RIG-I and MDA5. Importantly, ORF3c also significantly reduced the expression of endogenous IFN-β in response to Sendai virus infection, a known inducer of RIG-I sensing (Strahle *et al.*, 2006). Since ORF3c exerts its inhibitory effect independently of the receptor, it most likely targets a factor further downstream in the signaling cascade. In line with this, co-immunoprecipitation experiments revealed an interaction of ORF3c with the mitochondrial signaling adaptor MAVS. We found no evidence for an interaction with other components of the sensing cascade (i.e. RIG-I, MDA5, or TBK1). Notably, ORF3c failed to prevent *IFNB1* promoter activation if MAVS itself was used as an activator. This lack of inhibition is not merely the result of a saturation effect since MAVS induced the *IFNB1* promoter less efficiently than RIG-I (Fig. 2). MAVS is targeted by proteins from different viruses. For example, Influenza A Virus PB1-F2 inhibits innate sensing by binding to MAVS (Varga et al., 2011) and decreasing the

mitochondrial membrane potential (Varga et al., 2012). Another example is the NS3/4A serine

protease of Hepatitis C virus (HCV), which cleaves MAVS, thereby inhibiting downstream

immune activation (Anggakusuma et al., 2016). Similarly, SARS-CoV-2 ORF10 was recently

shown to induce the degradation of MAVS via mitophagy (Li et al., 2022). In contrast to HCV

NS3/4A and SARS-CoV-2 ORF10, however, we found no evidence for altered total protein

levels, proteolytic cleavage or other post-translational modifications of MAVS in the presence

of ORF3c. One possible mode of action may involve a competitive binding of ORF3c and RIG-

I/MDA5 to MAVS. In the presence of ORF3c, the CARD domain of MAVS may be blocked

and thus not be bound and activated by active RIG-I or MDA5. Notably, however, co-

immunoprecipitation experiments showed that the CARD domain of MAVS in dispensable for

an interaction of ORF3c with MAVS.

In line with a relevant role of ORF3c in viral replication, its immunosuppressive activity is

conserved in orthologs of other sarbecovirus species, including the SARS-CoV reference virus

Tor2. While all orthologs tested inhibited *IFNB1* promoter activation, those of SARS-CoV Tor2

and batCoV BANAL20-52 were less active than their counterparts from SARS-CoV-2 Wuhan-

Hu-1 and batCoV ZXC21. The reduced activity of BANAL-20-52 ORF3c can be ascribed to a

single amino acid change (L11P) distinguishing it from Wuhan-Hu-1 ORF3c (Fig. 3B). While

L11 is largely conserved in the SARS-CoV-2 cluster, most of the viruses in the SARS-CoV

cluster (including Tor2) harbor a glutamine at this position (Fig. 3B) (Cagliani *et al.*, 2020;

Firth, 2020). Thus, polymorphisms at position 11 can affect the inhibitory activity of ORF3c.

In addition to this, our alanine scanning approach revealed that Leu2/Leu3 and Ile6/Leu7 are

also contributing to the immunosuppressive effect of ORF3c. Most of these residues are

conserved among different sarbecoviruses. One notable exception is Ile6 (Fig. 3B). Viruses

from the SARS-CoV cluster harbor a Valine at this residue. These include SARS-CoV Tor2

ORF3c, which was less active than its SARS-CoV-2 counterpart.

ORF3c is not the only SARS-CoV-2 protein that interferes with RIG-I- and/or MDA-5-

mediated immune activation. As already mentioned above, ORF10 suppresses innate sensing

by targeting MAVS (Li *et al.*, 2022). Moreover, ORF3b, nucleocapsid, ORF6 and ORF8 have

all been shown to suppress IFN-β expression (Hayn et al., 2021; Kimura *et al.*, 2021; Konno *et*

al., 2020; Kopecky-Bromberg et al., 2007; Li *et al.*, 2020), highlighting the selection pressure

exerted by this pathway. The convergent evolution of viral proteins exerting overlapping

immune evasion activities may represent a backup mechanism that allows viral replication even

if one of the IFN-β suppressing proteins is lost. In line with this, SARS-CoV-2 variants

expressing a C-terminally truncated, inactive ORF6 protein have emerged several times during

the pandemic and spread via human-to-human transmission (Kimura *et al.*, 2021). Similarly,

natural SARS-CoV-2 variants lacking an intact ORF3c gene still efficiently spread in the human

population. In fact, more than 80% of the sequenced genomes of B.1.617.1, B.1.617.2 and

B.1.630 harbor premature stop codons at positions 5 and 15, respectively (Fig. 4). We

hypothesized that the loss of ORF3c in these viruses may be compensated by the activity of

ORF6. However, replication kinetics in CaCo-2 and CaLu-3 cells revealed that loss of ORF3c

does not affect viral replication in the absence of ORF6 either (Fig. 5). Intriguingly, IFN-β

mRNA was not detectable in cells infected with the SARS-CoV-2 double mutant lacking

ORF3c and ORF6 (data not shown). Thus, yet another viral inhibitor of IFN-β expression (e.g.

ORF8 or N) may be able to rescue efficient viral replication in this case.

Notably, the emergence of premature stop codons in small reading frames such as *ORF3c* may

also be tolerated or even be beneficial if they provide a fitness advantage by optimizing

overlapping reading frames. For example, the ORF3c Q5* mutation is accompanied by an S26L

change in ORF3a. However, experimental disruption of ORF3c without changing the amino

acid sequence of ORF3a (Fig. 5A) or upon deletion of ORF3a (Fig. 5B) did not affect viral

replication *in vitro* either.

One intriguing observation is the emergence of a nucleotide change in a subfraction (~3%) of B.1.617.2 viruses that reverts the stop codon at position 5 to a tyrosine (*5Y). Thus, it is tempting to speculate that the loss of ORF3c was initially just carried along with mutations elsewhere in the genome (e.g. in Spike) that conferred a major fitness advantage to the virus, before ORF3c expression was reverted by another point mutation.

In summary, our study identifies ORF3c as an immune evasion factor of SARS-CoV-2 and other sarbecoviruses. While an intact ORF3c gene is clearly dispensable for viral replication *in vitro* and *in vivo*, the conservation of this short open reading frame and the pseudo-reversion of premature stop codons suggests that it may still contribute to efficient viral replication *in vivo*. The emergence of future SARS-CoV-2 variants may help to fully decipher the role of this enigmatic ORF and its co-evolution with other viral genes.

ACKNOWLEDGMENTS

We thank Isabell Haußmann for excellent technical assistance and would like to thank the following persons for providing helpful reagents: Peter D Burbelo (LIPS assay plasmids), Konstantin MJ Sparrer (RIG-I, MDA5 and TBK1 expression plasmids, IRF luciferase reporter plasmid), Harmit Malik (MAVS expression plasmid), Rongtuan Lin (IRF3 expression plasmid), Bernd Baumann (NF-κB luciferase reporter plasmid, *Gaussia* luciferase control vector, IKKβ expression plasmid), Michael Gale Jr (IFN-β promoter reporter plasmid), Prof. Takasuke Fukuhara (CPER plasmids and cell lines) and Prof. Takashi Irie (pCAGGS Flag-IRF3 5d). SARS-CoV-2 mNeon Green was kindly provided by Prof. Michael Schindler. This work was funded by the Federal Ministry of Education and Research Germany (BMBF; grant ID: FKZ 01KI20135), the Canon Foundation Europe, the Heisenberg Program of the German Research Foundation (DFG; grant ID: SA 2676/3-1) and grants of the COVID-19 program of the Ministry of Science, Research and the Arts Baden-Württemberg (MWK; grants IDs: MWK K.N.K.C.014 (U21) and MWK K.N.K.C.015 (U22)) to D.S. AE and AH were supported by BMBF

320 SenseCoV2 01KI20172A; DFG Fokus COVID-19, EN 423/7-1; and Coronavirus research

321 grants by the Bavarian State Ministry of Science and the Arts and Bavarian State Ministry of

322 Health, Bay-VOC.

AUTHOR CONTRIBUTIONS

M.M. and D.S. conceptualized the study and designed the experiments. M.M. performed most

of the experiments. A.H. and A.E. generated and provided BAC clones for SARS-CoV-2

ΔORF6 YFP (ORF3c wild type and ΔORF3). S.F. and D.S. generated SARS-CoV-2 mutants

via CPER that were characterized by D.S., S.F. and K.U., C.K. and J.E.K. performed some of

the luciferase reporter assays. M.S. sequenced the YFP reporter viruses. A.S. and J.I. performed

in silico analyses to identify premature stop codons in ORF3c. M.M. and D.S. prepared the

figures, and D.S. wrote the initial draft of the manuscript. All authors reviewed and edited the

manuscript. A.E., K.S. and D.S. supervised the experiments, provided resources and acquired

funding.

DECLARATION OF INTERESTS

336 The authors declare no competing interests.

MATERIAL AND METHODS

<u>Cell lines</u>

HEK293T were maintained in Dulbecco's modified Eagle medium (DMEM) supplemented

with 10% heat-inactivated fetal calf serum (FCS), L-glutamine (2 mM), streptomycin (100

mg/ml) and penicillin (100 U/ml) and were cultured at 37°C, 90% humidity and 5% CO_2. They

were isolated from a female fetus. HEK293T cells were transfected using a standard calcium

phosphate method. HEK293-C34 cells, IFNAR1 KO HEK293 cells expressing human ACE2

and TMPRSS2 by doxycycline treatment (Torii *et al.*, 2021), were maintained in DMEM (high

glucose) containing 10% FBS, 10 mg/ml blasticidin (InvivoGen) and 1% PS. Caco-2 (human

colorectal adenocarcinoma, male) cells were grown in Dulbecco's modified Eagle's medium

(DMEM) supplemented with 10% heat-inactivated FCS, 100 units/ml penicillin, 100 µg/ml

streptomycin, 2 mM L-glutamine, and 1x non-essential amino acids (NEAA). Calu-3 cells (lung

adenocarcinoma, male) were cultured in Eagle's minimum essential medium (EMEM)

supplemented with 20% heat-inactivated FCS, 100 units/ml penicillin, and 100 µg/ml

streptomycin. Medium was changed every day. Vero E6 (*Cercopithecus aethiops* derived

epithelial kidney, female) cells were grown in Dulbecco's modified Eagle's medium (DMEM)

supplemented with 2.5% heat-inactivated FCS, 100 units/ml penicillin, 100 µg/ml

streptomycin, 2 mM L-glutamine, 1 mM sodium pyruvate, and 1x non-essential amino acids.

Derivatives thereof stably expressing TMPRSS2 were cultured in (DMEM) supplemented with

5% heat-inactivated FCS, 100 units/ml penicillin, 100 µg/ml streptomycin, and 100 µg/ml

G418.

<u>Virus strains and virus propagation</u>

SARS-CoV-2 ΔORF6-YFP and SARS-CoV-2 ΔORF6-YFP ΔORF3c were propagated by

inoculation of Vero E6/TMPRSS2 cells in 75 cm² cell culture flasks in medium containing 2%

FCS. Cells were incubated at 37°C and supernatants were harvested 2 to 4 days post inoculation.

Supernatants were centrifuged for 5 min at 1,000 × g to remove cellular debris, and then

aliquoted and stored at -80°C as virus stocks. Infectious titer was determined in Vero

E6/TMPRSS2 cells as Tissue Culture Infection Dose 50 ($TCID_{50}$)/ml (see Method Details).

In silico prediction of translation initiation sites

Translation initiation sites (TIS) and Kozak context were determined using TIS predictor

(https://www.tispredictor.com/) (Gleason *et al.*, 2022).

In silico prediction of ORF3c secondary structure and transmembrane domains

To predict the secondary structure of ORF3c and its R36I variant PEP-FOLD 3 (Camproux et al., 2004) was used.

The effect of point mutations on a transmembrane domain in ORF3c was predicted using the 'Prediction of transmembrane helices in proteins' tool TMHMM2.0 (Sonnhammer *et al.*, 1998).

Generation of expression plasmids

ORF3c genes were PCR-amplified using viral cDNA as a template and subsequently inserted into a pCG expression vector co-expressing GFP via an IRES using unique XbaI and MluI restriction sites. To facilitate protein detection, a C-terminal HA-tag (TACCCATACGATGTTCCAGATTACGCT) was added by extension PCR. To generate ORF3c-HA alanine mutants as well as ORF3c-R36I, -BANAL-20-52, -Tor2 and SL-CoVZXC21, point mutations were introduced by site-directed mutagenesis using the wild type Wuhan-Hu-1 ORF3c-HA expression plasmid as template. pRen2-ORF3c was generated by conventional cloning, using the unique EcoRI and XhoI restriction sites in pRen2. IRF3 5D was PCR-amplified using pCAGGS Flag-IRF3 5D as a template and subsequently inserted into pEGFP-C1 via XhoI and EcoRI. All constructs were sequenced to verify their integrity. PCR primers are listed in the Key Resources Table.

Generation and recovery of a recombinant SARS-CoV-2 ORF3 mutant

Stop mutations within ORF3c were introduced into the bacmid pBSCoV2_d6-YFP harboring the SARS-CoV-2 backbone (Herrmann et al., 2021) using 2-step Red Recombination (Tischer et al., 2006). For this purpose, the KanS cassette was amplified from pEP-KanS with the following oligonucleotides:

(1) mut3c-fwd: caattggaactgtaactttgaagcaaggtgaaatcaaggaCgctactccttcagattttgAGGATGACG

ACGATAAGTAGGG

(2) mut3c-rev: gtatcgttgcagtagcgcgaacaaaatctgaaggagtagcGtccttgatttcaccttgctCAACCAATT

AACCAATTCT GATTAG

Integrity of the obtained bacterial artificial chromosomes (BAC) and presence of desired stop

mutations were confirmed by restriction digestion and next generation sequencing.

Recombinant SARS-CoV-2 viruses expressing EYFP instead of the viral ORF6 protein and

containing mutations within ORF3c were recovered by transfection of the BACs into HEK293T

cells overexpressing viral N protein, ACE2 receptor, and T7 RNA polymerase as described

previously (Herrmann *et al.*, 2021). The obtained reporter viruses were further passaged on

CaCo-2 cells and viral titers were determined by endpoint titration (see $TCID_{50}$).

<u>Generation of recombinant SARS-CoV-2 mutants by circular polymerase extension reaction</u>

To generate recombinant SARS-CoV-2 by circular polymerase extension reaction (CPER)

(Torii *et al.*, 2021) nine DNA fragments comprising parts of SARS-CoV-2 (WK-521, PANGO

lineage A; GISAID ID: EPI ISL 408667) (Matsuyama et al., 2020) were generated by PCR

using PrimeSTAR GXL DNA polymerase. A linker fragment comprising hepatitis delta virus

ribozyme, the bovine growth hormone poly A signal and the cytomegalovirus promoter was

also prepared by PCR. The ten obtained DNA fragments were mixed and used for

CPER. ORF3c mutations were inserted in fragment 9/10 by site-directed overlap extension

PCR with the primers listed in the key resources table.

To produce chimeric recombinant SARS-CoV-2, Tetracycline-inducible ACE2 and TMPRSS-

expressing IFNAR1-deficient HEK293 (HEK293-C34) cells were transfected with the CPER

products using TransIT-LT1 according to the manufacturer's protocol. One day post

transfection, the culture medium was replaced with Dulbecco's modified Eagle's medium (high

glucose) containing 2% FCS, 1% PS and doxycycline. At 7 d post transfection, the culture

medium was harvested and centrifuged, and the supernatants were collected as the seed virus.

To remove the CPER products (i.e., any SARS-CoV-2 DNA), 1 ml of the seed virus was treated

with 2 µl TURBO DNase (Thermo Fisher Scientific, Cat# AM2238) and incubated at 37°C for

1 h. Complete removal of the CPER products (i.e., SARS-CoV-2-related DNA) from the seed

virus was verified by PCR.

To prepare virus stocks for infection, VeroE6/TMPRSS2 cells (5,000,000 cells in a T-75 flask)

were infected with 20-50 µl of the seed virus. One-hour post infection, the culture medium was

replaced with DMEM (low glucose) containing 2% FBS and 1% PS. Two to four days post

infection, the culture medium was harvested and centrifuged, and the supernatants were

collected. Viral titers were determined by $TCID_{50}$. To verify the sequence of chimeric

recombinant SARS-CoV-2, viral RNA was extracted from the virus stocks using the QIAamp

viral RNA mini kit and viral genomes were sequenced as described before (Kimura et al., 2022).

<u>Tissue culture infectious dose ($TCID_{50}$)</u>

Viral titers were determined as the 50% tissue culture infectious dose. Briefly, one day before

infection, VeroE6/TMPRSS2 cells (10,000 cells) were seeded into 96-well plates. Cells were

inoculated with serially diluted virus stocks and incubated at 37°C. Four days later, cells were

checked microscopically for cytopathic effects (CPE), and $TCID_{50}$/ml was calculated using the

Reed–Muench method.

<u>SARS-CoV-2 replication kinetics in Vero E6, Caco-2 and Calu-3 cells</u>

One day before infection with CPER-derived SARS-CoV-2 clones, Caco-2 cells (10,000

cells/well) or CaLu-3 cells (20,000 cells/well) were seeded into a 96-well plate. Cells were

infected with SARS-CoV-2 at an MOI of 0.1 and incubated at 37°C. One hour later, the infected

cells were washed and 180 µl of culture medium was added. The culture supernatants and cells

were harvested at the indicated timepoints and used for RT–qPCR to quantify the viral RNA

copy number. For replication kinetics of BAC-derived SARS-CoV-2 ΔORF6-YFP and SARS-

CoV-2 ΔORF6-YFP ΔORF3c, Caco-2 cells (10,000/well) were seeded into a 96-well plate one

day prior to infection. Cells were infected in triplicates at an MOI of 0.1 & 0.01 for 1 hour at

37°C. After washing and addition of 100 µl fresh culture medium the plates were placed in an

Incucyte plate reader and images were taken at the indicated time points for up to 96 hours.

The 'Basic Analysis Mode' was applied to quantify virus growth as green area normalized to

phase area. Supernatants and cells were harvested at the indicated time points to determine

cytokine levels by Cytokine Array and RT-qPCR respectively.

<u>RT–qPCR</u>

5 µl culture supernatant was mixed with 5 µL of 2 x RNA lysis buffer [2% Triton X-100,

50 mM KCl, 100 mM Tris-HCl (pH 7.4), 40% glycerol, 0.8 U/µL recombinant RNase inhibitor

and incubated at room temperature for 10 minutes. RNase-free water (90 µL) was added, and

the diluted sample (2.5 µl) was used as the template for real-time RT-PCR performed according

to the manufacturer's protocol using the OneStep TB Green PrimeScript PLUS RT-PCR kit

and the following primers: Forward *N*, 5′-AGC CTC TTC TCG TTC CTC ATC AC-3′; and

Reverse *N*, 5′-CCG CCA TTG CCA GCC ATT C-3′. The viral RNA copy number was

standardized using a home-made standard.

IFNB1 and *IL6* RNA levels were determined in cell lysates collected from (1) SARS-CoV-2

ΔORF6-YFP and SARS-CoV-2 ΔORF6-YFP ΔORF3c infected CaCo-2 wild-type cells and (2)

transfected HEK293T cells infected with Sendai virus for 8 hours. Total RNA was isolated

using the Viral RNA Mini Kit (Qiagen) according to the manufacturer's instructions. Genomic

DNA was removed using the DNA-free kit (Thermo Fischer Scientific) and subsequent cDNA

synthesis was performed using the PrimeScript RT reagent Kit (TAKARA), both according to

the manufacturer's instructions. qPCR was performed using the Luna Universal Probe qPCR

Master Mix (NEB) together with primer probes for IFN-β, IL-6 and GAPDH (Thermo Fischer

Scentific). All reactions were run in duplicates and RNA levels were internally normalized to

GAPDH.

<u>Transfection of HEK293T cells</u>

For overexpression experiments, HEK293T cells were transfected using standard calcium

phosphate transfection protocols. 6x 10^5 cells were seeded in 6-well plates on the day before

and medium was changed 6 hours after transfection.

<u>Co-immunoprecipitation</u>

To investigate possible interactions between ORF3c and proteins of the Interferon signaling

pathway, co-immunoprecipitation with subsequent analysis by western blotting was performed.

Briefly, HEK293T cells were seeded in 6-well plates and co-transfected with expression

plasmids for HA-tagged ORF3c and Flag-tagged RIG-I, MDA5, MAVS or TBK1 (ratio 4:1;

5µg/well). One day post transfection, cells were lysed in 300 µl western blot lysis buffer and

cleared by centrifugation (see "Western blotting"). 45 µl of the lysate was used for whole-cell

lysate analysis and further prepared as described in "Western blotting", while 255 µl of the

lysate was used for co-immunoprecipitation. A pre-clearing step was performed to remove

unspecifically binding compounds from the lysate. Pierce Protein A/G Magnetic beads (Thermo

Fisher) were washed three times with 1 ml NP40 wash buffer (50 mM HEPES, 300 mM NaCl,

0.5% NP40, pH 7.4) and added to the lysate. After incubation for 1 h at 4°C, beads were

removed from the lysate using a magnetic rack. To precipitate protein complexes, the lysate

was incubated first with an anti-Flag antibody (Sigma, 1.5 µg/sample) for 1 hour followed by

addition of 15 µl washed Protein A/G Magnetic Beads for one additional hour at 4°C. After

incubation, the beads were washed three times in NP40 wash buffer before incubation with

80 µl 1 x Protein Sample Loading Buffer at 95°C for 10 minutes to recover bound proteins.

After addition of 1.75 ml β-mercaptoethanol, whole-cell lysates and precipitates were analyzed

by western blotting.

<u>Western blotting</u>

To determine expression of cellular and viral proteins, cells were washed in PBS, lysed in

western blot lysis buffer (150 mM NaCl, 50 mM HEPES, 5 mM EDTA, 0.1% NP40, 500 mM

Na3VO4, 500 mM NaF, pH 7.5) and cleared by centrifugation at 20,800 x g for 20 min at 4°C.

Lysates were mixed with Protein Sample Loading Buffer (LI-COR) supplemented with 10% β-

mercaptoethanol and heated at 95°C for 5 min. Proteins were separated on NuPAGE 4%–12%

Bis-Tris Gels (Thermo Fischer Scientific), blotted onto Immobilon-FL PVDF membranes and

stained using primary antibodies directed against HA-tag, Flag-tag, GAPDH, RIG-I, MDA5,

MAVS, TBK1, IRF3 and Infrared Dye labeled secondary antibodies (LI-COR IRDye). Proteins

were detected using a LI-COR Odyssey scanner and band intensities were quantified using LI-

COR Image Studio Lite Version 5.2.

<u>Immunofluorescence microscopy</u>

Confocal immunofluorescence microscopy was used to determine the subcellular localization

of ORF3c and ORF3c R36I. Briefly, 150,000 HEK293T cells were seeded on 13 mm diameter

glass coverslips coated with poly-L-lysine (Sigma-Aldrich) in 24-well plates. On the following

day, cells were transfected with an expression plasmid for ORF3c, ORF3c R36I or an empty

vector control (500 ng) using Lipofectamin2000 (Invitrogen). One day post transfection, cells

were fixed in 4% PFA for 20 min at RT, permeabilized in PBS 0.5% Triton X-100 for 15 min

at RT and blocked in 5% BSA/PBS supplemented with 0,1% Triton X-100 for 15 min at RT.

ORF3c was stained using a primary antibody against the HA-tag and secondary goat anti-mouse

AF555. Nuclei were stained in parallel using 4′,6-Diamidine-2′-phenylindole (DAPI; Thermo

Fischer Scientific). Coverslips were mounted on glass slides using Mowiol (CarlRoth)

mounting medium and confocal microscopy was performed using an LSM710 (Carl Zeiss).

<u>Firefly luciferase assay</u>

HEK293T cells were seeded in 96 well plates at $3x10^4$ cells /well. After 24 h, cells were

transfected with a mix of expression vectors containing firefly luciferase reporter constructs for

the IFN-β promoter or a mutant thereof lacking NF-κB biding sites (IFN-β ΔNF-κB) (reporter,

10 ng), a *Gaussia* luciferase expression plasmid (normalization control, 5 ng), expression

plasmids for RIG-I-CARD, MDA5, MAVS or IRF3 5D (stimulus, 5 ng), different amounts of

ORF3 expression constructs (12.5 - 100 ng) and empty vector (pCG_HIV-1 M NL4-3 *nef* stop

ΔIRES-eGFP) to adjust total DNA amounts across all conditions to 200 ng/well. After 24 h,

supernatants were harvested, and cells were lysed in 100µl 1x Passive Lysis Buffer (Promega).

Gaussia luciferase activity in the supernatants was measured by addition of Coelenterazine

(PJK Biotech). Firefly luciferase activity was measured in the cells using the Luciferase Assay

System (Promega) according to the manufacturers instruction.

<u>LIPS assay</u>

Luciferase fusion protein (10^6 RLU) in 50 µl Buffer A (50 mM Tris, 150 mM NaCl, 0.1% Triton

X-100, pH 7.5) and 1 µl sample serum in 49 µl Buffer A was added to 1.5 ml tubes and

incubated with shaking at 300 rpm for 1 h at room temperature. Pierce Protein A/G Magnetic

beads were added to each condition as a 30% suspension in PBS for an additional hour and

shaking at room temperature. Samples were placed on a magnetic rack, and supernatant was

removed after 1-minute incubation. Magnetic beads were washed twice with 150 µl Buffer A

followed by 2 washes with 150 µl PBS. Samples were transferred into a 96-well opaque nunc-

plate (VWR), and 50 µl Coelenterazine (PJK Biotech) was added to each condition. Samples

were measured immediately on a TriStar² S LB 942 Multimode Reader (Berthold Technologies)

with an integration time of 0.1 seconds and a read height of 1 mm.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses were performed using GraphPad PRISM 9.4.1. For statistical testing

between two means P values were calculated using paired or unpaired Student's t test. For

comparison within one group, we used one-way analysis of variation (ANOVA) with Dunnett's

multiple comparison test and for comparison between two or more groups we used two-way

ANOVA with Sidak's multiple comparison test. Unless otherwise stated, data are shown as the

mean of at least three independent experiments ± SD. Significant differences are indicated as:

* p ≤ 0.05; ** p ≤ 0.01; *** p ≤ % 0.001 and **** p ≤ 0.0001. Statistical parameters are

specified in the figure legends.

FIGURE LEGENDS

Figure 1: Open reading frames in the *ORF3* gene locus and inhibition of IFN-β induction

by SARS-CoV-2 ORF3c. (A) Genome organization of SARS-CoV-2 is illustrated on top;

overlapping ORFs in the ORF3 locus are shown at the bottom. Vertical grey lines indicate

internal ATG codons. Experimentally confirmed translation initiation sites (Finkel *et al.*, 2021)

are highlighted by grey triangles. **(B)** Kozak sequences of the *ORF3c* initiation codon and

upstream ATG codons in *ORF3a* are shown. ATG context was determined using TIS predictor

(https://www.tispredictor.com/) (Gleason *et al.*, 2022). **(C)** Western blot analysis of HEK293T

cells transfected with two different concentrations of expression plasmids for the indicated

ORF3 proteins and peptides. ORF3a to ORF3e were detected via a C-terminal HA tag. GAPDH

served as loading control. **(D)** HEK293T cells were co-transfected with the indicated ORF3

expression plasmids, a reporter plasmid expressing firefly luciferase under the control of the

IFNB1 promoter and a construct expressing *Gaussia* luciferase under the control of a minimal

promoter. To induce immune signaling, half of the samples were additionally co-transfected

with an expression plasmid for the CARD domain of RIG-I. One day post transfection, firefly

luciferase activity was determined and normalized to *Gaussia* luciferase activity. Mean values

of three independent experiments measured in triplicates (±SD) are shown. **(E)** HEK293T cells

were transfected with an expression plasmid for SARS-CoV-2 ORF3c or an empty vector

control. 24 hours post transfection, cells were infected with increasing amounts of Sendai virus

(SeV) for an additional 8 hours. Cells were lysed to perform either RNA extraction and

subsequent qPCR for IFN-β (left panel) or western blot analysis (right panel). Mean values of

three independent experiments measured in duplicates (±SD) are shown. Multiple comparison

within individual reporter assays (D) were determined by one-way ANOVA with Dunnett's

test; * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq \% 0.001$ and **** $p \leq 0.0001$. Multiple comparison

between groups (E) were determined by two-way ANOVA with Sidak's multiple comparison

test; * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq \% 0.001$ and **** $p \leq 0.0001$.

Figure 2. ORF3c interacts with MAVS and inhibits IFN-β induction independently of the

pattern recognition receptor. (A) Cartoon illustrating IRF3- and NF-κB-mediated activation

of the *IFNB1* promoter upon RIG-I- or MDA5-mediated sensing. **(B)** HEK293T cells were co-

transfected with increasing amounts of an expression plasmid for SARS-CoV-2 ORF3c, a

construct expressing *Gaussia* luciferase under the control of a minimal promoter and a reporter

plasmid expressing firefly luciferase under the control of the *IFNB1* promoter (left panel) or a

mutant thereof lacking the NF-κB binding site (right panel). Immune signaling was induced by

co-transfecting an expression plasmid for the CARD domain of RIG-I. One day post

transfection, firefly luciferase activity was determined and normalized to *Gaussia* luciferase

activity. Mean values of three independent experiments measured in triplicates (±SD) are

shown. **(C)** HEK293T cells were transfected and analyzed essentially as described in (B).

Instead of RIG-I CARD, however, immune signaling was introduced by co-transfecting

expression plasmids for MDA5 (left panel), MAVS (central panel) or a constitutively active

mutant of IRF3 (right panel). Mean values of three independent experiments measured in

triplicates (±SD) are shown. **(D, E)** HEK293T cells were co-transfected with expression

plasmids for (D) Flag-tagged RIG-I, MDA5, MAVS, TBK1, (E) MAVS or a mutant thereof

lacking its CARD domain (MAVSΔCARD) and an expression plasmid for HA-tagged SARS-

CoV-2 ORF3c. One day post transfection, cells were lysed. Cell lysates were analyzed by

Western blotting, either directly ("input") or upon pull-down using a Flag-specific antibody

("IP"). **(F)** HEK293T cells were co-transfected with expression plasmids as described in (B,

left panel). Immune signaling was induced by co-transfecting an expression plasmid for the

CARD domain of RIG-I as well as the indicated ORF3c alanine mutants. Multiple comparison

within individual reporter assays (B, C, F) were determined by one-way ANOVA with

Dunnett's test; * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq \% 0.001$ and **** $p \leq 0.0001$.

Figure 3. Conservation of ORF3c and its immunosuppressive activity in sarbecoviruses.

(A) Simplified cartoon illustrating the *ORF3* locus of randomly selected members of the

Sarbeco-, *Hibeco*- and *Nobecovirus* genera. Open reading frames with a length of at least 30

nucleotides are indicated as rectangles. ORF3c is highlighted in dark red. **(B)** Alignment of

ORF3c amino acid sequences of the indicated viral isolates. Members of the SARS-CoV-2

cluster are shown on top, members of the SARS-CoV cluster at the bottom. For the underlined

ORF3c sequences, expression plasmids were generated and analyzed for their ability to inhibit

IFNB1 promoter activation in **(C)**. Briefly, HEK293T cells were co-transfected with increasing

amounts of the indicated ORF3c expression plasmids, a reporter plasmid expressing firefly

luciferase under the control of the *IFNB1* promoter and a construct expressing *Gaussia*

luciferase under the control of a minimal promoter. An expression plasmid for Influenza A virus

non-structural protein 1 (NS1) served as positive control. Immune signaling was induced by co-

transfecting an expression plasmid for the CARD domain of RIG-I. One day post transfection,

firefly luciferase activity was determined and normalized to *Gaussia* luciferase activity. Mean

values of three independent experiments measured in triplicates (±SD) are shown. Multiple

comparison between groups (C) were determined by two-way ANOVA with Sidak's multiple

comparison test; ns not significant, * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq \% 0.001$ and **** $p \leq$

0.0001.

Figure 4. Characterization of naturally occurring variants of SARS-CoV-2 ORF3c. (A)

Cartoon illustrating non-synonymous changes in ORF3a, c and d as a result of the naturally

occurring polymorphisms G25563T. **(B)** Secondary structure of Wuhan Hu-1 ORF3c (red) and

the respective R36I variant thereof (blue) as predicted using PEP-FOLD 3 (Camproux et al.,

2004). **(C)** The presence of transmembrane domains in Wuhan Hu-1 ORF3c (left panel) and

the respective R36I variant thereof (right panel) was predicted using TMHMM - 2.0 (Krogh et

al., 2001). **(D)** HEK293T cells were transfected with expression plasmids for Wuhan Hu-1

ORF3c or ORF3c R36I. One day post transfection, cells were stained for ORF3c (anti-HA,

green) and nuclei (DAPI, blue) (scale bar = 20 µm). **(E)** HEK293T cells were co-transfected

with increasing amounts of the indicated ORF3c expression plasmids, a reporter plasmid

expressing firefly luciferase under the control of the *IFNB1* promoter and a construct expressing

Gaussia luciferase under the control of a minimal promoter. Immune signaling was induced by

co-transfecting an expression plasmid for the CARD domain of RIG-I. One day post

transfection, firefly luciferase activity was determined and normalized to *Gaussia* luciferase

activity. Mean values of three independent experiments measured in triplicates (±SD) are

shown. **(F)** Mutations introducing premature stop codons in ORF3c that can be found in at least

20% of the samples of at least one PANGO (sub)lineage. **(G)** Frequency of the mutations shown

in (F) in the PANGO (sub)lineages B.1.617.1 (delta), B.1.617.2 (kappa), B.1.617 and B.1.630.

Multiple comparison between groups (E) were determined by two-way ANOVA with Sidak's

multiple comparison test; ns, not significant.

Figure 5. Disruption of ORF3c does not affect SARS-CoV-2 replication in CaCo-2 or CaLu-3 cells. (A) Circular polymerase extension reaction (CPER) was used to disrupt the start codon of ORF3c (M1T) in SARS-CoV-2 without affecting the amino acid sequence of ORF3a (left panel). CaCo-2 and CaLu-3 cells were infected with ORF3c wild type (red) or ORF3c-mutated (blue) SARS-CoV-2 at an MOI of 0.1. Viral replication was monitored over 72 hours by determining viral RNA copies in the culture supernatants (right panels). Mean values of four independent experiments (±SD) are shown. **(B)** CPER was used to introduce a premature stop codon in ORF3c (Q5stop). To avoid any bias by simultaneously changing the protein sequence of ORF3a (S26L), a premature stop codon was also inserted in ORF3a (R6*) (left panel). Viral replication (right panels) was monitored in CaCo-2 and CaLu-3 cells as described in (A). Mean values of four independent experiments (±SD) are shown. **(C)** A SARS-CoV-2 BAC clone harboring a disrupted ORF3c (M1T) and expressing YFP instead of ORF6 was generated (left panel). CaCo-2 cells were infected with SARS-CoV-2 ΔORF6-YFP (red) or SARS-CoV-2 ΔORF6-YFP ΔORF3c (blue) at an MOI of 0.1 (middle panel) or 0.01 (right panel). Cells were placed in a live cell imaging device, and the area of YFP positive cells over the total area of cells was quantified every 4 hours for 96 hours. Images below graphs show continuous virus spread (green) over the indicated time points from one randomly chosen well. Graphs show mean values of 3 independent experiments performed in triplicates (±SD). Multiple comparison between groups (A-C, right panel) were determined by two-way ANOVA with Sidak's multiple comparison test; * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq \% 0.001$ and **** $p \leq 0.0001$.

SUPPLEMENTAL INFORMATION

Figure S1: Detection of SARS-CoV-2-specific antibodies via Luciferase Immunoprecipitation System (LIPS). (A) Principle of the LIPS assay: HEK293T cells are transfected with expression plasmids for a viral protein of interest fused to *Renilla* luciferase.

Subsequently, transfected cells are lysed and incubated with serum samples and magnetic

beads. Antibodies against viral proteins of interest will cross-link the luciferase-containing

proteins with beads and allow magnet-assisted pull-down of both beads and luciferase activity.

(B) LIPS-mediated quantification of antibodies against SARS-CoV-2 N (left panel) and ORF3c

(right panel) in sera from SARS-CoV-2 naïve and convalescent sera (RLU, relative light units).

Each dot represents one independent serum sample. Differences in antibody levels between

SARS-CoV-2 naïve and convalescent sera was determined by unpaired student's t-test with

Welch's correction; * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$.

REFERENCES

Anggakusuma, Brown, R.J.P., Banda, D.H., Todt, D., Vieyres, G., Steinmann, E., and
Pietschmann, T. (2016). Hepacivirus NS3/4A Proteases Interfere with MAVS Signaling in both
Their Cognate Animal Hosts and Humans: Implications for Zoonotic Transmission. J Virol *90*,
10670-10681. 10.1128/JVI.01634-16.
Cagliani, R., Forni, D., Clerici, M., and Sironi, M. (2020). Coding potential and sequence
conservation of SARS-CoV-2 and related animal viruses. Infect Genet Evol *83*, 104353.
10.1016/j.meegid.2020.104353.
Camproux, A.C., Gautier, R., and Tuffery, P. (2004). A hidden markov model derived structural
alphabet for proteins. J Mol Biol *339*, 591-605. 10.1016/j.jmb.2004.04.005.
Finkel, Y., Mizrahi, O., Nachshon, A., Weingarten-Gabbay, S., Morgenstern, D., Yahalom-
Ronen, Y., Tamir, H., Achdout, H., Stein, D., Israeli, O., et al. (2021). The coding capacity of
SARS-CoV-2. Nature *589*, 125-130. 10.1038/s41586-020-2739-1.
Firth, A.E. (2020). A putative new SARS-CoV protein, 3c, encoded in an ORF overlapping
ORF3a. J Gen Virol *101*, 1085-1089. 10.1099/jgv.0.001469.
Firth, A.E., and Brierley, I. (2012). Non-canonical translation in RNA viruses. J Gen Virol *93*,
1385-1409. 10.1099/vir.0.042499-0.
Gleason, A.C., Ghadge, G., Chen, J., Sonobe, Y., and Roos, R.P. (2022). Machine learning
predicts translation initiation sites in neurologic diseases with nucleotide repeat expansions.
PLoS One *17*, e0256411. 10.1371/journal.pone.0256411.
Hachim, A., Gu, H., Kavian, O., Mori, M., Kwan, M.Y.W., Chan, W.H., Yau, Y.S., Chiu, S.S.,
Tsang, O.T.Y., Hui, D.S.C., et al. (2022). SARS-CoV-2 accessory proteins reveal distinct
serological signatures in children. Nat Commun *13*, 2951. 10.1038/s41467-022-30699-5.
Hayn, M., Hirschenberger, M., Koepke, L., Nchioua, R., Straub, J.H., Klute, S., Hunszinger,
V., Zech, F., Prelli Bozzo, C., Aftab, W., et al. (2021). Systematic functional analysis of SARS-
CoV-2 proteins uncovers viral innate immune antagonists and remaining vulnerabilities. Cell
Rep *35*, 109126. 10.1016/j.celrep.2021.109126.
Herrmann, A., Jungnickl, D., Cordsmeier, A., Peter, A.S., Uberla, K., and Ensser, A. (2021).
Cloning of a Passage-Free SARS-CoV-2 Genome and Mutagenesis Using Red Recombination.
Int J Mol Sci *22*. 10.3390/ijms221910188.
Jungreis, I., Nelson, C.W., Ardern, Z., Finkel, Y., Krogan, N.J., Sato, K., Ziebuhr, J., Stern-
Ginossar, N., Pavesi, A., Firth, A.E., et al. (2021a). Conflicting and ambiguous names of

overlapping ORFs in the SARS-CoV-2 genome: A homology-based resolution. Virology *558*,
145-151. 10.1016/j.virol.2021.02.013.

Jungreis, I., Sealfon, R., and Kellis, M. (2021b). SARS-CoV-2 gene content and COVID-19
mutation impact by comparing 44 Sarbecovirus genomes. Nat Commun *12*, 2642.
10.1038/s41467-021-22905-7.

Kimura, I., Konno, Y., Uriu, K., Hopfensperger, K., Sauter, D., Nakagawa, S., and Sato, K.
(2021). Sarbecovirus ORF6 proteins hamper induction of interferon signaling. Cell Rep *34*,
108916. 10.1016/j.celrep.2021.108916.

Kimura, I., Yamasoba, D., Nasser, H., Zahradnik, J., Kosugi, Y., Wu, J., Nagata, K., Uriu, K.,
Tanaka, Y.L., Ito, J., et al. (2022). The SARS-CoV-2 spike S375F mutation characterizes the
Omicron BA.1 variant. iScience *25*, 105720. 10.1016/j.isci.2022.105720.

Konno, Y., Kimura, I., Uriu, K., Fukushi, M., Irie, T., Koyanagi, Y., Sauter, D., Gifford, R.J.,
Consortium, U.-C., Nakagawa, S., and Sato, K. (2020). SARS-CoV-2 ORF3b Is a Potent
Interferon Antagonist Whose Activity Is Increased by a Naturally Occurring Elongation
Variant. Cell Rep *32*, 108185. 10.1016/j.celrep.2020.108185.

Kopecky-Bromberg, S.A., Martinez-Sobrido, L., Frieman, M., Baric, R.A., and Palese, P.
(2007). Severe acute respiratory syndrome coronavirus open reading frame (ORF) 3b, ORF 6,
and nucleocapsid proteins function as interferon antagonists. J Virol *81*, 548-557.
10.1128/JVI.01782-06.

Krogh, A., Larsson, B., von Heijne, G., and Sonnhammer, E.L. (2001). Predicting
transmembrane protein topology with a hidden Markov model: application to complete
genomes. J Mol Biol *305*, 567-580. 10.1006/jmbi.2000.4315.

Li, J.Y., Liao, C.H., Wang, Q., Tan, Y.J., Luo, R., Qiu, Y., and Ge, X.Y. (2020). The ORF6,
ORF8 and nucleocapsid proteins of SARS-CoV-2 inhibit type I interferon signaling pathway.
Virus Res *286*, 198074. 10.1016/j.virusres.2020.198074.

Li, X., Hou, P., Ma, W., Wang, X., Wang, H., Yu, Z., Chang, H., Wang, T., Jin, S., Wang, X.,
et al. (2022). SARS-CoV-2 ORF10 suppresses the antiviral innate immune response by
degrading MAVS through mitophagy. Cell Mol Immunol *19*, 67-78. 10.1038/s41423-021-
00807-4.

Matsuyama, S., Nao, N., Shirato, K., Kawase, M., Saito, S., Takayama, I., Nagata, N., Sekizuka,
T., Katoh, H., Kato, F., et al. (2020). Enhanced isolation of SARS-CoV-2 by TMPRSS2-
expressing cells. Proc Natl Acad Sci U S A *117*, 7001-7003. 10.1073/pnas.2002589117.

Min, Y.Q., Huang, M., Sun, X., Deng, F., Wang, H., and Ning, Y.J. (2021). Immune evasion
of SARS-CoV-2 from interferon antiviral system. Comput Struct Biotechnol J *19*, 4217-4225.
10.1016/j.csbj.2021.07.023.

Miorin, L., Kehrer, T., Sanchez-Aparicio, M.T., Zhang, K., Cohen, P., Patel, R.S., Cupic, A.,
Makio, T., Mei, M., Moreno, E., et al. (2020). SARS-CoV-2 Orf6 hijacks Nup98 to block STAT
nuclear import and antagonize interferon signaling. Proc Natl Acad Sci U S A *117*, 28344-
28354. 10.1073/pnas.2016650117.

Nelson, C.W., Ardern, Z., Goldberg, T.L., Meng, C., Kuo, C.H., Ludwig, C., Kolokotronis,
S.O., and Wei, X. (2020). Dynamically evolving novel overlapping gene as a factor in the
SARS-CoV-2 pandemic. Elife *9*. 10.7554/eLife.59633.

Oulas, A., Zanti, M., Tomazou, M., Zachariou, M., Minadakis, G., Bourdakou, M.M., Pavlidis,
P., and Spyrou, G.M. (2021). Generalized linear models provide a measure of virulence for
specific mutations in SARS-CoV-2 strains. PLoS One *16*, e0238665.
10.1371/journal.pone.0238665.

Pavesi, A. (2020). New insights into the evolutionary features of viral overlapping genes by
discriminant analysis. Virology *546*, 51-66. 10.1016/j.virol.2020.03.007.

Strahle, L., Garcin, D., and Kolakofsky, D. (2006). Sendai virus defective-interfering genomes
and the activation of interferon-beta. Virology *351*, 101-111. 10.1016/j.virol.2006.03.022.

Tischer, B.K., von Einem, J., Kaufer, B., and Osterrieder, N. (2006). Two-step red-mediated
recombination for versatile high-efficiency markerless DNA manipulation in Escherichia coli.
Biotechniques *40*, 191-197. 10.2144/000112096.
Torii, S., Ono, C., Suzuki, R., Morioka, Y., Anzai, I., Fauzyah, Y., Maeda, Y., Kamitani, W.,
Fukuhara, T., and Matsuura, Y. (2021). Establishment of a reverse genetics system for SARS-
CoV-2 using circular polymerase extension reaction. Cell Rep *35*, 109014.
10.1016/j.celrep.2021.109014.
Varga, Z.T., Grant, A., Manicassamy, B., and Palese, P. (2012). Influenza virus protein PB1-
F2 inhibits the induction of type I interferon by binding to MAVS and decreasing mitochondrial
membrane potential. J Virol *86*, 8359-8366. 10.1128/JVI.01122-12.
Varga, Z.T., Ramos, I., Hai, R., Schmolke, M., Garcia-Sastre, A., Fernandez-Sesma, A., and
Palese, P. (2011). The influenza virus protein PB1-F2 inhibits the induction of type I interferon
at the level of the MAVS adaptor protein. PLoS Pathog *7*, e1002067.
10.1371/journal.ppat.1002067.
Xia, H., Cao, Z., Xie, X., Zhang, X., Chen, J.Y., Wang, H., Menachery, V.D., Rajsbaum, R.,
and Shi, P.Y. (2020). Evasion of Type I Interferon by SARS-CoV-2. Cell Rep *33*, 108234.
10.1016/j.celrep.2020.108234.

Fig. 1

Fig. 2

Fig. 3

A

SARS-CoV Tor2

bat CoV Rs4231

civet CoV 007

Sarbecovirus

SARS-CoV-2 Wuhan Hu-1

bat CoV BANAL-20-52

pangolin CoV GX-P4L

Hibecovirus

bat CoV Hp-betacoronavirus/Zhejiang2013

Nobecovirus

bat CoV GCCDC1

B

```
                1                                                 41
Wuhan Hu-1      MLLLQILFAL LQRYRYKPHS LSDGLLLALH FLLFFFRALPK S

      RmYN02    .......... .......... .......... .......... .
 BANAL-20-103   .......... .......... .......... ...K..... .
 BANAL-20-236   .......... .......... .......... ...K..... .
       RaTG13   .......... .......... .......... ...K...R. .
    RShSTT200   ..H....... .......... .......... .......... .
    RShSTT182   ..H....... .......... .......... .......... .
  BANAL-20-52   .......... ..P....... .......... .......... .
        ZXC21   .......... ..Q....... .......... ..Q..K..Q. -

          P2S   .......... .......... .......... ...K..... -
          P1E   .......... .......... .......... .......... -
          P4L   .......... .......... .......... ....T.... -

         Tor2   ....V..M.. Q......... .......... .......... -
          SZ3   ....V..M.. Q......... .......... .......... -

       Rs4231   ....V..M.. Q....R.... .......... .......... -

          007   ...M.V..V. Q......... .......... .......... -
```

SARS-CoV-2 cluster

SARS-CoV cluster

C

IFN-β promoter activity (%)

- Tor2
- Wuhan Hu-1
- BANAL-20-52
- ZXC21

vector   NS1   ORF3c

ns   ****   ns   ****
***
**   ****   ***   ****
***   ****   ****   ****
****   ****   ****   ****

Fig. 4

Fig. 5

Fig. S1