Stress Recovery Triggers Rapid Transcriptional Reprogramming and Activation of Immunity in

Plants

Natanella Illouz-Eliaz[1,2], Jingting Yu[3], Joseph Swift[1,2], Kathryn Lande[3], Bruce Jow[2], Za Khai Tuang[4],

Travis Lee[1,2,6], Adi Yaaran[5], Rosa Gomez Castanon[2], Joseph R. Nery[2], Tatsuya Nobori[1,2], Yotam Zait[5], Saul

Burdman[4], and Joseph R. Ecker[1,2,4*]

[1]Plant Biology Laboratory, The Salk Institute for Biological Studies, La Jolla, CA 92037, United States

[2]Genomic Analysis Laboratory, The Salk Institute for Biological Studies, La Jolla, CA 92037, United States

[3]The Razavi Newman Integrative Genomics and Bioinformatics Core Facility, The Salk Institute for

Biological Studies, La Jolla, CA92037, United States

[4]Department of Plant Pathology and Microbiology, Institute of Environmental Sciences, The Robert H.

Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem,

Rehovot 7610001, Israel

[5]Institute of Plant Sciences and Genetics in Agriculture, The Robert H. Smith Faculty of Agriculture, Food

and Environment, The Hebrew University of Jerusalem, Rehovot 7610001, Israel

[6]Howard Hughes Medical Institute, The Salk Institute for Biological Studies, La Jolla, CA 92037, United

States

* Corresponding author: ecker@salk.edu

Keywords: Stress Recovery, Drought Stress, Plant Immunity, Plant Genomics, Single-Cell.

Summary

All organisms experience stress as an inevitable part of life, from single-celled microorganisms to complex

multicellular beings. The ability to recover from stress is a fundamental trait that determines the overall

resilience of an organism, yet stress recovery is understudied. To begin unraveling the stress recovery

process we studies recovery from drought stress in *Arabidopsis thaliana*. We performed a fine-scale time

series of bulk RNA sequencing starting 15 minutes after rehydration following moderate drought. We reveal

that drought recovery is a rapid process involving the activation of thousands of recovery-specific genes.

To capture these rapid recovery responses in different leaf cell types, we performed single-nucleus

transcriptome analysis at the onset of post-drought recovery, identifying a cell type-specific transcriptional

state developing within 15 minutes of rehydration independently across cell types. Furthermore, we reveal

a recovery-induced activation of the immune system that occurs independent of infection, which enhances

pathogen resistance *in vivo* in *A. thaliana*, wild tomato (*Solanum pennellii)* and domesticated tomato

(*Solanum lycopersicum* cv. M82)*. Since rehydration promotes microbial proliferation and thereby increases

the risk of infection[1-2], drought recovery-induced immunity may be crucial for plant survival in natural

environments. These findings indicate that drought recovery coincides with a preventive defense response,

unraveling the complex regulatory mechanisms that facilitate stress recovery in different plant cell types.

Main

In most plants, extended periods of water deficit result in reduced growth, premature flowering, flower abortion, fruit abscission, and, ultimately, decreased yield[3-4]. Plant responses to drought have therefore been studied extensively as part of efforts to develop strategies or genetic manipulations that could mitigate the economic and agricultural consequences of future droughts.

Previous studies have identified and functionally characterized numerous genes that respond to water deficit, including various transcription factors (TFs) that regulate plant drought responses[4-6]. In general, drought stress in plants induces dramatic changes in the transcriptional landscape[7-8]. For example, the rapid up-regulation of genes involved in osmolyte accumulation enables water retention by adjusting cellular osmotic potential[9]. However, efforts to enhance drought tolerance through genetic manipulation have frequently resulted in undesired growth inhibition under non-drought conditions, thereby constraining the widespread engineering and adoption of drought-resilient crops[6,10-11].

Given these limitations, we considered an alternative approach to investigating stress resilience in plants by focusing on post-drought recovery rather than drought resistance. Indeed, understanding a plant's ability to recover from drought is central to a comprehensive understanding of drought resilience, as the potential for recovery defines whether the system can return to a stable state of function[12]. For example, drought recovery has been established as an indicator of drought tolerance in annual crops such as maize[13-14], wheat[15-16] and rice[17-18]. This extends to other abiotic stresses as well, for instance, the rates of submergence recovery in *Arabidopsis thaliana* (Arabidopsis) accessions were found to correlate with their submergence tolerance[19]. This finding suggests that the ability to recover after flooding is critical for plant survival and reproductive success. One of the well-characterized plant responses to drought alleviation is the downregulation of the drought responsive genes. For example, most drought-regulated genes recover to normal expression levels within three hours of rehydration[20]. However, Oono et al.[21] identified 82 "recovery-specific" genes whose expression was drought-invariable but altered by subsequent rehydration.

Despite these known transcriptional responses during drought recovery, it remains unclear whether these responses constitute a conserved drought recovery mechanism activated throughout the entirety of

plant cell types. Here, we explore the transcriptomic landscape in Arabidopsis throughout the drought recovery process using a high-resolution time series of RNA sequencing data. We identified over 3,000 recovery-specific genes that were differentially expressed between 15 minutes and 6 hours after rehydration. Using single-nucleus RNA-seq, we analyzed cell type-specific transcriptional signatures immediately after inducing drought recovery. We identified a recovery unique transcriptional signature common between sub-population of epidermal, hydathode, sieve element and mesophyll cells that is found only once drought recovery is initiated. Within the gene modules enriched in sub-populations we found many immune-associated genes. Based on these observations, we propose that some cells enter a recovery cell state upon rehydration. The results indicated that one of the first steps of the recovery process is the activation of a rapid preventive immune response. We therefore tested whether short-term recovery from moderate drought activated functional immunity to block pathogen proliferation in plant leaves. In agreement with this hypothesis, we found that drought recovery-induced activation of immune system genes leads to reduced disease severity and bacterial load in infected leaves of both Arabidopsis and two tomato species.

Drought recovery-specific genes

To study the transcriptional response to drought recovery, we performed a fine-scale RNA-seq time series using vermiculite-grown Arabidopsis rosettes. We collected Arabidopsis rosettes during moderate drought and at seven subsequent recovery time points within the six hours following rehydration (Fig. a-b). At each time point, well-watered rosettes were also collected to control for diurnal transcriptional changes (Fig. 1b). We then conducted a differential expression (DE) analysis of the treatments and time points and defined two groups of genes: drought-responsive genes and recovery-specific genes. Drought-responsive genes were defined as genes that were DE compared to well-watered plants during the drought stress i.e., at time 0 (Extended Data Table 1). Conversely, recovery-specific genes were not DE during the drought period but were DE in any of the post-drought recovery time points (Extended Data Table 2).

Based on these definitions, we identified 1,248 drought-responsive genes, of which 662 (53%) were

also DE during recovery. After only six hours of rehydration, 97% of these genes returned to normal

expression levels (Fig. 1c-d). We additionally identified over 3,000 recovery-specific genes across all

recovery time points (Fig. 1c and e), suggesting that drought recovery responses cannot be solely explained

by the alleviation of drought induced transcriptional responses. We used K-means clustering to characterize

gene expression patterns during drought and recovery. The drought-responsive genes exhibited five distinct

expression patterns (Fig. 1d): (1) drought-induced but downregulated during recovery, (2) drought-induced

but not DE during recovery, (3) drought- and recovery-induced; (4) drought-downregulated but recovery-

induced; and (5) drought-downregulated but not DE during recovery. Recovery-specific genes exhibited

three types of expression patterns based on whether they were induced in (1) early recovery or (2) late-

recovery or (3) downregulated by rehydration (Fig. 1e). These results demonstrate that drought recovery

involves the activation of thousands of recovery specific genes. These genes are timely regulated, with a

set of genes being upregulated from as early as 15 mins after rehydration, and a later set of genes starting

to accumulate after 30-60 mins.

Global and cell type-specific recovery induced transcriptional reprogramming

To create study the leaf cell type-specific transcriptional responses upon drought recovery, we

performed single-nucleus RNA-seq (snRNA-seq). For this experiment, we processed two replicates

undergoing long-term moderate drought and an additional two replicates after 15 minutes of rehydration,

as well as equivalent samples from well-watered controls. In this case, the well-watered controls were also

provided with 15 minutes of additional irrigation to ensure that any changes in gene expression were

specific to drought recovery and not due to mechanical root stress (Fig. 2a). After data quality control and

filtering, our integrated dataset included over 144,000 single-nuclei transcriptomes. The median number of

unique molecular identifiers (UMI) per nucleus was between 2,534 to 3,995 across the eight samples, and

the median number of genes detected per nucleus was between 1,213 to 1,698 (average 1,530; see Extended

Data Table 3).

Unsupervised clustering of the snRNA-seq profiles resulted in 27 clusters representing unique

molecular cell identities (Fig. 2b, Extended Data Figs. 1-2). All independent samples and treatment

conditions integrated into the same clusters (Extended Data Fig. 1). We used three major practices to

annotate cell identities for these clusters. First, we overlapped each cluster's marker genes (genes uniquely

or highly expressed in a certain cluster) with previously validated marker genes for all leaf cell types and

tissues (Extended Data Tables 4-5 and Extended Data Figs. 3-4). Second, we projected our dataset on

published single-cell datasets preformed on Arabidopsis leaves (Extended Data Fig. 5). Third, we manually

investigated the top marker genes in each cluster and used Gene Ontology (GO) enrichment analysis to see

enriched biological processes (Unknown clusters; Extended Data Fig. 6). Based on the genes expressed in

each cluster and their expression levels, we were able to confidently assign cell types to 22 of the 27 clusters.

Once the clusters had been assigned to cell types, we investigated cell type-specific changes during

the onset of drought recovery. A transcription factor (TF) expression analysis across the different treatments

revealed that TFs are rapidly induced by rehydration across all cell types. These TFs had many genes from

the *ETHYLENE RESPONSE FACTOR* (*ERF*) and *WRKY DNA-BINDING PROTEIN* (*WRKY*) families (Full

list in: Extended Data Table 6, GO term enrichment analysis in: Extended Data Fig. 7). In contrast, some

cell types, such as myrosin and vascular cells exhibited cell type-specific changes in TF expression upon

recovery. An interesting group of TFs were rapidly activated in a contextual manner that was both treatment

specific and cell state specific. These TFs were upregulated only in dividing cells at the onset of recovery

(Fig. 2d). Four gene families dominated this set of genes: MADS box, B3, NO APICAL MERISTEM

(NAC), and PAIRED AMPHIPATHIC HELIX (PAH2) families (Full list in: Extended Data Table 7).

Based on a sequence similarity analysis, the highly activated genes within each family were

phylogenetically clustered. For example, the upregulated MADS box genes were all type I MADS box (M-

type) genes: *AGAMOUS-LIKE43* (*AGL43*), *AGL89*, *AGL77*, *AGL75*, *AGL74*, *AGL84*, and *AGL50*.

Similarly, all the B3 TFs induced by drought recovery in dividing cells were in the REPRODUCTIVE

MERISTEM (REM) branch[22], despite the B3 family containing multiple distinct gene subfamilies[23]. These

findings suggest that drought recovery triggers cell type-specific transcriptional reprogramming.

Recovery cell states in distinct cell types

To study heterogeneity within cell types during drought recovery, we further subclustered cells within each cell type based on their gene expression profiles (Extended Data Figs. 8-9). In epidermal, mesophyll, hydathode, and sieve-element cells, some of the cell subpopulations were enriched (i.e., over 50% of cells in the subcluster) with cells from the drought recovery treated plants (Fig. 3a, Extended Data Fig. 9). We hypothesized that these cell subpopulations form distinct subclusters due to their transition into a recovery cell state (RcS). To test this hypothesis, we performed a motif enrichment analysis on each of the suspected RcS subclusters to determine whether we could identify regulatory motifs that were common to the putative RcS across all cell types. We found that all the subclusters that we identified as having an activated RcS were enriched with the *CALMODULIN BINDING TRANSCRIPTION ACTIVATOR 1* (*CAMTA1*) and *CAMTA5* binding motifs (Fig. 3b). In Arabidopsis, the CAMTA family of TFs regulate plant defense and stress responses by modulating the expression of genes involved in pathogen defense, abiotic stress tolerance, and plant development. These TFs interact with calcium-bound calmodulin to mediate downstream signaling pathways[24]. To investigate if CAMTAs specifically regulate the RcS, we performed motif enrichment analysis on all subclusters to see if the *CAMTA1* and *CAMTA5* motifs were enriched in other subclusters. Out of 162 subclusters, only 9 subclusters were enriched with *CAMTA* motifs (Extended Data Table 8, and Extended Data Fig. 10), suggesting that *CAMTA1* and *CAMTA5* are nearly exclusively enriched in RcS subclusters, and could be the regulators of the formation of this cell state.

We predicted that *CAMTA1* and *CAMTA5* regulate the RcS; however, neither the CAMTA genes nor the cell type-specific TFs predicted to regulate the RcS were upregulated in response to rehydration (Fig. 3c). Since the RcS is rapidly activated, we hypothesized that *CAMTA* transcripts may be differentially expressed during the early stages of drought and would therefore not be observed in our data, which was collected after moderate drought had established. To test this hypothesis, we performed an additional snRNA-seq experiment on Arabidopsis plants harvested each day over the course of five days starting from the onset of drought (Fig. 3d, Extended Data Fig. 11). Cell identities were identified using unsupervised

clustering, as before. We then investigated the expression of *CAMTA* genes in different cell types during

early drought. In most cell types, *CAMTA* genes were upregulated in the days following dehydration (Fig.

3e, Extended Data Fig. 12), suggesting that the regulation of *CAMTA* is very sensitive to changes in water

availability and that *CAMTA* transcripts accumulate early in response to drought.

In addition to testing for a common regulatory pathway, we investigated whether the RcS

subpopulations within the different cell types expressed similar gene networks by performing gene co-

expression network analysis[25] for the seven cell clusters with an activated RcS subpopulation in the drought

recovery experiment (Fig. 3a). For each cluster, this analysis identified gene modules (Extended Data Fig.

13-14). We next searched for common hub genes (Genes with high module membership (kME) and high

intramodular connectivity (kIN)) among the modules that were enriched in RcS cells from different cell

types to identify genes that have common key roles in the formation of recovery cell states or its function.

Overall, we found 212 hub genes, of which 50 were shared by at least two modules (Extended Data Table

9). Interestingly, in the immediate recovery snRNAseq, 70% of the top marker genes for cluster "unknown

1" overlap with the hub genes identified in the RcS gene network. This finding led us to hypothesize that

cluster "unknown 1" represents a cell state that is superimposed across different cell types at the onset of

drought recovery (Extended Data Fig. 15).

Overall, the suite of common hub genes among the six RcS subclusters from different cell types

suggest that the functional onset of drought recovery in RcS subpopulations is comprised of cell wall

modifications and the regulation of nutrient uptake, as well as cytoplasmic detoxification processes and

DNA repair. The most prevalent hub genes shared among RcS subclusters are genes that play a critical role

in cell growth and plant development; these genes included *XYLOGLUCAN*

ENDOTRANSGLUCOSYLASE/HYDROLASE 22 (*XTH22*)[26], *SLAC1 HOMOLOGUE 3* (*SLAH3*)[27], and

EXORDIUM-LIKE1 (*EXL1*)[28]. Since toxins accumulate in plant cells during drought stress[29], we suspected

that detoxification may also be an important part of recovery. Indeed, our cross-cluster hub gene analysis

implicated *ARABIDOPSIS THALIANA DETOXIFICATION 1* (*AtDTX1*), which is localized in the plasma

membrane of plant cells and mediates the efflux of plant-derived or exogenous toxic compounds from the

cytoplasm[30]. An additional hub gene common across RcS cell populations was *CHROMATIN ASSEMBLY*

FACTOR 1 (*CAF-1*) *AtCAF1a*, which facilitates the incorporation of histones H3 and H4 onto newly

synthesized DNA. The absence of the CAF-1 chaperone complex results in mitotic chromosome

abnormalities and changes in the expression profiles of genes involved in DNA repair[31]. In addition,

AtCAF1 proteins regulate mRNA deadenylation and defenses against pathogen infections[32]. Taken

together, these finding suggest that drought recovery triggers the formation of a specific cell state imposed

within several leaf cell types.

To identify more general processes involved in drought recovery, we performed a cell type-specific

differential expression analysis comparing plants harvested 15 minutes after the onset of drought recovery

to (i) plants experiencing drought and (ii) the well-watered controls that received additional irrigation for

15 min. Both controls were used for DE analysis to ensure that DE genes were not simply maintained from

their changes during drought or induced by the mechanical stress of additional watering. In nearly all cell

types, photosynthesis and carbon fixation were suppressed during the onset of drought recovery. The only

exception was in one of the seven mesophyll cell clusters, which maintained photosynthetic activity but

downregulated the expression of genes associated with bacterial responses (Extended Data Fig. 16a).

Conversely, genes that were upregulated at the onset of recovery were associated with diverse stressors

including wounding, cold, heat, light, and reactive oxygen species (i.e., oxidative stress). The upregulation

of such a broad suite of stress response genes suggests that the plant perceives the onset of recovery as a

state of stress, perhaps because recovery presents a change in the current homeostasis that was established

during drought.

Recovery-induced immune responses

We found evidence for the activation of multiple immunity-related genes in the top shared hub

genes among RcS subclusters. Among these genes for example, *TETRASPANIN 8* (*TET8*) has been shown

to be upregulated in response to the Flg22 and Elongation Factor Tu (EF-Tu) elicitors, which mimic

pathogen infection[33]. Another shared hub gene *SALT-INDUCIBLE ZINC FINGER 1* (*SZF1*) also regulates

plant immunity, as *szf1,2* knock-out mutants show increased susceptibility to *Pseudomonas syringae* pv.

tomato DC3000 (*Pst* DC3000)[34]. Three other shared hub genes (*AT5G41750, AT5G41740* and *AT4G19520*)

encode TOLL INTERLEUKIN RECEPTOR (TIR)–type NB-LRR proteins and thus belong to the most

common class of disease resistance genes in plants[35]. Another example is *ACTIVATED DISEASE*

RESISTANCE 2 (*ADR2*) that enhances resistance to biotrophic pathogens[36]. Finally, we also found *EARLY*

RESPONSIVE TO DEHYDRATION 15 (*ERD15*) as a top shared hub genes within the RcS enriched

modules, which belongs to a small and highly conserved protein family that is ubiquitous but specific to

the plant kingdom. Overexpression of ERD15 proteins in response to various pathogen elicitors was shown

to improve resistance to known pathogens and has also been shown to impair drought tolerance[37]. Thus, the

activation of a gene such as *ERD15* specifically upon rehydration maybe beneficial to contribute to plant

defense without compromising the drought response.

Other genes upregulated during the recovery initiation were implicated in processes associated with

plant immunity, including genes involved in response to bacterial pathogens, immune system processes,

responses to jasmonic acid (JA), and cell death. We found additional processes associated with plant

defense that were upregulated in specific cell types; for example, genes involved in the responses to

oomycetes or ethylene response were upregulated in almost all cell types during recovery, whereas defense

responses against insects were only upregulated in one-third of epidermal cell clusters, 50% of vascular cell

clusters, and two of the "unknown" clusters (Extended Data Fig. 16b).

We suspected that the transcriptomic upregulation of immune system processes during drought

recovery is an innate response to post-drought rehydration rather than a proactive response to microbial

infection following rehydration. To test this hypothesis, we examined the upregulation of recovery induced

genes under axenic conditions. Arabidopsis seedlings were grown on sterile agar plates for 14 days and

then transferred to low-water-content agar plates[38], which we produced by decreasing the water content of

the media to 50% (thus imposing moderate osmotic stress) (Extended Data Fig. 17a-b). After 14 days on

the low-water-content plates, plants were rehydrated and collected at different recovery time points (0, 15,

30, 90 and 120 min) for bulk RNA-seq analysis. The upregulation of the known drought marker genes

RESPONSIVE TO DESICCATION 29A (RD29A), RD20, and *DELTA1-PYRROLINE-5-CARBOXYLATE*

SYNTHASE 1 (P5CS)[39-41] confirmed that the plants were experiencing moderate drought stress (Extended

Data Fig. 17c). We additionally found that immune-related genes were upregulated after 15 minutes of

rehydration even in the axenic system (Extended Data Fig. 17d). These results support the hypothesis that

plants activate a prophylactic defense response upon recovery from moderate drought, independent of

pathogens in their environment.

Given the strength of this immunity-related transcriptional response during drought recovery, we

used data from Bjornson et al.[42] to evaluate how the genes implicated in our analysis correlated with known

transcriptional responses to biotic elicitors. Bjornson et al.[42] performed transcriptomic analysis across a

fine-scale time series to study rapid signaling transcriptional outputs induced by well-characterized elicitors

of pattern-triggered-immunity (PTI) in Arabidopsis. Their results showed that the transcriptional responses

to diverse microbial- or plant-derived molecular patterns are highly conserved. When we aligned these

identified transcriptional responses with the recovery specific genes identified by the bulk RNA-seq data

from the first two hours of our drought recovery time series (Fig. 1), we found that 50% of recovery-specific

genes upregulated after 15 minutes of rehydration overlap with the core responses to biotic elicitors in the

Bjornson et al.[42] dataset. More generally, this analysis shows a rapid and robust peak of immune-relevant

gene expression that gradually decreased with time since rehydration (Fig. 4a).

Drought recovery-induced immunity *in vivo*

To test the functionality of post-recovery immune activation, we examined whether the activation

of immunity-related genes during short-term recovery from moderate drought promotes pathogen resistance

in vivo. We used the sterile low-water-content plate system and examined recovery from both moderate

(40% water-content) and severe stress (25% water-content), as described above. Plants from the control

and moderate or severe osmotic stress conditions were rehydrated for 90 minutes and then inoculated with

Pst DC3000 via submersion in a bacterial suspension. Whole rosettes were collected and weighed 48 hours

following infection and used to quantify bacterial growth. Plants recovering from drought had significantly

lower bacterial concentrations than control plants, and recovery from moderate stress suppressed bacterial

growth better than recovery from severe stress (Fig. 4b-c). These results suggest that recovery from

moderate drought enhances resistance to *Pst* DC3000.

To further validate these results, we grew plants on soil (non-sterile conditions) for 30 days under

short-day conditions. Drought-treated plants were transferred to dry trays and dehydrated for one week

down to 30% relative soil water content, while control plants continued to receive regular irrigation. We

rehydrated drought-treated plants for 90 minutes and then sprayed the leaves of both control and recovered

plants with *Pst* DC3000. Leaf discs were collected from each treatment 4 and 24 hours after inoculation,

surface sterilized, and then used to measure bacterial load. The bacterial load 4 hours after inoculation did

not differ between well-watered and drought-recovered plants. However, 24 hours after inoculation, the

bacterial load of drought-recovered plants was significantly lower than that of controls, indicating increased

resistance to *Pst* DC3000 in rehydrated plants (Fig. 4d-e). We call this response, which was consistent

across our two separate empirical tests, drought recovery-induced immunity (DRII).

Given the consistency of the DRII response in Arabidopsis, we further tested whether DRII is

conserved among various plant species and in response to different pathogens. First, we examined pathogen

proliferation in wild tomato plants recovering from moderate drought and infected with either *Pst* DC3000

or an additional tomato pathogen *Xanthomonas perforans* strain 97-2, which causes the tomato spot

disease[43]. For this test, we used the drought-tolerant tomato species *Solanum pennellii*[44]. We grew *S.*

pennellii plants at 25°C with a photoperiod of 12 hours light and 12 hours dark. When the plants had two

to three true leaves (~4 weeks old), we exposed them to moderate drought by stopping irrigation until the

soil reached 30% water content relative to saturated pots, as we had done for Arabidopsis. Drought was

then alleviated by irrigating pots to full saturation, and plants were infected 90 minutes after rehydration.

Both the drought-recovered plants and well-watered controls were syringe-infiltrated with a suspension of

either *X. perforans* 97-2 or *Pst* DC3000.

We first assessed disease severity in the well-watered and drought-recovered infected leaves by

imaging inoculated leaves 5 days post infection and calculating the percentage of symptomatic area relative

to the entire leaf surface. Leaves inoculated 90 minutes after recovery from moderate drought were

significantly less symptomatic than leaves from well-watered controls (Fig. 5a-b). To directly measure

bacterial concentrations in leaves, we compared leaf samples at an initial time point 4 hours after inoculation

to samples collected 5 days post-inoculation. Although a similar number of bacteria entered the leaves of

all plants, the plants that recovered from drought had a significantly lower bacterial load after five days

(Fig. 5c). Our results were consistent in both tested pathogens: regardless of the pathogen, tomato leaves

that recovered from drought exhibited lower disease severity and reduced bacterial concentrations than

control plants. Our observation that short-term recovery from moderate drought increases pathogen

resistance across all treatments further suggests that DRII is a preventive immune response that enhances

pathogen resistance during the initial period of drought recovery.

To further investigate DRII in an agriculturally relevant plant species, we repeated our post-drought

immune challenge experiment using a cultivar of domesticated tomato, *Solanum lycopersicum* cv. M82

(M82), and the same two pathogens as before (*Pst* DC3000 and *X. perforans* 97-2). Because breeders focus

on traits such as fruit size, shelf life, and brix, among others, we suspected that the DRII response could be

lost during domestication. However, M82 plants showed similar results to *S. pennellii* upon infection with

either pathogen (Fig. 6d-f), indicating that the DRII response we observed in Arabidopsis and *S. pennellii*

following short-term recovery from moderate drought is retained in domesticated tomato. We think that the

robustness of the DRII trait to the domestication processes underscores its potential significance in

agricultural contexts.

Conclusion

As large regions of the world gradually become drier, drought has become one of the most pressing

threats to global food security[45-46]. Plant drought responses have therefore been studied extensively over the

past decades, but the molecular mechanisms for drought recovery have been largely overlooked. We

advanced the understanding of drought recovery processes by using bulk RNA-seq over a six-hour time

series to identify recovery-specific genes. The resulting data enriches the pool of potential targets for

developing drought-tolerant crops. For example, the genes whose expression was drought-invariable but

downregulated during recovery ("cluster 3" in Fig. 1e) could be viable candidates for mutation screens

aimed at improving post-drought effects and recovery dynamics without affecting drought resistance.

Furthermore, the construction of a single-cell RNA-seq-based cell atlas of plants recovering from moderate

drought allowed us to explore cell type-specific transcriptional changes in response to drought and

immediate recovery. We were able to identify the major cell types in our dataset, and our findings suggest

that the five unidentified clusters (Fig. 2b) most likely contain multiple cell types that clustered together

because they were engaging in similar molecular or biochemical process during drought recovery, and

therefore cluster by the transcriptional signature of a specific cellular state. As plants must constantly adapt

to environmental changes, the molecular phenotypes of individual cells might vary to facilitate the

metabolic and biochemical changes essential for adaptation to the current stressor.

Using our drought recovery single cell atlas, we identified the first regulatory steps underlying the

transcriptional plasticity that allows plants to restore normal growth following drought. One of these steps

involves upregulating a set of TFs specifically in dividing cells within 15 minutes of rehydration. However,

little is known about the TFs identified by this analysis, although they largely belonged to four known TF

families—the MADS box, B3, NAC, and PAH2 families. The MADS box family of TFs are well known

for regulating flowering[47-48]. However, *AGL16* was shown to negatively regulate drought resistance via its

effects on stomata[49]. The B3 family members contains a highly conserved DNA-binding domain and are

involved in a variety of biological processes, many of which are related to the regulation of flowering

through polycomb silencing[50-52]. The NAC TFs play important roles in development as well as abiotic and

biotic stress responses[53]. Overall, the within-family phylogenetic proximity of the specific TFs upregulated

by recovery suggest that these TFs act together to play a functional role in drought recovery, though

determining the specific role of each TF will require further investigation of higher-order mutants.

Recently, Oliva et al.[55] characterized a highly active transcriptional state that affects a subset of

cells in different cell types across the developmental axis of the Arabidopsis root tip. The conserved cell

state was named environmentally responsive state (ERS), since many ERS enriched genes were previously

identified as responsive to various environmental cues. Here, we identified a transcriptional recovery cell

state (RcS) that is triggered by environmental stimuli. This state emerges in different cell types rapidly and

independently. We hypothesize that the RcS formation is governed by a small set of TFs, some of which

are common among cell types such as *CAMTA1* and *CAMTA5*. It is possible that some cell type-specific

TFs are also involved in the regulation of RcS depending on the cell type that the state was imposed on.

Our findings also show that transcripts for the calcium dependent *CAMTA* TFs accumulate during

early drought, and that, although the upregulation of *CAMTAs* does not persist during drought recovery,

many of the genes abundant during the RcS are regulated by *CAMTA1* and *CAMTA5*. Previous studies have

shown that calcium flux can be affected by changes in water availability (both water deficit and water

uptake)[54]. We therefore hypothesize that *CAMTA1 and CAMTA5* expression is induced during the early

stages of soil dehydration and that CAMTA proteins are post-translationally stabilized. CAMTAs activate

gene expression by interacting with calcium-bound calmodulin (CaM), and the influx of calcium upon

rehydration leads to an increase in calcium-bound calmodulin that then activates CAMTA proteins to bind

DNA and hence regulate the subsequent rehydration response. CaM levels may therefore be a limiting

factor in the regulation of the rapid CAMTA1-dependent gene activation that we observed in the initial

stages of drought recovery.

Our analysis suggest that the onset of drought recovery is coupled with the activation of a

preventive immune response. Plants have evolved complex defense networks in response to microbial

pathogens[35]. Aerial pathogens like *Pst* DC3000 or *X. perforans* mainly enter plants through stomatal pores,

but the stomata close in response to drought stress[56], making it harder for shoot pathogens to successfully

attack plants experiencing drought. It should be noted that some pathogenic bacteria have evolved specific

virulence factors to cause stomatal opening[57]. Moreover, plant immune responses can be sporadically

induced by the circadian clock even in the absence of a pathogenic threat, a process driven largely by daily

oscillations in humidity[58-59]. Such responses allow plants to prepare for the increased risk of infection when

microbes are anticipated to be most infectious. Because rehydration promotes both pathogen proliferation

and stomatal opening at a time when the immune system has already been suppressed by drought, plants

are particularly vulnerable to pathogen attack during the initial stages of drought recovery[60-61]. The rapid

upregulation of immunity genes that we observed in our study may therefore be crucial for ensuring plant

survival in water-fluctuating environments. We accordingly propose that DRII is a preventive immune

response that evolved to confer resistance against pathogens during drought recovery.

More broadly, our research shifts the focus from the well-studied drought responses to plant

transcriptional activity during the recovery period. Although plant responses to drought have been

extensively studied for over a century, these findings have yet to be successfully translated to the

development of drought-tolerant crops[62]. Our results highlight the presence and importance of recovery-

specific mechanisms that could be targeted in future experiments and engineering projects. We suggest a

new avenue for research in which efforts to improve crop resilience should focus not only on survival during

drought but also on rapid and effective recovery. By targeting recovery-specific genes, we can now

endeavor to develop crops that not only withstand drought but also recover more swiftly, ensuring minimal

yield loss and sustainable food production in an era of increasing climate unpredictability.

Main references

1. Palmero, D. *et al.* Fungal microbiota from rain water and pathogenicity of Fusarium species isolated from atmospheric dust and rainfall dust. *J. Ind. Microbiol. Biotechnol.* **38**, 13–20 (2011).
2. Schwartz, H. F., Otto, K. L. & Gent, D. H. Relation of temperature and rainfall to development of Xanthomonas and Pantoea leaf blights of onion in Colorado. *Plant Dis.* **87**, 11–14 (2003).
3. Campos, H., Cooper, M., Habben, J. E., Edmeades, G. O. & Schussler, J. R. Improving drought tolerance in maize: A view from industry. *F. Crop. Res.* **90**, 19–34. (2004).
4. Bruce, W. B., Edmeades, G. O. & Barker, T. C. Molecular and physiological approaches to maize improvement for drought tolerance. *J. Exp. Bot.* **53**, 13–25. (2002).
5. Singh, D. & Laxmi, A. Transcriptional regulation of drought response: A tortuous network of transcriptional factors. *Front. Plant Sci.* **6**, 1–11. (2015).
6. Shim, J. S. *et al.* Overexpression of OsNAC14 improves drought tolerance in rice. *Front. Plant Sci.* **9**, 1–14. (2018).
7. Shinozaki, K., Yamaguchi-Shinozaki, K. & Seki, M. Regulatory network of gene expression in the drought and cold stress responses. *Curr. Opin. Plant Biol.* **6**, 410–417 (2003).
8. Soma, F., Takahashi, F., Yamaguchi-Shinozaki, K. & Shinozaki, K. Cellular phosphorylation signaling and gene expression in drought stress responses: Aba-dependent and aba-independent regulatory systems. *Plants* **10**, 1–16. (2021).
9. Zhang, J., Nguyen, H. T. & Blum, A. Genetic analysis of osmotic adjustment in crop plants. *J. Exp. Bot.* **50**, 291–302. (1999).
10. Qin, F. *et al.* Regulation and functional analysis of *ZmDREB2A* in response to drought and heat stresses in *Zea mays* L. **50**, 54–69. (2007).
11. Illouz-Eliaz, N. *et al.* Mutations in the tomato gibberellin receptors suppress xylem proliferation and reduce water loss under water-deficit conditions. *J. Exp. Bot.* **71**, 3603–3612. (2020).
12. Gessler, A., Bottero, A., Marshall, J. & Arend, M. The way back: recovery of trees from drought and its implication for acclimation. *New Phytol.* **228**, 1704–1709. (2020).
13. Hayano-Kanashiro, C., Calderón-Vázquez, C., Ibarra-Laclette, E., Herrera-Estrella, L. & Simpson, J. Analysis of gene expression and physiological responses in three Mexican maize landraces under drought stress and recovery irrigation. *PLoS One.* **4**(10), e7531. (2009).
14. Chen, D. *et al.* Genotypic variation in growth and physiological response to drought stress and re-watering reveals the critical role of recovery in drought adaptation in maize seedlings. *Front Plant Sci.* **6**(1241), 1-15. (2016).
15. Vassileva, V., Signarbieux, C., Anders, I. & Feller, U. Genotypic variation in drought stress response and subsequent recovery of wheat (*Triticum aestivum L.*). *J Plant Res.* **124**(1), 147-154. (2011).
16. Abid, M. *et al.* Adaptation to and recovery from drought stress at vegetative stages in wheat (*Triticum aestivum*) cultivars. *Funct Plant Biol.* **43**(12), 1159-1169. (2016).
17. Kamoshita, A., Rodriguez, R., Yamauchi, A., & Wade, L. Genotypic variation in response of rainfed lowland rice to prolonged drought and rewatering. *Plant Production Science.* **7**(4), 406-420. (2004).
18. Efisue, A., Tongoona, P., Derera, J. & Ubi, B.E. Screening early generation progenies of interspecific rice genotypes for drought stress tolerance during vegetative phase. *J Crop Improv.* **23**, 174–193. (2009).
19. Yeung, E. *et al.* A stress recovery signaling network for enhanced flooding tolerance in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E6085–E6094 (2018).
20. Huang, D., Wu, W., Abrams, S. R. & Cutler, A. J. The relationship of drought-related gene expression in *Arabidopsis thaliana* to hormonal and environmental factors. *J. Exp. Bot.* **59**, (2008).
21. Oono, Y. *et al.* Monitoring expression profiles of Arabidopsis gene expression during rehydration process after dehydration using ca. 7000 full-length cDNA microarray. *Plant J.* **34**, 868–887

(2003).

22. Romanel, E. A. C., Schrago, C. G., Couñago, R. M., Russo, C. A. M. & Alves-Ferreira, M.
Evolution of the B3 DNA binding superfamily: New insights into REM family gene
diversification. *PLoS One* **4**, (2009).

23. Wang, Y. *et al.* Systematic analysis of plant-specific B3 domain-containing proteins based on the
genome resources of 11 sequenced species. *Mol. Biol. Rep.* **39**, 6267–6282 (2012).

24. Iqbal, Z., Shariq Iqbal, M., Singh, S. P. & Buaboocha, T. Ca2+/Calmodulin Complex Triggers
CAMTA Transcriptional Machinery Under Stress in Plants: Signaling Cascade and Molecular
Regulation. *Front. Plant Sci.* **11**, 1–16 (2020).

25. Morabito, S., Reese, F., Rahimzadeh, N., Miyoshi, E. & Swarup, V. hdWGCNA identifies co-
expression networks in high-dimensional transcriptomics data. *Cell Rep Methods*. **3**(6), 100498.
(2023).

26. Xu, W. *et al.* Arabidopsis TCH4, regulated by hormones and the environment, encodes a
xyloglucan endotransglycosylase. *Plant Cell* **7**, 1555–1567 (1995).

27. Liu, B. *et al.* The anion channel SLAH3 interacts with potassium channels to regulate nitrogen-
potassium homeostasis and the membrane potential in Arabidopsis. *Plant Cell* **35**, 1259–1280
(2023).

28. Smith, A. M. & Stitt, M. Coordination of carbon supply and plant growth. *Plant, Cell Environ.*
30, 1126–1149 (2007).

29. Ahluwalia, O., Singh, P. C. & Bhatia, R. A review on drought stress in plants: Implications,
mitigation and the role of plant growth promoting rhizobacteria. *Resour. Environ. Sustain.* **5**,
6267-6282 (2021).

30. Li, L., He, Z., Pandey, G. K., Tsuchiya, T. & Luan, S. Functional cloning and characterization of
a plant efflux carrier for multidrug and heavy metal detoxification. *J. Biol. Chem.* **277**, 5360–
5368 (2002).

31. Varas, J., Santos, J. L. & Pradillo, M. The absence of the arabidopsis chaperone complex CAF-1
produces mitotic chromosome abnormalities and changes in the expression profiles of genes
involved in DNA repair. *Front. Plant Sci.* **8**, 1–11 (2017).

32. Liang, W. *et al.* The Arabidopsis homologs of CCR4-associated factor 1 show mRNA
deadenylation activity and play a role in plant defence responses. *Cell Res.* **19**, 307–316 (2009).

33. Wang, F. *et al.* Functional analysis of the Arabidopsis tetraspanin gene family in plant growth and
development. *Plant Physiol.* **169**, 2200–2214 (2015).

34. Kong, L. *et al.* Noncanonical mono(ADP-ribosyl)ation of zinc finger SZF proteins counteracts
ubiquitination for protein homeostasis in plant immunity. *Mol. Cell* **81**, 4591-4604.e8 (2021).

35. Jones, J.D.J., & Dangl, J.L. The plant immune system. *Nature* **444**, 323–329 (2006).

36. Aboul-Soud, M.A.M. *et al.* Activation tagging of ADR2 conveys a spreading lesion phenotype
and resistance to biotrophic pathogens. *New Phytol.* **183**, 1163–1175 (2009).

37. Kariola, T. *et al.* Early responsive to dehydration 15, a negative regulator of abscisic acid
responses in arabidopsis. *Plant Physiol.* **142**, 1559–1573 (2006).

38. Gonzalez, S. & Swift, J. *et al*. Arabidopsis transcriptome responses to low water potential using
high-throughput plate assays. *eLife*. **12**, RP84747 (2023).

39. Msanne, J., Lin, J., Stone, J. M. & Awada, T. Characterization of abiotic stress-responsive
Arabidopsis thaliana RD29A and RD29B genes and evaluation of transgenes. *Planta* **234**, 97–
107 (2011).

40. Aubert, Y. *et al.* RD20, a stress-inducible caleosin, participates in stomatal control, transpiration
and drought tolerance in *Arabidopsis thaliana*. *Plant Cell Physiol.* **51**, 1975–1987 (2010).

41. Székely, G. *et al.* Duplicated P5CS genes of Arabidopsis play distinct roles in stress regulation
and developmental control of proline biosynthesis. *Plant J.* **53**, 11–28 (2008).

42. Bjornson, M., Pimprikar, P., Nürnberger, T. & Zipfel, C. The transcriptional landscape of
Arabidopsis thaliana pattern-triggered immunity. *Nat. Plants.* **7**, 579–586 (2021).

43. Jones, J. B., Lacy, G. H., Bouzar, H., Stall, R. E. & Schaad, N. W. Reclassification of the

xanthomonas associated with bacterial spot disease of tomato and pepper. *Syst. Appl. Microbiol.* **27**, 755–762 (2004).

44. Bolger, A. *et al.* The genome of the stress-tolerant wild tomato species *Solanum pennellii*. *Nat. Genet.* **46**, 1034–1038 (2014).

45. Gupta, A., Rico-Medina, A. & Caño-Delgado, A. I. The physiology of plant responses to drought. *Science.* **368**, 266–269 (2020).

46. CRED. Climate in action Executive Summary. *EM-DAT | Int. disasters database* 8 (2022).

47. Pinyopich, A. *et al.* Assessing the redundancy of MADS-box genes during carpel and ovule development. *Nature* **424**, 85–88 (2003).

48. Callens, C., Tucker, M. R., Zhang, D. & Wilson, Z. A. Dissecting the role of MADS-box genes in monocot floral development and diversity. *J. Exp. Bot.* **69**, 2435–2459 (2018).

49. Zhao, P. X., Miao, Z. Q., Zhang, J., Chen, S. Y., Liu, Q. Q., & Xiang, C. B. Arabidopsis MADS-box factor AGL16 negatively regulates drought resistance via stomatal density and stomatal movement. *J. Exp. Bot*. **71**(19), 6092–6106 (2020).

50. Sung, S. & Amasino, R. M. Vernalization in *Arabidopsis thaliana* is mediated by the PHD finger protein VIN3. *Nature* **427**, 159–164 (2004).

51. Franco-Zorrilla, J. M. *et al.* AtREM1, a member of a new family of B3 domain-containing genes, is preferentially expressed in reproductive meristems. *Plant Physiol.* **128**, 418–427 (2002).

52. Levy, Y. Y., Mesnage, S., Mylne, J. S., Gendall, A. R. & Dean, C. Multiple roles of Arabidopsis VRN1 in vernalization and flowering time control. *Science.* **297**, 243–246 (2002).

53. Welner D.H., Deeba F., Lo Leggio L., Skriver K. Chapter 13—NAC Transcription Factors: From Structure to Function in Stress-Associated Networks. In: Gonzalez D.H., editor. *Plant Transcription Factors.* pp. 199–212. Academic Press; Boston, MA, USA. (2016).

54. Shao, H. B., Song, W. Y., & Chu, L. Y. Advances of calcium signals involved in plant anti-drought. *C. R. Biol.* **331** (8), 587–596. (2008).

55. Oliva, M. *et al*. An environmentally responsive transcriptional state modulates cell identities during root development. *bioRxiv* doi: https://doi.org/10.1101/2022.03.04.483008 (2022).

56. Cai, S. *et al.* Evolutionary Conservation of ABA Signaling for Stomatal Closure. *Plant Physiol.* **174**(2), 732-747. (2017).

57. Melotto, M. *et al*. Plant stomata function in innate immunity against bacterial invasion. *Cell.* **126**(5), 969-980. (2006).

58. Wang, W. *et al.* Timing of plant immune responses by a central circadian regulator. *Nature* **470**, 110–115 (2011).

59. Zhou, M. *et al.* Redox rhythm reinforces the circadian clock to gate immune response. *Nature* **523**, 472–476 (2015).

60. Palmero, D. *et al.* Fungal microbiota from rain water and pathogenicity of Fusarium species isolated from atmospheric dust and rainfall dust. *J. Ind. Microbiol. Biotechnol.* **38**, 13–20 (2011).

61. Schwartz, H. F., Otto, K. L. & Gent, D. H. Relation of temperature and rainfall to development of Xanthomonas and Pantoea leaf blights of onion in Colorado. *Plant Dis.* **87**, 11–14 (2003).

62. Dalal, A., Attia, Z. & Moshelion, M. To produce or to survive: How plastic is your crop stress physiology? *Front. Plant Sci.* **8**, 1–8 (2017).

 Figures

Figure 1. A fine-scale RNA-seq time course of drought recovery reveals recovery-specific genes.
a, Illustration of the recovery time-course experimental design. Plants were grown in vermiculite under a
short-day photoperiod. After 30 days, irrigation was stopped for drought-treated plants until they reached
30% relative soil water content (SWC). We then rehydrated the drought-treated plants by saturating the
vermiculite to initiate drought recovery. We collected 3 biological samples from each time point of
recovery, and well-watered control at each time point (n=3) **b,** Illustration of the samples collected for

RNA-seq. We collected samples during moderate drought (t=0 at 30% SWC) and at seven additional
time-points during the recovery process, from 15 min to six hours after rehydration. All samples from
drought-treated plants were collected alongside equivalent samples from well-watered controls, with three
replicates per treatment (drought or control) per time point. **c,** Number of differentially expressed genes
(DEGs) at different time points during recovery. For each time point, we show the total number of DEGs
as well as the numbers of drought-responsive and recovery-specific genes. DEGs were identified by
comparing drought-treated samples to well-watered controls at each time point. **d-e,** K-means clustering
and expression patterns of **d,** drought-responsive and **e,** recovery-specific genes.

Figure 2. Drought recovery is initiated via transcriptional reprogramming in proliferating cells.
a, Illustration of the experimental design used to generate a single-nucleus gene expression map. We collected
two biological samples (n=2) from each of four conditions: (1) well-watered plants prior to drought [W]; (2)
plants experiencing moderate drought [D]; (3) drought-treated plants after 15 min of rehydration [R15]; and (4)
well-watered plants after 15 min of additional irrigation [W15]. **b,** Uniform Manifold Approximation and
Projection (UMAP) of the integrated single-nucleus RNA-seq dataset revealed 27 cell identities encompassing
the major leaf cell types: epidermal 1-3 (Epi), guard cell 1-2 (GC), trichome (Tr), hydathode (Hy), myrosin (M),
mesophyll 1-7 (Mes), phloem parenchyma (Pp), phloem companion (Pc), sieve element (Se), and dividing 1-2

(Div). Five clusters could not be assigned to a cell type (Unknown 1-5). **c,** Cell identities recovered from each of
the four conditions, expressed as a percent of the total number of recovered cells. Plants experiencing drought
maintain similar cell identities, but the initiation of recovery leads to enrichment of one of the unidentified
clusters (Unknown 1). **d,** Gene expression of Arabidopsis TFs during drought and immediate recovery
represented by Z-score. In green square, cluster 1 genes, representing TFs globally induced by rehydration in
most cell types. In red square, cluster 10 genes that were strongly induced by rehydration, specifically in
dividing cells.

Figure 3. Recovery cell states (RcS) appear in sub-populations of epidermal, mesophyll, hydathode, and vascular cells. a, Subclustering of cells within each cell type from Fig. 2C revealed unique subclusters enriched in cells from plants recovering from drought. Drought-recovered cells are indicated in red. **b,** Putative regulators of the RcS, identified using *de novo* motif enrichment analysis performed for each recovery-enriched subcluster. The gene name above each motif is the predicted TF that binds the enriched motif. **c,** Z-score representation showing the expression levels of the TFs that putatively regulate the formation of the post-drought RcS. **d,** Uniform Manifold Projection and Approximation (UMAP) of the snRNA-seq data generated from *Arabidopsis* rosettes over the course of a five-day dehydration experiment. We collected multiple rosettes for each day and processed them as one biological replicate (n=1). **e,** Z-score representation showing the expression levels of predicted CAMTA TFs during the early stages of plant dehydration.

Figure 4. Recovery from moderate drought enhanced pathogen resistance in *Arabidopsis*.
a, Overlap between upregulated recovery-specific genes in our bulk RNA-seq data and genes upregulated by different biotic elicitors as reported by Bjornson et al. (2021). For each recovery time point, bars show both the percentage of upregulated recovery-specific genes that are also known to be activated by biotic elicitors as well as the total number. **b,** Bacterial load ($\log_{10}$CFU) of plants that were grown on sterile low-water-content plates and then submerged in a suspension of *Pseduomonas syringae* pv. *tomato* DC3000 (*Pst* DC3000, OD_{600}=0.005) or a control solution. Bacterial growth was measured two days post-inoculation, and well-watered controls (W) were compared against plants recovering from either moderate (R-MD) or severe drought (R-SD). For this analysis, we merged two independent experiments for a total of n=12 per treatment. Significance values were calculated with a two-way ANOVA of treatment and batch followed by Tukey's *post hoc* test. P-values were FDR-corrected. **c,** Representative images of plants taken four days post-inoculation. **d,** Bacterial load ($\log_{10}$CFU)

of soil-grown plants sprayed with *Pst* DC3000 (OD_{600}=0.05) or a control solution after experiencing moderate
drought. Bacterial growth was measured 4 and 24 hours post-inoculation and compared against controls using a
two-way student's t-test. We collected 16 leaves (n=16) to measure bacterial load at the initial time point, and 24
leaves (n=24) to measure bacterial growth 24 hours post infection. **e,** Representative images of plants taken 14
days post-inoculation. In **b** and **d**: ns = P > 0.05, * = P ≤ 0.05, ** = P ≤ 0.01, *** = P ≤ 0.001, **** = P ≤
0.0001. Boxplots middle line shows the median, the lower and upper hinges are the 25th and 75th percentile,
respectively. The whiskers extend from the hinges to the most distant value within 1.5 * IQR of the hinge, where
IQR is the inter-quartile range, or distance between the first and third quartiles.

Figure 5. Drought recovery-induced immunity enhanced resistance to *X. perforans* and *Pst* DC3000 in wild (*Solanum pennellii*) and domesticated (*S. lycopersicum* cv. M82) tomato species. a, Lesion percentage analysis of *S. pennellii* leaves five days after being syringe infiltrated with either *X. perforans* OD_{600}= 0.02 or *Pst* DC3000 OD_{600}= 0.02. Inoculation was performed after 90 mins of recovery from moderate drought, and drought-treated samples were compared to well-watered controls. Results are shown for three independent experiments, 9 leaves were analyzed in each experiment (n=9). **b,** Representative images of well-watered or drought-recovered *S. pennellii* leaves infected with one of *Pst* DC3000, *X. perforans*, or a mock solution. Scale bar = 1 cm. **c,** Bacterial load (log_{10}CFU) of the *S. pennellii* leaves measured five days post-infection. Results are shown for three independent experiments. 2 leaf discs (0.5 cm in diameter) were prepared for each sample, 5 samples were used for each independent experiment (n=5). **d,** Lesion percentage analysis of M82 leaves after undergoing the same post-drought infection experiment described for *S. pennelli*. 9 leaves were analyzed in each

experiment (n=9). **e,** Representative images of M82 leaves infected with *Pst* DC3000, *X. perforans*, or a mock
solution. Scale bar = 2 cm. **f,** Bacterial load ($\log_{10}$CFU) of M82 leaves measured five days post-infection.
Results are shown for three independent experiments. 2 leaf discs (0.5 cm in diameter) were prepared for each
sample, 5 samples were used for each independent experiment (n=5). Significant differences in all panels were
identified using student's t-test (ns = $P > 0.05$, * = $P \leq 0.05$, ** = $P \leq 0.01$, *** = $P \leq 0.001$, **** = $P \leq 0.0001$).
Boxplots middle line shows the median, the lower and upper hinges are the 25th and 75th percentile,
respectively. The whiskers extend from the hinges to the most distant value within 1.5 * IQR of the hinge, where
IQR is the inter-quartile range, or distance between the first and third quartiles.

Methods

Plant materials and growth conditions

Arabidopsis thaliana Columbia-0 (Col-0) background was used throughout this study. Col-0 seeds were

sterilized with chlorine fumes generated by mixing 100 ml bleach and 4 ml hydrochloric acid (HCl 1M). Sterilized

seeds were sown on large square petri dishes with ¾ Linsmaier & Skoog with buffer (LS) media per liter and 2%

agar, and then stratified at 4 °C for three days. Plates were placed in a growth chamber and grown in short-day

conditions (8h light: 16h dark) at 22 °C with a light intensity of ~110–130 µmol m^{-2} s^{-1}. For drought treatments,

seedlings were transferred to vermiculite pots two weeks after germination on plates, with 6 seedlings per pot.

Tray size was 27.79 cm width, 54.46 cm in length, and 6.2 cm depth. Vermiculite was saturated with ¾ LS liquid

media before the seedling transfer, 2L media per tray. After two days, each tray was watered with 2L ultra-pure

water, and this continued for two weeks until the drought treatment started. When plants were 30 days old, each

pot was weighed and transferred to a dry tray. The pots in the tray were weighed daily, and relative water content

was calculated. The experiment started when the pots reached 30% relative soil-water content (SWC). Three whole

Arabidopsis rosettes per treatment were collected separately as an independent sample, time 0 was 9 am (one hour

after the lights turn on in the chambers). Three samples per treatment were collected alongside a well-watered

control at each time point. For sterile low-water agar experiments, plates were prepared modified from Gonzales

et al., 2023; 100% - 4L DDW, 3 LS bags, 80g agar – 120ml per plate, 50% (moderate stress) - 2L DDW, 3 LS

bags, 80g agar – 60ml per plate, and 25% (severe stress) - 1L DDW, 3 LS bags, 80g agar – 30ml per plate.

For the drought time course experiment, plants were grown on plates and transferred to vermiculite as

described above with the notable exception that transfer from plates took place 17 days after germination and

remained on saturated vermiculite for 12 days before drought initiation. Drought was initiated by draining excess

media from the pot and equilibrating to 100% relative SWC. Whole rosettes (n= 3 – 6) were sampled and frozen

4.5 hours after subjective dawn each day for 5 days.

Col-0 seeds were sterilized and seeded on plates. Plates were kept in the dark at 4C for three days. After

three days, plates were moved to a growth chamber with short-day photoperiod light and 22 °C temperature.

Tomato seeds of *Solanum pennellii* and *Solanum lycopersicum* cv. M82 were germinated in a petri dish

with 1/2 Murashige & Skoog (MS) with vitamins and FeNaEDTA (Cat# 07190008) media (no sucrose added).

Ten-days-old seedlings were transferred to tray pots containing a mixture of vermiculite (Agrekal Moshav

Habonim) and commercial soil (Tuff A.C.S) (1:1 by volume) and grown in a 12-hour light-dark cycle (12h light:

12h dark) at 25°C.

Bacterial inoculation

Pseudomonas syringae pv. *tomato (Pst)* DC3000 was grown overnight in 10 ml liquid King's B medium

containing rifampicin (40 µg/ml) (Cat#: R3501-250MG, MilliporeSigma, MA) and tetracyclin (10 mg/ml) (Cat#:

T3258, MilliporeSigma, MA) at 28°C for 24 h. Before adjusting the density, bacterial cells were washed two times

with autoclaved water followed by centrifugation at 6,000 rpm for 2 min and re-suspension in water. For bacterial

growth assays, well-watered and 90 min recovered Arabidopsis plants were inoculated with *Pst* DC3000 (OD_{600}

= 0.005-0.5, depending on the experiment and inoculation method). For spray treatment Silwett77 (Cat#: S7777,

PhytoTachLabs, KS) was added to bacteria before spraying. Bacterial titer was assessed as the log_{10} transformed

colony forming units (CFU) per plant weight when collecting whole plants from sterile plates.

For experiments performed on tomato, syringe infiltration was used for bacterial inoculation. For

bacterium counting, ten leaf discs (0.5 cm in diameter) were prepared from the second and third true leaves using

a hole puncher (two discs per leaf) for each sample, five samples were used for each independent experiment. Each

sample was separately homogenized in 10 mM $MgCl_2$. The homogenate was serially diluted in 10 mM $MgCl_2$.

10µl of diluted homogenate were plated on antibiotics and LB agar media on a square petri dish and incubated at

28°C for 48 h for determination of bacterial concentrations in leaves ($log^{10} CFU/cm^2$).

Assessment of disease severity in in tomato leaves

Disease severity in tomato (*S. pennellii* and *S. lycopersicum* cv. M82) leaves was assessed by calculating

the percentage of the symptomatic area in the leaves relative to the whole leaf area. The calculation was done

based on threshold color and color space HSB in ImageJ version 1.54d, according to Tuang et al.[1] Nine leaves per

experiment were analyzed for the lesion percentage (%).

RNA extraction, bulk RNA library construction

Total RNA was extracted from three independent biological replicates of each time point using RNeasy Plant Mini Kits (Cat#79254, Qiagen, CA). Tape Station checked RNA quantity for quality control. Library construction was performed using Illumina Stranded mRNA Prep (Cat#20040534, Illumina, CA).

Nuclei extraction and single-nuclei library construction

Seedlings were transferred to vermiculite pots two weeks after germination in trays, 6 seedlings per pot. Vermiculite was saturated with ¾ LS liquid media before the seedling transfer, 2L media per tray. After two days each tray was watered with 2L ultra-pure water, and this continued for two weeks until the drought treatment started. When plants were 30 days old, each pot was weighed and transferred to a dry tray. The experiment began when each pot reached 30% SWC. Between 12-18 whole rosettes were collected for each time point * condition sample. We used mortar and pestle to grind the frozen tissue. Powdered tissue was then placed in a nuclei extraction buffer [NEB; 500ul 1M TRIS pH=7.4 (Cat# 15567027, Thermo Fisher Scientific, MA), 150µl 1M MgCl$_2$ (Cat# AM9530G, Fisher Scientific, MA), 100µl 1M NaCl (Cat# AM9760G Fisher Scientific, MA), 50ml nuclease-free water (Cat# AM9937 Thermo Fisher Scientific, MA), 25µl 1M spermine (Cat# 85590-5G, MilliporeSigma, MA), 10µl 1M spermidine (Cat# S2626-5G, MilliporeSigma, MA), 500µl proteinase inhibitor (PI; Cat# P9599-5ML MilliporeSigma, MA), 250µl BSA (Cat# B2518-100MG, MilliporeSigma, MA), 250µl SUPERase-In (Cat# AM2696, Thermo Fisher Scientific, MA)] and incubated for 10 min. After incubation, tissue was filtered through a 40µm filter (Cat# 43-57040-51, pluriSelect, Germany). We then centrifuged at 500 rcf for 5 min at 4°C. Supernatant was aspirated out, and nuclei were resuspended with NEB + 500µl 10% Triton (Cat# 93443-100ML, MilliporeSigma, MA) and no PI. We incubated for 15 min, filtered through a 40µm filter, and spun at 500 rcf for 5 min. We washed until the pellet was clear. We prepared the density gradient using the Density Buffer [DB; 120 mM Tris-Cl pH=8 (Cat# AM9855G, Fisher Scientific, MA), 150 mM KCl (Cat# AM9640G, Fisher Scientific, MA), 30 mM MgCl2, 35mL H$_2$0 per 50mL] and filter sterilized it. We mixed 5 volumes of Optiprep (Cat# D1556-250ML, MilliporeSigma, MA) and the buffer volume to create a 50% stock. We also made a dilutant stock [400 mM Sucrose, 25 mM KCl, 5 mM MgCl$_2$, 10 mM Tris-Cl pH 8, 28mL H$_2$0 per 50mL] that we filter sterilized. A 45% solution was made by mixing 9ml 50% solution and 1ml dilutant, and a 15% solution by combining 1.5ml

50% stock and 3.5ml dilutant. We created a 45% solution and a 15% solution. We gently poured 2ml of the nuclei

solution at the top of the density gradient and then spun the tubes at 1,500 rcf for 5 min with no breaks. After the

spin, we pipette off nuclei and place them into a 15 ml tube with ~ 6mL of NEB (with no triton or PI). We note

that for vermiculite drought stressed nuclei, nuclei preparations were FACS purified. After counting the nuclei,

the nuclei suspension was loaded onto microfluidic chips (10X Genomics) with HT-v3.1 chemistry to capture

~20,000 nuclei/sample. Cells were barcoded with a Chromium X Controller (10X Genomics). mRNA was reverse

transcribed, and Illumina libraries were constructed for sequencing with reagents from a 3' Gene Expression HT-

v3.1 kit (10X Genomics) according to the manufacturer's instructions. cDNA and final library quality were

assessed using Tape-Station High Sensitivity DNA Chip (Agilent, CA).

Each of the single-nuclei samples was processed twice, to get a higher number of single nuclei

transcriptomes.

Bulk RNA-sequencing and analysis

Sequencing was performed with a NovaSeq 6000 instrument (Illumina, CA). About 40 million reads were

obtained for each sample. Raw reads were processed at the IGC bioinformatics core at Salk. Alignments were

performed using OSA4 and mapped to the Arabidopsis genome (TAIR 10) using Tophat2 software with default

settings. Mapped reads per library were counted using HTSeq software. Differentially expressed genes were

quantified in two ways. Firstly, differentially expressed genes were identified using a spline regression model in

splineTimeR v1.18.0, which were then sorted into time points using k-means clustering. Differential expression

was also quantified at each time point individually using DESeq2 v1.30.0.[2] For each pairwise comparison, genes

with fewer than 32 total raw counts across all samples were discarded before normalization. Genes with an absolute

$\log_2$foldchange > 1 and an FDR-corrected p-value ≤ 0.01 were pulled as significant. For functional enrichment,

genes were queried for time-specific functional enrichment using over-representation analysis (ORA) in

WebGestaltR v0.4.4.[3] Differentially expressed genes in each pairwise comparison were queried against the

biological process non-redundant ontology, and a significance threshold of FDR-corrected p-value ≤ 0.05 was

used.

snRNA-seq analysis

For the snRNA-seq libraries, CellRanger (v.6.0.1) was used to perform sample-demultiplexing, barcode

processing, and single-nuclei gene-UMI counting.[4] Each experiment's expression matrix was obtained by aligning

to Arabidopsis transcriptome reference (TAIR 10) using CellRanger with default parameters. For initial quality-

control filtering, aligned cell and transcript counts from each treatment (well-watered, drought, recovery, 2

replicates each) were processed by Seurat (Version 4.2).[5] The data was filtered in the following two ways: (1) Pre-

filtering each replicate by removing the low-quality and outlier cells containing a high abundance of chloroplast

reads (>40% of total transcripts) and mitochondrial reads (>1% of total transcripts), a low abundance of detected

genes (<300 detected genes) and a relatively high abundance of unique molecular identifiers (UMIs) (>10K for

D0; >15K and >10K for two W15, respectively; >25K and >20K for two W0 , respectively; >10K for R15)

(Extended Data Table 3). (2) Identifying possible doublet in each replicate using the method SCDS. SCDS

implements two complementary approaches to identify doublets: one is co-expression-based doublet scoring, and

the other is binary classification-based doublet scoring. Additionally, they provide a hybrid score by combing these

two approaches. SCDS showed relatively high detection accuracy and computational efficiency when

benchmarking with other computational methods.[6] We applied the hybrid scores for doublet estimation using R

package scds() to identify likely doublet and then removed them from downstream integration. Expression data of

cells passing these thresholds were log normalized with NormalizeData() function, and the top2K variable genes

were identified with the FindVariableFeatures() function. Next, data from all conditions were integrated using

Seurat's reciprocal PCA (RPCA) and FindIntegrationAnchors() functions to identify integration features and

correct for potential batch effects. The integrated data were then scaled with ScaleData() function. Principal

component analysis (PCA) was carried out with RunPCA() function, and the top 30 principal components (PCs)

were retained. Clusters were identified with the FindClusters() function using the shared nearest neighbor

modularity optimization with a clustering resolution set to 0.8. Clusters with only one cell were removed. This

resulted in 27 initial clusters with a total of 144,494 cells. We identified a median number of 1,500 genes, and

3,146 UMIs (representing unique transcripts), per nuclei. We detected between 25,064 – 26,342 genes in each

sample. Cell-type identity of initial clusters was determined with canonical markers and by referring to public

datasets, followed by sub-clustering each cluster using the same strategy described above.

766   Single cell differential gene expression analysis was conducted using Seurat FindMarker() and

FindAllmarkers() functions with the default Wilcoxon Rank Sum test. Marker genes per cell cluster and

differentially expressed genes (DEGs) among conditions were both identified by setting parameter min.pct as 0.1

and picked the top expressed markers ranking by average log_2FC. Functional enrichment analysis was carried out

by over-representation analysis (ORA) using the top100 markers ranked by average log_2FC. Using the GO

biological process as a reference, ORA was performed with WebGestalt.[3] Pathways with FDR <0.05 were

considered significantly enriched and visualized as plots with normalized enrichment scores. For recovery-

/drought-specific DEGs, enriched known motifs were discovered with HOMER findmotifs.pl

(http://homer.ucsd.edu/homer/), searching within 1000bp upstream to 1000kb downstream of their transcript start

sites (TSSs). For the sub-clusters, recovery-enriched cell clusters were identified with >=50% cells from recovery

experiment, followed with weighted gene co-expression network analysis (WGCNA) with R package hdWGCNA

(https://smorabit.github.io/hdWGCNA/). Co-expression networks were constructed for each initial cell cluster

with top 20 hub genes. By checking the average expression of hub genes in sub-clusters, the recovery-specific

networks were picked as those with higher expression in recovery-enriched cell clusters.

Statistical analysis

bulk RNA-seq data, statistical analysis was performed in R using a mixed linear model function (lmer)

from the package lme4 unless otherwise described. Standard errors were calculated from variance and covariance

values after model fitting. The Benjamini-Hochberg method was applied for correcting of multiple testing in

figures showing all pairwise comparisons of the mean estimates. For bacterial colonies count, merged two

independent experiments of plants grown on plates, a total of n=12 per treatment. Significance values

for log10(CFU) on the plates were calculated with a two-way ANOVA of treatment and batch followed by a Tukey

test. P-values are FDR-corrected. For bacterial colonies count, soil grown plants experiment, significance values

for log10(CFU) were calculated with a two-way student's t-test.

Data availability

Data is available here: GSE220278 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE220278).

Currently open only for reviewers.

Code availability

The code used to analyze both bulk and single-nuclei RNA-Seq data is available at:

https://github.com/NatanellaIE/DroughtRecovery.

Methods references

1. Tuang, Z. K. *et al.* Pst DC3000 infection alleviates subsequent freezing and heat injury to host plants via
a salicylic acid-dependent pathway in Arabidopsis. *Plant Cell Environ.* **43**, 801–817 (2020).
2. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).
3. Liao, Y., Wang, J., Jaehnig, E. J., Shi, Z. & Zhang, B. WebGestalt 2019: gene set analysis toolkit with
revamped UIs and APIs. *Nucleic Acids Res.* **47**, 199–205 (2019).
4. Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nat. Commun.* **8**,
(2017).
5. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29 (2021).
6. Xi, N. M. & Li, J. J. Benchmarking Computational Doublet-Detection Methods for Single-Cell RNA
Sequencing Data. *Cell Syst.* **12**, 176-194 (2021).

Extended data references

1. Abe, M., Takahashi, T. & Komeda, Y. Cloning and characterization of an L1 layer-specific gene in
Arabidopsis thaliana. *Plant Cell Physiol.* **40**, 571–580 (1999).
2. Abe, M., Takahashi, T. & Komeda, Y. Identification of a cis-regulatory element for L1 layer-specific
gene expression, which is targeted by an L1-specific homeodomain protein. *Plant J.* **26**, 487–494 (2001).
3. Alvarez-Buylla, E. R. *et al.* MADS-box gene evolution beyond flowers: Expression in pollen, endosperm,
guard cells, roots and trichomes. *Plant J.* **24**, 457–466 (2000).
4. Baima, S. *et al.* The expression of the Athb-8 homeobox gene is restricted to provascular cells in
Arabidopsis thaliana. *Development* **121**, 4171–4182 (1995).
5. Kang, J. & Dengler, N. Vein pattern development in adult leaves of Arabidopsis thaliana. *Int. J. Plant
Sci.* **165**, 231–242 (2004).
6. Scarpella, E., Francis, P. & Berleth, T. Stage-specific markers define early steps of procambium
development in Arabidopsis leaves and correlate termination of vein formation with mesophyll
differentiation. *Development* **131**, 3445–3455 (2004).
7. Kim, J. Y. *et al.* Distinct identities of leaf phloem cells revealed by single cell transcriptomics. *Plant Cell*
33, 511–530 (2021).
8. Barth, C. & Jander, G. Arabidopsis myrosinases TGG1 and TGG2 have redundant function in
glucosinolate breakdown and insect defense. *Plant J.* **46**, 549–562 (2006).
9. Bonke, M., Thitamadee, S., Mähönen, A. P., Hauser, M. T. & Helariutta, Y. APL regulates vascular
tissue identity in Arabidopsis. *Nature* **426**, 181–186 (2003).
10. Truernit, E. & Sauer, N. The promoter of the Arabidopsis thaliana SUC2 sucrose-H+ symporter gene

directs expression of β-glucuronidase to the phloem: Evidence for phloem loading and unloading by
SUC2. *Planta An Int. J. Plant Biol.* **196**, 564–570 (1995).

11. Bulankova, P., Akimcheva, S., Fellner, N. & Riha, K. Identification of Arabidopsis meiotic cyclins
reveals functional diversification among plant cyclin genes. *PLoS Genet.* **9**(5), e1003508 (2013).

12. Chauvin, A., Caldelari, D., Wolfender, J. L. & Farmer, E. E. Four 13-lipoxygenases contribute to rapid
jasmonate synthesis in wounded Arabidopsis thaliana leaves: A role for lipoxygenase 6 in responses to
long-distance wound signals. *New Phytol.* **197**, 566–575 (2013).

13. Nguyen, C. T., Kurenda, A., Stolz, S., Chételat, A. & Farmer, E. E. Identification of cell populations
necessary for leaf-to-leaf electrical signaling in a wounded plant. *Proc. Natl. Acad. Sci. U. S. A.* **115**,
10178–10183 (2018).

14. Chen, L. Q. *et al.* Sucrose efflux mediated by SWEET proteins as a key step for phloem transport.
Science. **335**, 207–211 (2012).

15. Cui, H., Kong, D., Liu, X. & Hao, Y. SCARECROW, SCR-LIKE 23 and SHORT-ROOT control bundle
sheath cell fate and function in Arabidopsis thaliana. *Plant J.* **78**, 319–327 (2014).

16. De Rybel, B. *et al.* A bHLH complex controls embryonic vascular tissue establishment and indeterminate
growth in Arabidopsis. *Dev. Cell* **24**, 426–437 (2013).

17. Endo, M., Shimizu, H., Nohales, M. A., Araki, T. & Kay, S. A. Tissue-specific clocks in Arabidopsis
show asymmetric coupling. *Nature* **515**, 419–422 (2014).

18. Fisher, K. & Turner, S. PXY, a receptor-like kinase essential for maintaining polarity during plant
vascular-tissue development. *Curr. Biol.* **17**, 1061–1066 (2007).

19. Funk, V., Kositsup, B., Zhao, C. & Beers, E. P. The Arabidopsis xylem peptidase XCP1 is a tracheary
element vacuolar protein that may be a papain ortholog. *Plant Physiol.* **128**, 84–94 (2002).

20. Glaring, M. A. *et al.* An extra-plastidial α-glucan, water dikinase from Arabidopsis phosphorylates
amylopectin in vitro and is not necessary for transient starch degradation. *J. Exp. Bot.* **58**, 3949–3960
(2007).

21. Gotor, C., Cejudo, F. J., Barroso, C. & Vega, J. M. Tissue-specific expression of ATCYS-3A, a gene
encoding the cytosolic isoform of O-acetylserine(thiol)lyase in Arabidopsis. *Plant J.* **11**, 347–352 (1997).

22. Guo, W. J., Bundithya, W. & Goldsbrough, P. B. Characterization of the Arabidopsis metallothionein
gene family: Tissue-specific expression and induction during senescence and in response to copper. *New*
Phytol. **159**, 369–381 (2003).

23. Hsu, P. K. & Tsay, Y. F. Two phloem nitrate transporters, NRT1.11 and NRT1.12, are important for
redistributing xylem-borne nitrate to enhance plant growth. *Plant Physiol.* **163**, 844–856 (2013).

24. Kondo, T. *et al.* Stomatal density is controlled by a mesophyll-derived signaling molecule. *Plant Cell*
Physiol. **51**, 1–8 (2010).

25. Sugano, S. S. *et al.* Stomagen positively regulates stomatal density in Arabidopsis. *Nature* **463**, 241–244
(2010).

26. Lopez-Anido, C. B. *et al.* Single-cell resolution of lineage trajectories in the Arabidopsis stomatal lineage
and developing leaf. *Dev. Cell* **56**, 1043-1055 (2021).

27. Johnson, C. S., Kolevski, B. & Smyth, D. R. Transparent Testa Glabra2, a trichome and seed coat
development gene of arabidopsis, encodes a WRKY transcription factor. *Plant Cell* **14**, 1359–1375
(2002).

28. Kang, B. H., Busse, J. S. & Bednarek, S. Y. Members of the arabidopsis dynamin-like gene family,
ADL1, are essential for plant cytokinesis and polarized cell growth. *Plant Cell* **15**, 899–913 (2003).

29. Kasahara, R. D., Portereiko, M. F., Sandaklie-Nikolova, L., Rabiger, D. S. & Drews, G. N. MYB98 is
required for pollen tube guidance and synergid cell differentiation in Arabidopsis. *Plant Cell* **17**, 2981–
2992 (2005).

30. Wenzel, C. L., Schuetz, M., Yu, Q. & Mattsson, J. Dynamics of MONOPTEROS and PIN-FORMED1
expression during leaf vein pattern formation in Arabidopsis thaliana. *Plant J.* **49**, 387–398 (2007).

31. Jammes, F. *et al.* MAP kinases MPK9 and MPK12 are preferentially expressed in guard cells and
positively regulate ROS-mediated ABA signaling. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 20520–20525
(2009).

32. Ji, J. *et al.* Wox4 promotes procambial development. *Plant Physiol.* **152**, 1346–1356 (2010).

33. Zhang, T. Q., Chen, Y. & Wang, J. W. A single-cell analysis of the Arabidopsis vegetative shoot apex.

Dev. Cell **56**, 1056-1074.e8 (2021).

34.   Procko, C. *et al.* Leaf cell-specific and single-cell transcriptional profiling reveals a role for the palisade
layer in UV light protection. *Plant Cell* **34**, 3261–3279 (2022).

35.   Kirschner, S. *et al.* Expression of SULTR2;2, encoding a low-affinity sulphur transporter, in the
Arabidopsis bundle sheath and vein cells is mediated by a positive regulator. *J. Exp. Bot.* **69**, 4897–4906
(2018).

36.   Kurata, T. *et al.* The YORE-YORE gene regulates multiple aspects of epidermal cell differentiation in
Arabidopsis. *Plant J.* **36**, 55–66 (2003).

37.   Mathur, J. *et al.* Transcription of the Arabidopsis CPD gene, encoding a steroidogenic cytochrome P450,
is negatively controlled by brassinosteroids. *Plant J.* **14**, 593–602 (1998).

38.   Müller, R., Borghi, L., Kwiatkowska, D., Laufs, P. & Simon, R. Dynamic and compensatory responses of
Arabidopsis shoot and floral to meristems to CLV3 signaling. *Plant Cell* **18**, 1188–1198 (2006).

39.   Chen, L. Q. *et al.* Sugar transporters for intercellular exchange and nutrition of pathogens. *Nature* **468**,
527–532 (2010).

40.   Mustroph, A. *et al.* Profiling translatomes of discrete cell populations resolves altered cellular priorities
during hypoxia in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 18843–18848 (2009).

41.   Zhang, L. *et al.* Altered xylem-phloem transfer of amino acids affects metabolism and leads to increased
seed yield and oil content in Arabidopsis. *Plant Cell* **22**, 3603–3620 (2010).

42.   Ohashi-Ito, K. & Bergmann, D. C. Arabidopsis FAMA controls the final proliferation/differentiation
switch during stomatal development. *Plant Cell* **18**, 2493–2505 (2006).

43.   Okumoto, S. *et al.* High affinity amino acid transporters specifically expressed in xylem parenchyma and
developing seeds of Arabidopsis. *J. Biol. Chem.* **277**, 45338–45346 (2002).

44.   Pilot, G. *et al.* Overexpression of GLUTAMINE DUMPER1 leads to hypersecretion of glutamine from
hydathodes of arabidopsis leaves. *Plant Cell* **16**, 1827–1840 (2004).

45.   Pommerrenig, B. *et al.* Phloem-Specific expression of Yang cycle genes and identification of novel Yang
cycle enzymes in Plantago and Arabidopsis. *Plant Cell* **23**, 1904–1919 (2011).

46.   Qiu, J. L., Jilk, R., Marks, M. D. & Szymanski, D. B. The Arabidopsis SPIKE1 gene is required for
normal cell shape control and tissue development. *Plant Cell* **14**, 101–118 (2002).

47.   Susek, R. E., Ausubel, F. M. & Chory, J. Signal transduction mutants of arabidopsis uncouple nuclear
CAB and RBCS gene expression from chloroplast development. *Cell* **74**, 787–799 (1993).

48.   Ranjan, A., Fiene, G., Fackendahl, P. & Hoecker, U. The Arabidopsis repressor of light signaling SPA1
acts in the phloem to regulate seedling de-etiolation, leaf expansion and flowering time. *Development*
138, 1851–1862 (2011).

49.   Stadler, R. & Sauer, N. The Arabidopsis thaliana AtSUC2 gene is specifically expressed in companion
cells. *Bot. Acta* **109**, 299–306 (1996).

50.   Rouse, D. T., Marotta, R. & Parish, R. W. Promoter and expression studies on an Arabidopsis thaliana
dehydrin gene. *FEBS Lett.* **381**, 252–256 (1996).

51.   Zimmermann, I., Saedler, R., Mutondo, M. & Hulskamp, M. The Arabidopsis GNARLED gene encodes
the NAP125 homolog and controls several actin-based cell shape changes. *Mol. Genet. Genomics* **272**,
290–296 (2004).

52.   Zhu, Y., Liu, L., Shen, L. & Yu, H. NaKR1 regulates long-distance movement of FLOWERING LOCUS
T in Arabidopsis. *Nat. Plants* **2**, (2016).

53.   Zhang, X., Dyachok, J., Krishnakumar, S., Smith, L. G. & Oppenheimer, D. G. IRREGULAR
TRICHOME BRANCH1 in arabidopsis encodes a plant homolog of the actin-related protein2/3 complex
activator Scar/WAVE that regulates actin and microtubule organization. *Plant Cell* **17**, 2314–2326
(2005).

54.   Yoshimoto, N., Inoue, E., Saito, K., Yamaya, T. & Takahashi, H. Phloem-localizing sulfate transporter,
Sultr1;3, mediates re-distribution of sulfur from source to sink organs in arabidopsis. *Plant Physiol.* **131**,
1511–1517 (2003).

55.   Yephremov, A., *et al*. Characterization of the FIDDLEHEAD gene of Arabidopsis reveals a link between
adhesion response and cell differentiation in the epidermis. *Plant cell*, **11**(11), 2187–2201. (1999).

56.   Xu, W. *et al.* Arabidopsis TCH4, regulated by hormones and the environment, encodes a xyloglucan
endotransglycosylase. *Plant Cell* **7**, 1555–1567 (1995).

57. Wu, H. *et al.* Molecular and biochemical characterization of the Fe(III) chelate reductase gene family in
Arabidopsis thaliana. *Plant Cell Physiol.* **46**, 1505–1514 (2005).
58. Werner, T. *et al.* Cytokinin-deficient transgenic Arabidopsis plants show multiple developmental
alterations indicating opposite functions of cytokinins in the regulation of shoot and root meristem
activity. *Plant Cell* **15**(11), 2532–2550 (2003).
59. Uemoto, K., Araki, T. & Endo, M. Isolation of arabidopsis palisade and spongy mesophyll cells. *Methods*
Mol. Biol. **1830**, 141–148 (2018).
60. Takemiya, A. *et al.* Phosphorylation of BLUS1 kinase by phototropins is a primary step in stomatal
opening. *Nat. Commun.* **4**, 2094. (2013).
61. Takada, S., Takada, N. & Yoshida, A. ATML1 promotes epidermal cell differentiation in Arabidopsis
shoots. *Dev.* **140**, 1919–1923 (2013).
62. Szymanski, D. B., Jilk, R. A., Pollock, S. M. & Marks, M. D. Control of GL2 expression in Arabidopsis
leaves and trichomes. *Development* **125**, 1161–1171 (1998).
63. Zhang, F., Gonzalez, A., Zhao, M., Payne, C. T. & Lloyd, A. A network of redundant bHLH proteins
functions in all TTG1-dependent pathways of Arabidopsis. *Development* **130**, 4859–4869 (2003).
64. Xia, Y., Nikolau, B. J. & Schnable, P. S. Developmental and hormonal regulation of the Arabidopsis
CER2 gene that codes for a nuclear-localized protein required for the normal accumulation of cuticular
waxes. *Plant Physiol.* **115**, 925–937 (1997).
65. Wallner, E. S. *et al.* Strigolactone- and Karrikin-Independent SMXL Proteins Are Central Regulators of
Phloem Formation. *Curr. Biol.* **27**(8), 1241–1247 (2017).
66. Walker, A. R. *et al.* The TRANSPARENT TESTA GLABRA1 locus, which regulates trichome
differentiation and anthocyanin biosynthesis in arabidopsis, encodes a WD40 repeat protein. *Plant Cell*
11, 1337–1349 (1999).
67. Vanzin, G. F. *et al.* The mur2 mutant of Arabidopsis thaliana lacks fucosylated xyloglucan because of a
lesion in fucosyltransferase AtFUT1. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 3340–3345 (2002).
68. Van Leene, J. *et al.* Targeted interactomics reveals a complex core cell cycle machinery in Arabidopsis
thaliana. *Mol. Syst. Biol.* **6**, (2010).
69. Vahisalu, T. *et al.* SLAC1 is required for plant guard cell S-type anion channel. *Nature* **452**, 487–491
(2008).
70. Takahashi, K. *et al.* Ectopic expression of an esterase, which is a candidate for the unidentified plant
cutinase, causes cuticular defects in arabidopsis thaliana. *Plant Cell Physiol.* **51**, 123–131 (2010).
71. Ohashi, Y., Oka, A., Ruberti, I., Morelli, G. & Aoyama, T. Entopically additive expression of GLABRA2
alters the frequency and spacing of trichome initiation. *Plant J.* **29**, 359–369 (2002).
72. Swaminathan, K., Yang, Y., Grotz, N., Campisi, L. & Jack, T. An enhancer trap line associated with a D-
class cyclin gene in Arabidopsis. *Plant Physiol.* **124**, 1658–1667 (2000).
73. Sistrunk, M. L., Antosiewicz, D. M., Purugganan, M. M. & Braam, J. Arabidopsis TCH3 encodes a novel
Ca2+ binding protein and shows environmentally induced and tissue-specific regulation. *Plant Cell* **6**,
1553–1565 (1994).
74. Sinlapadech, T., Stout, J., Ruegger, M. O., Deak, M. & Chapple, C. The hyper-fluorescent trichome
phenotype of the brt1 mutant of Arabidopsis is the result of a defect in a sinapic acid:UDPG
glucosyltransferase. *Plant J.* **49**, 655–668 (2007).
75. Schlereth, A. *et al.* MONOPTEROS controls embryonic root initiation by regulating a mobile
transcription factor. *Nature* **464**, 913–916 (2010).
76. Schellmann, S. *et al.* TRIPTYCHON and CAPRICE mediate lateral inhibition during trichome and root
hair patterning in Arabidopsis. *EMBO J.* **21**, 5036–5046 (2002).
77. Sawchuk, M. G., Donner, T. J., Head, P. & Scarpella, E. Unique and overlapping expression patterns
among members of photosynthesis-associated nuclear gene families in Arabidopsis. *Plant Physiol.* **148**,
1908–1924 (2008).
78. Sawa, S. *et al.* The HAT2 gene, a member of the HD-Zip gene family, isolated as an auxin inducible gene
by DNA microarray screening, affects auxin response in Arabidopsis. *Plant J.* **32**, 1011–1022 (2002).
79. Saedler, R., Zimmermann, I., Mutondo, M. & Hülskamp, M. The Arabidopsis KLUNKER gene controls
cell shape changes and encodes the AtSRA1 homolog. *Plant Mol. Biol.* **56**, 775–782 (2004).
80. Endo, M., Mochizuki, N., Suzuki, T. & Nagatani, A. CRYPTOCHROME2 in vascular bundles regulates

flowering in Arabidopsis. *Plant Cell* **19**, 84–93 (2007).

81.   Redovniković, I. R., Textor, S., Lisnić, B. & Gershenzon, J. Expression pattern of the glucosinolate side
chain biosynthetic genes MAM1 and MAM3 of Arabidopsis thaliana in different organs and
developmental stages. *Plant Physiol. Biochem.* **53**, 77–83 (2012).

82.   Pilot, G., Gaymard, F., Mouline, K., Chérel, I. & Sentenac, H. Regulated expression of Arabidopsis
Shaker K+ channel genes involved in K+ uptake and distribution in the plant. *Plant Mol. Biol.* **51**, 773–
787 (2003).

83.   Oppenheimer, D. G. *et al.* Essential role of a kinesin-like protein in Arabidopsis trichome morphogenesis.
Proc. Natl. Acad. Sci. U. S. A. **94**, 6261–6266 (1997).

84.   Kirik, V. *et al.* Ectopic expression of the Arabidopsis AtMYB23 gene induces differentiation of trichome
cells. *Dev. Biol.* **235**, 366–377 (2001).

85.   Dinkeloo, K., Boyd, S. & Pilot, G. Update on amino acid transporter functions and on possible amino
acid sensing mechanisms in plants. *Semin. Cell Dev. Biol.* **74**, 105–113 (2018).

86.   Negi, J. *et al.* A dof transcription factor, SCAP1, is essential for the development of functional stomata in
arabidopsis. *Curr. Biol.* **23**, 479–484 (2013).

87.   Geisler, M., Nadeau, J. & Sack, F. D. Oriented asymmetric divisions that generate the stomatal spacing
pattern in arabidopsis are disrupted by the too many mouths mutation. *Plant Cell* **12**, 2075–2086 (2000).

88.   Weichert, A. *et al.* AtPTR4 and AtPTR6 are differentially expressed, tonoplast-localized members of the
peptide transporter/nitrate transporter 1 (PTR/NRT1) family. *Planta* **235**, 311–323 (2012).

89.   Schachtman, D. P. *et al.* Expression of an inward-rectifying potassium channel by the Arabidopsis KAT1
cDNA. *Science (New York, N.Y.).* **258**, 5088 (1992).

90.   Menges, M. & Murray, J. A. H. Synchronous Arabidopsis suspension cultures for analysis of cell-cycle
gene activity. *Plant J.* **30**, 203–212 (2002).

91.   Mayer, K F et al. Role of WUSCHEL in regulating stem cell fate in the Arabidopsis shoot meristem.
Cell. **95**(6), 805-815. (1998).

92.   MacAlister, C. A., Ohashi-Ito, K. & Bergmann, D. C. Transcription factor control of asymmetric cell
divisions that establish the stomatal lineage. *Nature* **445**, 537–540 (2007).

93.   Pillitteri, L. J., Sloan, D. B., Bogenschutz, N. L. & Torii, K. U. Termination of asymmetric cell division
and differentiation of stomata. *Nature* **445**, 501–505 (2007).

94.   Schuster, J., Knill, T., Reichelt, M., Gershenzon, J. & Binder, S. BRANCHED-CHAIN
AMINOTRANSFERASE4 is part of the chain elongation pathway in the biosynthesis of methionine-
derived glucosinolates in Arabidopsis. *Plant Cell* **18**, 2664–2679 (2006).

95.   Lu, L., Lee, Y. R. J., Pan, R., Maloof, J. N. & Liu, B. An internal motor kinesin is associated with the
golgi apparatus and plays a role in trichome morphogenesis in Arabidopsis. *Mol. Biol. Cell* **16**, 811–823
(2005).

96.   Li, S. F., Higginson, T. & Parish, R. W. A novel MYB-related gene from Arabidopsis thaliana expressed
in developing anthers. *Plant Cell Physiol.* **40**, 343–347 (1999).

97.   Lai, L. B. *et al.* The Arabidopsis R2R3 MYB proteins FOUR LIPS and MYB88 restrict divisions late in
the stomatal cell lineage. *Plant Cell* **17**, 2754–2767 (2005).

98.   Kubo, M. *et al.* Transcription switches for protoxylem and metaxylem vessel formation. *Genes Dev*.
19(16), 1855-1860. (2005)

99.   Kirik, V. *et al.* Functional diversification of MYB23 and GL1 genes in trichome morphogenesis and
initiation. *Development* **132**, 1477–1485 (2005).

100. O'Malley, R. C. *et al*. Cistrome and epicistrome features shape the regulatory DNA landscape.
Cell. **165(**5), 1280-1292. (2016).

101. Bar-Peled, M. & O'Neill, M. A. Plant nucleotide sugar formation, interconversion, and salvage by
sugar recycling. *Annu Rev Plant Biol*. **62**,127-155. (2011).

102. Lalonde, S., Wipf, D. & Frommer, W. B. Transport mechanisms for organic forms of carbon and
nitrogen between source and sink. *Annu Rev Plant Biol*. **55**, 341-372. (2004).

103. Cobbett, C. & Goldsbrough, P. Phytochelatins and metallothioneins: roles in heavy metal
detoxification and homeostasis. *Annu Rev Plant Biol*. **53**, 159-182. (2002).

Acknowledgments

We want to thank T. Nolan for insightful comments to this manuscript. This research was supported by a Postdoctoral Award No. FI-601-2020 from the United States–Israel Binational Agricultural Research and Development Fund. N.I.-E. is a research fellow at the George E. Hewitt Foundation for Medical Research and an Awardee of the Weizmann Institute of Science – Israel National Postdoctoral Award Program for Advancing Women in Science and the Women's Postdoctoral Career Development Award. J.R.E. is an Investigator of the Howard Hughes Medical Institute.

Author contributions

Conceptualization – N.I-E

Data curation – N.I-E, R.G-C, J.S, B.J, T.Z-K, A.Y

Formal analysis – N.I-E, K.L, J.Y, J.R.N, T.Z-K

Funding acquisition - N.I-E, J.R.E

Investigation - N.I-E, R.G-C

Methodology - N.I-E, T.L, J.S, S.B, Y.Z

Project administration - N.I-E

Resources – J.R.E

Writing – original draft - N.I-E, J.R.E.

Declaration of interests

The authors declare no competing interests.