A molecularly distinct accumbal-to-lateral hypothalamic circuit modulates food seeking and consumption

Yiqiong Liu [1,2,3], Zheng-dong Zhao[1,2,3],

Guoguang Xie [1,2,3], Renchao Chen[1,2,3], and Yi Zhang[1,2,3,4,5*]

[1]Howard Hughes Medical Institute, Boston Children's Hospital, Boston, Massachusetts 02115, USA; [2]Program in Cellular and Molecular Medicine, Boston Children's Hospital, Boston, Massachusetts 02115, USA; [3]Division of Hematology/Oncology, Department of Pediatrics, Boston Children's Hospital, Boston, Massachusetts 02115, USA; [4]Department of Genetics, Harvard Medical School, Boston, Massachusetts 02115, USA; [5]Harvard Stem Cell Institute, WAB-149G, 200 Longwood Avenue, Boston, Massachusetts 02115, USA.

[*]To whom correspondence should be addressed

E-mail: yzhang@genetics.med.harvard.edu

Running title: NAc *Serpinb2*[+] D1 MSN in food reward

Keywords: Food intake, Nucleus accumbens, *Serpinb2*-expressing D1 neuron, motivation, Lateral hypothalamus, Leptin

Manuscript information: 31 pages, 6 figures, 6 supplemental figures

Abstract

Understanding the mechanism of energy homeostasis is expected to lead to effective treatment to obesity and metabolic diseases[1,2]. However, energy homeostasis is a complicated process largely controlled by neuronal circuits in the hypothalamus and brainstem[3-5], whereas reward and motivation of food intake are mainly controlled by the limbic regions[6] and cerebral cortex[7,8]. Although the limbic and hypothalamus connection like Nucleus Accumbens shell (NAcSh) to the lateral hypothalamus (LH) circuit has been reported to regulate feeding[9,10], the neuron subtypes involved, and how do the humoral/neuronal signals coordinate to direct feeding behavior remain unknown. Here we show that the projection from dopamine receptor D1(Drd1)- and *Serpinb2*-expressing subtype to leptin receptor (LepR) expressing neurons in LH modulates food seeking and consumption. We demonstrate that the *Serpinb2*$^+$ neuronal activity is dynamically modulated during feeding. Conversely, chemo/optogenetics-mediated modulation of *Serpinb2*$^+$ neurons bidirectionally regulate food seeking and consumption. Importantly, circuitry stimulation revealed the NAcSh^{Serpinb2}→LH^{LepR} projection controls refeeding and overcomes leptin-mediated feeding suppression. Ablation of NAcSh^{Serpinb2} neurons could decrease body weight. Together, our study reveals a molecularly distinct accumbal-to-lateral hypothalamic neural circuit that controls internal state-dependent food consumption, which provides a promising therapeutic target for anorexia and obesity.

Main

The hypothalamus, with highly heterogenous neuronal composition[11], plays a critical role in controlling feeding behavior[12,13], where feeding related hormones, such as ghrelin and leptin, coordinately produce sensations of appetite and satiety leading to behavioral response[14,15]. Traditionally, people regard the arcuate nucleus (Arc) as a major location where LepR performs anorexic function by acting on leptin receptors (LepR) to suppress food intake and bodyweight gain[16]. In addition to the Arc, lateral hypothalamus (LH)[17] also highly expresses LepR to play a similar role. However, the specific neuronal subtype targeting LH^{LepR} neurons beyond hypothalamus to regulate feeding remain to be elucidate. In recent years, several studies have analyzed the role of mediodorsal NAcSh in feeding[10,18,19] and revealed that activation of dopamine receptor 1 expressing medium spiny neurons (D1-MSNs)[20,21] projections to LH[9,10] stops ongoing food consumption. However, other studies showed that D1-MSNs activity was

enhanced during appetitive phase[22] as well as during consumption[23]. Although temporally

distinct phases of feeding behavior, such as food seeking, food evaluation and consumption,

could potentially account for such discrepancy, the neuronal heterogeneity of NAc[24] could

underlie the seemingly conflict feeding behavior as the different studies might have manipulated

different neuron subtypes with opposing functions. With the application of single cell RNA-seq

and spatial transcriptome techniques to decipher the neuron heterogeneity of different brain

regions[25-29], we could focus on different neuron subtypes located in the NAcSh implicated in

feeding behavior[9,10,19,22,30,31] to identify the neuron subtype(s) responsible for regulating feeding

behavior.

***Serpinb2*[+] neurons are activated in refeeding process**

Using iSpatial[32], an algorithm that integrates single cell transcriptome and spatial transcriptomic

information[24], we analyzed NAc MSN subtypes with medial dorsal NAcSh distribution. We

found that the *Tac2*, *serpinb2* and *Upb1* MSN subtypes exhibit distinct distribution patterns in

medial dorsal NAcSh (Fig. 1a). Since D1-MSNs, but not D2-MSNs, provide the dominant source

of accumbal inhibition to LH with rapid control over feeding via LH GABA neurons[19,22], we

focused our effort on the *Tac2* and *Serpinb2* D1-MSN subtypes. tSNE plots of scRNA-seq result

indicated that the *Tac2*[+] neurons are mainly enriched in the D1-MSN subclusters 6 and 8, while

Serpinb2[+] neurons are enriched in D1-MSN subcluster 2[24] (Fig. 1b). RNA-FISH further

confirmed that both *Tac2* and *Serpinb2* subtypes belong to D1-MSN (co-express Drd1) and are

mainly localized to medial dorsal NAcSh (Fig. 1c). Consequently, *Tac2*[+] and *Serpinb2*[+] MSN

subtypes are good candidates with potential in mediating feeding behavior.

To determine whether *Tac2*[+] and *Serpinb2*[+] MSN subtypes can mediate feeding behavior, we

first asked whether the activities of these neurons respond to feeding behavior by monitoring the

cFos expression under three conditions: ad libitum access to food, after 18 hours of fast, and

refeeding (Fig. 1d). By counting *cFos*[+] neurons that co-express *Serpinb2* or *Tac2* in the medial

dorsal NAcSh under Ad libitum, fast and refed conditions (Fig. 1e), we determined whether the

Serpinb2 and *Tac2* neurons respond to the different feeding status. We found fasting and

refeeding both increased neuronal activities compared to Ad libitum as indicated by the

increased *cFos*[+] neuron numbers (Fig. 1f). Importantly, most of the *Serpinb2*[+] neurons (~70%)

were activated by refeeding, but not fasting (Fig. 1g). In contrast, the *Tac2*[+] neurons do not

respond to refeeding or fasting (Fig. 1h). Collectively, these data indicate that the majority of

Serpinb2[+] neurons respond to refeeding process.

The *Serpinb2*[+] neurons respond to eating behavior

To facilitate studying the role of *Serpinb2*[+] neurons in the feeding process, we generated a

Serpinb2-Cre mouse line (Extended Data Fig. 1a,b). We validated this mouse model by injecting

a Cre-dependent AAV vector expressing light-gated cation channel channelrhodopsin (ChR2)

and observed about 90% colocalization of *Serpinb2*::ChR2-eYFP signal with endogenous

Serpinb2 mRNA signal (Extended Data Fig. 1c), which is consistent with endogenous *Serpinb2*

expression in the NAcSh as shown by Allen Brain Atlas. Thus, our *Serpinb2*-Cre mouse line can

be used for studying *Serpinb2*[+] neurons.

To test whether *Serpinb2*[+] neurons are involved in regulating feeding, we used fiber photometry

to monitor *Serpinb2*[+] neuronal activity during food seeking and consumption. To this end, we

inserted an optic cannula into the NAcSh to record the total *Serpinb2*[+] neuronal activity

(reflected by calcium reporter fluorescence intensity) by stereotaxic injection of a Cre-dependent

AAV expressing the seventh-generation calcium reporter GCaMP7s into the NAcSh region of

the *Serpinb2*-Cre mice. In parallel, we also implanted cannula to the *Drd1*-Cre mice for

comparative study (Fig. 2a). Three weeks after the viral injection, we performed fluorescence

recording during the feeding process. To monitor the activity dynamics of the *Serpinb2*[+] neurons,

we designed 3-chamber food seeking and food consumption assay (Fig. 2d) and aligned calcium

traces of mice with behavioral events that include habituation, food zone approaching, eating,

and leaving (Fig. 2b, c). First, we recorded the Ca^{2+} signals during the different feeding phases

after the mice were fasted overnight. In habituation phase, mice were allowed to freely travel

among the 3 chambers and we detected negligible response when mice approach the two empty

food cups (Fig. 2d, first graph). In food approaching phase, when mice enter the food zone to

interact with caged food pellets, the activity of the *Serpinb2*[+] neurons increased immediately

when mice entered the food zone and lasted for seconds (Fig. 2d, second graph), indicating that

Serpinb2[+] neurons respond to appetitive food seeking. In the eating phase, we observed a

significant increase of Ca^{2+} signals when mice start eating the food (Fig. 2d, 3[rd] graph), and

subsequently, declined to lower than baseline level after eating finished. Usually, mice turned

away and left the food zone (Fig. 2d, 4th graphs).

To quantify the total *Serpinb2*$^+$ neuronal activity at different phases, we calculated the area under

curve index (AUC) of the calcium signal curve for each trial. We find that the AUC is

significantly higher in the food zone approaching and eating phases and lower in the post eating

phases compared with that of habituation (Fig. 2d, right panel). In parallel experiments with the

Drd1-Cre mice, we observed Ca^{2+} signal tended to increase when approach to food zone (Fig. 2e,

second graph) and reduce during consumption (Fig. 2e, 3rd graph). After eating, *Drd1*$^+$ neuronal

activity increased concomitantly (Fig. 2e, 4th graphs). Similar results were also observed in ad

libitum status (Extended Data Fig. 2). These results indicate that *Serpinb2*$^+$ neurons function

differently from other *Drd1*$^+$ neurons especially in appetitive and consumption phases.

Serpinb2$^+$ neurons bidirectionally regulate food intake in hungry state

To determine whether *Serpinb2*$^+$ neuronal activity plays a causal role in regulating feeding

behavior, we asked whether the feeding behavior can be changed by manipulating *Serpinb2*$^+$

neurons' activity. To this end, we applied chemogenetic techniques[33] to the *Serpinb2*-Cre mice

by injecting the activating AAV encoding a modified human M3 muscarinic receptor (hM3Dq)

or the inhibitory vector AAV-DIO-hM4Di-mCherry into the NAcSh region. As controls, we also

performed parallel experiments using *Tac2*-Cre and *Drd1*-Cre mice (Extended Data Fig. 3a). We

first confirmed the accuracy of the injection site (Extended Data Fig. 3b), then the efficiency of

the activation and inhibition with *cFos* expression (Extended Data Fig. 3c).

We then tested whether activation or inhibition of the *Serpinb2*$^+$ neurons affect feeding and

reward-related behaviors (Fig. 3a). Interestingly, under Ad libitum conditions, manipulation of

the *Serpinb2*$^+$ neuronal activity does not affect food intake (Fig. 3b**,** left panel), indicating that

the eating behavior under Ad libitum maybe controlled by other neurons. Next, we performed the

same test using fasted mice to analyze total food consumption during refeeding (Fig. 3a-c). We

found that activation of the *Serpinb2*$^+$ neurons increased food consumption, while inhibition of

the *Serpinb2*$^+$ neurons decreased food consumption (Fig. 3b**,** right panel). However, similar

manipulation on *Tac2*-Cre or *Drd1*-Cre mice did not affect food consumption (Fig. 3c). In

addition to the total food consumption, we also analyzed food preference indicated by the time

spent in the food zone (Fig. 3c). As expected, fasted mice would significantly increase their time

spent in the food zone for the control mouse group (Fig. 3d). Importantly, activation (hM3Dq) or

inhibition (hM4Di) the *Serpinb2*[+] neurons respectively increased or decreased the time the mice

spent in the food zone compared to that of the control mice (Fig. 3e). In contrast, similar

manipulation of the *Tac2*-Cre or *Drd1*-Cre mice showed no such effect compared to that in the

control mice (Fig. 3e). These results indicate that manipulation the *Serpinb2*[+] neuronal activity

can regulate food seeking and intaking behavior in hungry state.

To further determine whether the neuronal activity of the *Serpinb2*[+] neurons has a causal role in

regulating food motivation, we next carried out food operant chamber test (Fig. 3f, g). To

maintain a similar food motivation status, all mice were food restricted to reduce body weight to

around 90% of their original value. After trained to operantly respond to sweetened chow pellets

on fixed ratio (FR) 1, 3, 5 schedule, the animals were then tested for lever pressing upon CNO-

induced chemogenetic manipulation. For the FR5 test, *Serpinb2*[+] neuron activation significantly

increased the active lever pressing and pellet reward (Fig. 3h), while *Serpinb2*[+] neuron inhibition

elicited the opposite effect (Fig. 3h). For the progressive ratio (PR) 5 test, *Serpinb2*[+] neuron

activation showed a tendency of increased number of active lever pressing and significantly

increased pellet reward (Fig. 3i), while *Serpinb2*[+] neuron inhibition decreased both the total

active lever pressing and the reward (Fig. 3i). Taken together, these results demonstrate that the

Serpinb2[+] neurons are involved in bidirectional control of goal-direct food seeking behavior.

NAc shell is also known to regulate anxiety[34] and drug reward[35,36]. Thus, we asked whether

Serpinb2[+] neurons also regulate these behaviors under the same chemogenetic manipulation. We

found that manipulation of *Serpinb2*[+] neuronal activity does not affect locomotion in open field

test (Extended Data Fig. 4a,b), or drug reward in cocaine conditioned place preference (CPP) test

(Extended Data Fig. 4c), or anxiety in elevated plus maze (EPM) test (Extended Data Fig. 4d,e).

Taken together, these results support that *Serpinb2*[+] neurons are specifically involved in food

refeeding process, but not involved in locomotion, anxiety or drug seeking behaviors.

***Serpinb2*[+] neurons mediate food consumption via LH projection**

Thus far, characterization of the *Serpinb2*[+] neurons was at the somatic level in NAcSh. Next, we

attempt to understand how the *Serpinb2*[+] neurons regulate food taking behavior at the circuit

level. Previous studies have indicated that NAcSh D1-MSNs could project to multiple brain

regions, including ventral tegmental area (VTA)[37], ventral pallidum (VP)[38], and bed nucleus stria

terminalis (BNST)[39]. To determine the projection site of *Serpinb2*[+] neurons, we performed

anterograde tracing by injecting Cre-dependent AAVs expressing ChR2(H134R) into the NAcSh

of the *Serpinb2*-Cre mice (Fig. 4a). Analysis of the brain slices after 3 weeks of virus injection

revealed that the *Serpinb2*[+] neurons only project to lateral hypothalamus (LH) (Fig. 4a). To

further validate this projection, we injected the widely used retrograde tracer Cholera toxin

subunit B (CTB) [40] into the LH region (Fig. 4b), and observed colocalization of CTB with

mCherry in the NAcSh after immunostaining of NAcSh from the *Serpinb2*-Cre mice (Fig. 4b, c),

supporting the NAcSh to the LH projection.

To demonstrate that the NAcSh to the LH projection of the *Serpinb2*[+] neurons is functionally

relevant to food intaking, we asked whether optogenetic manipulation of the *Serpinb2*[+] neuron

terminals in LH can change the feeding behavior of the *Serpinb2*-Cre mice. To this end, DIO-

ChR2-eYFP or DIO-NpHR-eYFP AAV viruses were injected to the NAcSh of the *Serpinb2* -Cre

mice with optic cannula implanted into their LH region (Fig. 4d). After fasting the mice for

overnight, we activated the *Serpinb2*[+] neuron terminals in the LH with blue light on-off

stimulation (20 Hz, 2-ms pulses). We found that mice with the *Serpinb2*[+] neuron terminal

stimulation significantly increased their food intake in 20 mins compared to the control mice that

express eYFP (Fig. 4e, left panel). Conversely, *Serpinb2*[+] neuron terminal inhibition in the LH

with yellow on-off stimulation (20 Hz, 2-ms pulses) decreased total food intake when compared

with the control (Fig. 4e, right panel). Collectively, viral tracing and circuit manipulation

demonstrate that the NAcSh to LH projecting *Serpinb2*[+] neurons have an important function in

regulating food intaking behaviors.

***Serpinb2*[+] neurons form a circuit with LH LepR[+] GABA[+] neurons**

Having demonstrated the functional importance of the NAcSh to LH projection, we next

attempted to determine the neuron types in LH that receive signals from the *Serpinb2*[+] neurons.

The LH is a highly heterogeneous brain region controlling food intake, energy expenditure, and

many other physiological functions[41]. Since neural peptides orexin/hypocretin and melanin-

concentrating hormone (MCH) are associated with feeding[42,43], and are mainly express in LH,

we first asked whether they are the down-stream targets of the NAcSh *Serpinb2*$^+$ neurons. To this

end, we injected the AAV-DIO-ChR2 anterograde viruses to the NAcSh of the *Serpinb2* -Cre

mice and performed immunostaining of candidate neural peptides or transmitters on slices

covering the LH (Fig. 5a). We found very few MCH- or orexin-expressing neurons in LH

overlapped with *Serpinb2*$^+$ terminals (Fig. 5b, indicated by arrow heads). Given that GABAergic

neuron is the major subtype in the LH and has been reported to be involved in feeding and leptin-

regulated energy homogenesis [11,44,45], we next checked whether leptin receptor (LepR) positive

GABAergic neurons[17] receive projections from NAcSh *Serpinb2*$^+$ neurons by similar

immunostaining. We found over 70% *Serpinb2*$^+$ neuron terminals overlap with LepR$^+$ GABA$^+$

neurons (Fig. 5c). This data indicates that the NAcSh *Serpinb2*$^+$ neurons mainly project to LepR$^+$

GABA$^+$ neurons in LH.

To identify the brain regions that response to refeeding, we performed whole brain *cFos*

mapping on the refeeding mice. By postmortem immunostaining, we found *cFos* expression was

increased in multiple brain regions compared with that of ad libitum, like NAc, Olfactory

tubercle (OT), paraventricular nucleus of the thalamus (PVT), Arc, Dorsomedial nucleus of the

hypothalamus (DMH), LH and Zona incerta (ZI) (Fig. 5d, Extended Data Fig. 5). We next

attempted to identify inputs for the NAcSh *Serpinb2*$^+$ neurons by performing a modified rabies

tracing experiment. To this end, the Cre-inducible avian sarcoma leucosis virus glycoprotein

EnvA receptor (TVA) and rabies virus envelope glycoprotein (RG) were injected unilaterally to

the NAcSh of *Serpinb2*-Cre mice (Fig. 5e) to allow monosynaptic retrograde transportation and

rabies virus infection in the starter neurons, respectively[37,46,47]. Two weeks later, the modified

rabies virus SADDG-EGFP (EnvA) was injected unilaterally into the NAcSh and slices of the

whole brain were imaged one week later. Confocal imaging results indicated that EGFP-labeled

neurons can be found in anterior cingulate area (ACA), PVT, LH, and lateral preoptic area (LPO)

(Fig. 5f, Extended Data Fig. 6), indicating that neurons in these regions send monosynaptic

projection to NAcSh to form a network regulating food consumption (Fig. 5g).

Combining the major efferent regions with *cFos* mapping (Fig. 5d, f), PVT may serve as a major

input for *Serpinb2*[+] neurons in NAcSh to regulate food seeking and taking. Collectively, our

study uncovered a neuronal network where the NAcSh *Serpinb2*[+] neurons may receive signals

from PVT neurons to inhibit the LepR[+]GABA[+] neurons in LH, to regulate food seeking and

eating behaviors.

Modulating *Serpinb2*[+] neuronal activity can overcome leptin effect and alter bodyweight

As an adipose-derived hormone, leptin plays a central role in regulating energy homeostasis[48-50].

Leptin performs most of its functions, including suppression of food intaking, by activating the

LepR on central nerve system (CNS) neurons[51,52]. Since the NAcSh *Serpinb2*[+] neurons are

projected to LepR[+] GABAergic neurons in LH (Fig. 5c), we anticipate that both leptin and the

NAcSh *Serpinb2*[+] neurons have shared neuron targets and consequently they should have

functional interaction. To analyze their functional interaction in food intaking, we implanted

catheter in the LH for leptin delivery (catheter administration) on the *Serpinb2*-Cre mice that

were also injected with hM3Dq-mCherry-expressing AAV into the NAcSh so that the NAcSh

Serpinb2[+] neurons can be activated by CNO by i.p. injection. First, we established that 1 μg of

bilateral intra-LH leptin cannula delivery[53] significantly decreased food intake in 3 hours

compared to the control with saline treatment (Fig. 6a). Then we used 1 μg of leptin for all the

following tests. As we have shown previously (Fig. 3b), CNO-induced *Serpinb2*[+] neuron

activation increased food intake (Fig. 6b). Importantly, although leptin delivery reduced the food

intake, the leptin effect can be at least partly overcome by CNO-induced *Serpinb2*[+] neuron

activation (Fig. 6b). This data indicates that *Serpinb2*[+] neurons' innervation to LH can at least

partly overcome leptin's inhibitory effect on food intake.

To access whether loss function of the *Serpinb2*[+] neurons can exert a long-term effect on energy

homeostasis, we selectively ablated NAcSh *Serpinb2*[+]neurons in *Serpinb2*-Cre mice by injecting

a flex-taCasp3-TEVp AAV expressing caspase-3 which eliminates the neurons by inducing cell

death (Fig. 6c). NAcSh *Serpinb2*[+] neurons ablation decreased food intake (Fig. 6d) as well as

reduced body weight gain by 10% in 7 weeks (Fig. 6e). Together, these results demonstrate that

transient change in *Serpinb2*[+] neuronal activity could alter food seeking and consumption

behaviors. Modulating *Serpinb2*[+] neuronal activity could partly overcome leptin effect to
maintain energy homeostasis, and ablation of *Serpinb2*[+] neurons can lead to bodyweight loss.

Discussion

Feeding is an essential goal-directed behavior that is heavily influenced by homeostatic state and
motivation. The accumbal-to-lateral hypothalamic pathway has been implicated in regulating
feeding behavior, but the specific neuron subtypes and precise neuronal circuit in LH are not
clear. In this study, we filled in this knowledge gap by delineating a circuit which integrates
neuronal and humoral signal to regulate food consumption in an innate energy state-dependent
manner. Specifically, we identified a D1-MSN subtype located in the NAcSh and expresses
Serpinb2 to regulate the feeding behavior through a neuronal circuit involving
$NAcSh^{Serpinb2} \rightarrow LH^{LepR}$ that is beyond the hypothalamus. We demonstrate that the *Serpinb2*[+]
neurons bidirectionally modulate food motivation and consumption specifically in hungry state.
Importantly, *Serpinb2*[+] neurons target the LepR expressing GABAergic neurons in LH and their
activation can at least partly overcome the suppressive effect of leptin on food intaking, and
whose ablation can chronically cause bodyweight loss.

The *Serpinb2*[+] neurons are functionally distinct from the pan D1-MSNs in NAcSh

Previous studies have observed reduced D1 firing during food consumption, and consistently,
suppressing D1-MSNs activity prolonged food intake[19]. Using *Serpinb2*-Cre and *Drd1*-Cre mice,
we compared the *Serpinb2*[+] MSNs neurons and the D1-MSNs in regulating feeding behaviors
(Fig. 2, 3), and found their manipulation have different outcomes. First, *Serpinb2*[+] neurons are
activated by both food approaching and food consumption, while *D1*[+] neurons show bi-phasic
response, activated during food approaching but suppressed during food consumption. Second,
Serpinb2[+] neurons bidirectionally regulate food seeking and intake, particularly during refeeding,
while *D1*[+] neuron manipulation does not significantly alter feeding behavior. On the other hand,
a previous study showed that *D1*[+] neuron inhibition promoted liquid fat food intake[19]. This
difference might be due to the different feeding assays used in the two studies, while we used
free-access chow food intake, the previous study used a head-fixed mice licking liquid fat food
as the assay[19]. Third, *Serpinb2*[+] neuron ablation significantly reduced food intake (Fig. 6), which

is consistent with our finding that *SerpinB2*[+] neuron activation positively regulates food intake

(Fig. 3). However, a previous study indicated that lesions or inactivation of the NAc neurons do

not significantly alter the food consumption[54]. We do not consider these results to be in conflict

as NAc is composed of many D1- and D2-MSN neuron subtypes, of which many are not

involved in regulating food intake, while others can positively or negatively regulate food intake.

Consequently, manipulating *Serpinb2*[+] neurons and the entire NAc neurons can have totally

different outcomes. This further indicates that finer granularity and cell type-specific approaches

are needed to dissect the function of different neuron subtypes in NAc. For example, although

the *D2*[+] neuronal activity as a whole is not altered during food consumption[19,22], the D2 receptors

are indeed downregulated in obese rodent as well as human[55,56], whether certain D2-MSN

subtypes are involved in regulating food intake remains to be determined.

***Serpinb2*[+] and *Tac2*[+] MSNs respectively modulate food and drug reward**

The NAcSh has long been implicated in regulating reward-related behaviors that are associated

with food, social, and drug. Previous studies were mainly based on dichotomous MSNs subtypes

that express dopamine receptor 1 or dopamine receptor 2 (D1- or D2-MSNs)[57]. Increasing

evidence suggest that the NAcSh is highly heterogeneous in terms of the molecular features and

anatomical connections of the neurons located in this region. This raises an interesting question

that whether these reward-related behaviors are regulated by distinct or overlap neuron subtypes

and/or projections. By combining iSpatial analysis, *cFos* mapping, neuronal activity

manipulation of different subtypes, and behavioral tests, we found that the *Serpinb2*[+] D1-MSNs

of NAcSh specifically regulate food reward, but not drug reward or other emotional and

cognitive functions (Fig. 1, 3, Extended Data Fig. 4). On the other hand, we have previously

showed that the *Tac2*[+] D1-MSNs of NAcSh specifically regulate cocaine reward[58]. These studies

indicate that different reward behaviors are at least partly regulated by distinct MSNs subtypes.

An important task for future studies will be to identify the relevant neuron subtypes regulating

the various reward-related behaviors, and eventually to link the cellular heterogeneity to

functional diversity of each brain regions. This way, the cellular and circuit mechanisms

underlying the various behaviors can be elucidated.

The PVT- NAcSh $^{Serpinb2+}$ -LH $^{LepR+}$ circuit controls feeding in hungry state

Previous studies have shown that either LH GABA or Vglut2 neuronal subpopulation can

receive NAc innervation[19,59]. However, the LH GABA and Vglut2 neurons are extremely

heterogeneous, and can be further divided into 15 distinct populations, respectively[60]. Thus, the

specific cell types that receive NAc innervation were unknown. Previous studies also showed

that NAcSh D1- MSN to LH inhibitory transmission stops eating, and endocannabinoids

mediated suppression of this projection promotes excessive eating of highly palatable chow[61],

but the D1 subtype involved in this projection was not known. Using viral tracing, we discovered

that the NAcSh *Serpinb2*$^+$ D1-MSNs project to *LepR*$^+$ neurons in LH underlying the *Serpinb2*$^+$

neuron function in food intake (Fig. 6c). Distributed in numerous regions involved in the

regulation of energy balance, the *LepR*$^+$ neurons lie in the mediobasal hypothalamic (MBH)

"satiety centers" and in LH that is regarded as the "feeding center"[50,62]. Leptin treatment induced

cFos expression and 100 nM of leptin depolarized 34% of LepR-expressing neurons in LH[17].

Unilateral intra-LH leptin decreased food intake and bodyweight[17]. In our study, we found

activation of the *Serpinb2*$^+$ neurons increased the inhibition of the *LepR*$^+$ neurons excitability,

resulting in increased food consumption; while inhibition of *Serpinb2*$^+$ neurons decreased the

inhibition of *LepR*$^+$ neurons excitability, leading to decreased food consumption even after

fasting (Fig. 3b, 4e). Our results are consistent with previous reports demonstrating that LH

LepR neuron activation decreased chow intake[63]. Importantly, manipulating *Serpinb2*$^+$ neuronal

activity could override leptin's effect in LH to modulate food consumption (Fig. 6b). It is not

clear whether the lack of effect of the *Serpinb2*$^+$ neurons on feeding behavior in ad libitum is due

to the lack of food taking motivation or the relatively high leptin level masked the *Serpinb2*$^+$

neuron effect. Alternatively, it is possible that the endocannabinoid and leptin signaling may

interact in LH, where activation of NAcSh *Serpinb2*$^+$ neurons suppresses LepR neurons in the

LH, which may increase the synthesis and release of endocannabinoids[64] and thus promote

feeding. Our study thus reveals a parallel and compensatory circuit from NAcSh to LH^{LepR},

which is beyond the hypothalamus circuit that directly modulates food intake, to maintain energy

homeostasis.

For the efferent site, *Serpinb2*$^+$ neurons majorly receive input from PVT based on the GFP$^+$ cell

numbers (Fig. 5f), which is believed to be an integration hub processing information and sending

"command" to the downstream targets[65-67]. Previous studies revealed that the *Slc2a2*$^+$ neurons *in*

PVT are activated by hypoglycemia and their activation by optogenetics increases motivated

sucrose seeking behavior[68]. On the other hand, the *Gck*[+] neurons in PVT have the opposite

glucose sensing property as their optogenetic activation decreased sucrose seeking behavior[69].

Taken together, we believe that in hungry state, PVT receives the "hungry signal" and send it to

Serpinb2[+] neurons. The activated *Serpinb2*[+] neurons then instruct the LH *LepR*[+] neurons to

promote eating. This PVT- NAcSh *Serpinb2+* -LH *LepR+* circuit controls feeding in hungry stage and

strengthens the sentinel role of NAcSh[19,70].

Serpinb2[+] neurons connect energy homeostasis and motivation

Feeding is an essential goal-directed behavior that is influenced by cellular homeostasis state and

appetitive motivation. Interestingly, we found that *Serpinb2*[+] neurons promote feeding in hunger

state, rather than in normal feeding state, suggesting that *Serpinb2*[+] neuron function is regulated

in an internal metabolic state-dependent manner. Consistently, ablation of *Serpinb2*[+] neurons

significantly reduced food intake and leading to bodyweight loss (Fig. 6e,f). The *Serpinb2*[+]

neurons are activated by both appetitive food approaching and food consumption. To measure

appetitive food motivation, we conducted operant food intake assay, where mice need to press

levers to earn food pellets. We found that *Serpinb2*[+] neuronal activity bi-directionally regulates

active lever presses and the earned reward (Fig. 3h,i). These studies suggested that *Serpinb2*[+]

neurons regulate both energy homeostasis and appetitive motivation which is consistent with the

demonstrated function of NAc in integrating descending signals pertaining to homeostatic needs

and goal-related behaviors[3,71]. Collectively, these data indicate that lose function of the NAcSh

Serpinb2[+] neurons can disrupt energy homeostasis and appetitive motivation, which provides a

potential therapeutic target for obese treatment. Conversely, activation of the NAcSh *Serpinb2*[+]

neurons can be a potential strategy for anorexia treatment.

In conclusion, we identified a molecularly defined neuron population with crucial functions in

regulating food intake via neuron-hormone axis. From a therapeutic point of view, our findings

are highly relevant because activating or ablating a small population of molecularly defined

neurons could respectively rescue food intake at low energy status or lead to long-term

bodyweight loss. Given its function, we believe the small population of NAcSh *Serpinb2*[+]

neurons is an ideal entry point for understanding the complex brain-metabolism regulatory

network underlying eating and bodyweight control.

Methods

404
Mice

All experiments were conducted in accordance with the National Institute of Health Guide for
Care and Use of Laboratory Animals and approved by the Institutional Animal Care and Use
Committee (IACUC) of Boston Children's Hospital and Harvard Medical School. The *Serpinb2-*
Cre mice were generated as described below. For behavioral assays, 12 -16 weeks old male mice
were used. The mice were housed in groups (3-5 mice/cage) in a 12-hr light/dark cycle, with
food and water ad libitum unless otherwise specified. The Tac2-Cre knock-in mouse line was a
gift from Q. Ma at Dana-Farber Cancer Institute and Harvard Medical School. The D1-cre mouse
line was obtained from Jackson Laboratory (JAX: 037156).

Fluorescence *in situ* hybridization (FISH) and immunofluorescence (IF) staining
Mice were transcardially perfused with PBS followed by 4% paraformaldehyde. Brains were
then placed in a 30% sucrose solution for 2 days. The brains were frozen in Optimal Cutting
Temperature (OCT) embedding media and 16 μm (for FISH) or 40 μm (for IF) coronal sections
were cut with vibratome (Leica, no. CM3050 S). For FISH experiments, the slices were mounted
on SuperFrost Plus slides, and air dried. The multi-color FISH experiments were performed
following the instructions of RNAscope Fluorescent Multiplex Assay (ACD Bioscience). For IF,
cryostat sections were collected and incubated overnight with blocking solution (1×PBS
containing 5% goat serum, 5% BSA, and 0.1% Triton X-100), and then incubated with the
following primary antibodies, diluted with blocking solution, for 1 day at 4 °C. Samples were
then washed three times with washing buffer (1×PBS containing 0.1% Tween-20) and incubated
with the Alexa Fluor conjugated secondary antibodies for 2 h at room temperature. The sections
were mounted and imaged using a Zeiss LSM800 confocal microscope or an Olympus VS120
Slide Scanning System. Antibodies used for staining were as follows: rabbit anti-cFos (1:2000,
Synaptic systems, #226003), chicken anti-GFP (1:2000, Aves Labs, no. GFP-1010), chicken
anti-mCherry (1:2000, Novus Biologicals, no. NBP2-25158), Orexin-A (KK09) antibody, (1:500,
Santa Cruz Biotechnology, Cat# sc-80263), anti-MCH, Ab1-pmch antibody (1:2000, Phoenix
Pharmaceuticals, Cat# H-070-47), Anti-GABA antibody (1:1000, Sigma, Cat# A2052) , Leptin
Receptor antibody (Abcam, Cat# ab104403), Goat anti-chicken Alexa Fluor 488(Thermo Fisher

Scientific, Cat# A11039), Donkey anti-rabbit Alexa Fluor 488(Thermo Fisher Scientific, Cat#

A21206), Donkey anti-rabbit Alexa Fluor 568(Thermo Fisher Scientific, Cat# A10042).

AAV vectors

The following AAV vectors (with a titer of $>10^{12}$) were purchased from UNC Vector Core:

AAV5-EF1a-DIO-hChR2(H134R)-EYFP,   AAV5-EF1a-DIO-EYFP,   AAV-DJ-

EF1a-DIO-GCaMP6m. The following AAV vectors were purchased from Addgene: AAV5-

hSyn-DIO-hM3D(Gq)-mCherry   (#44361),   AAV5-hSyn-DIO-hM4D(Gi)-mCherry   (#44362),

AAV5-hSyn-DIO-mCherry (#50459), pAAV-flex-taCasp3-TEVp (#45580), pAAV-Ef1α-DIO

eNpHR 3.0-EYFP (#26966), EnvA G-deleted Rabies-EGFP (Salk Institute).

Stereotaxic brain surgeries

The AAV vectors were injected through a pulled-glass pipette and the nanoliter injector

(Nanoject III, Drummond Scientific - 3-000-207). The injection was performed using a small-

animal stereotaxic instrument (David Kopf Instruments, model 940) under general anesthesia by

isoflurane (0.8 liter/min, isoflurane concentration 1.5%) in oxygen. A feedback heater was used

to keep mice warm during surgeries. Mice were allowed to recover in a warm blanket before

they were transferred to housing cages for 2–4 weeks before behavioral evaluation was

performed. For chemogenetics experiments, 0.1~0.15 μl of AAV vector was bilaterally delivered

into target regions. For optogenetics experiments, following viral injection, the fiber optic

cannula (200 μm in diameter, Inper Inc.) were implanted 0.1 mm above viral injection site and

were secured with dental cement (Parkell, #S380). For the drug delivery cannula implantation,

the cannula (Guide cannula: C.C 2.0mm, C=4.5mm; Injector: G1=0.5mm; Dummy cannula:

G2=0, RWD Life Science) were directly implanted 0.5 mm above the LH and were secured with

dental cement (Parkell, #S380). The coordinates of viral injection sites are based on previous

literature as follow: NAc (anterior-posterior [AP] +1.2, medial-lateral [ML] ± 0.6, dorsal-ventral

[DV] -4.5 mm) and LH (AP -1.3, ML ± 1.2, DV -5.0 mm).

Neuronal Tracing

For CTB tracing, mice were injected with 100–200 μl CTB-647 (AF-CTB, all from Life

Technologies) unilaterally into the LH (AP –1.2, ML +1.2, DV –4.75 mm). For rabies tracing,

Serpinb2-cre mice were first unilaterally injected in the NAcSh with the starter AAV. After 14 d,

the same mice were injected with the rabies virus. Then, 11 days after CTB injections and 7 days

after rabies injections, brains tissue was collected and processed for confocal imaging. To aid

visualization, images were adjusted for brightness and contrast using ImageJ, but alterations

always were applied to the entire image.

Behavioral assays

Open-field tests (OFT). A clear box (square 27.3 cm x 27.3 cm square base with 20.3 cm high

walls) used for the open field test, and the center zone was 50% of the total area. Prior to testing,

mice were habituated to the test room for at least 20 minutes. Mice were placed in the center of

the box at the start of the assay. Movement was recorded using a measurement (Med Associates,

St. Albans, VT, ENV-510) 1 hour in 5 mins bins. In addition to regular parameters related to

locomotor activity (such as total travel distance, velocity, ambulatory time, resting time), time

spent, and distance travelled in the center area of the testing arena were also recorded and

analyzed.

Elevated plus maze (EPM). EPM was used to measure anxiety effect. Before EPM test, mice

were brought to the testing room for environmental habituation for at least 30 min. The EPM

apparatus is consisted of an elevated platform (80 cm above the floor), with four arms (each arm

is 30 cm in length and 5 cm in width), two opposing closed arms with 14 cm walls and two

opposing open arms. Mice were attached into the fiber-optic patch cord and were individually

placed in the center of the EPM apparatus, towards one of the open arms. The mice trajectories

were tracked for 5 minutes, and the time spent in the open arms was analyzed using Ethovision

XT11 (Noldus).

Cocaine conditioned place preference (cocaine-CPP). Mice were allowed to freely explore

both sides of a custom-made (Med Associates) CPP training apparatus ($25 \times 19 \times 17$ cm for L $\times$

D $\times$ H) for 30 min. Trajectories were tracked by infra-red photobeam detectors, and the travel

distance and the duration were recorded to assess their baseline place preference. Mice that

showed strong bias (< 25% preference) were excluded from the experiments. Then, for

chemogenetic activation or inhibition during CPP formation, these mice were injected with

saline (i.p.) and confined to their preferred side of the chamber for 30 min before returned to

their home cage. At least four hours later, the same mice received CNO at least 15 min before an

i.p. injection of 15 mg/Kg cocaine and were confined to their non-preferred side of the chamber

for 30 minutes. They were then returned to their home cage. The same training with saline and

CNO injection were performed for three consecutive days. Twenty-four hours after the final

training session, mice were re-exposed to the CPP chamber and allowed to explore both sides of

the chamber for 30 min.

Post-fasted food intake. Mice were individually placed in the home cage and fasted overnight

(18 hours). Mice received N-clozapine (CNO, 2mg/mL for hM3Dq group and 5mg/mL for

hM4Di group) via i.p. injection and then regular chow pallets (3 g per pellet) were put in the

hopper. Three hours later, the remaining food pallets were collected and measured to calculate

total amount of food consumed (g). For the leptin treatment test, 15 mins after CNO injection, 1

µg of leptin (R&D, Cat# 498-OB) was delivered through cannula by pump for 5 mins and waited

for another 5 mins before adding pellets. For the Ta-Casp3 treatment group, the test was carried

out 3 weeks after virus injection.

Food place preference. Animals were placed in a custom three-chamber, $45 \times 60 \times 35$ cm arena

to assess the amount of food consumed and time spent in a designated food zone area. The arena

contained two 64-cm^2 food cups in two outer corners of separate chambers. One cup contained

standard grain-based chow (Harlan, Indianapolis, IN), while the other cup remained empty. Mice

were allowed to explore the arena freely, and spatial locations were tracked using EthoVision XT

10 (Noldus, Leesburg, VA) and CCD cameras (SuperCircuits, Austin, TX).

Operant behavior. Animals were first given access to 20 mg sweetened chow pellets in their

home cage before testing. Animals were then trained to enter the chamber to retrieve a pellet.

Each pellet was delivered 10 s after the prior pellet retrieval. After at least 2 days training and

until >30 pellets earned in a single session, animals were trained for the fixed ration 1 (FR1) task,

in which each active lever pressing was rewarded with a pellet. A new trial does not begin until

animals entered the magazine to retrieve the pellet. Retrieval was followed by a 5 s intertrial

interval, after which the levers were reactivated, indicated by a cue light. Training continued
until >40 pellets were earned in a single 60 mins session.
529
530 **Progressive ratio**. After FR1, FR3, FR5 training sessions, all mice were tested with CNO
531 treatment. For progressive ratio (PR) task, a schedule of reinforcement, each subsequent reward
532 required exponentially more lever pressing based on the formula $(5 \times e^{0.2n}) - 5$, rounded to the
533 nearest integer, where n = number of rewards earned[72]. 60 mins per session.
534
Optogenetic modulations of post-fasted food intake
Mice received 20 min laser stimulation (4×5 min, On-Off-On-Off), and then the remaining food
pallets were collected and food intake was measured. For photostimulating ChR2, a 473-nm laser
(OEM Lasers/OptoEngine) was used to generate laser pulses (10-15 mW at the tip of the fiber, 5
ms, 20 Hz) throughout the behavioral session, except when noted otherwise, controlled by a
waveform generator (Keysight). For NpHR photostimulation, a 532-nm laser (OEM
Lasers/OptoEngine) generated constant light of 8-10 mW power at each fiber tip.
542
Fiber photometry during feeding
The *Serpinb2*[+] neuronal dynamics during feeding was measured using fiber photometry.
Following injection of an AAV1-hSyn-FLEX-GCaMP7s vector into NAcSh of *Serpinb2*-Cre
mice, an optical cannula (Ø200 μm core, 0.37 numerical aperture) was implanted 100 μm above
the viral injection site. Mice were allowed to recover for 3 weeks and then subjected to
behavioral test. GCaMP fluorochrome was excited, and emission fluorescence was acquired with
the RZ10X fiber photometry system, which has built-in LED drivers, LEDs, and photosensors
(Tucker-Davis Technologies). The LEDs include 405 nm (as isosbestic control) and 465 nm (for
GCaMP excitation). Emitted light was received through the Mini Cube (Doric Lenses) and split
into two bands, 420 to 450 nm (autofluorescence) and 500 to 550 nm (GCaMP7 signal). Mice
with optical cannula were attached to recording optic cables, and the LED power at the tip of the
optical cables was adjusted to the lowest possible (~20 μW) to minimize bleaching. Mice were
placed in the 3- chamber for food preference and food consumption test. Mice behaviors were
recorded using EthoVision XT 10 (Noldus, Leesburg, VA) and CCD cameras (Super Circuits,
Austin, TX).

For the 3-chamber food seeking and food consumption, mice were fasted overnight, and were

then habituated for 10-min in the chamber that contain 2 empty food cups, after that, we put a

non-eatable object and food pellets in the two food cups, during this phase, food pellets are caged

so that mice can sense the food but are unable to eat them. After 10-min recording, we then

removed the barrier of food and mice have free-access to the food, mice eating events were then

recorded. Behavioral events, such as baseline of free moving, entering food zone, food

consumption and post eating were scored manually and synchronized with fluorescence signal

based on recorded videos. The voltage signal data stream was acquired with Synapse software

(Tucker-Davis Technologies) and were exported, filtered, and analyzed with custom-written

Matlab code. To calculate $\Delta F/F$, a polynomial linear fitting was applied to isosbestic signal to

align it to the GCaMP7 signal, producing a fitted isosbestic signal that was used to normalize the

GCaMP7 as follows: $\Delta F/F = (\text{GCaMP7signal} - \text{fitted isosbestic})/\text{fitted isosbestic signal}$.

Lead Contact

Further information and requests for reagents should be directed to and will be fulfilled by the

Lead Contact, Yi Zhang (yzhang@genetics.med.harvard.edu).

Data and Code Availability

The custom code that supports the findings from this study are available from the Lead Contact

upon request.

Reference

Giel, K. E. *et al.* Binge eating disorder. *Nat Rev Dis Primers* **8**, 16, doi:10.1038/s41572-022-00344-y (2022).

Kaye, W. H., Fudge, J. L. & Paulus, M. New insights into symptoms and neurocircuit function of anorexia nervosa. *Nature Reviews Neuroscience* **10**, 573-584, doi:10.1038/nrn2682 (2009).

Rossi, M. A. & Stuber, G. D. Overlapping Brain Circuits for Homeostatic and Hedonic Feeding. *Cell Metab* **27**, 42-56, doi:10.1016/j.cmet.2017.09.021 (2018).

Zhan, C. *et al.* Acute and long-term suppression of feeding behavior by POMC neurons in the brainstem and hypothalamus, respectively. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **33**, 3624-3632, doi:10.1523/JNEUROSCI.2742-12.2013 (2013).

Li, Y. *et al.* Serotonin neurons in the dorsal raphe nucleus encode reward signals. *Nat Commun* **7**, 10503, doi:10.1038/ncomms10503 (2016).

Dietrich, M. O. *et al.* AgRP neurons regulate development of dopamine neuronal plasticity and nonfood-associated behaviors. *Nature neuroscience* **15**, 1108-1110, doi:10.1038/nn.3147 (2012).

Kolb, B. & Nonneman, A. J. Prefrontal cortex and the regulation of food intake in the rat. *J Comp Physiol Psychol* **88**, 806-815, doi:10.1037/h0076397 (1975).

Davidson, T. L. *et al.* Contributions of the hippocampus and medial prefrontal cortex to energy and body weight regulation. *Hippocampus* **19**, 235-252, doi:10.1002/hipo.20499 (2009).

Pecina, S. & Berridge, K. C. Hedonic hot spot in nucleus accumbens shell: where do mu-opioids cause increased hedonic impact of sweetness? *The Journal of neuroscience : the official journal of the Society for Neuroscience* **25**, 11777-11786, doi:10.1523/JNEUROSCI.2329-05.2005 (2005).

Thoeni, S., Loureiro, M., O'Connor, E. C. & Luscher, C. Depression of Accumbal to Lateral Hypothalamic Synapses Gates Overeating. *Neuron* **107**, 158-172 e154, doi:10.1016/j.neuron.2020.03.029 (2020).

Chen, R., Wu, X., Jiang, L. & Zhang, Y. Single-Cell RNA-Seq Reveals Hypothalamic Cell Diversity. *Cell reports* **18**, 3227-3241, doi:10.1016/j.celrep.2017.03.004 (2017).

Li, Y. *et al.* Hypothalamic Circuits for Predation and Evasion. *Neuron* **97**, 911-924 e915, doi:10.1016/j.neuron.2018.01.005 (2018).

Waterson, M. J. & Horvath, T. L. Neuronal Regulation of Energy Homeostasis: Beyond the Hypothalamus and Feeding. *Cell Metab* **22**, 962-970, doi:10.1016/j.cmet.2015.09.026 (2015).

Wren, A. M. *et al.* Ghrelin enhances appetite and increases food intake in humans. *J Clin Endocrinol Metab* **86**, 5992, doi:10.1210/jcem.86.12.8111 (2001).

Myers, M. G., Cowley, M. A. & Munzberg, H. Mechanisms of leptin action and leptin resistance. *Annual review of physiology* **70**, 537-556, doi:10.1146/annurev.physiol.70.113006.100707 (2008).

Cowley, M. A. *et al.* Leptin activates anorexigenic POMC neurons through a neural network in the arcuate nucleus. *Nature* **411**, 480-484, doi:10.1038/35078085 (2001).

Leininger, G. M. *et al.* Leptin acts via leptin receptor-expressing lateral hypothalamic neurons to modulate the mesolimbic dopamine system and suppress feeding. *Cell Metab* **10**, 89-98, doi:10.1016/j.cmet.2009.06.011 (2009).

Loureiro, M. *et al.* Social transmission of food safety depends on synaptic plasticity in the prefrontal cortex. *Science* **364**, 991-995, doi:10.1126/science.aaw5842 (2019).

O'Connor, E. C. *et al.* Accumbal D1R Neurons Projecting to Lateral Hypothalamus Authorize Feeding. *Neuron* **88**, 553-564, doi:10.1016/j.neuron.2015.09.038 (2015).

Kreitzer, A. C. & Malenka, R. C. Striatal plasticity and basal ganglia circuit function. *Neuron* **60**, 543-554, doi:10.1016/j.neuron.2008.11.005 (2008).

Gerfen, C. R. & Surmeier, D. J. Modulation of striatal projection systems by dopamine. *Annual review of neuroscience* **34**, 441-466, doi:10.1146/annurev-neuro-061010-113641 (2011).

22 Matikainen-Ankney, B. A. *et al.* Enhanced food motivation in obese mice is controlled by D1R
expressing spiny projection neurons in the nucleus accumbens. *bioRxiv*, 2022.2001.2012.476057,
doi:10.1101/2022.01.12.476057 (2022).
23 Tan, B. *et al.* Dynamic processing of hunger and thirst by common mesolimbic neural ensembles.
Proceedings of the National Academy of Sciences of the United States of America **119**,
e2211688119, doi:10.1073/pnas.2211688119 (2022).
24 Chen, R. *et al.* Decoding molecular and cellular heterogeneity of mouse nucleus accumbens.
Nature neuroscience **24**, 1757-1771, doi:10.1038/s41593-021-00938-x (2021).
25 Sylwestrak, E. L. *et al.* Cell-type-specific population dynamics of diverse reward computations.
Cell **185**, 3568-3587 e3527, doi:10.1016/j.cell.2022.08.019 (2022).
26 Osterhout, J. A. *et al.* A preoptic neuronal population controls fever and appetite during sickness.
Nature **606**, 937-944, doi:10.1038/s41586-022-04793-z (2022).
27 Bhattacherjee, A. *et al.* Cell type-specific transcriptional programs in mouse prefrontal cortex
during adolescence and addiction. *Nat Commun* **10**, 4169, doi:10.1038/s41467-019-12054-3
(2019).
28 Ilanges, A. *et al.* Brainstem ADCYAP1(+) neurons control multiple aspects of sickness behaviour.
Nature **609**, 761-771, doi:10.1038/s41586-022-05161-7 (2022).
29 Moffitt, J. R. *et al.* Molecular, spatial, and functional single-cell profiling of the hypothalamic
preoptic region. *Science* **362**, doi:10.1126/science.aau5324 (2018).
30 Bond, C. W. *et al.* Medial Nucleus Accumbens Projections to the Ventral Tegmental Area
Control Food Consumption. *The Journal of neuroscience : the official journal of the Society for
Neuroscience* **40**, 4727-4738, doi:10.1523/JNEUROSCI.3054-18.2020 (2020).
31 Durst, M., Konczol, K., Balazsa, T., Eyre, M. D. & Toth, Z. E. Reward-representing D1-type
neurons in the medial shell of the accumbens nucleus regulate palatable food intake. *Int J Obes
(Lond)* **43**, 917-927, doi:10.1038/s41366-018-0133-y (2019).
32 Zhang, C., Chen, R. & Zhang, Y. Accurate inference of genome-wide spatial expression with
iSpatial. *Sci Adv* **8**, eabq0990, doi:10.1126/sciadv.abq0990 (2022).
33 Alexander, G. M. *et al.* Remote control of neuronal activity in transgenic mice expressing
evolved G protein-coupled receptors. *Neuron* **63**, 27-39, doi:10.1016/j.neuron.2009.06.014 (2009).
34 Kirouac, G. J. The Paraventricular Nucleus of the Thalamus as an Integrating and Relay Node in
the Brain Anxiety Network. *Front Behav Neurosci* **15**, 627633, doi:10.3389/fnbeh.2021.627633
(2021).
35 West, E. A. & Carelli, R. M. Nucleus Accumbens Core and Shell Differentially Encode Reward-
Associated Cues after Reinforcer Devaluation. *The Journal of neuroscience : the official journal
of the Society for Neuroscience* **36**, 1128-1139, doi:10.1523/JNEUROSCI.2976-15.2016 (2016).
36 Di Chiara, G. *et al.* Dopamine and drug addiction: the nucleus accumbens shell connection.
Neuropharmacology **47 Suppl 1**, 227-241, doi:10.1016/j.neuropharm.2004.06.032 (2004).
37 Watabe-Uchida, M., Zhu, L., Ogawa, S. K., Vamanrao, A. & Uchida, N. Whole-brain mapping of
direct inputs to midbrain dopamine neurons. *Neuron* **74**, 858-873,
doi:10.1016/j.neuron.2012.03.017 (2012).
38 Pardo-Garcia, T. R. *et al.* Ventral Pallidum Is the Primary Target for Accumbens D1 Projections
Driving Cocaine Seeking. *The Journal of neuroscience : the official journal of the Society for
Neuroscience* **39**, 2041-2051, doi:10.1523/JNEUROSCI.2822-18.2018 (2019).
39 Brog, J. S., Salyapongse, A., Deutch, A. Y. & Zahm, D. S. The patterns of afferent innervation of
the core and shell in the "accumbens" part of the rat ventral striatum: immunohistochemical
detection of retrogradely transported fluoro-gold. *J Comp Neurol* **338**, 255-278,
doi:10.1002/cne.903380209 (1993).
40 Conte, W. L., Kamishina, H. & Reep, R. L. The efficacy of the fluorescent conjugates of cholera
toxin subunit B for multiple retrograde tract tracing in the central nervous system. *Brain Struct
Funct* **213**, 367-373, doi:10.1007/s00429-009-0212-x (2009).

41   Berthoud, H. R. & Munzberg, H. The lateral hypothalamus as integrator of metabolic and
environmental needs: from electrical self-stimulation to opto-genetics. *Physiol Behav* **104**, 29-39,
doi:10.1016/j.physbeh.2011.04.051 (2011).
42   Sweet, D. C., Levine, A. S., Billington, C. J. & Kotz, C. M. Feeding response to central orexins.
Brain research **821**, 535-538, doi:10.1016/s0006-8993(99)01136-1 (1999).
43   Qu, D. *et al.* A role for melanin-concentrating hormone in the central regulation of feeding
behaviour. *Nature* **380**, 243-247, doi:10.1038/380243a0 (1996).
44   Kopp, W. *et al.* Low leptin levels predict amenorrhea in underweight and eating disordered
females. *Mol Psychiatry* **2**, 335-340, doi:10.1038/sj.mp.4000287 (1997).
45   Li, Z., Kelly, L., Heiman, M., Greengard, P. & Friedman, J. M. Hypothalamic Amylin Acts in
Concert with Leptin to Regulate Food Intake. *Cell Metab* **23**, 945,
doi:10.1016/j.cmet.2016.04.014 (2016).
46   Jennings, J. H., Rizzi, G., Stamatakis, A. M., Ung, R. L. & Stuber, G. D. The inhibitory circuit
architecture of the lateral hypothalamus orchestrates feeding. *Science* **341**, 1517-1521,
doi:10.1126/science.1241812 (2013).
47   Wickersham, I. R., Finke, S., Conzelmann, K. K. & Callaway, E. M. Retrograde neuronal tracing
with a deletion-mutant rabies virus. *Nat Methods* **4**, 47-49, doi:10.1038/nmeth999 (2007).
48   Friedman, J. M. The function of leptin in nutrition, weight, and physiology. *Nutr Rev* **60**, S1-14;
discussion S68-84, 85-17, doi:10.1301/002966402320634878 (2002).
49   Morton, G. J., Cummings, D. E., Baskin, D. G., Barsh, G. S. & Schwartz, M. W. Central nervous
system control of food intake and body weight. *Nature* **443**, 289-295, doi:10.1038/nature05026
(2006).
50   Myers, M. G., Jr., Munzberg, H., Leinninger, G. M. & Leshan, R. L. The geometry of leptin
action in the brain: more complicated than a simple ARC. *Cell Metab* **9**, 117-123,
doi:10.1016/j.cmet.2008.12.001 (2009).
51   Cohen, P. *et al.* Selective deletion of leptin receptor in neurons leads to obesity. *J Clin Invest* **108**,
1113-1121, doi:10.1172/JCI13914 (2001).
52   de Luca, C. *et al.* Complete rescue of obesity, diabetes, and infertility in db/db mice by neuron-
specific LEPR-B transgenes. *J Clin Invest* **115**, 3484-3493, doi:10.1172/JCI24059 (2005).
53   Kim, K. S. *et al.* Enhanced hypothalamic leptin signaling in mice lacking dopamine D2 receptors.
The Journal of biological chemistry **285**, 8905-8917, doi:10.1074/jbc.M109.079590 (2010).
54   Floresco, S. B. The nucleus accumbens: an interface between cognition, emotion, and action.
Annu Rev Psychol **66**, 25-52, doi:10.1146/annurev-psych-010213-115159 (2015).
55   Johnson, P. M. & Kenny, P. J. Dopamine D2 receptors in addiction-like reward dysfunction and
compulsive eating in obese rats. *Nature neuroscience* **13**, 635-641, doi:10.1038/nn.2519 (2010).
56   Wang, G. J. *et al.* Brain dopamine and obesity. *Lancet* **357**, 354-357, doi:10.1016/s0140-
6736(00)03643-6 (2001).
57   Gerfen, C. R. Segregation of D1 and D2 dopamine receptors in the striatal direct and indirect
pathways: An historical perspective. *Front Synaptic Neurosci* **14**, 1002960,
doi:10.3389/fnsyn.2022.1002960 (2022).
58   Zhao, Z.-d. *et al.* A molecularly defined D1 medium spiny neuron subtype negatively regulates
cocaine addiction. *Science Advances* **8**, eabn3552 (2022).
59   Zhou, K. *et al.* Reward and aversion processing by input-defined parallel nucleus accumbens
circuits in mice. *Nat Commun* **13**, 6244, doi:10.1038/s41467-022-33843-3 (2022).
60   Mickelsen, L. E. *et al.* Single-cell transcriptomic analysis of the lateral hypothalamic area reveals
molecularly distinct populations of inhibitory and excitatory neurons. *Nature neuroscience* **22**,
642-656, doi:10.1038/s41593-019-0349-8 (2019).
61   Thoeni, S., Loureiro, M., O'Connor, E. C. & Lüscher, C. Depression of Accumbal to Lateral
Hypothalamic Synapses Gates Overeating. *Neuron*, doi:10.1016/j.neuron.2020.03.029 (2020).
62   Elmquist, J. K., Bjorbaek, C., Ahima, R. S., Flier, J. S. & Saper, C. B. Distributions of leptin
receptor mRNA isoforms in the rat brain. *J Comp Neurol* **395**, 535-547 (1998).

63   de Vrind, V. A. J., Rozeboom, A., Wolterink-Donselaar, I. G., Luijendijk-Berg, M. C. M. & Adan,
R. A. H. Effects of GABA and Leptin Receptor-Expressing Neurons in the Lateral Hypothalamus
on Feeding, Locomotion, and Thermogenesis. *Obesity (Silver Spring)* **27**, 1123-1132,
doi:10.1002/oby.22495 (2019).
64   Jo, Y.-H., Chen, Y.-J. J., Chua, S. C., Talmage, D. A. & Role, L. W. Integration of
endocannabinoid and leptin signaling in an appetite-related neural circuit. *Neuron* **48**, 1055-1066
(2005).
65   Kirouac, G. J. Placing the paraventricular nucleus of the thalamus within the brain circuits that
control behavior. *Neurosci Biobehav Rev* **56**, 315-329, doi:10.1016/j.neubiorev.2015.08.005
(2015).
66   Millan, E. Z., Ong, Z. & McNally, G. P. in *Progress in Brain Research* Vol. 235 (eds Tanya
Calvey & William M. U. Daniels) 113-137 (Elsevier, 2017).
67   Otis, J. M. *et al.* Paraventricular Thalamus Projection Neurons Integrate Cortical and
Hypothalamic Signals for Cue-Reward Processing. *Neuron* **103**, 423-431 e424,
doi:10.1016/j.neuron.2019.05.018 (2019).
68   Labouebe, G., Boutrel, B., Tarussio, D. & Thorens, B. Glucose-responsive neurons of the
paraventricular thalamus control sucrose-seeking behavior. *Nature neuroscience* **19**, 999-1002,
doi:10.1038/nn.4331 (2016).
69   Kessler, S. *et al.* Glucokinase neurons of the paraventricular nucleus of the thalamus sense
glucose and decrease food consumption. *iScience* **24**, 103122, doi:10.1016/j.isci.2021.103122
(2021).
70   Krause, M., German, P. W., Taha, S. A. & Fields, H. L. A pause in nucleus accumbens neuron
firing is required to initiate and maintain feeding. *The Journal of neuroscience : the official
journal of the Society for Neuroscience* **30**, 4746-4756, doi:10.1523/JNEUROSCI.0197-10.2010
(2010).
71   Kelley, A. E. Ventral striatal control of appetitive motivation: role in ingestive behavior and
reward-related learning. *Neurosci Biobehav Rev* **27**, 765-776,
doi:10.1016/j.neubiorev.2003.11.015 (2004).
72   Richardson, N. R. & Roberts, D. C. Progressive ratio schedules in drug self-administration
studies in rats: a method to evaluate reinforcing efficacy. *Journal of neuroscience methods* **66**, 1-
11, doi:10.1016/0165-0270(95)00153-0 (1996).
762

Acknowledgements

The authors thank Dr. Chao Zhang for his help in generating Fig. 1a; Dr. Jeffrey M. Friedman

for discussion on some experiments; Dr. Aritra Bhattachejee for critical reading of the

manuscript. We thank the Mouse Behavior Core of Harvard Medical School and its director Dr.

Barbara Caldarone for her help. This project was partly supported by 1R01DA042283,

1R01DA050589, and HHMI. Y.Z. is an investigator of the Howard Hughes Medical Institute.

This article is subject to HHMI's Open Access to Publications policy. HHMI lab heads have

previously granted a nonexclusive CC BY 4.0 license to the public and a sublicensable license to

HHMI in their research articles. Pursuant to those licenses, the author-accepted manuscript of

Author information

Authors and Affiliations

Howard Hughes Medical Institute, Boston Children's Hospital, Boston, MA, USA

Yiqiong Liu, Zheng-dong Zhao, Guoguang Xie, Renchao Chen and Yi Zhang

Program in Cellular and Molecular Medicine, Boston Children's Hospital, Boston, MA, USA

Yiqiong Liu, Zheng-dong Zhao, Guoguang Xie, Renchao Chen and Yi Zhang

Division of Hematology/Oncology, Department of Pediatrics, Boston Children's Hospital, Boston, MA, USA

Yiqiong Liu, Zheng-dong Zhao, Guoguang Xie, Renchao Chen and Yi Zhang

Department of Genetics, Harvard Medical School, Boston, MA, USA

Yi Zhang

Harvard Stem Cell Institute, WAB-149G, 200 Longwood Avenue, Boston, MA, USA

Yi Zhang

Contributions

Y.Z. conceived the project; Y.L., and Y.Z. designed the experiments; Y.L. performed most of the experiments. Z.-D.Z. helped with the fiber photometry. G.X. helped with the catheter administration. R.C. initiated the Serpinb2-Cre mouse generation. Y.L., Z.-D.Z., R.C. and Y.Z. interpreted the data; Y.L. and Y.Z. wrote the manuscript with input from Z.-D.Z and R. C.

Ethics declarations

Competing interests

The authors declare no competing interests.

Figure Legends

Fig. 1: NAcSh *Serpinb2*[+] neurons and NAc *Drd1*[+]-MSNs are activated in refeeding process

a, Inferred spatial expression patterns from MERFISH database of *Tac2*, *Serpinb2* and *Ubp1* whose expression is highly enriched in medial dorsal NAc shell. Expression level is color-coded. Doted line circle anterior commissure olfactory limb (aco) and NAc core. The dorsal-ventral (DV) and medial-lateral (ML) axes are indicated.

b, Left, tSNE plot showing the 8 NAc D1-MSN subtypes. Middle and right panels indicate the expression of *Tac2* and *Serpinb2* in the NAc D1-MSNs, respectively. Expression level is color-coded.

c, RNA *in situ* hybridization showing *Tac2* and *Seprinb2* expression in the medial part of the NAc shell. Scale bar: 500 μm (left), 20 μm (right).

d, Schematic representation of the experimental design for the ad libitum, fast and refed groups.

e, Coexpression of *Serpinb2* mRNA with *cFos* mRNA in NAc of ad libitum, fasted and refed states. Representative images showing the colocalization of *cFos* (green), *Serpinb2* (red) and *Tac2* (magenta) expressing neurons. Scale bar: 50 μm.

f, The average number of *cFos*[+] neurons in the dorsal medial NAcSh at ad libitum, fasted and refed states. (n = 5 sections from three mice of each group, one-way ANOVA, with Tukey's multiple comparisons).

g, The percentages of activated *Serpinb2*-expressing neurons in the total *Serpinb2* neurons under different feeding states (n = 5 sections from three mice of each group, one-way ANOVA, with Tukey's multiple comparisons).

h, The percentages of activated *Tac2*-expressing neurons in the total *Tac2* neurons under different feeding states (n = 5 sections from three mice of each group, one-way ANOVA, with Tukey's multiple comparisons). All error bars represent mean ± SEM. ns, not significant, *P < 0.05.

Fig. 2: The activity of *Serpinb2*[+] neurons respond to feeding states

a, Left, illustration of light pathways of fiber photometry. Right, schematic illustration of the GCaMP7s injection and optic cannula implantation.

b, c, Validation of GCaMP7s expression and implantation of optic cannula in *Serpinb2*-Cre mice

(b) or *Drd1*-Cre mice (c) (left). Representative trace of real-time monitoring of *Serpinb2*$^+$

neurons (b) or *Drd1*$^+$ neurons (c) (right) during feeding process. Scale bar: 200 μm.

d, Ca^{2+} signals at different phases of feeding process of fasted mice (top). Average Ca^{2+} signal at

different feeding phases of the *Serpinb2*-Cre mice. Elevated Ca^{2+} signals were observed after

entering food zone in approaching and eating phases. Declined Ca^{2+} signals were observed

post eating. Quantification of AUC in the four phases is shown in bar graph (right, n=7).

e, The same as in panel D except *Drd1*-Cre mice were used. ***$P \leq 0.001$, **$P \leq 0.01$, *$P \leq 0.05$;

ns, $P > 0.05$, one-way ANOVA test. Data are represented as mean ± SEM.

Fig. 3: *Serpinb2*$^+$ neurons bidirectionally regulate food seek and intake in hungry state

a, Experimental scheme of the food consumption and food preference assays.

b, Total food consumption during the 3-hour test. Chemogenetic activation (hM3Dq) or

chemogenetic inhibition (hM4Di) of *Serpinb2*$^+$ neurons at ad libitum (left) or fasting state

(right). **, $p<0.01$; ns, $p>0.05$, unpaired t-test.

c, The same as in panel B except the *Tac2*-Cre (left) or *Drd1*-Cre (right) mice were used. ns,

$p>0.05$, unpaired t-test.

d, Color map encoding spatial location of a fasted mouse using the free access feeding paradigm.

e, Percentage of time that mice spent in food zone. Chemogenetic activation (hM3Dq) or

chemogenetic inhibition (hM4Di) of *Serpinb2*$^+$ neurons (left), *Tac2*$^+$ neurons (middle), and

Drd1$^+$ neurons (right). *, $p<0.05$; ns, $p>0.05$, unpaired t-test.

f, Experimental timeline of the food operant chamber assay.

g, Diagrammatic illustration of the food operant chamber paradigm. Mice were trained to press

the lever to get food; pressing the active lever is followed by the delivery of food pellet,

while pressing the inactive lever yields no outcome. The behavioral training includes

habituation phase and fixed ratio (FR) training phase. Mice received CNO injection (2 mg/kg

for the hM3Dq group and 5 mg/kg for the hM4Di group) 15 mins before they were placed

into the operant chamber to start the FR and progressive ratio (PR) tests.

h, Results of FR=5 test. The total number of active lever pressing (left) and total number of

reward (right) after chemogenetic manipulations. **, $p<0.01$; *, $p<0.05$; ns, $p>0.05$; one-way

ANOVA test.

i, Results of PR=5 test. The total number of active lever pressing (left) and total number of

reward (right) after chemogenetic manipulations. *, p<0.05; ns, p>0.05; one-way ANOVA

test.

Data are represented as mean ± SEM.

Fig. 4: *Serpinb2*$^+$ neurons mediate food intake via LH projection

a, Diagram of antegrade tracing of the NAc *Serpinb2*$^+$ neurons with ChR2-EYFP (left), the

expression of DIO-ChR2-EYFP in the NAcSh (2^{nd} panel), and *Serpinb2*$^+$ neuron projection

to LH (the right 3 panels). Scale bar: 100 μm.

b, Diagram (left) and image (right) of retrograde tracing of LH neurons with CTB 647, with the

Serpinb2$^+$ neurons labeled by DIO-mCherry. Scale bar: 50 μm.

c, Percentage of *Serpinb2*$^{+\ LH}$ projecting neurons over the total mCherry labeled *Serpinb2*$^+$

neurons.

d, Diagram illustrate the indicated AAV injection into NAc and optic cannulas implantation in

LH area (left), and histology validating virus expression and cannula implantation site (right).

Scale bar: 200 μm.

e, Optogenetic activation (left) or inhibition (right) of *Serpinb2*$^+$ NAc→LH neurons respectively

increased or decreased food intake.

Data in (e) is presented as mean ± SEM. TH, thalamus, CP, caudoputamen, AHNc: anterior

hypothalamic nucleus, central part, HPF, hippocampal formation, VL, lateral ventricle. ***P

≤ 0.001; *P ≤ 0.05; ns, P > 0.05, unpaired t-test.

Fig. 5: *Serpinb2*$^+$ neurons project to LH LepR$^+$ GABA$^+$ neurons and receive input related

to energy homeostasis

a, Diagram indicating the MCH$^+$, Orexin-A$^+$, GABA$^+$ and LepR$^+$ neurons in LH.

b, Images (left two panels) showing colocalization of *Serpinb2*$^+$ neuron terminals (green) with

MCH$^+$, Orexin-A$^+$ neurons in LH, and their quantifications (right panel). Scale bar, 50 μm.

Percentage = eYFP$^+$MCH$^+$/eYFP$^+$DAPI$^+$ or eYFP$^+$Orexin-A$^+$/eYFP$^+$DAPI$^+$

c, Images showing the colocalization of *Serpinb2*$^+$ neuron terminals (green) with GABA$^+$ (red)

or LepR$^+$ neurons (red) as indicated, as well as their quantifications (right panel). Scale bar,

50 μm. Percentage = eYFP$^+$GABA$^+$/eYFP$^+$DAPI$^+$ or eYFP$^+$LepR$^+$/eYFP$^+$DAPI$^+$

d, Quantification of cFos[+] cells of different brain regions in ad libitum and refeed status. (n = 3

sections from three mice of each group, unpaired t-test)

e, Schematic presentation of modified rabies tracing (left) and representative image confirming

the expression of the indicated proteins at the injection site (right); scale bar, 100 μm

(left); 20μm (right).

f, Representative images showing the brain areas with positive signals indicating these regions

have neurons projecting to *Serpinb2*[+] neurons. Scale bar: 100 μm, arrow heads indicate

neurons.

g, Diagram illustrating brain regions upstream and downstream of *Serpinb2*[+] neurons.

ACAd: Anterior cingulate area, dorsal part; fa, corpus callosum, anterior forceps; LPO: lateral

preoptic area; SI: substantia innominata; PVT: paraventricular nucleus of the thalamus; VMH:

ventromedial hypothalamic nucleus; LH: lateral hypothalamus; V3: third ventricle.

Fig. 6: Modulating *Serpinb2*[+] neuron activity can overcome leptin effect and alter

bodyweight

a, Diagram showing bilateral cannula implantation in LH for leptin delivery (left). CNO delivery

was achieved via i.p. injection. Results of total food consumption in 3 hours by fasted mice

with different doses of leptin administration. Scale bar: 500 μm.

b, Same as panel A except food consumption is quantified under different conditions with or

without *Serpinb2*[+] neuron activation in the presence or absence of 1 μg of leptin delivery.

***P ≤ 0.001, **P ≤ 0.01, *P ≤ 0.05; ns, P > 0.05, unpaired t-test.

c, FISH and quantification verify *Serpinb2*[+] neuron ablation after AAV-DIO-taCasp3 injection.

Scale bar: 100 μm. ***P ≤ 0.001, unpaired t-test.

d, *Serpinb2*[+] neuron ablation decreased 3 hours total food consumption by mice. ***P ≤ 0.001,

unpaired t-test.

e, *Serpinb2*[+] neuron ablation has a long-time effect on bodyweight loss. ***P ≤ 0.001, **P ≤

0.01, *P ≤ 0.05; ns, P > 0.05, unpaired t-test.

Data are presented as means ± SEM.

Extended Data Figure Legends

Extended Data Fig. 1: Generation and validation of *Serpinb2*-Cre line

a, Diagrams showing the targeting strategy.

b, Genotyping by PCR. Homozygotes: 413 bp. Heterozygotes: 413 bp/772 bp.

c, Left, in situ hybridization (ISH) data of *Serpinb2* from Allen Brain Atlas. Scale bar: 1000 μm.

Middle, colocalization of *Serpinb2* RNA (red), AAV-DIO-ChR2-eYFP (green) and DAPI

(blue). Scale bar: 50 μm. Right, quantification of $Serpinb2^+$ and $eYFP^+$ neurons among all

$Serpinb2^+$ neurons.

Extended Data Fig. 2: $Serpinb2^+$ neuron activity during different phases of feeding in ad

libitum mice

a, Ca^{2+} signals at different phases of feeding process of ad libitum mice (top). Average Ca^{2+}

signal at different feeding phases of the *Serpinb2*-Cre mice. Elevated Ca^{2+} signals were

observed after entering food zone in the approaching and eating phases. Declined Ca^{2+}

signals were observed post eating. Quantification of AUC in the four phases is shown in bar

graph (right, n=7 mice).

b, The same as in panel D except *Drd1*-Cre mice were used.

**P ≤ 0.01, *P ≤ 0.05; ns, P > 0.05, one-way ANOVA test. Data are represented as mean ± SEM.

Extended Data Fig. 3: Validation of chemogenetic manipulation

a, Experimental scheme of chemogenetic manipulation.

b, Validation of virus expression in *Serpinb2*-Cre, *Tac2*-Cre and *Drd1*-Cre mice. Scale bar: 500

μm.

c, *cFos* induction after intraperitoneal injection of ligand CNO in mCherry-expressing, hM3Dq-

mCherry-expressing and hM4Di-mCherry-expressing mice. The ratio of $cFos^+/mCherry^+$

cells in all $mCherry^+$ cells was calculated and shown on the right panel. Scale bar: 100 μm. *,

p<0.05; **, p<0.01, one-way ANOVA test. Data are represented as mean ± SEM.

Extended Data Fig. 4: $Serpinb2^+$ neuronal activity does not affect anxiety or drug seeking

behavior

a, b, Open field test for the effect of *Sepinb2*[+] neuronal activation (hM3Dq) (a) or inhibition

(hM4Di) (b) on the total distance traveled in the 1-hour post-treatment period after

chemogenetic manipulation of *Sepinb2*[+] neurons (left) or the distance traveled in 5-min time

bin (right).

ns, p>0.05, left, unpaired t-test; right, two-way ANOVA.

c, Left, illustration of the two-chamber cocaine-CPP paradigm. Right, cocaine-CPP with

chemogenetic activation (hM3Dq) or inhibition (hM4Di) of *Sepinb2*[+] neurons. CPP scores

were calculated by subtracting the time spent in the preconditioning phase from the time

spent in the postconditioning phase. ns, p>0.05, unpaired t-test.

d, e, Elevated plus maze test for the effect of *Sepinb2*[+] neuronal activation (hM3Dq) (d) or

inhibition (hM4Di) (e) on the time spent (left) or distance traveled (right) in open arm and

closed arm of the 5-min post-treatment period after chemogenetic manipulation of *Sepinb2*[+]

neurons. ns, p>0.05, unpaired t-test.

Data are represented as mean ± SEM.

Extended Data Fig. 5: cFos staining of different brain regions from mice under Ad libitum

and refeed states

Shown are cFos FISH in different brain regions listed below:

ACB: Nucleus accumbens; aco: anterior commissure, olfactory limb; OT:

Olfactory tubercle; V3: third ventricle; ARH: Arcuate hypothalamic nucleus; AHNp: Anterior

hypothalamic nucleus, posterior part; VMH: ventromedial hypothalamic nucleus; fx: columns of

the fornix; LH: lateral hypothalamus; PVT: paraventricular nucleus of the thalamus; sm: stria

medullaris; BLA: Basolateral amygdala; LA: Lateral amygdala; RE: Nucleus of reuniens; DMH:

Dorsomedial nucleus of the hypothalamus; ZI: Zona incerta. Scale bar: 100 μm.

Extended Data Fig. 6: Brain regions that do not innervate *Serpinb2*[+] neurons

Shown are immunostaining of the various brain regions listed below:

PL: Prelimbic area; IL: Infralimibic area; fa: corpus callosum, anterior forceps; ccg: genu of

corpus

callosum; LSr: Lateral septal nucleus, rostral part; VL: lateral ventricle; ACB: Nucleus

accumbens; aco: anterior commissure, olfactory limb; OT: Olfactory tubercle; ADP:

Anterodorsal preoptic nucleus; BST: Bed nuclei of the stria terminalis; V3: third ventricle; ARH:

Arcuate hypothalamic nucleus; AHNp: Anterior hypothalamic nucleus, posterior part; VMH:

ventromedial hypothalamic nucleus; fx: columns of the fornix; LH: lateral hypothalamus; PVT:

paraventricular nucleus of the thalamus; sm: stria medullaris; BLA: Basolateral amygdala; LA:

Lateral amygdala; RE: Nucleus of reuniens; DMH: Dorsomedial nucleus of the hypothalamus;

ZI: Zona incerta; DG: Dentate gyrus; CA1: field CA1; CA3: field CA3; PAG: Periaqueductal

gray ; APN: Anterior pretectal nucleus; HPF: Hippocampal formation. Scale bar: 100 μm

Fig. 1: NAcSh *Serpinb2*+ neurons and NAc *Drd1*+-MSNs are activated in refeeding process.

Fig. 2: The activity of *Serpinb2*+ neurons respond to feeding states.

Fig. 3: *Serpinb2*⁺ neurons bidirectionally regulate food seek and intake in hungry state.

Fig. 4: *Serpinb2*⁺ neurons mediate food intake via LH projection.

Fig. 5: *Serpinb2*⁺ neurons project to LH LepR⁺ GABA⁺ neurons and receive input related to energy homeostasis.

Fig. 6: Modulating *Serpinb2*⁺ neuron activity can overcome leptin effect and alter bodyweight.

Extended Data Fig. 1: Generation and validation of *Serpinb2*-Cre line

a

Wildtype allele

Targeting vector

Targeted allele

Legends

— Homology arm   Exon   UTR   2A   Cre

b

c

Serpinb2

Serpinb2
DIO-ChR2-eYFP
DAPI

% Serpinb2⁺ eYFP⁺/ Serpinb2⁺

Extended Data Fig. 2: *Serpinb2*[+] **neuron activity during different phases of feeding in ad libitum mice**

a

b

Extended Data Fig. 3: Validation of chemogenetic manipulation

Extended Data Fig. 4: *Serpinb2*+ neuronal activity does not affect anxiety or drug seeking behavior

Extended Data Fig. 5: cFos staining of different brain regions from mice under Ad libitum and refeed states

Extended Data Fig. 6: Brain regions that do not innervate *Serpinb2*+ neurons

cFos DAPI