Title: Engineered Immunogens to Elicit Antibodies Against Conserved Coronavirus Epitopes

Authors: Brenda Kapingidza[1,2,#], Daniel J. Marston[1,2,#], Caitlin Harris[1,2], Daniel Wrapp[1,2], Kaitlyn Winters[1,2], Dieter Mielke[3,4], Lu Xiaozhi[1,2], Qi Yin[1,2], Andrew Foulger[1,2], Rob Parks[1,2], Maggie Barr[1,2], Amanda Newman[1,2], Alexandra Schäfer[5], Amanda Eaton[3], Justine Mae Flores[1,2], Austin Harner[1,2], Nicholas J Cantazaro Jr.[5], Michael L Mallory[5], Melissa D Mattocks[5], Christopher Beverly[1,2], Brianna Rhodes[1,2], Katayoun Mansouri[1], Elizabeth Van Itallie[1,2], Pranay Vure[1,2], Brooke Manness[3], Taylor Keyes[3], Sherry Stanfield-Oakley[3], Christopher W. Woods[1,2,7], Elizabeth A. Petzold[7], Emmanuel B Walter[1,8], Kevin Wiehe[1,2], Robert J. Edwards[1,2], David Montefiori[3], Guido Ferrari[1,3,4,6], Ralph Baric[5], Derek W. Cain[1,2], Kevin O. Saunders[1,3,6,9], Barton F. Haynes[1,2,9], Mihai L. Azoitei[1,2,*]

Affiliations:

[1]Duke Human Vaccine Institute, Duke University, Durham, NC

[2]Department of Medicine, Duke University, Durham, NC

[3]Department of Surgery, Duke University, Durham, NC

[4]Center for Human Systems Immunology, Duke University, Durham, NC, USA;

[5]Department of Epidemiology, University of North Carolina at Chapel Hill, Chapel Hill, NC

[6]Department of Molecular Genetics and Microbiology, Duke University, Durham, NC

[7]Center for Infectious Diseases and Diagnostic Innovation, Duke University Medical Center, Durham, NC

[8]Department of Pediatrics, Duke University, Durham NC

[9]Department of Immunology, Duke University, Durham, NC

[#]These authors contributed equally to this work.

[*] Corresponding author. Email: mihai.azoitei@duke.edu

Abstract:

Immune responses to SARS-CoV-2 primarily target the receptor binding domain of the

spike protein, which continually mutates to escape acquired immunity. Other regions in the spike

S2 subunit, such as the stem helix and the segment encompassing residues 815-823 adjacent to

the fusion peptide, are highly conserved across sarbecoviruses and are recognized by broadly

reactive antibodies, providing hope that vaccines targeting these epitopes could offer protection

against both current and emergent viruses. Here we employed computational modeling to design

scaffolded immunogens that display the spike 815-823 peptide and the stem helix epitopes

without the distracting and immunodominant RBD. These engineered proteins bound with high

affinity and specificity to the mature and germline versions of previously identified broadly

protective human antibodies. Epitope scaffolds interacted with both sera and isolated monoclonal

antibodies with broadly reactivity from individuals with pre-existing SARS-CoV-2 immunity. When

used as immunogens, epitope scaffolds elicited sera with broad betacoronavirus reactivity and

protected as "boosts" against live virus challenge in mice, illustrating their potential as

components of a future pancoronavirus vaccine.

Main Text:

The majority of immune responses elicited against SARS-CoV-2, by either natural infection, vaccination or a combination of both, are focused on the receptor binding domain (RBD)[1-4]. However, deep scanning mutagenesis studies[5] and the emergence of novel variants that evade pre-existing immunity have revealed the plastic nature of the RBD[2,6], suggesting that next-generation coronavirus vaccines designed to protect against both current and emergent CoVs will need to induce antibodies against CoV spike regions beyond the RBD.

Two regions in the S2 subunit of the SARS-CoV-2 spike, the segment encompassing residues 815-823 adjacent to the fusion peptide and the stem helix domain (residues 1145-1156), show high conservation across diverse coronaviruses. Isolated antibodies that target these sites cross-react with both alpha and beta human coronaviruses, and protect by either neutralization, Fc receptor mediated mechanisms or both[7-10]. For example, antibodies DH1058, DH1294, VN01H1, Cov44-79, and Cov44-62, bind to an epitope located around SARS-CoV-2 spike residues 815-823 and recognize all seven human CoVs[7,8,11,12]. Antibodies against the stem helix region, such as S2P6 and CC40.8[9,10,13,14], bound spike glycoproteins representative of all sarbecovirus clades, and protected against viral challenge by inhibiting S-mediated membrane fusion. Despite their attractiveness as vaccine targets, humoral responses against the stem helix or the spike 815-823 epitopes are not robustly induced by existing vaccines or by natural infection[7-10]. This is likely due to a combination of factors, including the occlusion of these epitopes on the pre-fusion spike, requiring ACE2 binding for exposure[8], as well as the presence of the prominently displayed, immunodominant RBD domain. Given their ability to recognize diverse coronaviruses, immunogens that induce strong responses against the spike 815-823 region and the stem helix could be a key component of a future "pan-coronavirus" vaccine offering broad protection against both currently circulating and emergent coronaviruses.

For the design of immunogens that expose occluded epitopes in novel molecular contexts,

we and others previously described the development of "epitope scaffold" proteins[15-17]. These

molecules are engineered by transplanting the structure of the antibody-bound epitopes from viral

molecules onto unrelated protein scaffolds. Unlike peptide-based immunogens that can sample

diverse epitope conformations, epitope scaffolds are meant to present only the antibody-bound

conformation of the target epitope on their surface, which typically leads to higher antibody affinity

and the elicitation of antibodies specific for the structure of the target epitope[18,19]. Here, we applied

epitope grafting to design stem helix or spike 815-823 epitope scaffolds that strongly interact with

broadly cross-reactive antibodies against these regions of spike. These antigens were used to

isolate diverse antibodies with broad reactivity against these conserved epitopes from subjects

with pre-existing SARS-CoV-2 immunity. When used as immunogens in mice, epitope scaffolds

elicited sera with broad betacoronavirus reactivity. In a viral challenge model, boosting responses

induced by spike mRNA with stem helix epitope scaffolds offered protection against a pre-

emergent coronavirus, thus illustrating the potential of the immunogens engineered here as next-

generation vaccine candidates to preferentially boost antibodies that may protect against existing

and emerging coronaviruses.

Results

Design of epitope scaffolds that bind broadly cross-reactive antibodies against the

spike 815-823 epitope.

The region containing residues 815-823 of the SARS-CoV-2 spike, thereafter referred to

as spike[815-823], was initially thought to be part of the fusion peptide domain. However, recent

structural analysis of the post-fusion spike in a lipid environment revealed that the actual fusion

peptide is located upstream of this region (residues 867-909). While the functional role of the

spike[815-823] region remains to be determined, multiple broadly protective antibodies, such as

DH1058, DH1294, VN01H1, Cov44-79, and Cov44-62 target this site, with key contacts made

with virus residues R815, E819 and F823 (**Figure 1a**, **Supp. Fig. 1**)[7,8,11,12]. These residues are

occluded in the structure of the prefusion SARS-CoV-2 spike, likely limiting their immune

recognition (**Figure 1a**). To improve the accessibility and immune recognition of this epitope, we

developed spike[815-823] epitope scaffolds (ES) using "side chain" grafting computational methods

that we and others previously described (**Supp. Fig. 2**)[15-17]. A large library (~10,000) of small

monomeric scaffolding proteins was queried computationally to identify proteins with exposed

backbone regions that closely matched (<0.5Å backbone RMSD) the structure of the spike[815-823]

epitope (**Supp. Fig. 2**). In regions with high structural mimicry to the DH1058-bound epitope, the

epitope sequence replaced that of the parent scaffold and additional mutations were introduced

to accommodate the grafted sequence (**Supp. Fig. 2**).

106   Fifteen engineered spike[815-823] epitope scaffolds were expressed recombinantly (**Figure**

1b for representative designs), and six of them produced soluble and stable proteins that bound

DH1058 mAb by ELISA (**Supp. Fig. 3**, **Supp. Table 1**). Equilibrium dissociation constants for

mAbs DH1058 and DH1294 were determined by surface plasmon resonance (SPR) for the

designs that showed tight binding by ELISA (**Figure 1c; Supp. Fig. 4**). FP-2, FP-10, and FP-15

bound to DH1058 with picomolar affinities and to DH1294 with K_Ds of 1221nM, 0.1nM and 219nM

respectively. These values compare favorably with measured affinities of the full-length WA-2

spike to DH1058 and DH1294 mAbs (1.4nM and 25nM respectively). To probe the breadth of

their recognition, the binding of spike[815-823] epitope scaffolds was measured against antibodies

VN01H1, C77G12, VP12E7, COV44-79, and COV44-62[7,8], which were not explicitly considered

at the design stage (**Figure 1d; Supp. Fig. 5a**). Five designs bound at least 6 out of the 7

antibodies tested, with FP-12 and FP-15 binding all of them tighter than spike. In addition to the

mature antibodies, binding was also measured to their inferred unmutated common ancestors

(UCAs) or germline-reverted forms (iGLs). Spike only showed measurable binding to the DH1058

UCA, while FP-12 and FP-2 were bound by DH1058 UCA, DH1294 UCA and VP12E7 iGL (**Figure**

1d; Supp. Fig. 5b). Taken together, these data show that the engineered FP epitope scaffolds

have broad recognition of genetically diverse FP antibodies and their precursors.

DH1058 mAb bound three spike[815-823] epitope scaffolds with affinities 10-fold higher than

that of spike. To ensure that these affinity gains were primarily mediated by the grafted epitope

residues and not by contacts between the scaffold and the antibody that are not present in the

native DH1058-epitope complex, interface hotspot residues corresponding to SARS-CoV-2 spike

amino acids R815, E819, and F823 were mutated to alanine in designs FP-2, FP-10, and FP-15.

Each of the three alanine substitutions eliminated DH1058 mAb binding to a synthetic peptide

encompassing the spike[815-823] epitope, and to designs FP-2 and FP-10, demonstrating that the

grafted epitope is the major site of antibody interaction (**Figure 1e**). Two of the three substitutions

similarly eliminated DH1058 binding to FP-15, with the F823A equivalent retaining some low-level

antibody binding (**Figure 1e**).

To further confirm binding specificity and to validate the computational design process, a

high resolution (2.2Å) crystal structure of FP-15 in complex with DH1058 mAb was solved (**Figure**

1f, Supp. Table 2). The experimentally determined structure was in close agreement with that of

the computationally generated antibody-ES complex, both overall and over the epitope scaffold

only (backbone RMSD <1Å). The epitope conformation engaged by DH1058 on FP-15 was

essentially identical to that induced by the antibody on the spike[815-823] peptide (backbone

RMSD=0.3Å)[11], confirming that the antibody interacts with the epitope presented by the ES in the

same manner it binds to its natural target.

Design of epitope scaffolds to engage S2P6-like antibodies against the stem helix

region of the CoV spike.

Next, we engineered epitope scaffolds to engage antibodies against the stem helix region

of the CoV spike. Antibodies such as S2P6, DH1057.1, and CC40.8[9,10,20] bind to an epitope

located between residues 1144 and 1158, which is almost completely occluded by the

trimerization interface in the pre-fusion spike structure (**Figure 2a**)[9,10,20]. While both CC40.8 and

S2P6 target the same spike region and make critical contacts with residues Phe1148 and

Phe1156, CC40.8 makes additional interactions with Leu1145 and induces a more extended

epitope conformation upon binding (**Figure 2a**)[10]. Because the CC40.8 epitope is longer and

includes the S2P6 epitope, we first attempted to engineer epitope scaffolds that display the

antibody-bound conformation of the CC40.8 epitope, with the expectation that these molecules

would also be capable of binding to S2P6-class antibodies. However, the "side chain" grafting

protocol used for FP epitope scaffolds above, did not identify any candidate scaffolds with

exposed backbones similar to that of the CC40.8 epitope, likely due to the unusual conformation

of this epitope that adopts a distorted alpha-helical shape near the N-terminus (**Figure 2a**). In

contrast, multiple candidate epitope scaffolds that displayed the S2P6-bound conformation of the

epitope segment 1148-1156 were successfully designed computationally using side chain

grafting. This was possible because the S2P6 epitope adopts a canonical alpha-helical structure,

with most of the antibody contacting residues contained within three helical turns (**Figure 2a**).

Fifteen S2P6-targeted designs, named S2hlx-1 to 15, were chosen for recombinant

production in *E. coli* (**Figure 2a** for representative designs; **Supp. Table 1**) and seven yielded

soluble protein. In an initial ELISA screen, all seven designs bound both S2P6 and DH1057.1

mAbs comparable to or better than a synthesized SARS-CoV-2 stem helix peptide (residues

1147-1161). As expected, neither this peptide nor the epitope scaffolds bound CC40.8 mAb due

to the lack of epitope residue L1145 (**Supp. Fig. 6**). Stem helix epitope scaffolds had measured

equilibrium dissociation constants between 0.2nM and 83nM for S2P6 mAb and between 0.6nM

and 4.5uM for DH1057.1 mAb. In the same assay, the equilibrium dissociation constants of spike

for S2P6 and DH1057.1 were 0.4nM and 11.5nM respectively (**Figure 2b, c; Supp. Fig. 7 and**

8). The designs with the highest affinities, S2hlx-7 and S2hlx-15, had similar binding of S2P6 to

that of spike and bound DH1057.1 >10-fold better, thus demonstrating the successful

transplantation of the target epitope onto the scaffolds.

The binding specificity of S2hlx-4, S2hlx-7 and S2hlx-15 was confirmed by mutating

epitope residues equivalent to spike Phe1148, Leu1152 and Phe1156. Alanine mutations limited

the binding to both S2P6 and DH1057.1 mAbs in S2hlx-4 and -15 to a similar extent to what as

observed for synthesized stem helix peptides (**Figure 2d**)[9]. For S2hlx-7, the individual epitope

mutations greatly decreased DH1057.1 mAb interactions but had almost no effect on S2P6 mAb

recognition. However, an S2hlx-7 variant that contained alanine substitutions at all three epitope

positions completely abrogated binding, thus confirming that antibody interactions were mediated

by the grafted epitope residues. In addition, none of the parent scaffolds these designs were

based on showed binding to S2P6 or DH1057.1, further demonstrating epitope specificity (**Supp.**

Fig. 9).

High resolution (1.9Å) structural determination by x-ray crystallography of the S2hlx-

7/DH1057.1 complex revealed for the first time the structure and interaction mode of the

DH1057.1 antibody with the stem helix epitope. DH1057.1 engaged the epitope at the same angle

as S2P6 mAb and similarly interacted with residues Phe1148, Leu1152, and Phe1156 (SARS-

CoV-2 spike numbering) (**Figure 2e**), as predicted by the alanine mutagenesis (**Figure 2d**). Unlike

S2P6, DH1057.1 interacts with residue K1149 through hydrogen bonds mediated by heavy chain

residues Asp95 and Asp100C (**Figure 2e**). The structure of S2hlx-7 in the complex was closely

aligned with that of the computational model both overall (backbone RMSD=0.5Å) and, most

importantly, over the grafted epitope region (backbone RMSD=0.25Å) (**Figure 2f**). The epitope

conformation displayed on S2hlx-7 closely matched that of the antibody-bound S2P6 epitope and

that of the corresponding region on the prefusion spike (backbone RMSD<0.2Å, **Figure 2f**). These

data validated the computational design of antigens that have high affinity and specificity for

S2P6-like mAbs by mimicking the conformation of their epitope on native spikes.

Design of epitope scaffolds that engage stem helix antibodies with different

specificities.

Given the failure of side chain grafting to produce epitope scaffolds that engage the

CC40.8 mAb, we employed an alternative design technique, termed backbone grafting. Here,

rather than transplanting only the epitope-specific residues onto a preexisting scaffold backbone

which is otherwise unchanged, an entire region of the parent scaffold backbone is replaced with

the backbone of the secondary structure containing the desired epitope. This allows the grafting

of more complex epitopes that have no existing structural match, such as the CC40.8 S2-helix

epitope. Candidate scaffolds were identified by aligning the N- and C-termini of the stem helix

epitope in its CC40.8-bound conformation with the N- and C-termini of possible insertion sites on

candidate scaffolds (**Figure 3a, b**). For scaffolds where the backbone RMSD of this alignment

was below 0.75Å, the epitope replaced the overlapping backbone of the parent scaffold, and

additional mutations were introduced to integrate the epitope into the scaffold and to ensure

productive interactions with CC40.8 mAb (**Figure 3a**). The structures of candidate designs were

predicted using Alphafold2[21], and additional mutations were introduced to ensure proper epitope

conformation. Final designs contained between 16 to 32 mutations relative to the parent scaffolds,

and the grafted epitope was 3.1 to 6.7Å away by RMSD from the backbone it replaced.

Six epitope scaffolds, named S2hlx-Ex1 to 6 (**Figure 3b** for representative models; **Supp.**

Table 1) were tested for bacterial expression and analyzed for binding to S2P6 mAb in an initial

screen. While four of the six designs, S2hlx-Ex2, -Ex3, -Ex4 and -Ex6, bound to S2P6, three of

them those had poor expression (S2hlx-Ex2, -Ex4 and -Ex6) (**Supp. Fig. 10**). To improve

expression, we tested different protein expression tags and evaluated structural homologs of the

parent scaffolds as alternative design templates with limited success (**Supp. Figs. 11, 12**). Next,

we optimized the sequence of the S2hlx-Ex designs using ProteinMPNN[22], a recently described

deep learning algorithm that takes a protein structure as input and generates amino acid

sequences that are predicted to generate the same fold. All four backbone grafting designs that

bound S2P6, S2hlx-Ex2, -Ex3, Ex4, and -Ex6, were optimized with this approach, and sequences

that expressed with high yield (40mg/L of culture on average, **Supp. Fig. 13**) were identified for

all of them. For a given structure, ProteinMPNN generated designs that had highly different

sequences outside the epitope (**Figure 3b**). For example, S2hlx-Ex2 based designs -Ex15, -Ex17,

-Ex19, -Ex20 had between 42% and 55% amino acid sequence identity with the S2hlx-Ex2

template (**Figure 3c**), and only 58-75% identity with each other, yet all bound to the stem helix

mAbs by ELISA (**Supp. Figure 14**).

The binding affinities for the various ProteinMPNN designs based on the backbones of

S2hlx-Ex2 (*S2hlx-Ex15, -Ex17, -Ex19, and -Ex20*), S2hlx-Ex3 (*S2hlx-Ex46 and S2hlx-Ex54*),

S2hlx-Ex4 (*S2hlx-Ex8-Trx and S2hlx-Ex25*), and S2hlx-Ex6 (*S2hlx-Ex34 and -Ex37*) to CC40.8,

S2P6, and DH1057.1 mAbs were determined by SPR (**Supp. Fig. 15, 16)**. S2hlx-Ex2 derived

epitope scaffolds had the highest affinities with dissociation constants between 2nM and 30nM

for S2P6, 2nM and 49nM for DH1057.1, and 3nM to 27nM for CC40.8 mAb (**Figure 3d; Supp.**

Fig. 15). S2hlx-Ex15 and -Ex19 had affinities in a similar range to those measured for spike (**Supp**

Fig. 8) and those reported for the stem helix peptide[9,10]. S2hlx-Ex54, derived from the backbone

of S2hlx-Ex3, bound to S2P6, DH1057.1, and CC40.8 with equilibrium dissociation constants of

45nM, 222nM and 150nM respectively (**Supp. Fig. 16**), while S2hlx-Ex37, based on the fold of

S2hlx-Ex6, had affinities of 664nM, 220nM and 849nM to the three target mAbs (**Supp. Fig. 16**).

To test the ability of epitope scaffolds to engage diverse antibodies that may be elicited

as part of a polyclonal response against the stem helix by vaccination, the designed antigens

were tested against additional broadly reactive stem helix antibodies that were recently described

and were not explicitly targeted at the design stage[23]. The epitope scaffolds bound all tested

antibodies by ELISA (**Figure 3e; Supp. Fig. 17a)**, indicative of their ability to recognize diverse

antibodies that target the displayed epitope. In addition to the mature antibodies, binding was also

tested to five of their UCA or inferred germline precursors (**Figure 3e; Supp. Fig. 17b**). Kinetic

analysis revealed that S2hlx-Ex15 had low nanomolar affinities for DH1057 UCA and the iGLs for

S2P6 and CC40.8 (K_Ds of 22.4nM 1.4nM and 9.7nM), and also bound strongly to Cov44-26 UCA

and Cov89-22iGL by ELISA. Other S2hlx-Ex2 and S2hlx-Ex3 derived designs interacted similarly

with three to five of the antibody precursors tested (**Figure 3d, Supp. Fig. 15**). Four out of the

five antibody precursor recognized spike, but with significantly weaker affinity than S2hlx-Ex15

and S2hlx-Ex19 (**Supp. Fig. 17**).

To ensure that the S2hlx epitope scaffolds interacted with the target mAbs in a similar

fashion to spike, antibody binding was measured to eight epitope scaffolds that spanned all the

design families (**Figure 3b**) and that contained alanine mutations corresponding to spike epitope

sites Leu1145 and Phe1148 in the stem helix (**Supp. Fig. 18**). As anticipated based on structural

analysis and previously published observations[10], all the epitope scaffolds lost binding to CC40.8

when the Leu1145 equivalent residue was mutated, while binding to S2P6 and DH1057.1 was

not affected. When the epitope residue equivalent to spike Phe1148 was mutated, binding was

lost to DH1057.1 and S2P6 in all designs, confirming that antibody interactions are mediated by

the same residues on spike and the designed antigens.

Analysis of a high-resolution (2.3Å) crystal structure of unbound S2hlx-Ex19 further

confirmed the successful transplantation of the CC40.8 mAb epitope (**Figure 3f, Supplemental**

Table 2). As expected, the grafted epitope region in S2hlx-Ex19 diverged considerably from the

backbone region it replaced in the parent scaffold (backbone RMSD=5.8Å), with the epitope

adopting a more exposed conformation that allows for antibody engagement (**Figure 3f**). This

conformational change was likely facilitated by introducing two aromatic residues (Tyr62 and

Phe66) under the epitope helix, which effectively pushed the epitope segment away from the rest

of the protein core. Interestingly, the epitope conformation on S2hlx-Ex19 was almost identical to

that present on the prefusion stem helix, rather than the more extended one induced by CC40.8

(**Figure 3f**). Nevertheless, given the high affinity of S2hlx-Ex19 for CC40.8 (K_D=6.7nM), it is likely

that the antibody can readily induce the necessary conformation upon binding. Thus, by

combining epitope backbone grafting and ProteinMPNN, we engineered antigens that bind with

high affinity and specificity to the major classes of stem helix broadly reactive antibodies as well

as to their inferred precursors.

Cross-reactivity of engineered epitope scaffolds with pre-existing immune

responses induced by SARS-CoV-2 spike.

Next, we tested the ability of the engineered epitope scaffolds to cross-react with pre-

existing immune responses elicited by SARS-CoV-2 spike in order to evaluate their potential to

preferentially amplify spike[815-823] or stem helix directed responses by vaccination. Sera or plasma

samples isolated from three groups of 12 study participants who acquired SARS-CoV-2 immunity

either through vaccination, infection, or vaccination followed by infection respectively, were first

tested for binding to WA-2 spike, RBD and synthesized peptides encoding the spike[815-823] and

stem helix regions (see **Supp. Table 3** for full patient information). This analysis revealed that

only a small fraction of spike-directed humoral responses, regardless of their acquisition route,

targeted the spike[815-823] or stem helix regions (**Figure 4a**). On average, spike[815-823] and stem helix

peptide binding levels were 23.3 and 8.4-fold lower respectively than RBD-focused responses

across all groups, underscoring that the spike[815-823] and the stem helix domains are sub-dominant

targets of antibodies elicited by spike.

Both spike$^{815-823}$ and S2hlx-Ex epitope scaffolds showed significantly higher recognition of

human sera than the corresponding spike peptide domains (**Figure 4a, Supp. Fig. 19**). Among

the spike$^{815-823}$ epitope scaffolds, FP-15 had the best sera recognition, showing 11.3, 7.7, and 5.8-

fold higher IgG binding than the corresponding synthesized peptide in vaccinated, infected and

hybrid immunity samples respectively (**Figure 4b**). Importantly, these interactions were mediated

by the engineered epitope contacts, as binding to the parent scaffold of FP-15 that does not

contain the grafted epitope, was significantly reduced (3.5 -fold in vaccinated, 2.5-fold in infected

and 5.5-fold in hybrid, **Figure 4b**). A similar trend was observed for FP-10 and FP-2, but their

overall binding levels were lower than FP-15, although still above those of the synthesized

spike$^{815-823}$ peptide (**Supp. Fig. 19b**).

From the stem helix designs, S2hlx-Ex19 and S2hlx-Ex15 had the best recognition of

spike-induced immune responses **(Supp. Fig 19a)**. Compared to the synthesized stem helix

peptide, S2hlx-Ex19 bound on average 2.4 times better across all three groups (**Figure 4c**). The

binding was epitope specific, as evidenced by the decrease in interactions with S2hlx-Ex19

variants that contained alanine mutations at hotspot epitope residues Leu1145 (LA) and Phe1148

(FA), or that had the whole epitope replaced by the sequence of the parent scaffold (S2hlx-Ex19

PShlx, **Figure 4c**). As shown by the larger decrease in binding to the FA than LA mutants, cross-

reactive spike responses were more dependent on stem helix residue Phe1148 which is

recognized by both S2P6 and CC40.8 classes of antibodies (**Figure 4c, Supp. Fig 19**). Compared

to S2hlx designs, the ability of the S2hlx-Ex epitope scaffolds to recognize CC40.8 mAb translated

into a broader engagement of spike elicited humoral responses for these types of molecules

(**Figure 4a, Supp. Fig 19**).

Isolation of antibodies with broad coronavirus reactivity from humans pre-exposed

to SARS-CoV-2 using epitope scaffolds.

Next, we characterized the ability of the engineered epitope scaffolds to interact with B

cells that give rise to antibodies with broad activity against the spike[815-823] peptide or the stem

helix from people with preexisting SARS-CoV-2 immunity, as a way to evaluate the potential of

our immunogens to boost such responses by vaccination. Stem helix reactive BCRs were isolated

from IgD- B cells, a pool that included both memory B cells and plasmablasts, from the two

subjects with the highest sera titers against S2hlx-Ex19, using positive selection for S2hlx-Ex19

and negative selection for the parent scaffold S2hlx-Ex19-PS that lacks the epitope (**Figure 5a,b**).

Stem helix epitope-specific memory B cells were isolated at a frequency of ~1:3000 and ~1:2100

respectively from the two samples (**Figure 5b, Supp. Fig 20)**.

Antibody sequences were amplified from the isolated B cells and preliminarily screened

for spike binding[24] in order to select monoclonal antibodies for recombinant production and in-

depth functional characterization. Twelve recombinant antibodies selected from B cells that bound

S2hlx-Ex19 but not S2hlx-Ex19-PS were tested for binding against a panel of spike proteins from

SARS-CoV-2 variants as well as other human and animal betacoronaviruses. Ten antibodies

showed measurable affinity to multiple coronavirus spikes, with three of them displaying binding

on par to that of S2P6 and CC40.8 mAbs to all the spikes tested, including SARS-CoV-1, MERS,

OC43 and HKU-1 (**Figure 5c, Supp. Figs. 21-24**). Interestingly, these three antibodies,

DH1501.1, DH1501.2, and DH1501.3, were clonally related and their heavy and light chain V

gene segment pairing (IGHV1-46/IGKV3-20) matched those of S2P6 mAb. The inferred UCA of

these antibodies displayed tight binding to multiple spikes, similar to what was reported for the

iGL of S2P6 mAb (**Supp. Fig. 25**). Beyond the IGHV1-46/IGKV3-20 derived mAbs, multiple other

unique V_H/V_L pairings, not previously reported, were observed in the isolated antibodies (**Figure**

5c). This suggests the presence of a clonally diverse response against the stem helix epitope that

can be engaged by the engineered epitope scaffold. The best six antibodies by affinity and

breadth were tested side by side with S2P6 and CC40.8 for their ability to neutralize multiple

SARS-CoV-2 variants and SARS-CoV-1 in a pseudovirus neutralization assay. CC40.8 and two

isolated mAbs, DH1501.1 and DH1501.3, were the only ones that displayed neutralization activity,

albeit with high IC_{50} values above 1µg/mL that are typical for antibodies against this epitope.

DH1501.3 and CC40.8 neutralized all the pseudoviruses in the panel, including XBB1.5 and

SARS-CoV-1 (**Figure 5d**).

Given that stem helix antibodies were previously shown to protect against infection in

animal models despite limited neutralization potency[9,10], we next investigated whether the isolated

monoclonal antibodies could control the virus through antibody-mediated functions. To determine

the ability of these mAbs to mediate Fc effector functions, they were first tested for binding to

virus-infected cells, a prerequisite of antibody-dependent cellular cytotoxicity (ADCC). Three of

the five antibodies tested, DH1501.1, DH1501.2, and DH1501.3, showed binding to cells infected

with either the D614G or the BA.1 variant of SARS-CoV-2 in the same range as the previously

described stem helix antibodies CC40.8, S2P6 and DH1057.1 (**Figure 5e**). Binding of the stem

helix antibodies was lower that of the RBD-directed mAb DH1047, supporting the observation that

the stem helix epitope is more occluded on the viral spike. Next, we performed a Natural Killer

(NK) cell degranulation assay as a surrogate for assessing ADCC activity. Antibodies DH1489,

DH1501.1, DH1501.2, and DH1501.3 induced NK cell degranulation against all three variants of

concern tested, with potencies similar to those of known stem helix antibodies (**Figure 5f**).

Interestingly, all stem helix mAbs induced the most potent degranulation against BA.4/5 spike-

transfected cells. Overall, isolated antibodies with IGHV1-46/IGKV3-20 immunogenetics had the

highest activity in both the pseudovirus neutralization and ADCC assays. While other isolated

antibodies with different V_H/V_L pairings were not as broad or potent, it is possible that they could

be further matured to breadth by targeted vaccination and may be valuable targets for rational

vaccine design.

We similarly isolated antibodies against the spike[815-823] epitope by sorting B cells that

bound the FP-10 epitope scaffolds and lacked binding to the original parent scaffold that did not

contain the grafted epitope. The frequency of epitope specific B cells selected was ~1:4,000 in

two analyzed samples and two antibody sequences were recovered for recombinant expression

and characterizations after preliminary screening. These antibodies had distinct immunogenetics

not observed in other previously isolated antibodies against the spike[815-823] epitope (**Figure 5c**).

Nevertheless, they bound tightly to diverse coronavirus spikes, including those of human alpha

coronaviruses 229E and NL63 as is typical of antibodies with broad activity against this epitope

(**Figure 5c**). No binding was detected for SARS-CoV-2 variants BA.1 and XBB1.5, however these

recombinant spikes contained a "hexapro" mutation that is known to affect the presentation of the

spike[815-823] epitope and to limit the binding of antibodies against this site[8].

Antibody isolation from subjects with pre-existing SARS-CoV-2 immunity revealed that the

engineered epitope scaffolds interact with diverse monoclonals against the target epitopes. Some

of the antibodies are derived from the same V_H and V_L genes observed in antibodies previously

isolated[9]. While the epitope scaffold reactive mAbs against the stem helix had weak neutralization

potency, they exhibited strong ADCC activity *in vitro*, revealing a potential protection mechanism

consistent with previous observations for this antibody class. Taken together, the robust sera

binding and memory B cell engagement by epitope scaffolds demonstrate their strong recognition

of pre-existing SARS-CoV-2 immune response and support their use for boosting spike[815-823] and

stem helix antibodies by vaccination.

Engineered stem helix epitope scaffolds elicit antibodies with broad reactivity

against diverse CoVs by vaccination.

Next, we assessed the ability of the engineered epitope scaffolds to elicit antibodies

against the target epitope by vaccination. We focused on testing the stem helix immunogens

belonging to the S2hlx-Ex2 family of design, Ex15, Ex19, Ex20, and Ex17, because: 1) they have

high affinities for all the antibodies tested as well as their precursors; and 2) while they share the

same backbone structure, their sequence outside the epitope is highly divergent, which may help

focus immune responses on the epitope by vaccinating with combinations of different designs.

Epitope scaffolds were multimerized on mi03 nanoparticles (NPs) using the

SpyCatcher/Spytag[25] conjugation system in order to improve immune presentation **(Figure 6a)**.

Homotypic nanoparticles were developed that displayed 60 copies of designs Ex15, Ex19, or

Ex20 respectively. A "mosaic" NP was also engineered where Ex15, Ex19, Ex20, and Ex17 were

conjugated together. Antigenic and NSEM characterization confirmed that the NPs were well

formed and bound S2P6, DH1057.1 and CC40.8 (**Supp. Fig. 26**). Groups of eight BALB/c mice

were vaccinated three times, four weeks apart with either: 1) Ex15-NP; 2) Ex_mosaic-NP; 3) a

mixture of individual Ex15-NP, Ex19-NP and Ex20-NP; 4) a sequential regimen of Ex15-NP prime,

followed by Ex19-NP and Ex20-NP boosts; or 5) GLA-SE only, the adjuvant included in all the

immunizations, as control **(Figure 6b)**. After two immunizations, sera from all animals vaccinated

with epitope scaffolds showed strong reactivity to WA-2 and XBB SARS-CoV-2 spikes **(Figure

6c)**. Sera breadth was tested against spikes from all human beta coronaviruses as well as

RsSHC014 and GXP4L, two pre-emergent animal viruses from bats and pangolins. Binding was

observed against all the spikes tested, with the highest activity against SARS-CoV-1, RsSHC014,

and GXP4L **(Figure 6d)**. Importantly, sera cross-reacted with OC43, MERS, and HKU-1 spikes,

although at lower levels than those measured against SARS-CoV-2, but consistent with the

induction of a broad response against the stem helix epitope **(Figure 6d)**. Of the four immunization

regimens tested, the homotypic Ex15 NP elicited the lowest titers and breadth. Combinations of

epitope scaffolds whether administered together, in mosaic form or sequentially, performed

similarly well, although the Ex_mosaic-NP elicited the highest average titers against heterologous

human betacoronaviruses.

Epitope specificity was demonstrated by strong sera binding to an unrelated epitope

scaffold not used in the immunizations **(Supp. Fig. 27)**. Epitope specificity was further

demonstrated by measuring binding to synthesized peptides encoding the stem helix region as

well as mutated variants at sites known to be important for antibody recognition (**Figure 6e**).

Mutations at residues 1148, 1152, and 1156 decreased sera binding, while alanine substitution

at position 1145 had limited effect, consistent with the elicitation of antibodies that resemble the

binding mode of S2P6 or DH1057.1, rather than that of CC40.8 (**Figure 6e**). No neutralizing

activity was observed for the elicited sera, which was unsurprising given the limited activity of

S2P6-like recombinant monoclonals in our pseudovirus neutralization assay **(Figure 5d)**.

Next, we assessed the ability of the Ex_mosaic-NP to boost stem helix specific responses

in animals previously vaccinated with mRNA encoding the SARS-CoV-2 WA-2 spike and to

protect against a viral challenge with a heterologous virus. Assessing protection was particularly

important since sera elicited by stem helix epitope scaffolds did not neutralize in our assay;

however, antibodies that react with the grafted epitope can promote strong ADCC as shown

above (**Figure 5e,f**) and may protect through antibody-mediated functions. K18-ACE2 mice were

immunized every four weeks with either four shots of mRNA spike or twice with mRNA spike

followed by two boosts of the Ex_mosaic-NP. Two weeks after the last immunization, the animals

were challenged with live WIV-1 virus (**Figure 6f**). WIV-1 was chosen because its spike sequence

is significantly different from that of SARS-CoV-2 (78% over the whole spike and 75% over RBD),

while the stem helix epitope sequence is conserved, as in most sarbecoviruses. Animals

vaccinated with Ex_mosaic-NP had higher average titers of antibodies targeting the stem helix

epitope compared to those that received only spike mRNA, as measured by binding to an epitope

scaffold not used in the immunization; binding to the synthesized stem helix peptide was also

higher, but not statistically significant (**Supp. Fig. 28**). Both immunizations regimens provided

strong protection from WIV-1 based on the percentage of body weight the animals lost six days

after the challenge and when compared to animals that received adjuvant only (**Figure 6g**).

However, animals boosted with Ex_mosaic-NP scored better on two other clinical measures

compared to those that received spike mRNA only. No viral RNA was detected in the lungs of any

of the Ex_mosaic-NP immunized animals and their congestion scores were significantly lower

when compared to those of mRNA spike vaccinated animals (**Figure 6h**). These results

demonstrate that stem helix epitopes elicit sera with broad reactivity against beta coronaviruses

and can offer protection against divergent viruses when used to boost immune responses initially

induced by mRNA spike immunizations.

Discussion

While currently approved coronavirus vaccines and multiple vaccine candidates are based

on spike or RBD proteins[26], here we describe new types of immunogens that aim to elicit different

humoral responses, focused on the conserved stem helix or the spike[815-823] epitopes in the S2

domain of spike. The ability to elicit high titers of antibodies against these regions will likely be

critical to the development of a pan-coronavirus vaccine[27], a goal the molecules designed here

aim to contribute towards. The epitope scaffold proteins were engineered using a combination of

previously developed grafting methods[15,28] and more recently described machine learning

techniques[29] that proved critical for improving both immunogen stability and antigenicity.

Structural and mutational analysis confirmed that the epitope scaffolds engage multiple

antibodies, as well as their precursors, with high affinity and specificity. For each class of epitope

scaffolds, some designs showed tighter antibody binding than corresponding epitope peptides,

especially as it relates to the recognition of antibody precursors.

Spike[815-823] and stem helix epitope scaffolds that bound S2P6 and DH1058 mAbs, but not

CC40.8 mAb, were engineered using side chain grafting, an approach that modifies the sequence

of existing proteins to accommodate the epitope and that was successfully used in the past to

engineer epitope scaffolds against HIV[18,30,31] and RSV[32] antibodies. However, for conformationally

complex epitopes like the one induced by the CC40.8 mAb on spike, no known protein structures

may exist that are structurally similar to allow side chain grafting. To successfully transplant these

types of epitopes, different methods, called flexible backbone approaches, are needed to model

and control the conformation of the epitope on the target scaffolds. This can be achieved either

by engineering epitope scaffolds *de novo* by folding idealized protein backbones around the target

epitope, as was previously done for epitopes on RSV[19,33], or by modifying the backbone of existing

proteins to adopt the conformation of the grafted epitope, as illustrated before for HIV[28,30] and

here for designs that bind CC40.8 mAb. Previously, the success of flexible backbone epitope

grafting was limited; the majority of the computational designs failed to express recombinantly

and the ones that did typically required multiple rounds of directed evolution to achieve high

affinities to the target antibodies.

Here we combined previously described backbone grafting with recent machine learning

based protein engineering approaches to significantly improve the success rate of epitope

scaffolds designed using flexible backbone modeling. Structure prediction with AlphaFold2[34] was

used to validate the conformation of the grafted epitope in the structural context of the target

scaffold, while ProteinMPNN[22] was subsequently employed to identify optimal sequences that

fold into the desired epitope scaffold structure. With this approach, multiple proteins were

engineered *in silico* that bound tightly to CC40.8 mAb without needing additional experimental

optimization. Of note, the epitope was successfully transplanted on scaffolds with diverse

backbone structures, supporting our approach as a general way to present structurally complex

motifs in diverse molecular contexts. Development of stem helix epitope scaffolds that bound with

high affinity to all three target antibodies (S2P6, DH1057.1, CC40.8) proved challenging because

CC40.8 induces a different conformation on the epitope upon binding compared to the other two

antibodies. This suggests that, for a successful design, the epitope presentation must allow for

antibody-induced fit and some level of conformational changes. This was not explicitly modeled

here at the design stage but likely represents an area of improvement in the computational

workflow for transplanting epitopes that adopt different antibody-bound conformations in the

future. Interestingly, the stem helix immunogen S2hlx-Ex19, which bound with affinities for

CC40.8 and S2P6 similar to those of the native spike, displays the epitope at the N-terminus of

the scaffold. While a crystal structure of this unbound epitope scaffold revealed that the epitope

is well folded and adopts a helical conformation, it is possible that its positioning may more readily

allow for conformational changes upon antibody binding relative to other designs where the

epitope is grafted internally.

The ability of engineered immunogens to engage diverse antibodies against the displayed

epitope is likely important for their ability to elicit a robust polyclonal response *in vivo*, as has been

recently demonstrated for HIV vaccine candidates[35,36]. The epitope scaffolds designed here

engaged multiple spike[815-823] and stem helix antibodies with high affinity *in vitro*, which translated

to robust reactivity to polyclonal sera elicited by SARS-CoV-2 and the engagement of BCRs with

diverse immunogenetics against the target epitopes. The stem helix epitope scaffolds that also

bound CC40.8 mAb in addition to S2P6 and DH1057.1 mAbs, showed the highest level of binding

to human sera from subjects exposed to SARS-CoV-2 by vaccination, infection or both. The

S2hlx-Ex19 design was used to isolate multiple antibodies with broad reactivity against the stem

helix from people with pre-existing immunity. Some of these monoclonal antibodies had the same

V_H/V_L pairings (IGHV1-46/IGKV3-20) as previously isolated antibodies, suggesting that this type

of antibody is commonly induced by SARS-CoV-2 immunization and may be targeted for boosting

with a next-generation vaccine. However, as found here, other broad antibodies with different

immunogenetic characteristics exist against S2 and they can be engaged by the designed epitope

scaffolds as well. Isolated stem helix antibodies displayed limited neutralization against SARS-

CoV-2 variants, consistent with previous reports describing other antibodies against this

epitope[9,37]. Interestingly, despite their limited neutralization, stem helix antibodies were shown by

others to protect well against live virus challenges[9,37], likely through their antibody-mediated

functions which we demonstrated our antibodies also have.

In initial immunization studies, stem helix epitope scaffolds elicited sera with broad

reactivity against all human beta coronaviruses and towards two animal viruses tested. Sera

bound specifically to the target epitope in a manner consistent with the induction of S2P6-like

mAbs. Future studies will analyze the B cell repertoire of vaccinated animals in more detail to

better understand the specificity and protection ability of the elicited responses. The induced sera

did not neutralize SARS-CoV-2 pseudoviruses in our assay, consistent with our observations and

previous reports that antibodies against the stem helix are poor neutralizers, but can protect

through antibody mediated functions[9,37]. To demonstrate that the engineered immunogens can

protect against coronavirus infection, we vaccinated K18-ACE mice pre-exposed to spike mRNA

with stem helix immunogens, prior to a viral challenge with WIV-1. Compared to an adjuvant-only

group, mice boosted with our immunogens were protected against weight loss, had limited lung

congestion, and undetectable viral load in the lungs. While mice immunized with only mRNA spike

were also protected against weight loss, they exhibited increased congestion and higher titers of

lungs viremia compared to mice that received the stem helix epitope scaffolds. Taken together,

these results support the continued development and characterization of epitope scaffold

immunogens to boost responses against conserved S2 epitopes as part of a next-generation pan

betacoronavirus vaccine.

Acknowledgments

We thank Whitney Edwards Beck, Deborah Murray, Micah McClain, Thomas Burke, Jack Anderson, Lynn Harrington, Christopher Todd and Kathlene Chmielewski for help with acquiring and managing the human samples. We thank Aria Arus-Altuz for developing the fluorescent probes used for FACS. We thank Brian Watts, Ken Cronin and Alam Munir for their assistance at the DHVI Biomolecular Interaction Analysis (BIA) Core Facility. We thank Wes Rountree and Yunfei Wang for the statistical analysis. We thank Advaita Sing and Christopher O'Donnell from the Azoitei lab for their help. We thank Jacob Gater and Pahvie Chhan from the Duke Human Vaccine Institute Small Animal Team for their help with the immunization studies. We thank Priyamvada Acharya and Jared Lindenberger for their help with crystallization experiments. Structural results shown in this report are derived from data collected at Southeast Regional Collaborative Access Team (SER-CAT; 22 ID and 19 BM) beamlines at the Advanced Photon Source, Argonne National Laboratory. Use of the Advanced Photon Source was supported by the U. S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Contract No. W-31-109-Eng-38.

Funding: This work was supported by grants P01AI158571 (B.F.H) and R01AI155804 (M.L.A) from the National Institute of Allergy and Infectious Diseases of the National Institute of Health and by a Duke School of Medicine COVID Award (M.L.A).

Author Contribution. MLA, BK, and DJM: conceptualization, methodology, study design, and immunogen engineering; BK, CH, CB, KW, DJM, and BR performed *in vitro* characterization of epitope scaffold immunogens; BK, DW, and KW performed the crystallographic analysis; QY, LX, AF, RP, MB, KOS, and DWC performed the B cell isolation, sequencing and antibody identification; AE and DM performed the pseudovirus neutralization assay; DM, BM, TK, SSO, GF performed NK degranulation analysis; AN, CH, AS, JMF, AH, NJC performed the animal

immunization, the viral challenge experiments and their analysis; CW, EQP, EBW, KW, KOS, RB,

and BFH: resources; KW and EVI preformed unmutated common ancestor interference; KM and

RJE performed NSEM analysis; figure development; MLA, DJM, BK: writing, original draft; MLA,

DJM, BK, BFH, DW, KW, KOS, DM, GF, AE, DM, AN, AS, XL, EVI: writing, review and editing;

MLA, BFH, KW, KOS: funding acquisition; MLA: project administration and supervision.

Competing interests: MLA, BFH, DJM and BK submitted a patent application that covers

the engineered epitope scaffolds. The other authors declare no competing interests.

Data and materials availability: Crystal structures have been deposited in the PDB under

accession numbers 8F5I, 8FDO and 8F5H. The epitope scaffold can be obtained through an MTA

from the corresponding author. Other reagents are available upon requests.

Materials and methods

Computational design of epitope scaffolds.

Design of epitope scaffolds by side chain grafting. A database of 9884 candidate scaffolds was created by selecting from the Protein Data Bank (PDB) structures that were: 1) determined by x-ray crystallography; 2) high resolution (<2.8 Å); 3) monomeric; 4) expressed in *E. Coli*; 5) not of human origin and 6) that did not contain ligands. Epitope scaffolds were designed with Rosetta as previously reported, using the RosettaScripts for side chain grafting described in *Silva et. al.*, with the following parameters in the *MotifGraft* mover: RMSD_tolerance="0.3"; NC_points_RMSD_tolerance="0.5"; clash_test_residue="ALA"; clash_score_cutoff="5". For the design of spike$^{815-823}$ peptide (FP) epitope scaffolds, the structure of spike$^{815-823}$ peptide fragment ^{815}RSFIEDLLF^{823} as described in the crystal structure of the DH1058-FP peptide complex (PDBid: 7tow) was used as the target epitope to graft. With the exception of residues F817 and L821 which were allowed to change in the designed epitope scaffold because they did not contribute to antibody binding, the identity of all the other epitope residues was maintained. For the design of epitope scaffolds that bind S2P6 and DH1057.1 mAbs, the structure of the stem helix fragment ^{1148}FKEELDKYF$^{1156\ from}$ PDBid:7rnj was grafted. The identity of residues E1150 and K1154 was allowed to change during the design process while the other epitope residues were kept fixed. From the epitope scaffold generated by the automated protocol, the top 100 models by Rosetta ddG that were also smaller than 150 amino acids were visually examined to ensure appropriate epitope transplantation and antibody-scaffold interaction. At this stage, the suitability of a given parent scaffold in terms of function, conformational flexibility, and expression protocol was also investigated by referencing the publication it originated from. If necessary, additional changes were introduced in a candidate epitope scaffold using Rosetta fixed backbone design to remove antibody-epitope scaffold contacts that were not due to the target epitope and to ensure proper

interactions between the epitope and the rest of the scaffold. The best 15 designs for each of the

target epitopes were chosen for experimental characterization.

Design of stem helix epitope scaffolds by backbone grafting and MPNN optimization.

Epitope scaffolds that display the stem helix epitope recognized by the CC40.8 antibody

(PDBid: 7sjs) were designed by backbone grafting as previously reported and using the

RosettaScripts described by Silva et. al. The same set of curated parent scaffolds as the one

used for side chain grafting above was searched here. The structure of the stem helix fragment

[1144]ELDSFKEELDKYFK[1157] from PDBid:7sjs was grafted. The identity of residues E1144, D1146,

S1147, E1150, K1154, and K1157 were allowed to change during the design process while the

other epitope residues were kept fixed. The following parameters were used:

RMSD_tolerance="5.0" NC_points_RMSD_tolerance="0.75"
clash_score_cutoff="5" clash_test_residue="ALA"
hotspots="2:5:6:8:9:10:12:13"
combinatory_fragment_size_delta="1:1"
max_fragment_replacement_size_delta="-4:4"
full_motif_bb_alignment="0"
allow_independent_alignment_per_fragment="0"
graft_only_hotspots_by_replacement="0"
only_allow_if_N_point_match_aa_identity="0"
only_allow_if_C_point_match_aa_identity="0"
revert_graft_to_native_sequence="0"
allow_repeat_same_graft_output="0" />

The initial hits were filtered to remove scaffolds with (1) an average clash score greater

than 5 across the epitope, and (2) a Rosetta calculated ddg greater than 0. The top 100 models

by Rosetta ddG that were also smaller than 150 amino acids were then visually examined as

above and then Alphafold2 was used to predict the structure of the 10 best designs using the

following parameters-

--model_preset=monomer
--db_preset=full_dbs
--max_template_date=2021-11-01
--uniref90_database_path=/data/uniref90/uniref90.fasta
--mgnify_database_path=/data/mgnify/mgy_clusters.fa
--uniclust30_database_path=/data/uniclust30/uniclust30_2018_08/uniclust30_2018_08
--bfd_database_path=/data/bfd/bfd_metaclust_clu_complete_id30_c90_final_seq.sorted_opt
--template_mmcif_dir=/data/pdb_mmcif/mmcif_files
--obsolete_pdbs_path=/data/pdb_mmcif/obsolete.dat
--pdb70_database_path=/data/pdb70/pdb70

The scaffolds were then aligned via the grafted epitope to the stem helix fragment in the

CC40.8 structure PDBid:7sjs and inspected for clashes between the scaffold and antibody.

Scaffolds with clashes, such as where the antibody binding epitope was occluded or buried, were

modified such that key epitope residues above remained fixed while other residues in the epitope

and in contacting parts of the scaffold were allowed to vary through iterative rounds of fixed-

backbone rotamer-based sequence design in Rosetta. After each round, we checked the

Alphafold2 prediction and repeated until the epitope was predicted to be presented in a suitable

orientation for antibody binding (**Figure 3a**). Expression tags were screened by N-terminally

tagging S2hlx-Ex4 with GST, SUMO, Thioredoxin (Trx), Maltose binding protein (MBP), or F8H

tags. As Trx gave the highest fraction of full-length protein it was then used to tag S2hlx-Ex2 and

-Ex6 (**Supp. Fig. 11**).

The parent structure of S2hlx-Ex4 (PDBid:2QYW) was used to screen the VAST+

database   (https://structure.ncbi.nlm.nih.gov/Structure/VAST/vast.shtml)   for   homologous

structures, which were identified by RMSD (<2Å) and high fraction of alignment (>80%). Two

homologs identified were expressed with Trx tags (**Supp. Fig. 12**).

For ESs where protein expression was low the protein structures, either the Rosetta model

or   Alphafold   prediction,   were   entered   into   ProteinMPNN   at   the   website,

https://huggingface.co/spaces/simonduerr/ProteinMPNN. Initial designs for S2hlx-Ex2 and -4

used sampling temperature of 0.1 and backbone noise of 0.02 and the whole epitope sequence

was fixed, For S2hlx-Ex2, this approach produced six related soluble ESs but failed for S2hlx-

Ex4. The second set of S2hlx-Ex4 and S2hlx-Ex6 designs used the higher sampling temperature

(0.25) and higher backbone noise (0.2) to increase sequence diversity in designs, and only

L^{1145}xxFKxELDxYF^{1156} residues were fixed. ES structures were modelled using the integrated

Colabfold prediction and scored by RMSD and pLDDT, and the highest scoring were re-predicted

using Alphafold2. For the S2hlx-Ex6 designs, the epitope was occluded or buried in several

designs. In these cases, residues in the epitope or surrounding parts of the ES were manually

modified and the ES structure prediction repeated using Alphafold2 until the epitope was

predicted to be suitable to binding to S2P6.

Plasmids and DNA synthesis.

Genes encoding designed epitope scaffolds were commercially synthesized and cloned

into pET29b (Genscript). Genes encoding the antibody heavy and light chains were similarly

synthesized and cloned into the pcDNA3.1 vector (GenScript). Oligonucleotides were synthesized

by IDT. Mutations were introduced in the synthesized plasmids by Q5 mutagenesis (NEB).

Recombinant protein expression and purification.

Epitope scaffolds.

Plasmids encoding epitope scaffolds were transformed into E. coli BL-21 (DE3) (New

England Biolabs) cells and 5mL starter cultures were grown overnight at 37C in Lysogeny Broth

(LB) supplemented with 50µg/mL kanamycin. The cultures were diluted 1:100 in Terrific BrothTB

media (RPI) supplemented with kanamycin and grown at 37 °C to an OD600 of ~0.6. The

temperature was subsequently lowered to 16 °C and the cultures were grown to OD600 of ~0.8

and induced with isopropyl β-D-1-thiogalactopyranoside (IPTG) to a final concentration of 0.5mM.

The cultures were shaken for 16–18 hours at 200rpm. Cell pellets were collected by centrifugation

at 14,500x$g$xg and lysed in B-PER Reagent (ThermoFisher Scientific) according to the

manufacturer's protocol. The lysates were centrifuged at 14,000 xg for 30 minutes at 4C. The

supernatant was incubated with Ni-NTA beads (Qiagen) equilibrated with native binding buffer

(20mM NaH2PO4, 500M NaCl, 2% glycerol, 10mM imidazole, pH=7.5) for an hour at 4C. The

beads were settled by centrifugation and the supernatant was removed by pipetting. The beads

were washed with wash buffer (50mM NaH2PO4, 500M NaCl, 2% glycerol, 30mM imidazole,

pH=7.5) after which the protein was eluted with elution buffer (20mM NaH2PO4, 500M NaCl,

250mM imidazole, pH=7.5). Protein expression and purity was confirmed by SDS-PAGE analysis,

and quantified spectrophotochemically at 280nm on a Nanodrop 2000 (ThermoFisher Scientific).

The eluted protein was concentrated on 3kDa MW spinMW spin columns (ThermoFisher

Scientific) and further purified by size exclusion chromatography in 20mM NaH2PO4, 150mM

NaCl, pH=7.5, on an AKTA-Go FPLC (Cytiva) using a Superdex 200 Increase 10/300GL column.

Fractions containing monomeric protein were pooled and concentrated as above.

Monoclonal antibodies.

Expi293F cells (ThermoFisher Scientific) were split to a density of 2.5x10^6 cells/mL in

Expi293 Expression Medium with GlutaMAX (Gibco) and were transiently transfected with an

equimolar plasmid mixture of heavy and light chain using Expifectamine (Invitrogen). For a typical

100mL size transfection 100µg amount of total DNA and 270µL of lipofectamine were used. After

overnight incubation, enhancers were added as per the manufacturer's protocol and the cultures

were incubated with shaking for five days at 37C and 5% CO_2. The cell culture was centrifuged

to remove the cells and the supernatant was filtered with a 0.8-micron filter. The filtered

supernatant was incubated with equilibrated Protein A beads (ThermoFisher) for one hour at 4 °C

and washed with 20mM Tris, 350mM NaCl at pH=7. The antibodies were eluted with a 2.5%

Glacial Acetic Acid Elution Buffer and were buffer exchanged into 25mM Citric Acid, 125mM NaCl

buffer at pH=6. IgG expression was confirmed by reducing SDS-PAGE and quantified by

measuring absorbance at 280nmm (Nanodrop 2000).

Antigenic characterization of epitope-scaffolds.

Binding by Surface Plasmon Resonance.

To determine the dissociation constants (K_Ds) between epitope scaffolds and target

antibodies, binding was measured by Surface Plasmon Resonance (SPR) on a Biacore T200

instrument using Protein-A coated S series chips. The target antibodies were individually captured

for 60 sec at 30 µL/min to a level of 1000-2000RU onto this surface. The epitope scaffolds were

subsequently injected as analytes at five concentrations using the single cycle injection method.

The association phase was carried out for 180 seconds and the dissociation was done for 900

seconds with HBS-EP+ buffer flowing at 30µL/min. Regeneration of the binding surface was done

in 10mM glycine-HC, pH=2 for 30 seconds at 30µL/min with a 30 second baseline stabilization. A

1:1 Langmuir or Heterogenous ligand model was used for data fitting and analysis. For K_Ds

between Spike proteins and target antibodies, neutravidin (Thermo Scientific) was first captured

for 30-60 sec at 30 µL/min to a level of 1000-2000RU using CM5 S-series chips. Then biotinylated

Spike proteins (WA-2, 2P, R&D systems) were subsequently captured to a level of 3-500RU by

the neutravidin. FAB fragments of antibodies were then injected as analytes at five concentrations

using the single cycle method described above.

Antibody binding by ELISA.

Spike[815-823] peptide (residues 808-833) and stem helix (1140-1163) peptides and mutated

variants were commercially synthesized (Genscript). The following Spike proteins were used: WA-

2, 2P; SARS-CoV-1, 2P; MERS, 2P; OC43, 2P; HKU-1, 2P; GXP4L, 2P; SHC014, 2P; RaTG13,

2P, (all produced according to [38,39]); and Delta; BA1.1, Hexapro; XBB, Hexapro; 229E; NL63; (all

Sino Biological). Peptides and Spikes were diluted to 2µg/mL in 0.1M Sodium Bicarbonate and

epitope scaffolds were diluted to 100µg/mL. Antigens were coated onto 96 or 384 well high

binding enzyme-linked immunosorbent assay (ELISA) plates (Corning) by overnight incubation at

4C. Plates were washed with Superwash buffer (1X PBS supplemented with 0.1% Tween-20)

and blocked for 1 hour at room temperature with Superblock buffer supplemented with azide (80g

Whey Protein, 300mL Goat Serum, 20mL aqueous 5% Sodium Azide, 10mL Tween20, 80mL of

25X PBS, diluted to 2L). Antibodies starting at 100µg/mL were serially diluted in Superblock with

azide using a threefold serial dilution, added to the plates, and incubated at room temperature for

1 hour. Plates were then washed twice with Superwash and goat anti Human IgG-HRP secondary

antibody (Jackson Immunoresearch) diluted in SuperBlock (without Sodium azide) was

subsequently added. After incubating at room temperature for 1 hour, the plates were washed

four times in Superwash. Room temperature TMB (Tetramethylbenzidine) substrate was added

and after 5 minutes elapsed, the reaction was stopped by acid stop solution (0.33N HCl).

Absorption was measured at 405nm on a Cytation 1 plate reader (BioTek). Data was analyzed

and plotted with Prism version 10.0.0 (Graphpad).

Sera binding by ELISA.

Antigens were diluted to 2µg/mL in 0.1M Sodium Bicarbonate and coated onto 384 well

high binding enzyme-linked immunosorbent assay (ELISA) plates (Corning) by overnight

incubation at 4C. Plates were washed with Superwash buffer (1X PBS supplemented with 0.1%

Tween-20) and blocked for 1 hour at room temperature with Superblock buffer supplemented with

azide (80g Whey Protein, 300mL Goat Serum, 20mL aqueous 5% Sodium Azide, 10mL Tween20,

80mL of 25X PBS, diluted to 2L). Mouse sera starting at a dilution ratio of 1:50 and control

antibodies starting at 100µg/mL were serially diluted in Superblock with azide using a fivefold

serial dilution scheme. Human sera starting at a dilution ratio of 1:30 and control antibodies

starting at 100µg/mL were serially diluted in Superblock with azide using a threefold serial dilution

scheme. Sera and controls were added to the plates and incubated at room temperature for 1.5

hours. Plates were then washed twice with Superwash and the corresponding IgG-HRP

secondary antibody (goat anti mouse IgG-HRP or goat anti human IgG-HRP; Jackson

Immunoresearch) diluted in SuperBlock (without Sodium azide) was subsequently added. After

incubating at room temperature for 1 hour, the plates were washed four times in Superwash.

Room temperature TMB (Tetramethylbenzidine) substrate was added and after 15 minutes

elapsed, the reaction was stopped by acid stop solution (0.33N HCl). Absorption was measured

at 405nm on a SpectraMax Plus plate reader (Molecular Devices). Data was analyzed and plotted

with Prism version 10.0.0 (Graphpad).

Structural analysis by x-ray crystallography.

Crystallography experiments were performed using the sitting drop vapor diffusion

technique. 96-well crystallization plates were set, in which the protein of interest was mixed in a

1:1 ratio with reservoir solution. For *S2hlx-Ex*19, crystals were obtained when a 9.5mg/mL sample

was mixed with 0.1 M citric acid, 3.0 M sodium chloride, pH 3.5 (Index Screen) at 4℃. *S2hlx*-7

and antibody DH1057.1 were mixed in 1:1 molar ratio (300uM) and incubated for one hour at 4C.

Crystals for the complex were obtained in 25% PEG 3350 at room temperature. Prior to data

collection, crystals were cryoprotected in mother liquor supplemented with 20% glycerol before

being plunge-frozen into liquid nitrogen. A full description of the crystallographic data collection

and refinement statistics can be found in **Supplementary Table 2**. Diffraction data were indexed

in iMOSFLM[40] and scaled in AIMLESS[41]. Molecular replacement solutions for both *S2hlx-Ex*19

and the *S2hlx*-7-DH1057.1 complex were found in PhaserMR[42], using PDB ID: 3N1B as a search

ensemble for *S2hlx-Ex*19, PDB ID: 3LMO as a search ensemble for *S2hlx*-7 and PDB ID: 6UOE

as a search ensemble for DH1057.1. Coordinates for *S2hlx-Ex*19 and *S2hlx*-7-DH1057.1 were

iteratively built and refined using Coot[43], ISOLDE[44] and Phenix[45] to Rwork/Rfree values of 25.4/

27.5 and 21.1/24.5, respectively. The presence of translational non-crystallographic symmetry in

the *S2hlx*-7-DH1057.1 crystal likely accounts for its relatively high Rfree value and the L-test did

not indicate the presence of any twinning.

Reactivity of designed epitope scaffolds with human samples.

Human subject studies were approved by the Duke University Health System Institutional

Review Board (IRB) and conducted in agreement with the policies and protocols approved by the

Duke IRB, consistent with the Declaration of Helsinki. Written informed consent was obtained

from all research subjects or their legally authorized representatives. Study participants with

SARS-CoV-2 acute infection were enrolled in the Molecular and Epidemiological Study of

Suspected Infection protocol (MESSI, IRB Pro00100241) at Duke University, and were followed

longitudinally. Samples were selected for this study from participants who had seroconverted and

had symptom onset more than 10 days prior. From patient 26 and patient 29, PBMCs were

collected and analyzed by FACS at the same time point used for sera analysis for patient 26 and

7 days later for patient 29. Samples from vaccinated only participants were obtained from subjects

enrolled in the Study of Immune Response to COVID-19 Vaccines (IRB Pro00107929) at Duke

University. Supplementary Table 2 describes the vaccination protocol, SARS-CoV-2 variant

detected and the time of collection for the samples used in this analysis.

Human sera was analyzed as described above in *sera binding by ELISA* method.

Statistical differences were tested using the Wilcoxon signed rank test.

Isolation and analysis of epitope scaffold reactive B cells.

We prepared B cell tetramers of biotinylated S2hlx-Ex19 or FP-10 by mixing with

Streptavidin-VB515 (Miltenyi Biotec) and Streptavidin-AlexaFluor 647 (ThermoFisher Scientific)

at 4:1 molar ratio. A scaffold-only tetramer was also made with biotinylated S2hlx-Ex19-PShlx or

FP-10-wt mixed with Streptavidin-BV421 (Biolegend). These probes did not contain the grafted

epitope.

Cryopreserved PBMCs were thawed in warm RPMI-1640 containing 10% FBS, then

counted. Cells were stained with pre-optimized concentrations of the following antibodies: PE

anti-human IgD (clone IA6-2, BD Biosciences), PE-TXRD anti-human CD10 (clone HI10A, BD

Biosciences), PE-Cy5 anti-human CD3 (clone HIT3a, BD Biosciences), PE-Cy7 anti-human C27

(clone O323, ThermoFisher Scientific), AlexaFluor 700 anti-human CD38 (clone LS198-4-3,

Beckman Coulter), APC-Cy7 anti-human CD19 (clone SJ25C1, BD Biosciences), BV570 anti-

human CD16 (clone 3G8, Biolegend), BV605 anti-human CD14 (clone M5E2, Biolegend), and

BV711 anti-human IgM (clone G20-127, BD Biosciences). S2hlx-Ex19-VB515- and -AF647, and

S2hlx-Ex19-PShlx-BV421 were added to identify stem helix epitope-specific B cells. Cells were

incubated with Aqua Live/Dead (ThermoFisher Scientific) to exclude dead cells. Cells were

acquired on a BD S6 cell sorter (BD Biosciences). Data were analyzed using FlowJo v10.8 (BD

Biosciences).

Immunoglobulin heavy and light chain variable regions (VH and VK/L) from singly sorted

antigen-specific memory B cells were RT-PCR amplified using SuperScript III and AmpliTaq Gold

360 Master Mix (Thermo Fisher Scientific, Waltham, MA) under conditions previously described[46].

PCR products were purified in Biomek FX Laboratory Automation Workstation (Beckman Coulter,

Indianapolis, IN) and sequenced by Sanger sequencing. The V(D)J rearrangement, somatic

hypermutation frequency, CDR3 length of VH and VK/L chains, and antibody clonal lineages were

analyzed using the software Cloanalyst[47]. The heavy chains of mAbs DH1493, DH1501, and

DH1502 were assigned to the same clone by the software package Cloanalyst. Unmutated

common ancestor (UCA) inference was performed using the heavy and kappa chain pairs of the

three mAbs using the paired-chain inference implementation of the UCA inference part of the

software package Cloanalyst.

RT-PCR amplified sequences were transiently expressed as previously described[24].

Briefly, the linear expression cassettes were constructed by overlapping PCR to place the PCR-

amplified VH and VK/L chain genes under the control of a CMV promoter along with heavy chain

IgG1 constant region or light chain constant region and a BGH ploy A signal sequence. The linear

expression cassettes of heavy and light chains were then co-transfected into 293T cells in 6-well

plates. After 3 days, the cell culture supernatants were harvested and concentrated for binding

assays. For antibodies of interest, recombinant IgG1 monoclonals were expressed and purified

as described above.

Pseudovirus neutralization assay.

The pseudovirus neutralization assay performed at Duke has been described in detail[48]

and is a formally validated adaptation of the assay utilized by the Vaccine Research Center; the

Duke assay is FDA approved for D614G and other SARS-CoV-2 variants. For measurements of

neutralization, pseudovirus was incubated with 8 serial 5-fold dilutions of antibody samples in

duplicate in a total volume of 150 µl for 1 hr at 37C in 96-well flat-bottom culture plates.

293T/ACE2-MF cells were detached from T75 culture flasks using TrypLE Select Enzyme

solution, suspended in growth medium (100,000 cells/ml) and immediately added to all wells

(10,000 cells in 100 µL of growth medium per well). One set of 8 wells received cells + virus (virus

control) and another set of 8 wells received cells only (background control). After 71-73 hrs of

incubation, medium was removed by gentle aspiration and 30 µl of Promega 1X lysis buffer was

added to all wells. After a 10-minute incubation at room temperature, 100 µl of Bright-Glo

luciferase reagent was added to all wells. After 1-2 minutes, 110 µl of the cell lysate was

transferred to a black/white plate. Luminescence was measured using a GloMax Navigator

luminometer (Promega). Neutralization titers are the inhibitory dilution (ID) of serum samples at

which RLUs were reduced by 50% (ID50) compared to virus control wells after subtraction of

background RLUs. Serum samples were heat-inactivated for 30 minutes at 56C prior to assay.

Infected-cell antibody binding assay.

The binding of isolated anti-SARS-CoV-2 monoclonal antibodies to infected cells was

measured as previously reported[49]. Briefly, Vero E6 cells expressing TMPRSS2 and ACE2 and

infected with either D614G (Germany/BavPat1/2020) or BA.1 (hCoV-19/USA/MD-

HP20874/2021) variants were incubated with TrypLE Select (Gibco) for 15 minutes at 37°C to

detach cells and washed with PBS. Monoclonal antibodies were then added to infected cells at 2,

10 or 50 µg/mL. Approximately $2x10^5$ infected cells were incubated with the mAbs for 30 minutes

at room temperature, washed and then incubated with vital dye (Live/Dead Far Red Dead Cell

Stain, Invitrogen) for 15 minutes at room temperature to exclude nonviable cells from subsequent

analysis. Cells were then washed with Wash Buffer (1%FBS-PBS; WB), pelleted by centrifugation

and incubated with 1 mL of 4% Methanol-free Formaldehyde (Duke GHRB SOP 38; Attachment

17) for 30 minutes at room temperature. Cells were then washed twice with Wash Buffer,

permeabilized with CytoFix/CytoPerm (BD Biosciences) and stained with A568-conjugated anti-

SARS-COV-2 nucleocapsid antibody (40143-MM08, Sino Biological) and PE/Cy7-conjugated

secondary anti-Human IgG Fc antibody (Clone: HP6017, Biolegend) for 30 minutes at room

temperature. Cells were washed and resuspended in 250 µL PBS–1% paraformaldehyde.

Samples were acquired within 24 h using a BD Fortessa cytometer and a High Throughput

Sampler (HTS, BD Biosciences). Data analysis was performed using FlowJo 10 software (BD

Biosciences). Gates were set to include singlet, live, nucleocapsid+ (NC+) and IgG+ events.

Binding to mock infected cells was measured using the live cell gate as there were no NC+ events.

All final data represent specific binding, determined by subtraction of non-specific binding

observed in assays performed with mock-infected cells.

Antibody-dependent NK cell degranulation assays (infected and Spike-transfected).

Cell-surface expression of CD107a was used as a marker for NK cell degranulation and

performed as previously described[49]. Briefly, target cells were 293T cells 2-days post transfection

with a SARS-CoV-2 S protein (D614G) expression plasmid. Natural killer cells purified by negative

selection (Miltenyi Biotech) from peripheral blood mononuclear cells obtained by leukapheresis

from a healthy, SARS-CoV-2-seronegative individual (Fc-gamma-receptor IIIA (Fcy-RIIIA)158

V/F heterozygous) previously assessed for Fcy-RIIIA genotype and frequency of NK cells were

used as a source of effector cells. NK cells were incubated with target cells at a 1:1 ratio in the

presence of monoclonal antibodies, Brefeldin A (GolgiPlug, 1 µl/ml, BD Biosciences), monensin

(GolgiStop, 4µl/6mL, BD Biosciences), and anti-CD107a-FITC (BD Biosciences, clone H4A3) in

96-well flat bottom plates for 6 hours at 37ºC and 5% CO_2. NK cells were removed from the wells

and stained for viability prior to staining with CD56-PECy7 (BD Biosciences, clone NCAM16.2),

CD16-PacBlue (BD Biosciences, clone 3G8), and CD69-BV785 (Biolegend, Clone FN50). After

three washes, cells were resuspended in 115 µL PBS–1% paraformaldehyde. Flow cytometry

data analysis was performed using FlowJo software (v10). Data is reported as the area under the

curve (AUC) of %CD107a+ live NK cells (gates included singlets, lymphocytes, aqua blue-,

CD56+ and/or CD16+, CD107a+), calculated as previously described[49] at three concentrations of

monoclonal antibody: 2, 10 and 50 µg/mL. All final data represent specific activity, determined by

subtraction of non-specific activity observed in assays performed with mock-infected cells and in

the absence of antibodies, and in the presence of a non-specific monoclonal antibody, the anti-

HIV-1 antibody VRC01.

Development of mi03 nanoparticles conjugated with epitope scaffolds for immunizations.

SpyCatcher003-Mi3 Nanoparticles.

Plasmid encoding for the spycatcher003-mi3 particles was acquired from Addgene

(Plasmid #159995) and transformed into *E. coli* BL-21 (RIPL) (Agilent Technologies) cells. Five

milliliters starter cultures were grown overnight at 37C in Lysogeny Broth (LB) supplemented with

1% glucose and 50µg/mL kanamycin. Cultures were diluted 1:100 in Terrific Broth (TB) media

(RPI) supplemented with kanamycin and grown at 37C to an OD_{600} of ~0.6. The temperature was

subsequently lowered to 20C and the cultures were grown to OD_{600} of ~0.8 and induced with

isopropyl β-D-1-thiogalactopyranoside (IPTG) to a final concentration of 0.5mM. The cultures

were shaken for 16–18 hours at 250rpm. Cell pellets were collected by centrifugation at 6,000xg

and the pellets were stored overnight at -20C. Pellets were resuspended in 40 mL of 25mM Tris-

HCl, 300mM NaCl, pH=8.5 supplemented with a working concentration of 2mM PMSF dissolved

in isopropanol, 40mg lysozyme (0.1mg/mL) and 1 tablet of protease inhibitor. The lysates were

incubated at room temperature for 30 minutes on a platform shaker and later sonicated for 13

minutes with 10s on and 30s off 50% duty-cycle. The sonicated lysates were centrifuged for 45

minutes at 18,000xg at 4C. To precipitate the spycatcher003-mi3 nanoparticles, 170 mg/mL of

ammonium sulfate was added to the supernatant solution and subsequently shaken at 4C for 1

hour. Precipitated particles were collected by centrifuging at 16,000xg for 30 minutes at 4C. The

supernatant was discarded, and the pellet was washed with endotoxin-free water to remove

residual salty buffer. The nanoparticles were resolubilized in 8mL of 25mM Tris-HCl, 150mM

NaCl, pH=8.5. To remove insoluble aggregates, the solubilized nanoparticles were centrifuged at

18,000xg for 30 minutes. Nanoparticle expression and purity was confirmed by SDS-PAGE

analysis, and quantified spectrophotochemically at 280nm on a Nanodrop 2000 (ThermoFisher

Scientific). The spycatcher003-mi3 nanoparticles were concentrated in a 10kDa MW spin columns

(ThermoFisher Scientific) and further purified by size exclusion chromatography in 25mM Tris,

150mM NaCl, pH=8.0, on an AKTA-Go FPLC (Cytiva) using a Superose 6 Increase 10/300GL

column. Fractions containing the spycatcher003-mi3 nanoparticles were pooled and concentrated

as above. The nanoparticles were stored at -80C.

Conjugation of epitope scaffolds to Spycatcher003-mi3 nanoparticles.

Epitope scaffolds were mixed with spycatcher003-mi3 nanoparticles in a 1.2:1 molar ratio.

The complex was mixed thoroughly and incubated overnight on ice at 4C. Conjugation was

confirmed by SDS-PAGE. The conjugated nanoparticles were separated from excess monomeric

epitope scaffolds by size-exclusion chromatography in 25mM Tris, 150mM NaCl, pH 8.0, on an

AKTA-Go FPLC (Cytiva) using a Superose 6 Increase 10/300GL column. Fractions containing

the epitope-scaffold-mi3 nanoparticles were pooled and concentrated as above. Endotoxins were

removed by treating the nanoparticles with 2% triton X-100 as previously described[50]. Endotoxin

levels were tested by a chromogenic endotoxin quantification kit (ThermoFisher Scientific). The

endotoxin-free nanoparticles were stored at -80C.

Nanoparticles were analyzed by Negative Stain Electron Microscopy to confirm that they

were well formed. A frozen aliquot was thawed in RT water bath, then placed on ice. The sample

was then diluted to 400 µg/ml with 5 g/dl Glycerol in HBS (20 mM HEPES, 150 mM NaCl pH 7.4)

buffer containing 8 mM glutaraldehyde. After 5 min incubation, glutaraldehyde was quenched by

adding sufficient 1 M Tris stock, pH 7.4, to give 80 mM final Tris concentration and incubated for

5 min. Quenched sample was applied to a glow-discharged carbon-coated EM grid for 10-12

second, then blotted, and stained with 2 g/dL uranyl formate for 1 min, blotted and air-dried. Grids

were examined on a Philips EM420 electron microscope operating at 120 kV and nominal

magnification of 49,000x, and 30 images were collected on a 76 Mpix CCD camera at 2.4

Å/pixel. Images were analyzed by 2D and 3D class averages using standard protocols with Relion

3.0[51].

Mouse immunization and protection studies.

Immunization studies in BALB/c mice.

BALB/c (#028) female mice were purchased from Charles River. The Toll-like receptor 4

agonist glucopyranosyl lipid adjuvant–stable emulsion (GLA-SE) was used as the adjuvant for the

vaccine immunogens. Groups of mice (n=8) were vaccinates intramuscularly with GLA-SE-

adjuvanted Ex15-NP; GLA-SE-adjuvanted Ex_mosaic-NP; GLA-SE-adjuvanted mixture of equal

parts Ex15-NP, Ex17_NP, Ex19_NP and Ex20_NP; or GLA-SE-adjuvanted Ex15-NP (prime),

followed by Ex19_NP (boost 1) and Ex20_NP (boost 2). An adjuvant-only group (n=4) was

included for comparison. Vaccine immunogens were administered at 5µg and formulated with 5µg

of adjuvant. Mice were immunized at week 0, week 4 , and week 8. Blood samples were collected

either 7 days prior to immunization (pre-bleed), or 7 days after each immunization. Seven days

after the final dose mice were sacrificed for terminal analysis.

Live virus protection studies.

Nineteen to twenty-one female k18-hACE2 mice purchased from Jackson Laboratory

(B6.Cg-Tg(K18-ACE2)2Prlmn/J; JAX strain number #034860) were used for the WIV-1 viral

challenge studies. The Toll-like receptor 4 agonist glucopyranosyl lipid adjuvant–stable emulsion

(GLA-SE) was used as the adjuvant for the vaccine immunogens. Mouse vaccination studies

were performed intramuscularly twice with mRNA ACLNP 307 2019NCOV WUHAN S-2P spike

followed by two additional shots of either mRNA spike or GLA-SE-adjuvanted Ex_mosaic-NP.

Protein immunogens were administered at 10µg formulated with 5µg of adjuvant. mRNA

immunogens were administered at 20µg. Mice were immunized at week 0, week 4, week 8, and

week 12. Blood samples were collected either 7 days prior to immunization (pre-bleed), or 7 days

after each immunization. Mice were then moved into the BSL3 and acclimated. For infection,

vaccinated mice were anaesthetized with ketamine/xylazine and infected with 1 x 10^4 PFU WIV1-

CoV intranasally in a total volume of 50ul[52]. Infected mice were weighed daily.

Mouse studies were performed according to the recommendations for the care and use of

animals by the Office of Laboratory Animal Welfare (OLAW), National Institutes of Health, and

the Institutional Animal Care and Use Committee (IACUC) at the University of North Carolina

(UNC permit no. A-3410-01). All infectious work was performed in Biosafety Level 3 laboratories

(BSL-3) with approved standard operating procedures and safety conditions for SARS-CoV-2.

For virus titration by plaque assay, the caudal lobe of the right lung was homogenized in

phosphate-buffered saline (PBS), and the homogenate was serial diluted and inoculated onto

Vero E6 cells [American Type Culture Collection (ATCC), CRL1586], followed by agarose overlay.

Plaques were visualized with an overlay of neutral red dye on day 2 after infection. At indicated

time points, mice were euthanized and gross pathology (congestion score) of the lung tissue was

assessed and scored on a scale from 0 (no lung congestion) to 4 (severe congestion affecting all

lung lobes).

References

1 Piccoli, L. *et al.* Mapping Neutralizing and Immunodominant Sites on the SARS-CoV-2
Spike Receptor-Binding Domain by Structure-Guided High-Resolution Serology. *Cell* **183**,
1024-1042 e1021 (2020). https://doi.org:10.1016/j.cell.2020.09.037

2 Wang, Z. *et al.* mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants.
Nature **592**, 616-622 (2021). https://doi.org:10.1038/s41586-021-03324-6

3 Goel, R. R. *et al.* Distinct antibody and memory B cell responses in SARS-CoV-2 naïve and
recovered individuals after mRNA vaccination. *Science Immunology* **6**, eabi6950 (2021).
https://doi.org:10.1126/sciimmunol.abi6950

4 Robbiani, D. F. *et al.* Convergent antibody responses to SARS-CoV-2 in convalescent
individuals. *Nature* **584**, 437-442 (2020). https://doi.org:10.1038/s41586-020-2456-9

5 Starr, T. N. *et al.* Deep mutational scans for ACE2 binding, RBD expression, and antibody
escape in the SARS-CoV-2 Omicron BA.1 and BA.2 receptor-binding domains. *PLoS Pathog*
18, e1010951 (2022). https://doi.org:10.1371/journal.ppat.1010951

6 Qu, P. *et al.* Enhanced neutralization resistance of SARS-CoV-2 Omicron subvariants BQ.1,
BQ.1.1, BA.4.6, BF.7, and BA.2.75.2. *Cell Host Microbe* (2022).
https://doi.org:10.1016/j.chom.2022.11.012

7 Dacon, C. *et al.* Broadly neutralizing antibodies target the coronavirus fusion peptide.
Science **377**, 728-735 (2022). https://doi.org:10.1126/science.abq3773

8 Low, J. S. *et al.* ACE2-binding exposes the SARS-CoV-2 fusion peptide to broadly
neutralizing coronavirus antibodies. *Science* **377**, 735-742 (2022).
https://doi.org:10.1126/science.abq2679

9 Pinto, D. *et al.* Broad betacoronavirus neutralization by a stem helix–specific human
antibody. *Science* **373**, 1109-1116 (2021). https://doi.org:10.1126/science.abj3321

10 Zhou, P. *et al.* A human antibody reveals a conserved site on beta-coronavirus spike
proteins and confers protection against SARS-CoV-2 infection. *Science Translational*
Medicine **14**, eabi9215 (2022). https://doi.org:10.1126/scitranslmed.abi9215

11 Gobeil, S. M.-C. *et al.* Structural diversity of the SARS-CoV-2 Omicron spike. *Molecular Cell*
82, 2050-2068.e2056 (2022). https://doi.org:10.1016/j.molcel.2022.03.028

12 Stalls, V. *et al.* Cryo-EM structures of SARS-CoV-2 Omicron BA.2 spike. *Cell Reports* **39**,
111009 (2022). https://doi.org:10.1016/j.celrep.2022.111009

13   Dacon, C. *et al.* Rare, convergent antibodies targeting the stem helix broadly neutralize
diverse betacoronaviruses. *Cell Host & Microbe* **0** (2022).
https://doi.org:10.1016/j.chom.2022.10.010

14   Zhou, P. *et al.* Broadly neutralizing anti-S2 antibodies protect against all three human
betacoronaviruses that cause severe disease. (Immunology, 2022).

15   Azoitei, M. L. *et al.* Computation-guided backbone grafting of a discontinuous motif onto
a protein scaffold. *Science* **334**, 373-376 (2011). https://doi.org:10.1126/science.1209368

16   Correia, B. E. *et al.* Computational Design of Epitope-Scaffolds Allows Induction of
Antibodies Specific for a Poorly Immunogenic HIV Vaccine Epitope. *Structure* **18**, 1116-
1126 (2010). https://doi.org:10.1016/j.str.2010.06.010

17   Ofek, G. *et al.* Elicitation of structure-specific antibodies by epitope scaffolds. *Proceedings*
of the National Academy of Sciences **107**, 17880-17887 (2010).
https://doi.org:10.1073/pnas.1004728107

18   Ofek, G. *et al.* Elicitation of structure-specific antibodies by epitope scaffolds. *Proc Natl*
Acad Sci U S A **107**, 17880-17887 (2010). https://doi.org:10.1073/pnas.1004728107

19   Sesterhenn, F. *et al.* Trivalent cocktail of de novo designed immunogens
enables the robust induction and focusing of functional antibodies in vivo.
bioRxiv, 685867 (2019). https://doi.org:10.1101/685867

20   Li, D. *et al.* In vitro and in vivo functions of SARS-CoV-2 infection-enhancing and
neutralizing antibodies. *Cell* **184**, 4203-4219.e4232 (2021).
https://doi.org:10.1016/j.cell.2021.06.021

21   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021). https://doi.org:10.1038/s41586-021-03819-2

22   Dauparas, J. *et al.* Robust deep learning–based protein sequence design using
ProteinMPNN. *Science* **378**, 49-56 (2022). https://doi.org:10.1126/science.add2187

23   Dacon, C. *et al.* Rare, convergent antibodies targeting the stem helix broadly neutralize
diverse betacoronaviruses. *Cell Host & Microbe*
https://doi.org:10.1016/j.chom.2022.10.010

24   Liao, H.-X. *et al.* High-throughput isolation of immunoglobulin genes from single human B
cells and expression as monoclonal antibodies. *Journal of Virological Methods* **158**, 171-
179 (2009). https://doi.org:https://doi.org/10.1016/j.jviromet.2009.02.014

25   Bruun, T. U. J., Andersson, A.-M. C., Draper, S. J. & Howarth, M. Engineering a Rugged
Nanoscaffold To Enhance Plug-and-Display Vaccination. *ACS Nano* **12**, 8855-8866 (2018).
https://doi.org:10.1021/acsnano.8b02805

26   Saunders, K. O. *et al.* Neutralizing antibody vaccine for pandemic and pre-emergent
coronaviruses. *Nature* **594**, 553-559 (2021). https://doi.org:10.1038/s41586-021-03594-
0

27   Shah, P., Canziani, G. A., Carter, E. P. & Chaiken, I. The Case for S2: The Potential Benefits
of the S2 Subunit of the SARS-CoV-2 Spike Protein as an Immunogen in Fighting the
COVID-19 Pandemic. *Frontiers in Immunology* **12** (2021).
https://doi.org:10.3389/fimmu.2021.637651

28   Azoitei, M. L. *et al.* Computational design of high-affinity epitope scaffolds by backbone
grafting of a linear epitope. *J Mol Biol* **415**, 175-192 (2012).
https://doi.org:10.1016/j.jmb.2011.10.003

29   Dauparas, J. *et al.* Robust deep learning–based protein sequence design using
ProteinMPNN. *Science* **378**, 49-56 (2022). https://doi.org:doi:10.1126/science.add2187

30   Azoitei, M. L. *et al.* Computational design of protein antigens that interact with the CDR
H3 loop of HIV broadly neutralizing antibody 2F5. *Proteins* **82**, 2770-2782 (2014).
https://doi.org:10.1002/prot.24641

31   Correia, B. E. *et al.* Computational design of epitope-scaffolds allows induction of
antibodies specific for a poorly immunogenic HIV vaccine epitope. *Structure* **18**, 1116-
1126 (2010). https://doi.org:10.1016/j.str.2010.06.010

32   McLellan, J. S. *et al.* Design and characterization of epitope-scaffold immunogens that
present the motavizumab epitope from respiratory syncytial virus. *J Mol Biol* **409**, 853-
866 (2011). https://doi.org:10.1016/j.jmb.2011.04.044

33   Correia, B. E. *et al.* Proof of principle for epitope-focused vaccine design. *Nature* **507**, 201-
206 (2014). https://doi.org:10.1038/nature12966

34   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583-589 (2021). https://doi.org:10.1038/s41586-021-03819-2

35   Havenar-Daughton, C. *et al.* The human naive B cell repertoire contains distinct subclasses
for a germline-targeting HIV-1 vaccine immunogen. *Sci Transl Med* **10** (2018).
https://doi.org:10.1126/scitranslmed.aat0381

36   Leggat, D. J. *et al.* Vaccination induces HIV broadly neutralizing antibody precursors in
humans. *Science* **378**, eadd6502 (2022). https://doi.org:doi:10.1126/science.add6502

37   Zhou, P. *et al.* Broadly neutralizing anti-S2 antibodies protect against all three human
betacoronaviruses that cause deadly disease. *Immunity* **56**, 669-686.e667 (2023).
https://doi.org:10.1016/j.immuni.2023.02.005

38   Li, D. *et al.* Breadth of SARS-CoV-2 neutralization and protection induced by a nanoparticle
vaccine. *Nat Commun* **13**, 6309 (2022). https://doi.org:10.1038/s41467-022-33985-4

39   Martinez, D. R. *et al.* A broadly cross-reactive antibody neutralizes and protects against
sarbecovirus   challenge   in   mice.   *Sci   Transl   Med*   **14**,   eabj7125   (2022).
https://doi.org:10.1126/scitranslmed.abj7125

40   Battye, T. G., Kontogiannis, L., Johnson, O., Powell, H. R. & Leslie, A. G. iMOSFLM: a new
graphical interface for diffraction-image processing with MOSFLM. *Acta Crystallogr D Biol
Crystallogr* **67**, 271-281 (2011). https://doi.org:10.1107/S0907444910048675

41   Evans, P. R. & Murshudov, G. N. How good are my data and what is the resolution? *Acta
Crystallogr   D   Biol   Crystallogr*   **69**,   1204-1214   (2013).
https://doi.org:10.1107/S0907444913000061

42   McCoy, A. J. Solving structures of protein complexes by molecular replacement with
Phaser.   *Acta   Crystallogr   D   Biol   Crystallogr*   **63**,   32-41   (2007).
https://doi.org:10.1107/S0907444906045975

43   Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta
Crystallogr   D   Biol   Crystallogr*   **60**,   2126-2132   (2004).
https://doi.org:10.1107/S0907444904019158

44   Croll, T. I. ISOLDE: a physically realistic environment for model building into low-resolution
electron-density   maps.   *Acta   Crystallogr   D   Struct   Biol*   **74**,   519-530   (2018).
https://doi.org:10.1107/S2059798318002425

45   Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
structure   solution.   *Acta   Crystallogr   D   Biol   Crystallogr*   **66**,   213-221   (2010).
https://doi.org:10.1107/S0907444909052925

46   Liao, H. X. *et al.* High-throughput isolation of immunoglobulin genes from single human B
cells and expression as monoclonal antibodies. *J Virol Methods* **158**, 171-179 (2009).
https://doi.org:10.1016/j.jviromet.2009.02.014

47   Kepler, T. B. *et al.* Reconstructing a B-Cell Clonal Lineage. II. Mutation, Selection, and
Affinity   Maturation.   *Frontiers   in   Immunology*   **5**   (2014).
https://doi.org:10.3389/fimmu.2014.00170

48   Gilbert, P. B. *et al.* Immune correlates analysis of the mRNA-1273 COVID-19 vaccine
efficacy clinical trial. *Science* **375**, 43-50 (2022).
https://doi.org:doi:10.1126/science.abm3425

49   Mielke, D. *et al.* Development of flow cytometry-based assays to assess the ability of
antibodies to bind to SARS-CoV-2-infected and spike-transfected cells and mediate NK cell
degranulation. *Cytometry A* **101**, 483-496 (2022). https://doi.org:10.1002/cyto.a.24552

50   Teodorowicz, M. *et al.* Optimized Triton X-114 assisted lipopolysaccharide (LPS) removal
method reveals the immunomodulatory effect of food proteins. *PLOS ONE* **12**, e0173778
(2017). https://doi.org:10.1371/journal.pone.0173778

51   Zivanov, J. *et al.* New tools for automated high-resolution cryo-EM structure
determination in RELION-3. *eLife* **7**, e42166 (2018). https://doi.org:10.7554/eLife.42166

52   Menachery, V. D. *et al.* SARS-like WIV1-CoV poised for human emergence. *Proceedings of
the National Academy of Sciences* **113**, 3048-3053 (2016).
https://doi.org:doi:10.1073/pnas.1517719113

1119

Figure1

Figure 1. Design of epitope scaffolds that bind to broadly cross-reactive antibodies against the spike[815-823] peptide. (a) *Left*: Structure of the pre-fusion spike trimer (individual monomers: *blue, violet and pale green*; glycans: *grey*; PDBid:6xr8), with the RBD (*green*), spike[815-823] peptide (*red*) and stem helix (*orange*) domains highlighted. *Middle*: Zoom of spike monomer with spike[815-

[823] peptide highlighted (*red*) showing key residues engaged by antibodies. *Right*: Structure of

spike[815-823] peptide (*red*) bound to DH1058 mAb (*grey*) (PDBid:7tow). **(b)** Computational models

of DH1058 (*grey*) bound to ESs FP-2, FP-10 and FP-15 (*green*) with the grafted epitope shown

in *red sticks*. **(c)** Binding affinities of ESs and spike to DH1058 and DH1294 mAbs as determined

by SPR. **(d)** Binding of spike and ES to diverse spike[815-823]-targeting mAbs and their inferred

precursors. **(e)** ELISA binding of DH1058 mAb to synthetic spike[815-823] peptides and

representative ESs that contain alanine mutations at epitope residues critical for antibody

recognition. Numbers in brackets indicate the equivalent position of the mutated epitope residues

on spike. **(f)** Crystal structure of DH1058 mAb (*grey*) in complex with FP-15 (*salmon*) overlaid with

the computational model of the ESs (*green*). Epitope residues are shown in sticks (model: *red*;

crystal structure: *salmon*).

1137

Figure 2

Figure 2. Design of epitope scaffolds that bind to S2P6-like antibodies against the stem

helix. (a) The conformation of the stem helix epitope (*orange*) on the pre-fusion trimer, bound by

S2P6 and CC40.8mAbs (PDBids:7rnj and 7sjs), and displayed on representative epitope

scaffolds S2hlx-4 and -7. Key epitope residues are labeled and shown in sticks. **(b)** Binding

affinities of spike and S2hlx ESs to S2P6 and DH1057.1 mAbs **(c)** Representative single-cycle

binding SPR traces used to determine the dissociation constants in **(b)**. **(d)** The effect on S2P6

and DH1057 mAbs binding to epitope mutations in synthesized stem helix peptides (1140-1164)

and representative S2-hlx ESs. **(e)** Crystal structure of the DH1057.1 mAb (*grey*) bound to ES

S2hlx-7 (*orange*) with the epitope in *pink sticks*. Inset shows details of the epitope/mAb

interaction, with epitope residue K1149 highlighted. **(f)** Comparison of the S2hlx-7 conformation

in the computational model (*pink*) and the crystal structure (*blue*), both overall and in the epitope

region.

1152

1153   **Figure 3**

1154

Figure 3. Design of epitope scaffolds that engage two major classes of antibodies against

the stem helix. (a) Overview of the computational design process to graft the antibody-bound

CC40.8 epitope onto candidate scaffolds. ESs were designed by transplanting the epitope

backbone with Rosetta, validated with Alphafold2 and optimized for expression with

ProteinMPNN. **(b)** *Top*: Computational models of CC40.8 mAb (*gray and white*) bound to ESs

(*purple*) from the S2hlx-Ex2, S2hlx-Ex4, and S2hlx-Ex6 families of designs. The epitope is

highlighted in *orange sticks*. Residues designed by ProteinMPNN and that can differ between

designs from the same family are shown in *pink sticks*. *Bottom*: Overlay of the grafted epitope

conformation (*orange*) and the parent scaffold structure it replaced (*purple*). **(c)** Sequence identity

between S2hlx-Ex2 family of ESs designed by ProteinMPNN. **(d)** Dissociation constants of S2hlx-

Ex2 based designs to S2P6, DH1057 and CC40.8 mAbs and their precursors. *nb*=not binding. **(e)**

Binding of spike and S2hlx-Ex ESs to diverse stem helix mAbs and their precursors. **(f)** *Left*:

alignment of the S2hlx-Ex19 parent scaffold (*cyan*) with the crystal structure of the unbound

epitope scaffold (*pink*, with the epitope in *magenta sticks*). Residues Tyr62 and Phe66 that

stabilize the grafted epitope conformation are highlighted. *Right*: Comparison of the stem helix

epitope conformation in the pre-fusion spike (*grey*), induced by CC40.8 binding (*orange*) and

displayed on S2hlx-Ex19 (*magenta*).

1172

Figure 4

1174

Figure 4. Reactivity of engineered epitope scaffolds with sera from individuals with pre-existing immune responses to SARS-CoV-2 spike. (a) *Left*: ELISA binding of sera or plasma from patients with pre-existing SARS-CoV-2 immunity acquired by vaccination (*red*), infection (*green*), or vaccination followed by infection (*purple*) to recombinant WA-2 spike and RBD or to synthetic peptides encoding the spike[815-823] peptide (residues 808-833) or the stem helix peptide (1140-1164). *Middle* and *right:* Binding of the same samples in **(a)** to select stem helix peptide

and spike$^{815\text{-}823}$ peptide epitope scaffolds respectively. **(b)** Comparison between sera binding to

the synthesized spike$^{815\text{-}823}$ peptide with FP-15 ES, and the parent scaffold from which FP-15 was

derived and that lacks the grafted epitope (FP-15 PS). **(c)** Comparison between sera binding to

the synthesized stem helix peptide with S2hlx-Ex19 ES, and with versions of this design where

the epitope is mutated to reduce binding to either CC40.8 class antibodies (L1145A), to both

CC40.8 and S2P6 classes of antibodies (F1148A), or where the epitope is replaced with the

sequence from the parent scaffold it replaced (PShlx). Single lines represent measurements for

individual samples in the respective cohort. For all graphs *=$p<0.05$ and **=$p<0.01$.

1189

Figure 5

1191

Figure 5. Isolation and characterization of epitope scaffold reactive antibodies from subjects with pre-existing SARS-CoV-2 immunity. (a) Sera binding profile of subjects from which B cells were isolated. Binding was measured to the synthetic peptide epitope, the epitope scaffold used as the sorting probe, and the corresponding parent scaffold (PS) that lacks the

epitope and was employed for negative selection. **(b)** Representative FACS plots of memory B

cells labeled with epitope scaffold probes. Cells from Subject 26 were labeled with S2hlx-Ex19

conjugated to VB15 (x-axis) and AF647 (y-axis). A third bait comprising S2hlx-Ex19 PShlx

conjugated to BV421 was used to exclude scaffold-specific memory B cells. **(c)** Binding of the

isolated antibodies to diverse CoV spikes. **(d)** Neutralization of diverse pseudoviruses by isolated

stem helix antibodies. **(e)** Binding of isolated antibodies to cells infected with different SARS-CoV-

2 VOCs. Antibodies in *red* have been previously described, those depicted in *green* were isolated

here, while those in *blue* represent controls; DH1047 targets the RBD domain; DH1058 targets

the spike[815-823] peptide domain; anti-HIV antibody VRC01 was included as a negative control. **(d)**

NK cell degranulation ability of mAbs from **(e)**.

1206

Figure 6

Figure 6. Immunogenicity of stem helix epitope scaffolds. (a) Development and NSEM

analysis of the spycatcher/spytag mi03 nanoparticles displaying 60 epitope scaffold copies. **(b)**

Study design to determine the immunogenicity of stem helix nanoparticles in BALB/c mice. **(c)**

Binding of sera from animals immunized with the different epitope scaffold regimens against the

WA-2 spike at different study time points. **(d)** Sera binding after the second boost against diverse

CoV spikes. Values are calculated as the logarithm of the area under the curve for binding curves

measured as in (c). **(e)** Binding of S2P6 mAb and sera from animals immunized with Ex_mosaic

to the WT stem helix epitope peptide and mutated variants. **(f)** Study design to determine the

protection ability of Ex_mosaic-NP against live coronavirus challenge. **(g)** Percentage weight loss

in animals treated as in (f). Values represent the average per group, with error bars depicting the

standard deviation. **(h)**. Lung congestion score and plaque analysis of infected animals. Data

points correspond to individual animals. ns=not significant; **=p<0.01; ***=p<0.001 ****=p<0.0001

1221