TEAD1 is crucial for myelination, Remak bundles and functional regeneration of peripheral nerves

Matthew Grove[1], Hyukmin Kim[1], Shuhuan Pang[1], José Paz Amaya[2], Guoqing Hu[3], Jiliang Zhou[3], Michel Lemay[2], and Young-Jin Son[1]*

[1]Department of Neural Sciences, Shriners Hospitals Pediatric Research Center, Lewis Katz School of Medicine, Temple University, Philadelphia, PA 19140

[2]Department of Bioengineering, Temple University, Philadelphia, PA 19122, USA

[3]Department of Pharmacology & Toxicology, Medical College of Georgia, Augusta University, Augusta, GA

* Corresponding author: yson@temple.edu

ABSTRACT

The hippo pathway transcriptional effectors, YAP/TAZ, are crucial for Schwann cells (SCs) to myelinate axons but the mechanisms are poorly defined. Although TEAD1 has been implicated as a partner transcription factor, how it supports the transcriptional regulation of myelination and which aspects of the process it affects are unknown. Here, using conditional and inducible knockout mice, we report that TEAD1 is essential for SCs to develop, grow, and regenerate myelin sheaths. TEAD1/2/3/4 are present in SCs, but YAP/TAZ strongly favors TEAD1. It promotes myelination by positively and negatively regulating SC proliferation, enabling Krox20/Egr2 to upregulate myelin proteins, and upregulating the cholesterol biosynthetic enzymes FDPS and IDI1. We also show stage-dependent redundancy of TEAD1 and that non-myelinating SCs have a unique requirement for TEAD1 to enwrap nociceptive axons in Remak bundles. Our findings establish TEAD1 as a crucial partner of YAP/TAZ in developmental myelination and functional nerve regeneration and as a novel transcription factor regulating Remak bundle integrity.

ONE SENTENCE SUMMARY

TEAD1 in Schwann cells is a major transcriptional factor regulating peripheral nerve myelination and Remak bundle formation.

INTRODUCTION

In the mammalian peripheral nervous system, all motor and sensory axons are associated with Schwann cells (SCs). Axons larger than 1μm diameter are enveloped by myelinating SCs that form multilamellar myelin sheaths with a thickness proportional to axon caliber (*1, 2*). Axons smaller than 1μm diameter, the unmyelinated nociceptors (i.e., C-fibers), are organized in clusters called Remak bundles, in which each C-fiber is ensheathed by a cytoplasmic pocket formed by non-myelinating SCs, also known as Remak SCs (*3, 4*). Both myelinating and non-myelinating SCs provide metabolic and functional support essential for structural and functional integrity of peripheral nerves. Failure to initiate or maintain proper axon-SC association may lead to life-threatening paralysis and peripheral neuropathy (*5*). Myelinating SCs form and

maintain myelin via a set of key transcription factors (TFs) and regulators (*6-12*). However, the

transcriptional regulation of myelinating SCs remains incompletely understood, and that of non-

myelinating SCs is largely unknown (*4*).

Recent mouse genetic studies surprisingly revealed that YAP (Yes-associated protein)

and its paralogue, TAZ (Transcriptional coactivator with PDZ-binding motif), which are potent

oncoproteins and transcriptional regulators in the Hippo signaling pathway, play crucial roles in

developing, maintaining, and regenerating myelination of peripheral nerves. Both YAP and TAZ

(YAP/TAZ) are expressed initially in all immature SCs and then selectively in myelinating SCs

(*13*). The processes for which they are required include: timely sorting of large axons and

associated SC proliferation (*13, 14*); initiation of SC differentiation or myelination (*13, 15*);

myelin maintenance in adults (*13*); and remyelination following nerve injury (*16, 17*). YAP/TAZ

initiate and maintain myelination in part by upregulating Krox20 (a.k.a., Egr2) (*13, 18*), which is

widely recognized as the master TF promoting peripheral myelination. Krox20 is required to

initiate myelination, maintain adult myelin (*7, 19*), and likely to promote myelin sheath growth

(*20*). It acts through direct transcriptional upregulation of myelin proteins and membrane lipids

(*21-23*).

YAP/TAZ are transcriptional co-activators that contain a transcriptional activation

domain, but no DNA binding domain. They regulate gene expression by binding to partner TFs,

typically members of the TEA domain family (TEAD1-4) (*24*). Conversely, TEADs on their

own are unable to activate or repress transcription (*25*) and require the help of co-factors such as

YAP/TAZ to act (*26*). Partner TFs such as TEADs are required to maintain YAP/TAZ in the

nucleus, which keeps them transcriptionally active (*27-30*). Transcript data indicates that SCs

express all four TEADs and that their expression peaks at different times during development

(*31*). Moreover, TEAD1-4 show unique, redundant, and/or opposing roles in regulating

proliferation and differentiation of various cell types; these roles can also change as a function of

developmental stage (*26, 32-34*). It remains to be elucidated how and which TEAD(s) in SCs

mediate transcriptional regulation of YAP/TAZ for the development, homeostasis, and/or

regeneration of myelination.

In some cells, TEAD1 serves as the main partner TEAD and plays non-redundant roles

(*35, 36*). In SCs, ChiP, co-IP, and transcription assays have consistently indicated that TEAD1

associates with YAP/TAZ for upregulating myelin proteins (*13, 18, 37, 38*). Notably, unlike

YAP/TAZ, TEAD1 is expressed in all SC nuclei (*13*), suggesting that, independently of

YAP/TAZ, TEAD1 plays an unknown role in non-myelinating SCs. In this study, we generated

conditional and inducible knockout mice lacking TEAD1 in SCs to investigate the roles of

TEAD1 in myelinating and non-myelinating SCs. We found that TEAD1 is uniquely required for

myelinating SCs to initiate, grow and regenerate myelination, and for non-myelinating SCs to

enwrap C-fibers in Remak bundles during development and homeostasis. We also uncovered the

mechanistic basis of the non-redundancy of TEAD1 and found that YAP/TAZ-TEAD1 promotes

myelination by dually regulating SC proliferation, by enabling induced Krox20 to upregulate

myelin proteins and by upregulating cholesterol biosynthetic enzymes. Additional results suggest

that other TEADs may play redundant and opposing roles in a stage-dependent manner.

RESULTS

Multiple TEADs are expressed in Schwann cells of WT, *Tead1* cKO and iKO mice

To investigate the roles of SC TEAD1 in developing nerves, we conditionally ablated

Tead1 by crossing *Tead1*$^{fl/fl}$ mice (*39*) with mice carrying *Mpz*-Cre, which is active from E13.5 in

SCs (*40*). To investigate the roles of TEAD1 in maintaining and regenerating nerves in adults,

we inducibly inactivated SC *Tead1* by crossing *Tead1*$^{fl/fl}$ mice with mice carrying a tamoxifen-

inducible *Plp*-CreER (*41*). Immunostaining of P22 WT and *Mpz-Cre;Tead1*$^{fl/fl}$ (hereafter *Tead1*

cKO) sciatic nerves (SNs) showed that TEAD1 was expressed in the nuclei of 100% of WT SCs

and was undetectable in >97% *Tead1* cKO SCs (Figure 1A). Immunostaining of WT and *Plp* -

CreER*; Tead1*$^{fl/fl}$ (hereafter *Tead1* iKO) SNs after tamoxifen treatment showed that TEAD1 was

also present in 100% of SC nuclei in both intact and regenerating nerves of adult WT mice and

was deleted in almost all *Tead1* iKO SCs (Figure 1B, 1C). The specificity of the TEAD1

immunostaining and highly efficient deletion were further confirmed by Western blotting (Figure

1D). These results suggest potential roles for TEAD1 in both myelinating and non-myelinating

SCs during development, maintenance and after nerve injury.

To investigate the expression of other TEADs in WT, *Tead1* cKO and *Tead1* iKO SCs,

we used antibodies to TEAD2, 3 and 4, and a pan-TEAD antibody marketed to recognize all four

TEADs (TEAD1/2/3/4). Western blot analysis of SN lysates showed that anti-TEAD3 and anti-

pan TEAD both revealed a main ~ 50kD band that was decreased in *Tead1* cKO (Figure 1D).

The pan-TEAD antibody may therefore bind preferentially to TEAD3. Anti-TEAD4 recognized

three bands in WT and showed that the two faster-migrating bands were decreased in *Tead1* cKO

(Figure 1D). Anti-TEAD2 recognized a band of ~65 kD in *Tead1* cKO that was barely detectable

in WT lysates. This band corresponded to the size of TEAD2 reported in other cell types (*42*).

We also found that the ~65 kD band was present early postnatally, but rapidly diminished with

age (data not shown), consistent with previous transcript data from mouse SCs (*31*). Therefore,

the expression of TEAD2-4 was altered in *Tead1* cKO, confirming the specificity of the

antibodies and their expression in SCs. Of these anti-TEAD antibodies, only anti-pan TEAD

functioned effectively for immunofluorescence staining, verifying the presence of other TEADs

in SC nuclei in both intact (Figure 1E) and injured WT and *Tead1* iKO SNs (Figure 1F). These

results suggest that WT SCs express all four TEADs and that *Tead1* cKO and iKO SCs express

TEAD2-4 in their nuclei.

YAP/TAZ strongly favor TEAD1 but partner with other TEADs in the absence of TEAD1

In the absence of a partner TF, nuclear YAP/TAZ are decreased and phosphorylation of

YAP/TAZ increased, due to targeting by Large tumor suppressor kinase 1/2 (LATS1/2),

negative regulators of YAP/TAZ (*43*). To investigate if YAP/TAZ are in the nuclei of SCs and

therefore transcriptionally functional in the absence of TEAD1, we compared western blotting of

P50 WT vs *Tead1* cKO SN lysates and observed an overall decrease in YAP levels in *Tead1*

cKO but a > 2-fold increase in TAZ levels compared to WT (Figure 2A). This ratio of YAP to

TAZ in *Tead1* cKO resembled that observed during developmental myelination in WT SNs (*13,*

15). Furthermore, phosphorylation of YAP decreased in cKO, which would typically indicate

increased YAP activity. These results suggest that YAP/TAZ are not targeted for destruction in

the absence of TEAD1. Indeed, YAP/TAZ were retained in the SC nucleus of *Tead1* cKO

(Figure 2B) and both intact and injured *Tead1* iKO SNs (Figure 2C, 2D), as strongly supported

by quantitative analysis (Figure 2E).

To investigate whether other TEADs could be complexed with YAP/TAZ, thereby

retaining YAP/TAZ in SC nuclei in the absence of TEAD1, we carried out YAP/TAZ co-

immunoprecipitation (IP) experiments comparing WT vs *Tead1* iKO, and blotted for YAP/TAZ,

TEAD1 and pan-TEAD (Figure 2F). After verifying equal amounts of lysate samples and the

efficiency of IPs with post-IP lysates, we observed that TEAD1 strongly co-IPed with YAP/TAZ

from WT lysates and that levels were greatly reduced in *Tead1* iKO lysates. Blotting with anti-

pan-TEAD showed that multiple TEADs were co-IPed with YAP/TAZ from both WT and *Tead1*

iKO lysates. Notably, the pan-TEAD co-IPed from WT lysates, like TEAD1, migrated more

slowly than that from *Tead1* iKO lysates, whereas the *Tead1* iKO pan-TEAD migrated similarly

to the major pan-TEAD band in post-IP lysates. Longer exposure of the pan-TEAD film revealed

the presence of a fainter band migrating above the main pan-TEAD band in WT lysates (Figure

2F; arrow). Because this band co-migrated with TEAD1 and was not present in *Tead1* iKO

lysates, it was identified as TEAD1. These results suggest that TEAD1 makes up a very minor

portion of that revealed by pan-TEAD in WT lysates, but the overwhelming majority of that

revealed by pan-TEAD in YAP/TAZ IPs from these lysates. Therefore, although multiple

TEADs are present, TEAD1 is greatly favored as the binding partner for YAP/TAZ. However, in

the absence of TEAD1, other TEADs are associated with YAP/TAZ presumably to substitute for

some or all the functions of the YAP/TAZ-TEAD1 transcriptional complex.

TEAD1 is required for myelin sheath formation and growth

Tead1 cKO mice were smaller than WT mice at all ages, with labored breathing and

severely splayed fore- and hindlimbs, indicative of severe peripheral neuropathy (Figure 3A,

Supplementary Video 1). The hindlimbs never became completely paralyzed, however, in

contrast to *Yap/Taz* cDKO mice (*13*). *Tead1* cKO mice were humanely euthanized before P60.

Tead1 cKO SNs were noticeably translucent, indicative of reduced myelin content (Figure 3B).

Electrical stimulation of cKO SNs elicited low and dispersed compound muscle action potentials

(CMAPs) (Figure 3C). Nerve conduction velocity (NCV) in cKO was ~8 m/s, which was

markedly reduced from ~40 m/s in WT mice (Figure 3D).

Next, we used semi-thin and transmission electron microscopy (TEM) to examine the

processes of developmental myelination and Remak bundle formation. We first examined SNs at

P3, when, in WT, many large axons had already been radially sorted, establishing 1:1

relationship with promyelinating SCs, and myelination had begun. Compared to WT, *Tead1* cKO

SNs were thinner, unsorted bundles of axons were larger and more numerous, and myelination

was minimal (Figure 3E). Nevertheless, we frequently observed large axons that had been sorted

and established a 1:1 relationship with promyelinating SCs, as in WT. Therefore, radial sorting

was delayed in *Tead1* cKO, though less severely than in *Yap/Taz* cDKO mice, in which no axons
were sorted at P4 (*13*).

At P22, when WT SNs were fully myelinated, the phenotype of *Tead1* cKO SNs was
complex and included areas of unsorted axon bundles, sorted but unmyelinated axons, and thinly
myelinated axons (Figure 3F, Supplementary figure 1A). Notably, at P50, this phenotype was
unchanged from that at P22: areas of unsorted bundles, promyelination, and thin myelination
persisted (Figure 3G, Supplementary figure 1A). TEM analysis confirmed that myelin was
indeed thinner in *Tead1* cKO than in WT at both P22 and P50 (Figure 3F, 3G, Supplementary
figure 1B). The g-ratio did not significantly change in either WT or cKO between P22 and P50
(Figure 3H), suggesting that growth of the myelin sheath relative to axon diameter did not
increase over this period. These data suggest that myelination in *Tead1* cKO was arrested rather
than delayed. Plots of g-ratio versus axon diameter showed that the mutant defect was present in
axons of all diameters but was particularly notable in those of larger diameter (Figure 3I, 3J).
The mean axon diameter was also reduced in cKO at P22, though it increased with age (Figure
3K). These data show that TEAD1 is required, like YAP/TAZ (*13*), for timely radial sorting and
initiation of myelination during development. Although some axons become myelinated in the
absence of TEAD1, presumably due to other TEADs, they require TEAD1 for myelin sheath to
properly grow in proportion to axon diameter.

TEAD1 is required for proper formation of Remak bundles
Remak bundle formation was also abnormal in *Tead1* cKO mice. In WT SNs at P22 and
P50, small diameter axons were present in Remak bundles, with each C-fiber ensheathed by SC
cytoplasm (Figure 3F, 3L). In *Tead1* cKO, C-fibers were also present in bundles in which axons
were rarely larger than 1 μm in diameter, suggesting that they were almost fully sorted.
However, at both P22 and P50, the observed bundles were frequently small, often the size of a
large myelinated axon. Notably, a SC surrounded each bundle but individual C-fibers of most
Remak bundles were not ensheathed by SC cytoplasm (Figure 3L). Furthermore, these bundles
were frequently and abnormally thinly myelinated. Quantification showed that ~80% of C-fibers
were not ensheathed by SC cytoplasm in P50 *Tead1* cKO SNs (Figure 3M). Therefore, TEAD1
is required to develop proper association of C-fibers with non-myelinating SCs in Remak
bundles.

TEAD1 regulates SC proliferation both positively and negatively to promote myelination

For myelination and Remak bundle formation to occur properly, timely generation of sufficient numbers of SCs is essential. YAP/TAZ are necessary for optimal SC proliferation in the late embryonic and early postnatal periods (*13, 15, 44*). To determine if the defects of myelination and Remak bundles in *Tead1* cKO are due to defective SC proliferation, we counted SCs. At P3, SC number in cKO SNs was about half that of WT (Figure 4A), slightly more than in *Yap/Taz* cDKO at P4 (*13*). The number of SCs in the cell cycle (Ki67-positive; Figure 4B) and in S-phase (EdU-positive; Figure 4C) was greatly reduced in cKO at P3. Therefore, TEAD1 is required for optimal SC proliferation at P3, suggesting that the YAP/TAZ-TEAD1 complex promotes SC proliferation at early postnatal periods.

At P22 and P50, however, SC numbers in *Tead1* cKO SNs were greatly increased and became more than twice those of WT (Figure 4A); a similar increase was not observed in *Yap/Taz* cDKO (*13*). As *Tead1* cKO SC numbers were similar at P22 and P50, we reasoned that between P3 and P22, SC proliferation must occur at an abnormally high rate in this mutant. Indeed, the number of SCs that were Ki67-positive or EdU-positive was significantly increased in cKO at P22 (Figure 4B, 4C). The increased SC numbers were not due to proliferation of residual non-deleted SCs: whereas in WT, EdU-positive SCs contained TEAD1, TEAD1 was absent in most EdU-positive SCs in *Tead1* cKO (Figure 4D). During development, therefore, SCs switch from requiring TEAD1 for optimal proliferation, to requiring TEAD1 for timely exit from the cell-cycle, which is a prerequisite for SC differentiation and myelination (*45*).

TEAD1 is required for Krox20 to upregulate myelin proteins, but not for Krox20 upregulation

Our findings thus far show that, in the absence of TEAD1 in SCs, other TEADs can complex with YAP/TAZ. Nevertheless, only some axons are myelinated abnormally thinly in *Tead1* cKO which grossly phenocopied *Yap/Taz* iDKO. These defects are not because *Tead1* cKO mice lack SCs. *Yap/Taz* cDKO exhibited a complete failure of myelination due to arrest of their SCs as promyelinating SCs, in part because they fail to upregulate Krox20/Egr2 (Early Growth Response 2), which drives transcription of myelin proteins (*13*). To determine whether TEAD1 is required for Krox20 upregulation, we first investigated Oct6 (Pou3f1, Scip), whose

upregulation precedes and is required for Krox20 upregulation; Oct6 is subsequently

downregulated, possibly by Krox20 in a feedback mechanism (*46, 47*). Western blotting of P50

SN lysates showed minimal expression of Oct6 in WT, but greatly upregulated expression in

Tead1 cKO (Figure 5A). Immunohistochemistry of P50 cKO SNs showed intense Oct6

expression in ~80% of SC nuclei; Oct6 expression was minimal in WT (Figure 5B). Notably, at

both P11 and P50, a similar percentage of SCs expressed Krox20 in WT and *Tead1* cKO SNs

(Figure 5C). In Western blots, Krox20 was upregulated >4-fold in cKO compared to WT (Figure

5A). As adult *Tead1* cKO mice contain ~2-fold more SCs than WT (Figure 4A), these data

suggest that each SC in *Tead1* cKO contains ~2-fold more Krox20 than WT. Thus, in contrast to

SCs lacking YAP/TAZ (*13*), SCs lacking TEAD1 upregulate Krox20, suggesting that TEAD1 is

dispensable for YAP/TAZ to upregulate Krox20.

Consistent with the idea that Krox20 is pivotal for myelin gene transcription, expression

of myelin basic protein, MBP, and myelin protein zero, MPZ, was abundant in WT P50 SNs

(Figure 5A). In contrast, despite ample expression of Krox20, both MBP and MPZ were

markedly decreased in *Tead1* cKO, suggesting that Krox20 fails to properly upregulate myelin

proteins without TEAD1. To exclude the possibility that Krox20 expressed in cKO SCs is

defective, we took advantage of the earlier finding that loss of Krox20 or its co-factors, Nab1/2,

causes SCs to remain aberrantly in the cell-cycle (*12, 47*). Therefore, we studied whether EdU-

positive SCs in the cell-cycle of P11 cKO were also positive for Krox20, which would indicate

defective Krox20. We found that, although ~65% of SC nuclei contained Krox20, only a few

EdU-positive SCs exhibited even minimal Krox20 immunoreactivity in either WT or *Tead1* cKO

(Figure 5D). Thus, despite high levels of functional Krox20, myelin protein expression was

greatly reduced in cKO SCs, suggesting that TEAD1 is dispensable for Krox20 upregulation but

required for Krox20 to upregulate myelin proteins.

TEAD1 and YAP/TAZ upregulate cholesterol biosynthetic enzymes, FDPS and IDI1

Myelination of peripheral nerves by SCs requires a large amount of lipid and cholesterol

biosynthesis. De novo biosynthesis of fatty acids (integral building blocks of lipids) and

cholesterol is crucially important for SCs to initiate myelination and to grow a myelin sheath

(*48-51*). YAP/TAZ may regulate these pathways, as suggested by reduced mRNA expression in

Yap/Taz cDKO of SREBPs TFs (Sterol regulatory element binding proteins) and their target

enzymes involved in lipid/cholesterol biosynthesis (*14, 15*). We therefore investigated, at the

protein level, whether TEAD1 and YAP/TAZ are required for fatty acid and/or cholesterol

synthesis. To test this premise, we examined expression of SREBP1 and its target SCD1

(Stearoyl-CoA desaturase-1), which upregulates fatty acid synthesis, and of SREBP2 and its

targets HMGCR (HMG-CoA reductase), FDPS (Farnesyl Diphosphate Synthase) and IDI1

(Isopentenyl-diphosphate delta isomerase 1), which upregulate cholesterol synthesis (*51, 52*).

Because SREBP pathways are upregulated early postnatally for myelination in WT SNs, their

upregulation may simply be delayed in developmentally delayed *Tead1* cKO. We therefore

investigated SREBP pathways both at P8 and P40 in WT and *Tead1* cKO mice. Western blot

analysis showed that SREBP1 and SCID1 expression in cKO SNs was not significantly affected

by loss of TEAD1 at either P8 or P40 (Figure 6A). In contrast to SREBP1, SREBP2 was ~50%

reduced in *Tead1* cKO at both P8 and P40. Despite marked reduction in SREBP2, HMGCR was

not reduced at P8 or P40. However, both FDPS and IDI1 levels were greatly reduced at both P8

and P40 (Figure 6A).

It is possible that HMGCR was not reduced in cKO due to TEAD1 redundancy and

compensation by other TEADs. If so, then HMGCR levels should be reduced in *Yap/Taz* cDKO,

because YAP/TAZ are the primary partners for all TEADs (*24, 32*). To test this idea, we

examined SREBP pathways in *Yap/Taz* cDKO SNs at P60 because developmental delay is

greater in *Yap/Taz* cDKO than in *Tead1* cKO SNs (*13*). We found that, as in *Tead1* cKO,

SREBP1 and SCD1 levels were not noticeably reduced by the loss of YAP/TAZ in cDKO

(Figure 6B), suggesting that YAP/TAZ and TEAD1 do not directly affect fatty acid synthesis.

Unlike in *Tead1* cKO, levels of SREBP2 were not markedly reduced in Yap/Taz cDKO. As in

Tead1 cKO, however, HMGCR levels were not reduced in Yap/Taz cDKO, whereas levels of

FDPS and IDI1 were greatly reduced (Figure 6B). These results therefore suggest that

YAP/TAZ-TEAD1 transcriptional complex selectively upregulates IDI1 and FDPS, contributing

to cholesterol biosynthesis. Together with the lack of synthesis of myelin proteins, this reduction

in cholesterol biosynthesis may prevent myelin formation and growth in *Tead1* cKO.

Loss of TEAD1 in adult Schwann cells modestly affects myelin maintenance

Because TEAD1 regulates myelination and Remak bundle formation during

development, we next investigated its role in myelin and Remak bundle maintenance in adults.

We first verified efficient deletion of TEAD1 in almost all SCs of *Tead1* iKO (Figure 1B).

Unlike *Yap/Taz* iDKO mice (*13*), *Tead1* iKO mice appeared normal 2 weeks after tamoxifen

administration and continued to survive without obvious signs of peripheral neuropathy up to 5

months post-tamoxifen, the longest time of observation (Figure 7A). Analysis of iKO SNs

between 3 to 5 months post-tamoxifen revealed that they were grossly normal. However, we

occasionally observed large axons undergoing demyelination or completely demyelinated

(Figure 7B). Unlike *Yap/Taz* iDKO mice, which frequently exhibited segmental demyelination

and markedly reduced NCV (*13*), NCV of these iKO mice did not significantly differ from that

of WT (Figure 7C). Therefore, loss of TEAD1 alone in adult SCs evoked only modest

demyelination that does not cause significant functional impairment.

TEAD1 is required for maintaining structural integrity of Remak bundles

Next, we examined Remak bundles in these mice. As in WT SNs, C-fibers in WT mice

were individually ensheathed by SC cytoplasm. In contrast, in *Tead1* iKO, many c-fibers in most

Remak bundles were not separated by SC cytoplasm (Figure 7D). Furthermore, some Remak

bundles were abnormally myelinated, as in *Tead1* cKO SNs (Figure 7E). Thus, TEAD1 is

required for formation as well as maintenance of Remak bundles.

TEAD1 is dispensable for YAP/TAZ to maintain Krox20, FDP and IDI1 expression

To investigate why the demyelinating phenotype of *Tead1* iKO is much milder than that

of *Yap/Taz* iDKO, we first investigated expression of Krox20 which is required for myelin

maintenance (*7*). We previously observed that Krox20 is markedly downregulated in *Yap/Taz*

iDKO SCs, which correlated with frequent segmental demyelination and markedly reduced NCV

(*13*). As a control, we injected *PLP-CreER; Yap^{fl/fl}; Taz^{fl/fl}* mice with tamoxifen and reproduced

our previous result: Krox20 was dramatically downregulated in most *Yap/Taz* iDKO SCs only 2

weeks post-tamoxifen (Figure 7F). In contrast, in *Tead1* iKO, Krox20 expression was

indistinguishable from WT even at 6 weeks post-tamoxifen (Figure 7G). We then investigated

Oct6 expression, because aberrant overexpression of Oct6 in the presence of Krox20 inhibits

myelination (*46*). Unlike in *Yap/Taz* iDKO SNs (*13*), however, Oct6 was not elevated in *Tead1*

iKO SNs (Figure 7H). Therefore, TEAD1 is dispensable for maintaining Krox20 expression in

myelinating SCs, consistent with the only modest demyelinating phenotype of *Tead1* iKO.

Next, we investigated SREBP pathways, which are required for developmental myelination (*50, 51*). Whether these pathways are also involved in myelin maintenance has not been studied. We first assessed the effects of acute loss of YAP/TAZ on these pathways in adult SCs. Western blotting of WT and *Yap/Taz* iDKO SN lysates showed that the SREBP1 pathway, or expression of SREBP2 and HMGCR in the SREBP2 pathway, was unaffected in *Yap/Taz* iDKO (Figure 7I). However, FDPS and IDI1 levels were significantly downregulated in *Yap/Taz* iDKO (Figure 7I), recapitulating the effect of developmental loss of YAP/TAZ on these proteins (Figure 6B). Therefore, the acute downregulation of these key enzymes for cholesterol biosynthesis correlates with the acute demyelination previously observed in *Yap/Taz* iDKO mice. This result suggests that YAP/TAZ maintain myelination through both myelin gene expression and the cholesterol biosynthesis pathway.

Western blotting of *Tead1* iKO SN lysates, 6 weeks post-tamoxifen, revealed that the SREBP pathways were not significantly affected by loss of TEAD1 (Figure 7J), unlike *Tead1* cKO in which FDPS and IDI1 were markedly downregulated. This result correlated with the modest effect of loss of TEAD1 on myelin maintenance, suggesting that TEAD1 is largely dispensable for cholesterol synthesis in adult SCs.

TEAD1 is required for myelination of regenerating axons after nerve injury.

Next, we investigated whether TEAD1 is required for axon regeneration and remyelination after nerve injury. We previously showed that YAP/TAZ are not required for axon regeneration but are necessary for remyelination (*13*). Because *Yap/Taz* iDKO mice die ~2 weeks post-injury, it remains possible that remyelination is delayed rather than completely blocked. In contrast, *Tead1* iKO mice are not susceptible to early death. We, therefore, examined them 4 weeks after crushing SN, when hindlimb muscles are fully reinnervated in WT (Figure 8A). We first verified efficient deletion of TEAD1 in almost all SCs of *Tead1* iKO mice after nerve crush (Figure 1C). Longitudinal sections of iKO SNs showed numerous TuJ1-labeled axons that regenerated along the distal nerve stump, as in WT SNs (Figure 8B). However, MBP immunoreactivity was markedly reduced in the distal nerve stump of *Tead1* iKO, compared to WT, suggesting that axons regenerated but were not remyelinated. We also found that regenerated axons reached hindlimb muscles and reinnervated neuromuscular junctions (NMJs) in iKO, as in WT (Figure 8C). Moreover, *Tead1* iKO frequently exhibited features of paralyzed

muscles, such as axon terminals that grew past the synaptic area, fragmented clusters of

acetylcholine receptors (AChRs), and satellite extra-synaptic AChRs (*53*). Semithin and EM

analysis confirmed virtual absence of myelination in iKO SNs; although large axons were

associated with SCs, they were not myelinated (Figure 8D). Accordingly, electrical stimulation

of iKO SNs did not elicit CMAPs or NCV (Figure 8E). Thus, like YAP/TAZ, TEAD1 in SCs is

not required for axon regeneration but is required for remyelination after nerve injury.

369   After injury, SCs transdifferentiate into growth-promoting repair SCs, which then

proliferate and reach a peak around 3 dpi (*54-56*). Quantitative analysis of SC proliferation at 3

dpi by counting EdU$^+$/Sox10$^+$ SCs showed that a similar percentage of SCs were in S-phase in

both WT and *Tead1* iKO SNs (Figure 8F), suggesting that, like YAP/TAZ (*13*), TEAD1 is not

required for proliferation of repair SCs. Lastly, we counted Krox20$^+$ and Oct6$^+$ SCs at 4 wpi.

Notably, in contrast to *Yap/Taz* iDKO SCs, which fail to upregulate Krox20 after injury (*13*),

Krox20 was upregulated in *Tead1* iKO SCs, as in WT SCs (Figure 8G). Furthermore, Oct6 was

aberrantly elevated in *Tead1* iKO SCs, like in *Tead1* cKO SCs during development (Figure 8H).

These results suggest that, similar to developmental myelination, remyelination requires TEAD1

for Oct6 downregulation and for Krox20 to upregulate myelin proteins. Additionally, almost

complete absence of remyelination at 4 wpi highlights that TEAD1 is crucial for functional

regeneration of peripheral nerves.

DISCUSSION

The present study conclusively establishes TEAD1 as a crucial TF that partners with YAP/TAZ

to mediate myelin formation, growth, and regeneration. TEAD1 promotes myelination by dually

regulating SC proliferation, by enabling induced Krox20 to upregulate myelin proteins, and by

upregulating cholesterol biosynthetic enzymes. Our findings also suggest that other TEADs play

redundant and opposing roles in SCs, and that TEAD1 is critical in the proper formation and

maintenance of Remak bundles.

Unique, redundant, and opposing roles of TEAD1

Although SCs express all four TEADs, *Tead1* cKO mice grossly phenocopies *Yap/Taz* cDKO

mice. They exhibit abnormal SC proliferation, delayed radial sorting, blocked myelin formation

and remyelination, and normal axon regeneration. Furthermore, *Tead1* cKO mice, like *Yap/Taz*

cDKO, are defective in regulating Oct6, Krox20, myelin proteins, and enzymes that synthesize

cholesterol. These similarities show that YAP/TAZ regulate both SC proliferation and

differentiation, the two opposing processes crucial for myelination, by partnering mainly with

TEAD1 at least for developmental myelination and remyelination after injury.

We also observed multiple differences between *Tead1* cKO and *Yap/Taz* cDKO mutants. Unlike

Yap/Taz cDKO, *Tead1* cKO initiates myelin formation in some developing axons and maintain

myelination in most axons but fails to remyelinate almost all regenerating axons. Additionally, in

Tead1 cKO, Krox20 are upregulated and a large excess of SCs is generated. These differences

are likely due to stage-dependent redundancy and opposing roles of TEADs. Consistent with this

idea, YAP/TAZ functionally associate with TEAD2, 3 or 4, or an unrelated TF (*57*), in the

absence of TEAD1. Loss of TEAD1 does not affect nuclear localization of YAP/TAZ, which

depends mainly on interactions with TEADs (*27-30*). Our data and the results of others suggest

that TEAD1-4 are all co-expressed in SCs through multiple stages of nerve development,

homeostasis and after injury (*13, 34, 37*). Notably, nuclear retention of YAP/TAZ occurs in

Tead1 cKO when TEAD1 has a unique role in SCs (i.e., when *Tead1* cKO SCs fail to form and

re-form myelination). Therefore, other TEADs can retain nuclear YAP/TAZ, thereby supporting

transcriptional activity of YAP/TAZ, but cannot substitute for some or most of its functions with

TEAD1 in a stage-dependent manner.

It is also notable that, like YAP/TAZ, the unique roles of TEAD1 change dynamically in

successive steps of myelination, even though other TEADs are present. Thus, TEAD1 is

uniquely required for timely proliferation of SCs during radial sorting; uniquely opposes SC

proliferation to promote cell-cycle exit after radial sorting; redundantly promotes upregulation of

Krox20; and then uniquely promotes upregulation of myelin proteins and cholesterol

biosynthetic enzymes. Such unique and redundant roles for TEAD1 in the presence of other

TEADs are supported by cell-based transcription assays, in which YAP-TEAD4 can substitute

for YAP-TEAD1 in upregulating transcription from the Krox20 promoter, but not from the MPZ

intronic enhancer (*37*). Our data also show that even when TEAD1 is largely redundant because

other TEADs are present and can functionally substitute for TEAD1, such as in myelin

maintenance, YAP/TAZ preferentially bind to TEAD1. This situation resembles the dynamic

function of TEADs in cortical development, another multi-step differentiation program. As in

nerve development, multiple TEADs (TEAD1-3) are co-expressed throughout cortical

development. Depending on the differentiation stage, TEAD1 uniquely promotes proliferation,

redundantly promotes proliferation with TEAD2 and 3, or redundantly promotes differentiation

with TEAD3, when it is opposed by TEAD2 (*33*).

As *Yap/Taz* cDKO SCs arrest at the promyelination stage, we were unable to investigate the role

of YAP/TAZ in myelin sheath growth (*13*). In contrast, some *Tead1* cKO SCs can myelinate but

fail to grow myelin sheaths proportional in size to axon diameter. Importantly, such myelinating

SCs apparently lack TEAD1 because TEAD1 deletion is extremely efficient in *Tead1* cKO mice.

Support for a role of TEADs in the regulation of myelin sheath thickness comes from a study of

TEAD4 cKO mice, in which SCs lacking TEAD4 gradually upregulate myelin protein

expression, resulting in adult mice with abnormally thick myelin (*34*). Therefore, TEADs 1 and 4

oppose each other in the regulation of myelin sheath growth. Likewise, they may oppose each

other in the regulation of SC proliferation, resulting in a large excess of SCs we observed in P22

and P50 *Tead1* cKO.

TEAD1 regulation of Krox20 and myelin gene expression

Previous work suggests that YAP/TAZ-TEAD1 promote myelination in part through direct

upregulation of Krox20 (*13, 15*). Unlike *Yap/Taz* cKO mice, *Tead1* cKO mice upregulate

Egr2/Krox20, but they fail to upregulate myelin proteins and cholesterol biosynthetic enzymes.

Our findings extend previous work to suggest that TEAD1 functions redundantly in direct

upregulation of Krox20, but that its function is required for upregulated Krox20 to promote

myelination. Krox20 activity may be repressed in *Tead1* cKO by Oct6, which remains

abnormally elevated. In support of this possibility, persistently elevated Oct6 overrides Krox20

to inhibit myelin gene expression (*46, 58*). Other mice with defective lipid biosynthesis and a

phenotype like *Tead1* cKO also have abnormally elevated Oct6 after Krox20 upregulation (*49,*

59). An alternative or additional possibility is that a transcriptional co-activator that synergizes

with Krox20 is absent in *Tead1* cKO SCs. We suggest that TEAD1 itself may function as such a

factor. Supporting this idea are reports that gene regulatory elements function by co-binding

several TFs, which synergistically activate transcription (*60, 61*), as at the Krox20 MSE

(Myelinating Schwann cell element) which is the enhancer responsible for Krox20 expression (*9,*

11, 13, 62). Krox20 synergizes with Sox10 to activate myelin genes (*63*) and with SREBP to

activate lipid biosynthesis genes (*52*). Like Krox20, TEAD1 can directly promote transcription

from PMP22 and MPZ myelin gene enhancers (*18, 37*). A consistent finding of the present work

is that functional Krox20 is insufficient to drive myelin/cholesterol gene expression in the

absence of TEAD1. Alternatively, or additionally, TEAD1 may also function indirectly by

upregulating co-factors, such as SREBP2, which is reduced in *Tead1* cKO. Notably, lipid

biosynthesis is necessary to downregulate Oct6 and upregulate myelin gene transcription during

developmental myelination (*50, 51*). An interesting possibility, therefore, is that YAP/TAZ-

TEAD1 and Krox20 may cooperatively downregulate Oct6 and upregulate myelin gene

expression at least in part by promoting lipid biosynthesis.

YAP/TAZ-TEAD1 regulation of cholesterol biosynthesis

SREBP1 levels are not altered in *Tead1* cKO or *Yap/Taz* cDKO, suggesting that fatty acid

synthesis is not directly regulated by YAP/TAZ or TEAD1. We did not observe downregulation

of HMGCR in *Tead1* cKO or *Yap/Taz* cDKO mice at P8 or P40, which contradicts the report of

reduced mRNA expression of HMGCR in *Yap/Taz* mutants at P3 and P5 (*14, 15*); we suggest

this difference is due to developmental delay in *Tead1* cKO and *Yap/Taz* cDKO. However,

marked downregulation of IDI1 and FDPS in *Tead1* cKO, *Yap/Taz* cDKO, and *Yap/Taz* iDKO

suggests that cholesterol biosynthesis is defective in these mutants. In support of this notion, SC-

selective deletion of SCAP (SREBP cleavage activation protein), which is required to produce

active SREBP1 and SREBP2 (*51*), or of Squalene synthase, an essential enzyme in cholesterol

biosynthesis (*50*), evokes phenotypes which are remarkably similar to those of *Tead1* cKO mice:

both mutants have an increased number of SCs, a severe delay at the promyelination stage, and

persistently thin myelin. In addition, in SCAP cKO, Oct6 is upregulated continuously and

Squalene synthase cKO fails to upregulate myelin gene transcription, resulting in severely

reduced MBP, MPZ, and PMP22 protein levels.

How might YAP/TAZ-TEAD1 regulate IDI1 and FDPS? Multiple studies show that YAP/TAZ

can activate the mTORC1 pathway (*64, 65*), and IDI1 and FDPS levels are reduced in Raptor

cKO SNs (*49*). Furthermore, the Raptor cKO phenotype is like that of *Tead1* cKO, exhibiting an

increased number of SCs, normal upregulation of Krox20, persistent upregulation of Oct6, delay

at the promyelination stage, and thin myelin. However, HMGCR protein levels are reduced, and

aberrant upregulation of Oct6 is transient (*49*). Furthermore, after nerve injury, formation of

repair SCs is delayed in Raptor mutants (*66*), unlike TEAD1 and YAP/TAZ mutants. These

differences suggest that YAP/TAZ-TEAD1 are not directly upstream of mTORC1. Another

possibility is that YAP/TAZ-TEAD1 may directly regulate transcription of IDI1 and FDPS

genes. Such direct regulation of cholesterol biosynthetic enzymes has been observed for other

TFs in SCs (*52, 67*).

TEAD1 regulation of Remak bundles

In both *Tead1* cKO and iKO mice, there is defective ensheathment of C-fibers by SCs in Remak

bundles, which are occasionally thinly myelinated. This phenotype is distinct from those mutants

in which SCs surround and sometimes myelinate bundles filled with mixed-caliber axons, which

are considered radial sorting defects (*44*). Remak bundles can myelinate abnormally in WT SNs

during early postnatal development, but this defect is corrected and does not persist in adults

(*68*). Therefore, the defective Remak bundles that we observed in *Tead1* cKO are not due to

developmental delay, and the defects found in adult *Tead1* iKO appear unusual.

Remak bundles form normally at P60 in *Yap/Taz* cDKO, although many large bundles containing

mixed caliber axons are present during development (*13*). The difference between *Tead1* and

Yap/Taz mutants is supported by the presence of TEAD1 and absence of YAP/TAZ in non-

myelinating SC nuclei (*13*). Notably, defective SC ensheathment of C-fibers and myelinated

Remak bundles is observed in various SC developmental mutants, including those with defective

SC basal lamina (*68*) and overexpression of NRG1-IIIb in axons (*69*). Therefore, it is possible

that TEAD1 mediates extracellular signaling from the SC basal lamina and/or axons that is

responsible for forming and maintaining proper association of SCs with C-fibers. Notably, SC

ensheathment of C-fibers is required for normal NCV (*70*), and NCV with tg-NRG1-IIIb Remak

bundles is reduced compared to WT (*69*). Future studies will address NRG1 signaling and NCV

in cKO and iKO mice lacking TEAD1 and/or other TEADs.

MATERIALS AND METHODS

Animals

All surgical procedures and animal maintenance complied with the National Institute of Health guidelines regarding the care and use of experimental animals and were approved by the Institutional Animal Care and Use Committee of Temple University, Philadelphia, PA, USA. Generation and genotyping of mice carrying targeted loxP sites in the *Tead1* gene have been described previously (*39*). Targeted ablation of *Tead1* in Schwann cells of embryonic mice was achieved by crossing *Tead1*^{fl/fl} mice with mice transgenic for *Cre* driven by the *Mpz* promoter (*Tead1* cKO mice). Targeted ablation of *Tead1* in Schwann cells of adult mice was by crossing *Tead1*^{fl/fl} mice with mice transgenic for the inducible Cre recombinase *Cre-ERT2* under control of the *Plp1* promoter (*Tead1* iKO mice). Control mice contained floxed *Tead1* alleles but were not transgenic for *Cre* or *Cre-ERT2*. *Mpz-cre*;*Yap*^{fl/fl}; *Taz*^{fl/fl} (*Yap/Taz* cDKO) and *Plp1-creERT2*; *Yap*^{fl/fl}; *Taz*^{fl/fl} (*Yap/Taz* iDKO) mice used in this study were generated and genotyped as described previously (*13*). Both male and female mice were used in all experiments and were maintained on the C57BL/6 background.

Tamoxifen administration

Tamoxifen was prepared and injected into 6–8-week-old *Tead1* or *Yap/Taz* mutant mice as previously described (*13*). A 10 mg/ml solution of tamoxifen was made in 10:1 sunflower oil: 100% ethanol. This solution was injected intraperoneally at a concentration of 0.2 mg/g body weight. Injection was once daily for 5 days, followed by a 2-day break, then once daily for 5 consecutive days.

Nerve crush

Sciatic nerves of right hindlimbs were crushed 24 h after the final tamoxifen injection, as described previously (*17, 71*). A small skin incision was made in the posterior thigh and calf of the animals anesthetized by isoflurane. The sciatic nerve was crushed with a fine forceps (#5) for 10 seconds (3X) adjacent to the sciatic notch. The crush site was marked using charcoal-coated forceps, and the wound was closed. To identify proliferating Schwann cells, we intraperitoneally injected EdU (80 µg/g) eighty minutes before euthanasia, as previously described (*13*).

Western blotting and immunoprecipitation

After removal of epineurium and perineurium, sciatic nerves were snap-frozen in liquid nitrogen and stored at -80°C. For Western blotting, sciatic nerves were lysed in RIPA buffer (25 mM Tris pH 7.5, 150 mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, 1 mM EDTA, 0.1% SDS); for immunoprecipitation, lysis was in co-IP buffer (50 mM Tris pH 8.0, 150 mM NaCl, 0.5% Triton X-100, 10% glycerol, 1 mM EDTA), both containing protease and phosphatase inhibitors. Lysis was for 40 minutes on ice, followed by microcentrifugation at 14,000 rpm for 20 minutes. Protein concentration was determined using the BCA assay (ThermoFisher).

Immunoprecipitation was overnight at 4°C, followed by 1 hour with protein A sepharose (pre-blocked with 1% BSA), both with constant rotation. Lysates and immunoprecipitated complexes in 1x Laemmli buffer were resolved on 8 or 10% SDS-PAGE gels, transferred to nitrocellulose, and blotted with the appropriate antibodies. Primary antibodies were used at the following concentrations for Western blots or immunoprecipitations: rabbit anti-Krox20 (Covance; 1:400), rabbit anti-Oct6 (Abcam ab126746; 1:1000), mouse anti-Myelin Basic Protein (Covance SMI94; 1:1000), mouse anti-beta actin (SIGMA-Aldrich A5441; 1:5000), rabbit anti-TEAD1 (Cell Signaling D9X2L, #12292; 1:1000), rabbit anti-TEAD2 (Novus Biologicals, #NBP3-05267; 1:1000), rabbit anti-TEAD3 (Cell Signaling #13224; 1:1000), rabbit anti-TEAD4 (Santa Cruz, #sc-101184; 1:1000), rabbit anti-pan-TEAD (Cell Signaling, #13295; 1:1000), rabbit anti-YAP/TAZ (Cell Signaling D24E4, #8418; 1:1000, 1:100 for immunoprecipitations), rabbit anti-phospho-YAP(Ser127) (Cell Signaling D9W2I #13008; 1:1000), rabbit anti-phospho-YAP(Ser397) (Cell Signaling D1E7Y #13619; 1:1000), rabbit-anti-MPZ (Cell Signaling #57518; 1:1000), mouse anti-SREBP1 (Novus Biologicals, #NB600-582; 1:1000), rabbit anti-SCD1 (Cell Signaling, #2794; 1:1000), rabbit anti-SREBP2 (Bethyl laboratories, #A303-125A), rabbit anti-HMGCR (Novus Biologicals, #NBP2-66888; 1:1000), rabbit anti-FDPS (Proteintech, #16129-1-AP; 1:1000), rabbit anti-IDI1 (Proteintech, #11166-2-AP; 1:1000). Control whole rabbit IgG for immunoprecipitations was from Jackson Laboratories (011-000-003) and was used at 1μg/ml, which was several-fold greater than all other antibodies used for immunoprecipitations. Secondary antibodies for Western blotting were HRP-linked sheep anti-mouse IgG and donkey anti-rabbit (both GE Healthcare; 1:5000) and HRP-linked mouse anti-rabbit, light-chain specific (Jackson ImmunoResearch 211-032-171; 1:20,000). Blots were developed using ECL, ECL Plus (GE Healthcare) or Odyssey (LiCor).

Immunohistochemistry, confocal and widefield fluorescence microscopy

For immunostaining, sciatic nerves were removed and immediately fixed in 4%
paraformaldehyde in PBS for 1 hr. on ice. Nerves were washed 3 times in PBS, then stored in
15% sucrose in PBS overnight at 4°C for cryoprotection. Nerves were frozen-embedded in
cryoprotectant medium (Thermo Fisher Scientific, Waltham, MA) in isomethylbutane at -80°C.
7-10 μm sections from the nerves were cut using a cryostat (Leica Microsystems, Germany) and
collected directly onto glass slides. For immunolabeling, nerve sections were rehydrated in PBS,
permeabilized in 0.5% Triton/PBS for 20 min, washed with PBS, then blocked in 2% bovine
serum albumin (BSA) in PBS for 1 hr. Sections were incubated with primary antibodies in
blocking buffer overnight at 4°C in a hydrated chamber, washed with PBS, and incubated with
secondary antibodies in blocking buffer for 2 hrs. at room temperature. Sections were washed
with PBS, stained with DAPI for 10 min, and mounted with Vectashield mounting medium
(Vector Labs, Burlingame, CA).

Primary antibodies were used at the following concentrations for immunostaining of
cryosections: guinea pig anti-Sox10 (kind gift from Michael Wegner, University of Erlangen,
Bavaria, Germany; 1:1000), rabbit anti-Krox20 (Covance; 1:400), rabbit anti-Oct6 (Abcam
ab31766; 1:800), rabbit anti-Ki67 (Novocastra NCL-Ki67p; 1:200), rabbit anti-Neurofilament
200 (SIGMA-Aldrich N4142; 1:500), mouse anti-Neurofilament (Covance SMI 312; 1:1000),
mouse anti-Tubulin β3 (clone Tuj1, Covance #MMS-435P; 1:1000), mouse anti-SV2
(Developmental Studies Hybridoma Bank, Iowa; 1:10), rabbit anti-S100 (DAKO; 1:600), mouse
anti-Myelin Basic Protein (Covance SMI 94; 1:2000), rabbit anti-TEAD1 (Cell Signaling
D9X2L, #12292; 1:200), rabbit anti-pan-TEAD (Cell Signaling D3F7L, #13295; 1:200), rabbit
anti-YAP/TAZ (Cell Signaling D24E4, #8418; 1:200).

For immunostaining of extensor digitorum longus (EDL) and soleus muscles, mice were
perfused with 4% paraformaldehyde, and the muscles removed and post-fixed for 20 mins at
room temperature. Muscles were then incubated for 15 minutes with rhodamine-conjugated α-
bungarotoxin (Invitrogen, Carlsbad, CA), permeabilized in −20°C methanol for 5 min, and
blocked for 1 hr. in PBS containing 0.2% Triton and 2% BSA. The muscles were subsequently
incubated overnight at 4ºC in a cocktail of primary antibodies diluted in the blocking solution.
Axons and nerve terminals were labeled with mouse monoclonal antibodies to neurofilaments

(SMI 312; Sternberger Monoclonals, Baltimore, MD) and to a synaptic vesicle protein, SV2

(Developmental Studies Hybridoma Bank, Iowa City, IA). Schwann cells were labeled with

rabbit anti-cow S-100 polyclonal antibody (Dako, Carpentaria, CA). After incubation with the

primary antibodies, muscles were rinsed in PBS and incubated with secondary antibodies in the

blocking solution, for 1 hr. at room temperature. The muscles were then rinsed in PBS and

prepped for whole-mount by peeling the top layer of muscle fibers using microscissors and

forceps, as previously described (*71*). The resulting "muscle fillets" were mounted on a slide and

coverslipped.

Conventional fluorescence microscopy was performed using Olympus BX53 and Zeiss

Axio Imager 2 microscopes. Images were captured using Hamamatsu ORCA R2 C10600 or

Zeiss Axiocam HRm Rev 3 digital cameras, respectively, and Metamorph software. For confocal

microscopy, we used a Leica SP8 confocal microscope, with either a 40x 1.3 NA or 63x 1.4 NA

objective and Leica proprietary software. Acquired stacks were assembled using the maximum

projection tool, and figures were prepared using Adobe Photoshop.

Electron microscopy, histology, and morphometry

Mice were perfused intravascularly with 2.5% glutaraldehyde and 4% paraformaldehyde in 0.1M

cacodylate buffer pH 7.4. Sciatic nerves were removed and incubated in the same buffer for 90

minutes at room temperature, then overnight at 4°C. Specimens were post-fixed for 1 hr. with

OsO_4, dehydrated, then embedded in araldite. Semi-thin (0.5 μm) and ultrathin (70 nm) sections

were cut using an ultramicrotome and stained with toluidine blue or uranyl acetate and lead

citrate, respectively. Semi-thin sections were examined using bright field on an Olympus BX53

microscope, and images were captured using Hamamatsu ORCA R2 C10600 digital camera and

Metamorph software. For analysis of the number of myelin profiles in control vs mutant sciatic

nerves, semi-thin sections from proximal-, mid- and distal regions of the sciatic nerve were

taken, and 2 non- overlapping images of each section were obtained using the 60x objective. For

electron microscopy, stained ultrathin sections were examined using a JEOL 1010 electron

microscope fitted with a Hamamatsu digital camera and AMT Advantage image capture

software.

Proliferation assays

The percentage of Schwann cells in S-phase was measured by EdU incorporation. Mice were

injected intraperitoneally and subcutaneously with 80 μg/ g body weight of EdU dissolved in

PBS. Eighty minutes later, mice were sacrificed, and sciatic nerves removed and fixed for 1 hr.

in 4% paraformaldehyde; longitudinal 7μm cryosections were used for analysis. EdU

incorporation was identified using the Click-iT EdU Alexa Fluor 555 labeling kit (Molecular

Probes); the recommended protocol was followed, and labeling was for 1 hr. After EdU labeling,

sections were immunostained with anti-Sox10 to identify Schwann cells. The percentage of

Schwann cells (Sox10+ nuclei) that were EdU+ was determined by counting confocal images

using ImageJ software.

Electrophysiology

The procedure for measuring nerve conduction velocity was an adaption of a previously used

protocol (*72*). Mice were deeply anesthetized, and hair was completely removed from the

hindquarters. A 27g percutaneous needle electrode was inserted subcutaneously over the femur

to stimulate the sciatic nerve at two locations: one proximal to the spinal cord, at the hip joint,

and the other 1cm distal to this site. The stimulation reference electrode (26G percutaneous

needle) was inserted over the back. Bipolar EMG electrodes were inserted subcutaneously in the

digital interosseous muscles, to record compound action potentials in response to activation of

the sciatic nerve. Stimulation and recording were conducted on both hindlimbs. The muscle

response was first tested with single pulses (100 μs duration) to determine the threshold and level

of activation producing a stable compound action potential in the muscle. The level of

stimulation which produced stable responses was then used for two series of stimulations with 2-

20s long trains of biphasic square wave pulses (100 μs duration) delivered at 1-5 Hz. The nerve

was rested for 2 minutes between runs. Stimulation was delivered via an isolated pulse stimulator

(A-M Systems Model 2100, A-M Systems Inc, Carlsborg, WA). EMG compound muscle action

potential (CMAP) responses were recorded with a differential AC amplifier (A-M Systems

Model 1700; gain 1k-10k; pass band 10-50 KHz). Data was sampled at 50 KHz using LabVIEW

(National Instruments Corporation, Austin, TX) running on a personal computer. Multiple

recordings were taken at each location to ensure a minimum of 10 representative responses for

each location and animal. Electrophysiological data were analyzed using Igor Pro (Wavemetrics

Inc, Lake Oswego, OR).

Data Analysis

In each experiment, data collection and analysis were performed identically, regardless of mouse genotype. Data are presented as mean +/- SD. Statistical analysis was done using the student's unpaired t-test for two-group comparisons and analysis of variance (ANOVA) with Tukey's test for multiple comparisons, according to the number of samples and the analysis of mice at multiple ages. Sample sizes were similar to those conventional in the field and are indicated in the main text, methods, or figure legends. A p-value of 0.05 or less was considered statistically significant.

SUPPLMENTARY MATERIALS

Supplementary figure 1. Additional semi-thin and TEM images of WT and *Tead1* cKO

(A) Representative images of semi-thin sections of various areas of WT and Tead1 cKO SNs at P22 and P50. Scale bar: 20μm

(B) Representative TEM images of transverse sections of WT and Tead1 cKO SNs at P22 and P50. Scale bar: 1μm (P22 panels), 2μm (P50 panels)

Supplementary video 1. A movie showing a P55 WT and two littermates *Tead1* cKO mice

REFERENCES AND NOTES

1. K. R. Jessen, R. Mirsky, The origin and development of glial cells in peripheral nerves. *Nat Rev Neurosci* **6**, 671-682 (2005).
2. J. L. Salzer, Schwann cell myelination. *Cold Spring Harb Perspect Biol* **7**, a020529 (2015).
3. B. Beirowski *et al.*, Metabolic regulator LKB1 is crucial for Schwann cell-mediated axon maintenance. *Nat Neurosci* **17**, 1351-1361 (2014).
4. B. L. Harty, K. R. Monk, Unwrapping the unappreciated: recent progress in Remak Schwann cell biology. *Current opinion in neurobiology* **47**, 131-137 (2017).
5. S. S. Scherer, L. Wrabetz, Molecular mechanisms of inherited demyelinating neuropathies. *Glia* **56**, 1578-1589 (2008).

6.　　M. Bremer *et al.*, Sox10 is required for Schwann-cell homeostasis and myelin maintenance
in the adult peripheral nerve. *Glia* **59**, 1022-1032 (2011).

7.　　L. Decker *et al.*, Peripheral myelin maintenance is a dynamic process requiring constant
Krox20 expression. *The Journal of neuroscience : the official journal of the Society for*
Neuroscience **26**, 9771-9779 (2006).

8.　　M. Finzsch *et al.*, Sox10 is required for Schwann cell identity and progression beyond the
immature Schwann cell stage. *J Cell Biol* **189**, 701-712 (2010).

9.　　Y. He *et al.*, Yy1 as a molecular link between neuregulin and transcriptional modulation
of peripheral myelination. *Nat Neurosci* **13**, 1472-1480 (2010).

10.　　M. Jaegle *et al.*, The POU proteins Brn-2 and Oct-6 share important functions in Schwann
cell development. *Genes Dev* **17**, 1380-1391 (2003).

11.　　S. C. Kao *et al.*, Calcineurin/NFAT signaling is required for neuregulin-regulated Schwann
cell differentiation. *Science* **323**, 651-654 (2009).

12.　　N. Le *et al.*, Nab proteins are essential for peripheral nervous system myelination. *Nat*
Neurosci **8**, 932-940 (2005).

13.　　M. Grove *et al.*, YAP/TAZ initiate and maintain Schwann cell myelination. *Elife* **6**, (2017).

14.　　Y. Poitelon *et al.*, YAP and TAZ control peripheral myelination and the expression of
laminin receptors in Schwann cells. *Nat Neurosci* **19**, 879-887 (2016).

15.　　Y. Deng *et al.*, A reciprocal regulatory loop between TAZ/YAP and G-protein Galphas
regulates Schwann cell proliferation and myelination. *Nat Commun* **8**, 15161 (2017).

16.　　H. Jeanette *et al.*, YAP and TAZ regulate Schwann cell proliferation and differentiation
during peripheral nerve regeneration. *Glia* **69**, 1061-1074 (2021).

17.　　M. Grove, H. Lee, H. Zhao, Y. J. Son, Axon-dependent expression of YAP/TAZ mediates
Schwann cell remyelination but not proliferation after nerve injury. *Elife* **9**, (2020).

18.　　C. Lopez-Anido *et al.*, Tead1 regulates the expression of Peripheral Myelin Protein 22
during Schwann cell development. *Hum Mol Genet*, (2016).

19.　　P. Topilko *et al.*, Krox-20 controls myelination in the peripheral nervous system. *Nature*
371, 796-799 (1994).

20.　　N. Le *et al.*, Analysis of congenital hypomyelinating Egr2Lo/Lo nerves identifies Sox2 as
an inhibitor of Schwann cell differentiation and myelination. *Proc Natl Acad Sci U S A* **102**,
2596-2601 (2005).

735 21. S. W. Jang *et al.*, Locus-wide identification of Egr2/Krox20 regulatory targets in myelin
736   genes. *J Neurochem* **115**, 1409-1420 (2010).

737 22. S. W. Jang, J. Svaren, Induction of myelin protein zero by early growth response 2 through
738   upstream and intragenic elements. *J Biol Chem* **284**, 20111-20120 (2009).

739 23. S. E. LeBlanc, S. W. Jang, R. M. Ward, L. Wrabetz, J. Svaren, Direct regulation of myelin
740   protein zero expression by the Egr2 transactivator. *J Biol Chem* **281**, 5453-5460 (2006).

741 24. Y. Zheng, D. Pan, The Hippo Signaling Pathway in Development and Disease.
742   *Developmental cell* **50**, 264-282 (2019).

743 25. J. H. Xiao, I. Davidson, H. Matthes, J. M. Garnier, P. Chambon, Cloning, expression, and
744   transcriptional properties of the human enhancer factor TEF-1. *Cell* **65**, 551-568 (1991).

745 26. L. Currey, S. Thor, M. Piper, TEAD family transcription factors in development and
746   disease. *Development* **148**, (2021).

747 27. M. Diepenbruck *et al.*, Tead2 expression levels control the subcellular distribution of Yap
748   and Taz, zyxin expression and epithelial-mesenchymal transition. *J Cell Sci* **127**, 1523-
749   1536 (2014).

750 28. J. Kim *et al.*, MAML1/2 promote YAP/TAZ nuclear localization and tumorigenesis. *Proc*
751   *Natl Acad Sci U S A* **117**, 13529-13540 (2020).

752 29. M. Kofler *et al.*, Mediated nuclear import and export of TAZ and the underlying molecular
753   requirements. *Nature communications* **9**, 4966 (2018).

754 30. F. Yao *et al.*, SKP2- and OTUD1-regulated non-proteolytic ubiquitination of YAP
755   promotes YAP nuclear localization and activity. *Nature communications* **9**, 2269 (2018).

756 31. D. Gerber *et al.*, Transcriptional profiling of mouse peripheral nerves to the single-cell
757   level to build a sciatic nerve ATlas (SNAT). *Elife* **10**, (2021).

758 32. J. K. Holden, C. N. Cunningham, Targeting the Hippo Pathway and Cancer through the
759   TEAD Family of Transcription Factors. *Cancers (Basel)* **10**, (2018).

760 33. T. Mukhtar *et al.*, Tead transcription factors differentially regulate cortical development.
761   *Scientific reports* **10**, 4625 (2020).

762 34. X. He *et al.*, A histone deacetylase 3-dependent pathway delimits peripheral myelin growth
763   and functional regeneration. *Nature medicine* **24**, 338-351 (2018).

35.   T. Wen *et al.*, Transcription factor TEAD1 is essential for vascular development by promoting vascular smooth muscle differentiation. *Cell death and differentiation* **26**, 2790-2806 (2019).

36.   R. Liu *et al.*, Tead1 is required for perinatal cardiomyocyte proliferation. *PloS one* **14**, e0212017 (2019).

37.   R. N. Fernando *et al.*, Optimal myelin elongation relies on YAP activation by axonal growth and inhibition by Crb3/Hippo pathway. *Nat Commun* **7**, 12186 (2016).

38.   C. Lopez-Anido *et al.*, Differential Sox10 genomic occupancy in myelinating glia. *Glia*, (2015).

39.   T. Wen *et al.*, Characterization of mice carrying a conditional TEAD1 allele. *Genesis (New York, N.Y. : 2000)* **55**, (2017).

40.   M. L. Feltri *et al.*, P0-Cre transgenic mice for inactivation of adhesion molecules in Schwann cells. *Ann N Y Acad Sci* **883**, 116-123 (1999).

41.   D. P. Leone *et al.*, Tamoxifen-inducible glia-specific Cre mice for somatic mutagenesis in oligodendrocytes and Schwann cells. *Mol Cell Neurosci* **22**, 430-440 (2003).

42.   A. Beverdam *et al.*, Yap controls stem/progenitor cell proliferation in the mouse postnatal epidermis. *J Invest Dermatol* **133**, 1497-1505 (2013).

43.   Z. Meng, T. Moroishi, K. L. Guan, Mechanisms of Hippo pathway regulation. *Genes Dev* **30**, 1-17 (2016).

44.   M. L. Feltri, Y. Poitelon, S. C. Previtali, How Schwann Cells Sort Axons: New Concepts. *Neuroscientist* **22**, 252-265 (2016).

45.   C. Hindley, A. Philpott, Co-ordination of cell cycle and differentiation in the developing nervous system. *Biochem J* **444**, 375-382 (2012).

46.   E. J. Ryu *et al.*, Misexpression of Pou3f1 results in peripheral nerve hypomyelination and axonal loss. *J Neurosci* **27**, 11552-11559 (2007).

47.   T. S. Zorick, D. E. Syroid, A. Brown, T. Gridley, G. Lemke, Krox-20 controls SCIP expression, cell cycle exit and susceptibility to apoptosis in developing myelinating Schwann cells. *Development* **126**, 1397-1406 (1999).

48.   L. Montani *et al.*, De novo fatty acid synthesis by Schwann cells is essential for peripheral nervous system myelination. *J Cell Biol* **217**, 1353-1368 (2018).

49. C. Norrmen *et al.*, mTORC1 controls PNS myelination along the mTORC1-RXRgamma-
SREBP-lipid biosynthesis axis in Schwann cells. *Cell Rep* **9**, 646-660 (2014).

50. G. Saher *et al.*, Cholesterol regulates the endoplasmic reticulum exit of the major
membrane protein P0 required for peripheral myelin compaction. *J Neurosci* **29**, 6094-
6104 (2009).

51. M. H. Verheijen *et al.*, SCAP is required for timely and proper myelin membrane synthesis.
Proc Natl Acad Sci U S A **106**, 21383-21388 (2009).

52. S. E. Leblanc *et al.*, Regulation of cholesterol/lipid biosynthetic genes by Egr2/Krox20
during peripheral nerve myelination. *J Neurochem* **93**, 737-748 (2005).

53. M. C. Wright, W. J. Cho, Y. J. Son, Distinct patterns of motor nerve terminal sprouting
induced by ciliary neurotrophic factor vs. botulinum toxin. *J Comp Neurol* **504**, 1-16
(2007).

54. A. Clemence, R. Mirsky, K. R. Jessen, Non-myelin-forming Schwann cells proliferate
rapidly during Wallerian degeneration in the rat sciatic nerve. *J Neurocytol* **18**, 185-192
(1989).

55. K. R. Jessen, R. Mirsky, The repair Schwann cell and its function in regenerating nerves.
J Physiol **594**, 3521-3531 (2016).

56. N. Tricaud, H. T. Park, Wallerian demyelination: chronicle of a cellular cataclysm. *Cell
Mol Life Sci* **74**, 4049-4057 (2017).

57. X. Varelas, The Hippo pathway effectors TAZ and YAP in development, homeostasis and
disease. *Development* **141**, 1614-1626 (2014).

58. E. S. Monuki, R. Kuhn, G. Lemke, Repression of the myelin P0 gene by the POU
transcription factor SCIP. *Mech Dev* **42**, 15-32 (1993).

59. D. L. Sherman *et al.*, Arrest of myelination and reduced axon growth when Schwann cells
lack mTOR. *The Journal of neuroscience : the official journal of the Society for
Neuroscience* **32**, 1817-1825 (2012).

60. S. Heinz, C. E. Romanoski, C. Benner, C. K. Glass, The selection and function of cell type-
specific enhancers. *Nat Rev Mol Cell Biol* **16**, 144-154 (2015).

61. B. Ren, F. Yue, Transcriptional Enhancers: Bridging the Genome and Phenome. *Cold
Spring Harb Symp Quant Biol* **80**, 17-26 (2015).

62. J. Ghislain, P. Charnay, Control of myelination in Schwann cells: a Krox20 cis-regulatory
element integrates Oct6, Brn2 and Sox10 activities. *EMBO Rep* **7**, 52-58 (2006).

63. E. A. Jones *et al.*, Interactions of Sox10 and Egr2 in myelin gene regulation. *Neuron Glia*
Biol **3**, 377-387 (2007).

64. J. K. Hu *et al.*, An FAK-YAP-mTOR Signaling Axis Regulates Stem Cell-Based Tissue
Renewal in Mice. *Cell Stem Cell* **21**, 91-106 e106 (2017).

65. M. Kim, T. Kim, R. L. Johnson, D. S. Lim, Transcriptional co-repressor function of the
hippo pathway transducers YAP and TAZ. *Cell Rep* **11**, 270-282 (2015).

66. C. Norrmén *et al.*, mTORC1 Is Transiently Reactivated in Injured Nerves to Promote c-
Jun Elevation and Schwann Cell Dedifferentiation. *The Journal of neuroscience : the*
official journal of the Society for Neuroscience **38**, 4811-4828 (2018).

67. M. Kim *et al.*, Maf links Neuregulin1 signaling to cholesterol synthesis in myelinating
Schwann cells. *Genes Dev* **32**, 645-657 (2018).

68. K. Rasi *et al.*, Lack of collagen XV impairs peripheral nerve maturation and, when
combined with laminin-411 deficiency, leads to basement membrane abnormalities and
sensorimotor dysfunction. *J Neurosci* **30**, 14490-14501 (2010).

69. J. A. Gomez-Sanchez *et al.*, Sustained axon-glial signaling induces Schwann cell
hyperproliferation, Remak bundle myelination, and tumorigenesis. *The Journal of*
neuroscience : the official journal of the Society for Neuroscience **29**, 11304-11315 (2009).

70. C. Taveggia *et al.*, Neuregulin-1 type III determines the ensheathment fate of axons.
Neuron **47**, 681-694 (2005).

71. Y. J. Son, W. J. Thompson, Schwann cell processes guide regeneration of peripheral axons.
Neuron **14**, 125-132 (1995).

72. A. Schulz, C. Walther, H. Morrison, R. Bauer, In vivo electrophysiological measurements
on mouse sciatic nerves. *J Vis Exp*, (2014).

ACKNOWLEDGMENTS

We thank members of the Son laboratory for critical reading of the manuscript. We thank Dr.

Eric Olson for *Yap* and *Taz* floxed mice, Drs. Ueli Suter and Kelly Monk for Plp-creERT2 mice.

Plp-creERT2 mice were generated by Dr. Ueli Suter using a patented Cre-ERT2 construct

developed by Dr. Pierre Chambon at GIE-CERBM.

Figure 1. Expression of TEADs in Schwann cells (SCs) of WT, *Tead1* cKO, and iKO mice

(A) Immunostaining and quantification of TEAD1+ SCs on longitudinal sections of P22 WT and

Tead1 cKO SNs. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. n = 3

mice per genotype, ****P < 0.0001, Student's unpaired t-test. Scale bar: 20μm

(B) Immunostaining and quantification of TEAD1+ SCs on longitudinal sections of WT and

Tead1 iKO SNs 3 months after tamoxifen administration. SC nuclei are identified by Sox10. All

cell nuclei are marked by DAPI. n = 3 mice per genotype, ****$P < 0.0001$, Student's unpaired t-

test. Scale bar: 20μm

(C) Immunostaining and quantification of TEAD1+ SCs on longitudinal sections of WT and

Tead1 iKO SNs 28 days after nerve crush. SC nuclei are identified by Sox10. All cell nuclei are

marked by DAPI. n = 3 mice per genotype, ****$P < 0.0001$, Student's unpaired t-test. Scale bar:

20μm

(D) Western blotting of P50 WT and *Tead1* cKO SN lysates with anti-TEAD1, -TEAD2, -

TEAD3, -TEAD4, and -pan TEAD. TEAD expression is normalized to that of *β-actin* as an

internal control, and WT expression is arbitrarily given the value 1. n = 3 mice per genotype. ns,

$p > 0.05$; *$p \leq 0.05$, **$p \leq 0.005$, ***$p \leq 0.001$, unpaired Student's t-test.

(E) Immunostaining and quantification of pan-TEAD+ SCs on longitudinal sections of P100 WT

and *Tead1* iKO SNs. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. n =

3 mice per genotype. ns, $p > 0.05$, Student's unpaired t-test. Scale bar: 20μm

(F) Immunostaining and quantification of pan-TEAD+ SCs on longitudinal sections of P100 WT

and *Tead1* iKO SNs 28 days after nerve crush. SC nuclei are identified by Sox10. All cell nuclei

are marked by DAPI. n = 3 mice per genotype. ns, $p > 0.05$, Student's unpaired t-test. Scale bar:

20μm

Figure 2. YAP/TAZ strongly favor TEAD1 but partner with other TEADs

(A) Western blotting of P50 WT and *Tead1* cKO SN lysates with anti-YAP, -phospho-YAPs

(Ser127 and Ser397), and -TAZ. YAP/TAZ expression is normalized to that of *β-actin* as an

internal control, and WT expression is arbitrarily given the value 1. n = 3 mice per genotype. *p

≤ 0.05, ***p ≤ 0.001, ****P < 0.0001, Student's unpaired t-test.

(B) Immunostaining of YAP/TAZ+ SCs on longitudinal sections of P50 WT and *Tead1* cKO

SNs. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. Scale bar: 20μm

(C) Immunostaining of YAP/TAZ+ SCs on longitudinal sections of P100 WT and *Tead1* iKO

SNs. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. Scale bar: 20μm

(D) Immunostaining of YAP/TAZ+ SCs on longitudinal sections of P100 WT and *Tead1* iKO

SNs 28 days after nerve crush. SC nuclei are identified by Sox10. All cell nuclei are marked by

DAPI. Scale bar: 20μm

(E) Quantification of YAP/TAZ+ SCs in WT, Tead1 cKO, intact, and injured *Tead1* iKO SNs. n

= 3 mice per group. ns, p > 0.05, Student's unpaired t-test.

(F) Immunoprecipitation of P100 WT and *Tead1* iKO SN lysates with anti-YAP/TAZ and

control Rb IgG. Lysate samples, labeled post-IP lysate, were taken from tubes after

immunoprecipitated complexes attached to beads had been spun down to assess the efficiency.

Immunoprecipitates were blotted for YAP/TAZ, TEAD1, and pan-TEAD. Arrow denotes a faint

TEAD1 band revealed by pan-TEAD in WT lysate.

Figure 3. TEAD1 is required for myelin sheath formation and growth

(A) A living control (WT) and two *Tead1* cKO mice from one litter at P55. Arrows point to splayed paralyzed hindlimbs.

(B) Exposed SNs of WT and *Tead1* cKO mice during dissection. Scale bar: 1mm

(C) Representative images of CMAPs generated by stimulation of WT and *Tead1* cKO SNs.

(D) Quantification of NCV in WT and *Tead1* cKO. n = 6 mice per genotype. ***P < 0.001, Student's unpaired t-test.

(E) Representative images of semithin (left panels) and TEM (right panels) of transverse sections of WT and *Tead1* cKO SNs at P3. Scale bar: 20μm (left panels), 1μm (right panels)

(F) Representative images of semithin (left panels) and TEM (right panels) of transverse sections of WT and *Tead1* cKO SNs at P22. Scale bar: 20μm (left panels), 1μm (right panels)

(G) Representative images of semithin (left panels) and TEM (right panels) of transverse sections of WT and *Tead1* cKO SNs at P50. Scale bar: 20μm (left panels), 1μm (right panels)

(H) Quantification of G-ratio for axons in WT and *Tead1* cKO SNs at P22 and at P50. n = 3 mice per group. ns, p > 0.05 (P22 WT versus P50 WT, P22 cKO versus P50 cKO), ****P < 0.0001 (P22 WT versus P22 cKO, P50 WT versus P50 cKO).

(I) Scatter plot graph displaying G-ratio in relation to axon diameter in WT and *Tead1* cKO SNs at P22. n = 1063 axons from 3 mice per each genotype.

(J) Scatter plot graph displaying G-ratio in relation to axon diameter in WT and *Tead1* cKO SNs at P50. n = 880 axons from 3 mice per each genotype.

(K) Quantitative comparison of mean axon diameter in WT and *Tead1* cKO SNs at P22 and at P50. n = 3 mice per group. ns, p > 0.05, *p ≤ 0.05, 2-way ANOVA with Tukey's multiple comparisons test.

(L) Representative TEM images of transverse sections of P50 WT and *Tead1* cKO SNs. Right panels are enlarged images of the boxed area in left panels. Scale bar: 1μm

(M) Quantification of normal and abnormal C-fibers in P50 WT and *Tead1* cKO SNs. C-fibers normally ensheathed by SC cytoplasm are marked as 'ensheathed', whereas those not ensheathed are denoted as 'naked'. Naked C-fibers located in a myelinated bundle are denoted as 'naked myelinated'. Relative percentage of each type of C-fiber is plotted. N > 500 C-fibers from 3 mice per each genotype. *p ≤ 0.05, ***p ≤ 0.001, ****P < 0.0001, 2-way ANOVA Tukey's multiple comparisons test.

Figure 4. TEAD1 regulates SC proliferation both positively and negatively

(A) Immunostaining and quantification of SCs on transverse sections of WT and *Tead1* iKO SNs at P3, P22, and P52. SC nuclei and all cell nuclei are marked by Sox10 and DAPI. Axons are marked by Neurofilaments (NF). n = 3 mice per genotype, ns, p > 0.05, *p ≤ 0.05, **p ≤ 0.01, ****P < 0.0001, 2-way ANOVA with Tukey's multiple comparisons test. Scale bar: 30μm

(B) Immunostaining and quantification of Ki67+ SCs on longitudinal sections of WT and *Tead1* iKO SNs at P3, P11, and P22. SC nuclei and all cell nuclei are marked by Sox10 and DAPI. n = 3 mice per genotype, ns, p > 0.05, **p ≤ 0.01, ****P < 0.0001, 2-way ANOVA with Tukey's multiple comparisons test. Scale bar: 30μm (P3 panels), 20μm (P11 and P22 panels)

(C) Immunostaining and quantification of EdU+ SCs on longitudinal sections of WT and *Tead1* iKO SNs at P3, P11, and P22. SC nuclei and all cell nuclei are marked by Sox10 and DAPI. n = 3 mice per genotype, ns, p > 0.05, *p ≤ 0.05, ****P < 0.0001, 2-way ANOVA with Tukey's multiple comparisons test. Scale bar: 30μm (P3 panels), 20μm (P11 and P22 panels)

(D) Immunostaining and quantification of EdU+/TEAD1+ SCs on longitudinal sections of WT and *Tead1* iKO SNs at P11. SC nuclei are marked by Sox10. Asterisks denote *Tead1* cKO SCs in which TEAD1 is not deleted and which are not EdU+. Arrowheads denote EdU+ SCs which are TEAD1+ in WT but TEAD1- in *Tead1* cKO. n = 3 mice per genotype, ***P < 0.001, Student's unpaired t-test. Scale bar: 20μm

Figure 5. TEAD1 is required for Krox20 to upregulate myelin proteins during

development

(A) Western blotting of P50 WT and *Tead1* cKO SN lysates with anti-Oct6, -Krox20, -MPZ, and

-MBP. Protein expression is normalized to that of *β-actin* as an internal control, and WT

expression is arbitrarily given the value 1. n = 3 mice per genotype. *p ≤ 0.05, ***p ≤ 0.001,

***p ≤ 0.0001, Student's unpaired t-test.

(B) Immunostaining and quantification of Oct6+ SCs on longitudinal sections of P50 WT and

Tead1 cKO SNs. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. n = 3

mice per genotype, ****P < 0.0001, Student's unpaired t-test. Scale bar: 20µm

(C) Immunostaining and quantification of Krox20+ SCs on longitudinal sections of WT and

Tead1 cKO SNs at P11 and P50. SC nuclei are identified by Sox10. All cell nuclei are marked by

DAPI. n = 3 mice per each group, ns, p > 0.05, Student's unpaired t-test. Scale bar: 20µm

(D) Immunostaining and quantification of EdU+/Krox20+ SCs on longitudinal sections of WT

and *Tead1* cKO SNs at P11. SC nuclei are identified by Sox10. Arrowheads denote examples of

EdU+ SCs which are Krox20-. Asterisks denote rarely observed EdU+ SCs which are Krox20+.

n = 3 mice per genotype, ns, p > 0.05, Student's unpaired t-test. Scale bar: 20µm

Figure 6. TEAD1 and YAP/TAZ upregulate cholesterol synthesis enzymes: FDPS and IDI1

(A) Western blotting of P8 and P40 WT and *Tead1* cKO SN lysates with anti-SREBP1, -SCD1, -SREBP2, -HMGCR, -FDPS, and -IDI1. Protein expression is normalized to that of *β-actin* as an internal control, and WT expression is arbitrarily given the value 1. n = 3 or 4 mice per genotype. ns, p > 0.05, *p ≤ 0.05, **p ≤ 0.01, ****P < 0.0001, Student's unpaired t-test. **(B)** Western blotting of P60 WT and *Yap/Taz* cDKO SN lysates with anti-YAP, -TAZ, -SREBP1, -SCD1, -

SREBP2, -HMGCR, -FDPS, and -IDI1. Protein expression is normalized to that of *β-actin* as an

internal control, and WT expression is arbitrarily given the value 1. n = 3 or 4 mice per genotype.

ns, p > 0.05, *p ≤ 0.05, **p ≤ 0.01, ***P < 0.001, Student's unpaired t-test.

Figure 7. TEAD1 is largely dispensable for myelin maintenance, but required for preservation of Remak bundle integrity

(A) Cartoon showing timeline of tamoxifen injection and time of sacrifice of *Tead1* iKO mice.

(B) Representative TEM images of P150 WT and *Tead1* iKO SNs showing occasional large axons undergoing demyelination (arrowheads) or completely demyelinated (arrows). Scale bar: 6μm (left panels), 2μm (right panel)

(C) Quantification of NCV in P90-P150 WT and *Tead1* iKO mice. n = 5 or 6 mice per genotype. ns, $p > 0.05$, Student's unpaired t-test.

(D) Representative TEM images of P97 WT and *Tead1* iKO SNs showing Remak bundles. Asterisk denotes a bundle of thinly myelinated C-fibers. Arrow denotes demyelinated axon. Scale bar: 1μm

(E) Quantification of normal and abnormal C-fibers in P150 WT and *Tead1* iKO SNs. Relative percentage of each type of C-fiber is plotted. N > 500 C-fibers from 3 mice per each genotype. ns, $p > 0.05$, ****$P < 0.0001$, 2-way ANOVA Tukey's multiple comparisons test.

(F) Immunostaining and quantification of Krox20+ SCs on longitudinal sections of WT and *Yap/Taz* iKO SNs 2 weeks after 1st tamoxifen injection. SC nuclei are marked by Sox10. All cell nuclei are marked by DAPI. n = 3 mice per genotype, ***$P < 0.001$, Student's unpaired t-test. Scale bar: 20μm

(G) Immunostaining and quantification of Krox20+ SCs on longitudinal sections of WT and *Tead1* iKO SNs 6 weeks after 1st tamoxifen injection. SC nuclei are marked by Sox10. All cell nuclei are marked by DAPI. n = 3 mice per genotype, ns, $p > 0.05$, Student's unpaired t-test. Scale bar: 20μm

(H) Immunostaining and quantification of Oct6+ SCs on longitudinal sections of WT and *Tead1* iKO SNs 6 weeks after 1st tamoxifen injection. SC nuclei are marked by Sox10. All cell nuclei are marked by DAPI. n = 3 mice per genotype, ns, $p > 0.05$, Student's unpaired t-test. Scale bar: 20μm

(I) Western blotting of WT and *Yap/Taz* iDKO SN lysates with anti-YAP, -TAZ, -SREBP1, -SCD1, -SREBP2, -HMGCR, -FDPS, and -IDI1. Protein expression is normalized to that of *β-actin* as an internal control, and WT expression is arbitrarily given the value 1. n = 3 or 4 mice per genotype. ns, $p > 0.05$, **$p \leq 0.01$, ***$P < 0.001$, ****$P < 0.0001$, Student's unpaired t-test.

1013   **(J)** Western blotting of WT and *Tead1* iKO SN lysates with anti-SREBP1, -SCD1, -SREBP2, -

1014   HMGCR, -FDPS, and -IDI1. Protein expression is normalized to that of *β-actin* as an internal

1015   control, and WT expression is arbitrarily given the value 1. n = 3 or 4 mice per genotype. ns, p >

1016   0.05, Student's unpaired t-test.

1017

1018

1019   **Figure 8. TEAD1 is required for functional regeneration of peripheral nerve**

(A) Cartoon showing timeline of tamoxifen injection and time of sacrifice of *Tead1* iKO mice after nerve crush.

(B) Low magnification images of longitudinal sections of ~4mm long SN segments of WT and *Tead1* iKO mice 28 days after nerve crush. Arrowheads denote crush site. Axons and myelin are marked by Tuj1 and MBP, respectively. All cell nuclei are marked by DAPI. Yellow asterisks denote the distal nerve segment of *Tead1* iKO, which almost completely lacks myelin. White asterisks denote distal segments of both WT and *Tead1* iKO, which exhibit more cells than proximal segments due to SC proliferation after injury. Scale bar: 500µm

(C) Representative low and high magnification NMJ images of WT and Tead1 iKO 28 days after nerve crush. Axon and axon terminals are marked by neurofilament (NF) and SV2 staining. SCs are marked by S100. Muscle acetylcholine receptors are marked by α-Bungarotoxin. Arrows denote examples of axon terminals that grew past NMJs in iKO. Arrowheads denote examples of extrasynaptic AChR clusters in iKO. Scale bar: 50µm (low magnification panels), 20µm (high magnification panels)

(D) Representative semi-thin and TEM images of transverse sections of WT and *Tead1* iKO 28 days after nerve crush. Scale bar: 20µm (left semi-thin panels), 2µm (right TEM panels)

(E) Representative CMAP recordings and NCV measurements of WT and *Tead1* iKO at 28 days after nerve crush. n = 3 mice per genotype, **$p \leq 0.01$, Student's unpaired t-test.

(F) Immunostaining and quantification of EdU+ SCs on longitudinal sections of WT and *Tead1* iKO SNs at 28 days after nerve crush. SC nuclei are marked by Sox10. All cell nuclei are marked by DAPI. n = 4 mice per genotype, ns, $p > 0.05$, Student's unpaired t-test. Scale bar: 20µm

(G) Immunostaining and quantification of Krox20+ SCs on longitudinal sections of WT and *Tead1* iKO SNs at 28 days after nerve crush. SC nuclei are marked by Sox10. All cell nuclei are marked by DAPI. n = 3 mice per genotype, ns, $p > 0.05$, Student's unpaired t-test. Scale bar: 20µm

(H) Immunostaining and quantification of Oct6+ SCs on longitudinal sections of WT and *Tead1* iKO SNs at 28 days after nerve crush. SC nuclei are marked by Sox10. All cell nuclei are marked by DAPI. n = 3 mice per genotype, ns, ***$P < 0.001$, Student's unpaired t-test. Scale bar: 20µm

FUNDINGS

1050 This work was supported by grants from Shriners Children's (#84050 to Y-J.S.) and NIH NINDS

1051 (NS105796 to Y-J.S.).

1052

AUTHOR CONTRIBUTIONS

Conceptualization: MG, YS

Methodology: GH, JZ

Investigation: MG, HK, SP, JA, ML, YS

Funding acquisition: YS

Supervision: ML, YS

Writing – original draft: MG

Writing – review & editing: YS

1061

COMPETING INTERESTS

1063 The authors declare no competing interests.

1064

DATA AND MATERIALS AVAILABILITY

All data are available in the main text or the supplementary materials.

Supplementary figure 1. Additional semi-thin and TEM images of WT and *Tead1* cKO

(A) Representative images of semi-thin sections of various areas of WT and Tead1 cKO SNs at

P22 and P50. Scale bar: 20µm

(B) Representative TEM images of transverse sections of WT and Tead1 cKO SNs at P22 and

P50. Scale bar: 1µm (P22 panels), 2µm (P50 panels)

Supplementary video 1. A movie showing a P55 WT and two littermates *Tead1* cKO mice