TEAD1 is crucial for developmental myelination, Remak bundles, and functional regeneration of peripheral nerves

Matthew Grove[1], Hyukmin Kim[1], Shuhuan Pang[1], José Paz Amaya[2], Guoqing Hu[3], Jiliang Zhou[3], Michel Lemay[2], and Young-Jin Son[1]*

[1]Department of Neural Sciences, Shriners Hospitals Pediatric Research Center, Lewis Katz School of Medicine, Temple University, Philadelphia, PA 19140

[2]Department of Bioengineering, Temple University, Philadelphia, PA 19122, USA

[3]Department of Pharmacology & Toxicology, Medical College of Georgia, Augusta University, Augusta, GA

* Corresponding author: yson@temple.edu

ABSTRACT

Previously we showed that the hippo pathway transcriptional effectors, YAP and TAZ, are

essential for Schwann cells (SCs) to develop, maintain and regenerate myelin (Grove et al.,

2017; Grove, Lee, Zhao, & Son, 2020). Although TEAD1 has been implicated as a partner

transcription factor, the mechanisms by which it mediates YAP/TAZ regulation of SC

myelination are unclear. Here, using conditional and inducible knockout mice, we show that

TEAD1 is crucial for SCs to develop and regenerate myelin. It promotes myelination by both

positively and negatively regulating SC proliferation, enabling Krox20/Egr2 to upregulate

myelin proteins, and upregulating the cholesterol biosynthetic enzymes FDPS and IDI1. We also

show stage-dependent redundancy of TEAD1 and that non-myelinating SCs have a unique

requirement for TEAD1 to enwrap nociceptive axons in Remak bundles. Our findings establish

TEAD1 as a major partner of YAP/TAZ in developmental myelination and functional nerve

regeneration and as a novel transcription factor regulating Remak bundle integrity.

INTRODUCTION

In the mammalian peripheral nervous system, all motor and sensory axons are associated with Schwann cells (SCs). Axons larger than 1μm diameter are enveloped by myelinating SCs that form multilamellar myelin sheaths with a thickness proportional to axon caliber (Jessen & Mirsky, 2005; Salzer, 2015). Axons smaller than 1μm diameter, the unmyelinated nociceptors (i.e., C-fibers), are organized in clusters called Remak bundles, in which each C-fiber is ensheathed by a cytoplasmic pocket formed by non-myelinating SCs, also known as Remak SCs (Beirowski et al., 2014; Harty & Monk, 2017). Both myelinating and non-myelinating SCs provide metabolic and functional support essential for structural and functional integrity of peripheral nerves. Failure to initiate or maintain proper axon-SC association may lead to life-threatening paralysis and peripheral neuropathy (Scherer & Wrabetz, 2008). Myelinating SCs form and maintain myelin via a set of key transcription factors (TFs) and regulators (Bremer et al., 2011; Decker et al., 2006; Finzsch et al., 2010; Y. He et al., 2010; Jaegle et al., 2003; Kao et al., 2009; Le, Nagarajan, Wang, Svaren, et al., 2005). However, the transcriptional regulation of myelinating SCs remains incompletely understood, and that of non-myelinating SCs is largely unknown (Harty & Monk, 2017).

Recent mouse genetic studies surprisingly revealed that YAP (Yes-associated protein, also known as YAP1) and its paralogue, TAZ (Transcriptional coactivator with PDZ-binding motif, also known as WWTR1), which are potent oncoproteins and transcriptional regulators in the Hippo signaling pathway, play crucial roles in developing, maintaining, and regenerating myelination of peripheral nerves. Both YAP and TAZ (YAP/TAZ) are expressed initially in all immature SCs and then selectively in myelinating SCs (Grove et al., 2017). The processes for which they are required include: timely sorting of large axons and associated SC proliferation (Grove et al., 2017; Poitelon et al., 2016); initiation of SC differentiation or myelination (Deng et al., 2017; Grove et al., 2017); myelin maintenance in adults (Grove et al., 2017); and remyelination following nerve injury (Grove et al., 2020; Jeanette et al., 2021). YAP/TAZ initiate and maintain myelination in part by upregulating Krox20 (Egr2) (Grove et al., 2017; Lopez-Anido et al., 2016), which is widely recognized as the master TF promoting peripheral myelination by SCs. Krox20 is required to develop and maintain myelin (Decker et al., 2006; Topilko et al., 1994), and likely to promote myelin sheath growth (Le, Nagarajan, Wang, Araki,

et al., 2005). It acts through direct transcriptional upregulation of myelin proteins and membrane

lipids (Jang et al., 2010; Jang & Svaren, 2009; LeBlanc, Jang, Ward, Wrabetz, & Svaren, 2006).

YAP/TAZ are transcriptional co-activators that contain a transcriptional activation

domain, but no DNA binding domain. They regulate gene expression by binding to partner TFs,

typically members of the TEA domain family (TEAD1-4) (Wu, Liu, Zheng, Dong, & Pan, 2008;

Zhao et al., 2008). Conversely, TEADs on their own are unable to activate or repress

transcription and require the help of co-factors such as YAP/TAZ to act (Li et al., 2010; Tian,

Yu, Tomchick, Pan, & Luo, 2010). Partner TFs such as TEADs are required to maintain

YAP/TAZ in the nucleus, which keeps them transcriptionally active (Diepenbruck et al., 2014; J.

Kim et al., 2020; Kofler et al., 2018; Yao et al., 2018). Recent single cell transcriptional analysis

indicates that SCs express all four TEADs (TEAD1/2/3/4) and that their expression peaks at

different times during development (Gerber et al., 2021). Moreover, TEAD1-4 show unique,

redundant, and/or opposing roles in regulating proliferation and differentiation of various cell

types; these roles can also change as a function of developmental stage (X. He et al., 2018;

Mukhtar et al., 2020). It remains to be elucidated how and which TEAD(s) in SCs mediate

transcriptional regulation of YAP/TAZ for the development, maintenance, and/or regeneration of

myelination after nerve injury.

In some cell types, TEAD1 serves as the main partner TEAD and plays non-redundant

roles (Liu et al., 2019; Wen et al., 2019). In accordance, in SCs, TEAD1 associates with

YAP/TAZ for upregulating myelin proteins as indicated by ChiP, co-IP, and transcription assays

(Fernando et al., 2016; Grove et al., 2017; Lopez-Anido et al., 2016; Lopez-Anido et al., 2015).

Notably, however, unlike YAP/TAZ, which are barely detectable in non-myelinating SCs,

TEAD1 is clearly expressed in all SC nuclei (Grove et al., 2017), suggesting that, independently

of YAP/TAZ, TEAD1 plays an unknown role in non-myelinating SCs. In this study, we

generated conditional and inducible knockout mice lacking TEAD1 in SCs to investigate the

roles of TEAD1 in myelinating and non-myelinating SCs. We found that TEAD1 is required for

myelinating SCs to form, grow, and regenerate myelin, and for non-myelinating SCs to enwrap

C-fibers in Remak bundles during development and homeostasis. We also found that YAP/TAZ-

TEAD1 promotes myelination by dually regulating SC proliferation, by enabling induced

Krox20 to upregulate myelin proteins, and by upregulating cholesterol biosynthetic enzymes.

RESULTS

Multiple TEADs are expressed in Schwann cells of WT and *Tead1* cKO mice

To investigate the roles of SC TEAD1 in developing nerves, we conditionally ablated *Tead1* by crossing *Tead1*[fl/fl] mice (Wen et al., 2017) with mice carrying *Mpz*-Cre, which is active from E13.5 in SCs (Feltri et al., 1999). Immunostaining of P22 WT and *Mpz-Cre;Tead1*[fl/fl] (hereafter *Tead1* cKO) sciatic nerves (SNs) showed that TEAD1 was expressed in the nuclei of 100% of WT SCs and was undetectable in >97% *Tead1* cKO SCs (Figure 1A). The specificity of the TEAD1 immunostaining and highly efficient deletion were further confirmed by Western blotting (Figure 1B).

To investigate the protein expression of other TEADs, we used antibodies to TEAD2, 3, and 4, and a pan-TEAD antibody marketed as recognizing all four TEADs (TEAD1/2/3/4). Western blot analysis of SN lysates showed that anti-TEAD3 and anti-pan TEAD both contained a main ~ 50kD band that was decreased in *Tead1* cKO (Figure 1B). The pan-TEAD antibody may therefore bind preferentially to TEAD3. Anti-TEAD4 recognized three bands in WT and showed that the two faster-migrating bands were decreased in *Tead1* cKO. Anti-TEAD2 recognized a band of ~65 kD in *Tead1* cKO that was barely detectable in WT lysates. This band corresponded to the size of TEAD2 reported in other cell types (Beverdam et al., 2013). We also found that the ~65 kD band was present early postnatally, but rapidly diminished with age (Figure 1-figure supplement 1), in accordance with the transcript data from mouse SCs (Gerber et al., 2021). Thus, the expression of TEADs 2-4 was altered in *Tead1* cKO, likely reflecting their expression in SCs, consistent with the single cell transcriptomic analysis of SCs (Gerber et al., 2021). Of these anti-TEAD antibodies, only anti-pan TEAD functioned for immunofluorescence staining, verifying the presence of other TEADs in SC nuclei lacking TEAD1 (Figure 6-figure supplement 1C, 1E).

In the absence of a partner TF, nuclear YAP/TAZ are decreased and phosphorylation of YAP/TAZ increased, due to targeting by Large tumor suppressor kinase 1/2 (LATS1/2), negative regulators of YAP/TAZ (Hao, Chun, Cheung, Rashidi, & Yang, 2008; Zhao et al., 2007). To investigate if YAP/TAZ are in the nuclei of SCs and therefore remain transcriptionally functional in the absence of TEAD1, we compared Western blotting of P50 WT vs *Tead1* cKO SN lysates and observed an overall decrease in YAP levels in *Tead1* cKO but a > 2-fold increase in TAZ

levels compared to WT (Figure 1C). This ratio of TAZ to YAP in *Tead1* cKO resembled that

observed during developmental myelination in WT SNs (Deng et al., 2017; Grove et al., 2017).

Furthermore, phosphorylation of YAP decreased in *Tead1* cKO, which would typically indicate

increased YAP activity. Thus, YAP/TAZ are not targeted for destruction in the absence of

TEAD1 in SCs. Indeed, YAP/TAZ were retained in the SC nucleus of *Tead1* cKO (Figure 1D),

presumably due to other TEADs in the absence of TEAD1.

TEAD1 is required for myelin formation and growth

Tead1 cKO mice were smaller than WT mice at all ages, with labored breathing and

widely splayed fore- and hindlimbs, indicative of severe peripheral neuropathy (Figure 2A,

Video 1). Because of the neuropathy, *Tead1* cKO mice were humanely euthanized before P60.

Tead1 cKO SNs were noticeably translucent, indicative of reduced myelin content (Figure 2B).

Electrical stimulation of cKO SNs elicited low and dispersed compound muscle action potentials

(CMAPs) (Figure 2C). Nerve conduction velocity (NCV) in cKO was ~8 m/s, which was

markedly reduced from ~40 m/s in WT mice (Figure 2D).

Next, we used semi-thin and transmission electron microscopy (TEM) to examine the

processes of developmental myelination and Remak bundle formation. We first examined SNs at

P3, when, in WT, many large axons had already been radially sorted, establishing 1:1

relationship with promyelinating SCs, and myelination had begun. Compared to WT, *Tead1* cKO

SNs were thinner, unsorted bundles of axons were larger and more numerous, and myelination

was minimal (Figure 2E). Nevertheless, we frequently observed large axons that had been sorted

and established a 1:1 relationship with promyelinating SCs, as in WT. Therefore, radial sorting

was delayed, but not blocked, in *Tead1* cKO.

At P22, when WT SNs were fully myelinated, *Tead1* cKO SNs were largely

unmyelinated and exhibited areas of unsorted axon bundles, of sorted but unmyelinated axons,

and of thinly myelinated axons (Figure 2F, Figure 2-figure supplement 1A). Notably, at P50, this

phenotype was unchanged from that at P22: areas of unsorted bundles, promyelination, and thin

myelination persisted (Figure 2G, Figure 2-figure supplement 1A). TEM analysis confirmed that

myelin was indeed thinner in *Tead1* cKO than in WT at both P22 and P50 (Figure 2F, 2G, Figure

2-figure supplement 1B). The myelin thickness measured as g-ratio, i.e., the ratio between the

axon diameter and the fiber diameter (axon plus myelin sheath), did not significantly change in

cKO between P22 and P50 (Figure 2H), suggesting that growth of the myelin sheath relative to

axon diameter did not increase over this period. These data suggest that myelination in *Tead1*

cKO was arrested rather than delayed. Plots of g-ratio versus axon diameter showed that the

mutant defect was present in axons of all diameters but was particularly notable in those of larger

diameter (Figure 2I, 3J). The mean axon diameter was also reduced in cKO at P22, though it

increased with age (Figure 2K). Collectively, these data show that TEAD1 is required for timely

radial sorting, myelin formation and growth during development. Although some axons become

myelinated in *Tead1* cKO, their myelin sheaths fail to properly grow in proportion to axon

diameter in the absence of TEAD1.

TEAD1 is required for proper formation of Remak bundles

Remak bundle formation was also abnormal in *Tead1* cKO mice. In WT SNs at P22 and

P50, small diameter axons were present in Remak bundles, with each C-fiber ensheathed by SC

cytoplasm (Figure 2F, 2L). In *Tead1* cKO, C-fibers were also present in bundles in which axons

were rarely larger than 1 μm in diameter, suggesting that small diameter axons were almost fully

sorted. However, at both P22 and P50, the observed bundles were frequently small, often the size

of a large myelinated axon, and were occasionally thinly myelinated. Notably, a SC surrounded

each bundle but individual C-fibers of most Remak bundles were not ensheathed by SC

cytoplasm (Figure 2L). Quantification showed that ~80% of C-fibers were not ensheathed by SC

cytoplasm in P50 *Tead1* cKO SNs (Figure 2M). Thus, TEAD1 is required to develop proper

association of C-fibers with non-myelinating SCs in Remak bundles.

TEAD1 regulates SC proliferation both positively and negatively to promote myelination

For myelination and Remak bundle formation to occur properly, timely generation of

sufficient numbers of SCs is essential. To determine if the defects of myelination and Remak

bundles in *Tead1* cKO are due to defective SC proliferation, we counted SCs. At P3, SC number

in cKO SNs was about half that of WT (Figure 3A). Consistently, the number of SCs in the cell

cycle (Ki67-positive; Figure 3B) and in S-phase (EdU-positive; Figure 3C) was greatly reduced

in cKO at P3. Thus, TEAD1 is required for optimal SC proliferation during development.

Notably, at P22 and P50, SC numbers in cKO were markedly increased and became more

than twice those of WT (Figure 3A). As cKO SC numbers were similar at P22 and P50, we

reasoned that between P3 and P22, SC proliferation must occur at an abnormally high rate.

Indeed, the number of SCs that were Ki67-positive or EdU-positive was significantly increased

in cKO at P22 (Figure 3B, 3C). The increased SC numbers were not due to proliferation of

residual SCs that express TEAD1: TEAD1 was absent in most EdU-positive SCs in *Tead1* cKO

(Figure 3D). These data suggest that, during development, SCs switch from requiring TEAD1 for

optimal proliferation, to requiring TEAD1 for timely exit from the cell-cycle, which is a

prerequisite for SC differentiation and the onset of myelination (Hindley & Philpott, 2012).

TEAD1 is required for Krox20 to upregulate myelin proteins, but not for Krox20

upregulation

Our findings thus far show that, in the absence of TEAD1 in the SCs of *Tead1* cKO, most

axons are not myelinated, which grossly phenocopied *Yap1/Wwtr1* iDKO (*Yap1* and *Wwtr1*

encode the proteins commonly listed as YAP and TAZ, respectively). *Yap1/Wwtr1* cDKO

exhibited a complete failure of myelination due to arrest of their SCs as promyelinating SCs, in

part because they fail to upregulate Krox20/Egr2 (Early Growth Response 2), which drives

transcription of myelin proteins (Grove et al., 2017). To determine whether TEAD1 is required

for Krox20 upregulation, we first investigated Oct6 (Pou3f1, Scip), whose upregulation precedes

and is required for Krox20 upregulation; Oct6 is subsequently downregulated, possibly by

Krox20 in a feedback mechanism (Ryu et al., 2007; Zorick, Syroid, Brown, Gridley, & Lemke,

1999). Western blotting of P50 SN lysates showed minimal expression of Oct6 in WT, but

abnormally upregulated expression in *Tead1* cKO (Figure 4A). Immunohistochemistry of P50

cKO SNs also showed intense Oct6 expression in ~80% of SC nuclei; Oct6 expression was

minimal in WT (Figure 4B). Notably, at both P11 and P50, a similar percentage of SCs

expressed Krox20 in *Tead1* cKO, as in WT SNs (Figure 4C). In Western blots, Krox20 was

upregulated >4-fold in cKO compared to WT (Figure 4A). As adult *Tead1* cKO mice contain ~2-

fold more SCs than WT (Figure 3A), these data suggest that each SC in *Tead1* cKO contains ~2-

fold more Krox20 than WT. Thus, in contrast to SCs lacking YAP/TAZ, SCs lacking TEAD1

upregulate Krox20, suggesting that TEAD1 is dispensable for YAP/TAZ to upregulate Krox20.

Consistent with the idea that Krox20 is pivotal for myelin gene transcription, expression

of myelin basic protein (MBP), and myelin protein zero (MPZ), was abundant in WT P50 SNs

(Figure 4A). In contrast, despite ample expression of Krox20, both MBP and MPZ were

markedly decreased in *Tead1* cKO, indicating that Krox20 fails to properly upregulate myelin

proteins without TEAD1. To exclude the possibility that Krox20 expressed in cKO SCs is

defective, we took advantage of the earlier finding that loss of Krox20 or its co-factors, Nab1/2,

causes SCs to remain aberrantly in the cell-cycle (Le, Nagarajan, Wang, Svaren, et al., 2005;

Zorick et al., 1999). Therefore, we studied whether EdU-positive SCs in the cell-cycle of P11

cKO were also positive for Krox20, which would indicate defective Krox20. We found that,

although ~65% of SC nuclei contained Krox20, only a few EdU-positive SCs exhibited even

minimal Krox20 immunoreactivity in either WT or *Tead1* cKO (Figure 4D). Thus, despite high

levels of functional Krox20, myelin protein expression was greatly reduced in cKO SCs,

suggesting that TEAD1 is dispensable for Krox20 upregulation but required for Krox20 to

upregulate transcription of myelin proteins during development.

TEAD1 and YAP/TAZ upregulate cholesterol biosynthetic enzymes, FDPS and IDI1

Myelination of peripheral nerves by SCs requires a large amount of lipid and cholesterol

biosynthesis. De novo biosynthesis of fatty acids (integral building blocks of lipids) and

cholesterol is crucially important for SCs to initiate myelination and to grow a myelin sheath

(Montani et al., 2018; Norrmen et al., 2014; Saher et al., 2009; Verheijen et al., 2009). YAP/TAZ

may regulate these pathways, as suggested by reduced mRNA expression in *Yap1/Wwtr1* cDKO

of SREBPs TFs (Sterol regulatory element binding proteins) and their target enzymes involved

in lipid/cholesterol biosynthesis (Deng et al., 2017; Poitelon et al., 2016). We therefore

investigated, at the protein level, whether TEAD1 and YAP/TAZ are required for fatty acid

and/or cholesterol synthesis. To test this premise, we examined expression of SREBP1 and its

target SCD1 (Stearoyl-CoA desaturase-1), which upregulates fatty acid synthesis, and of

SREBP2 and its targets HMGCR (HMG-CoA reductase), FDPS (Farnesyl Diphosphate

Synthase) and IDI1 (Isopentenyl-diphosphate delta isomerase 1), which upregulate cholesterol

synthesis (Leblanc et al., 2005; Verheijen et al., 2009). Because SREBP pathways are

upregulated early postnatally for myelination in WT, their upregulation may simply be delayed

in developmentally delayed *Tead1* cKO. We therefore investigated SREBP pathways both at P8

and P40 in WT and *Tead1* cKO mice. Western blot analysis showed that levels of nuclear

SREBP1 were not affected by loss of TEAD1 at P8, and only slightly reduced at P40 (Figure

5A). SCD1 was reduced at P8, but slightly increased at P40. In contrast to SREBP1, SREBP2

was ~50% reduced in *Tead1* cKO at both P8 and P40. Despite marked reduction in SREBP2,

HMGCR was not reduced at P8 or P40. However, both FDPS and IDI1 levels were greatly

reduced at both P8 and P40 (Figure 5A).

It is possible that HMGCR was not reduced in cKO due to TEAD1 redundancy and

compensation by other TEADs. If so, then HMGCR levels should be reduced in *Yap1/Wwtr1*

cDKO, because YAP/TAZ are the primary partners for all TEADs (Holden & Cunningham,

2018; Zheng & Pan, 2019). To test this idea, we examined SREBP pathways in *Yap1/Wwtr1*

cDKO SNs at P60 because developmental delay is greater in *Yap1/Wwtr1* cDKO than in *Tead1*

cKO SNs (Grove et al., 2017). We found that, as in *Tead1* cKO, SREBP1 levels were unaffected

and SCD1 levels were markedly increased by the loss of YAP/TAZ in cDKO at P60 (Figure 5B).

Unlike in *Tead1* cKO, levels of SREBP2 were not markedly reduced in Yap1/Wwtr1 cDKO. As

in *Tead1* cKO, however, HMGCR levels were not reduced in Yap1/Wwtr1 cDKO, whereas

levels of FDPS and IDI1 were greatly reduced. These results therefore indicate that YAP/TAZ-

TEAD1 transcriptional complex selectively upregulates IDI1 and FDPS, contributing to

cholesterol biosynthesis. Together with the lack of synthesis of myelin proteins, this reduction in

cholesterol biosynthesis may prevent myelin formation and growth in *Tead1* cKO.

TEAD1 is largely dispensable for myelin maintenance in adult Schwann cells

Because TEAD1 is required for myelin formation and growth and Remak bundle

formation during development, we next investigated its role in maintenance and regeneration of

peripheral nerves in adults. We crossed *Tead1*^fl/fl mice with mice carrying a tamoxifen-inducible

Plp1-CreER (Leone et al., 2003) to generate *Plp1* -CreER*; Tead1*^fl/fl (hereafter *Tead1* iKO) and

inducibly inactivated SC *Tead1* in adult *Tead1* iKO mice. Western blotting showed markedly

reduced TEAD1 in *Tead1* iKO (Figure 6-figure supplement 1A). Residual TEAD1 is likely not

associated with SCs. Indeed, immunostaining of WT and induced iKO SNs consistently showed

that TEAD1 was deleted in >95% of SC nuclei in both intact and regenerating iKO SNs, whereas

TEAD1 was present in 100% of SC nuclei of WT SNs (Figure 6-figure supplement 1B, 1D). As

anticipated, iKO SC nuclei in both intact and regenerating nerves were strongly immunostained

by pan-TEAD antibody, presumably indicating the presence of other TEADs (Figure 6-figure

supplement 1C, 1E). We also found that YAP/TAZ were transcriptionally functional in adult SCs

lacking TEAD1, as indicated by strong nuclear expression of YAP/TAZ in SCs of both intact
and regenerated iKO SNs (Figure 6-figure supplement 1F, 1G).

        Tead1 iKO mice appeared normal without obvious signs of peripheral neuropathy up to 5
months post-tamoxifen, the longest time of observation (Figure 6A). TEM analysis of iKO SNs
between 3 to 5 months post-tamoxifen revealed, however, some large axons undergoing
demyelination or completely demyelinated, which were undetectable in WT SNs (Figure 6B).
The demyelinated axons were observed in discrete regions of SNs, indicating segmental
demyelination (<5% large axons on 12 TEM sections exhibiting demyelinated axons). NCV of
these iKO mice did not significantly differ from that of WT (Figure 6C). Therefore, loss of
TEAD1 in adult SCs evoked only modest focal demyelination in a small number of axons, which
did not cause significant functional impairment of peripheral nerves.

TEAD1 is required for maintaining structural integrity of Remak bundles

        Next, we examined Remak bundles in *Tead1* iKO mice. As in WT SNs, C-fibers in WT
mice were individually ensheathed by SC cytoplasm. In stark contrast, in *Tead1* iKO, many C-
fibers in most Remak bundles were not ensheathed by SC cytoplasm (Figure 6D). Furthermore,
some small Remak bundles were occasionally myelinated (Figure 6E), as in *Tead1* cKO SNs.
Thus, TEAD1 is required for formation as well as maintenance of Remak bundles.

TEAD1 is dispensable for YAP/TAZ to maintain Krox20, FDP, and IDI1 expression

        To investigate why the demyelinating phenotype of *Tead1* iKO is much milder than that
of *Yap1/Wwtr1* iDKO, we first investigated expression of Krox20, which is required for myelin
maintenance (Decker et al., 2006). We previously observed that Krox20 is markedly
downregulated in *Yap1/Wwtr1* iDKO SCs, which correlated with frequent segmental
demyelination and markedly reduced NCV (Grove et al., 2017). As a control, we injected *Plp1-
CreER; Yap1^{fl/fl}; Wwtr1^{fl/fl}* mice with tamoxifen and reproduced our previous result: Krox20 was
dramatically downregulated in most *Yap1/Wwtr1* iDKO SCs only 2 weeks post-tamoxifen
(Figure 7A). In contrast, in *Tead1* iKO, Krox20 expression was indistinguishable from WT even
at 6 weeks post-tamoxifen (Figure 7B). We then investigated Oct6 expression. Unlike in
Yap1/Wwtr1 iDKO SNs (Grove et al., 2017), Oct6 was not elevated in *Tead1* iKO SNs (Figure
7C). Thus, TEAD1 is dispensable for maintaining Krox20 expression in adult SCs.

Next, we investigated SREBP pathways, which are required for developmental

myelination (Saher et al., 2009; Verheijen et al., 2009). Whether these pathways are also

involved in myelin maintenance has not been studied. We first assessed the effects of acute loss

of YAP/TAZ on these pathways in adult SCs. Western blotting of WT and *Yap1/Wwtr1* iDKO

SN lysates showed that the SREBP1 pathway and expression of SREBP2 and HMGCR in the

SREBP2 pathway were unaffected in *Yap1/Wwtr1* iDKO (Figure 7D). However, FDPS and IDI1

levels were significantly downregulated in *Yap1/Wwtr1* iDKO, recapitulating the effect of

developmental loss of YAP/TAZ on these proteins (Figure 5B). Thus, the acute downregulation

of these key enzymes for cholesterol biosynthesis correlates with the acute demyelination

previously observed in *Yap1/Wwtr1* iDKO mice. This result suggests that YAP/TAZ maintain

myelination through both myelin gene expression and the cholesterol biosynthesis pathway.

Western blotting of *Tead1* iKO SN lysates revealed that the SREBP pathways were not

significantly affected by loss of TEAD1 (Figure 7E), unlike *Tead1* cKO in which FDPS and IDI1

were markedly downregulated (Figure 5A). This result suggests that TEAD1 is dispensable for

cholesterol synthesis in adult SCs.

TEAD1 is absolutely required for myelination of regenerating axons after nerve injury.

Next, we investigated whether TEAD1 is required for axon regeneration and

remyelination after nerve injury. YAP/TAZ are not required for axon regeneration but are crucial

for remyelination (Grove et al., 2017; Jeanette et al., 2021). Because *Yap1/Wwtr1* iDKO mice die

~2 weeks post-injury (wpi), it remains possible that remyelination is delayed rather than

completely blocked. In contrast, *Tead1* iKO mice are not susceptible to early death. We,

therefore, examined them 4 weeks after crushing SN (Figure 8A), when hindlimb muscles are

fully reinnervated in WT. We first verified efficient deletion of TEAD1 in >95% SCs of *Tead1*

iKO mice after nerve crush (Figure 6-figure supplement 1D). Longitudinal sections of iKO SNs

showed numerous Tuj1-labeled axons that regenerated along the distal nerve stump, as in WT

SNs (Figure 8B). However, MBP immunoreactivity was greatly reduced in the distal nerve

stump of *Tead1* iKO, compared to WT, suggesting that axons regenerated but were not

remyelinated. We also found that regenerated axons reached hindlimb muscles and reinnervated

neuromuscular junctions (NMJs) in iKO, as in WT (Figure 8C). Moreover, *Tead1* iKO

frequently exhibited features of paralyzed muscles, such as axon terminals that grew past the

synaptic area, fragmented clusters of acetylcholine receptors (AChRs), and satellite extra-

synaptic AChRs (Wright, Cho, & Son, 2007). Semithin and EM analysis confirmed virtual

absence of myelinated axons in iKO SNs; although large axons were associated with SCs, they

were not myelinated (Figure 8D). Accordingly, electrical stimulation of iKO SNs did not elicit

CMAPs or NCV (Figure 8E). Thus, like YAP/TAZ, TEAD1 in SCs is not required for axon

regeneration but is required for remyelination after nerve injury.

After injury, SCs transdifferentiate into growth-promoting repair SCs, which proliferate

and reach a peak around 3 days post-injury (dpi) (Clemence, Mirsky, & Jessen, 1989; Jessen &

Mirsky, 2016; Tricaud & Park, 2017). Quantitative analysis of SC proliferation at 3 dpi by

counting $EdU^+/Sox10^+$ SCs showed that a similar percentage of SCs were in S-phase in both WT

and *Tead1* iKO SNs (Figure 8F), suggesting that, like YAP/TAZ (Grove et al., 2017), TEAD1 is

not required for proliferation of repair SCs. Lastly, we counted $Krox20^+$ and $Oct6^+$ SCs at 4 wpi.

Notably, in contrast to *Yap1/Wwtr1* iDKO SCs, which fail to upregulate Krox20 after injury

(Grove et al., 2017), Krox20 was upregulated in *Tead1* iKO SCs, as in WT SCs (Figure 8G).

Furthermore, Oct6 was aberrantly elevated in *Tead1* iKO SCs, like in *Tead1* cKO SCs during

development (Figure 8H). These results suggest that, similar to developmental myelination,

remyelination requires TEAD1 for Oct6 downregulation and for Krox20 to upregulate myelin

proteins. Additionally, almost complete absence of remyelination at 4 wpi highlights the non-

redundant, crucial role of TEAD1 in the remyelination of regenerating nerves after injury.

DISCUSSION

The present study conclusively establishes TEAD1 as a crucial TF mediating myelin formation,

growth, and regeneration. TEAD1 promotes myelination by dually regulating SC proliferation,

by enabling induced Krox20 to upregulate myelin proteins, and by upregulating cholesterol

biosynthetic enzymes. Our findings also suggest that TEAD1 is critical in the proper formation

and maintenance of Remak bundles.

Unique, redundant, and opposing roles of TEAD1

Although SCs express other TEADs, *Tead1* cKO mice grossly phenocopy *Yap1/Wwtr1* cDKO

mice. They exhibit abnormal SC proliferation, delayed radial sorting, blocked myelin formation

and remyelination, and normal axon regeneration. Furthermore, *Tead1* cKO mice, like

Yap1/Wwtr1 cDKO, are defective in regulating Oct6, Krox20, myelin proteins, and enzymes that

synthesize cholesterol. These similarities show that YAP/TAZ regulate both SC proliferation and

differentiation, the two opposing processes crucial for myelination, by partnering mainly with

TEAD1 for developmental myelination and remyelination after injury. We also observed

multiple differences between *Tead1* cKO and *Yap1/Wwtr1* cDKO mice. Unlike *Yap1/Wwtr1*

cDKO, *Tead1* cKO initiates myelin formation in some developing axons and maintains

myelination in almost all adult axons. Additionally, in *Tead1* cKO, Krox20 is upregulated and a

large excess of SCs is generated. It is also notable that, whereas some developing axons are

myelinated in *Tead1* cKO, almost all axons regenerating after injury are unmyelinated. These

differences are likely due to stage-dependent redundancy and opposing roles of TEADs.

Consistent with this notion, loss of TEAD1 does not affect nuclear localization of YAP/TAZ,

which depends on interactions with TEADs (Diepenbruck et al., 2014; J. Kim et al., 2020; Kofler

et al., 2018; Yao et al., 2018). TEAD1-4 are all co-expressed in SCs through multiple stages of

nerve development, maintenance, and regeneration (Fernando et al., 2016; Grove et al., 2017; X.

He et al., 2018). Notably, nuclear retention of YAP/TAZ occurs in *Tead1* cKO and iKO SCs

when TEAD1 has a unique role in SCs for developmental myelination and remyelination. Thus,

other TEADs can retain nuclear YAP/TAZ, thereby supporting transcriptional activity of

YAP/TAZ, but cannot substitute for some or most of the functions of TEAD1 in a stage-

dependent manner.

It is also notable that, like YAP/TAZ, the unique roles of TEAD1 change dynamically in

successive steps of myelination, even though other TEADs are present. Thus, TEAD1 is

uniquely required for timely proliferation of SCs; uniquely opposes SC proliferation to promote

cell-cycle exit after radial sorting; redundantly promotes upregulation of Krox20; and then

uniquely promotes upregulation of myelin proteins and cholesterol biosynthetic enzymes. Such

unique and redundant roles for TEAD1 in the presence of other TEADs are supported by cell-

based transcription assays, in which YAP-TEAD4 can substitute for YAP-TEAD1 in

upregulating transcription from the Krox20 promoter, but not from the MPZ intronic enhancer

(Fernando et al., 2016). Stage-dependent function of TEADs was also reported in cortical

development, another multi-step differentiation program. As in nerve development, multiple

TEADs (TEAD1-3) are co-expressed throughout cortical development. Depending on the

differentiation stage, TEAD1 uniquely promotes proliferation, redundantly promotes

proliferation with TEAD2 and 3, or redundantly promotes differentiation with TEAD3, when it

is opposed by TEAD2 (Mukhtar et al., 2020).

As *Yap1/Wwtr1* cDKO SCs arrest at the promyelination stage, we were unable to investigate the

role of YAP/TAZ in myelin growth (Grove et al., 2017). In contrast, some *Tead1* cKO SCs can

myelinate but fail to grow myelin sheaths proportional in size to axon diameter. Importantly,

such myelinating SCs apparently lack TEAD1 because TEAD1 deletion is highly efficient in

Tead1 cKO mice. Support for a role of TEADs in the regulation of myelin sheath thickness

comes from a study of TEAD4 cKO mice, in which SCs lacking TEAD4 gradually upregulate

myelin protein expression, resulting in adult mice with abnormally thick myelin (X. He et al.,

2018). Thus, it is possible that TEADs 1 and 4 oppose each other in the regulation of myelin

sheath growth. Likewise, they may oppose each other in the regulation of SC proliferation during

development, resulting in a large excess of SCs at P22 and P50 in *Tead1* cKO.

TEAD1 regulation of Krox20 and myelin gene expression

Previous work suggests that YAP/TAZ-TEAD1 promote myelination in part through direct

upregulation of Krox20 (Deng et al., 2017; Grove et al., 2017). Unlike *Yap1/Wwtr1* cKO mice,

Tead1 cKO mice upregulate Egr2/Krox20, but they fail to upregulate myelin proteins and

cholesterol biosynthetic enzymes. Our findings extend previous work to suggest that TEAD1

functions redundantly in direct upregulation of Krox20, but that its function is required for

upregulated Krox20 to promote myelination. Krox20 activity may be repressed in *Tead1* cKO by

Oct6, which remains abnormally elevated. In support of this possibility, persistently elevated

Oct6 overrides Krox20 to inhibit myelin gene expression (Monuki, Kuhn, & Lemke, 1993; Ryu

et al., 2007). Other mice with defective lipid biosynthesis and a phenotype like *Tead1* cKO also

have abnormally elevated Oct6 after Krox20 upregulation (Norrmen et al., 2014; Sherman et al.,

2012). An alternative possibility is that a transcriptional co-activator that synergizes with Krox20

is absent in *Tead1* cKO SCs. We speculate that TEAD1 itself may function as such a factor.

Supporting this idea are reports that gene regulatory elements function by co-binding several

TFs, which synergistically activate transcription (Heinz, Romanoski, Benner, & Glass, 2015;

Ren & Yue, 2015), as at the Krox20 MSE (Myelinating Schwann cell element) which is the

enhancer responsible for Krox20 expression (Ghislain & Charnay, 2006; Grove et al., 2017; Y.

He et al., 2010; Kao et al., 2009). Krox20 synergizes with Sox10 to activate myelin genes (Jones

et al., 2007) and with SREBP to activate lipid biosynthesis genes (Leblanc et al., 2005). Like

Krox20, TEAD1 can directly promote transcription from PMP22 and MPZ myelin gene

enhancers (Fernando et al., 2016; Lopez-Anido et al., 2016). Alternatively, or additionally,

TEAD1 may function indirectly by upregulating co-factors, such as SREBP2, which is reduced

in *Tead1* cKO. Notably, lipid biosynthesis is necessary to downregulate Oct6 and upregulate

myelin gene transcription during developmental myelination (Saher et al., 2009; Verheijen et al.,

2009). An interesting possibility, therefore, is that YAP/TAZ-TEAD1 and Krox20 may

cooperatively downregulate Oct6 and upregulate myelin gene expression in part by promoting

lipid biosynthesis.

YAP/TAZ-TEAD1 regulation of cholesterol biosynthesis

SREBP1 levels are not altered in *Tead1* cKO or *Yap1/Wwtr1* cDKO, suggesting that fatty acid

synthesis is not directly regulated by TEAD1 or YAP/TAZ. It is likely that the reduced levels of

SCD1 at P8 and slight increase at P40 in *Tead1* cKO, and the large increase in SCD1 in

Yap1/Wwtr1 cDKO at P60 are due to developmental delay in these mutants. We did not observe

downregulation of HMGCR in *Tead1* cKO or *Yap1/Wwtr1* cDKO mice at P8 or P40, which

contradicts the report of reduced mRNA expression of HMGCR in *Yap1/Wwtr1* mutants at P3

and P5 (Deng et al., 2017; Poitelon et al., 2016); we suggest this difference is due to

developmental delay in *Tead1* cKO and *Yap1/Wwtr1* cDKO. However, marked downregulation

of IDI1 and FDPS in *Tead1* cKO, *Yap1/Wwtr1* cDKO, and *Yap1/Wwtr1* iDKO suggests that

cholesterol biosynthesis is defective in these mutants. In support of this notion, SC-selective

deletion of SCAP (SREBP cleavage activation protein), which is required to produce active

SREBP1 and SREBP2 (Verheijen et al., 2009), or of Squalene synthase, an essential enzyme in

cholesterol biosynthesis (Saher et al., 2009), evokes phenotypes which are remarkably similar to

those of *Tead1* cKO mice: both mutants have an increased number of SCs, a severe delay at the

promyelination stage, and persistently thin myelin. In addition, in SCAP cKO, Oct6 is

upregulated continuously and Squalene synthase cKO fails to upregulate myelin gene

transcription, resulting in severely reduced MBP, MPZ, and PMP22 protein levels.

How might YAP/TAZ-TEAD1 regulate IDI1 and FDPS? Multiple studies show that YAP/TAZ

can activate the mTORC1 pathway (Hu et al., 2017; M. Kim, Kim, Johnson, & Lim, 2015), and

that IDI1 and FDPS levels are reduced in SNs of Raptor cKO, a core component of mTORC1

(Norrmen et al., 2014). Furthermore, the Raptor cKO phenotype is like that of *Tead1* cKO,

exhibiting an increased number of SCs, normal upregulation of Krox20, persistent upregulation

of Oct6, delay at the promyelination stage, and thin myelin. However, HMGCR protein levels

are reduced, and aberrant upregulation of Oct6 is transient (Norrmen et al., 2014). After nerve

injury, formation of repair SCs is delayed in Raptor mutants (Norrmén et al., 2018), unlike

TEAD1 and YAP/TAZ mutants. These differences suggest that YAP/TAZ-TEAD1 are not

directly upstream of mTORC1. Another possibility is that YAP/TAZ-TEAD1 may directly

regulate transcription of IDI1 and FDPS genes. Such direct regulation of cholesterol biosynthetic

enzymes has been observed for other TFs in SCs (M. Kim et al., 2018; Leblanc et al., 2005).

TEAD1 regulation of Remak bundles

In both *Tead1* cKO and iKO mice, there is defective ensheathment of C-fibers by non-

myelinating SCs in Remak bundles. This phenotype is distinct from those mutants in which SCs

surround and occasionally myelinate bundles filled with large and small caliber axons, which are

considered radial sorting defects (Feltri, Poitelon, & Previtali, 2016). Remak bundles are

occasionally myelinated in WT SNs during early postnatal development, but this defect is

corrected and does not persist in adults (Rasi et al., 2010). Thus, the defective Remak bundles

that we observed in *Tead1* cKO are not due to developmental delay.

Remak bundles form normally at P60 in *Yap1/Wwtr1* cDKO, although many large bundles

containing mixed caliber axons are present during development (Grove et al., 2017). The

difference between *Tead1* and *Yap1/Wwtr1* mutants is supported by the presence of TEAD1 and

absence of YAP/TAZ in non-myelinating SC nuclei (Grove et al., 2017). Notably, defective SC

ensheathment of C-fibers is observed in various SC developmental mutants, including those with

defective SC basal lamina (Rasi et al., 2010) and overexpression of NRG1-IIIb (Neuregulin 1

type-IIIb) in axons (Gomez-Sanchez et al., 2009). Therefore, it is possible that TEAD1 mediates

extracellular signaling from the SC basal lamina and/or axons that is responsible for forming and

maintaining proper association of SCs with C-fibers. Notably, SC ensheathment of C-fibers is

required for normal NCV (Taveggia et al., 2005), and NCV in the mutants overexpressing

NRG1-IIIb is significantly reduced compared to WT (Gomez-Sanchez et al., 2009). Future

studies will address NRG1 signaling and NCV in cKO and iKO mice lacking TEAD1 and/or

other TEADs.

MATERIALS AND METHODS

Reagent type (species) or resource	Designation	Source or Reference	Identifiers	Additional Information
Strain, strain background (*Mus musculus*)	C57Bl/6	Jackson Laboratory	Stock #: 000664; RRID:IMSR JAX:000664	
Genetic reagent (*M. musculus*)	*Plp1*-Cre-ERT2		MGI:2663093	(Leone et al., 2003)
Antibody	anti-Yap/Taz (rabbit monoclonal)	Cell Signaling Technology	D24E4, #8418 RRID:AB_10950494	IHC 1:200 Western 1:1000
Antibody	anti-SCG10 (rabbit monoclonal)	Novus Biologicals	NBP1-49461 RRID:AB_10011569	IHC 1:5000
Antibody	anti-Yap (rabbit monoclonal)	Cell Signaling Technology	D8H1X, #14074 RRID:AB_2650491	IHC 1:200
Antibody	anti-Sox10 (goat polyclonal)	R&D Systems	#AF-2864 RRID:AB_442208	IHC 1:100
Antibody	anti-Sox10 (rabbit monoclonal)	Abcam	EPR4007, #ab155279 RRID:AB_2650603	IHC 1:250
Antibody	anti-Egr2 (rabbit polyclonal)	Professor Dies Meijer, University of Edinburgh		IHC 1:4000
Antibody	anti-Oct6 (rabbit monoclonal)	Abcam	EP5421, #ab126746 RRID:AB_11130256	WB 1:1000

Antibody	anti-Oct6 (rabbit polyclonal)	Abcam	#ab31766 RRID:AB_776899	IHC 1:800
Antibody	anti-c-Jun (mouse monoclonal)	BD Transduction Laboratories	#610326 RRID:AB_397716	IHC 1:500
Antibody	anti-c-Jun (rabbit monoclonal)	Cell Signaling Technology	60A8, #9165 RRID:AB_2130165	WB 1:1000
Antibody	anti-pS63-c-Jun (rabbit polyclonal)	Cell Signaling Technology	#9261 RRID:AB_2130162	IHC 1:100
Antibody	anti-Ki67 (rabbit polyclonal)	Abcam	#ab15580 RRID:AB_443209	IHC 1:200
Antibody	anti-p75NGFR (goat polyclonal)	Neuromics	#GT15057 RRID:AB_2737189	IHC 1:400
Antibody	anti-Tubulin β3 (rabbit polyclonal)	Biolegend	#802001 RRID:AB_2564645	IHC 1:1000
Antibody	IRDye-680 (goat anti-mouse)	LI-COR	#926-32220 RRID:AB_621840	WB 1:15,000
Antibody	HRP-Goat anti-mouse secondary antibody	Jackson Immunoresearch	#715-035-150 RRID:AB_2340770	WB 1:12,000
Antibody	HRP-Goat anti-rabbit secondary antibody	Jackson Immunoresearch	#115-055-062 RRID:AB_2338533	WB 1:12,000
Chemical compound, drug	Araldite 6005	EMS	#10920	
Chemical compound, drug	DDSA	EMS	#13710	
Chemical compound, drug	DBP	EMS	#13101	
Chemical compound, drug	BDMA	EMS	#11400-25	
other	Coated grids (100 mesh)	EMS	#FF100-Cu	
Chemical compound, drug	Osmium tetroxide (4% solution)	EMS	#19170	

Chemical compound, drug	Lead nitrate	EMS	#17900	
Chemical compound, drug	Sodium citrate	EMS	#21140	
Chemical compound, drug	Uranyl acetate	EMS	#22400	
Chemical compound, drug	Sodium borate	EMS	#21130	
Chemical compound, drug	Toluidine blue	EMS	#22050	
Chemical compound, drug	Paraformaldehyde	Sigma-Aldrich	#158127	
Commercial assay or kit	Click-It EdU Alexa Fluor 594 kit	ThermoFisher Scientific	#C10339	
Chemical compound, drug	EdU	ThermoFisher Scientific	#E10187	
Chemical compound, drug	Tamoxifen	Sigma-Aldrich	#T5648	
other	DAPI stain	Invitrogen	#D1306	IHC 1:250
Antibody	Alexa 488, 568 or 647 secondaries	Jackson Immunoresearch		IHC 1:250 to 1:1000
software, algorithm	Image Studio Lite	LI-COR, Inc		
software, algorithm	Prism	GraphPad Software, Inc		
software, algorithm	Stata	StataCorp LP		Mann-Whitney test

Animals

All surgical procedures and animal maintenance complied with the National Institute of Health

guidelines regarding the care and use of experimental animals and were approved by the

Institutional Animal Care and Use Committee (Protocol# 4920) of Temple University,

Philadelphia, PA, USA. Generation and genotyping of mice carrying targeted loxP sites in the

Tead1 gene have been described previously (Wen et al., 2017). Targeted ablation of *Tead1* in

Schwann cells of embryonic mice was achieved by crossing *Tead1*$^{fl/fl}$ mice with mice transgenic

for *Cre* driven by the *Mpz* promoter (*Tead1* cKO mice). Targeted ablation of *Tead1* in Schwann

cells of adult mice was by crossing *Tead1*$^{fl/fl}$ mice with mice transgenic for the inducible Cre

recombinase *Cre-ERT2* under control of the *Plp1* promoter (*Tead1* iKO mice). Control mice

contained floxed *Tead1* alleles but were not transgenic for *Cre* or *Cre-ERT2*. *Mpz-cre*;*Yap1*$^{fl/fl}$;

Wwtr1$^{fl/fl}$ (*Yap1/Wwtr1* cDKO) and *Plp1-creERT2*; *Yap1*$^{fl/fl}$; *Wwtr1*$^{fl/fl}$ (*Yap1/Wwtr1* iDKO) mice

used in this study were generated and genotyped as described previously (Grove et al., 2017).

Both male and female mice were used in all experiments and were maintained on the C57BL/6

background.

Tamoxifen administration

Tamoxifen was prepared and injected into 6–8-week-old *Tead1* or *Yap1/Wwtr1* mutant mice as

previously described (Grove et al., 2017). A 10 mg/ml solution of tamoxifen was made in 10:1

sunflower oil: 100% ethanol. This solution was injected intraperoneally at a concentration of 0.2

mg/g body weight. Injection was once daily for 5 days, followed by a 2-day break, then once

daily for 5 consecutive days.

Nerve crush

Sciatic nerves of right hindlimbs were crushed 24 h after the final tamoxifen injection, as

described previously (Grove et al., 2020; Son & Thompson, 1995). A small skin incision was

made in the posterior thigh and calf of the animals anesthetized by isoflurane. The sciatic nerve

was crushed with a fine forceps (#5) for 10 seconds (3X) adjacent to the sciatic notch. The crush

site was marked using charcoal-coated forceps, and the wound was closed. To identify

proliferating Schwann cells, we intraperitoneally injected EdU (80 μg/g) eighty minutes before

euthanasia, as previously described (Grove et al., 2017).

Western blotting

After removal of epineurium and perineurium, sciatic nerves were snap-frozen in liquid nitrogen

and stored at -80ºC. For Western blotting, sciatic nerves were lysed in RIPA buffer (25 mM Tris

pH 7.5, 150 mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, 1 mM EDTA, 0.1% SDS);

for immunoprecipitation, lysis was in co-IP buffer (50 mM Tris pH 8.0, 150 mM NaCl, 0.5%

Triton X-100, 10% glycerol, 1 mM EDTA), both containing protease and phosphatase inhibitors.

Lysis was for 40 minutes on ice, followed by microcentrifugation at 14,000 rpm for 20 minutes.

Protein concentration was determined using the BCA assay (ThermoFisher).

Primary antibodies were used at the following concentrations for Western blots: rabbit

anti-Krox20 (Covance; 1:400), rabbit anti-Oct6 (Abcam ab126746; 1:1000), mouse anti-Myelin

Basic Protein (Covance SMI94; 1:1000), mouse anti-beta actin (SIGMA-Aldrich A5441;

1:5000), rabbit anti-TEAD1 (Cell Signaling D9X2L, #12292; 1:1000), rabbit anti-TEAD2

(Novus Biologicals, #NBP3-05267; 1:1000), rabbit anti-TEAD3 (Cell Signaling #13224;

1:1000), rabbit anti-TEAD4 (Santa Cruz, #sc-101184; 1:1000), rabbit anti-pan-TEAD (Cell

Signaling, #13295; 1:1000), rabbit anti-YAP/TAZ (Cell Signaling D24E4, #8418; 1:1000, 1:100

for immunoprecipitations), rabbit anti-phospho-YAP(Ser127) (Cell Signaling D9W2I #13008;

1:1000), rabbit anti-phospho-YAP(Ser397) (Cell Signaling D1E7Y #13619; 1:1000), rabbit-anti-

MPZ (Cell Signaling #57518; 1:1000), mouse anti-SREBP1 (Novus Biologicals, #NB600-582;

1:1000), rabbit anti-SCD1 (Cell Signaling, #2794; 1:1000), rabbit anti-SREBP2 (Bethyl

laboratories, #A303-125A), rabbit anti-HMGCR (Novus Biologicals, #NBP2-66888; 1:1000),

rabbit anti-FDPS (Proteintech, #16129-1-AP; 1:1000), rabbit anti-IDI1 (Proteintech, #11166-2-

AP; 1:1000). Control whole rabbit IgG for immunoprecipitations was from Jackson Laboratories

(011-000-003) and was used at 1µg/ml, which was several-fold greater than all other antibodies

used for immunoprecipitations. Secondary antibodies for Western blotting were HRP-linked

sheep anti-mouse IgG and donkey anti-rabbit (both GE Healthcare; 1:5000) and HRP-linked

mouse anti-rabbit, light-chain specific (Jackson ImmunoResearch 211-032-171; 1:20,000). Blots

were developed using ECL, ECL Plus (GE Healthcare) or Odyssey (LiCor).

Immunohistochemistry, confocal and widefield fluorescence microscopy

For immunostaining, sciatic nerves were removed and immediately fixed in 4%

paraformaldehyde in PBS for 1 hr. on ice. Nerves were washed 3 times in PBS, then stored in

15% sucrose in PBS overnight at 4°C for cryoprotection. Nerves were frozen-embedded in

cryoprotectant medium (Thermo Fisher Scientific, Waltham, MA) in isomethylbutane at -80°C.

7-10 μm sections from the nerves were cut using a cryostat (Leica Microsystems, Germany) and

collected directly onto glass slides. For immunolabeling, nerve sections were rehydrated in PBS,

permeabilized in 0.5% Triton/PBS for 20 min, washed with PBS, then blocked in 2% bovine

serum albumin (BSA) in PBS for 1 hr. Sections were incubated with primary antibodies in

blocking buffer overnight at 4°C in a hydrated chamber, washed with PBS, and incubated with

secondary antibodies in blocking buffer for 2 hrs. at room temperature. Sections were washed

with PBS, stained with DAPI for 10 min, and mounted with Vectashield mounting medium

(Vector Labs, Burlingame, CA).

Primary antibodies were used at the following concentrations for immunostaining of

cryosections: guinea pig anti-Sox10 (kind gift from Michael Wegner, University of Erlangen,

Bavaria, Germany; 1:1000), rabbit anti-Krox20 (Covance; 1:400), rabbit anti-Oct6 (Abcam

ab31766; 1:800), rabbit anti-Ki67 (Novocastra NCL-Ki67p; 1:200), rabbit anti-Neurofilament

200 (SIGMA-Aldrich N4142; 1:500), mouse anti-Neurofilament (Covance SMI 312; 1:1000),

mouse anti-Tubulin β3 (clone Tuj1, Covance #MMS-435P; 1:1000), mouse anti-SV2

(Developmental Studies Hybridoma Bank, Iowa; 1:10), rabbit anti-S100 (DAKO; 1:600), mouse

anti-Myelin Basic Protein (Covance SMI 94; 1:2000), rabbit anti-TEAD1 (Cell Signaling

D9X2L, #12292; 1:200), rabbit anti-pan-TEAD (Cell Signaling D3F7L, #13295; 1:200), rabbit

anti-YAP/TAZ (Cell Signaling D24E4, #8418; 1:200).

For immunostaining of extensor digitorum longus (EDL) and soleus muscles, mice were

perfused with 4% paraformaldehyde, and the muscles removed and post-fixed for 20 mins at

room temperature. Muscles were then incubated for 15 minutes with rhodamine-conjugated α-

bungarotoxin (Invitrogen, Carlsbad, CA), permeabilized in −20°C methanol for 5 min, and

blocked for 1 hr. in PBS containing 0.2% Triton and 2% BSA. The muscles were subsequently

incubated overnight at 4ºC in a cocktail of primary antibodies diluted in the blocking solution.

Axons and nerve terminals were labeled with mouse monoclonal antibodies to neurofilaments

(SMI 312; Sternberger Monoclonals, Baltimore, MD) and to a synaptic vesicle protein, SV2

(Developmental Studies Hybridoma Bank, Iowa City, IA). Schwann cells were labeled with

rabbit anti-cow S-100 polyclonal antibody (Dako, Carpentaria, CA). After incubation with the

primary antibodies, muscles were rinsed in PBS and incubated with secondary antibodies in the

blocking solution, for 1 hr. at room temperature. The muscles were then rinsed in PBS and

prepped for whole-mount by peeling the top layer of muscle fibers using microscissors and

forceps, as previously described (Son & Thompson, 1995). The resulting "muscle fillets" were

mounted on a slide and coverslipped.

Conventional fluorescence microscopy was performed using Olympus BX53 and Zeiss

Axio Imager 2 microscopes. Images were captured using Hamamatsu ORCA R2 C10600 or

Zeiss Axiocam HRm Rev 3 digital cameras, respectively, and Metamorph software. For confocal

microscopy, we used a Leica SP8 confocal microscope, with either a 40x 1.3 NA or 63x 1.4 NA

objective and Leica proprietary software. Acquired stacks were assembled using the maximum

projection tool, and figures were prepared using Adobe Photoshop.

Electron microscopy, histology, and morphometry

Mice were perfused intravascularly with 2.5% glutaraldehyde and 4% paraformaldehyde in 0.1M

cacodylate buffer pH 7.4. Sciatic nerves were removed and incubated in the same buffer for 90

min. at room temperature, then overnight at 4°C. Specimens were post-fixed for 1 hr. with OsO_4,

dehydrated, then embedded in araldite. Semi-thin (0.5 μm) and ultrathin (70 nm) sections were

cut using an ultramicrotome and stained with toluidine blue or uranyl acetate and lead citrate,

respectively. Semi-thin sections were examined using bright field on an Olympus BX53

microscope, and images were captured using Hamamatsu ORCA R2 C10600 digital camera and

Metamorph software. For analysis of the number of myelin profiles in control vs mutant sciatic

nerves, semi-thin sections from proximal-, mid- and distal regions of the sciatic nerve were

taken, and 2 non- overlapping images of each section were obtained using the 60x objective. For

electron microscopy, stained ultrathin sections were examined using a JEOL 1010 electron

microscope fitted with a Hamamatsu digital camera and AMT Advantage image capture

software.

Proliferation assays

The percentage of Schwann cells in S-phase was measured by EdU incorporation. Mice were

injected intraperitoneally and subcutaneously with 80 μg/ g body weight of EdU dissolved in

PBS. Eighty min. later, mice were sacrificed, and sciatic nerves removed and fixed for 1 hr. in

4% paraformaldehyde; longitudinal 7μm cryosections were used for analysis. EdU incorporation

was identified using the Click-iT EdU Alexa Fluor 555 labeling kit (Molecular Probes); the

recommended protocol was followed, and labeling was for 1 hr. After EdU labeling, sections

were immunostained with anti-Sox10 to identify Schwann cells. The percentage of Schwann

cells (Sox10+ nuclei) that were EdU+ was determined by counting confocal images using

ImageJ software.

Electrophysiology

The procedure for measuring nerve conduction velocity was an adaption of a previously used

protocol (Schulz, Walther, Morrison, & Bauer, 2014). Mice were deeply anesthetized, and hair

was completely removed from the hindquarters. A 27g percutaneous needle electrode was

inserted subcutaneously over the femur to stimulate the sciatic nerve at two locations: at the hip

joint and 1cm distal to this site. The stimulation reference electrode (26G percutaneous needle)

was inserted over the back. Bipolar EMG electrodes were inserted subcutaneously in the digital

interosseous muscles to record compound action potentials in response to activation of the sciatic

nerve. Stimulation and recording were conducted on both hindlimbs. The muscle response was

first tested with single pulses (100 μs duration) to determine the threshold and level of activation

producing a stable compound action potential in the muscle. The level of stimulation which

produced stable responses was then used for two series of stimulations with 2-20s long trains of

biphasic square wave pulses (100 μs duration) delivered at 1-5 Hz. The nerve was rested for 2

min. between runs. Stimulation was delivered via an isolated pulse stimulator (A-M Systems

Model 2100, A-M Systems Inc, Carlsborg, WA). EMG compound muscle action potential

(CMAP) responses were recorded with a differential AC amplifier (A-M Systems Model 1700;

gain 1k-10k; pass band 10-50 KHz). Data was sampled at 50 KHz using LabVIEW (National

Instruments Corporation, Austin, TX) running on a personal computer. Multiple recordings were

taken at each location to ensure a minimum of 10 representative responses for each location and

animal. Electrophysiological data were analyzed using Igor Pro (Wavemetrics Inc, Lake

Oswego, OR).

Data Analysis

In each experiment, data collection and analysis were performed identically, regardless of mouse

genotype. We used the sample (replicate) size (number) typically used in earlier studies of

peripheral myelination. Data are presented as mean +/- SD. Statistical analysis was done using

Student's unpaired t-test for two-group comparisons and analysis of variance (ANOVA) with

Tukey's test for multiple comparisons, according to the number of samples and the analysis of

mice at multiple ages. Normal distribution of all data was tested using Shapiro-Wilk normality

test. Sample sizes were similar to those conventional in the field and are indicated in the main

text, methods, or figure legends. A p-value of 0.05 or less was considered statistically significant.

ACKNOWLEDGMENTS

We thank Alan Tessler and members of the Son laboratory for critical reading of the manuscript.

We thank Dr. Eric Olson for *Yap1* and *Wwtr1* floxed mice, Drs. Ueli Suter and Kelly Monk for

Plp1-creERT2 mice. *Plp1*-creERT2 mice were generated by Dr. Ueli Suter using a patented Cre-

ERT2 construct developed by Dr. Pierre Chambon at GIE-CERBM.

COMPETING INTERESTS

The authors declare no competing interests.

FUNDINGS

This work was supported by grants from Shriners Children's (#84050 to Y-J.S.) and NIH NINDS

(NS105796 to Y-J.S.).

REFERENCES

Beirowski, B., Babetto, E., Golden, J. P., Chen, Y. J., Yang, K., Gross, R. W., . . . Milbrandt, J.
(2014). Metabolic regulator LKB1 is crucial for Schwann cell-mediated axon maintenance.
Nat Neurosci, 17(10), 1351-1361. doi:10.1038/nn.3809
Beverdam, A., Claxton, C., Zhang, X., James, G., Harvey, K. F., & Key, B. (2013). Yap controls
stem/progenitor cell proliferation in the mouse postnatal epidermis. *J Invest Dermatol,*
133(6), 1497-1505. doi:10.1038/jid.2012.430
Bremer, M., Frob, F., Kichko, T., Reeh, P., Tamm, E. R., Suter, U., & Wegner, M. (2011). Sox10
is required for Schwann-cell homeostasis and myelin maintenance in the adult peripheral
nerve. *Glia, 59*(7), 1022-1032. doi:10.1002/glia.21173
Clemence, A., Mirsky, R., & Jessen, K. R. (1989). Non-myelin-forming Schwann cells proliferate
rapidly during Wallerian degeneration in the rat sciatic nerve. *J Neurocytol, 18*(2), 185-
192.
Decker, L., Desmarquet-Trin-Dinh, C., Taillebourg, E., Ghislain, J., Vallat, J. M., & Charnay, P.
(2006). Peripheral myelin maintenance is a dynamic process requiring constant Krox20
expression. *J Neurosci, 26*(38), 9771-9779. doi:10.1523/JNEUROSCI.0716-06.2006

Deng, Y., Wu, L. M. N., Bai, S., Zhao, C., Wang, H., Wang, J., . . . Lu, Q. R. (2017). A reciprocal regulatory loop between TAZ/YAP and G-protein Galphas regulates Schwann cell proliferation and myelination. *Nat Commun, 8*, 15161. doi:10.1038/ncomms15161

Diepenbruck, M., Waldmeier, L., Ivanek, R., Berninger, P., Arnold, P., van Nimwegen, E., & Christofori, G. (2014). Tead2 expression levels control the subcellular distribution of Yap and Taz, zyxin expression and epithelial-mesenchymal transition. *J Cell Sci, 127*(Pt 7), 1523-1536. doi:10.1242/jcs.139865

Feltri, M. L., D'Antonio, M., Previtali, S., Fasolini, M., Messing, A., & Wrabetz, L. (1999). P0-Cre transgenic mice for inactivation of adhesion molecules in Schwann cells. *Ann N Y Acad Sci, 883*, 116-123.

Feltri, M. L., Poitelon, Y., & Previtali, S. C. (2016). How Schwann Cells Sort Axons: New Concepts. *Neuroscientist, 22*(3), 252-265. doi:10.1177/1073858415572361

Fernando, R. N., Cotter, L., Perrin-Tricaud, C., Berthelot, J., Bartolami, S., Pereira, J. A., . . . Tricaud, N. (2016). Optimal myelin elongation relies on YAP activation by axonal growth and inhibition by Crb3/Hippo pathway. *Nat Commun, 7*, 12186. doi:10.1038/ncomms12186

Finzsch, M., Schreiner, S., Kichko, T., Reeh, P., Tamm, E. R., Bosl, M. R., . . . Wegner, M. (2010). Sox10 is required for Schwann cell identity and progression beyond the immature Schwann cell stage. *J Cell Biol, 189*(4), 701-712. doi:10.1083/jcb.200912142

Gerber, D., Pereira, J. A., Gerber, J., Tan, G., Dimitrieva, S., Yánguez, E., & Suter, U. (2021). Transcriptional profiling of mouse peripheral nerves to the single-cell level to build a sciatic nerve ATlas (SNAT). *Elife, 10*. doi:10.7554/eLife.58591

Ghislain, J., & Charnay, P. (2006). Control of myelination in Schwann cells: a Krox20 cis-regulatory element integrates Oct6, Brn2 and Sox10 activities. *EMBO Rep, 7*(1), 52-58. doi:10.1038/sj.embor.7400573

Gomez-Sanchez, J. A., Lopez de Armentia, M., Lujan, R., Kessaris, N., Richardson, W. D., & Cabedo, H. (2009). Sustained axon-glial signaling induces Schwann cell hyperproliferation, Remak bundle myelination, and tumorigenesis. *J Neurosci, 29*(36), 11304-11315. doi:10.1523/jneurosci.1753-09.2009

Grove, M., Kim, H., Santerre, M., Krupka, A. J., Han, S. B., Zhai, J., . . . Son, Y. J. (2017). YAP/TAZ initiate and maintain Schwann cell myelination. *Elife, 6*. doi:10.7554/eLife.20982

Grove, M., Lee, H., Zhao, H., & Son, Y. J. (2020). Axon-dependent expression of YAP/TAZ mediates Schwann cell remyelination but not proliferation after nerve injury. *Elife, 9*. doi:10.7554/eLife.50138

Hao, Y., Chun, A., Cheung, K., Rashidi, B., & Yang, X. (2008). Tumor Suppressor LATS1 Is a Negative Regulator of Oncogene YAP*. *Journal of Biological Chemistry, 283*(9), 5496-5509. doi:https://doi.org/10.1074/jbc.M709037200

Harty, B. L., & Monk, K. R. (2017). Unwrapping the unappreciated: recent progress in Remak Schwann cell biology. *Curr Opin Neurobiol, 47*, 131-137. doi:10.1016/j.conb.2017.10.003

He, X., Zhang, L., Queme, L. F., Liu, X., Lu, A., Waclaw, R. R., . . . Lu, Q. R. (2018). A histone deacetylase 3-dependent pathway delimits peripheral myelin growth and functional regeneration. *Nat Med, 24*(3), 338-351. doi:10.1038/nm.4483

He, Y., Kim, J. Y., Dupree, J., Tewari, A., Melendez-Vasquez, C., Svaren, J., & Casaccia, P. (2010). Yy1 as a molecular link between neuregulin and transcriptional modulation of peripheral myelination. *Nat Neurosci, 13*(12), 1472-1480. doi:10.1038/nn.2686

Heinz, S., Romanoski, C. E., Benner, C., & Glass, C. K. (2015). The selection and function of cell type-specific enhancers. *Nat Rev Mol Cell Biol, 16*(3), 144-154. doi:10.1038/nrm3949

Hindley, C., & Philpott, A. (2012). Co-ordination of cell cycle and differentiation in the developing nervous system. *Biochem J, 444*(3), 375-382. doi:10.1042/BJ20112040

Holden, J. K., & Cunningham, C. N. (2018). Targeting the Hippo Pathway and Cancer through the TEAD Family of Transcription Factors. *Cancers (Basel), 10*(3). doi:10.3390/cancers10030081

Hu, J. K., Du, W., Shelton, S. J., Oldham, M. C., DiPersio, C. M., & Klein, O. D. (2017). An FAK-YAP-mTOR Signaling Axis Regulates Stem Cell-Based Tissue Renewal in Mice. *Cell Stem Cell, 21*(1), 91-106 e106. doi:10.1016/j.stem.2017.03.023

Jaegle, M., Ghazvini, M., Mandemakers, W., Piirsoo, M., Driegen, S., Levavasseur, F., . . . Meijer, D. (2003). The POU proteins Brn-2 and Oct-6 share important functions in Schwann cell development. *Genes Dev, 17*(11), 1380-1391. doi:10.1101/gad.258203

Jang, S. W., Srinivasan, R., Jones, E. A., Sun, G., Keles, S., Krueger, C., . . . Svaren, J. (2010). Locus-wide identification of Egr2/Krox20 regulatory targets in myelin genes. *J Neurochem, 115*(6), 1409-1420. doi:10.1111/j.1471-4159.2010.07045.x

Jang, S. W., & Svaren, J. (2009). Induction of myelin protein zero by early growth response 2 through upstream and intragenic elements. *J Biol Chem, 284*(30), 20111-20120. doi:10.1074/jbc.M109.022426

Jeanette, H., Marziali, L. N., Bhatia, U., Hellman, A., Herron, J., Kopec, A. M., . . . Belin, S. (2021). YAP and TAZ regulate Schwann cell proliferation and differentiation during peripheral nerve regeneration. *Glia, 69*(4), 1061-1074. doi:10.1002/glia.23949

Jessen, K. R., & Mirsky, R. (2005). The origin and development of glial cells in peripheral nerves. *Nat Rev Neurosci, 6*(9), 671-682.

Jessen, K. R., & Mirsky, R. (2016). The repair Schwann cell and its function in regenerating nerves. *J Physiol, 594*(13), 3521-3531. doi:10.1113/jp270874

Jones, E. A., Jang, S. W., Mager, G. M., Chang, L. W., Srinivasan, R., Gokey, N. G., . . . Svaren, J. (2007). Interactions of Sox10 and Egr2 in myelin gene regulation. *Neuron Glia Biol, 3*(4), 377-387. doi:10.1017/S1740925X08000173

Kao, S. C., Wu, H., Xie, J., Chang, C. P., Ranish, J. A., Graef, I. A., & Crabtree, G. R. (2009). Calcineurin/NFAT signaling is required for neuregulin-regulated Schwann cell differentiation. *Science, 323*(5914), 651-654.

Kim, J., Kwon, H., Shin, Y. K., Song, G., Lee, T., Kim, Y., . . . Jho, E. H. (2020). MAML1/2 promote YAP/TAZ nuclear localization and tumorigenesis. *Proc Natl Acad Sci U S A, 117*(24), 13529-13540. doi:10.1073/pnas.1917969117

Kim, M., Kim, T., Johnson, R. L., & Lim, D. S. (2015). Transcriptional co-repressor function of the hippo pathway transducers YAP and TAZ. *Cell Rep, 11*(2), 270-282. doi:10.1016/j.celrep.2015.03.015

Kim, M., Wende, H., Walcher, J., Kuhnemund, J., Cheret, C., Kempa, S., . . . Birchmeier, C. (2018). Maf links Neuregulin1 signaling to cholesterol synthesis in myelinating Schwann cells. *Genes Dev, 32*(9-10), 645-657. doi:10.1101/gad.310490.117

Kofler, M., Speight, P., Little, D., Di Ciano-Oliveira, C., Szaszi, K., & Kapus, A. (2018). Mediated nuclear import and export of TAZ and the underlying molecular requirements. *Nat Commun, 9*(1), 4966. doi:10.1038/s41467-018-07450-0

Le, N., Nagarajan, R., Wang, J. Y., Araki, T., Schmidt, R. E., & Milbrandt, J. (2005). Analysis of congenital hypomyelinating Egr2Lo/Lo nerves identifies Sox2 as an inhibitor of Schwann

cell differentiation and myelination. *Proc Natl Acad Sci U S A, 102*(7), 2596-2601.
doi:10.1073/pnas.0407836102
Le, N., Nagarajan, R., Wang, J. Y., Svaren, J., LaPash, C., Araki, T., . . . Milbrandt, J. (2005). Nab
proteins are essential for peripheral nervous system myelination. *Nat Neurosci, 8*(7), 932-
940. doi:10.1038/nn1490
LeBlanc, S. E., Jang, S. W., Ward, R. M., Wrabetz, L., & Svaren, J. (2006). Direct regulation of
myelin protein zero expression by the Egr2 transactivator. *J Biol Chem, 281*(9), 5453-5460.
doi:10.1074/jbc.M512159200
Leblanc, S. E., Srinivasan, R., Ferri, C., Mager, G. M., Gillian-Daniel, A. L., Wrabetz, L., &
Svaren, J. (2005). Regulation of cholesterol/lipid biosynthetic genes by Egr2/Krox20
during peripheral nerve myelination. *J Neurochem, 93*(3), 737-748. doi:10.1111/j.1471-
4159.2005.03056.x
Leone, D. P., Genoud, S., Atanasoski, S., Grausenburger, R., Berger, P., Metzger, D., . . . Suter,
U. (2003). Tamoxifen-inducible glia-specific Cre mice for somatic mutagenesis in
oligodendrocytes and Schwann cells. *Mol Cell Neurosci, 22*(4), 430-440.
Li, Z., Zhao, B., Wang, P., Chen, F., Dong, Z., Yang, H., . . . Xu, Y. (2010). Structural insights
into the YAP and TEAD complex. *Genes Dev, 24*(3), 235-240. doi:10.1101/gad.1865810
Liu, R., Jagannathan, R., Li, F., Lee, J., Balasubramanyam, N., Kim, B. S., . . . Moulik, M. (2019).
Tead1 is required for perinatal cardiomyocyte proliferation. *PLoS One, 14*(2), e0212017.
doi:10.1371/journal.pone.0212017
Lopez-Anido, C., Poitelon, Y., Gopinath, C., Moran, J. J., Ma, K. H., Law, W. D., . . . Svaren, J.
(2016). Tead1 regulates the expression of Peripheral Myelin Protein 22 during Schwann
cell development. *Hum Mol Genet*. doi:10.1093/hmg/ddw158
Lopez-Anido, C., Sun, G., Koenning, M., Srinivasan, R., Hung, H. A., Emery, B., . . . Svaren, J.
(2015). Differential Sox10 genomic occupancy in myelinating glia. *Glia*.
doi:10.1002/glia.22855
Montani, L., Pereira, J. A., Norrmen, C., Pohl, H. B. F., Tinelli, E., Trotzmuller, M., . . . Suter, U.
(2018). De novo fatty acid synthesis by Schwann cells is essential for peripheral nervous
system myelination. *J Cell Biol, 217*(4), 1353-1368. doi:10.1083/jcb.201706010
Monuki, E. S., Kuhn, R., & Lemke, G. (1993). Repression of the myelin P0 gene by the POU
transcription factor SCIP. *Mech Dev, 42*(1-2), 15-32. doi:10.1016/0925-4773(93)90095-f
Mukhtar, T., Breda, J., Grison, A., Karimaddini, Z., Grobecker, P., Iber, D., . . . Taylor, V. (2020).
Tead transcription factors differentially regulate cortical development. *Sci Rep, 10*(1), 4625.
doi:10.1038/s41598-020-61490-5
Norrmen, C., Figlia, G., Lebrun-Julien, F., Pereira, J. A., Trotzmuller, M., Kofeler, H. C., . . . Suter,
U. (2014). mTORC1 controls PNS myelination along the mTORC1-RXRgamma-SREBP-
lipid biosynthesis axis in Schwann cells. *Cell Rep, 9*(2), 646-660.
doi:10.1016/j.celrep.2014.09.001
Norrmén, C., Figlia, G., Pfistner, P., Pereira, J. A., Bachofner, S., & Suter, U. (2018). mTORC1
Is Transiently Reactivated in Injured Nerves to Promote c-Jun Elevation and Schwann Cell
Dedifferentiation. *J Neurosci, 38*(20), 4811-4828. doi:10.1523/jneurosci.3619-17.2018
Poitelon, Y., Lopez-Anido, C., Catignas, K., Berti, C., Palmisano, M., Williamson, C., . . . Feltri,
M. L. (2016). YAP and TAZ control peripheral myelination and the expression of laminin
receptors in Schwann cells. *Nat Neurosci, 19*(7), 879-887. doi:10.1038/nn.4316
Rasi, K., Hurskainen, M., Kallio, M., Staven, S., Sormunen, R., Heape, A. M., . . . Pihlajaniemi,
T. (2010). Lack of collagen XV impairs peripheral nerve maturation and, when combined

with laminin-411 deficiency, leads to basement membrane abnormalities and sensorimotor dysfunction. *J Neurosci, 30*(43), 14490-14501. doi:10.1523/JNEUROSCI.2644-10.2010

Ren, B., & Yue, F. (2015). Transcriptional Enhancers: Bridging the Genome and Phenome. *Cold Spring Harb Symp Quant Biol, 80*, 17-26. doi:10.1101/sqb.2015.80.027219

Ryu, E. J., Wang, J. Y., Le, N., Baloh, R. H., Gustin, J. A., Schmidt, R. E., & Milbrandt, J. (2007). Misexpression of Pou3f1 results in peripheral nerve hypomyelination and axonal loss. *J Neurosci, 27*(43), 11552-11559. doi:10.1523/JNEUROSCI.5497-06.2007

Saher, G., Quintes, S., Mobius, W., Wehr, M. C., Kramer-Albers, E. M., Brugger, B., & Nave, K. A. (2009). Cholesterol regulates the endoplasmic reticulum exit of the major membrane protein P0 required for peripheral myelin compaction. *J Neurosci, 29*(19), 6094-6104. doi:10.1523/JNEUROSCI.0686-09.2009

Salzer, J. L. (2015). Schwann cell myelination. *Cold Spring Harb Perspect Biol, 7*(8), a020529. doi:10.1101/cshperspect.a020529

Scherer, S. S., & Wrabetz, L. (2008). Molecular mechanisms of inherited demyelinating neuropathies. *Glia, 56*(14), 1578-1589. doi:10.1002/glia.20751

Schulz, A., Walther, C., Morrison, H., & Bauer, R. (2014). In vivo electrophysiological measurements on mouse sciatic nerves. *J Vis Exp*(86). doi:10.3791/51181

Sherman, D. L., Krols, M., Wu, L. M., Grove, M., Nave, K. A., Gangloff, Y. G., & Brophy, P. J. (2012). Arrest of myelination and reduced axon growth when Schwann cells lack mTOR. *J Neurosci, 32*(5), 1817-1825. doi:10.1523/JNEUROSCI.4814-11.2012

Son, Y. J., & Thompson, W. J. (1995). Schwann cell processes guide regeneration of peripheral axons. *Neuron, 14*(1), 125-132. doi:0896-6273(95)90246-5 [pii]

Taveggia, C., Zanazzi, G., Petrylak, A., Yano, H., Rosenbluth, J., Einheber, S., . . . Salzer, J. L. (2005). Neuregulin-1 type III determines the ensheathment fate of axons. *Neuron, 47*(5), 681-694. doi:10.1016/j.neuron.2005.08.017

Tian, W., Yu, J., Tomchick, D. R., Pan, D., & Luo, X. (2010). Structural and functional analysis of the YAP-binding domain of human TEAD2. *Proc Natl Acad Sci U S A, 107*(16), 7293-7298. doi:10.1073/pnas.1000293107

Topilko, P., Schneider-Maunoury, S., Levi, G., Baron-Van Evercooren, A., Chennoufi, A. B., Seitanidou, T., . . . Charnay, P. (1994). Krox-20 controls myelination in the peripheral nervous system. *Nature, 371*(6500), 796-799. doi:10.1038/371796a0

Tricaud, N., & Park, H. T. (2017). Wallerian demyelination: chronicle of a cellular cataclysm. *Cell Mol Life Sci, 74*(22), 4049-4057. doi:10.1007/s00018-017-2565-2

Verheijen, M. H., Camargo, N., Verdier, V., Nadra, K., de Preux Charles, A. S., Medard, J. J., . . . Smit, A. B. (2009). SCAP is required for timely and proper myelin membrane synthesis. *Proc Natl Acad Sci U S A, 106*(50), 21383-21388. doi:10.1073/pnas.0905633106

Wen, T., Liu, J., He, X., Dong, K., Hu, G., Yu, L., . . . Zhou, J. (2019). Transcription factor TEAD1 is essential for vascular development by promoting vascular smooth muscle differentiation. *Cell Death Differ, 26*(12), 2790-2806. doi:10.1038/s41418-019-0335-4

Wen, T., Yin, Q., Yu, L., Hu, G., Liu, J., Zhang, W., . . . Zhou, J. (2017). Characterization of mice carrying a conditional TEAD1 allele. *Genesis, 55*(12). doi:10.1002/dvg.23085

Wright, M. C., Cho, W. J., & Son, Y. J. (2007). Distinct patterns of motor nerve terminal sprouting induced by ciliary neurotrophic factor vs. botulinum toxin. *J Comp Neurol, 504*(1), 1-16. doi:10.1002/cne.21439

Wu, S., Liu, Y., Zheng, Y., Dong, J., & Pan, D. (2008). The TEAD/TEF family protein Scalloped
mediates transcriptional output of the Hippo growth-regulatory pathway. *Dev Cell, 14*(3),
388-398. doi:10.1016/j.devcel.2008.01.007
Yao, F., Zhou, Z., Kim, J., Hang, Q., Xiao, Z., Ton, B. N., . . . Ma, L. (2018). SKP2- and OTUD1-
regulated non-proteolytic ubiquitination of YAP promotes YAP nuclear localization and
activity. *Nat Commun, 9*(1), 2269. doi:10.1038/s41467-018-04620-y
Zhao, B., Wei, X., Li, W., Udan, R. S., Yang, Q., Kim, J., . . . Guan, K. L. (2007). Inactivation of
YAP oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue
growth control. *Genes Dev, 21*(21), 2747-2761. doi:10.1101/gad.1602907
Zhao, B., Ye, X., Yu, J., Li, L., Li, W., Li, S., . . . Guan, K. L. (2008). TEAD mediates YAP-
dependent gene induction and growth control. *Genes Dev, 22*(14), 1962-1971.
doi:10.1101/gad.1664408
Zheng, Y., & Pan, D. (2019). The Hippo Signaling Pathway in Development and Disease. *Dev
Cell, 50*(3), 264-282. doi:10.1016/j.devcel.2019.06.003
Zorick, T. S., Syroid, D. E., Brown, A., Gridley, T., & Lemke, G. (1999). Krox-20 controls SCIP
expression, cell cycle exit and susceptibility to apoptosis in developing myelinating
Schwann cells. *Development, 126*(7), 1397-1406.

FIGURE LEGENDS

Figure 1. Multiple TEADs are expressed in Schwann cells of WT and *Tead1* cKO mice

(A) Immunostaining and quantification of TEAD1+ SCs on longitudinal sections of P22 WT and

Tead1 cKO SNs. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. n = 3

mice per genotype, ****P < 0.0001, Student's unpaired t-test. Scale bar: 20μm

(B) Western blotting of P50 WT and *Tead1* cKO SN lysates with anti-TEAD1, -TEAD2, -

TEAD3, -TEAD4, and -pan TEAD. TEAD expression is normalized to that of *β-actin* as an

internal control, and WT expression is arbitrarily given the value 1. n = 3 mice per genotype. ns,

p > 0.05; *p ≤ 0.05, **p ≤ 0.005, ***p ≤ 0.001, Student's unpaired t-test.

(C) Western blotting of P50 WT and *Tead1* cKO SN lysates with anti-YAP, -phospho-YAPs

(Ser127 and Ser397), and -TAZ. YAP/TAZ expression is normalized to that of *β-actin* as an

internal control, and WT expression is arbitrarily given the value 1. n = 3 mice per genotype. *p

≤ 0.05, ***p ≤ 0.001, ****P < 0.0001, Student's unpaired t-test.

(D) Immunostaining of YAP/TAZ+ SCs on longitudinal sections of P50 WT and *Tead1* cKO

SNs. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. n = 3 mice per

genotype, ns, p > 0.05, Student's unpaired t-test. Scale bar: 20μm

The following figure supplements are available for Figure 1.

Figure 1-figure supplement 1

Western blotting of P5, P38, P85 WT and P38 *Tead1* cKO SN lysates with anti-TEAD2

TEAD2 expression is normalized to that of *β-actin* as an internal control. Arrow denotes ~65 kD

TEAD2 band that was diminished with age.

Source data 1

This zip archive contains source files for graphs in figure 1A, 1B, 1C, 1D and uncropped labeled

or unlabeled blots of figure 1B and 1D.

Source data 2

This zip archive contains uncropped labeled or unlabeled blots in **Figure 1-figure supplement 1.**

Figure 2. TEAD1 is required for myelin sheath formation and growth

(A) A living control (WT) and two *Tead1* cKO mice from one litter at P55. Arrows point to splayed paralyzed hindlimbs.

(B) Exposed SNs of WT and *Tead1* cKO mice during dissection. Scale bar: 1mm

(C) Representative images of CMAPs generated by stimulation of WT and *Tead1* cKO SNs.

(D) Quantification of NCV in WT and *Tead1* cKO. n = 6 mice per genotype. ***P < 0.001, Student's unpaired t-test.

(E) Representative images of semithin (left panels) and TEM (right panels) of transverse sections of WT and *Tead1* cKO SNs at P3. Scale bar: 20μm (left panels), 1μm (right panels)

(F) Representative images of semithin (left panels) and TEM (right panels) of transverse sections

of WT and *Tead1* cKO SNs at P22. Scale bar: 20μm (left panels), 1μm (right panels)

(G) Representative images of semithin (left panels) and TEM (right panels) of transverse

sections of WT and *Tead1* cKO SNs at P50. Scale bar: 20μm (left panels), 1μm (right panels)

(H) Quantification of G-ratio for axons in WT and *Tead1* cKO SNs at P22 and at P50. n = 3

mice per group. ns, p > 0.05 (P22 WT versus P50 WT, P22 cKO versus P50 cKO), ****P <

0.0001 (P22 WT versus P22 cKO, P50 WT versus P50 cKO).

(I) Scatter plot graph displaying G-ratio in relation to axon diameter in WT and *Tead1* cKO SNs

at P22. n = 1063 axons from 3 mice per each genotype.

(J) Scatter plot graph displaying G-ratio in relation to axon diameter in WT and *Tead1* cKO SNs

at P50. n = 880 axons from 3 mice per each genotype.

(K) Quantitative comparison of mean axon diameter in WT and *Tead1* cKO SNs at P22 and at

P50. n = 3 mice per group. ns, p > 0.05, *p ≤ 0.05, 2-way ANOVA with Tukey's multiple

comparisons test.

(L) Representative TEM images of transverse sections of P50 WT and *Tead1* cKO SNs. Right

panels are enlarged images of the boxed area in left panels. Scale bar: 1μm

(M) Quantification of normal and abnormal C-fibers in P50 WT and *Tead1* cKO SNs. C-fibers

normally ensheathed by SC cytoplasm are marked as 'ensheathed', whereas those not ensheathed

are denoted as 'naked'. Naked C-fibers located in a myelinated bundle are denoted as 'naked

myelinated'. Relative percentage of each type of C-fiber is plotted. N > 500 C-fibers from 3 mice

per each genotype. *p ≤ 0.05, ***p ≤ 0.001, ****P < 0.0001, 2-way ANOVA Tukey's multiple

comparisons test.

The following figure supplements are available for Figure 2.

Figure 2-figure supplement 1

Additional semi-thin and TEM images of WT and *Tead1* cKO

(A) Representative images of semi-thin sections of various areas of WT and Tead1 cKO SNs at

P22 and P50. Scale bar: 20μm

(B) Representative TEM images of transverse sections of WT and Tead1 cKO SNs at P22 and

P50. Scale bar: 1μm (P22 panels), 2μm (P50 panels)

Source data 1

This zip archive contains source files for graphs in Figure 2D, 2H, 2I, 2J, 2K, 2M.

Figure 3. TEAD1 regulates SC proliferation both positively and negatively

(A) Immunostaining and quantification of SCs on transverse sections of WT and *Tead1* cKO

SNs at P3, P22, and P52. SC nuclei and all cell nuclei are marked by Sox10 and DAPI. Axons

are marked by Neurofilaments (NF). n = 3 mice per genotype, ns, p > 0.05, *p ≤ 0.05, **p ≤

0.01, ****P < 0.0001, 2-way ANOVA with Tukey's multiple comparisons test. Scale bar: 30μm

(B) Immunostaining and quantification of Ki67+ SCs on longitudinal sections of WT and *Tead1*

cKO SNs at P3, P11, and P22. SC nuclei and all cell nuclei are marked by Sox10 and DAPI. n =

3 mice per genotype, ns, p > 0.05, **p ≤ 0.01, ****P < 0.0001, 2-way ANOVA with Tukey's

multiple comparisons test. Scale bar: 30μm (P3 panels), 20μm (P11 and P22 panels)

(C) Immunostaining and quantification of EdU+ SCs on longitudinal sections of WT and *Tead1*

cKO SNs at P3, P11, and P22. SC nuclei and all cell nuclei are marked by Sox10 and DAPI. n =

3 mice per genotype, ns, p > 0.05, *p ≤ 0.05, ****P < 0.0001, 2-way ANOVA with Tukey's

multiple comparisons test. Scale bar: 30μm (P3 panels), 20μm (P11 and P22 panels)

(D) Immunostaining and quantification of EdU+/TEAD1+ SCs on longitudinal sections of WT

and *Tead1* cKO SNs at P11. SC nuclei are marked by Sox10. Asterisks denote *Tead1* cKO SCs

in which TEAD1 is not deleted and which are not EdU+. Arrowheads denote EdU+ SCs which

are TEAD1+ in WT but TEAD1- in *Tead1* cKO. n = 3 mice per genotype, ***P < 0.001,

Student's unpaired t-test. Scale bar: 20μm

1006

1007   The following figure supplements are available for Figure 3.

1008

1009   **Source data 1**

1010   This zip archive contains source files for graphs in Figure 3A, 3B, 3C, 3D.

1011

1012

1013

Figure 4. TEAD1 is required for Krox20 to upregulate myelin proteins during

development

(A) Western blotting of P50 WT and *Tead1* cKO SN lysates with anti-Oct6, -Krox20, -MPZ, and

-MBP. Protein expression is normalized to that of *β-actin* as an internal control, and WT

expression is arbitrarily given the value 1. n = 3 mice per genotype. *p ≤ 0.05, ***p ≤ 0.001,

***p ≤ 0.0001, Student's unpaired t-test.

(B) Immunostaining and quantification of Oct6+ SCs on longitudinal sections of P50 WT and

Tead1 cKO SNs. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. n = 3

mice per genotype, ****P < 0.0001, Student's unpaired t-test. Scale bar: 20μm

(C) Immunostaining and quantification of Krox20+ SCs on longitudinal sections of WT and

Tead1 cKO SNs at P11 and P50. SC nuclei are identified by Sox10. All cell nuclei are marked by

DAPI. n = 3 mice per each group, ns, p > 0.05, Student's unpaired t-test. Scale bar: 20μm

(D) Immunostaining and quantification of EdU+/Krox20+ SCs on longitudinal sections of WT

and *Tead1* cKO SNs at P11. SC nuclei are identified by Sox10. Arrowheads denote examples of

EdU+ SCs which are Krox20-. Asterisks denote rarely observed EdU+ SCs which are Krox20+.

n = 3 mice per genotype, ns, p > 0.05, Student's unpaired t-test. Scale bar: 20μm

1030

1031 The following figure supplements are available for Figure 4.
1032

1033 **Source data 1**

1034 This zip archive contains source files for graphs in Figure 4A, 4B, 4C, 4D and uncropped labeled

1035 or unlabeled blots of Figure 4A.

1036

Figure 5. TEAD1 and YAP/TAZ upregulate cholesterol synthesis enzymes: FDPS and IDI1

(A) Western blotting of P8 and P40 WT and *Tead1* cKO SN lysates with anti-SREBP1, -SCD1, -SREBP2, -HMGCR, -FDPS, and -IDI1. Protein expression is normalized to that of *β-actin* as an internal control, and WT expression is arbitrarily given the value 1. n = 3 or 4 mice per genotype. ns, p > 0.05, *p ≤ 0.05, **p ≤ 0.01, ****P < 0.0001, Student's unpaired t-test.

1044 **(B)** Western blotting of P60 WT and *Yap1/Wwtr1* cDKO SN lysates with anti-YAP, -TAZ, -

1045 SREBP1, -SCD1, -SREBP2, -HMGCR, -FDPS, and -IDI1. Protein expression is normalized to

1046 that of *β-actin* as an internal control, and WT expression is arbitrarily given the value 1. n = 3 or

1047 4 mice per genotype. ns, p > 0.05, *p ≤ 0.05, **p ≤ 0.01, ***P < 0.001, Student's unpaired t-test.

1048

1049 The following figure supplements are available for Figure 5.

1050

1051 **Source data 1**

1052 This zip archive contains uncropped labeled or unlabeled blots of P8 WT and Tead1 cKO SNs in

1053 Figure 5A.

1054

1055 **Source data 2**

1056 This zip archive contains source files for graphs in Figure 5A and 5B, and uncropped labeled or

1057 unlabeled blots of P40 WT and Tead1 cKO SNs in Figure 5A, and P60 SNs of WT and

1058 YAP/TAZ cDKO in Figure 5B.

1059

1060

1061

Figure 6. TEAD1 is largely dispensable for myelin maintenance, but required for Remak

bundle integrity

(A) Cartoon showing timeline of tamoxifen (TAM) injection and time of sacrifice of *Tead1* iKO

mice.

(B) Representative TEM images of P150 WT and *Tead1* iKO SNs showing occasional large

axons undergoing demyelination (arrowheads) or completely demyelinated (arrows). Scale bar:

6μm (left panels), 2μm (right panel)

(C) Quantification of NCV in P90-P150 WT and *Tead1* iKO mice. n = 5 or 6 mice per genotype.

ns, $p > 0.05$, Student's unpaired t-test.

(D) Representative TEM images of P97 WT and *Tead1* iKO SNs showing Remak bundles.

Asterisk denotes a bundle of thinly myelinated C-fibers. Arrow denotes demyelinated axon.

Scale bar: 1μm

(E) Quantification of normal and abnormal C-fibers in P150 WT and *Tead1* iKO SNs. Relative

percentage of each type of C-fiber is plotted. N > 500 C-fibers from 3 mice per each genotype.

ns, $p > 0.05$, ****$P < 0.0001$, 2-way ANOVA with Tukey's multiple comparisons test.

1077

1078 The following figure supplements are available for Figure 6.
1079

1080

1081

Figure 6-figure supplement 1

Expression of TEAD1, pan-TEAD and YAP/TAZ in intact and regenerating nerves of WT and

Tead1 **iKO mice**

(A) Western blotting of adult WT and *Tead1* iKO SN lysates with anti-TEAD1. TEAD1

expression is normalized to that of *β-actin* as an internal control, and WT expression is

arbitrarily given the value 1. n = 3 mice per genotype. **p ≤ 0.01, Student's unpaired t-test.

(B) Immunostaining and quantification of TEAD1+ SCs on longitudinal sections of P100 WT

and *Tead1* iKO SNs 3 months after tamoxifen administration. SC nuclei are identified by Sox10.

All cell nuclei are marked by DAPI. n = 3 mice per genotype, ****$P < 0.0001$, Student's

unpaired t-test. Scale bar: 20μm

(C) Immunostaining and quantification of pan-TEAD+ SCs on longitudinal sections of P100 WT

and *Tead1* iKO SNs. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. n =

3 mice per genotype. ns, $p > 0.05$, Student's unpaired t-test. Scale bar: 20μm

(D) Immunostaining and quantification of TEAD1+ SCs on longitudinal sections of WT and

Tead1 iKO SNs 28 days after nerve crush. SC nuclei are identified by Sox10. All cell nuclei are

marked by DAPI. n = 3 mice per genotype, ****$P < 0.0001$, Student's unpaired t-test. Scale bar:

20μm

(E) Immunostaining and quantification of pan-TEAD+ SCs on longitudinal sections of P100 WT

and *Tead1* iKO SNs 28 days after nerve crush. SC nuclei are identified by Sox10. All cell nuclei

are marked by DAPI. n = 3 mice per genotype. ns, $p > 0.05$, Student's unpaired t-test. Scale bar:

20μm

(F) Immunostaining of YAP/TAZ+ SCs on longitudinal sections of P100 WT and *Tead1* iKO

SNs. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. n = 3 mice per

group. ns, $p > 0.05$, Student's unpaired t-test. Scale bar: 20μm

(G) Immunostaining of YAP/TAZ+ SCs on longitudinal sections of WT and *Tead1* iKO SNs 28

days after nerve crush. SC nuclei are identified by Sox10. All cell nuclei are marked by DAPI. n

= 3 mice per group. ns, $p > 0.05$, Student's unpaired t-test. Scale bar: 20μm

1109

1110 **Source data 1**

1111 This zip archive contains source files for graphs in Figure 6C and 6E.

1112

1113 **Source data 2**

1114 This zip archive contains source files for graphs in Figure 6A-6G and uncropped labeled or

1115 unlabeled blots of Figure 6A.

1116

1117

1118

1119

1120

Figure 7. TEAD1 is dispensable for YAP/TAZ to maintain Krox20, FDP and IDI1 expression

(A) Immunostaining and quantification of Krox20+ SCs on longitudinal sections of WT and *Yap1/Wwtr1* iDKO SNs 2 weeks after 1ˢᵗ tamoxifen injection. SC nuclei are marked by Sox10. All cell nuclei are marked by DAPI. n = 3 mice per genotype, ***P < 0.001, Student's unpaired t-test. Scale bar: 20μm

(B) Immunostaining and quantification of Krox20+ SCs on longitudinal sections of WT and *Tead1* iKO SNs 6 weeks after 1ˢᵗ tamoxifen injection. SC nuclei are marked by Sox10. All cell nuclei are marked by DAPI. n = 3 mice per genotype, ns, p > 0.05, Student's unpaired t-test. Scale bar: 20μm

(C) Immunostaining and quantification of Oct6+ SCs on longitudinal sections of WT and *Tead1* iKO SNs 6 weeks after 1ˢᵗ tamoxifen injection. SC nuclei are marked by Sox10. All cell nuclei are marked by DAPI. n = 3 mice per genotype, ns, p > 0.05, Student's unpaired t-test. Scale bar: 20μm

(D) Western blotting of WT and *Yap1/Wwtr1* iDKO SN lysates with anti-YAP, -TAZ, -SREBP1,

-SCD1, -SREBP2, -HMGCR, -FDPS, and -IDI1. Protein expression is normalized to that of *β-*

actin as an internal control, and WT expression is arbitrarily given the value 1. n = 3 or 4 mice

per genotype. ns, $p > 0.05$, **$p \leq 0.01$, ***$P < 0.001$, ****$P < 0.0001$, Student's unpaired t-test.

(E) Western blotting of WT and *Tead1* iKO SN lysates with anti-SREBP1, -SCD1, -SREBP2, -

HMGCR, -FDPS, and -IDI1. Protein expression is normalized to that of *β-actin* as an internal

control, and WT expression is arbitrarily given the value 1. n = 3 or 4 mice per genotype. ns, $p >$

0.05, Student's unpaired t-test.

1143

1144 The following figure supplements are available for Figure 7.

1145

1146 **Source data 1**

1147 This zip archive contains source files for graphs in Figure 7A-7E and uncropped labeled or

1148 unlabeled blots of Figure 7D and 7E.

1149

Figure 8. TEAD1 is required for functional regeneration of peripheral nerve

(A) Cartoon showing timeline of tamoxifen (TAM) injection and time of sacrifice of *Tead1* iKO mice after nerve crush.

(B) Low magnification images of longitudinal sections of ~4mm long SN segments of WT and *Tead1* iKO mice 28 days after nerve crush. Arrowheads denote crush site. Axons and myelin are marked by Tuj1 and MBP, respectively. All cell nuclei are marked by DAPI. Yellow asterisks denote the distal nerve segment of *Tead1* iKO, which almost completely lacks myelin. White asterisks denote distal segments of both WT and *Tead1* iKO, which exhibit more cells than proximal segments due to SC proliferation after injury. Scale bar: 500μm

(C) Representative low and high magnification NMJ images of WT and Tead1 iKO 28 days after nerve crush. Axon and axon terminals are marked by neurofilament (NF) and SV2 staining. SCs

are marked by S100. Muscle acetylcholine receptors are marked by α-Bungarotoxin. Arrows

denote examples of axon terminals that grew past NMJs in iKO. Arrowheads denote examples of

extrasynaptic AChR clusters in iKO. Scale bar: 50μm (low magnification panels), 20μm (high

magnification panels)

(D) Representative semi-thin and TEM images of transverse sections of WT and *Tead1* iKO 28

days after nerve crush. Scale bar: 20μm (left semi-thin panels), 2μm (right TEM panels)

(E) Representative CMAP recordings and NCV measurements of WT and *Tead1* iKO at 28 days

after nerve crush. n = 3 mice per genotype, **$p \leq 0.01$, Student's unpaired t-test.

(F) Immunostaining and quantification of EdU+ SCs on longitudinal sections of WT and *Tead1*

iKO SNs at 28 days after nerve crush. SC nuclei are marked by Sox10. All cell nuclei are marked

by DAPI. n = 4-5 mice per genotype, ns, $p > 0.05$, Student's unpaired t-test. Scale bar: 20μm

(G) Immunostaining and quantification of Krox20+ SCs on longitudinal sections of WT and

Tead1 iKO SNs at 28 days after nerve crush. SC nuclei are marked by Sox10. All cell nuclei are

marked by DAPI. n = 3 mice per genotype, ns, $p > 0.05$, Student's unpaired t-test. Scale bar:

20μm

(H) Immunostaining and quantification of Oct6+ SCs on longitudinal sections of WT and *Tead1*

iKO SNs at 28 days after nerve crush. SC nuclei are marked by Sox10. All cell nuclei are marked

by DAPI. n = 3 mice per genotype, ns, ***$P < 0.001$, Student's unpaired t-test. Scale bar: 20μm

The following figure supplements are available for Figure 8.

Source data 1

This zip archive contains source files for graphs in Figure 8E, 8F, 8G, 8H.

Video 1. A movie showing a P55 WT and two littermates *Tead1* cKO mic