Induction of viral mimicry upon loss of DHX9 and ADAR1 in breast cancer cells

Kyle A. Cottrell [1,4,5‡*], Sua Ryu [1,4‡], Luisangely Soto Torres [1,4], Angela M. Schab [1,4], Jason D.

Weber [1,2,3,4*]

[1]Department of Medicine, Division of Molecular Oncology, [2]Department of Cell Biology and

Physiology, and [3]Department of Biology, Siteman Cancer Center, [4]ICCE Institute, Washington

University School of Medicine, Saint Louis, Missouri, USA

[5]Department of Biochemistry, Purdue University, West Lafayette, IN, USA

[‡]Authors contributed equally

*Co-Corresponding authors

The authors declare no potential conflicts of interest.

Correspondence:
Kyle A Cottrell, Ph.D.
Department of Biochemistry
Purdue University
201 S University St.
West Lafayette, IN
Email: cottrellka@pudue.edu
Telephone: 765-494-6941
19
Jason D. Weber, Ph.D.
Department of Medicine
Division of Molecular Oncology
Washington University School of Medicine
660 South Euclid Avenue
Campus Box 8069
St. Louis, MO 63110 USA
Email: jweber@wustl.edu
Telephone: 314-747-3896
Fax: 314-362-0152
30

Summary

Detection of viral double-stranded RNA (dsRNA) is an important component of innate immunity.

However, many endogenous RNAs containing double-stranded regions can be misrecognized

and activate innate immunity. The interferon inducible ADAR1-p150 suppresses dsRNA

sensing, an essential function for ADAR1 in many cancers, including breast. Although ADAR1-

p150 has been well established in this role, the functions of the constitutively expressed

ADAR1-p110 isoform are less understood. We used proximity labeling to identify putative

ADAR1-p110 interacting proteins in breast cancer cell lines. Of the proteins identified, the RNA

helicase DHX9 was of particular interest. Knockdown of DHX9 in ADAR1-dependent cell lines

caused cell death and activation of the dsRNA sensor PKR. In ADAR1-independent cell lines,

combined knockdown of DHX9 and ADAR1, but neither alone, caused activation of multiple

dsRNA sensing pathways leading to a viral mimicry phenotype. Together, these results reveal

an important role for DHX9 in suppressing dsRNA sensing by multiple pathways.

46
47

Introduction

RNA editing enhances protein diversity and modulates multiple aspects of RNA metabolism[1–3]. A-to-I editing is carried out by adenosine deaminase acting on RNA 1 (ADAR1), an RNA editase that binds double-stranded RNA (dsRNA) and converts adenosines to inosines[4,5]. The main domains of ADAR1 include the Z-DNA binding domains (ZBD), the double-stranded RNA (dsRNA) binding domains (dsRBD), and the deaminase domain[6,7]. There are two isoforms, p110 and p150, produced by alternative transcriptional start sites[8]. They share the same deaminase domain, dsRBDs, and a ZBD, but exhibit a distinct subcellular localization[9,10]. ADAR1-p150 is predominantly cytoplasmic, whereas ADAR1-p110 is nuclear-localized[9,11].

An important role for ADAR1 is to suppress dsRNA sensing[12–15]. Many endogenously encoded RNAs can form large double-stranded regions, often through base-pairing between inverted Alu elements[16]. ADAR1 edits the majority of human genes, with most editing occurring within inverted Alu repeats[3,4]. By binding, editing, and thus altering the structure of dsRNA, ADAR1 suppresses the detection of dsRNA by various cytoplasmic sensors such as MDA5, RIG-I, and PKR[12,13,15,17,18]. These RNA sensors are part of innate immunity against viral infections[16,19–21]. Thus, ADAR1 prevents the activation of innate immunity pathways by endogenous immunogenic RNAs[16,20,22]. Some mutations of ADAR1 in humans cause inappropriate dsRNA sensing and activation of the type I IFN pathway which manifests as the interferonopathy Aicardi–Goutières syndrome[23].

The cell-intrinsic antiviral response against foreign dsRNA - or misrecognized endogenous dsRNA - involves multiple pathways: (1) Recognition of dsRNA by MDA5 or RIG-I results in the activation of type I interferon (IFN-I) signaling[16,20,22]. (2) Activation of the dsRNA-binding kinase PKR triggers translational shutdown by phosphorylation of the translation initiation factor eIF2α[12]. (3) Detection of dsRNA by OAS proteins activates RNase L, which

carries out non-specific cleavage of RNA and triggers cell death[24]. There is significant crosstalk

between the three pathways, and the aforementioned dsRNA sensors are transcriptionally

controlled by IFN-I, known as interferon-stimulated genes (ISGs)[25,26]. The ADAR1-p150 isoform

is itself an ISG and is the isoform responsible for suppressing the activation of dsRNA

sensors[8,10,14,27]. ADAR1-p110, however, is constitutively expressed, though its functions are less

established[28,29].

Because IFN-I signaling is often cytotoxic and antiproliferative, ADAR1's ability to

suppress IFN-I signaling was shown to exert pro-tumor effects[22,30,31]. As such, ADAR1 has been

proposed as a potential therapeutic target for various cancers, including breast cancer[30,32–34].

ADAR1 mRNA expression is elevated in breast cancer and is correlated with a poor

prognosis[30,35]. Furthermore, ADAR1 is essential for a subset of breast cancer cells

overrepresented by triple-negative breast cancer (TNBC)[35]. Known as 'ADAR1-dependency,'

depletion of ADAR1-p150 leads to IFN-I signaling and global translational repression in cells

that are sensitive to depletion of ADAR1[34,35]. What makes cancer cells sensitive or refractory to

ADAR1 loss is not yet determined[35].

ADAR1 interacts with numerous RNA binding proteins including RNA helicases,

transcription machinery, and DNA repair proteins[29,36–39]. The influence of ADAR1-interacting

proteins on A-to-I editing has previously been reported[37,40–43]. In this study, we evaluated

components of the ADAR1 interactome in breast cancer cells and identified the RNA helicase

DHX9 as a redundant suppressor of immunogenic dsRNA in ADAR1-independent breast cancer

cells. We demonstrate that co-depletion of ADAR1 and DHX9 is sufficient to trigger a viral

mimicry phenotype in ADAR1-independent cells.

Results

Proximity Labeling by APEX2 Reveals ADAR1 Interacting Proteins

To better understand the role of ADAR1-p110 in breast cancer, we turned to a proximity

labeling approach using APEX2 to identify putative ADAR1-p110-interacting proteins. Proximity

labeling by APEX2 allows for the identification of proteins within 20 nm of an APEX2 fusion

protein via biotin-mediated pulldown[44]. An APEX2-ADAR1-p110 fusion construct or APEX2

alone was expressed in MDA-MB-231, MCF-7, and SK-BR-3 (Figure 1a). Following proximity

labeling, biotinylated proteins were purified by streptavidin pulldown and subsequently identified

by mass spectrometry (Fig 1b-c, Figure S1a, Supplementary Table 2). In total, we identified

over one hundred enriched proteins across the three cell lines, with many identified in all three

lines (Figure 1d). Over-representation analysis of gene ontology (GO) terms revealed that many

of the proteins identified by proximity labeling have roles in multiple aspects of RNA metabolism

and localize to the nucleus and nucleolus (Figure 1e, Supplementary Table 3). This finding was

supported by a comparison to proteins previously observed to localize to the nucleus and

nucleolus within MCF-7 in the SubCellBarcode dataset[45] (Figure 1f and 1g). These findings are

largely consistent with the localization of ADAR1 within the cell lines studied.

Immunofluorescence for ADAR1, which largely reflects the localization of ADAR1-p110 as the

predominant isoform, showed that ADAR1 was localized in both the nucleus and nucleolus

(Figure 1h).

Validation of Protein Interactions Identified by Proximity Labeling

Although proximity labeling by APEX2 is a powerful technique for identifying putative

protein-protein interactions, it does not distinguish between interactions and close

associations[44]. To validate the proximity labeling findings and provide supporting evidence for

direct protein-protein interactions, we performed co-immunoprecipitation followed by

immunoblotting for five proteins identified by proximity labeling. ADAR1 could be

immunoprecipitated by antibodies against the helicases DHX9, DDX17, and DDX54 (Figure 2a-

g). ADAR1 could also be immunoprecipitated with antibodies against XRN2 and PARP (Figure

S1b-c). To assess whether these potential interactions depended on RNA, we treated the

lysates with RNase A to degrade RNA prior to immunoprecipitation (Figure S1d).

Immunoprecipitation of ADAR1 by antibodies against DHX9, DDX17, and DDX54 was possible

even in the presence of RNase A and improved in some cases (Figure 2a-g). For DDX17 and

DDX54, RNase A treatment did not change the co-immunoprecipitation results. However, DHX9

immunoprecipitated with both isoforms of ADAR1 in the absence of RNase A, but only with

ADAR1-p110 when treated with RNase A. These findings suggest that DHX9, DDX17, and

DDX54 directly interact with ADAR1-p110, and DHX9 interacts with ADAR1-p150 through an

RNA bridge/scaffold. The co-immunoprecipitation findings are also consistent with the

localization of the proteins studied. Much like ADAR1, the helicases DHX9, DDX17, and DDX54

localized to the nucleus or nucleolus (Figure 2h-j). Interestingly, in the cell lines used in this

study, ADAR1-p150 is localized to the nucleus and cytosol (Figure S1e), consistent with

previous reports of ADAR1-p150 shuttling between these two compartments[46,47]. Together,

these findings validate the results of the proximity labeling described in Figure 1 and provide

evidence for direct interactions between ADAR1 and multiple helicases.

DHX9 is overexpressed in breast cancer

Of the identified helicases, DHX9 was of particular interest for several reasons. First,

DHX9 is the only helicase in humans that has a dsRBD. The dsRBD that is present in DHX9 is

of the same class present in ADAR1 and PKR (Figure 3a). Analysis of publicly available RNA-

seq data for human cell lines and tumors revealed that DHX9 expression closely correlates with

ADAR1 expression (Figure 3b-d, Figure S2a-c). The correlation is stronger between DHX9 and

the transcript encoding ADAR1-p110, than between DHX9 and ADAR1-p150 (Figure 3c-d,

Figure S2d-e). Furthermore, the expression of DHX9 and ADAR1 correlates better than any

other helicase identified by proximity labeling (Figure 3b, Figure S2b). Consistent with this

correlated expression, and much like ADAR1, DHX9 is highly expressed in breast cancer and is

correlated with a poor prognosis[35] (Figure 3c, Figure S2f-h).

DHX9 is essential in TNBC cell lines and suppresses PKR activation

The similarities between ADAR1 and DHX9 led us to further study the role of DHX9 in

breast cancer. Analysis of publicly available data from DepMap

(https://depmap.org/portal/download/custom/) revealed that DHX9 is commonly essential in

breast cancer cell lines (Figure 3f). We validated this finding by knocking down DHX9 in four

TNBC cell lines previously shown to be ADAR1-dependent[35]. In all four lines, knockdown of

DHX9 caused cell death, likely through caspase-mediated apoptosis as indicated by cleaved

PARP (Figure 3g and h). Given the presence of the common dsRBD in DHX9 and PKR, and the

role of DHX9 in regulating the abundance of dsRNA[48], we asked whether DHX9 could influence

activation of PKR. To our surprise, we found that in three of the TNBC cell lines studied,

knockdown of DHX9 caused activation of PKR (Figure 3g and 3i), this finding was confirmed for

one cell line by CRISPR-Cas9 knockout of DHX9 (Figure S3i-k). Together these findings show

that DHX9 is essential in breast cancer cell lines and that in some ADAR1-dependent cell lines,

DHX9 suppresses PKR activation.

DHX9 and ADAR1 redundantly suppress PKR activation

The experiments above were performed in ADAR1-dependent TNBC cell lines - cell

lines that activate PKR following ADAR1 knockdown[35]. We were curious if DHX9 knockdown

would also cause PKR activation in ADAR1-independent cell lines - cell lines that do not

activate PKR following ADAR1 knockdown. Using shRNAs, we knocked down DHX9 in two

ADAR1-independent breast cancer lines, MCF-7 and SK-BR-3 (Figure 4a-b and 4f-g, Figure

S4a-b and 4g-h). Unlike in ADAR1-dependent cell lines, knockdown of DHX9 did not cause

activation of PKR in ADAR1-independent cell lines (Figure 4a, 4d, 4f, 4i). Next, we asked

whether combined knockdown of ADAR1 and DHX9 in these cell lines would lead to activation

of PKR. As we had previously observed, knockdown of ADAR1 in SK-BR-3 and MCF-7 did not

cause PKR activation[35]. However, combined knockdown of DHX9 and ADAR1 caused robust

activation of PKR in both cell lines (Figure 4a, 4d, 4f, 4i). Consistent with PKR activation, we

observed increased phosphorylation of the PKR substrate eIF2☐ following combined

knockdown of ADAR1 and DHX9 (Figure 4a and 4d, Figure S4c and S4i). Like in ADAR1-

dependent TNBC cell lines, knockdown of DHX9 caused reduced proliferation of MCF-7 and

SK-BR-3 (Figure 4c, 4d, 4h, 4j), likely through caspase-dependent apoptosis, as indicated by

elevated cleaved PARP (Figure 4a, 4d, Figure S4e, S4k). Although PARP cleavage was

increased upon combined knockdown of DHX9 and ADAR1, this did not statistically reduce the

proliferation of the cells compared to single knockdown of ADAR1 and DHX9 as measured by

foci formation. Together, these results reveal that ADAR1 and DHX9 redundantly suppress PKR

activation in ADAR1-independent breast cancer cell lines.

DHX9 and ADAR1 redundantly suppress RNase L activation

Next, we wanted to evaluate the activation of other dsRNA sensing pathways in ADAR1-

independent cell lines after the combined knockdown of ADAR1 and DHX9. To assess whether

the IFN-I pathway, or other pathways, is activated after combined knockdown of ADAR1 and

DHX9 we turned to analysis of differential gene expression by RNA-seq. In the process of

preparing RNA for sequencing, we were surprised to find specific degradation of rRNA in MCF-7

cells following combined knockdown of ADAR1 and DHX9 (Figure 4k). Single knockdown of

either DHX9 or ADAR1 did not cause rRNA degradation. The observed rRNA degradation in the

combined knockdown cells is consistent with the degradation products caused by RNase L, as

indicated in Figure 4k[49,50]. Transfection with poly(I:C), which activates the IFN-I pathway and

RNase L[51], created an identical band pattern to that of combined knockdown of DHX9 and

ADAR1, indicating that the degradation of rRNA observed in these cells is likely caused by

RNase L activity (Figure 4k). We performed the same experiment with the ADAR1-dependent

TNBC cell lines described above. In these cells, we did not see activation of RNase L after the

knockdown of DHX9 alone (Figure S3h).

DHX9 and ADAR1 redundantly suppress multiple innate immunity pathways

RNA sequencing revealed that many more RNAs were differentially expressed after

combined knockdown of DHX9 and ADAR1, compared to single knockdown of ADAR1 or DHX9

(Figure 5a, Figure S5a-f). Analysis of differential gene expression by gene set enrichment after

combined knockdown of ADAR1 and DHX9 in MCF-7 revealed activation of multiple pathways

involved in the innate response to viral infection and repression of several pathways involved in

translation (Figure 5b and Supplemental Table 14). An enrichment map showed that the

activated pathways associated with innate immunity formed one cluster and the depressed

pathways formed a separate cluster (Figure S5g). Of the upregulated pathways, several were

associated with activation of IFN signaling (Figure 5b, and Supplementary Table 14). Analysis of

core ISG expression revealed significant upregulation of ISGs in MCF-7 after combined

knockdown of ADAR1 and DHX9 (Figure 5c and Figure S5c). On the contrary, knockdown of

DHX9 or ADAR1 alone did not increase ISG expression. Activation of RNase L described above

also indicated that the type I IFN pathway was active in MCF-7 after double knockdown of

DHX9 and ADAR1, because the activators of RNase L, the OAS proteins, are ISGs[24]. In fact, a

GO term associated with OAS activity was upregulated in MCF-7 after combined knockdown of

ADAR1 and DHX9 (Supplementary Table 14). Consistent with activation of PKR, we also

observed increased expression of ATF4 targets (Figure 5d) and NF-KB targets (Figure 5e).

Interestingly, we did not observe activation of any of these pathways or RNase L in SK-BR-3

following combined ADAR1 and DHX9 knockdown. It is not clear from these data what causes

this difference between SK-BR-3 and MCF-7. SK-BR-3 expresses MDA-5, MAVS, and PKR to a

similar level as MCF-7 (Supplemental Figure 6g). Furthermore, both cell lines respond to

transfection of p(I:C) by activation of PKR and the type I IFN pathway as indicated by increased

expression of PKR and MDA5 (Figure S6h).

The finding that combined knockdown of ADAR1 and DHX9 did not induce ISG

expression in SK-BR-3 is consistent with our findings in TNBC cell lines. While knockdown of

DHX9 alone caused activation of PKR in several TNBC cell lines, we observed no activation of

the type I IFN pathway, as indicated by no change in ISG15 expression (Figure S3f-g). ISG15

was found to be highly upregulated at the RNA and protein level in MCF-7 after double

knockdown of ADAR1 and DHX9 (Figure S4f, S4m, Supplementary Table 8). Like ISG

expression overall, ISG15 expression in SK-BR-3 was not changed by knockdown of DHX9 and

ADAR1 (Figure S4l, S4n).

Given the previously described role of DHX9 in the control of Alu containing RNAs[48], we

next sought to assess if increased expression of transposable elements, especially Alus, could

explain the activation of PKR or the IFN pathway upon combined knockdown of DHX9 and

ADAR1. Analysis of our RNA-seq data revealed that transposable element expression was

generally unchanged upon either single knockdown of ADAR1 or DHX9, or combined

knockdown of both proteins (Figure S6a-f).

The dsRBDs of DHX9 are sufficient to suppress PKR activation in the absence of ADAR1

Having shown through knockdown studies that ADAR1 and DHX9 function redundantly

to suppress dsRNA sensing, we next wanted to assess which functions of DHX9 and which

isoforms of ADAR1 are important for this role. To determine which functions of DHX9 are

sufficient to suppress PKR activation, we performed a rescue experiment with wild-type DHX9,

a helicase deficient mutant DHX9, and a truncated DHX9 that possess the N-terminal dsRBDs

fused to EGFP (Figure 6a, Figure S6a). Overexpression of wild-type DHX9 rescued the PKR

activation phenotype caused by double knockdown of ADAR1 and DHX9, confirming that the

observed phenotypes are not an off-target effect of the shRNAs used for knockdown (Figure 6b-

6e). Interestingly, the DHX9[K417R] mutant, which lacks helicase activity due to its inability to bind

ATP[52], was also capable of suppressing PKR activation. The same was true for a construct

which contained the dsRBDs of DHX9 fused to EGFP (dsRBD-EGFP). Rescue experiments in

MCF-7 revealed that expression of the DHX9 dsRBD-EGFP fusion protein was sufficient to

suppress not only activation of PKR, but also the RNA expression pathways described in Figure

5 and activation of RNase L (Figure 6f, g, h and I, Figure S8) These findings indicate that the

DHX9 dsRBDs are likely sufficient to suppress activation of PKR, the type I IFN pathway and

RNase L in the absence of ADAR1 and DHX9. However, only wild-type DHX9 could rescue the

reduced foci formation observed after DHX9 knockdown (Figure 6c, e, j and k). This finding

indicates that the PKR activation phenotype and the reduced proliferation phenotype are

uncoupled.

Consistent with the observation that the DHX9 dsRBDs are sufficient to suppress PKR

activation, knockdown of DDX17, which lacks dsRBDs, did not cause substantial PKR activation

in SK-BR-3 (Figure S9). Unlike DHX9, knockdown of DDX17 had no effect on cell proliferation

as measured by the foci formation assay (Figure S9b and S9c). While combined knockdown of

DHX9 and ADAR1 in SK-BR-3 caused a 5-10 fold increase in PKR phosphorylation, the

combined knockdown of DDX17 and ADAR1 caused only a modest 1.5-fold increase in PKR

phosphorylation (Figure S9a and S9d). This finding underscores the novel role of the DHX9

helicase and its dsRBD, a unique domain among this large family of RNA helicases.

The p110 and p150 isoforms of ADAR1 suppress PKR activation in the absence of DHX9

Next, we turned to ADAR1, and asked which isoform of ADAR1 is sufficient to suppress

PKR activation in the absence of DHX9. We used the same approach as above, a rescue

experiment with overexpression of ADAR1-p110 or ADAR1-p150. Interestingly, we found that

both ADAR1 isoforms were sufficient to suppress PKR activation upon loss of DHX9 (Figure 7a

and c, Figure S10). However, overexpression of neither ADAR1 isoform was able to rescue the

foci formation phenotype, again indicating that the PKR activation and cell proliferation

phenotypes are uncoupled (Figure 7b and 7d).

Discussion

In recent years, ADAR1 has become an important therapeutic target for breast and other

cancers. It is clear from the literature that depletion of ADAR1 in ADAR1-dependent cell lines

leads to activation of dsRNA sensors and innate immunity programs that lead to cell

death[22,34,35]. Yet unclear is what distinguishes ADAR1-dependent cell lines from ADAR1-

independent cell lines - those that are insensitive to ADAR1 depletion. Elevated ISG expression

has been proposed as a potential prerequisite for ADAR1-dependency, but some ADAR1-

independent cell lines exhibit elevated ISG expression[22,35]. As such, more information is needed

to identify the factors that establish ADAR1-dependency or ADAR1-independency. To begin to

fill in some of the knowledge gaps surrounding ADAR1, we utilized proximity labeling to identify

putative ADAR1 interacting proteins, specifically focusing on the less studied ADAR1-p110

isoform.

Of the proteins identified by proximity labeling, the DEAH box helicase DHX9 was of

particular interest. Like ADAR1 and PKR, DHX9 possesses dsRBDs, a singularly unique trait

among the DEAD/DEAH box RNA helicase family. Like many RNA helicases, DHX9 is

important in numerous cellular processes, ranging from mRNA processing to resolution of R-

Loops[53,54], and has been shown to promote antiviral immunity by activation of the type I IFN,

though those studies were not performed using cancer cell lines[55,56]. DHX9 expression is

strongly correlated with ADAR1-p110 expression in breast cancer. Both genes are located on

the q-arm of chromosome 1, though they are separated by 28 mb and are thus unlikely to be

physically co-regulated. Consistent with other reports in the literature, we show here that

ADAR1 and DHX9 likely interact directly[40]. While future experiments are needed to verify that

ADAR1 interacts directly with DHX9, the data described here support an RNA-independent

interaction between ADAR1-p110 and DHX9, and an RNA-dependent interaction between

ADAR1-p150 and DHX9. This result contradicts previous findings, in which ADAR1-p150 and

DHX9 co-immunoprecipitated after RNase A treatment[48]. This discrepancy may be due to

different cell lines used. The importance of the ADAR1-DHX9 interaction is unclear from this

work. Given that knockdown of either DHX9 or ADAR1 alone can induce PKR activation in

ADAR-dependent cell lines, and that knockdown of both proteins is required to do the same in

ADAR-independent cell lines, it seems unlikely that the interaction between ADAR1 and DHX9

is required for suppression of dsRNA sensing. That of course does not preclude other functions

for a DHX9-ADAR1 complex, possibly in resolution of R-loops as both proteins have been

implicated in R-loop homeostasis[29,53]. More studies are needed to structurally assess the

interaction and directly perturb the interaction to understand what function it may have.

Here we report that in addition to being an essential gene in breast cancer, DHX9

suppresses dsRNA sensing. In ADAR1-dependent cell lines, knockdown of DHX9 alone - much

like knockdown of ADAR1 alone - caused activation of the dsRNA sensor PKR[35]. Like ADAR1

knockdown, DHX9 knockdown had no effect on PKR activation in ADAR1-independent cell

lines. However, combined knockdown of DHX9 and ADAR1 caused robust activation of PKR in

those cells. This finding indicates that ADAR1 and DHX9 function redundantly to suppress PKR

activation in ADAR1-independent cell lines and provides an explanation for why PKR is not

activated in ADAR1-independent cells upon ADAR1 knockdown. In addition to suppression of

PKR activation, we also observed that DHX9 and ADAR1 redundantly suppress activation of

several other dsRNA sensing pathways in MCF-7. Knockdown of both proteins caused

activation of IFN-I signaling, likely via MDA5 activation, as previously shown for ADAR1[13,15]. We

also observed activation of OAS-RNaseL, and increased expression of the ATF4 and NF-KB

targets, likely downstream of PKR activation[57–59]. Taken together, combined knockdown of

DHX9 and ADAR1 in MCF-7 creates a viral mimicry phenotype, where multiple innate immunity

pathways against RNA viruses have been activated . In contrast to MCF-7, we only observed

activation of PKR in other cell lines after either DHX9 knockdown alone or combined knockdown

with ADAR1. Transfection of the same cell lines with poly(I:C) revealed that each cell line was

capable of activating PKR, the type I IFN pathway and RNase L (Figures S6h and 4k), which

suggests that other variables are preventing activation of those pathways upon loss of DHX9

and ADAR1 in unresponsive cell lines. Given that PKR can be activated by much shorter

dsRNAs (>30 bp[60]) than MDA5 (>500 bp[61]), it is possible that a different set of RNAs is

responsible for activation of each pathway. Further studies are needed to identify other proteins

that function to suppress dsRNA sensing, and to identify the dsRNAs that are activating the

various pathways.

Rescue experiments revealed that the helicase activity of DHX9 was dispensable for

suppression of PKR activation, and that the N-terminal dsRBDs of DHX9 were sufficient to

suppress PKR activation in the absence of ADAR1 in ADAR1-independent cell lines. These

data are consistent with a model in which DHX9 competes with PKR for dsRNA binding. This

competition could come in the form of DHX9 binding directly to PKR and preventing its binding

to dsRNA, or DHX9 binding dsRNA and preventing PKR binding. While DHX9 and PKR interact,

and in fact DHX9 is phosphorylated by PKR [62], it is our opinion that a protein-protein interaction

model is less likely considering that DHX9 is nuclear localized whereas PKR is largely localized

to the cytoplasm [54,63–65]. Conversely, DHX9 may compete with PKR for binding of dsRNA and do

so by sequestering some dsRNAs within the nucleus. Sequestration of some dsRNAs within

the nucleus is an important means of preventing dsRNA sensors activation. During mitosis,

nuclear dsRNAs diffuse into the cytosol and activate PKR[66]. Interestingly, we did not observe a

global shift in dsRNA localization following knockdown of DHX9 and ADAR1 in MCF-7 (Figure

S11e-f). This finding may suggest that only a small subset of dsRNAs are responsible for

activation of dsRNA sensors following loss of ADAR1 and DHX9 in those cells. This hypothesis

is consistent with previous studies of ADAR1 depletion in which it was observed that only 2% of

A-to-I edits are required to prevent MDA5 activation[67]. The hypothesis that only a subset of

endogenous dsRNAs is responsible for activation PKR or MDA5 following loss of ADAR1 and/or

DHX9 could also explain the varied results seen here across different cell lines. Some cell lines

may be more sensitive to loss of ADAR1 and/or DHX9 based on the expression of specific

endogenous dsRNAs.

Previously, the ADAR1 p150 isoform, and not the p110 isoform, was shown to be

responsible for suppression of PKR activation[19,34]. Through rescue experiments, we show here

that both isoforms are sufficient to suppress PKR activation in the absence of DHX9 in ADAR1-

independent cell lines. It is important to note that the role of ADAR1-p110 in suppressing PKR

activation has likely been missed previously due to the expression of DHX9. The finding that

ADAR1-p110 can suppress PKR activation in the absence DHX9 highlights a redundant role of

these nuclear dsRNA binding proteins. An article that was published during the preparation of

this manuscript showed that the dsRBDs of ADAR1, ADAR2, and STAU1 were sufficient to

suppress PKR activation[68]. Additionally, we and others have previously shown that the

phenotype of ADAR1 depletion in ADAR1-dependent cells could at least be partially rescued by

overexpression of an editing deficient ADAR1[34,35]. Based on these findings and our rescue

experiment with the DHX9 dsRBDs, it is likely that ADAR1-p150 and ADAR1-p110 suppress

dsRNA sensing by competing with PKR for dsRNA binding. For the ADAR1-p150 isoform, this

competition with PKR for dsRNA binding is likely to be direct due to the cytoplasmic localization

of ADAR1-p150 and PKR. However, for the nuclear localized ADAR1-p110 (Figure S11d), we

propose that like DHX9, ADAR1-p110 may function to sequester PKR activating dsRNAs in the

nucleus. Interestingly, the DHX9-dsRBD-EGFP fusion protein, which we showed was capable of

suppressing PKR activation, localizes to the cytoplasm (Figure S11c). Together these findings

suggest that there may be multiple mechanisms to prevent PKR activation – direct competition,

like ADAR1-p150; and sequestration of dsRNAs, as may be the case for ADAR1-p110 and

DHX9.

Our rescue experiments revealed that while the helicase activity of DHX9 was

dispensable for suppression of dsRNA sensing, it was required for cell viability. Given that

DHX9 has been shown to have an important role in many cellular processes, ranging from

processing of mRNAs to resolution of R-Loops, we suspect the reduced viability associated with

loss of DHX9 helicase activity is related to one or more of these additional DHX9 roles[53,69]

Induction of viral mimicry has great potential as a therapeutic approach for multiple

cancers, including TNBC[70–73]. In addition to the cell intrinsic effects of activating innate immune

pathways within the tumor, the signaling that occurs after activation of those pathways can

promote anti-tumor immunity, especially when combined with checkpoint inhibitors[74–76].

Combined therapies targeting ADAR1 and DHX9 may serve as an effective means of treating

breast and potentially other cancers by inducing viral mimicry.

Materials and Methods

Cell culture

Breast cancer cell lines (MCF-7 (RRID:CVCL_0031), SK-BR-3 (RRID:CVCL_0033), BT-

549 (RRID: CVCL_1092), MDA-MB-231 (RRID: CVCL_0062), HCC1806 (RRID: CVCL_1258),

MDA-MB-468 (RRID: CVCL_0063) and 293T (RRID: CVCL_0063) were obtained from

American Type Culture Collection. All cell lines were cultured in Dulbecco's modified Eagle's

medium (DMEM) (Hyclone) with 10% fetal bovine serum (BioTechne), 2 mM glutamine

(Hyclone), 0.1 mM nonessential amino acids (Hyclone), and 1 mM sodium pyruvate (Hyclone).

Viral Production and Transduction

Lentivirus was produced by Turbo DNAfection 3000 or LipoFexin (Lamda Biotech)

transfection of 293T cells with pCMV-VSV-G, pCMV-ΔR8.2, and the appropriate plasmid for

expression of genes of interest or shRNAs. Virus was harvested 48 hours post-transfection.

Cells were transduced with lentivirus for 16 hours in the presence of 10 µg/mL protamine sulfate

(Sigma-Aldrich). The cells were selected with puromycin at 2 µg/mL (Sigma-Aldrich), 150 µg/mL

hygromycin (Invitrogen) or 10 µg/mL Blasticidin (Invitrogen).

Plasmids

APEX2 was PCR amplified from pcDNA3-APEX2-NES, a kind gift from the laboratory of

Kendall Blumer at Washington University in St. Louis. ADAR1-p110 was PCR amplified from

pLVX-p110-ADAR1 described previously[35]. APEX2 and p110 were cloned into pLVX-IRES-puro

(Takara, 632183) via a series of restriction enzyme digests and ligations. The final plasmids

pLVX-3xFLAG-APEX2 and pLVX-3xFLAG-APEX2-linker-p110 were confirmed by digestion and

sequencing. The linker consists of three repeats of Gly-Gly-Gly-Gly-Ser. Lentiviral shRNA

constructs in the pLKO.1-puro vector were purchased as glycerol stocks from Millipore Sigma.

For shADAR1, the shRNA was subcloned into pLKO.1-hygro, a gift from Bob Weinberg

(Addgene, #24150). The sequences for the shRNA-scramble (shSCR) and shRNA-ADAR1

(shADAR1) were described and validated previously[35]. The sequences for shRNAs against

DHX9 and DDX17 are in Supplementary Table 1. For sgDHX9, oligos encoding the sgRNAs

(Supplemental Table 1) were cloned into lentiCRISPR v2, a gift from Feng Zhang (Addgene,

#52961)[77]. Overexpression constructs for DHX9 and ADAR were generated by PCR

amplification and ligation into pLV-EF1a-IRES-Blast vector, a gift from Tobias Meyer (Addgene,

#85133)[78]. For DHX9 overexpression, wobble mutants were made to reduce shRNA targeting.

Mutagenesis primers for DHX9 K417R and shRNA-resistant codons (designed using the

Synonymous Mutation Generator[79]) are included in Supplemental Table 1. The DHX9-dsRBD-

EGFP construct was generated by digestion of pLV-EF1-DHX9 with SpeI and EcoRI and

ligation of EGFP in place of the 3' portion of DHX9. The resulting construct codes for the first

344 amino acids of DHX9 fused to EGFP.

Immunoblot

Cell pellets were lysed and sonicated in RIPA Buffer (50 mM Tris pH 7.4 (Ambion), 150

mM NaCl (Ambion), 1% Triton X-100 (Sigma-Aldrich), 0.1% sodium dodecyl sulfate (Promega)

and 0.5% sodium deoxycholate (Sigma-Aldrich) with 1x HALT Protease Inhibitor (Pierce).

Protein was quantified using the DC Assay kit (Bio-Rad) and diluted in SDS Sample Buffer (125

mM Tris pH 6.8, 30% glycerol, 10% sodium dodecyl sulfate, 0.012% bromophenol blue) prior to

denaturation by heating to 95 °C for 7 minutes. Thirty micrograms of protein lysate were

resolved on 4-12% TGX Acrylamide Stain-Free gels (Bio-Rad). Stain-Free gels were imaged

prior to transfer to PVDF membrane (Millipore or Bio-Rad) by TransBlot Turbo (Bio-Rad). The

blots were then probed with the appropriate primary antibodies: ADAR1 (Santa Cruz, sc-73408;

Bethyl Laboratories, A303-883A), DDX17 (Thermo Scientific, PA5-84585), DHX9 (Bethyl, A300-

855A), DDX54 (Novus Biologicals, NB100-60678), eIF2a (Abcam, ab5369), eIF2a-Ser-51-P

(Abcam, ab32157), Fibrillarin (Santa Cruz, sc-25397), beta-tubulin (Abcam, ab6046), ISG15

(Santa Cruz, sc-166755), cleaved PARP (Cell Signaling, #9541), PKR (Cell Signaling, #3072),

PKR Thr-446-P (Abcam, ab32036), MDA5 (Cell Signaling, #5321S), MAVS (Cell Signaling,

#3993S). Primary antibodies were detected with horseradish-peroxidase conjugated secondary

antibodies (Jackson ImmunoResearch) and detection was carried out with Clarity Western ECL

Substrate (Bio-Rad). Densitometry was performed using Image Lab (Bio-Rad). Band intensity

was normalized to total protein measured by imaging of the Stain-Free gel.

Proximity Labeling by APEX2

SK-BR-3, MCF-7, or MDA-MB-231 cells expressing pLVX-puro-FLAG-APEX2 or pLVX-

puro-FLAG-APEX2-ADAR1p110 were grown to ~80% confluency in a 15 cm dish. Quencher

solution (10 mM sodium azide, 10 mM sodium ascorbate, and 5 mM Trolox (Sigma-Aldrich) in

1X PBS) was prepared at 1X and 2X concentrations. Prior to labeling, cells were incubated in

10 mL of culture media containing 500 µM biotin phenol (Toronto Research Chemicals,

B397770) for 30 min at 37 °C. Next, hydrogen peroxide was added to the cells at 1 mM and

incubated at room temperature for 1 min. Immediately one volume of 2X quencher solution was

added to the cells to stop labeling. The cells were washed twice with 1X quencher solution.

Cells were harvested by scraping in 1X quencher solution and lysed in RIPA buffer containing

10 mM sodium azide, 10 mM sodium ascorbate, 5 mM Trolox and 1x HALT. Biotin labeling was

verified by immunoblotting with HRP-streptavidin (Abcam, ab7403). Biotinylated proteins were

purified using streptavidin magnetic beads (Thermo Fisher Scientific, 88816). Streptavidin

magnetic beads were washed twice with RIPA containing HALT and quenching agents. The

lysate from above was incubated with the beads for 1 hour at room temperature. The beads

were washed in the following order: once with RIPA containing HALT and quenching agents,

once with RIPA, once with 1 M KCl, once with 2 M urea pH 8.0, twice in RIPA and once with

water. Elution was performed in 1X SDS-Sample buffer by heating at 95 °C for 10 minutes. The

eluate was analyzed by LC-MS-MS, see below.

Mass Spectrometry

Liquid-chromatography and tandem mass spectrometry was performed by MSBioWorks

(Ann Arbor, MI). The eluates from the streptavidin pulldown above were processed by SDS-

PAGE using 10% Bis-Tris NuPage Mini-gel (Invitrogen) with the MES buffer system. The gel

was run 2cm. The mobility region was excised and processed by in-gel digestion with trypsin

using a robot (ProGest, DigiLab). For the trypsin digestion, the gel slices were washed with

25mM ammonium bicarbonate followed by acetonitrile. The samples were reduced with 10mM

dithiothreitol at 60°C followed by alkylation with 50mM iodoacetamide at room temperature.

Subsequently proteins were digested with trypsin (Promega) at 37°C for 4 h. The trypsin

digestion was quenched with formic acid and the supernatant was analyzed directly without

further processing. The digested sample was analyzed by nano LC-MS/MS with a Waters M-

Class HPLC system interfaced to a ThermoFisher Fusion Lumos mass spectrometer. Peptides

were loaded on a trapping column and eluted over a 75 μm analytical column at 350 nL/min;

both columns were packed with Luna C18 resin (Phenomenex). The mass spectrometer was

operated in data-dependent mode, with the Orbitrap operating at 60,000 FWHM and 15,000

FWHM for MS and MS/MS respectively. APD was enabled and the instrument was run with a 3

s cycle for MS and MS/MS. Five hours of instrument time was used for the analysis of each

sample.

Analysis of Mass Spectrometry Data

Data were searched using a local copy of Mascot (Matrix Science) with the following

parameters: Enzyme - Trypsin/P; Database - SwissProt Human (concatenated forward and

reverse plus common contaminants); Fixed modification – Carbamidomethyl; Variable

modifications - Oxidation; Acetyl; Pyro-Glu; Deamidation; Mass values – Monoisotopic; Peptide

Mass Tolerance - 10 ppm; Fragment Mass Tolerance - 0.02 Da; Max Missed Cleavages - 2.

Mascot DAT files were parsed into Scaffold (Proteome Software) for validation, filtering and to

create a non-redundant list per sample. Data were filtered using a 1% protein and peptide FDR

and requiring at least two unique peptides per protein. Fold change of protein abundance was

determined by DESeq2 using spectral counts (see Data Availability below for scripts).

Overrepresentation analysis was performed using 'enrichR' in R[80]. The cutoff used for

enrichment for the overrepresentation analysis was an FDR < 0.05 and a log2 fold change of >

0.5.

Immunoprecipitation

Cell lysates prepared in RIPA with 1X HALT. RNaseOUT (Thermo Fisher) RNase

inhibitor was added to the lysis buffer at 0.5 U/µL when RNase A was not used. One milligram

of protein lysate was mixed with 2-10 µg of IgG or specific antibody overnight at 4 °C with

rotation. For samples treated with RNase A, 20 µg of RNase A (Invitrogen) was added to the

lysate during overnight mixing with the antibody. Protein G Dynabeads (Thermo Fisher, 25 µL

per sample) were prepared by washing twice in the lysis buffer. Prepared beads were mixed

with lysates for 30 min at 4 °C with rotation. Supernatants were collected and beads were

washed three times in the lysis buffer, and eluted by mixing the beads in SDS sample buffer and

incubating at 95 °C for 7 min. Antibodies: Rabbit IgG (Jackson ImmunoResearch, 011-000-003),

Mouse IgG (Jackson ImmunoResearch, 015-000-003), DHX9 (Bethyl, A300-855A), PARP (Cell

Signaling, 9532S), XRN2 (Novus, NB100-57541), DDX54 (Novus, NB100-60678), DDX17

(Thermo Scientific, PA5-84585).

Immunofluorescence

Cells were plated on glass coverslips (Corning) two days prior to fixation for

immunofluorescence. Cells were washed in PBS prior to fixation with 4% paraformaldehyde

(Thermo Scientific) and permeabilization with 0.15% Triton-X100 in PBS. Following

permeabilization the cells were washed three times with PBS then blocked with Protein Block

(Aligent/Dako, X090930-2). Primary antibodies (ADAR1 (Santa Cruz, sc-73408), Fibrillarin

(Santa Cruz, sc-25397), DDX54 (Novus Biologicals, NB100-60678) or DDX17 (Thermo

Scientific, PA5-84585), DHX9 (Bethyl, A300-855A), ADAR1-p150 (Abcam, Ab126745), dsRNA-

J2 (Millipore-Sigma, MABE1134) and secondary antibodies (Thermo Scientific, A21207,

A21203, A21202, A21206) were diluted in Antibody Diluent (Agilent/Dako, S302283-2).

Antibody binding was performed in a humidity chamber for 1.5 hours for primaries and 30

minutes for secondaries. Between primary and secondary antibodies, and after secondary

antibody binding the coverslips were washed in PBS. The coverslips were washed once in

water before mounting on glass slides with Vectashield Antifade Mounting Media with DAPI

(Vector Laboratories, H-1200-10). Fluorescence microscopy images were obtained with an

Eclipse 90i microscope (Nikon) using a Plan Apochromatic 20x/NA 0.75 objective (Nikon) and a

CoolSNAP ES2 monochrome digital camera cooled to 0°C (Photometrics). Fluorescence

images were captured with MetaMorph version 7.8.0.0 software (Molecular Devices) and

resized and formatted with Fiji.

Transfection of poly(I:C)

The cell line indicated was transfected with high-molecular weight poly(I:C) (Invivogen)

with Lipofectamine LTX (Invitrogen). Three microliters of Lipofectamine LTX was used per

microgram of poly(I:C). Sixteen to 24 hours after transfection, cells were harvested in RIPA with

1X HALT or the RNA lysis buffer from the Nucleospin RNA kit (Macherey-Nagel).

RNA Purification and RNA sequencing

RNA-sequencing was performed for two replicates of ADAR1 and/or DHX9 knockdown

in MCF-7 and SK-BR-3. RNA was purified using the Nucleospin RNA kit (Macherey-Nagel).

Assessment of rRNA integrity and RNA-sequencing was performed by the Genome Technology

Access Center at Washington University in St. Louis. Total RNA integrity was determined using

Agilent TapeStation 4200. Library preparation was performed with 500 ng to 1 ug of total RNA.

Ribosomal RNA was removed by an RNase-H method using RiboErase kit (Kapa Biosystems).

After rRNA depletion, the remaining RNA was then fragmented in reverse transcriptase buffer

(Life Technologies) by heating to 94 degrees for 8 minutes. The RNA was reverse transcribed to

yield cDNA using SuperScript III RT and random hexamers (Life Technologies) per

manufacturer's instructions. A second strand reaction was performed with DNA Polymerase I

and RNase H (Qiagen) to yield ds-cDNA. The cDNA was then blunted with T4 DNA

Polymerase, Polynucleotide Kinase and Klenow DNA Polymerase (Qiagen). An A base was

added to the 3' ends with Klenow (3'-4' exo-) (Qiagen). The processed ds-cDNA was then

ligated to Illumina sequencing adapters with T4 DNA Ligase (Qiagen). Ligated fragments were

then amplified for 12-15 cycles using primers incorporating unique dual index tags with VeraSeq

polymerase (Qiagen). Fragments were sequenced on an Illumina NovaSeq-6000 using paired

end reads extending 150 bases.

RNA Sequencing Analysis

The Illumina bcl2fastq software was used for base calling and demultiplexing, allowing

for one mismatch in the indexing read. STAR version 2.7.9a1 was used for read alignment to

RNA-seq to the Ensembl GRCh38.101 primary assembly. Gene counts were determined using

Subread:featureCount version 2.0.32, only uniquely aligned unambiguous reads were counted.

Differential gene expression was determined using DESeq2 (see Data Availability below for

scripts)[81]. The experimental design for DESeq2 analysis included an interaction term between

the shRNAs used for knockdown ('shrna1 + shrna2 + shrna1:shrna2'; where 'shrna1' was either

shSCR or shADAR and 'shrna2' was either shSCR or shDHX9-3). Contrasts were used to

assess differential expression after singular knockdown of either ADAR1 or DHX9. Fold

changes were shrunken using the 'apeglm' approach from DESeq2[82]. Gene set enrichment

analysis was performed using 'clusterProfiler'[83]. For analysis of transposable element

expression, 'TEcount' from TEtranscripts[84] was used to determine family level counts for

transposable elements using a GTF file containing transposable element information from

RepeatMasker (http://www.repeatmasker.org). (see Data Availability section for more

information about the GTF file).

Foci Formation Assay

Five thousand cells were plated for each condition in a 10 cm culture dish. After 10 (BT-

549, MDA-MB-MB231, HCC1806, MDA-MB-468 and SK-BR-3) to 20 (MCF-7) days the cells

were washed briefly with 1x PBS prior to fixation in 100% methanol for 5 min. After drying, the

cells were stained with 0.005% Crystal Violet solution containing 25% methanol (Sigma-Aldrich)

prior to washing excess stain away with deionized water. The plates were scanned using an

ImageScanner III (General Electric). Foci area was calculated using ImageJ.

Quantitative PCR

572   Reverse transcription and qPCR were performed as described previously using iScript

573   Supermix for cDNA synthesis and iTaq for qPCR (Bio-Rad)[85]. Fold change of RNA expression

574   was determined using the $\Delta\Delta C_t$ method with normalization to PSMA5 and OAZ1. Primers are

575   shown in Supplemental Table 1.

Analysis of CCLE and TCGA data

577   RNA-seq normalization and calculation of z-scores was performed as described

578   previously[35]. Molecular subtypes of breast cancer cell lines and TCGA samples were defined

579   previously[86]. Breast cancer survival analysis was performed using the R packages RTCGA and

580   survminer[87,88]. The DHX9 expression level used for stratification in survival analysis was

581   determine by the surv_cutpoint function of survminer.

Data and Code Availability

Scripts used for analysis of mass spectrometry, RNA-seq and generation of all plots are

available at (https://github.com/cottrellka/Cottrell-Ryu-et-al-2023), raw sequencing data and

count files were deposited at the Gene Expression Omnibus (GEO) under accession

GSE224677. RNA-seq data for cancer cell lines (CCLE_expression_full.csv,

CCLE_RNAseq_rsem_transcripts_tpm_20180929.txt) were obtained from the DepMap portal

(https://depmap.org/portal/download/custom/)[89]. RNAi based dependency data for DHX9

(D2_combined_gene_dep_scores) was obtained from DepMap Portal

Cottrell, Ryu et al., 2023   24

(https://depmap.org/portal/download/custom/)[90]. RNA-seq data for TCGA BRCA samples

(illuminahiseq_rnaseqv2-RSEM_genes, illuminahiseq_rnaseqv2-RSEM_isoforms_normalized)

and clinical data (Merge_Clinical) were obtained from the Broad Institute FireBrowse and are

available at http://firebrowse.org/. The GTF file used for TEcount

(GRCh38_Ensembl_rmsk_TE.gtf) is available at

https://www.dropbox.com/sh/1ppg2e0fbc64bqw/AACUXf-TA1rnBIjvykMH2Lcia?dl=0.

Acknowledgements:

The authors would like to thank Leonard B. Maggi, Jr., Raleigh Kladney, Louis Kerestes,

Che-Pei Kung, and Anbrielle Blake for technical assistance, Elliot Klotz and Toni Sinnwell

(Genome Technology Access Center at Washington University) for RNA sequencing, Joshua B.

Rubin, Jason C. Mills, Jieya Shao, and Alessandro Vindigni for discussion and suggestions. The

data described here is in part based on data generated by the TCGA Research

Network: https://www.cancer.gov/tcga. This work was supported by R01CA262804 (J.D.W.),

K99MD016946 and R00MD016946 (K.A.C) from the National Institutes of Health, W81XWH-18-

1-0025 (J.D.W.) and W81XWH-21-10466 (J.D.W.) from the Department of Defense, and a

Centene Corporation Contract (P19-00559) for the Washington University–Centene ARCH

Personalized Medicine Initiative (J.D.W.). L.S.T. supported by R25HG006687 and

T32GM148405.

Author Contributions

K.A.C, S.R. and J.D.W. conceived the project. K.A.C., S.R., L.S.T., and A.M.S.

performed the experiments. K.A.C., S.R., and L.S.T. performed the data analysis. K.A.C. and

S.R. wrote the manuscript. All authors edited the manuscript.

Declaration of interests

The authors declare no competing interests.

Figure 1: Identification of putative ADAR1 interacting proteins by APEX2 proximity labeling

a Representative immunoblot showing expression of the constructs used for proximity labeling in MCF-7, MDA-MB-231 and SK-BR-3, MDA-MB-231 is shown. **b** Representative fluorescently stained gel image showing proteins purified by streptavidin-biotin pulldown following proximity labeling in MCF-7. **c** Volcano plot summarizing the proteins identified by mass spectrometry following streptavidin pulldown subsequent to proximity labeling. Differential abundance of all proteins in each cell line can be found in Supplementary Table 2. **d** Venn diagram showing overlap between enriched proteins from all three cell lines. The cut-off for enriched proteins was an FDR adjusted p-value of less than 0.05 and a Log2 fold change of greater than 0.5. **e** GO terms found to be overrepresented in the list of the enriched proteins. Only the top ten GO terms, by FDR, are shown for each category. All other significant GO terms are available in the Supplementary Table 3-4. **f** and **g** Venn diagrams showing overlap between the enriched proteins identified and those proteins found previously to localize to the nucleus or nucleolus in MCF-7[45]. **h** Representative indirect immunofluorescence micrographs showing localization of ADAR1, fibrillarin (a nucleolar marker) and DAPI.

Figure 2: Validation of putative protein-protein interactions identified by proximity

labeling

Immunoprecipitation of DHX9 **a-c**, DDX54 **d-e** or DDX17 **f-g** followed by immunoblot for ADAR

in breast cancer cell lines**.** Immunoblot of immunoprecipitation eluates and inputs from SK-BR-3

a, **d** and **f**, MCF-7 **b**, **e** and **g**, and MDA-MB-231 **c**. Input represents 5% of the lysate used for

immunoprecipitation. The IgG lanes represent immunoprecipitation eluates from pulldown with

anti-rabbit IgG antibody. The lanes labeled DHX9, DDX54 and DDX17 indicate the eluates from

immunoprecipitation with antibodies against those proteins respectively. The IgG^{HC} label

indicates the band corresponding to the IgG heavy chain from the antibody used for

immunoprecipitation. Uncropped immunoblots for panels **a-g** can be found in Source Data

Figures. Immunofluorescence for ADAR1 and DHX9 **h**, DDX54 **i**, or DDX17 **j** in SKBR3 or MCF-

7.

Figure 3: DHX9 is overexpressed in breast cancer and suppresses PKR activation

a Schematic showing the domain structure of PKR, ADAR1, DHX9 and other helicases

identified by proximity labeling in Figure 1, dsRBD refers to the dsRNA Binding Domain. **b**

Pearson and Spearman correlation coefficients for the correlation between ADAR1 expression

at the RNA level and the expression of each indicated helicase at the RNA level, data from

breast tumors within TCGA. Scatterplots showing the correlation between ADAR1-p110 **c**, or

ADAR1-p150 **d**, and DHX9 expression in normal breast or breast tumors. **e** Expression of DHX9

at the RNA level in normal breast, non-TNBC or TNBC tumors. **f** Waterfall plot showing DHX9

dependency of breast cancer cell lines using data from DepMap, ER = estrogen receptor

positive cell lines, ERRB2 = HER2 positive cell lines. **g** Representative immunoblot following

knockdown of DHX9 with two different shRNAs in four TNBC cell lines. Immunoblots for other

replicates and uncropped blots can be found in Source Data Figures. **h** Foci formation assay for

the same cells used in **g** for immunoblot. **i** Quantification of PKR phosphorylation as determined

by the immunoblot in **g**. Quantification of protein expression for other proteins of interest can be

found in Figure S3a-e. Bars represent the average of at least three replicates, error bars are +/-

SD. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. P-values determined by Dunnett's test.

 Figure 4: DHX9 and ADAR1 redundantly suppress dsRNA sensing in ADAR1-independent

cell lines

Representative immunoblot showing the phenotype of ADAR1 and/or DHX9 knockdown in

MCF-7 **a**, or SK-BR-3 **f**. Immunoblots for other replicates and uncropped blots can be found in

Source Data Figures. Protein abundance from the immunoblot in **a** and **f** was normalized by

total protein abundance by quantification of the Stain Free gel in **b** and **g** respectively. Fold

change of PKR phosphorylation at Thr-446 in MCF-7 **d** or SK-BR-3 **i** as determined by the

immunoblots in **a** or **f** respectively. Quantification of protein expression for other proteins of

interest can be found in Figure S4a-l. Representative foci formation phenotype of ADAR1 and/or

DHX9 knockdown in MCF-7 **c** or SK-BR-3 **h**, quantification of relative foci area is shown in **e** or **j**

respectively. **k** Analysis of rRNA integrity upon knockdown of ADAR1 and/or DHX9 in MCF-7 or

SK-BR-3. Additionally, panel **k** shows the effect of poly(I:C) (p(I:C)) transfection on rRNA

integrity in MCF-7. Arrows indicate canonical RNase L cleavage products[50]. Bars represent the

average of at least three replicates, error bars are +/- SD. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. P-

values determined by one-way ANOVA with post-hoc Tukey. Comparisons between the two

different shRNAs targeting DHX9 (shDHX9-3 and shDHX9-5) were not included for clarity.

Figure 5: Induction of a viral mimicry phenotype upon knockdown of DHX9 and ADAR1 in

MCF-7

A Volcano plot showing changes in RNA expression upon knockdown of DHX9 and ADAR1 in

MCF-7, a volcano plot for SK-BR-3 is in Figure S5f. Fold-change of RNA expression shown in **a**

was determined using an interaction term between ADAR1 and DHX9 knockdown, volcano plots

for fold change of RNA expression for single knockdown of ADAR1 or DHX9 is in Figure S5a-b

and S5d-e. **b** GO terms identified by gene set enrichment analysis of the RNA-seq data in **a**. **c-e**

Heatmaps and summary box plots showing RNA expression changes in MCF-7 and SK-BR-3

upon knockdown of ADAR1 and/or DHX9. Panel **c** shows RNA expression for core ISGs[22,35],

panel **d** shows ATF4 targets and panel **e** shows NF-KB targets with ISGs removed.

Figure 6: Rescue of PKR activation by ADAR1-p110, ADAR-p150, DHX9 and DHX9 mutants

a Schematic showing the domain structure of ADAR1 isoforms, DHX9 and mutants of DHX9, dsRBD refers to the dsRNA Binding Domain. Representative immunoblot showing the phenotype of ADAR1 and DHX9 knockdown with DHX9, DHX9^{K417R} or dsRBD-EGFP overexpression in SK-BR-3 **b**, or MCF-7 **f**. Immunoblots for other replicates and uncropped blots can be found in Source Data Figures. Representative foci formation phenotype of ADAR1 and DHX9 knockdown with DHX9, DHX9^{K417R} or dsRBD-EGFP overexpression in SK-BR-3 **c**, or MCF-7 **j**. Fold change of PKR phosphorylation at Thr-446 upon ADAR1 and DHX9 knockdown with DHX9, DHX9^{K417R} or dsRBD-EGFP overexpression in SK-BR-3 **d**, or MCF-7 **g**, quantified from immunoblots in **b** and **f** respectively. Quantification of protein expression for other proteins of interest can be found in Figure S7b-f and S8b-g. Quantification of relative foci area is shown in **e** SK-BR-3 and **k** for MCF-7. **h** RNA expression of representative genes from upregulated pathways in Figure 5 as determined by qRT-PCR for the MCF-7 rescue experiment. Statistical analysis for each pathway is shown in Figure S8h-j. **i** Analysis of rRNA integrity upon knockdown of ADAR1 and DHX9 in MCF-7 with overexpression of EGFP or dsRBD-EGFP. Arrows indicate canonical RNase L cleavage products[50]. Bars represent the average of at least three replicates, error bars are +/- SD. * p <0.05, ** p <0.01, *** p < 0.001. P-values determined by Dunnett's test.

Figure 7: Rescue of PKR activation by ADAR1-p110 and ADAR-p150

Representative immunoblot showing the phenotype of ADAR1 and DHX9 knockdown with

ADAR1-p110 or ADAR1-p150 overexpression in SK-BR-3 **a**. Immunoblots for other replicates

and uncropped blots can be found in Source Data Figures. Representative foci formation

phenotype of ADAR1 and DHX9 knockdown ADAR1 isoform overexpression **b**. Quantification of

relative foci area is shown in **d**. Fold change of PKR phosphorylation at Thr-446 upon ADAR1

and DHX9 knockdown with ADAR1 isoform overexpression **c**, quantified from immunoblots in **a**.

Quantification of protein expression for other proteins of interest can be found in Figure S10a-f.

Bars represent the average of at least three replicates, error bars are +/- SD. *** $p < 0.001$. P-

values determined by Dunnett's test.

References

1.   Paz-Yaacov, N., Bazak, L., Buchumenski, I., Porath, H.T., Danan-Gotthold, M., Knisbacher,
B.A., Eisenberg, E., and Levanon, E.Y. (2015). Elevated RNA Editing Activity Is a Major
Contributor to Transcriptomic Diversity in Tumors. Cell Rep. *13*, 267–276.
10.1016/j.celrep.2015.08.080.

2.   Walkley, C.R., and Li, J.B. (2017). Rewriting the transcriptome: adenosine-to-inosine RNA
editing by ADARs. Genome Biol. *18*, 205. 10.1186/s13059-017-1347-3.

3.   Li, M., Wang, I.X., Li, Y., Bruzel, A., Richards, A.L., Toung, J.M., and Cheung, V.G. (2011).
Widespread RNA and DNA Sequence Differences in the Human Transcriptome. Science
333, 53–58. 10.1126/science.1207018.

4.   Bazak, L., Haviv, A., Barak, M., Jacob-Hirsch, J., Deng, P., Zhang, R., Isaacs, F.J., Rechavi,
G., Li, J.B., Eisenberg, E., et al. (2014). A-to-I RNA editing occurs at over a hundred million
genomic sites, located in a majority of human genes. Genome Res. *24*, 365–376.
10.1101/gr.164749.113.

5.   Wagner, R.W., Smith, J.E., Cooperman, B.S., and Nishikura, K. (1989). A double-stranded
RNA unwinding activity introduces structural alterations by means of adenosine to inosine
conversions in mammalian cells and Xenopus eggs. Proc. Natl. Acad. Sci. *86*, 2647–2651.
10.1073/pnas.86.8.2647.

6.   Kim, U., Wang, Y., Sanford, T., Zeng, Y., and Nishikura, K. (1994). Molecular cloning of
cDNA for double-stranded RNA adenosine deaminase, a candidate enzyme for nuclear
RNA editing. Proc. Natl. Acad. Sci. *91*, 11457–11461. 10.1073/pnas.91.24.11457.

7.   Herbert, A., Alfken, J., Kim, Y.-G., Mian, I.S., Nishikura, K., and Rich, A. (1997). A Z-DNA
binding domain present in the human editing enzyme, double-stranded RNA
adenosine⬜deaminase. Proc. Natl. Acad. Sci. *94*, 8421–8426. 10.1073/pnas.94.16.8421.

8.   George, C.X., and Samuel, C.E. (1999). Human RNA-specific adenosine deaminase
ADAR1 transcripts possess alternative exon 1 structures that initiate from different
promoters, one constitutively active and the other interferon inducible. Proc. Natl. Acad. Sci.
96, 4621–4626. 10.1073/pnas.96.8.4621.

9.   Desterro, J.M.P., Keegan, L.P., Lafarga, M., Berciano, M.T., O'Connell, M., and Carmo-
Fonseca, M. (2003). Dynamic association of RNA-editing enzymes with the nucleolus. J.
Cell Sci. *116*, 1805–1818. 10.1242/jcs.00371.

10.   Patterson, J.B., and Samuel, C.E. (1995). Expression and regulation by interferon of a
double-stranded-RNA-specific adenosine deaminase from human cells: evidence for two
forms of the deaminase. Mol. Cell. Biol. *15*, 5376–5388. 10.1128/MCB.15.10.5376.

11.   Eckmann, C.R., Neunteufl, A., Pfaffstetter, L., and Jantsch, M.F. (2001). The human but not
the Xenopus RNA-editing enzyme ADAR1 has an atypical nuclear localization signal and
displays the characteristics of a shuttling protein. Mol. Biol. Cell *12*, 1911–1924.
10.1091/mbc.12.7.1911.

12. Chung, H., Calis, J.J.A., Wu, X., Sun, T., Yu, Y., Sarbanes, S.L., Dao Thi, V.L., Shilvock,
A.R., Hoffmann, H.-H., Rosenberg, B.R., et al. (2018). Human ADAR1 Prevents
Endogenous RNA from Triggering Translational Shutdown. Cell *172*, 811-824.e14.
10.1016/j.cell.2017.12.038.

13. Liddicoat, B.J., Piskol, R., Chalk, A.M., Ramaswami, G., Higuchi, M., Hartner, J.C., Li, J.B.,
Seeburg, P.H., and Walkley, C.R. (2015). RNA editing by ADAR1 prevents MDA5 sensing
of endogenous dsRNA as nonself. Science *349*, 1115–1120. 10.1126/science.aac7049.

14. Pestal, K., Funk, C.C., Snyder, J.M., Price, N.D., Treuting, P.M., and Stetson, D.B. (2015).
Isoforms of the RNA editing enzyme ADAR1 independently control nucleic acid sensor
MDA5-driven autoimmunity and multi-organ development. Immunity *43*, 933–944.
10.1016/j.immuni.2015.11.001.

15. Mannion, N.M., Greenwood, S.M., Young, R., Cox, S., Brindle, J., Read, D., Nellåker, C.,
Vesely, C., Ponting, C.P., McLaughlin, P.J., et al. (2014). The RNA-Editing Enzyme ADAR1
Controls Innate Immune Responses to RNA. Cell Rep. *9*, 1482–1494.
10.1016/j.celrep.2014.10.041.

16. Chen, Y.G., and Hur, S. (2022). Cellular origins of dsRNA, their recognition and
consequences. Nat. Rev. Mol. Cell Biol. *23*, 286–301. 10.1038/s41580-021-00430-1.

17. Yang, S., Deng, P., Zhu, Z., Zhu, J., Wang, G., Zhang, L., Chen, A.F., Wang, T., Sarkar,
S.N., Billiar, T.R., et al. (2014). ADAR1 Limits RIG-I RNA Detection and Suppresses IFN
Production Responding to Viral and Endogenous RNAs. J. Immunol. Baltim. Md 1950 *193*,
3436–3445. 10.4049/jimmunol.1401136.

18. George, C.X., Ramaswami, G., Li, J.B., and Samuel, C.E. (2016). Editing of Cellular Self-
RNAs by Adenosine Deaminase ADAR1 Suppresses Innate Immune Stress Responses*. J.
Biol. Chem. *291*, 6158–6168. 10.1074/jbc.M115.709014.

19. Clerzius, G., Gélinas, J.-F., Daher, A., Bonnet, M., Meurs, E.F., and Gatignol, A. (2009).
ADAR1 Interacts with PKR during Human Immunodeficiency Virus Infection of Lymphocytes
and Contributes to Viral Replication. J. Virol. *83*, 10119–10128. 10.1128/JVI.02457-08.

20. Quin, J., Sedmík, J., Vukić, D., Khan, A., Keegan, L.P., and O'Connell, M.A. (2021). ADAR
RNA Modifications, the Epitranscriptome and Innate Immunity. Trends Biochem. Sci. *46*,
758–771. 10.1016/j.tibs.2021.02.002.

21. Montavon, T.C., Baldaccini, M., Lefèvre, M., Girardi, E., Chane-Woon-Ming, B., Messmer,
M., Hammann, P., Chicher, J., and Pfeffer, S. (2021). Human DICER helicase domain
recruits PKR and modulates its antiviral activity. PLOS Pathog. *17*, e1009549.
10.1371/journal.ppat.1009549.

22. Liu, H., Golji, J., Brodeur, L.K., Chung, F.S., Chen, J.T., deBeaumont, R.S., Bullock, C.P.,
Jones, M.D., Kerr, G., Li, L., et al. (2019). Tumor-derived IFN triggers chronic pathway
agonism and sensitivity to ADAR loss. Nat. Med. *25*, 95–102. 10.1038/s41591-018-0302-5.

23. Nakahama, T., Kato, Y., Shibuya, T., Inoue, M., Kim, J.I., Vongpipatana, T., Todo, H., Xing,
Y., and Kawahara, Y. (2021). Mutations in the adenosine deaminase ADAR1 that prevent

endogenous Z-RNA binding induce Aicardi-Goutières-syndrome-like encephalopathy.
Immunity *54*, 1976-1988.e7. 10.1016/j.immuni.2021.08.022.

24. Li, Y., Banerjee, S., Goldstein, S.A., Dong, B., Gaughan, C., Rath, S., Donovan, J.,
Korennykh, A., Silverman, R.H., and Weiss, S.R. (2017). Ribonuclease L mediates the cell-
lethal phenotype of double-stranded RNA editing enzyme ADAR1 deficiency in a human cell
line. eLife *6*, e25687. 10.7554/eLife.25687.

25. Rath, S., Prangley, E., Donovan, J., Demarest, K., Wingreen, N.S., Meir, Y., and Korennykh,
A. (2019). Concerted 2-5A-Mediated mRNA Decay and Transcription Reprogram Protein
Synthesis in the dsRNA Response. Mol. Cell *75*, 1218-1228.e6.
10.1016/j.molcel.2019.07.027.

26. Schoggins, J.W., and Rice, C.M. (2011). Interferon-stimulated genes and their antiviral
effector functions. Curr. Opin. Virol. *1*, 519–525. 10.1016/j.coviro.2011.10.008.

27. George, C.X., Das, S., and Samuel, C.E. (2008). Organization of the mouse RNA-specific
adenosine deaminase Adar1 gene 5'-region and demonstration of STAT1-independent,
STAT2-dependent transcriptional activation by interferon. Virology *380*, 338–343.
10.1016/j.virol.2008.07.029.

28. Sun, T., Yu, Y., Wu, X., Acevedo, A., Luo, J.-D., Wang, J., Schneider, W.M., Hurwitz, B.,
Rosenberg, B.R., Chung, H., et al. (2021). Decoupling expression and editing preferences
of ADAR1 p150 and p110 isoforms. Proc. Natl. Acad. Sci. *118*, e2021757118.
10.1073/pnas.2021757118.

29. Shiromoto, Y., Sakurai, M., Minakuchi, M., Ariyoshi, K., and Nishikura, K. (2021). ADAR1
RNA editing enzyme regulates R-loop formation and genome stability at telomeres in cancer
cells. Nat. Commun. *12*, 1654. 10.1038/s41467-021-21921-x.

30. Fumagalli, D., Gacquer, D., Rothé, F., Lefort, A., Libert, F., Brown, D., Kheddoumi, N.,
Shlien, A., Konopka, T., Salgado, R., et al. (2015). Principles Governing A-to-I RNA Editing
in the Breast Cancer Transcriptome. Cell Rep. *13*, 277–289. 10.1016/j.celrep.2015.09.032.

31. Kung, C.-P., Maggi, L.B., and Weber, J.D. (2018). The Role of RNA Editing in Cancer
Development and Metabolic Disorders. Front. Endocrinol. *9*.

32. Paz, N., Levanon, E.Y., Amariglio, N., Heimberger, A.B., Ram, Z., Constantini, S., Barbash,
Z.S., Adamsky, K., Safran, M., Hirschberg, A., et al. (2007). Altered adenosine-to-inosine
RNA editing in human cancer. Genome Res. *17*, 1586–1595. 10.1101/gr.6493107.

33. Chan, T.H.M., Lin, C.H., Qi, L., Fei, J., Li, Y., Yong, K.J., Liu, M., Song, Y., Chow, R.K.K.,
Ng, V.H.E., et al. (2014). A disrupted RNA editing balance mediated by ADARs (Adenosine
DeAminases that act on RNA) in human hepatocellular carcinoma. Gut *63*, 832–843.
10.1136/gutjnl-2012-304037.

34. Gannon, H.S., Zou, T., Kiessling, M.K., Gao, G.F., Cai, D., Choi, P.S., Ivan, A.P.,
Buchumenski, I., Berger, A.C., Goldstein, J.T., et al. (2018). Identification of ADAR1
adenosine deaminase dependency in a subset of cancer cells. Nat. Commun. *9*, 5450.
10.1038/s41467-018-07824-4.

35. Kung, C.-P., Cottrell, K.A., Ryu, S., Bramel, E.R., Kladney, R.D., Bao, E.A., Freeman, E.C.,
Sabloak, T., Maggi Jr., L., and Weber, J.D. (2021). Evaluating the therapeutic potential of
ADAR1 inhibition for triple-negative breast cancer. Oncogene *40*, 189–202.
10.1038/s41388-020-01515-5.

36. Ota, H., Sakurai, M., Gupta, R., Valente, L., Wulff, B.-E., Ariyoshi, K., Iizasa, H., Davuluri,
R.V., and Nishikura, K. (2013). ADAR1 forms a complex with Dicer to promote microRNA
processing and RNA-induced gene silencing. Cell *153*, 575–589.
10.1016/j.cell.2013.03.024.

37. Freund, E.C., Sapiro, A.L., Li, Q., Linder, S., Moresco, J.J., Yates, J.R., and Li, J.B. (2020).
Unbiased Identification of trans Regulators of ADAR and A-to-I RNA Editing. Cell Rep. *31*,
107656. 10.1016/j.celrep.2020.107656.

38. Sakurai, M., Shiromoto, Y., Ota, H., Song, C., Kossenkov, A.V., Wickramasinghe, J.,
Showe, L.C., Skordalakes, E., Tang, H.-Y., Speicher, D.W., et al. (2017). ADAR1 controls
apoptosis of stressed cells by inhibiting Staufen1-mediated mRNA decay. Nat. Struct. Mol.
Biol. *24*, 534–543. 10.1038/nsmb.3403.

39. Jimeno, S., Prados-Carvajal, R., Fernández-Ávila, M.J., Silva, S., Silvestris, D.A., Endara-
Coll, M., Rodríguez-Real, G., Domingo-Prim, J., Mejías-Navarro, F., Romero-Franco, A., et
al. (2021). ADAR-mediated RNA editing of DNA:RNA hybrids is required for DNA double
strand break repair. Nat. Commun. *12*, 5512. 10.1038/s41467-021-25790-2.

40. Hong, H., An, O., Chan, T.H.M., Ng, V.H.E., Kwok, H.S., Lin, J.S., Qi, L., Han, J., Tay,
D.J.T., Tang, S.J., et al. (2018). Bidirectional regulation of adenosine-to-inosine (A-to-I) RNA
editing by DEAH box helicase 9 (DHX9) in cancer. Nucleic Acids Res. *46*, 7953–7969.
10.1093/nar/gky396.

41. Rajendren, S., Manning, A.C., Al-Awadi, H., Yamada, K., Takagi, Y., and Hundley, H.A.
(2018). A protein–protein interaction underlies the molecular basis for substrate recognition
by an adenosine-to-inosine RNA-editing enzyme. Nucleic Acids Res. *46*, 9647–9659.
10.1093/nar/gky800.

42. Sapiro, A.L., Freund, E.C., Restrepo, L., Qiao, H.-H., Bhate, A., Li, Q., Ni, J.-Q., Mosca,
T.J., and Li, J.B. (2020). Zinc Finger RNA-Binding Protein Zn72D Regulates ADAR-
Mediated RNA Editing in Neurons. Cell Rep. *31*, 107654. 10.1016/j.celrep.2020.107654.

43. Shamay-Ramot, A., Khermesh, K., Porath, H.T., Barak, M., Pinto, Y., Wachtel, C.,
Zilberberg, A., Lerer-Goldshtein, T., Efroni, S., Levanon, E.Y., et al. (2015). Fmrp Interacts
with Adar and Regulates RNA Editing, Synaptic Density and Locomotor Activity in Zebrafish.
PLOS Genet. *11*, e1005702. 10.1371/journal.pgen.1005702.

44. Lam, S.S., Martell, J.D., Kamer, K.J., Deerinck, T.J., Ellisman, M.H., Mootha, V.K., and
Ting, A.Y. (2015). Directed evolution of APEX2 for electron microscopy and proximity
labeling. Nat. Methods *12*, 51–54. 10.1038/nmeth.3179.

45. Orre, L.M., Vesterlund, M., Pan, Y., Arslan, T., Zhu, Y., Fernandez Woodbridge, A., Frings,
O., Fredlund, E., and Lehtiö, J. (2019). SubCellBarCode: Proteome-wide Mapping of Protein
Localization and Relocalization. Mol. Cell *73*, 166-182.e7. 10.1016/j.molcel.2018.11.035.

46. Poulsen, H., Nilsson, J., Damgaard, C.K., Egebjerg, J., and Kjems, J. (2001). CRM1
879   mediates the export of ADAR1 through a nuclear export signal within the Z-DNA binding
880   domain. Mol. Cell. Biol. *21*, 7862–7871. 10.1128/MCB.21.22.7862-7871.2001.

881   47. Strehblow, A., Hallegger, M., and Jantsch, M.F. (2002). Nucleocytoplasmic distribution of
882   human RNA-editing enzyme ADAR1 is modulated by double-stranded RNA-binding
883   domains, a leucine-rich export signal, and a putative dimerization domain. Mol. Biol. Cell *13*,
884   3822–3835. 10.1091/mbc.e02-03-0161.

885   48. Aktaş, T., Avşar Ilık, İ., Maticzka, D., Bhardwaj, V., Pessoa Rodrigues, C., Mittler, G.,
886   Manke, T., Backofen, R., and Akhtar, A. (2017). DHX9 suppresses RNA processing defects
887   originating from the Alu invasion of the human genome. Nature *544*, 115–119.
888   10.1038/nature21715.

889   49. Iordanov, M.S., Paranjape, J.M., Zhou, A., Wong, J., Williams, B.R., Meurs, E.F., Silverman,
890   R.H., and Magun, B.E. (2000). Activation of p38 mitogen-activated protein kinase and c-Jun
891   NH(2)-terminal kinase by double-stranded RNA and encephalomyocarditis virus:
892   involvement of RNase L, protein kinase R, and alternative pathways. Mol. Cell. Biol. *20*,
893   617–627. 10.1128/MCB.20.2.617-627.2000.

894   50. Karasik, A., Jones, G.D., DePass, A.V., and Guydosh, N.R. (2021). Activation of the antiviral
895   factor RNase L triggers translation of non-coding mRNA sequences. Nucleic Acids Res. *49*,
896   6007–6026. 10.1093/nar/gkab036.

897   51. Burke, J.M., Gilchrist, A.R., Sawyer, S.L., and Parker, R. (2021). RNase L limits host and
898   viral protein synthesis via inhibition of mRNA export. Sci. Adv. *7*, eabh2479.
899   10.1126/sciadv.abh2479.

900   52. Nakajima, T., Uchida, C., Anderson, S.F., Lee, C.G., Hurwitz, J., Parvin, J.D., and
901   Montminy, M. (1997). RNA helicase A mediates association of CBP with RNA polymerase II.
902   Cell *90*, 1107–1112. 10.1016/s0092-8674(00)80376-1.

903   53. Cristini, A., Groh, M., Kristiansen, M.S., and Gromak, N. (2018). RNA/DNA Hybrid
904   Interactome Identifies DXH9 as a Molecular Player in Transcriptional Termination and R-
905   Loop-Associated DNA Damage. Cell Rep. *23*, 1891–1905. 10.1016/j.celrep.2018.04.025.

906   54. Lee, T., and Pelletier, J. (2016). The biology of DHX9 and its potential as a therapeutic
907   target. Oncotarget *7*, 42716–42739. 10.18632/oncotarget.8446.

908   55. Ren, X., Wang, D., Zhang, G., Zhou, T., Wei, Z., Yang, Y., Zheng, Y., Lei, X., Tao, W.,
909   Wang, A., et al. (2023). Nucleic DHX9 cooperates with STAT1 to transcribe interferon-
910   stimulated genes. Sci. Adv. *9*, eadd5005. 10.1126/sciadv.add5005.

911   56. Zhang, Z., Yuan, B., Lu, N., Facchinetti, V., and Liu, Y.-J. (2011). DHX9 Pairs with IPS-1 To
912   Sense Double-Stranded RNA in Myeloid Dendritic Cells. J. Immunol. Baltim. Md 1950 *187*,
913   4501–4508. 10.4049/jimmunol.1101307.

914   57. Harding, H.P., Novoa, I., Zhang, Y., Zeng, H., Wek, R., Schapira, M., and Ron, D. (2000).
915   Regulated Translation Initiation Controls Stress-Induced Gene Expression in Mammalian
916   Cells. Mol. Cell *6*, 1099–1108. 10.1016/S1097-2765(00)00108-8.

58. Pakos-Zebrucka, K., Koryga, I., Mnich, K., Ljujic, M., Samali, A., and Gorman, A.M. (2016).
The integrated stress response. EMBO Rep. *17*, 1374–1395. 10.15252/embr.201642195.

59. Zamanian-Daryoush, M., Mogensen, T.H., DiDonato, J.A., and Williams, B.R.G. (2000). NF-
κB Activation by Double-Stranded-RNA-Activated Protein Kinase (PKR) Is Mediated through
NF-κB-Inducing Kinase and IκB Kinase. Mol. Cell. Biol. *20*, 1278–1290.

60. Lemaire, P.A., Anderson, E., Lary, J., and Cole, J.L. (2008). Mechanism of PKR Activation
by dsRNA. J. Mol. Biol. *381*, 351–360. 10.1016/j.jmb.2008.05.056.

61. Kato, H., Takeuchi, O., Mikamo-Satoh, E., Hirai, R., Kawai, T., Matsushita, K., Hiiragi, A.,
Dermody, T.S., Fujita, T., and Akira, S. (2008). Length-dependent recognition of double-
stranded ribonucleic acids by retinoic acid-inducible gene-I and melanoma differentiation-
associated gene 5. J. Exp. Med. *205*, 1601–1610. 10.1084/jem.20080091.

62. Sadler, A.J., Latchoumanin, O., Hawkes, D., Mak, J., and Williams, B.R.G. (2009). An
Antiviral Response Directed by PKR Phosphorylation of the RNA Helicase A. PLOS Pathog.
5, e1000311. 10.1371/journal.ppat.1000311.

63. Fuchsová, B., and Hozák, P. (2002). The Localization of Nuclear DNA Helicase II in
Different Nuclear Compartments Is Linked to Transcription. Exp. Cell Res. *279*, 260–270.
10.1006/excr.2002.5617.

64. Jeffrey, I.W., Kadereit, S., Meurs, E.F., Metzger, T., Bachmann, M., Schwemmle, M.,
Hovanessian, A.G., and Clemens, M.J. (1995). Nuclear localization of the interferon-
inducible protein kinase PKR in human cells and transfected mouse cells. Exp. Cell Res.
218, 17–27. 10.1006/excr.1995.1126.

65. García, M.A., Gil, J., Ventoso, I., Guerra, S., Domingo, E., Rivas, C., and Esteban, M.
(2006). Impact of protein kinase PKR in cell biology: from antiviral to antiproliferative action.
Microbiol. Mol. Biol. Rev. MMBR *70*, 1032–1060. 10.1128/MMBR.00027-06.

66. Kim, Y., Lee, J.H., Park, J.-E., Cho, J., Yi, H., and Kim, V.N. (2014). PKR is activated by
cellular dsRNAs during mitosis and acts as a mitotic regulator. Genes Dev. *28*, 1310–1322.
10.1101/gad.242644.114.

67. Kim, J.I., Nakahama, T., Yamasaki, R., Costa Cruz, P.H., Vongpipatana, T., Inoue, M.,
Kanou, N., Xing, Y., Todo, H., Shibuya, T., et al. (2021). RNA editing at a limited number of
sites is sufficient to prevent MDA5 activation in the mouse brain. PLoS Genet. *17*,
e1009516. 10.1371/journal.pgen.1009516.

68. Hu, S.-B., Heraud-Farlow, J., Sun, T., Liang, Z., Goradia, A., Taylor, S., Walkley, C.R., and
Li, J.B. (2023). ADAR1p150 prevents MDA5 and PKR activation via distinct mechanisms to
avert fatal autoinflammation. Mol. Cell *0*. 10.1016/j.molcel.2023.09.018.

69. Lee, T., and Pelletier, J. (2016). The biology of DHX9 and its potential as a therapeutic
target. Oncotarget *7*, 42716–42739. 10.18632/oncotarget.8446.

70. Roulois, D., Loo Yau, H., Singhania, R., Wang, Y., Danesh, A., Shen, S.Y., Han, H., Liang,
G., Jones, P.A., Pugh, T.J., et al. (2015). DNA-Demethylating Agents Target Colorectal

Cancer Cells by Inducing Viral Mimicry by Endogenous Transcripts. Cell *162*, 961–973.
10.1016/j.cell.2015.07.056.

71. Chen, R., Ishak, C.A., and De Carvalho, D.D. (2021). Endogenous Retroelements and the
Viral Mimicry Response in Cancer Therapy and Cellular Homeostasis. Cancer Discov. *11*,
2707–2725. 10.1158/2159-8290.CD-21-0506.

72. Jansz, N., and Faulkner, G.J. (2021). Endogenous retroviruses in the origins and treatment
of cancer. Genome Biol. *22*, 147. 10.1186/s13059-021-02357-4.

73. Wu, Q., Nie, D.Y., Ba-alawi, W., Ji, Y., Zhang, Z., Cruickshank, J., Haight, J., Ciamponi,
F.E., Chen, J., Duan, S., et al. (2022). PRMT inhibition induces a viral mimicry response in
triple-negative breast cancer. Nat. Chem. Biol. *18*, 821–830. 10.1038/s41589-022-01024-4.

74. Zhou, X., Singh, M., Sanz Santos, G., Guerlavais, V., Carvajal, L.A., Aivado, M., Zhan, Y.,
Oliveira, M.M.S., Westerberg, L.S., Annis, D.A., et al. (2021). Pharmacologic Activation of
p53 Triggers Viral Mimicry Response Thereby Abolishing Tumor Immune Evasion and
Promoting Antitumor Immunity. Cancer Discov. *11*, 3090–3105. 10.1158/2159-8290.CD-20-
1741.

75. Sheng, W., LaFleur, M.W., Nguyen, T.H., Chen, S., Chakravarthy, A., Conway, J.R., Li, Y.,
Chen, H., Yang, H., Hsu, P.-H., et al. (2018). LSD1 Ablation Stimulates Anti-tumor Immunity
and Enables Checkpoint Blockade. Cell *174*, 549-563.e19. 10.1016/j.cell.2018.05.052.

76. Goel, S., DeCristo, M.J., Watt, A.C., BrinJones, H., Sceneay, J., Li, B.B., Khan, N.,
Ubellacker, J.M., Xie, S., Metzger-Filho, O., et al. (2017). CDK4/6 inhibition triggers anti-
tumour immunity. Nature *548*, 471–475. 10.1038/nature23465.

77. Sanjana, N.E., Shalem, O., and Zhang, F. (2014). Improved vectors and genome-wide
libraries for CRISPR screening. Nat. Methods *11*, 783–784. 10.1038/nmeth.3047.

78. Hayer, A., Shao, L., Chung, M., Joubert, L.-M., Yang, H.W., Tsai, F.-C., Bisaria, A., Betzig,
E., and Meyer, T. (2016). Engulfed cadherin fingers are polarized junctional structures
between collectively migrating endothelial cells. Nat. Cell Biol. *18*, 1311–1323.
10.1038/ncb3438.

79. Ong, J.Y. (2021). Synonymous Mutation Generator: a web tool for designing RNAi-resistant
sequences. Preprint at bioRxiv, 10.1101/2021.01.02.425100 10.1101/2021.01.02.425100.

80. Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev,
S., Jenkins, S.L., Jagodnik, K.M., Lachmann, A., et al. (2016). Enrichr: a comprehensive
gene set enrichment analysis web server 2016 update. Nucleic Acids Res. *44*, W90–W97.
10.1093/nar/gkw377.

81. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550. 10.1186/s13059-014-
0550-8.

82. Zhu, A., Ibrahim, J.G., and Love, M.I. (2019). Heavy-tailed prior distributions for sequence
count data: removing the noise and preserving large differences. Bioinforma. Oxf. Engl. *35*,
2084–2092. 10.1093/bioinformatics/bty895.

83. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., et
al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. Innov.
Camb. Mass *2*, 100141. 10.1016/j.xinn.2021.100141.

84. Jin, Y., Tam, O.H., Paniagua, E., and Hammell, M. (2015). TEtranscripts: a package for
including transposable elements in differential expression analysis of RNA-seq datasets.
Bioinforma. Oxf. Engl. *31*, 3593–3599. 10.1093/bioinformatics/btv422.

85. Cottrell, K.A., Soto-Torres, L., Dizon, M.G., and Weber, J.D. (2021). 8-Azaadenosine and 8-
Chloroadenosine are not Selective Inhibitors of ADAR. Cancer Res. Commun. *1*, 56–64.
10.1158/2767-9764.CRC-21-0027.

86. Marcotte, R., Sayad, A., Brown, K.R., Sanchez-Garcia, F., Reimand, J., Haider, M.,
Virtanen, C., Bradner, J.E., Bader, G.D., Mills, G.B., et al. (2016). Functional Genomic
Landscape of Human Breast Cancer Drivers, Vulnerabilities, and Resistance. Cell *164*,
293–309. 10.1016/j.cell.2015.11.062.

87. Kosinski, M., Biecek, P., and Chodor, W. (2022). RTCGA: The Cancer Genome Atlas Data
Integration. Version 1.28.0 (Bioconductor version: Release (3.16)).
10.18129/B9.bioc.RTCGA 10.18129/B9.bioc.RTCGA.

88. Kassambara, A., Kosinski, M., Biecek, P., and Fabian, S. (2020). survminer: Drawing
Survival Curves using "ggplot2." Version 0.4.8.

89. Ghandi, M., Huang, F.W., Jané-Valbuena, J., Kryukov, G.V., Lo, C.C., McDonald, E.R.,
Barretina, J., Gelfand, E.T., Bielski, C.M., Li, H., et al. (2019). Next-generation
characterization of the Cancer Cell Line Encyclopedia. Nature *569*, 503–508.
10.1038/s41586-019-1186-3.

90. DepMap 22Q4 Public (2022). (figshare). 10.6084/m9.figshare.21637199.v2
10.6084/m9.figshare.21637199.v2.

1018

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

a

| | shADAR1 | |
| shSCR | shDHX9-3 |

EV ADAR1-p110 ADAR1-p150 EV ADAR1-p110 ADAR1-p150

DHX9
ADAR1
p-PKR
Total PKR
p-eIF2α
Total eIF2α
c-PARP
β-tubulin
Total Protein

b

shADAR1

EV ADAR1-p110 ADAR1-p150

shSCR

shDHX9-3

c

Relative p-PKR/PKR Abundance

***

shSCR EV
shSCR ADAR1-p110
shSCR ADAR1-p150
shDHX9-3 EV
shDHX9-3 ADAR1-p110
shDHX9-3 ADAR1-p150

d

Relative Foci Area

***
***
***

shSCR EV
shSCR ADAR1-p110
shSCR ADAR1-p150
shDHX9-3 EV
shDHX9-3 ADAR1-p110
shDHX9-3 ADAR1-p150