Tile: Repetitive Spreading Depolarization induces the activation of cell differentiation, synaptic plasticity and neuroprotective pathways.

Michela Dell'Orco [a], Jordan E. Weisend [a], Nora I. Perrone-Bizzozero [a], Andrew P. Carlson [b], Russell A. Morton [a] , David N Linsenbardt [a] & C. William Shuttleworth [a].

[a] Department of Neurosciences, [b] Department of Neurosurgery; University of New Mexico School of Medicine, Albuquerque, New Mexico, 87131, USA.

Correspondence to:

C. William Shuttleworth, Ph.D.

Department of Neurosciences

1 University of New Mexico, MSC08 4740

Albuquerque, NM 87131

USA

Email:bshuttleworth@salud.unm.edu

Phone: (505) 272 4290

KEYWORDS: spreading depression, preconditioning, gene expression, pathway analysis, brain injury, stroke.

Abstract

Spreading depolarization (SD) is a slowly propagating wave of profound depolarization that sweeps through

cortical tissue. While much emphasis has been placed on the damaging consequences of SD, it is possible

that SDs also activate compensatory pathways related to cell survival and plasticity. The present study

aimed to provide an unbiased assessment of gene expression changes following SD, as well as novel

molecular networks associated with these transcriptional alterations. SD clusters were induced with either

focal application of KCl or with optogenetic stimulation in healthy mice, and then 2 hours later cortical

RNA was extracted and sequenced. SD's significantly increased the expression of 21 genes - no genes were

significantly downregulated. Notable top hits included the immediate early genes FOS, ARC, and JUN, the

cell proliferation-related gene DUSP6, the plasticity-related gene HOMER1, and inflammation related

genes PTGS2, EGR2, and NR4A1. Pathway analysis identified the recruitment of genes associated with

axonogenesis, branching, neuritogenesis, and dendritic growth, as well as inhibition of pathways associated

with cell death, apoptosis, and neuronal degeneration. These results identify the induction of plasticity

and/or circuit modification as an important consequence of SDs in healthy tissue, as well as specific gene

targets and pathways amenable to manipulation in follow up studies.

Introduction

Spreading Depolarization (SD) is a slowly propagating wave of extended depolarization that is followed by suppression of neural activity, first described by Leao (Leao 1944). SD is implicated as a mechanism of migraine with aura (Pietrobon and Moskowitz 2014; Charles and Brennan 2009) and otherwise healthy brain is able to fully recover from SD (Nedergaard and Hansen 1988). However, the long duration of depolarization (up to a minute or more) can lead to severe energy depletion (Dreier et al. 2017; Lindquist and Shuttleworth 2014), particularly when coupled with neurovascular disruptions in ischemic brain (Hartings et al. 2017; Ayata and Lauritzen 2015; Dreier 2011). In addition, transient extracellular glutamate surges caused by SD can be sufficient to cause neuronal injury in metabolically compromised brain tissue (Aiba et al. 2012; Hinzman et al. 2015). Recent work has identified SDs as contributors to tissue loss following stroke, trauma, and other acute brain injuries (Dreier et al. 2017; Lauritzen et al. 2011) and initial efforts have begun targeting SD in efforts to improve outcomes in brain injury (Helbok et al. 2020; Carlson et al. 2018).

In contrast to the large body of literature characterizing electrophysiological and vascular responses to SD, only a relatively small number of studies have examined SD-induced gene expression changes and their relationship to outcomes in different experimental and clinical conditions. Initial studies identified increases in brain derived neurotrophic factor (BDNF), c-Fos and heat shock protein 70 (HSP70) (Dietrich et al. 2000; Kokaia et al. 1993; Karikó et al. 1998; Matsushima, Hogan, and Hakim 1996; Rangel et al. 2001). Subsequent studies showed increased expression of pro-inflammatory and apoptotic genes, including tumor necrosis factor (TNF), C-C motif chemokine ligand 2 (CCL2), interleukin-1 (IL-1) and -6, and cyclooxygenase-2 (COX-2), and B-cell lymphoma 2 (BCL-2) (Eising et al. 2017; Jander et al. 2001; Kaido et al. 2012; Takizawa et al. 2020). While most of these studies focused predetermined genes, a relatively recent microarray study has applied gene ontology (GO) and pathway analysis (PA) to the study of SD in rats (Sintas et al. 2017). Pathway analysis allows the identification of altered cellular and molecular

pathways from data obtained from high-throughput technologies; leading to the identification of novel sets

of related genes that are altered in different conditions (García-Campos, Espinal-Enríquez, and Hernández-

Lemus 2015). Sintas and colleagues focused their analysis on SD's influence on pathways linked to

oxidative stress and cellular injury, but less is known of other cellular adaptations that may occur. A lack

of comprehensive transcriptomic analysis has limited the understanding of potentially beneficial pathways

that could be activated following SD.

In the current study, we provide the first unbiased RNA sequencing (RNAseq) analysis following SD. We

applied pathway analysis approaches to uncover candidate networks and novel genes altered by SD. To

increase generalizability, we combined data from the two most common methods of SD initiation (focal

depolarization with either topical KCl administration or optogenetic stimulation) and generated clusters of

SDs that are analogous to patterns of SDs that have been recorded from injured human brain. The results

identify the upregulation of multiple molecular pathways that could regulate potentially beneficial effects

of SD, such as neuritogenesis and synaptic plasticity.

RESULTS

1) Identification of gene expression induced by repetitive spreading depolarizations

Multiple SDs (4 over a period of 2 hours) were generated to match clusters of SDs in clinical settings (Dreier

et al. 2017) and this procedure has previously been shown to induce robust transcriptional changes (Sintas

et al. 2017; Urbach, Bruehl, and Witte 2006; Takizawa et al. 2020). Thirty minutes after the last SD or sham

stimulus, the neocortex was removed, and RNA extracted from tissue subjected to SD or shams for

subsequent RNA sequencing (see Materials and Methods and **Supplementary Figure 1**).

As both focal KCl and optogenetic stimulation are routinely used in studies of SD, data from these two SD-

stimulation methods were combined for differential expression (DE) analysis here to increase

generalizability. For the combined DE analysis, we performed DESeq2 analysis (Ge, Son, and Yao 2018)

and identified 26 genes that were also significant after false discovery rate correction (FDR <0.05) when

comparing the SD group with the sham tissues (**Figure 1, red, FDR <0.05**).

Consistent with previous studies (Kaido et al. 2012; Rangel et al. 2001; Kokaia et al. 1993; Dietrich et al.

2000), we found significant changes intermediate early genes ARC, FOS and JUN proto-oncogenes, and

early growth response (EGR) genes. Among other DEGs with significantly increased levels were

adrenoceptor beta 1 (ADRB1), dual specificity phosphatase (DUSP) gene family, the nuclear receptor

subfamily 4 group A member 1 (NR4A1), and prostaglandin-endoperoxide synthase 2 (PTGS2).

2) Molecular networks regulated by repetitive SDs

We next used Ingenuity Pathway Analysis (IPA) to identify molecular networks and biological pathways

regulated by DEGs. This information was then used to obtain a prediction of physiological and pathological

brain processes altered by SD. **Figure 2A** and **B** show the top two scored networks (score 63 and 36)

generated by IPA with the combined DEGs upregulated in the SD group as compared to sham animals. The

intensity of the color red for each molecule is proportional to the fold change (FC) of the labeled gene. Full

arrows indicate a direct relationship between molecules, while dotted arrows indicate a predicted indirect

relationship. Orange arrows indicate the predicted activation of a gene, while yellow indicates predicted

inhibition. These networks implicated SD target genes associated with cell proliferation and synaptic

activity, including cAMP responsive element binding protein 1 (CREB1), mitogen-activated protein kinase

1 and 3 (MAPK1 and MAPK3), and HOMER1 (**Figure 2B**).

Predictions of SD-induced changes associated with known disease states and functions are shown in **Figure**

2C and they include: neurological diseases, psychological disorders, nervous system development and

function, tissue morphology, and tissue development. The complete list of diseases and functions and the

genes associated with each pathway are reported in **Supplementary Table 2**. **Figure 2C** also shows the

number of genes associated with each pathway and associated p value range. They are shown together so

that different strengths of the network elements can be compared. A complete list of networks and

associated DEGs is reported in **Supplementary Table 3.**

Figure 3 illustrates pathways related to cellular functions induced by repetitive spreading depolarization.

The top common pathways identified include: cellular assembly and organization, cellular function and

maintenance, and cellular development and proliferation (**Supplementary Table 4**). The heatmaps in

Figure 3 show the p value and the activation score for each pathway, and a graphical representation of the

cellular functions and genes involved, respectively. As shown in Figure 2, each gene expression FC is

proportional to their color intensity; orange arrows and pathways indicate a predicted activation, whereas

blue indicates increased levels of genes which lead to inhibition of the pathway. IPA analysis predicted an

activation of axonogenesis, axon branching, dendritic growth and branching and growth of neurites, based

on increased levels of genes such as VGF, FOS, PTGS2, JUN, BDNF, gamma-aminobutyric acid type A

receptor subunit beta3 (GABRB3), activating transcription factor 3 (ATF3), and cyclin dependent kinase

like 3 (CDKL3). We also found predicted activation of pathways regulating neuronal development, long-

term potentiation, and neurotransmission due to increased expression of genes such as FOS, PTGS2, EGR2,

BDNF, JUN, KCNJ2, ARC, and HOMER1.

Interestingly, we also found predicted inhibition of neuronal cell death, apoptosis and neuronal degeneration

pathways, together with increased survival of cerebral cortex cells, regeneration of axons, and cell viability,

due to increased expression of genes such as FOS, BDNF, ATF3, JUNB, DUSP1, neuronal PAS domain

protein 4 (NPAS4), nuclear factor interleukin 3 regulated (NFIL3), and SERPINE1.

3) Comparison between contralateral and ipsilateral hemispheres

Many studies utilize the contralateral neocortex as a convenient comparison tissue for DEG studies

(Volobueva et al. 2023), as SD does not directly propagate across the corpus collosum. However, even if

SD does not directly propagate through the contralateral cortex, the profound and widespread disruption of

neuronal network activity produced by SD might be expected to have some impact on neuronal activity and

gene expression in the contralateral hemisphere. The assumption that contralateral tissue is equivalent to

tissue from sham animals for DE analysis has not previously been tested.

We first performed DE analysis between these 2 tissues (i.e., contralateral hemispheres in animals exposed

to SD in the ipsilateral hemisphere vs. sham tissues). As showed in Supplementary Figure 2 no genes were

significant after FDR correction, confirming a high degree of similarity between gene expression in these

tissues.

We next performed the same analyses described above (**Figure 1**), but comparing tissue exposed to SD

with the contralateral cortex in the same animals, rather than shams. Of the 12893 mapped 33 significantly

upregulated after FDR correction (FDR <0.05) (**Figure 4A, red, FDR <0.05**). IPA analysis confirmed a

high similarity between the sham and contralateral cortex when used as control compared to the SD group.

Figure 4B shows the top scored network (score 65) generated from the SD vs contralateral DEGs list and

the top diseases and function are reported in **Figure 4B** and **Supplementary Table 2 (compare with**

Figure 2). Likewise, **Figure 5** shows cellular and molecular pathways that are comparable with DEG

analysis with sham tissues (compared with Figure 3)**.** These results imply that, at least at the level of gene

expression changes at 2hours, the ipsilateral hemisphere that is not directly invaded by SD can likely serve

as an unaffected control.

4) SD target genes control cell differentiation and proliferation, synaptic plasticity and stress

responses

To validate the predicted activation of neuroprotective pathways, we analyzed the expression levels of

selected SD target genes by qRT-PCR. Based on the IPA results above, we focused on genes regulating 3

cellular functions: 1) cell differentiation and proliferation, 2) synaptic plasticity and 3) stress responses.

qPCR was performed on the same RNA that was subjected to the RNA-seq analyses, as well as 3 additional

animals for each group. As shown in **Figure 6**, multiple SDs resulted in significant increases in the

expression of genes activating cell proliferation and differentiation such as BNDF, DUPS6, FOS and JUNB

(**Figure 6**). Considering the predicted activation of neurons and long-term potentiation, as well the

increased axonogenesis, growth of neurites, and axon and neurites branching, we then analyzed the

expression of genes known for their regulation of synaptic plasticity. As shown in **Figure 6** we found

significantly increased expression of Homer1a and ARC. Finally, we examined levels of PTGS2, NR4A1,

and EGR2 as genes regulating stress and inflammatory responses.

5) Target genes showed differential expression depending on the distance from the SD initiation site

Lastly, we tested whether the levels of gene expression varied with distance from the SD initiation site. In

this set of studies, KCl was used as the stimulus, as the site of initiation is clearly restricted to small zone

of the application of KCl through the burrhole. Thus, two hours after onset of the initial SD, RNA was

extracted from either total contralateral cortex, or from 3 different regions of the ipsilateral hemisphere: 1)

SD initiation site; 2) intermediate site, and 3) remote site (see Methods and **Supplementary Figure 3**). By

qRT-PCR we analyzed the levels of the previously identified target genes (see section 3 above) and

compared these to the levels in the contralateral hemisphere of SHAM animals. For the majority of the

genes examined (BDNF, DUSP6, PTGS2 ARC, and Homer 1a), highest expression was detected in the

intermediate area (>3mm from the initiation site; **Figure 7**) while FOS and JUNB genes showed

significantly increased expression in the remote site (>6 mm from the initiation site). Interestingly, when

analyzing Homer1a levels, we found a decrease in their expression at the SD initiation site (**Figure 7**) when

compared to the contralateral and intermediate sites (**Figure 7**). The difference between the contralateral

and remote sites for NR4A1 did not reach statistical significance (p value=0.09) and EGR2 did not show

differential expression across regions (**Supplementary Figure 3**).

Discussion

As noted above (Introduction) previous work has identified molecular pathways linking SD to inflammation and apoptosis, but there is interest in determining a wider spectrum of possible consequences of SD, including possible contributions to plasticity and repair (Shuttleworth et al. 2020). Previous studies have shown that SD leads to the increased synaptic activity (Sadowska et al. 2021; Kriz and Lalancette-Hébert 2009) as well as adult neurogenesis (Urbach et al. 2017; Urbach, Bruehl, and Witte 2006; Carmichael and Chesselet 2002) and preconditioning (Matsushima et al. 1998; Yanamoto et al. 2004; Shen et al. 2016). However, there has been limited investigation of specific molecular pathways underlying these processes. Here, through RNA-seq and pathway analysis, we showed enrichment in genes regulating neurological disorders, psychological disorders, and behavior (**Figure 2 and 4**). Furthermore, there were significant increases in predicted pathways involving processes expected to be important for brain plasticity and repair, including neuronal differentiation, and neurite growth and branching (**Figures 3 and 5**).

Prior exposure to SD has been shown to confer protection against subsequent ischemic challenges, and increases in BDNF have previously been implicated in the mechanism (Matsushima et al. 1998; Yanamoto et al. 2004). Consistent with previous work (Yanamoto et al. 2004; Rangel et al. 2001; Kokaia et al. 1993; Matsushima et al. 1998), BDNF was one of the genes significantly increased by both KCl and optogenetic stimuli (**Figure 6 and 7**) and was identified in SD-induced pathways related to neuronal differentiation (**Figure 3 and 5**).

This study identified novel gene expression changes that could be used to test whether clusters of SD may enhance plasticity. Importantly, the upregulation of SD specific target genes can lead to predicted inhibition of pathways controlling neuronal cell death, apoptosis, and neuronal degeneration, together with increased survival of cerebral cortex cells, regeneration of axons, and cell viability. The increased expression of genes such as FOS, BDNF, JUNB, and neuronal PAS domain protein 4 (NPAS4), (**Figure 3 and 5),** lead to higher rate of cellular development and differentiation compared to cell death and apoptosis. These genes have

been previously associated to increased neuronal tolerance for ischemic stroke (Yanamoto et al. 2004;

Matsushima et al. 1998; Takahashi et al. 2021).

Finally, we found increased expression of the mRNA for the plasticity-associated protein ARC, which

mediates AMPA receptors endocytosis and is required for protein synthesis dependent LTP and memory

formation (Chowdhury et al. 2006). Consistent with our findings, prior work in brain slices and rodent

cortex has reported LTP-like synaptic strengthening following SD (Footitt and Newberry 1998; Berger et

al. 2008), and it will be of interest to determine whether the time course of associated gene expression

changes correspond to variations in physiological recordings and how the expression of these genes change

under different metabolic conditions surrounding ischemic infarcts.

Although our results predict activation of neuroprotective pathways, they are also consistent with prior

reports of robust increases of the pro-inflammatory mediator COX2 (Takizawa et al. 2020; Kaido et al.

2012; Volobueva et al. 2023). We did not see significant increases in some other inflammatory mediators

(e.g., TNF-α, IL-1β, IL6), perhaps because the time course for peak increases in these mediators is later

than the 2-hour time point used in the present study (Takizawa et al. 2020). This time point was chosen

based on 1) prior expression analysis confirming peak increases at 2 hours for most genes of interest

(Kokaia et al. 1993; Karikó et al. 1998; Rangel et al. 2001) and 2) clinical recordings establishing clusters

of SDs (usually defined as three or more events within 2 hours) as being clinically significant (Hartings et

al. 2020; Dreier et al. 2006). Prior work (Takizawa et al. 2020) has shown that some inflammatory

mediators can be detected following a single SD (measured at 1 hour), while others peak at 24 h (Jander et

al. 2001; Matsushima et al. 1998; Volobueva et al. 2023). It remains to be determined which of the genes

seen here may peak earlier or at later times points.

To evaluate differences in gene expression at initiation sites vs. more remote sites, we micro-dissected

regions to compare differential expression of a number of key genes at the initiation, intermediate, and

remote sites using RT-qPCR analyses (**Figure 7 and Supplemental Figure 3**). We found that most genes

examined had largest increases at the intermediate site, >3 mm from the initiation site. If this is generally

true for other genes in the RNAseq data set, then it suggests that the initiation site with KCl stimulation

might not be a major contributor to DEG changes observed here. It is also possible that sites of SD initiation

and propagation could be different in SDs generated "spontaneously"/ naturalistically in patients or animal

models of migraine or brain injury. Thus, an SD initiation site required an area of coordinated

depolarization (Tang et al. 2014) from ischemia or synaptic stimulation with inadequate perfusion (von

Bornstädt et al. 2015). Gene expression changes in this environment may be quite different from changes

seen in remote areas into which the SD propagates. Spatial genomics approaches (Ibarra-Lopez et al. 2022)

will be useful to further investigate the distribution of changes of larger gene sets with distance from SD

initiation, and with respect to differential vascular perfusion, for example.

While most validated genes showed higher expression levels in the intermediate or remote sites and no

change in the initiation site, Homer1a showed a significant decrease in the SD initiation site (**Figure 7**).

Homer1 scaffolding protein controls the insertion of postsynaptic glutamate receptors (Ango et al. 2002;

Peng Luo et al. 2012). Homer1a is increased by synaptic activity through the MAP kinase pathway. This

type of regulation creates a negative feedback loop in which activity-dependent Homer1a inhibits

Homer1b/c ability to scaffold and diminishes synaptic strength by modulation of dendritic spine

morphology and AMPA/NMDA receptor activity (Bottai et al. 2002; Sato, Suzuki, and Nakanishi 2001).

This loop has been associated with TBI and epilepsy, particularly, and is thought to block postsynaptic

Ca^{2+}-mediated excitotoxicity (Cavarsan et al. 2012; P. Luo et al. 2014). Here, we found significantly

decreased levels of Homer1a in the initiation site, while expression was significantly increased in the

intermediate site. These results may suggest a negative feedback mechanism to reduce downstream

excitotoxicity, possibly reducing the number of glutamate postsynaptic receptors after repeated SDs.

Conclusions

Spreading depolarization (SD) is common in acute brain injury and contributes to expansion of lesions.

The potential role of SD in the recovery of surviving brain tissue is not yet clear. For the first time we

combined RNA-seq with predicted pathway analysis to identify a new range of molecular targets pathways

that could potentially be activated and contribute to circuit adaptation and recovery after brain **injury**

(Figure 8). While it is very clear that SDs are direct contributors to injury in compromised brain tissue,

our findings suggest that multiple SDs could also activate a range of neuroprotective pathways including

increased cell proliferation, decreased cell death, cell differentiation, and neuronal plasticity (**Figure 2-5**).

Whether or not such pathways make significant contributions to brain recovery and repair requires explicit

testing that can be guided, in part, by modification of key regulatory mediators identified in the present

study.

Materials and Methods

Animal model

All experimental procedures were performed in accordance with the National Institutes of Health Guide for

Care and Use of Laboratory Animals and were approved by the University of New Mexico Health Sciences

Center Institutional Animal Care and Use Committee (IACUC). SDs were induced repetitively (4 SDs in 2

hours at 30 min intervals) in healthy 7-8 week old female mice and SDs were confirmed with intrinsic

optical signal imaging (Supplementary Figure 1). SDs were induced in 6 mice with either focal application

of KCl through a burr hole in C57Bl6J mice (n=3) or with optogenetic stimulation (3 mW for 20 seconds)

in Thy1-ChR2-YFP positive mice (n=3). Thy1-ChR2-YFP breeders were initially purchased from The

Jackson   Laboratory   (B6.Cg-Tg(Thy1-COP4/EYFP)18Gfng/J,   Strain   #:007612,

RRID:IMSR_JAX:007612) and crossed to maintain the line, as previously described (Arenkiel et al. 2007).

With either stimulus, evoked SDs propagated throughout the neocortex ipsilateral to the stimulus, without

crossing to the contralateral hemisphere **(Supplementary Figure 1).**   A matched number of sham controls

was used with focal NaCl administration through a burrhole, in C57Bl/6 mice (n=3), or 0.4 mW for 20

seconds in Thy1-ChR2-YFP positive mice (n=3). Thirty minutes after the last SD, the anesthetized animal

was decapitated and the brain extracted, washed with ice cold PBS and kept on ice.

RNA extraction and qRT-PCR validation

The contralateral and ipsilateral cortices were rapidly dissected from underlying striatum and the

hippocampus and total RNA was extracted using TRIzol (Invitrogen, ThermoFisher, Waltham, MA. USA).

Extractions were either from intact ipsilateral or contralateral cortices (**Figures 2-7**) or from regions ~18

mm^2 rapidly micro-dissected on ice (**Figure 7**). Where indicated, microdissections were made from 3

different regions of the ipsilateral hemisphere: 1) SD initiation site; 2) intermediate site > 3mm from

initiation, 3) remote site >5 mm from initiation (**Supplementary Figure 3**). All collected tissues were then

flash frozen and kept at –80°C until RNA extraction. RNA was quantified using Qubit (Bio-Rad) and its

quality was determined using a NanoDrop 1000 (ThermoFisher), using absorbance at 260, 280 and 230 nm.

Aliquots of 2 μg RNA were sent to Arraystar, Inc. for Illumina paired-end RNA-seq as described below.

To validate gene expression from our RNA-seq data set, separate aliquots containing 1 μg of total RNA

were reverse transcribed using SuperScript II reverse transcriptase (Life Technologies) following the

manufacturer's protocol and qPCR carried out with a CFX96 Touch Real-Time PCR Detection System

using SYBR Green mix (Life Technologies). Primer sequences for qRT-PCR for mouse mRNAs were

obtained from PrimerBank (https://pga.mgh.harvard.edu/primerbank/).   The GAPDH gene (FW

TGTGATGGGTGTGAACCACGAGAA, RV GAGCCCTTCCACAATGCCAAAGTT) was used to

normalize values. As shown in **Supplementary Figure 3**, GAPDH expression itself was unchanged by SD

with either KCl or optogenetic induction of SD. The relative expression of target genes was determined

using the comparative 2−ΔCt method (Livak and Schmittgen 2001).

Values were expressed as means ± SEM. Statistical analysis was performed using GraphPad Prism version

9 (La Jolla, CA, USA). The data were analyzed by unpaired t-test followed by Welch's correction or

analysis of variance (ANOVA) followed by Dunnett's Multiple Comparison tests. Differences were

considered statistically significant when p values were < 0.05 (*p<0.05, **p<0.01, ***p<0.001).

RNA-seq

RNA sequencing analysis was performed by Arraystar Inc. (Rockville, MD). Total RNA (2 µg) was used

to prepare the sequencing library in the following steps: total RNA was enriched by oligo (dT) magnetic

beads (rRNA removed); RNA-seq library was prepared using KAPA Stranded RNA-Seq Library Prep Kit

(Illumina), which incorporates dUTP into the second cDNA strand and renders the RNA-seq library strand-

specific. The completed libraries were qualified with an Agilent 2100 Bioanalyzer and quantified by using

the absolute quantification qPCR method. To sequence the libraries on an Illumina NovaSeq 6000

instrument, the barcoded libraries were mixed, denatured to single stranded DNA in NaOH, captured on

Illumina flow cell, amplified *in situ,* and subsequently sequenced for 150 cycles for both ends on an Illumina

NovaSeq 6000 instrument. Sequence quality was examined using the FastQC software. The trimmed reads

(trimmed 5', 3'-adaptor bases using cutadapt) were aligned to a reference genome (Mouse genome

GRCm38) using Hisat2 software.

The data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus (Edgar,

Domrachev, and Lash 2002) and are accessible through GEO Series accession number GSE214623

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE214623).

Differential expression analysis

The expression levels of known genes and transcripts were calculated using the StringTie and Ballgown

software package (Pertea et al. 2016) and expressed as fragments per kilobase of transcript per million

mapped reads (FPKM). Only genes with minimal counts per million (CPM) in at least one library were kept

for further analysis. Differentially expressed gene (DEG) and transcript analyses were performed using an

unpaired t-test followed by Welch's correction; a fold change (FC) threshold of 1.25 and p value < 0.05

were used together to classify genes and transcripts as differentially expressed. We subsequently performed

DESeq2 analysis (Ge, Son, and Yao 2018) with a false discovery rate (FDR) correction cutoff of 0.05.

Ingenuity Pathway Analysis

Predicted molecular functions and biological pathways enriched in genes differentially expressed after SD

were identified using Ingenuity Pathway software (IPA, Content version: 62089861 (Release Date: 2021-

02-17), Qiagen, https://www.qiagenbioinformatics.com/products/ingenuity-pathway-analysis). The

analysis was set to filter nervous system expressed genes only; reference set: Ingenuity Knowledge Base

(Genes Only); and including direct and indirect relationships.

Acknowledgments: We thank the Animal Research Facility (ARF) of the Health Science Center (HSC),

University of New Mexico School of Medicine. We also thank the UNM Preclinical Core of the Center

from Brain Recovery and Repair (University of New Mexico) for technical support.

Funding: This work was supported by the National Institute of Health grants NS106901 and

P206GM109089.

351 **References**

352 Aiba, Isamu, Andrew P. Carlson, Christian T. Sheline, and Shuttleworth C. William. 2012. "Synaptic

353 Release and Extracellular Actions of Zn 2+ Limit Propagation of Spreading Depression and Related

354 Events in Vitro and in Vivo." *Journal of Neurophysiology* 107 (3): 1032–41.

355 https://doi.org/10.1152/jn.00453.2011.

356 Ango, Fabrice, David Robbe, Jian Cheng Tu, Bo Xiao, Paul F Worley, Jean-Philippe Pin, Joël Bockaert,

357  and Laurent Fagni. 2002. "Homer-Dependent Cell Surface Expression of Metabotropic Glutamate

358  Receptor Type 5 in Neurons." *Molecular and Cellular Neurosciences* 20 (2): 323–29.

359  http://www.ncbi.nlm.nih.gov/pubmed/12093163.

360 Arenkiel, Benjamin R., Joao Peca, Ian G. Davison, Catia Feliciano, Karl Deisseroth, George J J.

361  Augustine, Michael D. Ehlers, and Guoping Feng. 2007. "In Vivo Light-Induced Activation of

362  Neural Circuitry in Transgenic Mice Expressing Channelrhodopsin-2." *Neuron* 54 (2): 205–18.

363  https://doi.org/10.1016/J.NEURON.2007.03.005.

364 Ayata, Cenk, and Martin Lauritzen. 2015. "Spreading Depression, Spreading Depolarizations, and the

365  Cerebral Vasculature." *Physiological Reviews* 95 (3): 953–93.

366  https://doi.org/10.1152/PHYSREV.00027.2014.

367 Berger, M., E. J. Speckmann, H. C. Pape, and A. Gorji. 2008. "Spreading Depression Enhances Human

368  Neocortical Excitability in Vitro." *Cephalalgia : An International Journal of Headache* 28 (5): 558–

369  62. https://doi.org/10.1111/J.1468-2982.2008.01556.X.

370 Bornstädt, Daniel von, Thijs Houben, Jessica L. Seidel, Yi Zheng, Ergin Dilekoz, Tao Qin, Nora Sandow,

371  et al. 2015. "Supply-Demand Mismatch Transients in Susceptible Peri-Infarct Hot Zones Explain

372  the Origins of Spreading Injury Depolarizations." *Neuron* 85 (5): 1117–31.

373  https://doi.org/10.1016/J.NEURON.2015.02.007.

374 Bottai, Daniele, John F. Guzowski, Martin K. Schwarz, Shin H. Kang, Bo Xiao, Anthony Lanahan, Paul

375  F. Worley, and Peter H. Seeburg. 2002. "Synaptic Activity-Induced Conversion of Intronic to

376  Exonic Sequence in Homer 1 Immediate Early Gene Expression." *The Journal of Neuroscience* 22

377  (1): 167. https://doi.org/10.1523/JNEUROSCI.22-01-00167.2002.

Carlson, Andrew P., Mohammad Abbas, Robert L. Alunday, Fares Qeadan, and C. William Shuttleworth.

2018. "Spreading Depolarization in Acute Brain Injury Inhibited by Ketamine: A Prospective,

Randomized, Multiple Crossover Trial." *Journal of Neurosurgery* 130 (5): 1513–19.

https://doi.org/10.3171/2017.12.JNS171665.

Carmichael, S. Thomas, and Marie Françoise Chesselet. 2002. "Synchronous Neuronal Activity Is a

Signal for Axonal Sprouting after Cortical Lesions in the Adult." *Journal of Neuroscience* 22 (14):

6062–70. https://doi.org/10.1523/JNEUROSCI.22-14-06062.2002.

Cavarsan, C. F., F. Tescarollo, C. Tesone-Coelho, R. L.T. Morais, F. L.T. Motta, M. M. Blanco, and L. E.

Mello. 2012. "Pilocarpine-Induced Status Epilepticus Increases Homer1a and Changes MGluR5

Expression." *Epilepsy Research* 101 (3): 253–60.

https://doi.org/10.1016/J.EPLEPSYRES.2012.04.011.

Charles, A., and K. C. Brennan. 2009. "Cortical Spreading Depression-New Insights and Persistent

Questions." *Cephalalgia : An International Journal of Headache* 29 (10): 1115–24.

https://doi.org/10.1111/J.1468-2982.2009.01983.X.

Chowdhury, Shoaib, Jason D. Shepherd, Hiroyuki Okuno, Gregory Lyford, Ronald S. Petralia, Niels

Plath, Dietmar Kuhl, Richard L. Huganir, and Paul F. Worley. 2006. "Arc Interacts with the

Endocytic Machinery to Regulate AMPA Receptor Trafficking." *Neuron* 52 (3): 445.

https://doi.org/10.1016/J.NEURON.2006.08.033.

Dietrich, W D, J Truettner, R Prado, N E Stagliano, W Zhao, R Busto, M D Ginsberg, and B D Watson.

2000. "Thromboembolic Events Lead to Cortical Spreading Depression and Expression of C-Fos,

Brain-Derived Neurotrophic Factor, Glial Fibrillary Acidic Protein, and Heat Shock Protein 70

MRNA in Rats." *Journal of Cerebral Blood Flow and Metabolism : Official Journal of the*

International Society of Cerebral Blood Flow and Metabolism 20 (1): 103–11.

https://doi.org/10.1097/00004647-200001000-00014.

Dreier, Jens P. 2011. "The Role of Spreading Depression, Spreading Depolarization and Spreading

Ischemia in Neurological Disease." *Nature Medicine* 17 (4): 439–47.

https://doi.org/10.1038/nm.2333.

Dreier, Jens P., Martin Fabricius, Cenk Ayata, Oliver W. Sakowitz, C. William Shuttleworth, Christian

Dohmen, Rudolf Graf, et al. 2017. "Recording, Analysis, and Interpretation of Spreading

Depolarizations in Neurointensive Care: Review and Recommendations of the COSBID Research

Group." *Journal of Cerebral Blood Flow and Metabolism* 37 (5): 1595–1625.

https://doi.org/10.1177/0271678X16654496.

Dreier, Jens P., Johannes Woitzik, Martin Fabricius, Robin Bhatia, Sebastian Major, Chistoph

Drenckhahn, Thomas Nicolas Lehmann, et al. 2006. "Delayed Ischaemic Neurological Deficits after

Subarachnoid Haemorrhage Are Associated with Clusters of Spreading Depolarizations." *Brain : A*

Journal of Neurology 129 (Pt 12): 3224–37. https://doi.org/10.1093/BRAIN/AWL297.

Edgar, Ron, Michael Domrachev, and Alex E. Lash. 2002. "Gene Expression Omnibus: NCBI Gene

Expression and Hybridization Array Data Repository." *Nucleic Acids Research* 30 (1): 207–10.

https://doi.org/10.1093/NAR/30.1.207.

Eising, Else, Reinald Shyti, Peter A.C. 't Hoen, Lisanne S. Vijfhuizen, Sjoerd M.H. Huisman, Ludo A.M.

Broos, Ahmed Mahfouz, et al. 2017. "Cortical Spreading Depression Causes Unique Dysregulation

of Inflammatory Pathways in a Transgenic Mouse Model of Migraine." *Molecular Neurobiology* 54

(4): 2986–96. https://doi.org/10.1007/S12035-015-9681-5.

Footitt, David R., and Nigel R. Newberry. 1998. "Cortical Spreading Depression Induces an LTP-like

Effect in Rat Neocortex in Vitro." *Brain Research* 781 (1–2): 339–42.

https://doi.org/10.1016/S0006-8993(97)01359-0.

García-Campos, Miguel A., Jesús Espinal-Enríquez, and Enrique Hernández-Lemus. 2015. "Pathway

Analysis: State of the Art." *Frontiers in Physiology* 6 (DEC): 383.

https://doi.org/10.3389/FPHYS.2015.00383/BIBTEX.

Ge, Steven Xijin, Eun Wo Son, and Runan Yao. 2018. "IDEP: An Integrated Web Application for

Differential Expression and Pathway Analysis of RNA-Seq Data." *BMC Bioinformatics* 19 (1): 1–

24. https://doi.org/10.1186/s12859-018-2486-6.

Hartings, Jed A., Norberto Andaluz, M. Ross Bullock, Jason M. Hinzman, Bruce Mathern, Clemens Pahl,

Ava Puccio, et al. 2020. "Prognostic Value of Spreading Depolarizations in Patients With Severe

Traumatic Brain Injury." *JAMA Neurology* 77 (4): 489–99.

https://doi.org/10.1001/JAMANEUROL.2019.4476.

Hartings, Jed A., C. William Shuttleworth, Sergei A. Kirov, Cenk Ayata, Jason M. Hinzman, Brandon

Foreman, R. David Andrew, et al. 2017. "The Continuum of Spreading Depolarizations in Acute

Cortical Lesion Development: Examining Leão's Legacy." *Journal of Cerebral Blood Flow and*

Metabolism 37 (5): 1571–94. https://doi.org/10.1177/0271678X16654495.

Helbok, Raimund, Jed A. Hartings, Alois Schiefecker, Baptiste Balança, Sharon Jewel, Brandon

Foreman, Ari Ercole, et al. 2020. "What Should a Clinician Do When Spreading Depolarizations

Are Observed in a Patient?" *Neurocritical Care* 32 (1): 306–10. https://doi.org/10.1007/s12028-019-

00777-6.

Hinzman, Jason M., Vince A. DiNapoli, Eric J. Mahoney, Greg A. Gerhardt, and Jed A. Hartings. 2015.

"Spreading Depolarizations Mediate Excitotoxicity in the Development of Acute Cortical Lesions."

Experimental Neurology 267 (May): 243–53. https://doi.org/10.1016/j.expneurol.2015.03.014.

Ibarra-Lopez, Veronica, Sangeeta Jayakar, Yeqing Angela Yang, Ciara Martin, Zora Modrusan, and

Sandra Rost. 2022. "Spatial Profiling of Protein and RNA Expression in Tissue: An Approach to

Fine-Tune Virtual Microdissection." *Journal of Visualized Experiments : JoVE*, no. 185 (July).

https://doi.org/10.3791/62651.

Jander, S, M Schroeter, O Peters, O W Witte, and G Stoll. 2001. "Cortical Spreading Depression Induces

Proinflammatory Cytokine Gene Expression in the Rat Brain." *J. Cereb. Blood Flow Metab.* 21 (3):

218–25. https://doi.org/10.1097/00004647-200103000-00005.

Kaido, Kempski, Heimann, Heers, and Bartsch. 2012. "Cluster Analysis of MRNA Expression Levels

Identifies Multiple Sequential Patterns Following Focal Cerebral Ischemia." *Turkish Neurosurgery*

22 (4). https://doi.org/10.5137/1019-5149.JTN.5523-11.0.

Karikó, Katalin, Valerie A. Harris, Yolanda Rangel, Monica E. Duvall, and Frank A. Welsh. 1998.

"Effect of Cortical Spreading Depression on the Levels of MRNA Coding for Putative

Neuroprotective Proteins in Rat Brain." *Journal of Cerebral Blood Flow and Metabolism* 18 (12):

1308–15. https://doi.org/10.1097/00004647-199812000-00005.

Kokaia, Zaal, Gunilla Gidö, Tomas Ringstedt, Johan Bengzon, Merab Kokaia, Bo K. Siesjö, Håkan

Persson, and Olle Lindvall. 1993. "Rapid Increase of BDNF MRNA Levels in Cortical Neurons

Following Spreading Depression: Regulation by Glutamatergic Mechanisms Independent of Seizure

Activity." *Molecular Brain Research* 19 (4): 277–86. https://doi.org/10.1016/0169-328X(93)90126-

A.

Kriz, Jasna, and Mélanie Lalancette-Hébert. 2009. "Inflammation, Plasticity and Real-Time Imaging after

Cerebral Ischemia." *Acta Neuropathologica* 117 (5): 497–509. https://doi.org/10.1007/S00401-009-

0496-1.

Lauritzen, Martin, Jens Peter Dreier, Martin Fabricius, Jed A. Hartings, Rudolf Graf, and Anthony John

Strong. 2011. "Clinical Relevance of Cortical Spreading Depression in Neurological Disorders:

Migraine, Malignant Stroke, Subarachnoid and Intracranial Hemorrhage, and Traumatic Brain

Injury." *Journal of Cerebral Blood Flow and Metabolism* 31 (1): 17–35.

https://doi.org/10.1038/jcbfm.2010.191.

Leao, Aristides A. P. 1944. "SPREADING DEPRESSION OF ACTIVITY IN THE CEREBRAL

473 CORTEX." *Journal of Neurophysiology* 7 (6): 359–90. https://doi.org/10.1152/jn.1944.7.6.359.

474 Lindquist, Britta E., and C. William Shuttleworth. 2014. "Spreading Depolarization-Induced Adenosine

475 Accumulation Reflects Metabolic Status in Vitro and in Vivo." *Journal of Cerebral Blood Flow and*

476 *Metabolism* 34 (11): 1779–90. https://doi.org/10.1038/jcbfm.2014.146.

477 Livak, Kenneth J., and Thomas D. Schmittgen. 2001. "Analysis of Relative Gene Expression Data Using

478 Real-Time Quantitative PCR and the 2-ΔΔCT Method." *Methods* 25 (4): 402–8.

479 https://doi.org/10.1006/meth.2001.1262.

480 Luo, P., T. Chen, Y. Zhao, L. Zhang, Y. Yang, W. Liu, S. Li, et al. 2014. "Postsynaptic Scaffold Protein

481 Homer 1a Protects against Traumatic Brain Injury via Regulating Group I Metabotropic Glutamate

482 Receptors." *Cell Death & Disease* 5 (4). https://doi.org/10.1038/CDDIS.2014.116.

483 Luo, Peng, Xia Li, Zhou Fei, and Waisang Poon. 2012. "Scaffold Protein Homer 1: Implications for

484 Neurological Diseases." *Neurochemistry International* 61 (5): 731–38.

485 https://doi.org/10.1016/j.neuint.2012.06.014.

486 Matsushima, Kazushi, Matthew J Hogan, and Antoine M Hakim. 1996. "Cortical Spreading Depression

487 Protects against Subsequent Focal Cerebral Ischemia in Rats." *J. Cereb. Blood Flow Metab.* 16 (2):

488 221–26. https://doi.org/10.1097/00004647-199603000-00006.

489 Matsushima, Kazushi, Rainald Schmidt-Kastner, Matthew J. Hogan, and Antoine M. Hakim. 1998.

490 "Cortical Spreading Depression Activates Trophic Factor Expression in Neurons and Astrocytes and

491 Protects against Subsequent Focal Brain Ischemia." *Brain Res.* 807 (1,2): 47–60.

492 https://doi.org/10.1016/s0006-8993(98)00716-1.

493 Nedergaard, Maiken, and Anker Jon Hansen. 1988. "Spreading Depression Is Not Associated with

494 Neuronal Injury in the Normal Brain." *Brain Research* 449 (1–2): 395–98.

495 https://doi.org/10.1016/0006-8993(88)91062-1.

Pertea, Mihaela, Daehwan Kim, Geo M. Pertea, Jeffrey T. Leek, and Steven L. Salzberg. 2016.

"Transcript-Level Expression Analysis of RNA-Seq Experiments with HISAT, StringTie and

Ballgown." *Nature Protocols 2016 11:9* 11 (9): 1650–67. https://doi.org/10.1038/nprot.2016.095.

Pietrobon, Daniela, and Michael A. Moskowitz. 2014. "Chaos and Commotion in the Wake of Cortical

Spreading Depression and Spreading Depolarizations." *Nature Reviews. Neuroscience* 15 (6): 379–

93. https://doi.org/10.1038/NRN3770.

Rangel, Yolanda M., Katalin Karikó, Valerie A. Harris, Monica E. Duvall, and Frank A. Welsh. 2001.

"Dose-Dependent Induction of MRNAs Encoding Brain-Derived Neurotrophic Factor and Heat-

Shock Protein-72 after Cortical Spreading Depression in the Rat." *Molecular Brain Research* 88 (1–

2): 103–12. https://doi.org/10.1016/S0169-328X(01)00037-7.

Sadowska, M, Mehlhorn C, Średniawa W, Szewczyk ŁM, Szlachcic A, Urban P, Winiarski M, and

Jabłonka JA. 2021. "Spreading Depressions and Periinfarct Spreading Depolarizations in the

Context of Cortical Plasticity." *Neuroscience* 453 (January): 81–101.

https://doi.org/10.1016/J.NEUROSCIENCE.2020.10.032.

Sato, Masaaki, Kazunori Suzuki, and Shigetada Nakanishi. 2001. "NMDA Receptor Stimulation and

Brain-Derived Neurotrophic Factor Upregulate Homer 1a MRNA via the Mitogen-Activated Protein

Kinase Cascade in Cultured Cerebellar Granule Cells." *The Journal of Neuroscience : The Official*

Journal of the Society for Neuroscience 21 (11): 3797–3805.

https://doi.org/10.1523/JNEUROSCI.21-11-03797.2001.

Shen, Ping Ping, Shuai Hou, Di Ma, Ming Ming Zhao, Ming Qin Zhu, Jing Dian Zhang, Liang Shu Feng,

Li Cui, and Jia Chun Feng. 2016. "Cortical Spreading Depression-Induced Preconditioning in the

Brain." *Neural Regeneration Research* 11 (11): 1857–64. https://doi.org/10.4103/1673-

5374.194759.

Shuttleworth, C. William, R. David Andrew, Yama Akbari, Cenk Ayata, Ramani Balu, K. C. Brennan,

Martyn Boutelle, et al. 2020. "Which Spreading Depolarizations Are Deleterious To Brain Tissue?" *Neurocritical Care* 32 (1): 317–22. https://doi.org/10.1007/s12028-019-00776-7.

Sintas, Cèlia, Noèlia Fernàndez-Castillo, Marta Vila-Pueyo, Patricia Pozo-Rosich, Alfons Macaya, and Bru Cormand. 2017. "Transcriptomic Changes in Rat Cortex and Brainstem After Cortical Spreading Depression With or Without Pretreatment With Migraine Prophylactic Drugs." *The Journal of Pain* 18 (4): 366–75. https://doi.org/10.1016/j.jpain.2016.11.007.

Takizawa, Tsubasa, Tao Qin, Andreia Lopes de Morais, Kazutaka Sugimoto, Joon Yong Chung, Liza Morsett, Inge Mulder, et al. 2020. "Non-Invasively Triggered Spreading Depolarizations Induce a rapidpro-Inflammatory Response in Cerebral Cortex." *Journal of Cerebral Blood Flow & Metabolism* 40 (5): 1117. https://doi.org/10.1177/0271678X19859381.

Tang, Yujie T., Jorge M. Mendez, Jeremy J. Theriot, Punam M. Sawant, Héctor E. López-Valdés, Y. Sungtaek Ju, and K. C. Brennan. 2014. "Minimum Conditions for the Induction of Cortical Spreading Depression in Brain Slices." *Journal of Neurophysiology* 112 (10): 2572–79. https://doi.org/10.1152/JN.00205.2014.

Urbach, Anja, Eileen Baum, Falko Braun, and Otto W. Witte. 2017. "Cortical Spreading Depolarization Increases Adult Neurogenesis, and Alters Behavior and Hippocampus-Dependent Memory in Mice." *Journal of Cerebral Blood Flow and Metabolism* 37 (5): 1776–90. https://doi.org/10.1177/0271678X16643736.

Urbach, Anja, Claus Bruehl, and Otto W. Witte. 2006. "Microarray-Based Long-Term Detection of Genes Differentially Expressed after Cortical Spreading Depression." *European Journal of Neuroscience* 24 (3): 841–56. https://doi.org/10.1111/j.1460-9568.2006.04862.x.

Volobueva, Maria N., Elena M. Suleymanova, Maria P. Smirnova, Alexey P. Bolshakov, and Lyudmila V. Vinogradova. 2023. "A Single Episode of Cortical Spreading Depolarization Increases MRNA Levels of Proinflammatory Cytokines, Calcitonin Gene-Related Peptide and Pannexin-1 Channels in

the Cerebral Cortex." *International Journal of Molecular Sciences* 24 (1).

https://doi.org/10.3390/ijms24010085.

Yanamoto, Hiroji, Jing Hui Xue, Susumu Miyamoto, Izumi Nagata, Yoshikazu Nakano, Kenichi Murao,

and Haruhiko Kikuchi. 2004. "Spreading Depression Induces Long-Lasting Brain Protection against

Infarcted Lesion Development via BDNF Gene-Dependent Mechanism." *Brain Res.* 1019 (1,2):

178–88. https://doi.org/10.1016/j.brainres.2004.05.105.

Figure Legends

Figure 1. Differential Expression (DE) analysis. We performed unpaired t-test followed by Welch's correction; significant genes (FC > 1.25, p value <0.05) are represented in black. Consequently, we performed DESeq2 analysis (Ge, Son, and Yao 2018); genes that were also significant after False Rate Discovery correction (FDR <0.05) are represented in red.

Figure 2. A) and B) top Molecular networks (score 63 and 32) generated by IPA analysis of DEGs upregulated in SD animals vs Sham. The intensity of the color red for each molecule is proportional to the FC of the gene. Full arrows indicate a direct relationship between molecules, while dotted arrows indicate an indirect relationship between genes. Orange arrows indicate the predicted activation of a gene, while yellow indicates predicted inhibition. **C) Top Diseases and Functions.**

Figure 3. Top Cellular functions generated by IPA analysis of DEGs upregulated in SD animals vs Sham. Top pathways identified include: cellular assembly and organization, cellular function and maintenance, and cellular development and proliferation**.** The heatmaps show the p value and the activation score for each pathway, and a graphical representation of the networks and the genes respectively involved. In particular, the intensity of the color red for each molecule is proportional to the FC of the gene. Orange arrows indicate a positive control of the gene on the related pathway, and the pathway's name highlighted in orange indicates its activation.

Figure 4. A) Differential Expression (DE) analysis. We performed unpaired t-test followed by Welch's correction; significant genes (FC > 1.25, p value <0.05) are represented in black. Consequently, we

performed DESeq2 analysis (Ge, Son, and Yao 2018); genes that were also significant after False Rate

Discovery correction (FDR <0.05) are represented in red, in green we highlighted the genes in common

with SD vs Sham DE analysis. **B) top molecular network (score 65) generated by IPA analysis of DEGs**

upregulated in SD animals vs Contralateral. The intensity of the color red for each molecule is

proportional to the FC of the gene. Full arrows indicate a direct relationship between molecules, while

dotted arrows indicate an indirect relationship between genes. Orange arrows indicate the predicted

activation of a gene, while yellow indicates predicted inhibition. **C) Top Diseases and Functions.**

Figure 5. Top Cellular functions generated by IPA analysis of DEGs upregulated in SD animals vs

Contralateral. Top pathways identified include: cellular assembly and organization, cellular function and

maintenance, and cellular development and proliferation**.** The heatmaps show the p value and the activation

score for each pathway, and a graphical representation of the networks and the genes respectively involved.

In particular, the intensity of the color red for each molecule is proportional to the FC of the gene. Orange

arrows indicate a positive control of the gene on the related pathway, and the pathway's name highlighted

in orange indicates its activation.

Figure 6. RT-qPCR validation of SD target genes identified by RNA-seq. mRNA levels of the target

genes were measured as described in Material and Methods. Data were analyzed by unpaired t-test,

followed by Welch's correction and are represented as mean ± SEM; $p<0.05$*, **$p<0.01$, ***$p<0.001$; n=6.

Figure 7. Target genes showed differential expression depending on the distance from the SD

initiation site. mRNA levels of the target genes were measured as described in Material and Methods.

Specifically, we determined the mRNA levels in 4 different regions: 1) SD initiation site, 2) SD intermediate

site, 3) SD remote site, and 4) the contralateral hemisphere. Data were analyzed by 2-way ANOVA

followed by Tukey's Multiple Comparison test, and are represented as mean ± SEM; p<0.05*, **p<0.01,

***p<0.001; and unpaired t-test, followed by Welch's correction and are represented as mean ± SEM;

#p<0.05, ##p<0.01, ###p<0.001; n=3.

Figure 8. The combination of unbiased RNA-seq analysis with predicted pathway analysis showed that

clusters of SDs could activate a range of neuroprotective pathways including increased cell proliferation,

decreased cell death, cell differentiation, and neuronal plasticity. The identification of molecular markers

responsible for the activation of this pathways could lead to the discovery of new therapeutic targets.

Supplementary Figure 1. Schematic representation of the SDs induction: SDs were induced

repetitively (4 SDs in 2 hours at 30 min intervals) in healthy, female mice and SDs were confirmed with

Intrinsic Optical Signal imaging. SD induction was with either focal application of KCl (NaCl for sham

controls, n=6) in C57Bl/6 mice or with optogenetic stimulation (2 mW for 20 seconds) in Thy1-ChR2-YFP

mice (n=6). **C) iOS images of the SDs propagating through the left hemisphere**; after KCl application,

or Optogenetic stimulation.

Supplementary Figure 2 Differential Expression (DE) analysis. We performed unpaired t-test followed

by Welch's correction; significant genes (FC > 1.25, p value <0.05) are represented in black. Consequently,

we performed DESeq2 analysis (Ge, Son, and Yao 2018); no genes significant after False Rate Discovery

correction (FDR <0.1).

Supplementary Figure 3. A) Representation of the localization of the micro dissected tissues collected for

qRT-PCR **B)** GAPDH levels in RNA sequencing samples are expressed as FPKM. No differences in

GAPDH were found after either KCl- or optogenetic- induced SDs. Thus, GAPDH represents a valid hkg

for qRT-PCR validation. Data were analyzed by unpaired t-test, followed by Welch's correction and

represented as mean ± SEM, n=3. **C)** Target genes showed differential expression depending on the distance

from the SD initiation site. mRNA levels of the target genes were measured as described in Material and

Methods. Specifically, we determined the mRNA levels in 4 different regions: 1) SD initiation site, 2) SD

intermediate site, 3) SD remote site, and 4) the contralateral hemisphere. Data were analyzed by 2-way

ANOVA followed by Tukey's Multiple Comparison test, and are represented as mean ± SEM; n=3.

RNAseq SD vs Sham

-log10 (p-value)

FDR < 0.05

p value < 0.05

log2(Fold Change)

DEGs

Fold Change

FOS
PTGS2
FOSB
ARC
DUSP6
EGR2
JUNB
EGR4
DUSP5
RGS2
ARL4D
BTG2
GADD45G
EGR1
DUSP1
TRIB1
IER2
KLF6
GM13889
ADRB1
KCNJ2
ATF3
FLRT3
JUN

2A.

2B.

2C.

Disease and Disorders	p-value range	# Molecules
Neurological Diseases	3.69E-02 - 7.77E-09	42
Organismal Injury and Abnormalities	3.69E-02 - 6.10E-09	43
Psychological Disorders	2.65E-02 - 6.10E-09	29
Hereditary Disorders	2.65E-02 - 6.10E-09	18
Skeletal and Muscular Disorders	2.65E-02 - 2.70E-08	17

Physiological System Development and Functions	p-value range	# Molecules
Nervous System Development and Function	3.69E-02 - 7.62E-07	51
Tissue Morphology	3.69E-02 - 4.54E-08	28
Tissue Development	3.46E-02 - 7.62E-07	40
Organismal Development	3.46E-02 - 4.43E-06	37
Behavior	3.17E-02 - 1.11E-05	12

Legend:
- Enzyme
- Growth factor
- Ion Channel
- Kinase
- Ligand-dependent Nuclear Receptor
- Transcription Regulator
- Other
- Disease
- Function
- — — Relationship

	p-value	Activation Score
Cell death of hippocampal neurons	8.02E-05	-2.410
Apoptosis	4.95E-07	-2.003
Neuronal cell death	1.68E-05	-1.974
Cell death of cerebral cortex cells	0.000146	-1.589
Degeneration of neurons	0.0309	-0.660
Cell death of neuroglia	0.00234	-0.339
Cell death of cortical neurons	0.033	-0.283
Apoptosis of granule cells	6.06E-05	-0.266
Apoptosis of neuroglia	0.00413	-0.152
Apoptosis of sympathetic neuron	0.000254	-0.145
Cell viability of hippocampal neurons	0.00023	1.124
Cell viability of cerebral cortex cells	0.000519	1.413
Survival of cephalic neurons	0.000519	1.413
Survival of neural cells	0.0241	1.774
Cell viability	0.0263	1.774

	p-value	Activation Score
Proliferation of neuroglia	0.00193	-0.314
Formation of dendritic spines	0.0147	0.577
Formation of dendrites	0.0284	0.577
Growth of dendrites	0.00487	1.082
Dendritic growth/branching	0.000413	1.090
Density of dendritic spines	0.02	1.165
Axonogenesis	0.0156	1.412
Branching of neurons	5.32E-05	1.587
Proliferation of neural cells	0.00537	1.627
Neuritogenesis	5.03E-06	1.913
Branching of axons	0.00933	1.922
Branching of neurites	0.000164	1.924
Outgrowth of neurites	0.0169	1.988
Growth of neurites	0.0122	2.238
Development of neurons	1.27E-06	2.286

4A.

RNAseq SD vs Contralateral

NPAS4
FOS
PTGS2
FOSB
DUSP6
ARC
NR4A1
JUNB
EGR4
EGR2
EGR4
DUSP5
RGS2
BTG2
TRIB1
EGR1
DUSP1
ARL4D
GADD45G
IER2
NPTX2
TRIB1
ADRB1
GM13889
KLF6
KCNJ2
JUN
CITED2
ATF3

4B.

4C.

Disease and Disorders	p-value range	# Molecules
Neurological Diseases	4.56E-02 - 2.49E-08	73
Organismal Injury and Abnormalities	4.56E-02 - 1.91E-08	73
Psychological Disorders	2.85E-02 - 1.91E-08	35
Hereditary Disorders	1.54E-02 - 1.91E-08	21
Skeletal and Muscular Disorders	7.75E-03 - 6.30E-08	20

Physiological System Development and Functions	p-value range	# Molecules
Nervous System Development and Function	4.56E-02 - 7.04E-07	61
Tissue Morphology	4.56E-02 - 1.07E-09	39
Tissue Development	4.56E-02 - 1.03E-06	49
Organismal Development	4.56E-02 - 5.03E-06	44
Organ Morphology	4.56E-02 - 5.96E-05	16

	p-value	Activation Score	
Cell death of hippocampal neurons	0.0015	-1.958	
Apoptosis	9.12E-08	-1.197	
Neuronal cell death	8.64E-06	-0.903	
Cell death of cerebral cortex cells	0.00133	-0.838	
Apoptosis of neuroglia	0.00107	-0.152	
Apoptosis of sympathetic neuron	6.08E-05	-0.145	
Apoptosis of granule cells	0.000199	0.063	
Cell death of cortical neurons	0.0346	0.173	
Cell viability of cerebral cortex cells	0.00112	1.126	
Survival of cephalic neurons	0.00112	1.126	
Survival of neural cells	0.0188	1.475	
Cell viability	0.0203	1.475	

	p-value	Activation Score	
Proliferation of neuroglia	0.00204	0.784	
Differentiation of neuroglia	0.00708	1.090	
Dendritic growth/branching	0.000278	1.412	
Axonogenesis	0.00923	1.587	
Branching of neurons	0.000015	1.772	
Outgrowth of neurites	0.0255	1.912	
Development of neural cells	7.62E-07	1.922	
Branching of axons	0.00252	1.924	
Branching of neurites	6.03E-05	1.985	
Growth of dendrites	0.00127	2.048	
Neuritogenesis	4.43E-06	2.183	
Differentiation of neural cells	0.000945	2.183	
Development of neurons	1.3E-06	2.224	

Cluster of Cortical Spreading Depolarizations (SDs)

Differential Expression (DE) & Pathway Analysis (PA)

Increased expression of genes inducing Cellular Development and Differentiation

Preconditioning Synaptic Plasticity activation

Molecular Markers Therapeutic Targets

1A.

1B.

1. SD Initiation site
2. SD Intermediate site
3. SD remote site
Burr Hole
4. Contralateral

GAPDH

NR4A1

0.1185

0.0848

EGR2

☐ Sham
■ KCl-SD