Syndapin and GTPase RAP-1 Control Endocytic Recycling via RHO-1 and Non-Muscle Myosin II

Wilmer R. Rodriguez-Polanco[1], Anne Norris[1], Agustin B. Velasco[1], Adenrele M. Gleason[1], and Barth D. Grant[1, 2]*

[1]Department of Molecular Biology and Biochemistry Rutgers University, Piscataway, New Jersey 08854, United States of America.

[2]Rutgers Center for Lipid Research

* barthgra@dls.rutgers.edu

Abstract

After endocytosis, many plasma membrane components are recycled via narrow-diameter membrane tubules that emerge from early endosomes to form recycling endosomes, eventually leading to their return to the plasma membrane. We previously showed that the F-BAR and SH3 domain Syndapin/PACSIN-family protein SDPN-1 is required *in vivo* for basolateral endocytic recycling in the *C. elegans* intestine. Here we sought to determine the significance of a predicted interaction between the SDPN-1 SH3 domain and a target sequence in PXF-1/PDZ-GEF1/RAPGEF2, a known exchange factor for Rap-GTPases. We found that endogenous mutations we engineered into the SDPN-1 SH3 domain, or its binding site in the PXF-1 protein, interfere with recycling *in vivo*, as does loss of the PXF-1 target RAP-1. Rap-GTPases have been shown in several contexts to negatively regulate RhoA activity. Our results show that RHO-1/RhoA is enriched on SDPN-1 and RAP-1 positive endosomes in the *C. elegans* intestine, and loss of SDPN-1 or RAP-1 elevates RHO-1(GTP) levels on intestinal endosomes. Furthermore, we found that depletion of RHO-1 suppressed *sdpn-1* mutant recycling defects, indicating that control of RHO-1 activity is a key mechanism by which SDPN-1 acts to promote endocytic recycling. RHO-1/RhoA is well-known for controlling actomyosin contraction cycles, although little is known of non-muscle myosin II on endosomes. Our analysis found that non-muscle myosin II is enriched on SDPN-1 positive endosomes, with two non-muscle myosin II heavy chain isoforms acting in apparent opposition. Depletion of *nmy-2* inhibited recycling like *sdpn-1* mutants, while depletion of *nmy-1* suppressed *sdpn-1* mutant recycling defects, indicating actomyosin contractility in controlling recycling endosome function.

Introduction

Cells internalize extracellular material, receptor-associated ligands, membrane proteins and lipids by endocytosis (1). Uptake mechanisms can be generally divided into clathrin-dependent and clathrin-independent endocytosis (CDE and CIE, respectively), after which internalized cargo is sorted within endosomes for delivery to distinct intracellular destinations (2). As membrane and fluid are added to the early endosome by incoming vesicles, narrow-diameter recycling tubules form, collecting a subset of cargo (3). Most plasma membrane lipids, and many types of internalized transmembrane proteins, are recycled back to the plasma membrane either directly (rapid recycling), indirectly via the endocytic recycling compartment (ERC) (slow recycling), or Golgi (retrograde recycling) (3, 4). As early endosomes mature they become late endosomes, eventually fusing with lysosomes, promoting degradation of remaining cargo. Many cellular processes depend upon accurate and efficient endocytic recycling, including cell migration, cell division, cell-cell fusion events, and control of neuronal signaling during learning and memory (3, 4). Trafficking pathways are often more elaborate in the context of polarized cells such as epithelia and neurons that maintain multiple functionally distinct plasma membrane domains, e.g. basolateral versus apical in epithelia, and dendrite versus axon in neurons (4).

The *C. elegans* intestine is a polarized epithelial tube consisting of exactly 20 cells that form nine donut-like rings, the most anterior ring consisting of four symmetrical cells and all other rings comprised of two cells (5, 6). The intestinal cells have a basolateral surface that faces the pseudocoelom and controls the exchange of molecules between the intestine and the rest of

the body (5). The microvilli-rich apical surface faces the central lumen and is important for

nutrient uptake (6). We previously developed the *C. elegans* intestine as a model system for the

study of polarized epithelial trafficking mechanisms, allowing analysis of transport in the native

context of a polarized-epithelial tube within the intact living animal (7-10). Taking advantage of

the *C. elegans* intestinal system, our laboratory previously showed that *C. elegans* SDPN-1, a

Syndapin/PACSIN family protein, is important for basolateral endocytic recycling *in vivo,* a

finding confirmed independently by others (9, 11).

Syndapin/PACSIN family proteins bear a single N-terminal F-BAR domain and a single C-terminal

SH3 domain. BAR domains in general, including the F-BAR subclass, are known to sense

membrane curvature and to promote positive curvature, properties likely to help drive

membrane budding and tubulation (12). Our previous work showed that *C. elegans* SDPN-1 is

essential for the basolateral recycling of cargo proteins that are transported via the early

endosome–to–recycling endosome route, but not other routes (Gleason et al., 2016). Affected

cargos accumulate at much higher-than-normal levels in the endosomes of *sdpn-1* mutants

(Gleason et al., 2016). These include well characterized model cargo proteins hTAC (human IL-2

receptor α-chain), hTFR (human transferrin receptor), and DAF-4 (type-II TGF-β receptor), but

not MIG-14/Wls or SMA-6 (type-I TGF-β receptor) that recycle via the retromer pathway, or

CD4-LL that does not recycle (Gleason et al., 2016). SDPN-1 protein is highly enriched on early

and basolateral recycling endosomes of the intestinal cells, and *sdpn-1 null* mutants display

abnormal accumulation of early and basolateral recycling endosomes, suggesting a direct role

in their regulation (Gleason et al. 2016). Mammalian Syndapin2 is also well known to associate

with recycling endosome proteins and to promote endocytic recycling, but many questions

remain about the mechanism by which Syndapin/PACSIN proteins regulate endosomal function

(13-15).

To better understand how SDPN-1 promotes recycling we sought to identify partner proteins

relevant to recycling, with a particular focus on results from the *C. elegans* SH3-ome project

(16). Here we investigated one high-confidence hit for the SDPN-1 SH3 domain, PXF-1/PDZ-GEF,

an exchange factor for the small Ras-like GTPase RAP-1/Rap (17, 18). Importantly we show that

PXF-1 and its target RAP-1 also regulate recycling like SDPN-1, and interfering with the

interaction of SDPN-1 and PXF-1 disrupts recycling. We trace the activity of RAP-1 on

endosomes to negative regulation of RHO-1 and myosin II heavy chain isoform NMY-1, and

show that reduction in RHO-1 or NMY-1 greatly ameliorates *sdpn-1* mutant trafficking defects.

This function appears to work in opposition to myosin II heavy chain isoform NMY-2 that we

find is required for recycling. Taken together our results indicate that myosin II regulation is a

key output of endosomal SDPN-1 that is essential to maintain recycling function.

Materials and Methods

General methods and strains

All *C. elegans* strains were derived from Bristol strain N2 and cultured at 20°C on NGM plates

seeded with the *Escherichia coli* strain OP50. A list of strains used in this study can be found in

Supplemental Table 1.

Transgenic Strains

To construct GFP or RFP fusion transgenes for expression in the worm intestine, Gateway

destination vectors were used. Each vector contained the promoter region of the intestine-

specific gene *vha-6*, followed by GFP, mScarlet, or tagRFP coding sequences, a Gateway

cassette (Invitrogen, Carlsbad, CA), and *let-858* 3' UTR sequences. The genomic or cDNA

sequences of *C. elegans* genes were cloned individually into entry vector pDONR221. Mutant

forms of the coding regions were obtained using site-directed mutagenesis using the

Quikchange II kit (Stratagene). Coding regions were transferred into P*vha-6*-GFP or P*vha-6*-RFP

vectors by LR reaction (Invitrogen). Constructs were bombarded into *unc-119(ed3)* mutant

animals to establish low copy integrated transgenic lines, or integrated in single copy using the

MiniMos system and Hygromycin or G418 selection (19).

RNAi

C. elegans RNAi-mediated interference was performed as described (20). Feeding constructs in

this study were from the Ahringer library, except for *rho-1*, for which a cDNA was cloned into

the L4440/pPD129.36 vector between unique SacI and HindIII sites (21). RNAi clones for *nmy-1*

and *nmy-2* were kindly provided by Adriana Dawes (Ohio State University) (22). For most RNAi

experiments, synchronized L4 stage animals were grown on RNAi plates at 20°C. After the first

generation, we transferred about 60 L4 stage animals to fresh RNAi plates at 20°C for another

24 hours prior to imaging as young adults.

Protein Expression, Pulldown Assays, and Western Analysis

N-terminally hemagglutinin (HA)-tagged PXF-1 protein, amino acids 1-477, with or without

mutation of the predicted SDPN-1 binding site, was synthesized *in vitro* using the TNT-coupled

transcription-translation system (Promega, Madison, WI) using DNA templates pcDNA3.1–

2xHA-PXF-1 and pcDNA3.1–2xHA-PXF-1(K191/P195/P198->AAA), respectively (1.6 ug/each 50-

ul reaction). The reaction cocktail was incubated at 30°C for 90 min. Control glutathione *S*-

transferase (pGEX-2T), GST-SDPN-1 SH3 (final 112 amino acids), and GST-SDPN-1 SH3 (P479L)

fusion proteins were expressed in the *ArcticExpress* strain of *Escherichia coli* (Stratagene, La

Jolla, CA). Bacterial pellets were lysed in 10 ml B-PER Bacterial Protein Extraction Reagent

(Pierce, Rockford, ILL) with Complete Protease Inhibitor Cocktail Tablets (Roche, Indianapolis,

IN). Extracts were cleared by centrifugation, and supernatants were incubated with

glutathione-Sepharose 4B beads (Amersham Pharmacia, Piscataway, NJ) at 4°C overnight.

Beads were then washed six times with cold STET buffer (10 mM Tris-HCl, pH 8.0, 150 mM NaCl,

1 mM EDTA, 0.1% Tween-20). In vitro–synthesized HA-tagged protein (10 ul TNT mix diluted in

500 ul STET) was added to the beads and allowed to bind at 4°C for 2 h. After six additional

washes in STET the proteins were eluted by boiling in 30 ul 2X-SDS-PAGE sample buffer. Eluted

proteins were separated on SDS-PAGE, blotted to nitrocellulose, visualized with Ponceau S and

scanned, then washed to remove Ponceau staining, and probed with anti-HA (16B12)

antibodies and HA-conjugated secondary antibodies (Pierce). Detection was achieved using

Super Signal West Pico detection system (Pierce).

CRISPR

sdpn-1 P479L and *pxf-1* K191/P195/P198->AAA mutations were introduced into the

endogenous genes in *C. elegans* using the *dpy-10* co-CRISPR technique as described by Paix and

Seydoux (23), creating mutant alleles *sdpn-1(pw33)* and *pxf-1(pw34)*. 3XFLAG-GFP was inserted

into the *pxf-1* gene between the last codon and stop codon using the SEC selection technique

described by Dickinson as modified by Schwartz (24, 25).

Microscopy and image analysis

Imaging was performed on live first-day adult hermaphrodites animals mounted on slides using

5% agarose pads with 10 mM levamisole. Most single channel data in Figure 2 was captured on

a Zeiss LSM 710 confocal microscope. All other single channel data, including Figure **2B**-**B'**, was

captured on a Zeiss LSM 800 confocal microscope. For both microscopes, the LSM 710 and LSM

800, GFP tagged protein image acquisition used the 488 nm laser line and the 63X (1.4 NA)

objective and ZEISS ZEN 2.3 imaging software. True GFP signal was separated from the broad-

spectrum auto-fluorescence displayed by the lysosome-related organelles (gut granules) of the

C. elegans intestine using the Spectral Fingerprinting and Linear Unmixing functions in ZEN. The

images were exported as TIF files for analysis in Metamorph version 7.7 software.

The "Integrated Morphometry Analysis" function of Metamorph was used to measure

fluorescence intensity, puncta number (structure count), and fluorescence area (total area)

within unit regions. At least 9 animals were used for each analysis. Fluorescence signal in the

basolateral membrane between two neighboring cells was avoided for the analysis.

For multiwavelength fluorescence colocalization studies, images were obtained using a

spinning-disk confocal imaging system: Axiovert Z1 microscope (Carl Zeiss Microimaging)

equipped with a X-Light V2 Spinning Disk Confocal Unit (CrestOptics), 7-line LDI Laser Launch

(89 North), Prime 95B Scientific CMOS camera (Photometrics) and 63X and 100X oil-immersion

objectives. Data were captured as an image stack with 0.2 µm step size using Metamorph

version 7.7 software (Universal Imaging), and deconvolved using Autoquant X3 (Media

Cybernetics). In Metamorph, a single plane from each image stack (either top plane or middle

plane) was chosen in which both markers were clearly visible. In the middle plane of the worm

intestine, the apical membranes and the basolateral membranes are visible, while on top plane

images only the basolateral membrane is visible. Images taken in the DAPI channel were used

to identify broad-spectrum intestinal autofluorescence caused by lysosome-like organelles. The

DAPI channel image was subtracted from the green and red channel images to remove

autofluorescent signal from the colocalization analysis. The DAPI subtraction was done using

the Metamorph "Arithmetic" function at 16-bit depth. Quantification of colocalization

(Manders coefficients) in autofluorescence subtracted images was performed using open

source Fiji (Image J) software and the "Colocalization Threshold" function. At least 9 animals

were used for each analysis. For middle plane images, the fluorescence signal in the apical

membrane and the basolateral membrane between two neighboring cells were avoided.

Statistical analysis

Quantified data obtained in this study was graphed and statistically analyzed using GraphPad

Prism 7.04 software.

Results

The SDPN-1 SH3 domain binds to PXF-1 via a specific proline-rich sequence

Syndapin family proteins, also known as PACSINs, contain an N-terminal F-BAR domain (FCH-

BAR, Fes/CIP4 homology-Bin/Amphiphysin/Rvs) and a C-terminal SH3 (Src-homology 3) domain

(14). *C. elegans* has a single Syndapin family member SDPN-1, while the situation in mammals is

more complex, with three Syndapin family members Syndapin1/PACSIN1, Syndapin2/PACSIN2,

and Syndapin3/PACSIN3 (26). Syndapins form dimers through their F-BAR domains, with the F-

BAR dimer forming a mildly curved structure that binds directly to membrane phospholipids

(14). Higher order assemblies of these proteins scaffold the membrane to promote bending and

remodeling, in many cases producing membrane tubulation (26). Indeed, our previous work

showed that full-length recombinant *C. elegans* SDPN-1 tubulates acidic membranes *in vitro* (9).

In vivo SDPN-1 localizes to early endosomes and basolateral recycling endosomes in the *C.*

elegans intestine, and is required for basolateral recycling, but the molecular mechanism(s) by

which it mediates endocytic recycling remain poorly understood (9). To gain a better

understanding of how SDPN-1 promotes endosomal transport we sought to identify partner

proteins that work with SDPN-1 in this process.

The SH3 domain is a peptide recognition module for proline-rich sequences that is well-known

for mediating specific protein-protein interactions (27). Thus, we focused on the SDPN-1 SH3

domain as a molecular entry point to identify direct SDPN-1 binding partners. We were aided in

this goal by the *C. elegans* SH3-interactome project (16), a large-scale analysis that used peptide

phage display to define the binding preferences for most SH3-domains encoded in the *C.*

elegans genome, including generating a weighted SDPN-1 SH3 consensus target sequence (Fig.

1A). The SH3 interactome project also screened directly for SDPN-1 SH3-domain interacting

proteins using a stringent yeast-2-hybrid approach. Importantly, both approaches identified the

protein PXF-1 as a prime SDPN-1 SH3 domain candidate. The project recovered PXF-1 via

genome-wide yeast-2-hybrid with a SDPN-1 SH3 bait, but not any of the other SH3 domain

baits, and independently identified PXF-1 amino acids 191-198 as the second-best match for

the SDPN-1 SH3-binding consensus in the entire predicted *C. elegans* proteome (16).

PXF-1 (PDZ-GEF or RAGEF in mammals) is a well-characterized exchange factor for the small

Ras-like GTPases RAP-1/Rap1 and RAP-2/Rap2, and Rap-GTPases have been implicated in some

endocytic recycling events in other organisms, further suggesting that a SDPN-1/PXF-1

interaction could be relevant to understanding the role of SDPN-1 in recycling (17, 28-30). PXF-1

has a complex domain structure, with two RCBD-like domains: one near the N-terminus, and

the other more centrally located(17, 31). While RCBD domains in some proteins bind to cAMP,

biochemical analysis of human PDZ-GEF2, a PXF-1 ortholog, failed to identify any cyclic-

nucleotide binding for these RCBD-like domains (32). The predicted SDPN-1 SH3 binding site at

residues 191-198 resides in a predicted unstructured region between the two RCBD-like

domains (https://alphafold.ebi.ac.uk/entry/G5EDB9). PXF-1 also contains REM (Ras Exchange

Motif), PDZ (PSD-95, Dlg, and ZO-1 homology), RA (Ras Association), and GEF (Guanine

nucleotide Exchange Factor) domains, as well as a C-terminal VSRV sequence that may interact

with PDZ-domains (Fig 1B) (17).

To better analyze the ability of the SDPN-1 SH3 domain to bind to PXF-1, and test the

importance of the predicted binding site identified as a strong match to the phage-display

consensus sequence, we performed GST pulldown experiments. As baits we compared GST,

GST-SH3[sdpn-1](+), and GST-SH3[sdpn-1](P478L). The P478L mutation in the SDPN-1 SH3 domain is

equivalent to mammalian Syndapin I P434L, which was previously shown to abolish the ability

of the Syndapin I SH3 domain to bind partner proteins (33). These baits were paired with *in*

vitro translated HA-PXF-1(1-478) bearing the wild-type sequence, or with the consensus-

matching PXF-1(191-198) SH3 binding site mutated (**K**GPA**P**LP**P** to **A**GPA**A**LP**A**) (Fig 1A-B). We

found that HA-PXF-1(+) interacted robustly with GST-SH3[sdpn-1](+), but not GST or GST-SH3[sdpn-]

[1](P478L), indicating the specificity of the interaction. Moreover, we found that mutation of HA-

PXF-1 at the predicted interaction sequence (191-198) abolished the interaction with GST-

SH3[sdpn-1](+), despite other proline-rich sequences remaining in the PXF-1 prey protein. Taken

together these results indicate that the SDPN-1 SH3 domain can bind to PXF-1, and the

predicted binding site at aa191-198 is crucial for this interaction.

The SDPN-1/PXF-1 interaction is important for endocytic recycling

Given our binding results we hypothesized that the interaction of the SDPN-1 SH3-domain with

PXF-1 is important for SDPN-1-mediated regulation of endocytic recycling. To interfere with this

interaction *in vivo* we used CRISPR-mediated genome editing to create *C. elegans* mutants

sdpn-1(pw33) and *pxf-1(pw34)*, in which the same point mutations that interfered with the

SDPN-1/PXF-1 binding interaction *in vitro* were engineered into the endogenous *sdpn-1* and

pxf-1 genes *in vivo* (Fig. 1b). While *pxf-1* null mutations are lethal (17), *pxf-1(pw34)* is a viable

allele.

In particular we found that the interaction-defective mutants *sdpn-1(pw33)* and *pxf-1(pw34)*

displayed gross over-accumulation of intestinal GFP::RME-1 marked basolateral recycling

endosomes and GFP::RAB-5 marked early endosomes in the central cytoplasm (middle plane

Fig. 2K), very similar to phenotypes we had previously observed in null mutants for *sdpn-1* (Fig.

2, panel A-A'' and B-B''; Fig. S1 A, A") (9). For GFP::RME-1 labeled recycling endosomes we

measured endosome fluorescence intensity and number. In the case of GFP::RAB-5, the

mutants accumulated so many early endosomal compartments that it was not possible to

segment the images to quantify endosome number, so in addition to fluorescence intensity we

measured the the percent of total cytoplasmic area labeled by GFP::RAB-5.

We also tested the importance of the SDPN-1/PXF-1 interaction in cargo trafficking, focused on

well-characterized model endocytic recycling cargoes markers hTAC (human IL2-receptor alpha-

chain) and hTFR (human transferrin receptor) tagged with GFP, both of which depend upon

SDPN-1 for their recycling (9, 10). hTAC is a clathrin-independent cargo that recycles via the

recycling endosome in an ARF-6-dependent pathway (34, 35), while hTFR is clathrin-dependent

in its endocytosis and also recycles via recycling endosomes (10).   Previous studies showed that

sdpn-1 null mutants display dramatic endosomal accumulation of hTAC and hTFR within the

intestinal cells (9). Importantly, we observed that the interaction-defective mutants caused

intracellular accumulation of recycling cargoes similar to that previously observed upon the

complete loss of SDPN-1 (Fig. 2, panel C-C'' and D-D''). Taken together, our results indicate that

a SDPN-1/PXF-1 interaction is important for the function of the basolateral recycling pathway.

The RAP-1 GTPase cycle is required for endocytic recycling

Since *C. elegans* PXF-1 and its mammalian homologs are specific GDP/GTP exchange factors for

Rap-GTPases *in vivo* and *in vitro* (17, 36, 37), we surmised that the role of the SDPN-1/PXF-1

interaction is to activate Rap-GTPases to promote endocytic recycling. To test this hypothesis,

we analyzed the effect of a *rap-1(0)* mutant on relevant markers *in vivo*. Indeed, we found that

rap-1 null mutants displayed significantly increased accumulation of RAB-5 marked early

endosomes and RME-1 marked recycling endosomes, as well as increased intracellular

accumulation of recycling cargo markers hTAC::GFP and hTFR::GFP, supporting this hypothesis

(Fig. 2, panel A''', B''', C''', D''', F, H-J; Fig. S1 A, A''). Interestingly, we also identified abnormal

accumulation defects for recycling cargo and endosome markers in a GTPase-defective *rap-*

1(re180) mutant (38), bearing the G12V mutation at the endogenous gene (Fig. 2B',C'''', D'''',

E'''', F, G, I-J; Fig. S1 A'). These results indicate a requirement for the full RAP-1 GTP/GDP-cycle,

rather than a simple requirement for active RAP-1. We also tested for effects of *sdpn-1(0)* and

rap-1(0) mutants on GFP-tagged *C. elegans* GLUT1 homolog FGT-1, a glucose transporter known

to be naturally expressed in the *C. elegans* intestine (39). Loss of SDPN-1 and RAP-1 produced

similar intracellular retention of FGT-1::GFP, also suggesting a defect in recycling (Fig. S2). Taken

together these results support an important role for RAP-1 in regulating recycling downstream

of SDPN-1 and PXF-1.

PXF-1 and RAP-1 are enriched on early and basolateral recycling endosomes

To further interrogate SDPN-1, PXF-1, and RAP-1 for shared function together in recycling, we

compared the localization of PXF-1 and RAP-1 with SDPN-1 in the context of the intestinal

epithelium within the intact whole animal. We also compared RAP-1 localization to

independent early endosome and basolateral recycling endosome markers with which SDPN-1

is known to colocalize (9). We found that endogenously tagged PXF-1::GFP colocalized with

intestinal SDPN-1::tagRFP indicating significant residence of PXF-1 with SDPN-1 on basolateral

endosomes (Fig 3 A-A''''and E). We also identified strong colocalization of intestinal SDPN-

1::GFP with tagRFP::RAP-1, indicating that at any given time most SDPN-1 overlaps with RAP-1

on the same endosomes (Fig 3 B-B''''and E). We did note that RAP-1 distribution appeared

broader than that of SDPN-1, as reflected in a lesser but still substantial Manders' coefficient

for the reverse comparison (Fig 3E). RAP-1 labeling of structures that were SDPN-1-negative

could reflect endosomal microdomains lacking SDPN-1, and/or additional RAP-1 positive

organelle types such as late endosomes (40). We also observed considerable overlap between

tagRFP::RAP-1 and basolateral recycling endosome marker GFP::RME-1 (Fig 3C-C'''), and early

endosome marker GFP::RAB-5 (Fig 3D-D'''), further supporting the interpretation that SDPN-1

and RAP-1 are present on the same early and basolateral recycling endosomes where they

could function together to regulate endocytic recycling.

RHO-1 is enriched on SDPN-1 and RAP-1 positive endosomes

Given our findings indicating that RAP-1 signaling functions downstream of SDPN-1 to promote

recycling, we sought to better understand the relevant target of RAP-1 for this process. While

previous studies of Rap-GTPases had not identified a target of Rap-signaling for regulation of

recycling, some reports indicated that mammalian Rap1 or Rap2 can negatively regulate RhoA

to promote mammalian epithelial cell spreading, endothelial barrier function, or ECM stiffness

mechanotransduction, in part via ARHGAP29 (41-45). We hypothesized that misregulation of

endosomal Rho-GTPase activity could explain *sdpn-1/pxf-1/rap-1* associated phenotypes.

Supporting this notion, we found that mutation of *spv-1*, the closest *C. elegans* homolog of

ARFGAP29, resulted in an increase in RME-1 labeled recycling endosome number, similar to the

defect displayed in *pxf-1* and *rap-1* mutants (Fig. S3).

Given these results, we tested for a role of SDPN-1 and RAP-1 in the regulation of endosomal

RHO-1, the only *C. elegans* homolog of RhoA (46). First, we assayed for enrichment of RHO-1 on

the relevant endosomes in the *C. elegans* intestine, the expected location for direct function as

part of the basolateral recycling pathway. Indeed we found that nearly all SDPN-1::GFP positive

structures contained tagRFP::RHO-1 (Fig 4 A-A'''' and 4C), while tagRFP::RHO-1 was also present

on SDPN-1 negative structures (Fig 4 A-A'''' and 4C). Importantly, GFP::RHO-1 and tagRFP::RAP-

1 colocalized very well, with most endosomes positive for one also positive for the other (Fig.4

B-B''' and 4C). Taken together, these results indicate that RHO-1 is enriched on the same

endosomes as SDPN-1 and RAP-1, pointing to the potential for a SDPN-1/PXF-1/RAP-1 module

influencing endosomal RHO-1 activity.

SDPN-1 and RAP-1 negatively regulate RHO-1

If RHO-1 is regulated by SDPN-1 and RAP-1, we might expect to find a change in RHO-1

subcellular localization in *sdpn-1* and *rap-1* mutants. Indeed, we found an increased

accumulation of GFP::RHO-1 in intracellular puncta in both mutants (Figure S4). This result was

suggestive, but was not sufficient to define any changes in RHO-1 GTP-level.

To directly test for effects of *sdpn-1* and *rap-1* mutants on intestinal endosomal RHO-1

activation state, we co-expressed a RHO-1(GTP) biosensor (GFP::AHPH) (47) with tagRFP::RHO-

1, assaying for changes in GFP::AHPH recruitment to tagRFP::RHO-1 labeled endosomes in the

intestine. If SDPN-1 normally acts to negatively regulate RHO-1 activation via RAP-1 signaling,

we would expect *sdpn-1* and *rap-1* mutants to show increased RHO-1 activation, leading to

increased GFP::AHPH recruitment. Indeed, we measured a significant increase in the

colocalization of RHO-1 and its active-state biosensor on intracellular structures in *sdpn-1* or

rap-1 null mutant backgrounds, indicating abnormally high RHO-1(GTP) levels on intestinal

endosomes (Fig. 4D-G). Importantly, there was no change in GFP::AHPH colocalization and

tagRFP::RHO-1 on the apical plasma membrane, indicating the specificity of this regulation (Fig

4H). This data suggests a model in which a critical function of the SDPN-1/PXF-1/RAP-1 pathway

is to prevent the inappropriate overactivation of RHO-1 on endosomes that could interfere with

recycling function.

RHO-1 functions downstream of SDPN-1 to regulate endocytic recycling

A corollary to this model is that reducing RHO-1 levels could be expected to compensate for

elevated RHO-1 activation in *sdpn-1* mutants, leading to rescue of *sdpn-1* endocytic recycling

defects. To test this hypothesis directly, we used RNAi to deplete RHO-1 in a *sdpn-1* mutant

background and assayed for effects on recycling endosomes and endocytic recycling cargo.

Since *rho-1* is an essential gene, we established RNAi conditions that strongly depleted RHO-1,

as verified in a tagRFP::RHO-1 strain (Fig. S5), but allowed development to the adult stage

typically used in our assays. Remarkably, this RNAi-mediated depletion of RHO-1 fully

suppressed *sdpn-1* null mutant defects that normally result in abnormal accumulation of

GFP::RME-1-labeled recycling endosomes and hTAC::GFP recycling cargo (Fig. 6, panel A-B''''').

These results strongly support regulation of endosomal RHO-1 as a key factor in the SDPN-1-

dependent regulation of endocytic recycling.

Non-muscle myosin II is a key regulator of recycling downstream of SDPN-1, RAP-1 and RHO-1

Given our results with RHO-1, we sought greater insight into which RHO-1-mediated activities

influence recycling. *C. elegans* RHO-1 and mammalian RhoA are well known for regulating

actomyosin contractility through non-muscle myosin II (48, 49). In support of a roll for myosin II

in this pathway, we found that SDPN-1 colocalized with MLC-4, the myosin II regulatory light

chain reported to function with non-muscle myosin II heavy chain isoforms NMY-1 and NMY-2

(Fig. 4, panel C-C''' and D). We also found that loss of SDPN-1 increased intracellular

accumulation of punctate MLC-4 intracellular signal suggesting increased recruitment to

endosomal membranes (50).

If regulation of myosin II contraction cycling is a key output of SDPN-1/PXF-1/RAP-1 signaling

for endocytic recycling, forced activation of myosin II activity might be expected interfere with

endosome function. Indeed, we found that forced activation of non-muscle myosin II in the

intestine via expression of a constitutively active phosphomimetic form of MLC-4 (MLC-4(DD))

(50), increased intracellular retention of recycling cargo hTAC::GFP, and resulted in intracellular

accumulation of RAB-5 and RME-1 marked endosomes (Fig. 6, panel E-H; Fig S1C'-C''). These

results support a requirement for myosin II regulation on endosomes for successful endocytic

recycling.

To investigate myosin II requirements further, we assayed for effects of depletion of non-

muscle myosin II isoforms NMY-1 and NMY-2. Importantly, depletion of myosin II isoform NMY-

2 caused intracellular accumulation of recycling endosome marker RME-1 and recycling cargo

hTAC::GFP similar to *sdpn-1* mutants, while depletion of NMY-1 did not (Fig 7 H-J, G). By

contrast, depletion of myosin II isoform NMY-1, but not depletion of NMY-2, strongly

suppressed *sdpn-1* null mutant intracellular accumulation of GFP::RME-1-labeled recycling

endosomes and recycling cargo hTAC::GFP, similar to the effects of RHO-1 depletion (Fig. 7).

Taken together these results strongly support a key role for myosin II regulation on endosomes

in recycling function, with NMY-1 as a likely target of SDPN-1/PXF-1/RAP-1/RHO-1 regulation

(Fig.8).

Discussion

Syndapin/PACSIN proteins have long been implicated in endocytic recycling in multiple

organisms, but a clear understanding of the mechanism of action of these proteins in

endosomal regulation has remained unclear (Braun et al., 2005; Farmer et al., 2021; Gleason et

al., 2016). Some clues are provided by the syndapin/PACSIN domain organization that is

characterized by a single N-terminal F-BAR domain that links these proteins to membranes, and

a single C-terminal SH3 domain that interacts with other proteins. Our new *in vivo* data

reinforces the essential nature of the SH3 domain to syndapin/PACSIN function in recycling,

finding that a point mutation in the SH3 domain of endogenous *sdpn-1* produced endocytic

recycling defects in the *C. elegans* intestine equivalent to those we previously established for

sdpn-1 null mutants. Thus, identifying the key physiologically relevant binding partners of the

SDPN-1 SH3 domain is essential to understanding how SDPN-1 controls recycling.

The *C. elegans* SH3-ome project predicted a high likelihood binding site for the SDPN-1 SH3 in

RAP-GEF PXF-1, an interaction that we confirmed biochemically. We found the importance of

the interaction *in vivo* to be clear, as mutation of this binding site in endogenous *pxf-1* also

severely impaired endocytic recycling, causing intracellular accumulation of intestinal

basolateral recycling cargo and abnormal accumulation of markers of early and basolateral

recycling endosomes. The role of PXF-1 in recycling appears to match its known biochemical

activity as a Rap-GEF, as our results further showed that loss of PXF-1 target RAP-1 also

impaired intestinal recycling, and PXF-1 and RAP-1 are enriched on SDPN-1 labeled endosomes

as would be expected if they function together. Although Rap GTPases have not been well

studied in the context of recycling, mammalian Rap1 and Rap2 have been reported to be

enriched on recycling endosomes, with Rap1 supporting recycling of the GABA(B) receptor in

neurons, and Rap2 supporting recycling of integrin LFA-1 in T lymphocytes (28, 51). Taken

together with our data in *C. elegans* we conclude that a requirement for Rap-GTPases in

endocytic recycling is evolutionarily conserved. Thus, gaining insight into how Rap GTPases are

regulated on recycling endosomes, and understanding the mechanisms by which Rap activation

influences recycling function in *C. elegans* should be broadly applicable to other organisms, and

has important implications for higher order processes known to require Rap signaling, including

cell adhesion, cell migration, cell polarity, and cell fate specification.

What then is the function of RAP-1 signaling from endosomes during recycling? One known

output of Rap signaling is the negative regulation of RhoA, a function first shown during neurite

outgrowth, and also prominent during endothelial barrier formation (41, 52). Indeed, we found

that *C. elegans* RHO-1/RhoA was highly enriched on early and basolateral recycling endosomes

in the adult *C. elegans* intestine, and RHO-1 activation was increased upon loss of SDPN-1 or

RAP-1 as judged by a RHO-1(GTP) biosensor, indicating that SDPN-1 and RAP-1 act as negative

regulators of RHO-1 activity on endosomes. Importantly, our results further revealed RHO-1

negative regulation as a key output of SDPN-1 signaling relevant to recycling, since a reduction

in RHO-1 levels by RNAi strongly suppressed the recycling defects of *sdpn-1* mutants. We note

that recent work in mammalian epithelial cells found cortical recycling endosomes as a major

site of RhoA activation, in this case relevant to stress fiber generation and front-rear polarity

acquisition, although this work did not test for effects of RhoA activation on recycling function

(53). These similarities further suggest that our findings will be relevant in a broad context

across species.

We further sought to determine which output of RHO-1 signaling was most relevant to SDPN-1-

mediated recycling regulation. RhoA is a well-established regulator of actomyosin contractility

that functions through effectors mDia1 (a diaphanous-related formin) and ROCK (Rho-

associated serine/threonine kinase), with ROCK activating non-muscle myosin II and actin

dynamics (48). Our experiments found that depletion of the NMY-1 isoform of non-muscle

myosin II was sufficient to significantly suppress *sdpn-1* mutant defects. These results indicate

that inappropriately high levels of active myosin II, or the inability to move through complete

myosin contractility cycles, explains at least part of the effects of *sdpn-1* mutants on recycling

(Gleason et al., 2016; Yan et al., 2021). This interpretation is supported by our results showing

that constitutive activation of myosin II by expression of a myosin light chain phosphomimetic

also impairs recycling function, and our findings that myosin light chain is enriched on SDPN-1

positive endosomes.

Surprisingly, we found the opposite phenotype for non-muscle myosin isoform NMY-2.

Depletion of NMY-2 caused intracellular accumulation of recycling cargo, indicating a positive

role for NMY-2 in recycling. Interestingly, previous work on NMY-1 and NMY-2 documented

opposing effects in the regulation of the contractile ring channels of the *C. elegans* gonad (22).

Mathematical modeling in that work suggested that these opposing effects could be explained

by NMY-1 and NMY-2 contractile fibers exerting forces in opposing directions, one normal to

the ring, and the other tangential (22). If these principles apply to the recycling endosome case,

NMY-2 derived forces might increase tension on endosomal tubules to promote their fission,

while NMY-1 contraction in an opposing direction might reduce tubule tension and fission.

Normal recycling tubule release might require cycles of increased and decreased tension, or

forces regulating recycling rates might be modulated in response to changes in cellular

requirements, such as changes in cargo load or metabolism.

Interestingly, three putative Rho effectors, formin CYK-1/mDia, LET-502/ROCK, and RTKN-

1/Rhotekin, are known to function in basolateral recycling in the *C. elegans* intestine, although

current models posit that these proteins function independently of RHO-1 in recycling (11, 54,

55). Branched actin formed by Arp2/3 is traditionally considered the relevant F-actin type

associated with endosomes, but recycling may require a balance of branched and linear actin

for proper function. Actin tracks produced by CYK-1 could be substrates upon which cycles of

myosin II contractility would act, potentially producing cycles of actomyosin-driven endosomal

membrane tension that could promote cycles of tubular cargo carrier fission.

We note that non-muscle myosin II was previously shown to promote mitochondrial fission and

could contribute to endosomal fission during recycling in a similar way (56-59). Myosin II,

Spire1C, and endoplasmic reticulum-bound formin INF2 have been proposed to create

actomyosin tension that deforms the outer mitochondrial membrane during "prefission",

promoting curvature-sensing-based recruitment of dynamin-related protein Drp1 and other

factors that complete mitochondrial constriction (59). Further work will be required to

determine if a similar process is operative at the early endosome – recycling endosome

interface, and if there is a relationship between SDPN-1/RAP-1/RHO-1/NMY-1 and previously

identified functions for LET-502/RTKN-1/CYK-1 during recycling.

Syndapin/PACSIN proteins are often thought of as acting directly in endosomal tubule

generation or tubule severing, since isolated BAR domains from several of these proteins are

able to generate tubules from liposomes *in vitro*, and in some cases enhance endosomal

tubulation when overexpressed *in vivo*. However, it remains unclear if an F-BAR-driven

constricting/tubulating activity is a key output of syndapin/PACSINs that promotes recycling

function. An alternate hypothesis is that the key function of the F-BAR domain in

syndapin/PACSINs is to provide membrane binding and curvature-sensing that brings

syndapin/PACSIN to endosomal tubules elaborated by other means, such as cytoskeletal

pushing and pulling forces. In the *C. elegans* intestine and human HeLa cells RAB-10/Rab10 is

essential for the extensive tubulation found in the recycling endosome network (10, 60-64). In

the *C. elegans* case microfilament and microtubule integrity is required for network structure,

and RAB-10 is known to regulate association of endosomal membranes with actin filaments via

the EHBP-1 protein (60, 63, 64). In the HeLa case Rab10-mediated regulation of Kif13/kinesin

pulling forces have also been implicated (61).

Our new data clearly points to an essential role for Syndapin in signaling via Rap and Rho to

control myosin II on endosomes (Fig.8), but we cannot rule out an additional mechanical

contribution of SDPN-1 to endosomal function via direct membrane bending. However, the high

efficiency of *sdpn-1* mutant recycling phenotype suppression by *rho-1* depletion suggests that

RHO-1 regulation is the main output relevant to recycling. We also note that in some systems

syndapin/PACSIN proteins associate with other trafficking regulators such as MICAL-L1, OCRL1,

Dynamin, and EHD1/RME-1 that could promote tubulation and/or severing (62). It will be of

great interest to revisit the question of relative contributions to the tubule-based recycling

mechanism as new genetic and cell biological tools evolve, allowing more complete, acute, and

fine-tuned control of experimental conditions, as we further elaborate our understanding of

this complex system.

Acknowledgments

We would like to thank the Grant lab for critical comments and suggestions. We thank Peter

Schweinsberg and Ge Bai for technical assistance in making plasmid clones and biolistic

transgenic lines, and Helen Ushakov for expert microinjection. We thank Adriana Dawes for

nmy-1 and *nmy-2* RNAi clones, Michel Labouesse for *mlc-4* wild-type and phosphomimetic DNA

clones, and David Reiner for *rap-1* mutant strains. We also thank Michael Pierce, Noriko

Goldsmith and Jessica Shivas for help with confocal microscopy, and Gary Bader for SH3

weighted match analysis. This work was supported by NIH Grant 5R01GM135326 to B.D.G.

References

1.   G. J. Doherty, H. T. McMahon, Mechanisms of endocytosis. *Annu Rev Biochem* **78**, 857-902
        (2009).

2.   L. Maldonado-Báez, C. Williamson, J. G. Donaldson, Clathrin-independent endocytosis: a cargo-
        centric view. *Exp Cell Res* **319**, 2759-2769 (2013).

3.   J. A. Solinger, A. Spang, Sorting of cargo in the tubular endosomal network. *Bioessays* **44**,
        e2200158 (2022).

4.   B. D. Grant, J. G. Donaldson, Pathways and mechanisms of endocytic recycling. *Nat Rev Mol Cell
        Biol* **10**, 597-608 (2009).

5.   J. D. McGhee, The C. elegans intestine. *WormBook* 10.1895/wormbook.1.133.1, 1-36 (2007).

6.   B. Leung, G. J. Hermann, J. R. Priess, Organogenesis of the Caenorhabditis elegans intestine. *Dev*

Biol **216**, 114-134 (1999).

7.   B. Grant *et al.*, Evidence that RME-1, a conserved C. elegans EH-domain protein, functions in

endocytic recycling. *Nat Cell Biol* **3**, 573-579 (2001).

8.   K. Sato, A. Norris, M. Sato, B. D. Grant, C. elegans as a model for membrane traffic. *WormBook*

10.1895/wormbook.1.77.2, 1-47 (2014).

9.   A. M. Gleason, K. C. Nguyen, D. H. Hall, B. D. Grant, Syndapin/SDPN-1 is required for endocytic

recycling and endosomal actin association in the C. elegans intestine. *Mol Biol Cell* **27**, 3746-

3756 (2016).

10.   C. C. Chen *et al.*, RAB-10 is required for endocytic recycling in the Caenorhabditis elegans

intestine. *Mol Biol Cell* **17**, 1286-1297 (2006).

11.   Y. Yan *et al.*, RTKN-1/Rhotekin shields endosome-associated F-actin from disassembly to ensure

endocytic recycling. *J Cell Biol* **220** (2021).

12.   E. Casal *et al.*, The crystal structure of the BAR domain from human Bin1/amphiphysin II and its

implications for molecular recognition. *Biochemistry* **45**, 12917-12928 (2006).

13.   A. Braun *et al.*, EHD proteins associate with syndapin I and II and such interactions play a crucial

role in endosomal recycling. *Mol Biol Cell* **16**, 3642-3658 (2005).

14.   A. Quan, P. J. Robinson, Syndapin--a membrane remodelling and endocytic F-BAR protein. *Febs j*

280, 5198-5212 (2013).

15.   J. Kovár, Hybridoma cultivation in defined serum-free media: growth-supporting substances. II.

Insulin, other hormones, and growth factors. *Folia Biol (Praha)* **32**, 304-310 (1986).

16.   X. Xin *et al.*, SH3 interactome conserves general function over specific form. *Mol Syst Biol* **9**, 652

(2013).

17.   W. Pellis-van Berkel *et al.*, Requirement of the Caenorhabditis elegans RapGEF pxf-1 and rap-1

524   for epithelial integrity. *Mol Biol Cell* **16**, 106-116 (2005).

525   18.   M. R. Kooistra, N. Dubé, J. L. Bos, Rap1: a key regulator in cell-cell junction formation. *J Cell Sci*

526   **120**, 17-22 (2007).

527   19.   C. Frøkjær-Jensen *et al.*, Random and targeted transgene insertion in Caenorhabditis elegans

528   using a modified Mos1 transposon. *Nat Methods* **11**, 529-534 (2014).

529   20.   L. Timmons, A. Fire, Specific interference by ingested dsRNA. *Nature* **395**, 854 (1998).

530   21.   R. S. Kamath, J. Ahringer, Genome-wide RNAi screening in Caenorhabditis elegans. *Methods* **30**,

531   313-321 (2003).

532   22.   V. C. Coffman, T. M. Kachur, D. B. Pilgrim, A. T. Dawes, Antagonistic Behaviors of NMY-1 and

533   NMY-2 Maintain Ring Channels in the C. elegans Gonad. *Biophys J* **111**, 2202-2213 (2016).

534   23.   A. Paix, A. Folkmann, G. Seydoux, Precision genome editing using CRISPR-Cas9 and linear repair

535   templates in C. elegans. *Methods* **121-122**, 86-93 (2017).

536   24.   G. Huang *et al.*, Improved CRISPR/Cas9 knock-in efficiency via the self-excising cassette (SEC)

537   selection method in C. elegans. *MicroPubl Biol* **2021** (2021).

538   25.   M. L. Schwartz, M. W. Davis, M. S. Rich, E. M. Jorgensen, High-efficiency CRISPR gene editing in

539   C. elegans using Cas9 integrated into the genome. *PLoS Genet* **17**, e1009755 (2021).

540   26.   Q. Wang *et al.*, Molecular mechanism of membrane constriction and tubulation mediated by the

541   F-BAR protein Pacsin/Syndapin. *Proc Natl Acad Sci U S A* **106**, 12700-12705 (2009).

542   27.   H. Meng *et al.*, PACSIN 2 represses cellular migration through direct association with cyclin D1

543   but not its alternate splice form cyclin D1b. *Cell Cycle* **10**, 73-81 (2011).

544   28.   P. Stanley, S. Tooze, N. Hogg, A role for Rap2 in recycling the extended conformation of LFA-1

545   during T cell migration. *Biol Open* **1**, 1161-1168 (2012).

546   29.   F. Balzac *et al.*, E-cadherin endocytosis regulates the activity of Rap1: a traffic light GTPase at the

547   crossroads between cadherin and integrin function. *J Cell Sci* **118**, 4765-4783 (2005).

548   30.   L. Li *et al.*, A unique interplay between Rap1 and E-cadherin in the endocytic pathway regulates

549   self-renewal of human embryonic stem cells. *Stem Cells* **28**, 247-257 (2010).

550   31.   Y. Liao *et al.*, RA-GEF, a novel Rap1A guanine nucleotide exchange factor containing a

551   Ras/Rap1A-associating domain, is conserved between nematode and humans. *J Biol Chem* **274**,

552   37815-37820 (1999).

553   32.   H. B. Kuiperij *et al.*, Characterisation of PDZ-GEFs, a family of guanine nucleotide exchange

554   factors specific for Rap1 and Rap2. *Biochim Biophys Acta* **1593**, 141-149 (2003).

555   33.   B. Qualmann, J. Roos, P. J. DiGregorio, R. B. Kelly, Syndapin I, a synaptic dynamin-binding protein

556   that associates with the neural Wiskott-Aldrich syndrome protein. *Mol Biol Cell* **10**, 501-513

557   (1999).

558   34.   A. Shi *et al.*, RAB-10-GTPase-mediated regulation of endosomal phosphatidylinositol-4,5-

559   bisphosphate. *Proc Natl Acad Sci U S A* **109**, E2306-2315 (2012).

560   35.   R. J. Gleason, A. M. Akintobi, B. D. Grant, R. W. Padgett, BMP signaling requires retromer-

561   dependent recycling of the type I receptor. *Proc Natl Acad Sci U S A* **111**, 2578-2583 (2014).

562   36.   E. W. Frische *et al.*, RAP-1 and the RAL-1/exocyst pathway coordinate hypodermal cell

563   organization in Caenorhabditis elegans. *Embo j* **26**, 5083-5092 (2007).

564   37.   J. F. Rebhun, A. F. Castro, L. A. Quilliam, Identification of guanine nucleotide exchange factors

565   (GEFs) for the Rap1 GTPase. Regulation of MR-GEF by M-Ras-GTP interaction. *J Biol Chem* **275**,

566   34901-34908 (2000).

567   38.   N. R. Rasmussen, D. J. Dickinson, D. J. Reiner, Ras-Dependent Cell Fate Decisions Are Reinforced

568   by the RAP-1 Small GTPase in Caenorhabditiselegans. *Genetics* **210**, 1339-1354 (2018).

569   39.   Y. Feng, B. G. Williams, F. Koumanov, A. J. Wolstenholme, G. D. Holman, FGT-1 is the major

570   glucose transporter in C. elegans and is central to aging pathways. *Biochem J* **456**, 219-229

571   (2013).

572   40.   V. Pizon, M. Desjardins, C. Bucci, R. G. Parton, M. Zerial, Association of Rap1a and Rap1b

573   proteins with late endocytic/phagocytic compartments and Rap2a with the Golgi complex. *J Cell*

574   *Sci* **107 (Pt 6)**, 1661-1670 (1994).

575   41.   A. Post, W. J. Pannekoek, B. Ponsioen, M. J. Vliem, J. L. Bos, Rap1 Spatially Controls ArhGAP29 To

576   Inhibit Rho Signaling during Endothelial Barrier Regulation. *Mol Cell Biol* **35**, 2495-2502 (2015).

577   42.   A. Post *et al.*, Rasip1 mediates Rap1 regulation of Rho in endothelial barrier function through

578   ArhGAP29. *Proc Natl Acad Sci U S A* **110**, 11427-11432 (2013).

579   43.   G. A. Smolen *et al.*, A Rap GTPase interactor, RADIL, mediates migration of neural crest

580   precursors. *Genes Dev* **21**, 2131-2136 (2007).

581   44.   C. W. Wilson *et al.*, Rasip1 regulates vertebrate vascular endothelial junction stability through

582   Epac1-Rap1 signaling. *Blood* **122**, 3678-3690 (2013).

583   45.   Z. Meng *et al.*, RAP2 mediates mechanoresponses of the Hippo pathway. *Nature* **560**, 655-660

584   (2018).

585   46.   B. Saenz-Narciso, E. Gomez-Orte, A. Zheleva, I. Gastaca, J. Cabello, Control of developmental

586   networks by Rac/Rho small GTPases: How cytoskeletal changes during embryogenesis are

587   orchestrated. *Bioessays* **38**, 1246-1254 (2016).

588   47.   Y. C. Tse *et al.*, RhoA activation during polarization and cytokinesis of the early Caenorhabditis

589   elegans embryo is differentially dependent on NOP-1 and CYK-4. *Mol Biol Cell* **23**, 4020-4031

590   (2012).

591   48.   M. Chircop, Rho GTPases as regulators of mitosis and cytokinesis in mammalian cells. *Small*

592   *GTPases* **5** (2014).

593   49.   B. Sáenz-Narciso, E. Gómez-Orte, A. Zheleva, I. Gastaca, J. Cabello, Control of developmental

594   networks by Rac/Rho small GTPases: How cytoskeletal changes during embryogenesis are

595   orchestrated. *Bioessays* **38**, 1246-1254 (2016).

596   50.   C. Gally *et al.*, Myosin II regulation during C. elegans embryonic elongation: LET-502/ROCK,

597   MRCK-1 and PAK-1, three kinases with different roles. *Development* **136**, 3109-3119 (2009).

598   51.   Z. Zhang *et al.*, GABAB receptor promotes its own surface expression by recruiting a Rap1-

599   dependent signaling cascade. *J Cell Sci* **128**, 2302-2313 (2015).

600   52.   T. Yamada, T. Sakisaka, S. Hisata, T. Baba, Y. Takai, RA-RhoGAP, Rap-activated Rho GTPase-

601   activating protein implicated in neurite outgrowth through Rho. *J Biol Chem* **280**, 33026-33034

602   (2005).

603   53.   C. Gaston *et al.*, EpCAM promotes endosomal modulation of the cortical RhoA zone for

604   epithelial organization. *Nat Commun* **12**, 2226 (2021).

605   54.   T. Gong *et al.*, PTRN-1/CAMSAP promotes CYK-1/formin-dependent actin polymerization during

606   endocytic recycling. *Embo j* **37** (2018).

607   55.   W. Zhang *et al.*, LET-502/ROCK Regulates Endocytic Recycling by Promoting Activation of RAB-5

608   in a Distinct Subpopulation of Sorting Endosomes. *Cell Rep* **32**, 108173 (2020).

609   56.   F. Korobova, T. J. Gauvin, H. N. Higgs, A role for myosin II in mammalian mitochondrial fission.

610   *Curr Biol* **24**, 409-414 (2014).

611   57.   U. Manor *et al.*, A mitochondria-anchored isoform of the actin-nucleating spire protein regulates

612   mitochondrial division. *Elife* **4** (2015).

613   58.   A. L. Hatch, P. S. Gurel, H. N. Higgs, Novel roles for actin in mitochondrial fission. *J Cell Sci* **127**,

614   4549-4560 (2014).

615   59.   C. Yang, T. M. Svitkina, Ultrastructure and dynamics of the actin-myosin II cytoskeleton during

616   mitochondrial fission. *Nat Cell Biol* **21**, 603-613 (2019).

60.   A. Shi *et al.*, EHBP-1 functions with RAB-10 during endocytic recycling in Caenorhabditis elegans.

Mol Biol Cell **21**, 2930-2943 (2010).

61.   K. Etoh, M. Fukuda, Rab10 regulates tubular endosome formation through KIF13A and KIF13B

motors. *J Cell Sci* **132** (2019).

62.   T. Farmer *et al.*, Defining the protein and lipid constituents of tubular recycling endosomes. *J*

Biol Chem **296**, 100190 (2021).

63.   P. Wang *et al.*, RAB-10 Promotes EHBP-1 Bridging of Filamentous Actin and Tubular Recycling

Endosomes. *PLoS Genet* **12**, e1006093 (2016).

64.   S. Chen *et al.*, SEC-10 and RAB-10 coordinate basolateral recycling of clathrin-independent cargo

through endosomal tubules in Caenorhabditis elegans. *Proc Natl Acad Sci U S A* **111**, 15432-

15437 (2014).

Figure S1. Additional quantification related to main figures 2, 5, and 6. (A-A'') Increased intracellular area labeled by GFP::RAB-5 and increased number of intracellular recycling compartments in interaction-defective mutants *sdpn-1(pw33)* and *pxf-1(pw34)*, as well as null *rap-1(tm861)* and activated *rap-1(re180)* (*rap-1 G12V*) mutants (related to Main Fig. 2A-C''''). **(B)** Quantification of the number of intracellular mScarlet::MLC-4 regulatory myosin light chain puncta (related to Main Fig. 6C-C'). **(C-C')** Quantification of GFP::RME-1 labeled recycling compartments and intracellular area positive for early endosome marker GFP::RAB-5 (related to main Fig. 6E-H). Error bars represent SEM: *p< 0.05, **p< 0.01, **** p< 0.0001 (Two-tailed Welch's *t* test).

Figure S2. Intracellular accumulation of glucose transporter FGT-1 in *sdpn-1* and *rap-1 null* mutants. All micrograph images were obtained by confocal laser-scanning microscopy in intact living animals expressing intestine-specific GFP-tagged protein. **(A-A'')**. Representative images are shown for FGT-1::GFP in wild-type, *sdpn-1(ok1667),* and *rap-1(tm861)* mutants. Whole image scale bar = 10 μm. The panel at the bottom left of the image is the magnified section inside the white rectangle, scale bar = 1 μm. **(B)** Quantification of total intracellular fluorescence intensity. There is a significant increase in intracellular FGT-1::GFP accumulation in a *sdpn-1(0)* and *rap-1(0)* mutants compared to *wild-type*. Each data point represents one animal. Error bars represent SEM: ** p<0.01, **** p< 0.0001 (Two-tailed Welch's *t* test).

Figure S3. Loss of the SPV-1 Rho-GAP causes intracellular accumulation of RME-1-positive recycling endosomes. All micrograph images were obtained by confocal laser-scanning microscopy in intact living animals expressing intestine-specific GFP-tagged protein. **(A-A')** Representative images are shown for GFP::RME-1 labeled recycling endosome in wild-type and *spv-1(ok1498)* mutant animals. Whole image scale bar = 10 μm. The panel at the bottom left of the image is the magnified section inside the white rectangle, scale bar = 1 μm. **(B)** Quantification of total intracellular fluorescence intensity of recycling endosomes. **(C)** Quantification of the number of intracellular recycling endosomes. There is a significant increase in RME-1-positive recycling endosomes in *spv-1* mutants. Each data point represents one animal. Error bars represent SEM: * p< 0.05, **** p< 0.0001 (Two-tailed Welch's *t* test).

Figure S4. Intracellular RHO-1 signal increases in *sdpn-1* and *rap-1 null* mutants. All micrograph images were obtained via confocal laser-scanning microscopy in intact living animals expressing intestine-specific GFP-tagged protein. **(A-A'')** Whole image scale bar = 10 µm. The panel at the bottom left of the image is the magnified section inside the white rectangle, scale bar = 1 µm. **(B)** Quantification of total intracellular fluorescence intensity of RHO-1. **(C)** Quantification of the number of intracellular RHO-1 puncta. Intracellular puncta signal for GFP::RHO-1 increases significantly in *sdpn-1(ok1667)* and *rap-1(tm861)* mutants. Each data point represents one animal. Error bars represent SEM: * p< 0.05, ** p< 0.01, *** p< 0.001, **** p< 0.0001 (Two-tailed Welch's *t* test).

Figure S5. Intestinal RHO-1 was depleted by RHO-1 RNAi. All micrographs are from confocal image stacks acquired in intact living animals expressing intestine-specific RFP-tagged protein. **(A)** tagRFP::RHO-1 expression on a wild-type background. **(A')** tagRFP::RHO-1 is depleted by rho-1 RNAi. Scale bar = 10 μm. In each image, the blue channel denotes autofluorescent lysosome-like organelles, whereas the red channel denotes RFP. The red channel signal that does not overlap with the blue channel signal is considered bona fide RFP signal respectively. **(B)** Quantification of total fluorescence intensity. Each data point represents one animal. Error bars represent SEM: **** $p < 0.0001$ (Two-tailed Welch's *t* test).

Figure 1. The SDPN-1 SH3 domain binds to PXF- 1. (A) SDPN-1 SH3 domain consensus target sequence; red arrows indicate conserved residues that were mutated to alanine (*pxf-1(pw34)*) to disrupt interaction with the SDPN-1 SH3 domain. **(B)** Schematic representation of SDPN-1 and PXF-1 domain structure from the N-terminal to the C-terminal; the blue arrow indicates the *sdpn-1(pw33)* mutation site and the red arrow indicates the *pxf-1(pw34)* mutation site.   **(C)** Glutathione beads loaded with recombinant GST or the GST-SDPN-1 SH3 domain were incubated with *in vitro* translated HA-tagged PXF-1 *wild-type* or aa191-198 (**K**GPA**P**LP**P** to **A**GPA**A**LP**A**). Bound proteins were eluted and analyzed by Western blot probed with anti-HA antibody. The *wild-type* HA-tagged PXF-1 was bound by *wild-type* GST-SDPN-1-SH3, but not by mutant (P478L), or unfused GST. The (KGPAPLPP to AGPAALPA) mutant HA-tagged PXF-1 failed to bind GST alone or GST-SDPN-1-SH3 *wild-type* and mutant domains. Total GST and GST-SDPN-1-SH3 bait proteins were visualized by Ponceau Staining prior to antibody staining.

Figure 2. SDPN-1 interaction with PXF-1, and RAP-1 is important for proper endocytic recycling. All images were obtained by confocal laser-scanning microscopy in intact living animals expressing intestine-specific GFP-tagged proteins. Representative confocal images of GFP::RAB-5 expressed in a *wild-type* **(A)**, *sdpn-1(pw33)* **(A')**, *pxf-1(pw34)* **(A'')**, and *rap-1(0)* **(A''')** background are shown. SDPN-1 and PXF-1 interacting-defective mutations, and loss of RAP-1, causes intracellular accumulation of compartments labeled by early endosome marker GFP::RAB-5 when compared with wild-type. In addition, GFP::RAB-5 expressed in a wild-type **(B)**, and *rap-1(re180)* GTP-locked mutant **(B')** background also showed an increase in the intracellular compartments labeled by early endosome marker GFP::RAB-5. Confocal images of GFP::RME-1 basolateral recycling endosome marker expressed in a *wild-type* **(C)**, *sdpn-1(pw33)* **(C')**, *pxf-1(pw34)* **(C'')**, *rap-1(0)* **(C''')**, and *rap-1(re180)* **(C'''')** background are shown. There is an increase of intracellular recycling compartments signal when SDPN-1 and PXF-1 interaction is lost, and when RAP-1 is missing or is GTP-locked. Micrographs showing representative confocal images of model GFP-tagged recycling cargo protein, the human interleukin-2 receptor (hTAC::GFP), expressed in a *wild-type* **(D)**, *sdpn-1(pw33)* **(D')**, *pxf-1(pw34)* **(D'')**, *rap-1(0)* **(D''')**, and *rap-1(re180)* **(D'''')** background. Micrographs showing representative confocal images of another model GFP-tagged recycling cargo protein, the human transferrin receptor (hTFR::GFP), expressed in a *wild-type* **(E)**, *sdpn-1(pw33)* **(E')**, *pxf-1(pw34)* **(E'')**, *rap-1(0)* **(E''')**, and *rap-1(re180)* **(E'''')** background. Losing the interaction between SDPN-1 and PXF-1, and when RAP-1 is missing or is GTP-locked, displays a significant increase in the intracellular accumulation of both recycling cargoes hTAC::GFP and hTFR::GFP. Main panel scale bars = 10 μm. Magnified inset panel scale bars = 1 μm. **(F)** Quantification of total fluorescence intensity for **(A-A''')**. **(G)** Quantification of total fluorescence intensity for **(B-B')**. **(H)** Quantification of total fluorescence intensity for **(C-C'''')**. **(I)** Quantification of total fluorescence intensity for **(D-D'''')**. **(J)** Quantification of total fluorescence intensity for **(E-E'''')**. **(K)** *C. elegans* intestinal diagram. Each data point represents one animal. Error bars represent SEM: *p< 0.05, **p< 0.01, *** p< 0.001, **** p< 0.0001 (Two-tailed Welch's *t* test).

Figure 3. PXF-1 and RAP-1 are enriched on SDPN-1 positive endosomes. All micrographs are from three-dimensional (3D) confocal image stacks acquired in intact living animals expressing intestine-specific GFP and RFP-tagged proteins. **(A-A'')** GFP::PXF-1 is found on SDPN-1 positive compartments. **(A''')** Magnified image from the section designated by the white rectangle in panel A''. **(B-B'')** tagRFP::RAP-1 colocalizes with SDPN-1 positive endosomes. **(B''')** Magnified image from the section designated by the white rectangle in panel B''. **(C-C'')** tagRFP::RAP-1 is present in GFP::RME-1-labeled basolateral recycling endosomes. **(C''')** Magnified image from the section designated by the white rectangle in panel C''. **(D-D'')** There is colocalization between tagRFP::RAP-1 and GFP::RAB-5-labeled early endosomes. Scale bar = 10 μm. **(D''')** Magnified image from the section designated by the white rectangle in panel D''. Scale bar = 1 μm. In each image, the blue channel denotes autofluorescent lysosome-like organelles, whereas the green channel denotes GFP and the red channel denotes RFP. The green channel and red channel signal that do not overlap with the blue channel signal are considered *bona fide* GFP or RFP signals respectively. **(E)** Manders coefficient for colocalization of GFP::PXF-1 with tagRFP::SDPN-1, and tagRFP::RAP-1 with SDPN-1::GFP, GFP::RME-1 and GFP::RAB-5. Each data point represents one animal.

Figure 4. SDPN-1 and RAP-1 negatively regulate intracellular RHO-1 activation. All micrographs are from three-dimensional (3D) confocal image stacks acquired in intact living animals expressing intestine-specific GFP and RFP-tagged proteins. **(A-A'')** tagRFP::RHO-1 colocalizes extensively with SDPN-1 positive endosomes. **(A''')** Magnified image from the section designated by the white rectangle in panel **(A'')**. **(B-B'')** GFP::RHO-1 colocalizes with RAP-1 positive endosomes. Scale bar = 10 μm. **(B''')** Magnified image from the section designated by the white rectangle in panel **(B'')**. Scale bar = 1 μm. **(C)** Manders coefficient for colocalization of SDPN-1::GFP with tagRFP::RHO-1 and GFP::RHO-1 with tagRFP::RAP-1. **(D-F'')** Imaging of active RHO-1(GTP) via colocalization of RHO-1(GTP) biosensor AHPH with tagRFP::RHO-1 in wild-type, *sdpn-1(ok1667)*, and *rap-1(tm861)* backgrounds. Scale bar = 10 μm. **(D''-F'')** Magnified images from the section designated by the white rectangle in panel. Scale bar = 1 μm. **(D'''-F''')**. In each image, the blue channel denotes autofluorescent lysosome-like organelles, whereas the green channel denotes GFP and the red channel denotes RFP. The green channel and red channel signal that do not overlaps with the blue channel signal are considered bona fide GFP or RFP signals respectively. **(G)** Manders coefficient for intracellular colocalization of AHPH with RHO-1 in a *wild-type*, *sdpn-1 null* and *rap-1 null* background. Loss of SDPN-1 or RAP-1 results in an increased colocalization, indicating increased activation of RHO-1 in *sdpn-1* and *rap-1* mutants. **(H)** Manders coefficient for apical colocalization of AHPH with RHO-1 in a *wild-type*, *sdpn-1 null* and *rap-1 null* background. AHPH recruitment to apical RHO-1 is not increased in *sdpn-1* and *rap-1* mutants, supporting that the key site of action of SDPN-1 on RHO-1 is the endosomes and not the apical plasma membrane. Each data point represents one animal. Error bars represent SEM: ns (non-significant), **p< 0.01, **** p< 0.0001 (Two-tailed Welch's *t* test).

Figure 5. Depletion of RHO-1 suppresses *sdpn-1 null* mutant endocytic recycling phenotypes. All micrograph images were captured using confocal laser-scanning microscopy in intact living animals expressing intestine-specific GFP-tagged proteins. **(A-A')** GFP::RME-1 positive recycling endosomes visualized in otherwise *wild-type* animals after *empty-vector RNAi* or *rho-1* RNAi. **(A''-A''')** GFP::RME-1 positive recycling endosomes visualized in *sdpn-1(ok1667)* mutant animals after *empty-vector RNAi* or *rho-1* RNAi. **(B-B')** Recycling cargo hTAC::GFP was visualized in otherwise *wild-type* animals after *empty-vector RNAi* or *rho-1* RNAi. **(B''-B''')** Recycling cargo hTAC::GFP was visualized in *sdpn-1(ok1667)* mutant animals after *empty-vector RNAi* or *rho-1* RNAi. Whole image scale bar = 10 μm. The panel at the bottom left of the image is the magnified section inside the white rectangle, scale bar = 1 μm. **(C)** Quantification of intracellular total fluorescence intensity for **(A-A''')**. *sdpn-1(ok1667)* mutant animals display an intracellular increase of GFP::RME-1-positive recycling endosomes. This phenotype is suppressed by *rho-1 RNAi* knockdown. **(D)** Quantification of intracellular total fluorescence intensity for **(B-B''')**. *sdpn-1(ok1667)* mutant animals accumulate abnormally high levels of intracellular recycling cargo hTAC::GFP. This phenotype is suppressed by *rho-1 RNAi* knockdown. Each data point represents one animal. Error bars represent SEM: ns (non-significant), *p< 0.05, ***p< 0.001 (Two-tailed Welch's *t* test).

Figure 6. Non-muscle myosin II associates with SDPN-1-positive endosomes and modulates their function. (A-A'') Confocal images comparing the subcellular localization of SDPN-1::GFP-positive endosomes and myosin light chain mScarlet::MLC-4 in the intestine of intact living animals. Most of SDPN-1 positive compartments are enriched in myosin light chain (MLC-4). Scale bar = 10 μm. **(A''')** Magnified image from the section designated by the white rectangle in panel **(A'')**. Scale bar = 1 μm. In each image, the blue channel denotes autofluorescent lysosome-like organelles, whereas the green channel denotes GFP and the red channel denotes mScarlet. The green channel and red channel signal that does not overlaps with the blue channel signal are considered *bona fide* GFP or mScarlet signals respectively. **(B)** Manders coefficient for colocalization of SDPN-1::GFP with mScarlet::MLC-4 indicates strong enrichment for MLC-4 on endosomes. **(C-C')** The signal of intracellular myosin light chain (mScarlet::MLC-4) positive compartments increases in a *sdpn-1(ok1667)* mutant background. **(D)** Quantification intracellular total fluorescence intensity for **(C-C')**. **(E-H)** Animals expressing a constitutively active myosin light chain phospho-mimetic protein MLC-4DD in the intestine display increased intracellular accumulation of early and recycling endosomes. **(I-J)** Animals expressing a constitutively active myosin light chain phospho-mimetic protein MLC-4DD in the intestine display an increase in intracellular recycling cargo hTAC::GFP. Each data point represents one animal. Error bars represent SEM: **p< 0.01, **** p< 0.0001 (Two-tailed Welch's *t* test).

Figure 7. Opposing requirements for non-muscle myosin heavy chains on endocytic recycling. All micrograph images were captured using confocal laser-scanning microscopy in intact living animals expressing intestine-specific GFP-tagged proteins. **(A-C)** GFP::RME-1 positive recycling endosomes visualized in otherwise *wild-type* animals after *empty-vector* RNAi, *mny-1* RNAi or *nmy-2* RNAi. **(D-F)** GFP::RME-1 positive recycling endosomes visualized in *sdpn-1(ok1667)* mutant animals after *empty-vector* RNAi, *nmy-1* RNAi or *nmy-2* RNAi. Whole image scale bar = 10 μm. The panel at the bottom left of the image is the magnified section inside the white rectangle, scale bar = 1 μm. **(G)** Quantification of intracellular total fluorescence intensity for **(A-F)**. *sdpn-1(ok1667)* mutant animals display an increase of intracellular GFP::RME-1-positive recycling endosomes. This phenotype is suppressed by *nmy-1 RNAi* knockdown, but phenocopied by *nmy-2 RNAi* knockdown. **(H-J)** Recycling cargo hTAC::GFP visualized in otherwise *wild-type* animals after *empty-vector* RNAi, *mny-1* RNAi or *nmy-2* RNAi. **(K-M)** Recycling cargo hTAC::GFP visualized in *sdpn-1(ok1667)* mutant animals after *empty-vector* RNAi, *nmy-1* RNAi or *nmy-2* RNAi. Whole image scale bar = 10 μm. The panel at the bottom left of the image is the magnified section inside the white rectangle, scale bar = 1 μm. **(N)** Quantification of intracellular total fluorescence intensity for **(H-M)**. *sdpn-1(ok1667)* mutant animals accumulate abnormally high levels of intracellular recycling cargo hTAC::GFP. This phenotype is suppressed by *nmy-1 RNAi* knockdown, but phenocopied by *nmy-2 RNAi* knockdown. Each data point represents one animal. Error bars represent SEM: ns (non-significant), *p< 0.05, **p< 0.01, ***p< 0.001, ****p< 0.0001 (Two-tailed Welch's *t* test).

Figure 8. Model: SDPN-1 differentially regulates Non-Muscle Myosins via RAP-1 to RHO-1 signaling to regulate endocytic recycling. Recycling tubules pulled from sorting endosomes require non-muscle myosin NMY-2, likely for tubule fission. The F-BAR protein SDPN-1/Syndapin and its binding partner PXF-1 activate RAP-1/Rap to restrain the level of RHO-1/RhoA and non-muscle myosin NMY-1 activity on the endosome. Such negative regulation is required for successful recycling.