CRISPR/Cas9-based edition of frataxin gene in *Dictyostelium discoideum* for Friedreich's Ataxia disease modeling

Authors:
Gentili, Hernan G.[1]; Pignataro, María Florencia[1]; Olmos, Justo[1]; Pavan, Florencia[2]; Ibáñez, Itati[2]; Santos, Javier[1,3] * Velázquez Duarte, Francisco[1, *]

[1] Instituto de Biociencias, Biotecnología y Biología Traslacional (iB3). Departamento de Fisiología y Biología Molecular y Celular, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires. Intendente Güiraldes 2160, Ciudad Universitaria, C1428EGA, Buenos Aires, Argentina.

[2] Instituto de Química Física de los Materiales, Medio Ambiente y Energía (INQUI-MAE), CONICET, FCEN, UBA, Intendente Güiraldes 2160, Ciudad Universitaria, C1428EGA, Buenos Aires, Argentina.

[3] Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires. Intendente Güiraldes 2160, Ciudad Universitaria, C1428EGA, Buenos Aires, Argentina.

*Corresponding Authors: Francisco Velazquez Duarte at fvelazquez.ib3_@gmail.com and Javier Santos at javiersantosw@gmail.com

Running Title: Modeling Friedreich Ataxia in *Dictyostelium*

Keywords: Iron-Sulfur Cluster Assembly/ Frataxin/ Friedreich Ataxia/Rare Disease

Abbreviations: ACP, acyl carrier protein; Cas9, CRISPR associated protein 9; CD, circular dichroism; CTR, C-terminal region; CRISPR, clustered regularly interspaced short palindromic repeats; DLS, dynamic light scattering; Fe-S, iron-sulfur; FA, Friedreich's Ataxia; FXN, frataxin; DdFXN, *D. discoideum* frataxin; DTT, dithiothreitol; HPLC, high-performance liquid chromatography; HsFXN, human frataxin; ISCU, iron-sulfur cluster assembly enzyme; ISD11, NFS1 interacting protein; NFS1, mitochondrial L-cysteine desulfurase enzyme; NMR, nuclear magnetic resonance; PAGE, polyacrylamide gel electrophoresis; PDB, Protein Data Bank; SDS, sodium dodecyl sulfate; SEC, size exclusion chromatography.

Here we present the development of a new model system for Friedreich's Ataxia (FA)

using *D. discoideum* (Dd). FA is a rare disease caused by disfunction of frataxin

(FXN), a protein involved in Fe-S cluster assembly machinery. We firstly investigated

the conservation of function between human and *D. discoideum*. In this work we show

that DdFXN can substitute the human version in the interaction and activation of Fe-S

assembly supercomplex. DdFXN can in vitro displace HsFXN in competition assays

and also it can activate cysteine desulfurase activity in the context of human Fe-S

assembly supercomplex. We then manage to edit *fxn* locus and isolated clone 8, a

defective mutant with undetectable levels of frataxin. Clone 8 presents landmarks of

frataxin deficiency such as decrease in Fe-S cluster dependent enzymatic functions,

growth rate reduction and increase sensitivity to oxidative stress. Besides these

phenotypes, shared with other FA models, clone 8 presents defects in the multicellular

developmental program induced by starvation in this protist. We then assessed the

rescuing capacity of DdFXN G122V, a version that mimics a human variant presented

in some FA patients. While expression of DdFXN G122V rescues growth and

enzymatic activities defects as well as DdFXN does, multicellular development defects

were only partially rescued. This work opens the door to develop drug or treatment

screenings that would help to design, and/or evaluate therapeutical strategies. Besides

this biological model offers a wide range of possibilities to easily explore diverse

phenotypes in FA.

Introduction

Dictyostelium discoideum is a social amoeba that lives in the soil and feeds on bacteria

and other microbes. Dictyostelids belong to a separate branch of the eukaryotic

organisms, distinct from plants, fungi and animals. Its cells lack a cell wall resembling

animal cells in organization (*1*). *D. discoideum* has become a very attractive eukaryotic

non-mammalian model organism to study molecular mechanisms, cell physiology and

human pathology (*2*, *3*). The studies carried out with *D. discoideum* have provided great

insights in diverse areas such as bacterial infection, immune cell chemotaxis,

autophagy/phagocytosis as well as mitochondrial and neurological disorders (*4*).

This amoeba has been widely used to study human diseases including

neurodegenerative such as Alzheimer's, Parkinson's, Huntington's, obtaining important

discoveries of the pathophysiology of these pathologies (*5*). Furthermore, *D. discoideum*

has been used as a model to identify drug targets and discover new compounds with

therapeutic potential, and even these advances served as a platform for new clinical

trials (*6*, *7*). This organism allows to study the effect of very specific molecular

alterations, from different viewpoints, combining biochemistry and structural and cell

biology to generate a comprehensive picture concerning the outcomes of the alteration

at different levels of complexity. *D. discoideum* has a unique developmental life cycle

among eukaryotes, presenting both unicellular and multicellular phases. This allows the

study of different levels of cell organization and cell-cell interaction and communication

(**Figure 1**).

Figure 1. *D. discoideum* Life Cycle. *D. discoideum* grows as unicellular amoebae as
long as there are nutrients available (Vegetative Cycle). During this growth phase,
haploid amoeboid cells feed on bacteria and divide by a binary fission process. In
response to starvation *D. discoideum* triggers a multicellular developmental program
that leads to the formation of a fruiting body bearing a mass of spores for dispersal
(Social Cycle). During this program, starved cells gather by chemotaxis and form an
aggregate. This aggregate suffers several morphogenetic changes passing through
different stages. First, cells form the mound, enclosed within extracellular matrix. After
that, the mound develops into a standing structure (finger). The finger may develop into
the migrating slug, or it may directly progress through the culmination stages. Finally,
the fruiting body is formed. It will contain spores that will be released and eventually
germinate to produce growing amoeboid cells (*8*). Alternatively, to the multicellular
program *D. discoideum* can go through a sexual cycle if sexually compatible cells are
present and other environmental conditions are met (*9*).
97

Recently, CRISPR/Cas9-mediated technology has been implemented to *D.*

discoideum, allowing the rapid genome editing by transiently expressing sgRNA and

Cas9 using an all-in-one vector, and the generation of genomic mutants (*10–12*). This

possibility may accelerate the gaining of knowledge of specific components of the

molecular machinery underlying the complex life cycle in *D. discoideum*.

In addition to all stated above, *D. discoideum* helps to implement the "Three Rs

principle" in animal research (Replacement, Reduction, and Refinement); so

development of *Dictyostelium* based disease models will be highly beneficial in

preliminary drug screening (*13*). D. *discoideum* has been identified as an exceptional

model organism to study rare diseases (*14*, *15*). Rare diseases are more than 8000

different pathologies, affecting more than the 8% of the world human population; they

are generally underrepresented in scientific agenda and suffer from a lack of tools,

model organisms and therapeutics.

In this context, our laboratory is interested in the development of new tools to

facilitate the finding of therapeutic solutions for Friedreich's Ataxia. We focused on

Friedreich's Ataxia (FA), a neurodegenerative disease that involves 1:50000 people.

The main cause of this disease is the decrease in the expression or in the functionality of

a mitochondrial protein named frataxin, which is encoded in the nuclear genome.

Frataxin is involved in the biosynthesis of iron-sulfur cluster (Fe-S), in the

mitochondrial matrix. The deficiency of frataxin affects several enzymatic reactions that

depend on Fe-S clusters, which are essential cofactors involved in several enzymatic

functions. The Krebs cycle enzymes, aconitase and succinate dehydrogenase (*16*), the

respiratory chain complexes (*17*, *18*), and critical processes such as DNA repair (*19*),

chemical modification of transfer RNAs (*20*), and lipoic acid synthase, which catalyzes

the final step in the novo pathway for the biosynthesis of lipoic acid (key coenzyme of

pyruvate dehydrogenase and α-ketoglutarate dehydrogenase enzymes) (*21*), all depend

on Fe-S clusters.

Fe-S assembly reaction depends on a supercomplex formed by at least five

different subunits: i) the L-Cys desulfurase NFS1, a pyridoxal-phosphate (PLP)

dependent enzyme, which catalyzes the desulfurization of L-cysteine generating as

products the precursor sulfide attached as a persulfide group to a Cys residue (Cys-S-

SH) and L-alanine, ii) ISD11 protein, which is only present in eukaryotic organisms

(*16*, *22*), iii) the mitochondrial acyl carrier protein (ACP), which stabilizes ISD11(*23*,

24), iv) the scaffolding protein (ISCU) and v) frataxin.

NFS1 dimer is stabilized by the ACP-ISD11 heterodimer (*25*). The Fe-S cluster

assembly site is situated on the scaffolding protein ISCU, and frataxin is the kinetic

activator of the reaction but also its surface is part of the Fe-S cluster assembly. The

stoichiometry of the supercomplex is $(NFS1\text{-}ACP\text{-}ISD11\text{-}FXN\text{-}ISCU)_2$. Its architecture

is intricate, each heterodimer ACP-ISD11 also interacts with both NFS1 subunits as a

bridge and two assembly sites formed by ISCU/FXN and both NFS1 chains. That is,

frataxin simultaneously interacts with both NFS1 subunits and ISCU (*26*). Even though

it was demonstrated that frataxin binds iron, it is not clear whether it works as a

chaperon of this metal ions in the context of the supercomplex. It was also shown that

ISCU is able to bind this metal and this activity is crucial for Fe-S cluster assembly

(*27*).

Besides Friedreich's ataxia caused by mutations in frataxin gene, the alteration

of transcription, aberrant splicing or the presence of point mutations in other proteins

involved in cluster assembly affecting their expression or functionality, result in severe

human diseases. Thus, mutation of NFS1 results in an autosomal recessive

mitochondrial disease characterized by the respiratory chain complex II and III

deficiency and multisystem organ failure (*28*), mutation of ISCU results in ISCU

myopathy (*29*), whereas the mutation of ISD11 (R68L) is associated with the

development of a mitochondrial genetic disorder, an autosomal recessive disease,

known as Combined Oxidative Phosphorylation Deficiency 19 (COXPD19) (*17*).

The mutations of frataxin that result in FA affect the protein at different levels:

conformational stability (e.g., L198R, G137V, G130V) (*30–32*), mitochondrial import

and processing pathway (e.g., W168R and W173G) (*33*), iron binding affinity (e.g.,

D122Y) (*32*, *34*) or alterations at the assembly site architecture (e.g., W155R and

N146K and Q148R) (*35*, *36*). In addition, the truncation of frataxin at position 193

(deletion of the last stretch of residues named C-terminal region that conform a non-

periodic structure) resulted in a pathogenic variant (*37*). A very similar variant,

truncated at position 195, exhibited strong alterations in its internal motions and also a

reduced conformational stability, and a decrease in its iron binding capability (*30*).

Frataxin has been described as an essential protein in eukaryotic organisms since

the deletion of this protein is lethal in yeast and mammalian cells (*38*). Moreover, in

multicellular eukaryotic organisms (plants and mice), the complete deletion of frataxin

leads to early embryonic lethality (*39*, *40*), or to the arrest of the larval stage L2 / L3 in

Caenorhabditis elegans (*41*) and reduced larval viability along with a metamorphosis

failure in *Drosophila melanogaster* (*42*, *43*). Regarding the use of cell cultures, different

models have been described. Since the patient's cell line (fibroblasts or lymphocytes)

does not consistently express the biochemical phenotypes associated with FA under

basal culture conditions, RNA interference (RNAi) strategies have been developed to

reproduce partial frataxin deficiency in human and murine cell lines (*44*, *45*).

Furthermore, "humanized" murine cell models have been developed to eliminate

endogenous frataxin and expressing frataxin with pathogenic mutations (*46*). A murine

fibroblast cellular model, with the ability to deactivate frataxin transcription, was also

generated using the Cre /loxP recombination system. On the other hand, stem cells from

patients have been used and the use of inducible pluripotent stem cells that can mimic

tissues affected by FA is under development (*47*, *48*).

In a previous paper we examined the *D. discoideum* genome and found the

complete dotation of proteins involved in Fe-S cluster assembly was found (*49*).

Furthermore, by analyzing the sequences and structure models of these proteins, we

found that residues located in the protein-protein interaction surfaces are highly

conserved between the amoeba and human. In fact, the frataxin residues involved in FA

are fully conserved, with the exceptions of a core residue His183 of the human frataxin

that interacts with residues located in the CTR (the last stretch of the protein), whereas

in DdFXN, this residue is an Arg fully exposed to the solvent and a Tyr residue (in

DdFXN) is located at the position corresponding to the Trp168 in the human variant, as

inferred by means of a structure model of DdFXN.

In this paper, we investigated the effect of the functional deficiency in frataxin

on the amoeba *D. discoideum* metabolism and physiology. The short duplication time,

the simple genetic manipulation, the simplicity of creating knockout cell lines by

CRISPR/Cas9 due to its haploid genome and the simplicity of the characterization of

phenotypes were conceived as key advantages of this model.

The alterations observed in mammalian cells are also present in *D. discoideum*,

as reduced aconitase and succinate dehydrogenase or higher sensitivity to oxidative

stress. In addition, the amoeba cultures deficient in frataxin grew at lower rate and the

life cycle exhibited alterations. We have explored the rescue capacity of constitutive

frataxin expression.

These cell lines obtained by CRISPR/Cas9 will allow us to carry out therapeutic

screening of different compounds and drugs that can be used in the treatment of FA.

Material and Methods

Strains, Cell Culture, Plasmid Constructions, CRISPR/Cas9 Guide Design and D.

discoideum Transformation

D. discoideum strain AX2 was cultured at 22 °C on axenic culture on dishes, in

SM agar plates with *Klebsiella aerogenes*, or in Erlenmeyer flasks, in HL5 medium and

shaking (180 rpm). The parental plasmid used for CRISPR/Cas9 genome editing

(pTM1285) (*50*) was kindly provided by Dr. Kamimura. It was used to insert the DNA

fragment codifying for the RNA guide to target the *D. discoideum fxn* gene. For the

guide preparation, a pair of oligonucleotides designed using CRISPOR

(http://crispor.tefor.net) and Breaking-Cas

(https://bioinfogp.cnb.csic.es/tools/breakingcas/index.php) were adequately annealed,

and for the DNA ligation, the Golden Gate strategy was used. Construct checking was

performed by colony PCR of *E. coli* DH5α, using the RevGuide553 and tRNA_seq_3 as

primers.

For transformation of *D. discoideum* AX2 cells were harvested during the

exponential phase of growth, washed twice in ice-cold H50 buffer (20 mM HEPES, pH

7.0, 50 mM KCl, 10.0 mM NaCl, 1.0 mM $MgSO_4$, 5.0 mM $NaHCO_3$ and 1.0 mM

NaH_2PO_4), and re-suspended in H50 buffer at a concentration of 5×10^7 cells/ml. A

volume of 100 µl of cell suspension was electroporated using 10-20 µg of plasmid (0.75

kV, 25 µF, twice). Cells were transferred to culture dishes containing HL5 culture

medium an incubated for16 h. For clone selection G418 antibiotic (geneticin) that

blocks polypeptide synthesis was added (20 µl/mL) and the cells were incubated for

48h; the resistance to G418 is conferred by the neo gene located in pTM1285 vector.

227   After that, HL5 medium containing G418 was removed from the plastic dishes

228 and the cells were re-suspended in a volume of 400 µL of HL5 medium; a volume of 75

229 µL was mixed with 250 µL of bacteria *Klebsiella aerogenes* liquid culture and plated in

230 SM agar plates. After an incubation of 3-4 days at 22 °C, plaques were observed

231 (absence of bacterial lawn).

D. discoideum Clone Isolation and DNA Sequencing

To identify sgRNA/Cas9 editing, the isolated genomic DNA was directly carried

out by picking up cell material from corresponding *D. discoideum* plaque using a

sterilized tip. The material was re-suspended in a volume of 50 µL of Lysis Buffer (50

mM KCl, 10 mM Tris-HCl pH 8.3, 2.5 mM $MgCl_2$, 0.45% NP40, 0.45% Tween-20 and

Proteinase K (40 µg). The cell suspension was incubated at 20-24 °C for 20 min and

then heated at 95°C for 3 min to inactivate the protease. The cell lysate was used as

template for PCR. The DNA corresponding to *fxn* gene was amplified using KOD hot

start master mix (Cat. Nº 71842 Millipore) and FwDdFXNOE and RevDdFXNOE as

primers. The PCR cycle comprises denaturation at 95 °C for 30 s, annealing at 55 °C for

30 s an extension at 68 °C for 30 s, and 40 PCR cycles were programed. The amplified

DNA fragments were purified using Wizard SV Gel and PCR Clean-Up System (Cat.

Nº A9281 Promega), then the DNA was sequenced using the Macrogen facility

(https://dna.macrogen.com/).

Evaluations of D. discoideum Growth Rate and Life Cycle Alterations

To evaluate the generation time (*g*), *D. discoideum* growth curves (22 °C and

180 rpm) were made using flasks (250 mL) containing HL5 culture medium (75 mL)

and starting from a $1x10^5$ amoeba cells/mL inoculum. The cell cultures were grown

during a week, and small aliquots (0.1 mL) were regularly taken for cell counting using

a Neubauer chamber. The growth curves were analyzed, and the maximum (μ) was

obtained from the slope when the culture grows exponentially. The generation time was

calculated according to Equation 1.

$$g = \frac{\ln 2}{\mu} \,(1)$$

For the study of the *D. discoideum* life cycle and development, 6-cm petri dish

were prepared with 10 mL of 1.8% Oxoid L28 agar in KK2 medium (16.5 mM

KH_2PO_4, 3.9 mM K_2HPO_4, pH: 6.2) and 2.0 mM $MgSO_4$, 0.1 mM $CaCl_2$ (complete

KK2).

A volume of 5-15 mL of cell culture in log phase (2-5 $\times$ 10^6 cells/mL) was

centrifuged at 1400 rpm for 3 min under conditions, next two washes with KK2 were

carried out. Then, the cells were resuspended in complete KK2 to give 2.5 $\times$ 10^7

cell/mL. After that, a volume of 1.6 mL containing 4 $\times$ 10^7 cells were added to each

plate, and the plates were left in a levelled table for 15 min for cell adhesion, after that

medium was carefully removed by tilting the plates. The plates are incubated in

humidity at 22 °C for 18-22 h.

Aconitase and Succinate Dehydrogenase Activities Measurements

Aconitase (ACO) and Succinate dehydrogenase (SDH) enzymatic activity

measurements were carried out on the soluble fraction of cellular extracts of *D.*

discoideum AX2, **clones 8** and **J**.

For the SDH activity assay, 2 $\times$ 10^5 cells were used per reaction (in multi well

plates) and cell lysis was performed by three freeze-thaw cycles. The assay was carried

out according to the supplier's instructions (Abcam: ab 228560).

To measure ACO activity, 1.5×10^6 cells were used per reaction. In this case,

cell lysis was performed by means of a detergent solution included in the commercial

kit. The protocol was according to the supplier's instructions (Abcam: ab 109712).

Frataxin Variants Molecular Dynamics Simulation

Molecular Dynamics simulations and the analysis of production runs were

carried out using the YASARA Structure (*51*) on the following hardware: Processor

Intel CORE i7 10,700 10th generation; SOCKET 1200 2.9 GHZ (Max 4.8 GHZ) 16 M

cores/threads 8/16, 2; Memory Kingston HX426C16FB3/8G HyperX2666 MHz; Disc

SSD Kingston A400 240GB SATA 7 mm; Linux Ubuntu 20.04 LTS 64 bit. The

molecular models corresponding to wild-type DdFXN and DdFXN fragment from clone

8 strain were modeled using AlphaFold2 (*52*) and with SwissModel at

https://swissmodel.expasy.org/interactive.

The coordinates were solvated, and standard minimization protocols were

applied to remove steric clashes. The simulation cell was prepared by maintaining a 20

Å water-filled space around the protein with a density of 0.997 g/mL. The system (cubic

cell, periodic boundaries, and 8.0 Å cut-off for long-range coulomb electrostatics

forces) was neutralized with 0.9% NaCl, and the temperature was maintained at 298 K

with a pH of 7.4. After the initial steepest descent minimization, unrestrained replicas of

100 ns MD simulations using an ff14SB Amber force field were carried out with 2.50 fs

time steps (*53*). Snapshots were saved every 0.1 ns. The root-mean square deviation

(RMSD), root mean square fluctuation (RMSF), and secondary structure content were

calculated. Molecular dynamics simulations were performed using Yasara Structure

program (*54*).

Western Blotting Analysis

For frataxin detection in cell lysates of *D. discoideum*, we used a set of

nanobodies prepared in our laboratory using recombinant DdFXN as the target and

phage display technology. Three panning rounds with increasing washing steps (10, 15

and 25 for rounds 1, 2 and 3, respectively) were carried out.

For de detection of DdFXN-G122V detection an anti-FLAG monoclonal

antibody (Cell Signaling Technology, 8146S (9A3) Mouse mAb that binds to same

epitope as Sigma's Anti-FLAG M2 Antibody) was used to specifically identified the

expression of frataxin from plasmidic DNA. For this detection, the FLAG of sequence

DYKDDDDK was included in the C-terminal stretch of DdFXN.

Table 1. Oligonucleotides Used in this Work

Name	Oligonucleotide Sequence [4]
FwDdFXNOE [1]	AAagatctATGATTTTCAACTTTTTAAACAAAGC
RevDdFXNOE [1]	GTactagtAATTTCCATATCATATTTACAAAGTG
FwGuide 553 [2]	agcaTGGTGGTCATCACCATTGAG
RevGuide 553 [2]	aaacCTCAATGGTGATGACCACCA
tRNA_seq_3 [3]	GCTCGATTAGCTCAGTCGGCAG

311 [1] These primers anneal within *D. discoideum* frataxin coding sequence.
312 [2] These primers were used to clone in pTM1285 and define the guide RNA employed
313 during CRISPR/Cas9 edition that targets essential region of DdFXN. 3
314 [3] This primer anneals in pTM1285 and allows to check proper guideRNA cloning.
315 [4] Upper-case letter indicates match and lower-case mismatch with target template.
316

Oxidative Stress Sensitivity Assay

To assess sensitivity to oxidative stress cell viability was determined upon H2O2

treatment using crystal violet assay described by Feoktistova et al. (*55*). In summary

100000 cells were seeded in a 24-well plate and let to adhere for 1h. Then medium was

removed and fresh HL5 medium or supplemented with 2 mM H_2O_2 was added to each

well. Since adherent cells detach from cell culture plates during cell death, after

corresponding time of treatment, wells were gently washed to removed dying cells.

Remaining cells are fixed and stained with crystal violet. Culture plates were dried, and

crystal violet was measured after solubilization with acetic acid (A570 nm) as an

estimate of remaining cells. Viability was calculated referenced to the crystal violet at

time 0 for each strain. The experiment was performed at least 3 times for each strain and

contains 4 replicates of each time point.

Results
331

DdFXN Can Substitute Human Orthologue in Supercomplex for Fe-S cluster
Assembly

Since we aimed to establish *D. discoideum* as model organism for FA using

DdFXN to explore its roles within the cell and extrapolate to human context, we firstly

test the ability of DdFXN to substitute the human orthologue in the context of the

supercomplex.

As a prove of functional conservation, we assess the ability of DdFXN to

interact with human supercomplex NIAU (NFS1/ACP-ISD11/ISCU)$_2$. We carried out

fluorescence anisotropy assays using a variant of DdFXN (DdFXN_C35A) or the

human frataxin (HsFXN) labelled with Texas-red.

The labelled frataxin variant, in the presence of saturating ISCU concentration,

was incubated with increasing NIA concentrations. As it can be seen in **Figure 2A**,

DdFXN variant was able to bind to the human NIAU in a similar fashion as the mature

form of the human frataxin do, although with lower apparent affinity. It is worthy of

note that wild-type DdFXN was able to compete with the labeled-human-frataxin

variant (**Figure 2A, inset**), suggesting that the same binding site is involve for both

proteins.

We then tested if this binding is functional and DdFXN is also able to act as the

kinetic activator of L-Cys-desulfurase activity in human supercomplex context. **Figure**

2B shows that the interaction of DdFXN to human supercomplex was productive and

Cys-desulfurase activity increased upon addition of human or DdFXN.

Figure 2. *In vitro substitution of human frataxin by DdFXN.* (A) DdFXN interaction with the Human supercomplex L-Cys desulfurase. Texas red-Labeled DdFXN (blue and cyan correspond to two independent experiments) or human FXN (black and gray correspond to two independent experiments) were incubated with increasing concentrations of NIAU subcomplex and the fluorescence anisotropy was monitored. Fraction of bound frataxin was plotted as function of free NIAU complex. Inset shows the competition with non-labeled proteins of a preformed supercomplex in which human frataxin (hFXN) was labeled with Texas red followed by fluorescence anisotropy. The competition was carried out using non-labeled hFXN (black triangles) or non-labeled DdFXN (blue triangles); the fraction of bound frataxin was plotted as a function of the added FXN variant. (B) In vitro activity of the human supercomplex activated by human frataxin (hFXN) or DdFXN. The activity corresponding to NIA or NIAU complexes without frataxin are shown for the reference.

Fxn Locus Can Be Easily Edited in D. discoideum with CRISPR/Cas9 Technology

As noted above, we aim to generate a new experimental model to study the biochemical consequences of alterations in the functionality of frataxin in the cell. We intend to get mutant strains that were highly deficient in frataxin so we could explore strategies for reestablishing homeostasis and to evaluate the rescue capacity of wild type and disease-associate variants. With this aim in mind, we have carried out CRISPR/Cas9 editing of *D. discoideum* endogenous locus, using a guide RNA targeted at Trp146. This residue, corresponding to the human Trp155. It is extremely conserved in frataxin along the tree of life because of a functional role (*49*). When looking at the

structure of human supercomplex (NFS1/ACP-ISD11/ISCU/FXN)$_2$, Trp155 is located

near docking surface of ISCU and at Van der Waals distance of its [2Fe-2S] assembly

site, as well as from Leu386 of NFS1 what gives away its importance. Trp155 belongs

to a highly conserved region that contains the Motif 1 described by Gibson *et al.* (*56*).

Hence it would be more likely to obtain deleterious mutations in this region.

Using an all-in-one vector, pTM1285 to express the corresponding guide RNA we

obtained hundreds of clones and selected 14 presumably edited clones (**Figure S1**).

After the isolation of clones, the genomic DNA corresponding to the *fxn* gene

was successfully amplified and sequenced for 13 clones. The analysis of the genomic

DNA sequences showed a variety of mutations in the protein as the result of

CRISPR/Cas9 editing and the subsequent DNA repair (**Table 2**, **Figure S1**).

Table 2. Summary of CRISPR/Cas9 Edition of *fxn* Locus in *D. discoideum* in

Sequenced Clones.

Mutation	Number of Clones
Premature STOP codon	9 (7)[1]
Small Insertion	2
Point Mutation	1
Silence Mutation	1

[1] Seven different editions.

A D. discoideum Clone Lacking Frataxin is Viable

The amino acid sequence analysis of *D. discoideum* clones obtained by CRISPR/Cas9

genome editing (*e.g.*, **clone 8**) indicated that *fxn* gene functionality of DdFXN can be

obliterated without being lethal for the amoeba, at least in the experimental conditions

assayed in this work.

Edition in **clone 8** produces a large truncation of the protein chain because of

one base pair deletion generating a frame shift, and premature stop codon (**Figure S2**).

This truncation eliminates a stretch that includes more than 40% of the sequence, which

may completely destabilize the structure of the remaining protein fragment, comprising

residues 81-150 from DdFXN, using the numbering of the DdFXN precursor (**Figures**

3A and B). The truncation eliminated part of strand beta 4, strands beta 5 and 6, alpha

helix 2 and the CTR.

We studied the molecular motions of this fragment by molecular dynamics

simulations. Even though the global conformation of the fragment persisted during 100-

200ns MDs simulations, significantly higher RMSD values and atomic fluctuations

were observed, compared to the wild-type (**Figures 4A and B**). Alpha helix 1 presented

high distortions, stablishing non-native contacts with the beta sheet (**Figures 4E and F**)

and higher internal motions compared to the wild type frataxin, as judged by the

fluctuations.

All this suggests that the fragment should be highly dynamic and more likely

easily degraded in the cellular environment. However, we cannot discard that the

fragment might acquire frataxin-like secondary structure and even some kind of

packing, although residual activity is highly unlikely.

AlphaFold2 predictions suggested a conformation for a putative dimeric fold of

the fragment, in which the predicted accessible apolar surface for each monomer is

highly reduced by the interaction, and the remanent Cys residue (Cys112) might

establish an intermolecular disulfide bond stabilizing the hypothetical dimeric structure

(**Figure S3**). Experiments will be done to evaluate this possibility.

Figure 3. The alterations of Frataxin Structure. (A) and (B) Two views of truncated frataxin DdFXN_81-150 (clone 8). The topology shown is only for visualization purpose. (C) and (D) Two views of the wild-type DdFXN model. DdFXN models were constructed using AlphaFold2 (*52*). AlphaFold formed a disulfide bond between Cys112 and Cys187, whereas other programs like Swiss-model did not establish the -S-S- bond. (E) Amino acid sequence truncation predicted from the DNA sequences for **clone 8** (**Figure S2** shows the complete sequence alignment).

434
435
436
Figure 4. Molecular Dynamics Simulations of DdFXN Variant. (A) Root-mean-square deviation (RMSD) along the simulations (calculated for the alpha carbon atoms). (B) Root-mean-square fluctuations (RMSF) of the alpha carbon atoms. Snapshots of wild-type variant (C and D) and the fragment (E and F). Wild-type and clone 8 (fragment) FXN were constructed using AlphaFold2 (*52*). A disulfide bond between Cys112 and Cys187 is formed. For the simulations, three non-native-extra residues in the N-terminal stretch were included: **Met78-Gly79-Ser80-**Pro81-Ile82-Ser83, as for the recombinant proteins produced in our laboratory.

Edited Clone 8 Present Undetectable Frataxin Expression Levels

448   We firstly analyzed the expression levels of frataxin in wild-type AX2 and

449   edited clone 8 strains. Two different nanobodies that detected the recombinant DdFXN

450   in western blotting were used to study the frataxin expression in total *D. discoideum* cell

451   lysates (**Figure 5**).

Figure 5. *Analysis of Frataxin Expression by Western Blotting.* Total cell lysates corresponding to AX2 and clone 8 were analyzed by western blotting using two different nanobodies that carry a 6xHis-tag (NB41 and NB45) raised against DdFXN and a HRP conjugated secondary antibody antiHis tag (lower panel) and anti ß-actin as loading control (upper panel).

As it can be seen in **Figure 5**, cell lysate corresponding to the wild-type AX2

undoubtedly showed detectable levels of frataxin. On the other hand, clone 8 a band

corresponding to frataxin or to a frataxin fragment was not detectable. The absence of

detectable signal in clone 8 was not due to the inability of nanobodies to recognize the

mutated versions since when truncated version was heterologous expressed in bacteria

and purified it was readily detected by both nanobodies (**Figure S4**). This result

indicated the virtual absence of frataxin, in the CRISPR edited cells (**Figure 5**).

A D. discoideum Clone Lacking Frataxin Has Severe Defective Growth

As an initial global analysis on the effects of altering frataxin functionality in *D.*

discoideum biology, we evaluated the growth of the frataxin-deficient clone in the HL5

rich medium and on bacteria. As it is shown in **Figure 6 and Table 3**, clone 8 presents

a growth defect in both conditions. Wild-type strain AX2 exhibited a generation time of

9.5 ± 2.2 h, this is in accordance with the literature ([57]). On the other hand, the edited

clone 8 carrying the truncated frataxin, grew at lower rate, $(14.9 \pm 2.2$ h$)$.

Figure 6. *D. discoideum* Grow. Ax2 and clone 8 were grown on bacterial lawn (left) of
in HL5 culture medium (right), starting from a $\sim 1 \times 10^5$ amoeba cells mL^{-1} inoculum.
The cell cultures were grown during a week, and small aliquots were regularly taken for
cell counting using a Neubauer chamber. Three independent experiments were carried
out; a representative experiment is shown.

Table 3. Generation Time Summary.

Strain	Ax2	Clone 8	Ax2 (empty)	Ax2 (G122V)	Ax2 (FXN wt)	Clone 8 (empty)	Clone 8 (G122V)	Clone 8 (FXN wt)
Generation time (h)	9.5 ± 2.2	14.9 ± 2.2	12.6 ± 2.6	13.2 ± 2.8	12.7 ± 2.3	19.3 ± 2.4	12.8 ± 2.7	13.2 ± 2.8

A D. discoideum Clone Lacking Frataxin Presents a Strong Decrease in Fe-S cluster

Dependent Enzymatic Activities

Given that the typical phenotype of a reduction in frataxin functionality is the

decrease in Fe-S cluster dependent enzymatic activities, and that this is a feature shared

among the FA models, we studied the Krebs cycle enzymes aconitase (ACO) and

succinate dehydrogenase (SDH, the Complex II from the respiratory chain). We

compare activity of these mitochondrial enzymes in cellular extracts of strains AX2 and

clone 8 (**Figure 7**).

Clones 8 exhibited significant reduction of the enzymatic activities by

comparison to the wild-type AX2. The activity decreased to the 30-40% of the level

observed in wild type; these values are comparable to those obtained in other cell

models deficient in frataxin, where ~50% reduction of ACO and SDH activities were

detected (*46, 58*).

498
499
500
501 **Figure 7. Effect of Frataxin Alteration on Fe-S Cluster Dependent Enzymatic**
502 **Activities.** Aconitase (ACO) and succinate dehydrogenase (SDH) activities were
503 assayed on total lysate extract of *D. discoideum* cells from AX2 strain and clone 8.
504 Three independent experiments were carried out. The activity measurements are relative
505 to the total protein quantified by triplicate in each extract by the Bradford method.
506
507

A D. discoideum Clone Lacking Frataxin Presents Higher Sensitivity to Oxidative

Stress

Another hallmark of FA disease is an increased sensitivity to oxidative stress. The

evidence provided by several organism models supports the idea that the deficiency in

frataxin function causes a deregulation in the antioxidant response (*42, 59–64*). In a

similar fashion, one can think that clones 8, which exhibits a significant lower growth

rate, may show an inefficient handling of reactive oxygen species (ROS), with an

ineffective elimination of ROS by the antioxidant system. To study this issue, we

carried out a multi-well plate assay and the effect on cellular viability of a treatment

with H_2O_2 was determined. We observed a higher impact of hydroperoxide on clone 8

cells compared to the wild-type AX2 cells (**Figure 8**).

523
524
525 **Figure 8. Sensitivity to Oxidative Stress.** For determining viability of cultured cells, a
526 Crystal Violet assay was carried out. The differential effects of H_2O_2 on clone 8 cells
527 compared to the wild-type AX2 cells was evaluated. A treatment of frataxin-deficient
528 amoeba with hydroperoxide resulted in a significantly increase of the cell detachment
529 from the multi-well plate surface.
530

A D. discoideum Clone Lacking Frataxin Presents Defects in Multicellular

Development Progress.

To continue with the exploration of the frataxin deficiency effects on *D.*

discoideum biology, we monitor the progression of the multicellular development

program in the frataxin-deficient clone 8 (**Figure 9**). As we mentioned previously, *D.*

discoideum grows as isolated ameboid cells that divide mitotically in a nutrient rich

culture medium, or in the presence of bacterial lawn. On the other hand, starvation

induces a developmental program the result of which is the formation of a fruiting body

bearing spores that would disseminate and germinate when nutrients become available.

This program involves chemotaxis, cell differentiation, morphogenetic movements, and

integration of internal and external signals. In amoeba with mitochondrial deficiency, as

a compensatory mechanism for energy deficit, there is a chronic increase in AMPK

activity (AMP-activated protein kinase) that maintains mitochondrial mass and ATP

levels at normal levels; however, it affects other process that require energy such as:

growth, multicellular development, phototaxis and chemotaxis (*65*). Furthermore,

previous reports of *D. discoideum* with mitochondrial disease show that mitochondria

defective cells present poor growth both in fluid and in bacterial grasses and also

alterations in multicellular development (*66*).

As it can be seen in **Figure 9**, clone 8 has an altered development; after 18 h of

starvation, while wild type AX2 already showed as mature fruiting bodies, clone 8

presented immature stages (tipped mound). This significant delay seems to start from

the beginning of development as it is already evident after 12 h of starvation; at this

point wild type strain has reached mound stage while clone is only at early aggregate.

556
557
558
559 **Figure 9**. **Starvation-triggered Multicellular Development Progression.**
560 Exponentially growing cells of Ax2 (top) and clone 8 (bottom) were seeded in non-
561 nutrient agar plates to induce multicellular development and incubated at 22 °C.
562 Pictures were taken at different times (12, 18, 24 and 40 hours) a representative picture
563 of each timepoint is shown.
564
565
Constitutive Expression of Frataxin Wild type Rescues all Phenotypes in Clone 8

To undoubtedly establish that all defects reported in clone 8 were due to the lack

of functional frataxin we express an intron-less, C- terminal flag-tagged wild type

frataxin (DdFXNwt) version from a constitutive promoter.

We then evaluate the rescue capacity of the phenotypes reported above. As it can

be seen, both growth (**Table 3**) and development (**Figure 10**) of clone 8 expressing

DdFXNwt was not significantly different to wild type strain AX2 transformed with

empty vector or over-expressing DdFXNwt. On the other hand, clone 8 transformed

with empty vector recapitulates the reported alterations: increased generation time

(**Table 3**), decreased aconitase and succinate dehydrogenase enzymatic activities

(**Figure 11B**) and delayed development (**Figure 10**).

Interestingly the rescue of the decrease of the iron-sulfur cluster dependent

enzymatic activities was not complete (**Figure 11B**). The expression of DdFXNwt from

the plasmid was corroborated by western blotting analysis, using an anti-FLAG

monoclonal antibody (**Figure 11C**). As expected, no expression was detected when the

transformation was carried out with empty vector.

Figure 10. Starvation-triggered multicellular development progression. Exponentially growing cells of each strain were seeded in non-nutrient agar plates to induce multicellular development and incubated at 22 °C. Representative pictures after 14 hours (left column), 16 hours (middle column) and 20 hours (right column) of development are shown. Yellow arrows and red arrows mark immature mound/tipped mound and early culminants/fruiting bodies.

589 *Constitutive Expression of Frataxin G122V in Clone 8 Fully Rescues Growth Defect*
590 *but Produces only Partial Recovery of Developmental Impairment.*
591
Finally, we wondered whether a high expression of a FA frataxin variant can

rescue the wild-type features of *D. discoideum* cells. We transformed clone 8 cells with

plasmid pDM326-DdFXN-G122V, which encodes the precursor form of G122V

frataxin variant. This variant carries a mutation that produces a very specific and less

aggressive FA phenotype in humans (G130V). Noteworthy, clone 8 cells transformed

with pDM326-DdFXN-G122V recovered the growth rate of wild-type AX2 (**Table 3**,

Figure 11A).

We then evaluated whether the G122V frataxin might be able to rescue the

defects in multicellular development observed for clone 8. Although a small

improvement was detected when these cells were transformed with pDM326-DdFXN-

G122V, compared with the same cells transformed with empty pDM326, development

was not fully rescue.

At 20 h after starvation, we could still see immature structures in clone 8 overexpressing

G122V variant while none when complemented with wild type frataxin (**Figure 10**).

We also evaluated the rescue capacity of DdFXN-G122V on the enzymatic

activities decrease (**Figure 11B**); although rescue was not total the enzymatic levels

reached were like the obtained when DdFXNwt was expressed.

These results suggested that, because of the specific features of this variant, the

expression of DdFXN-G122V cannot fully compensate all deficiencies of lacking wild-

type frataxin. This fact also suggested different activities, which are finely tuned by

frataxin structure and dynamics, or different control over the metabolic pathways.

Figure 11. The Expression Frataxin Variant G122V Rescued Clone 8. (A) Growth rate and (B) aconitase and succinate dehydrogenase enzymatic activities of *D. discoideum* AX2 and **clone 8** cells transformed with pDM326 or pDM326 + DdFXN G122V encoding the precursor variant of DdFXN G122V (G122 position corresponds to G130 in human frataxin). Growing was carried out at 22 °C and 180 rpm in HL5 culture medium, starting from a 1×10^5 amoeba cells mL^{-1} inoculum. The cell cultures were grown during a week, and small aliquots were regularly taken for cell counting using a Neubauer chamber. The antibiotic blasticidin was present in culture medium to maintain the pDM326 plasmid in *D. discoideum* cells. For activity assays, the activities were corrected by the total protein mass, measured by Bradford. (C) Western blotting analysis of the expression of wild-type (WT) or DdFXN G122V variant from plasmid pDM304. For G122V, upper: and lower panels correspond to 30 s and 2min exposition,

627 respectively. + and – symbols indicate transformation of *D. discoideum* with plasmid
628 pDM304 with or without frataxin gene insert (wild type or DdFXN G122V variant).
629 The detection of the recombinant variant was carried out by using an anti-FLAG
630 monoclonal antibody that recognizes the FLAG sequence located in the C-terminal
631 stretch of the plasmidic frataxin.
632
633
Discussion and Conclusions

Friedreich's Ataxia is a rare disease caused by disfunction of frataxin, a

mitochondrial protein, coded in the nucleus. Frataxin is a key component of the

mitochondrial supercomplex responsible for Fe-S cluster assembly. Fe-S cluster

dependent activities are essential for the homeostasis of the cells. There is a high

diversity of cell functions that need the correct insertion of Fe-S cluster into protein

structure. The energetics of the cell, that is commanded by the mitochondrial

metabolism directly depends on Fe-S clusters because clusters are crucial in electron

transport reaction. Thus, the alteration of Fe-S cluster biosynthesis affects Complex I, II

and III. But Fe-S are also essential for other enzymatic activities like lipoic acid

synthesis, necessary for pyruvate dehydrogenase activity or substrate binding in

mitochondrial aconitase (Krebs cycle); ultimately given that pyruvate dehydrogenase

complex, Krebs cycle, the electron chain, and ATP synthesis are highly coupled

metabolisms, the alteration of Fe-S cluster assembly result in the impairment of

numerous ATP-dependent cellular processes.

In a previous work we showed that *D. discoideum* has a complete pathway

involved in the Fe-S cluster assembly and transferring, as found in mammalian cells.

Our inferences from protein sequence conservation and structure models suggested that

some proteins (human and *D. discoideum*) may be exchangeable. In fact, the residues

that when mutate result in FA, are practically fully conserved between human and *D.*

discoideum. In this study we observed that DdFXN can bind to the human

supercomplex NIAU with a high affinity, similarly to the human FXN variant.

Additionally, it was able to activate the L-Cys desulfurase function. This is indicative of

the role of protein-protein interaction surfaces and the high conservation score for the

residues involved in these interactions. This highlights the potential of *D. discoideum* as

a model system for FA.

We then showed that frataxin locus can be easily edited in *D. discoideum*. Taking

advantage of the haploid context of this amoeba, we constructed a completely frataxin-

deficient strain. In the edited *D. discoideum* clone 8, studied in this paper, a premature

STOP codon occurred. The frataxin fragment encoded in the genomic DNA seemed to

be extremely unstable as judged by its absence in clone 8 amoeba cellular lysates, a

result that matches with molecular dynamics simulations that suggested highly mobile

protein backbone compared to the wild type DdFXN. This strain presents a significant

decrease of the Fe-S cluster dependent enzymatic functions and an altered phenotype.

Reduced aconitase and succinate dehydrogenase activities, a decrease in growth rate,

and higher sensitivity to oxidative stress were consistently observed; being all these

landmarks of frataxin deficiency in other Friedreich's Ataxia cellular and organism

models. Furthermore, we explored other processes that have been linked to

mitochondrial disfunction in *D. discoideum* and found defects during the multicellular

development. Remarkably, we were able to rescue the phenotype alterations with the

constitutive expression of wild type DdFXN. The expression of wild-type protein was

able to fully rescue growth and multicellular behavior alterations while the enzymatic

activities decrease were only partially rescued. There are a few conceivable, not

mutually exclusive, explanations for this result. The frataxin expression levels obtained

when expressed from plasmids are different than endogenous. Furthermore, given that

frataxin is imported into the mitochondrial matrix, the higher expression levels of the

protein might also affect the importing and processing machinery, although no effect

was observed when constitutive expression was carried out in wild-type AX2 strain. On

the other hand, it is also worth noticing that the DdFXN sequence used for protein

expression from plasmids were engineered carrying a C-terminal FLAG and EPEA tags

for protein identity assignment that could affect some roles of frataxin but not others.

The fact that FLAG sequence includes a highly negatively charged stretch of residues,

makes feasible that it alters in some degree frataxin interaction with the *D. discoideum*

NIAU supercomplex. Experiments with untagged versions would help to test this

hypothesis.

We then use this set-up to assess the rescuing capacity of DdFXN G122V, a

frataxin version that mimics a pathogenic variant found in Friedreich's ataxia patients

(i.e.: G122V in *D. discoideum* corresponds to G130V in humans). This variant, when is

constitutively expressed, fully rescues the defects on growth but only marginally rescue

the defects on multicellular development. This points out that DdFXN G122V is not

merely a less stable frataxin variant, what should be bypassed by the increased

expression levels; so it may be that other defects, besides protein stability, in G122V are

more important for some roles of frataxin than for others; this would mean that using

clone 8 of *D. discoideum* the roles of frataxin within the cell can be dissected. The

enzymatic activities defects were rescue to the same levels as in the case of the

expression of the wild-type C-terminal tagged DdFXN.

In this context, this new biological model offers a wide range of options to easily

explore diverse phenotypes occurring in FA, making *D. discoideum* very attractive to

study in a very simple fashion the effect of the diversity of FA variants that have been

already found in a simple fashion. Moreover, this model may help to understand

whether frataxin works as a bottle neck for some processes, whereas in other processes,

it has lower flux control, at least under the experimental conditions assayed.

In summary we have generated a system where we can easily monitor the effects

of lacking frataxin activity; this opens the door to develop drug or treatment screenings

that would help to design, and/or evaluate therapeutical strategies. Besides this

biological model offers a wide range of possibilities to easily explore diverse

phenotypes in FA. To complete the list of desirable attributes for a disease model

organism, it has been proved that frataxin locus can be easily edited, what opens the

possibility of generating specific strains that express the variant of interest from

endogenous locus.

Acknowledgments

We are grateful to CONICET and Universidad de Buenos Aires. We especially

thanks Fundación Ciencias Exactas y Naturales (FUNDACEN) and ALAPA.

Funding Sources

This study was supported by the *Agencia Nacional de Promoción de la*

Investigación, el Desarrollo Tecnológico y la Innovación through grant No. PICT 2019-

0083, Universidad de Buenos Aires (UBACyT 20020190100338BA), the *Consejo*

Nacional de Investigaciones Científicas y Técnicas (CONICET), and the Friedreich's

Ataxia Research Alliance (FARA).

References

1.   L. Fets, R. Kay, F. Velazquez, Dictyostelium. *Current Biology*. **20**, R1008--
R1010 (2010).
2.   J. M. Goldberg, G. Manning, A. Liu, P. Fey, K. E. Pilcher, Y. Xu, J. L. Smith,
The Dictyostelium Kinome—Analysis of the Protein Kinases from a Simple
Model Organism. *PLoS Genet*. **2**, e38 (2006).
3.   J. D. Dunn, C. Bosmani, C. Barisch, L. Raykov, L. H. Lefrançois, E. Cardenal-
Muñoz, A. T. López-Jiménez, T. Soldati, Eat Prey, Live: Dictyostelium
discoideum As a Model for Cell-Autonomous Defenses. *Front Immunol*. **8**, 1906
(2018).
4.   J. Martín-González, J.-F. Montero-Bullón, J. Lacal, Dictyostelium discoideum as

736 a non-mammalian biomedical model. *Microb Biotechnol*. **14**, 111–125 (2021).

737 5. H. N. Haver, K. M. Scaglione, Dictyostelium discoideum as a Model for
738 Investigating Neurodegenerative Diseases. *Front Cell Neurosci*. **15** (2021),
739 doi:10.3389/FNCEL.2021.759532.

740 6. J. Schaf, J. Damstra-Oddy, R. S. B. Williams, Dictyostelium discoideum as a
741 pharmacological model system to study the mechanisms of medicinal drugs and
742 natural products. *International Journal of Developmental Biology*. **63**, 541–550
743 (2019).

744 7. K. Augustin, A. Khabbush, S. Williams, S. Eaton, M. Orford, J. H. Cross, S. J. R.
745 Heales, M. C. Walker, R. S. B. Williams, Mechanisms of action for the medium-
746 chain triglyceride ketogenic diet in neurological and metabolic disorders. *Lancet*
747 *Neurol*. **17**, 84–93 (2018).

748 8. R. H. Kessin, *Dictyostelium* (Cambridge University Press, Cambridge, 2001;
749 http://ebooks.cambridge.org/ref/id/CBO9780511525315).

750 9. G. Bloomfield, Sex and macrocyst formation in Dictyostelium. *International*
751 *Journal of Developmental Biology*. **63**, 439–446 (2019).

752 10. K. Yamashita, H. Iriki, Y. Kamimura, T. Muramoto, CRISPR Toolbox for
753 Genome Editing in Dictyostelium. *Front Cell Dev Biol*. **9** (2021),
754 doi:10.3389/FCELL.2021.721630.

755 11. T. Ogasawara, J. Watanabe, R. Adachi, Y. Ono, Y. Kamimura, T. Muramoto,
756 CRISPR/Cas9-based genome-wide screening of Dictyostelium. *Scientific Reports*
757 *2022 12:1*. **12**, 1–13 (2022).

758 12. T. Muramoto, H. Iriki, J. Watanabe, T. Kawata, Recent Advances in
759 CRISPR/Cas9-Mediated Genome Editing in Dictyostelium. *Cells*. **8**, 46 (2019).

760 13. G. P. Otto, M. Cocorocchio, L. Munoz, R. A. Tyson, T. Bretschneider, R. S. B.
761 Williams, "Employing Dictyostelium as an advantageous 3Rs model for
762 pharmacogenetic research" in *Methods in Molecular Biology* (Humana Press Inc.,
763 2016; https://pubmed.ncbi.nlm.nih.gov/27271898/), vol. 1407, pp. 123–130.

764 14. A. Tornero-Écija, L. C. Tábara, M. Bueno-Arribas, L. Antón-Esteban, C.
765 Navarro-Gómez, I. Sánchez, O. Vincent, R. Escalante, A Dictyostelium model
766 for BPAN disease reveals a functional relationship between the WDR45/WIPI4
767 homolog Wdr45l and Vmp1 in the regulation of autophagy-associated PtdIns3P
768 and ER stress. *Autophagy*. **18**, 661–677 (2022).

769 15. J. Rodriguez-Centeno, R. Perona, L. Sastre, Dyskerin Mutations Present in
770 Dyskeratosis Congenita Patients Increase Oxidative Stress and DNA Damage
771 Signalling in Dictyostelium Discoideum. *Cells 2019, Vol. 8, Page 1406*. **8**, 1406
772 (2019).

773 16. A. C. Adam, C. Bornhövd, H. Prokisch, W. Neupert, K. Hell, The Nfs1
774 interacting protein Isd11 has an essential role in Fe/S cluster biogenesis in
775 mitochondria. *EMBO Journal*. **25**, 174–183 (2006).

776 17. S. C. Lim, M. Friemel, J. E. Marum, E. J. Tucker, D. L. Bruno, L. G. Riley, J.
777 Christodoulou, E. P. Kirk, A. Boneh, C. M. DeGennaro, M. Springer, V. K.
778 Mootha, T. A. Rouault, S. Leimkühler, D. R. Thorburn, A. G. Compton,
779 Mutations in LYRM4, encoding iron-sulfur cluster biogenesis factor ISD11,
780 cause deficiency of multiple respiratory chain complexes. *Hum Mol Genet*. **22**,
781 4460–4473 (2013).

782 18. L. Böttinger, C. U. Martensson, J. Song, N. Zufall, N. Wiedemann, T. Becker,
783 Respiratory chain supercomplexes associate with the cysteine desulfurase
784 complex of the iron-sulfur cluster assembly machinery. *Mol Biol Cell*. **29**, 776–
785 785 (2018).

19.   J. K. Barton, R. M. B. Silva, E. O'Brien, Redox chemistry in the genome:
Emergence of the [4fe4s] cofactor in repair and replication. *Annu Rev Biochem*.
88, 163–190 (2019).
20.   A. K. Pandey, J. Pain, A. Dancis, D. Pain, Mitochondria export iron-sulfur and
sulfur intermediates to the cytoplasm for iron-sulfur cluster assembly and tRNA
thiolation in yeast. *Journal of Biological Chemistry*. **294**, 9489–9502 (2019).
21.   J. A. Mayr, R. G. Feichtinger, F. Tort, A. Ribes, W. Sperl, Lipoic acid
biosynthesis defects. *J Inherit Metab Dis*. **37**, 553–563 (2014).
22.   T. A. Richards, M. van der Giezen, Evolution of the Isd11-IscS complex reveals
a single alpha-proteobacterial endosymbiosis for all eukaryotes. *Mol Biol Evol*.
23, 1341–1344 (2006).
23.   M. Georgina Herrera, M. Florencia Pignataro, M. Ezequiel Noguera, K. Magalí
Cruz, J. Santos, M. G. Herrera, M. F. Pignataro, M. E. Noguera, K. M. Cruz, J.
Santos, Rescuing the Rescuer: On the Protein Complex between the Human
Mitochondrial Acyl Carrier Protein and ISD11. *ACS Chem Biol*. **13**, 1455–1462
(2018).
24.   K. Cai, R. O. Frederick, M. Tonelli, J. L. Markley, Mitochondrial Cysteine
Desulfurase and ISD11 Coexpressed in Escherichia coli Yield Complex
Containing Acyl Carrier Protein. *ACS Chem Biol*. **12**, 918–921 (2017).
25.   M. G. Herrera, M. E. Noguera, K. E. Sewell, W. A. Agudelo Suárez, L. Capece,
S. Klinke, J. Santos, Structure of the Human ACP-ISD11 Heterodimer.
Biochemistry. **58** (2019), doi:10.1021/acs.biochem.9b00539.
26.   N. G. Fox, X. Yu, X. Feng, H. J. Bailey, A. Martelli, J. F. Nabhan, C. Strain-
Damerell, C. Bulawa, W. W. Yue, S. Han, Structure of the human frataxin-bound
iron-sulfur cluster assembly complex provides insight into its activation
mechanism. **10**, 2210 (2019).
27.   S. Gervason, D. Larkem, A. ben Mansour, T. Botzanowski, C. S. Müller, L.
Pecqueur, G. le Pavec, A. Delaunay-Moisan, O. Brun, J. Agramunt, A. Grandas,
M. Fontecave, V. Schünemann, S. Cianférani, C. Sizun, M. B. Tolédano, B.
D'Autréaux, Physiologically relevant reconstitution of iron-sulfur cluster
biosynthesis uncovers persulfide-processing functions of ferredoxin-2 and
frataxin. *Nat Commun*. **10** (2019), doi:10.1038/S41467-019-11470-9.
28.   S. M. K. Farhan, J. Wang, J. F. Robinson, P. Lahiry, V. M. Siu, C. Prasad, J. B.
Kronick, D. A. Ramsay, C. Anthony Rupar, R. A. Hegele, Exome sequencing
identifies nfs1 deficiency in a novel fe-s cluster disease, infantile mitochondrial
complex ii/iii deficiency. *Mol Genet Genomic Med*. **2**, 73–80 (2014).
29.   F. Mochel, M. A. Knight, W. H. Tong, D. Hernandez, K. Ayyad, T. Taivassalo,
P. M. Andersen, A. Singleton, T. A. Rouault, K. H. Fischbeck, R. G. Haller,
Splice Mutation in the Iron-Sulfur Cluster Scaffold Protein ISCU Causes
Myopathy with Exercise Intolerance. *Am J Hum Genet*. **82**, 652–660 (2008).
30.   S. E. Faraj, E. A. Roman, M. Aran, M. Gallo, J. Santos, The alteration of the C-
terminal region of human frataxin distorts its structural dynamics and function.
FEBS Journal. **281**, 3397–3419 (2014).
31.   N. Faggianelli, R. Puglisi, L. Veneziano, S. Romano, M. Frontali, T. Vannocci,
S. Fortuni, R. Testi, A. Pastore, Analyzing the effects of a G137V mutation in the
FXN gene. *Front Mol Neurosci*. **8**, 1–8 (2015).
32.   A. R. Correia, C. Pastore, S. Adinolfi, A. Pastore, C. M. Gomes, Dynamics,
stability and iron-binding activity of frataxin clinical mutants. *FEBS J*. **275**,
3680–3690 (2008).
33.   P. Cavadini, C. Gellera, P. I. Patel, G. Isaya, Human frataxin maintains

mitochondrial iron homeostasis in Saccharomyces cerevisiae. *Hum Mol Genet*. **9**,
2523–2530 (2000).

34. D. Doni, L. Passerini, G. Audran, S. R. A. Marque, M. Schulz, J. Santos, P.
Costantini, M. Bortolus, D. Carbonera, Effects of Fe2+/Fe3+ Binding to Human
Frataxin and Its D122Y Variant, as Revealed by Site-Directed Spin Labeling
(SDSL) EPR Complemented by Fluorescence and Circular Dichroism
Spectroscopies. *International Journal of Molecular Sciences 2020, Vol. 21, Page
9619*. **21**, 9619 (2020).

35. C. L. Tsai, J. Bridwell-Rabb, D. P. Barondeau, Friedreich's ataxia variants I154F
and W155R diminish frataxin-based activation of the iron-sulfur cluster assembly
complex. *Biochemistry*. **50**, 6478 (2011).

36. J. Bridwell-Rabb, A. M. Winn, D. P. Barondeau, Structure - Function analysis of
friedreich's ataxia mutants reveals determinants of frataxin binding and
activation of the fe - S assembly complex. *Biochemistry*. **50**, 7265–7274 (2011).

37. F. Saccà, A. Marsili, G. Puorro, A. Antenora, C. Pane, A. Tessa, P. Scoppettuolo,
C. Nesti, V. Brescia Morra, G. de Michele, F. M. Santorelli, A. Filla, Clinical use
of frataxin measurement in a patient with a novel deletion in the FXN gene. *J
Neurol*. **260**, 1116–1121 (2013).

38. D. Poburski, J. B. Boerner, M. Koenig, M. Ristow, R. Thierbach, Time-resolved
functional analysis of acute impairment of frataxin expression in an inducible cell
model of Friedreich ataxia. *Biol Open*. **5**, 654 (2016).

39. M. Cossée, H. Puccio, A. Gansmuller, H. Koutnikova, A. Dierich, M. LeMeur,
K. Fischbeck, P. Dollé, M. Kœnig, Inactivation of the Friedreich ataxia mouse
gene leads to early embryonic lethality without iron accumulation. *Hum Mol
Genet*. **9**, 1219–1226 (2000).

40. M. v. Busi, M. v. Maliandi, H. Valdez, M. Clemente, E. J. Zabaleta, A. Araya, D.
F. Gomez-Casati, Deficiency of Arabidopsis thaliana frataxin alters activity of
mitochondrial Fe–S proteins and induces oxidative stress. *The Plant Journal*. **48**,
873–882 (2006).

41. N. Ventura, S. L. Rea, R. Testi, Long-lived C. elegans Mitochondrial mutants as
a model for human mitochondrial-associated diseases. *Exp Gerontol*. **41**, 974–
991 (2006).

42. J. v. Llorens, J. A. Navarro, M. J. Martínez-Sebastián, M. K. Baylies, S.
Schneuwly, J. A. Botella, M. D. Moltó, Causative role of oxidative stress in a
Drosophila model of Friedreich ataxia. *The FASEB Journal*. **21**, 333–344 (2007).

43. P. R. Anderson, K. Kirby, A. J. Hilliker, J. P. Phillips, RNAi-mediated
suppression of the mitochondrial iron chaperone, frataxin, in Drosophila. *Hum
Mol Genet*. **14**, 3397–3405 (2005).

44. I. Zanella, M. Derosas, M. Corrado, E. Cocco, P. Cavadini, G. Biasiotto, M. Poli,
R. Verardi, P. Arosio, The effects of frataxin silencing in HeLa cells are rescued
by the expression of human mitochondrial ferritin. *Biochimica et Biophysica Acta
(BBA) - Molecular Basis of Disease*. **1782**, 90–98 (2008).

45. C. Lu, G. Cortopassi, Frataxin knockdown causes loss of cytoplasmic iron–sulfur
cluster functions, redox alterations and induction of heme transcripts. *Arch
Biochem Biophys*. **457**, 111 (2007).

46. N. Calmels, S. Schmucker, M. Wattenhofer-Donzé, A. Martelli, N. Vaucamps, L.
Reutenauer, N. Messaddeq, C. Bouton, M. Koenig, H. Puccio, The first cellular
models based on frataxin missense mutations that reproduce spontaneously the
defects associated with Friedreich ataxia. *PLoS One*. **4** (2009),
doi:10.1371/journal.pone.0006379.

47. F. Lupoli, T. Vannocci, G. Longo, N. Niccolai, A. Pastore, The role of oxidative
stress in Friedreich's ataxia. *FEBS Lett*. **592**, 718–727 (2018).

48. F. Codazzi, A. Hu, M. Rai, S. Donatello, F. S. Scarzella, E. Mangiameli, I.
Pelizzoni, F. Grohovaz, M. Pandolfo, Friedreich ataxia-induced pluripotent stem
cell-derived neurons show a cellular phenotype that is corrected by a benzamide
HDAC inhibitor. *Hum Mol Genet*. **25** (2016), doi:10.1093/hmg/ddw308.

49. J. Olmos, M. F. Pignataro, A. B. B. dos Santos, M. Bringas, S. Klinke, L.
Kamenetzky, F. Velazquez, J. Santos, A highly conserved iron-sulfur cluster
assembly machinery between humans and amoeba dictyostelium discoideum:
The characterization of frataxin. *Int J Mol Sci*. **21**, 1–25 (2020).

50. R. Sekine, T. Kawata, T. Muramoto, CRISPR/Cas9 mediated targeting of
multiple genes in Dictyostelium OPEN. *Scientific RepoRts* |. **8**, 8471 (2018).

51. H. Land, M. S. Humble, YASARA: A Tool to Obtain Structural Guidance in
Biocatalytic Investigations. *Methods Mol Biol*. **1685**, 43–67 (2018).

52. R. Evans, M. O'Neill, A. Pritzel, N. Antropova, A. Senior, T. Green, A. Žídek, R.
Bates, S. Blackwell, J. Yim, O. Ronneberger, S. Bodenstein, M. Zielinski, A.
Bridgland, A. Potapenko, A. Cowie, K. Tunyasuvunakool, R. Jain, E. Clancy, P.
Kohli, J. Jumper, D. Hassabis, *bioRxiv*, in press, doi:10.1101/2021.10.04.463034.

53. J. A. Maier, C. Martinez, K. Kasavajhala, L. Wickstrom, K. E. Hauser, C.
Simmerling, ff14SB: Improving the Accuracy of Protein Side Chain and
Backbone Parameters from ff99SB. *J Chem Theory Comput*. **11**, 3696–3713
(2015).

54. E. Krieger, G. Vriend, New ways to boost molecular dynamics simulations. *J
Comput Chem*. **36**, 996 (2015).

55. M. Feoktistova, P. Geserick, M. Leverkus, *Cold Spring Harb Protoc*, in press,
doi:10.1101/PDB.PROT087379.

56. T. J. Gibson, E. v Koonin, G. Musco, A. Pastore, P. Bork, Friedreich's ataxia
protein: Phylogenetic evidence for mitochondrial dysfunction. *Trends Neurosci*.
19, 465–468 (1996).

57. J. M. Ashworth, D. J. Watts, Metabolism of the cellular slime mould
Dictyostelium discoideum grown in axenic culture. *Biochemical Journal*. **119**,
175 (1970).

58. T. Ast, J. D. Meisel, S. Patra, H. Wang, R. M. H. Grange, S. H. Kim, S. E. Calvo,
L. L. Orefice, F. Nagashima, F. Ichinose, W. M. Zapol, G. Ruvkun, D. P.
Barondeau, V. K. Mootha, Hypoxia Rescues Frataxin Loss by Restoring Iron
Sulfur Cluster Biogenesis. *Cell*. **177**, 1507-1521.e16 (2019).

59. S. Al-Mahdawi, R. M. Pinto, D. Varshney, L. Lawrence, M. B. Lowrie, S.
Hughes, Z. Webster, J. Blake, J. M. Cooper, R. King, M. A. Pook, GAA repeat
expansion mutation mouse models of Friedreich ataxia exhibit oxidative stress
leading to progressive neuronal and cardiac pathology. *Genomics*. **88**, 580–590
(2006).

60. K. Chantrel-Groussard, V. Geromel, H. Puccio, M. Koenig, A. Munnich, A.
Rötig, P. Rustin, Disabled early recruitment of antioxidant defenses in
Friedreich's ataxia. *Hum Mol Genet*. **10**, 2061–2067 (2001).

61. C. M. Gomes, R. Santos, Neurodegeneration in Friedreich's Ataxia: From
Defective Frataxin to Oxidative Stress. *Oxid Med Cell Longev*. **2013** (2013),
doi:10.1155/2013/487534.

62. R. Santos, N. Buisson, S. A. B. Knight, A. Dancis, J. M. Camadro, E. Lesuisse,
Candida albicans lacking the frataxin homologue: a relevant yeast model for
studying the role of frataxin. *Mol Microbiol*. **54**, 507–519 (2004).

63. R. P. Vázquez-Manrique, P. González-Cabo, S. Ros, H. Aziz, H. A. Baylis, F.
Palau, Reduction of Caenorhabditis elegans frataxin increases sensitivity to
oxidative stress, reduces lifespan, and causes lethality in a mitochondrial
complex II mutant. *The FASEB Journal*. **20**, 172–174 (2006).
64. A. Wong, J. Yang, P. Cavadini, C. Gellera, B. Lonnerdal, F. Taroni, G.
Cortopassi, The Friedreich's Ataxia Mutation Confers Cellular Sensitivity to
Oxidant Stress Which Is Rescued by Chelators of Iron and Calcium and
Inhibitors of Apoptosis. *Hum Mol Genet*. **8**, 425–430 (1999).
65. L. M. Francione, S. J. Annesley, S. Carilla-Latorre, R. Escalante, P. R. Fisher,
The Dictyostelium model for mitochondrial disease. *Semin Cell Dev Biol*. **22**,
120–130 (2011).
66. M. Kotsifas, C. Barth, A. de Lozanne, S. T. Lay, P. R. Fisher, Chaperonin 60 and
mitochondrial disease in Dictyostelium.
949