Title: Genetic Association Studies in Transgender Cohorts: A Systematic Review and Meta-

Analysis

Authors: Blake Ashley[1,2] & Vincent Harley[1,2,3]

Affiliations

[1]Sex Development Laboratory, Hudson Institute of Medical Research, Melbourne, Victoria,

Australia.

[2]Department of Molecular and Translational Science, Monash University, Melbourne, Victoria,

Australia.

[3]Corresponding Author.

Email Addresses and Institutional Addresses

blake.ashley@monash.edu

Vincent.harley@hudson.org.au

Phone: +61 3 85722527

Institution Address: 27-31 Wright Street, Clayton, VIC, 3168, Australia

Keywords

Transgender, gender incongruence, gender dysphoria, genetic, meta-analysis, association, sex

hormones

Abstract

According to twin studies, there is a heritable contribution to gender incongruence, but the

genetic mechanisms of this are unknown. Recent efforts to identify an aetiology of gender

incongruence have focused on the hypothesis that sex hormones establish gender identity

through influencing the development of neuroanatomy. Candidate gene studies that have

sought to elucidate whether polymorphisms in sex steroidogenesis genes are overrepresented

in transgender populations have been equivocal. A systematic search for case-control genetic

association studies in transgender populations was conducted. Mean (+SD) or allele

frequencies were extracted and combined quantitatively in random effects meta-analysis,

summarised as standardised mean difference for continuous alleles or odds ratios for allele

frequencies. Eight studies were included in the analysis. These studies spanned

polymorphisms in five genes; the CAG repeat in *androgen receptor*, the TA repeat in

estrogen receptor alpha, the CA repeat in *estrogen receptor beta*, the TTTA repeat in *CYP19,*

and the T>C SNP in *CYP17*. Pooled estimates indicated that transgender women have a

significant overrepresentation of short *ESR1* alleles compared to cisgender men (OR = 1.23,

95% CI: 1.06, 1.44, p = 0.0089). This may contribute an increased likelihood of developing

gender incongruence amongst natal males. Future investigations into gender incongruence

should use genome-wide methods.

Introduction

Gender identity refers to one's psychological experience of their own gender. For most, physical sex and gender identity are congruent. However, for some individuals the two can be incongruent; their gender identity does not match their physical sexual characteristics. This incongruence can be distressing, and at clinically relevant levels, can be diagnosed as gender dysphoria according to the Diagnostic Statistical Manual of Mental Disorders (DSM-V), or gender incongruence (GI) according to the International Classification of Diseases (ICD-11). GI will be used throughout this paper to refer to the state of having an incongruence between gender identity and biological sex. Transgender men are individuals who were born physically as female but identify as women. Transgender women are individuals who were born physically as male but identify as men.

Recent conceptions of an aetiology of GI have incorporated biological factors. The prevailing hypothesis is that gender identity is established during foetal/early post-natal development via sex hormones influencing the development of the brain. The hypothesis supposes that in transgender individuals, the process by which sex hormones influence the development of gender identity is altered (Guillamon et al., 2016). This idea has its basis in the organisational/activational dogma of hormonal impacts on the sexual differentiation of the brain. Under this framework, sex hormones prenatally organise brain tissues through receptor interactions, which later manifests in sex-specific behaviours when 'activated' by subsequent hormonal release (Arnold, 2017). Given that gender identity is one of few variables strongly correlated with biological sex (i.e., most biological males also have a male gender identity), the current hypothesis of biological contributions to gender identity posits that it may be a sex-typical behaviour that is, at least in part, prenatally instantiated by the influence of sex

hormones. Evidence for this hypothesis primarily stems from studies in people with

Differences of Sex Development (DSDs), where variants in sex steroidogenesis genes alter

the development of chromosomal or anatomical sex. DSDs are associated with increased

rates of cross-gender identification and gender incongruence, such as in congenital adrenal

hyperplasia (Dessens et al., 2005; Pasterski et al., 2015; Seneviratne et al., 2021), partial

androgen insensitivity syndrome (Mazur, 2005), and 5-alpha reductase type 2 deficiency

(Cohen-Kettenis, 2005). Neuroanatomical studies have also demonstrated that transgender

individuals displays differences to cisgender individuals in various brain regions (Mueller et

al., 2021), however the degree to which these findings support aetiological hypotheses or

whether they are merely neurobiological representations of the experience of gender

incongruence is up for debate.

Though direct evidence for genetic variants functionally affecting sex hormone signalling in

transgender participants is limited, twin data does suggest a substantial genetic component,

with concordance rates in monozygotic twins being greater (37%) than in dizygotic twins

(0%) (Heylens et al., 2012). Most genetic-oriented study in transgender populations have

been candidate gene studies comparing the frequency of polymorphisms in various sex

hormone signalling genes between cisgender and transgender participants. The selection of

these specific genes has been based on the hypothesis outlined above. These polymorphisms

are thought to be functional, given their associations to other phenotypes. For example, the

trinucleotide CAG repeat polymorphism in exon 1 of the *androgen receptor* gene

(*AR*(CAG)n), the dinucleotide CA repeat in intron 5 of the *estrogen receptor 2* gene

(*ESR2*(CA)n), the TTTA repeat in intron 4 of aromatase (CYP19(TTTA)n), and the T>C

SNP in the 5' UTR of the *CYP17AI* gene have been associated with hormone levels in

adulthood (Gennari et al., 2004, 2005; Scariano et al., 2004; Sharp et al., 2004; Westberg et

al., 2001). As well as the dinucleotide TA repeat in the promotor region of the *estrogen*

receptor 1 gene (*ESR1*(TA)n), these polymorphisms are also associated with fertility (Xiao et

al., 2016), prostate cancer risk (Weng et al., 2017), and bone mineral density (Gennari et al.,

2005; Langdahl et al., 2000; Ogawa et al., 2000). Further, *AR*(CAG)n correlates with male-

typical patterns of brain structure (Perrin et al., 2008; Raznahan et al., 2010). *ESR1*(TA)n has

also been associated with 2D:4D ratio (Vaillancourt et al., 2012), which shows sex

differences (Hönekopp & Watson, 2010) and is thought to be a proxy measure of prenatal

hormone exposure (McIntyre, 2006). This is especially relevant in the context of GI, as

transgender women show feminised 2D:4D ratios compared to controls (Siegmann et al.,

2020).

The results of candidate gene studies in transgender cohorts have often been contradictory.

Longer *AR*(CAG)n have been found more frequently in transgender women (Hare et al.,

2009), though this has failed to replicate in several other studies (Fernández, Esteva, Gómez-

Gil, et al., 2014; Fernández et al., 2018; Foreman et al., 2019; Henningsson et al., 2005;

Lombardo et al., 2013; Ujike et al., 2009). However, three studies have found that *AR*(CAG)n

was significantly associated to transgender women when in conjunction with alleles of other

sex steroidogenesis genes, including *CYP19* (Henningsson et al., 2005), *PGT* and *COMT*

(Foreman et al., 2019) and *ESR2* (Fernández et al., 2018; Foreman et al., 2019). Notably, a

previous meta-analysis comparing mean *AR*(CAG)n in transgender women and cisgender

men found that transgender women had significantly, though very slightly, longer repeat

lengths (D'Andrea et al., 2020). However, the analysis did not include data from the second

largest association study in transgender women (Foreman et al., 2019), and modelling was

used to estimate mean *AR*(CAG)n length in one of the studies (Fernández et al., 2018), even

though exact allele frequencies were available in the supplementary data, potentially causing

inaccuracies. Two studies examining *AR*(CAG)n in transgender men found no significant

relationships (Fernández et al., 2018; Ujike et al., 2009). Shorter *ESR1*(TA)n were found in

transgender women in one study (Foreman et al., 2019) but not others (Cortés-Cortés et al.,

2017; Fernández, Delgado-Zayas, et al., 2020; Ujike et al., 2009), whilst transgender men

showed no difference in *ESR1*(TA)n compared to cisgender women in three studies (Cortés-

Cortés et al., 2017; Fernández, Delgado-Zayas, et al., 2020; Ujike et al., 2009). CA repeat

length in *ESR2* has been associated to both transgender women (Henningsson et al., 2005)

and transgender men (Fernández et al., 2018), though these results have also failed to

replicate in other studies (Fernández et al., 2018; Foreman et al., 2019; Hare et al., 2009;

Ujike et al., 2009). No studies have demonstrated an independent association between *CYP19*

and gender incongruence in transgender women or transgender men (Fernández, Esteva,

Gómez-Gil, et al., 2014; Fernández, Esteva, Gómez☐Gil, et al., 2014; Fernández et al., 2018;

Foreman et al., 2019; Hare et al., 2009; Ujike et al., 2009). Only one study has shown an

association between *CYP17* and gender incongruence (Bentz et al., 2008), whilst others have

not (Fernández et al., 2015; Foreman et al., 2019).

The equivocal results of these studies are likely due to small sample sizes and study

heterogeneity. For example, the results of genetic association studies are contingent on the

ethnicity of the participants, which may differ between studies. Furthermore, transgender

individuals, even within the same natal sex, may not be a homogenous group with the same

developmental trajectories. A conceptual distinction has been made between transgender

individuals who had early onset of GI and tend to be attracted to members of their natal sex,

whilst transgender individuals who have later onset tend to be attracted to the sex of their

gender identities (Blanchard, 1985; Guillamon et al., 2016). Thus, studies may differ in their

results due to differences in the subtypes of transgender participants recruited. Inconsistency

between study results has also contributed to disagreement in the literature about the degree

to which candidate gene studies support the hypothesis that GI has a basis in altered sex

hormone signalling. For example, some cite genetic association studies as supporting the

hypothesis that GI has at least a partial basis in altered sex hormone signalling during early

development (Polderman et al., 2018; Rosenthal, 2021), whilst some claim that results are too

nascent and contradictory to offer such evidence (Levin et al., 2022).

To gain further insight into the role, if any, of sex steroidogenesis genes in a genetic

contribution to GI, we conducted a systematic review and meta-analysis of previous

candidate gene studies. In studying the mechanisms by which GI may occur, we also

investigate how gender identity develops in general. We hypothesised that specific

polymorphisms in genes related to sex hormone signalling would be overrepresented in

transgender populations compared to cisgender populations, and thus that sex steroidogenesis

genes play a role in the development of gender identity.

Methods and Materials

This study was conducted in accordance with the Preferred Reporting Items for Systematic

Reviews and Meta-Analysis guidelines (Page et al., 2021) and the Meta-Analysis of

Observational Studies in Epidemiology checklist (Brooke et al., 2021).

<u>Eligibility Criteria</u>

Eligibility for inclusion in this analysis were: a) case-control genetic association study, b)

comparison of allele length/frequency between transgender women (cases) and cisgender

men (controls), or between transgender men (cases) and cisgender women (control), and c)

availability of either mean length and standard deviation (SD), or allele frequencies, or

adequate information to derive these, for both cisgender and transgender cohorts. When

participants were included in multiple studies, the study with the largest sample size was

included.

<u>Search Strategy</u>

MedLine, Scopus, and the Cochrane Library databases were searched. Searches were done

for papers published between January 2000 and August 2022. No restriction was put on the

language of the publication. The database search strategy was designed and conducted by

B.A, with guidance from Monash Health librarians on incorporating MeSH terms. In all three

databases, a range of keywords were used; "transgender", "gender incongruence", "gender

dysphoria", "gender identity disorder", "transsex*", "MTF", "FTM", "assigned sex",

"trans?women", "trans?men", "transgender women", "transgender men", "polymorphism*",

"variant*", "allele*", "repeat", "SNP", "RFLP", "sex steroid*", "sex hormone*",

"hormone*", "receptor*", "androgen receptor*", "estrogen receptor" "oestrogen receptor*",

"metabolism", "enzyme", "androgen*", "testosterone", "estrogen", "oestrogen", "aromatase",

"CYP17". In the MedLine and Cochrane Library databases, MeSH terms were also used;

"Transgender Persons", "Gender Dysphoria", "Transsexualism", "Polymorphism, Genetic",

"Polymorphism, Restriction Fragment Length", "Polymorphism, Single Nucleotide",

"Alleles", "Gonadal Steroid Hormones", "Testosterone", "Estrogens", "Aromatase".

Reference lists of retrieved articles were also scrutinised to identify further potential studies,

but no additional studies were found. Once the search was complete, titles and abstracts were

retrieved and reviewed for relevance. The full texts of publications that passed this stage were

then read to deem eligibility.

<u>Data Extraction</u>

Data was extracted by both B.A and V.H. From the eligible studies, first author, year,

geographic location, age of participants, diagnostic tool used to define cases, the gene(s)

examined, and allelic information for meta-analysis were extracted. In some cases, studies

analysed polymorphisms in different manners and thus reported different statistical

endpoints. For example, differences in AR(CAG)n were analysed by both t-tests

(Henningsson et al., 2005; Lombardo et al., 2013), Mann Whitney tests (Fernández et al.,

2018; Foreman et al., 2019; Ujike et al., 2009), and through dichotomisation of the repeat into

short and long forms and subsequent calculation of odds ratios (Henningsson et al., 2005).

Thus, before data extraction took place, it was decided that a) normally distributed repeat

length fragment polymorphisms (RFLPs) would be meta-analysed via the calculation of a

pooled standardised mean difference estimate, b) non-normally distributed RFLPs would be

meta-analysed via calculation of a pooled odds ratio estimate, and c) single nucleotide

polymorphisms would be meta-analysed via calculation of a pooled odds ratio estimate. For

normally distributed RFLPs, it was therefore necessary to obtain the means and standard

deviations for both case and controls groups. In cases where these were not explicitly

reported, there were calculated using a method that estimates mean and standard deviation

from distribution information (Wan et al., 2014), or obtained via contact with the original

authors. For non-normally distributed RFLPs, dichotomised allele frequencies, where short

alleles were defined as equal to or less than the median allele length in the corresponding

control group, were either obtained directly from publications or via contact of the original

authors, or calculated based on allele distributions supplied in the publications. Inconsistency

in data extraction were resolved via discussion and re-extraction of the data to reach

consensus.

<u>Quality of Studies Assessment</u>

The Newcastle-Ottawa Scale (NOS) (Mamikutty et al., 2021; Wells et al., 2014) was used to

assess the quality and risk of bias of the included studies. This scale measures study quality

along three domains; selection of participants, comparability of groups, and the measurement

of outcome. Studies can receive a maximum of 9 points and a minimum of 0. Scores of 0-4

are deemed low quality, 5 to 6 moderate, and 7-9 high quality. Study quality was assessed by

both B.A and V.H. Consensus was reached through discussion after assessment was

conducted individually. Inter-rater reliability was assessed via calculation of Cohen's kappa.

<u>Statistical Analysis</u>

The association between AR(CAG)n and GI was assessed via calculation of the pooled

standardised mean difference (SMD) with a 95% confidence interval. The associations

between $ESR1$(TA)n, $ESR2$(CA)n, $CYP19$(TTTA)n, and $CYP17$ (T>C) were assessed via

calculation of a pooled odds ratios with 95% confidence intervals, quantifying the strength of

the association between form of the allele (short/long for $ESR1$, $ESR2$ and $CYP19$ and T/C for

$CYP17$) and GI. Heterogeneity between the studies was assessed using the I^2 test and the

Cochrane Chi-square X^2 (Cochrane Q) statistic, where heterogeneity was defined as $I^2 =$

>50%, and p = <0.1, respectively (Cordero & Dans, 2021; Petitti, 2001; Song et al., 2001). In

cases of significant heterogeneity, leave-one-out sensitivity analysis was conducted to assess

which studied contributed significant heterogeneity. This was only carried out in analyses

that contained >3 studies. Given the studies included participants from different ethnic

backgrounds, random effects models were used to calculate pooled effect estimates in all

instances. The results of fixed and random effects models are essentially identical in the

absence of heterogeneity (Petitti, 2001), but random effects models are necessitated in

instances where there may be systematic differences in study characteristics, such as in the

ethnicity of the participants (Munafò et al., 2004).

Publication bias was assessed through visual examination of funnel plots and statistically

with the use of a regression test (Egger et al., 1997), followed by the trim-and-fill method

(Duval & Tweedie, 2000) to attempt correction of any suspected publication bias. Given that

the sensitivity of statistical methods in assessing publication bias is low for meta-analysis

with few studies (Munafò et al., 2004), publication bias was only tested for in this study

where analysis involved ≥ 3 studies, and $p < 0.1$ was used as the significance cut off for the

regression test to account for the lack of power. All analysis and visualisation was conducted

using R Studio (v. 1.4.1717) with the 'metafor' package (v3.0.2) (Viechtbauer, 2010)

Results

Study Selection

Figure 1: PRISMA flow chart for the study selection process.

After database searches and screening for inclusion/exclusion criteria, eight studies were

included in the quantitative analysis (figure 1). Amongst these studies, polymorphisms in 5

genes were examined in multiple (>2) studies, presenting the opportunity for meta-analysis.

These polymorphisms were *AR*(CAG)n, *ESR1*(TA)n, *ESR2*(CA)n, *CYP19*(TTTA)n, and

CYP17 (T>C).

Study Characteristics

Table 1: Characteristics of studies included in meta-analysis.

First Author (Year)	Geographic Location	Ethnicity of Participants	Age of Participants Mean (SD) or age range	Sample Size	Diagnostic Tool	Gene(s) Examined
Henningsson *et al.* (2005)	Sweden	Caucasian	N/A	29 TW 229 CM	DSM-IV (GID)	*AR*(CAG)n *ESR2*(CA)n *CYP19*(TTTA)n
Bentz *et al.* (2007)	Austria	Caucasian	N/A	102 TW 756 CM 49 TM 915 CW	DSM-IV	*CYP17*(T>C)
Ujike *et al.* (2009)	Japan	Japanese	TW = 34.9 (10) CM = 28.1 (6.9) TM = 28.3 (7.6) CW = 35.7 (10.7)	74 TW 106 CM 168 TM 169 CW	DSM-IV (GID) ICD-10	*AR*(CAG)n *ESR1*(TA)n *ESR2*(CAG)n *CYP19*(TTTA)n
Lombardo *et al.* (2013)	Italy	23 Caucasian Italians 1 Caucasian Eastern European 6 South Americans	TW = 24 – 39 CM = 25 - 40	30 TW 40 CM	DSM-IV	*AR*(CAG)n
Fernández *et al.* (2016)	Spain	Caucasian	N/A	317 TW 358 CM 223 TM 264 CW	DSM-V DSM-IV-TR	*CYP17*(T>C)
Fernández *et al.* (2018)	Spain	Caucasian	TW/TM = 18 – 47	549 TW	DSM-V DSM-IV	*AR*(CAG)n *ESR2*(TA)n

			CM/CW = 18 - 65	728 CM 425 TM 599 CW		*CYP19*(TTTA)n
Foreman *et al.* (2019)	Australia /USA	Caucasian	N/A	380 TW 344 CM	DSM-IV DSM-V	*AR*(CAG)n *ESR1*(TA)n *ESR2*(CA)n *CYP19*(TTTA)n *CYP17*(T>C)
Fernández *et al.* (2020)	Spain	Caucasian	18 – 47	323 TW 285 CM 235 TM 252 CW	ICD-11 ICD-10 DSM-IV-TR	*ESR1*(TA)n

260 SD = Standard deviation; DSM = Diagnostic and Statistical Manual of Mental Disorders; TR

261 = Text revision; ICD-11 = International Classification of Diseases 11[th] edition; ICD-10 =

262 Internal Classification of Diseases 10[th] edition; N/A = Not available.

267 Quality of Included Studies

268 Table 2: Newcastle-Ottawa Assessment Scale assessment of included studies. Total scores of

269 0-4 are deemed low quality, 5 to 6 moderate, and 7-9 high quality.

	Selection				Comparability		Outcome			Total
Study (First Author)	Definition	Represent-	Selection of	Definition of	Controls for	Controls for	Assessment of	Same Ascertain	Non-Respo	

	of Cases	ativeness of Cases	Controls	Controls	Ethnicity	Other Factors	Exposure	ment Method of Cases/Controls	nse Rate	
Henningsson *et al.* (Henningsson et al., 2005)	1	0	1	0	1	0	1	1	0	5
Bentz *et al.* (Bentz et al., 2008)	1	0	0	0	1	0	1	1	0	4
Ujike *et al.* (Ujike et al., 2009)	1	0	0	1	1	0	1	1	0	5
Lombardo *et al.* (Lombardo et al., 2013)	1	0	0	1	0	0	1	1	0	4
Fernández *et al.* (Fernández et al., 2016)	1	0	1	1	1	1	1	1	0	7
Fernández *et al.* (Fernández et al., 2018)	1	0	1	1	1	1	1	1	0	7
Foreman *et al.* (Foreman et al., 2019)	1	0	0	1	0	0	1	1	0	4
Fernández *et al.* (Fernández, Delgado-Zayas, et al., 2020)	1	0	0	0	1	1	1	1	0	5

271   The quality ratings of studies included in the meta-analyses are shown in table 1. Three of the

272   studies were rated low quality, two were rated moderate quality, and two were rated high

273   quality. Inter-rater reliability was substantial (k = 0.76) before consensus.

274   <u>Synthesis of Association Studies</u>

Study	Transgender Women			Cisgender Men				Odds Ratio [95% CI]
ESR1	S	L*	n	S	L*	n		
Fernández et al. (2020)	296	200	248	199	159	179		1.18 [0.90, 1.56]
Foreman et al. (2018).1	376	272	324	288	268	278		1.29 [1.02, 1.62]
Ujike et al. (2009).1	90	58	74	118	94	106		1.24 [0.81, 1.89]
Total			646			563		
RE Model for ESR1 (Q = 0.21, df = 2, p = 0.90; I^2 = 0.0%, τ^2 = 0.00)								1.24 [1.06, 1.46]
p (Overall Model) = 0.0089								
ESR2	L	S*	n	L	S*	n		
Fernández et al. (2018).1	416	681	549	572	884	728		0.94 [0.80, 1.11]
Foreman et al. (2018).2	137	531	334	109	457	283		1.08 [0.82, 1.43]
Ujike et al. (2009).2	59	89	74	83	129	106		1.03 [0.67, 1.58]
Henningsson et al. (2005).1	19	34	27	87	371	229		2.38 [1.30, 4.38]
Total			984			1346		
RE Model for ESR2 (Q = 8.54, df = 3, p = 0.04; I^2 = 71.8%, τ^2 = 0.06)								1.15 [0.85, 1.56]
p (Overall Model) = 0.37								
CYP19	L	S*	n	L	S*	n		
Fernández et al. (2018).2	461	637	549	656	800	728		0.88 [0.75, 1.03]
Foreman et al. (2018).3	309	343	326	252	314	283		1.12 [0.90, 1.41]
Ujike et al. (2009).3	59	89	74	65	147	106		1.50 [0.97, 2.33]
Henningsson et al. (2005).2	31	27	29	225	233	229		1.19 [0.69, 2.06]
Total			978			1346		
RE Model for CYP19 (Q = 7.04, df = 3, p = 0.07; I^2 = 58.4%, τ^2 = 0.03)								1.08 [0.86, 1.36]
p (Overall Model) = 0.49								
CYP17	C	T*	n	C	T*	n		
Foreman et al. (2018).4	274	384	329	202	354	278		1.25 [0.99, 1.58]
Fernández et al. (2016)	246	388	317	283	433	358		0.97 [0.78, 1.21]
Bentz et al. (2008)	78	126	102	604	908	756		0.93 [0.69, 1.26]
Total			748			1392		
RE Model for CYP17 (Q = 3.29, df = 2, p = 0.19; I^2 = 39.9%, τ^2 = 0.01)								1.05 [0.87, 1.26]
p (Overall Model) = 0.59								

a

0.25   1   4

Study	Transgender Women			Cisgender Men				SMD [95% CI]
	Mean	SD	n	Mean	SD	n		
Henningsson et al. (2005)	18.86	2.79	29	18.61	2.59	229		0.10 [-0.29, 0.48]
Ujike et al. (2009)	21	2.18	74	20.87	3.41	106		0.04 [-0.25, 0.34]
Lombardo et al. (2013)	21.7	3.4	30	21.9	2.84	40		-0.06 [-0.54, 0.41]
Foreman et al. (2018)	244.64	9.9	328	243.91	8.76	282		0.08 [-0.08, 0.24]
Fernández et al. (2018)	19.05	3.13	550	19.01	2.78	728		0.01 [-0.10, 0.12]
Total			1011			1385		
RE Model (Q = 0.68, df = 4, p = 0.95; I^2 = 0.0%)								0.03 [-0.05, 0.12]
p (Overall Effect) = 0.41								

b

-0.9   -0.45   0   0.45   0.9

Standardized Mean Difference

Figure 2: Forest plots of meta-analysis of dichotomised *ESR1*(TA)n, *ESR2*(CA)n,

CYP19(TTTA)n and *CYP17* (T>C) (a) and mean *AR*(CAG)n (b) between transgender women

and cisgender men. Diamonds represent overall summary estimates. Width of the diamond

represents the 95% confidence interval. Boxes indicate the effect estimate obtained in the

individual studies, and box width indicates the weight given to each individual study based

on its sample size. Whiskers indicate 95% confidence intervals. Odds ratios are in reference

to the reference allele for each gene, indicated by *. *ESR1*, *ESR2*, *CYP19*, and *CYP17* are

autosomal genes, so the number of alleles is double the number of participants. For

AR(CAG), Foreman *et al*. (2018) recorded *AR*(CAG)n as total length of the repeat sequence,

rather than the number of trinucleotide repeats as other studies did. SD = standard deviation,

n = number of participants, SMD = standardised mean difference, CI = confidence interval,

RE = random effects, df = degrees of freedom

Figure 3: Forest plots of meta-analysis of dichotomised *ESR1*(TA)n, *ESR2*(CA)n,

CYP19(TTTA)n and *CYP17* (T>C) (a) and mean *AR*(CAG)n (b) between transgender women

and cisgender men.

Figures 2 and 3 display the results of the meta-analyses in transgender women/cisgender men,

and transgender men/cisgender women, respectively. Out of all analyses, one significant

association was identified; transgender women had a significant overrepresentation of short

ESR1(TA)n compared to cisgender men (p = 0.0089) (Figure 2a). Heterogeneity tests found

significant heterogeneity between the results of the individual studies in the analysis of *ESR2*

in transgender women (figure 2a). Subsequent sensitivity analysis revealed that all

heterogeneity disappeared when the Henningsson et al., (2005) study, which had a much

smaller sample size than the other studies, was removed. After removal, the model became

more insignificant, with the OR tightening around 1 (OR = 0.98, 95% CI = [0.86, 1.12], p =

0.86). Significant heterogeneity was also identified between the studies in the analysis of

CYP19 in transgender women (figure 2a). Removal of Fernández et al. (2018) eliminated

heterogeneity, but the model remained insignificant (OR = 1.19, 95% CI = 0.99, 1.43], p =

0.07).

<u>Publication Bias</u>

Figure 4: Funnel plots of meta-analyses of allele difference in *AR*(CAG)n (a), *ESR1*(TA)n

(b), *ESR2*(CA)n (c), *CYP19*(TTTA)n (d) and *CYP19* T>C (e) in transgender women and

cisgender men.

Figure 4 displays the funnel plots obtained from testing for publication bias in the analyses of

all five genes in transgender women and cisgender men. Publication bias was not tested for in

the studies of transgender men as there were too few studies. Across *AR*, *ESR1*, *ESR2*,

CYP19, and *CYP17*, regression tests for funnel plot asymmetry were not significant (p = 0.86,

0.76, 0.16, 0.17, 0.77, respectively). Trim-and-fill analysis did not result in any potentially

missing studies being added to the funnel plots for *AR*(CAG)n, *ESR2*(CA)n, or *CYP17* T>C.

Two putative missing studies were added to the right side of the funnel plot for *ESR1*(TA)n,

and two putative missing studies were filled in on the left side of the plot for

CYP19(TTTA)n. The model remained non-significant for *CYP19*(TTTA) after these putative

studies were added (OR = 0.98, 95% CI = [0.80, 1.22], p = 0.88). For *ESR1*(TA)n, the model

remained significant after these studies were added (OR = 1.28, 95% CI = [1.13, 1.46], p =

0.0002. Thus, no changes to the overall results from these meta-analyses occurred after

testing for publication bias.

Discussion

This is the largest genetic analysis of transgender individuals conducted to date. Mean

AR(CAG)n, frequency of dichotomised short/long *ESR1*(TA)n, *ESR2*(CA)n, *CYP19*(TTTA)n

and T>C SNP frequency in *CYP17* were compared between transgender women and

cisgender men, and transgender men and cisgender women. One significant association was

identified, where transgender women had a significantly higher frequency of short

ESR1(TA)n, supporting our earlier finding (Foreman et al., 2019).

ESR1 is highly expressed throughout the human brain (Laflamme et al., 1998). The length of

the TA repeat in the promotor region of *ESR1* (*ESR1*(TA)n) correlates with the volume and

density of many structures in the brain which display sex differences (Tan et al., 2020).

However, there has been considerable debate regarding the role of estrogen receptors in the

sexual differentiation of the brain. In rodents, masculinisation is typically accepted as

occurring through the binding of locally aromatised testosterone to estrogen receptors, which

does not occur in females due to the protective role of α-fetoprotein (Zuloaga et al., 2008).

The role of estrogen receptors in human brain sexual differentiation is less clear. It has been

suggested sexual brain differentiation in humans occurs primarily through direct binding of

testosterone to androgen receptors (Puts & Motta-Mena, 2018), with female brain

differentiation occurring merely due to the absence of testosterone. However, some argue that

female brain differentiation is a more active process than this (Nugent et al., 2015), which

involving the action of *ER*s (Bakker & Brock, 2010; Luoto & Rantala, 2018). If GI in natal

males is the result of under masculinisation/feminisation of brain structures, our results may

suggest that *ESR1* plays an active role in sexual brain differentiation.

The finding that short *ESR1*(TA)n are overrepresented in transgender women may support the

idea that GI has its basis in altered sex hormone signalling influencing the development of

sexually differentiated brain structures. However, it is currently difficult to tie in genetics

findings with results from neuroanatomical studies in transgender individuals. For example, a

diffusion tensor imaging study demonstrated the superior longitudinal fascicles are feminised

in transgender women (Rametti et al., 2011). Feminisation of the region is correlated with

short *ESR1*(TA)n (Tan et al., 2020), which seems in congruence with our results. However, in

other brain regions that are feminised in transgender women, such as the hypothalamus

(Garcia-Falgueras & Swaab, 2008; Zhou et al., 1995), short *ESR1*(TA)n is associated with

masculinisation (Tan et al., 2020). Therefore, whilst estrogen signalling affects brain sexual

differentiation, our ability to relate it to specific brain regions in transgender individuals is

limited.

Other types of genetic analysis of transgender individuals have also implicated estrogen

signalling. First, a whole-exome analysis identified 5 variants in estrogen signalling pathway

genes that were unique amongst transgender individuals (Theisen et al., 2019), though this

study only included 30 transgender individuals. Second, a candidate gene study of *SRC* genes

showed certain SNPs were overrepresented amongst transgender individuals (Ramírez et al.,

2021). SRCs are coactivators of estrogen receptors (Yore et al., 2010) and are influential in

the masculinisation of the brain (Auger et al., 2000). Third, an epigenetic study (Fernández,

Ramírez, et al., 2020) demonstrated sex differences in the degree of methylation in a

regulatory region *ESR1*, and that both transgender women and transgender men had

significantly lower levels of methylation in this region compared to cisgender men prior to

receiving gender-affirming hormone therapy.

Contrary to several of the individual studies, no single-gene associations to GI were

demonstrated for *AR*(CAG)n, *ESR2*(CA)n, *CYP19*(TTTA)n, and *CYP17* T>C. The results are

also contrary to a previous meta-analysis that found significantly longer *AR*(CAG)n in

transgender women compared to cisgender men (D'Andrea et al., 2020). Our *AR*(CAG)n

analysis differs from D'Andrea *et al.,* (2020) in that we included data from Foreman *et al.,*

(2019), and we obtained direct allelic information from the supplementary material of

Fernández *et al.*, (2018) instead of using mathematical estimation.

Assessment of the quality of these candidate gene studies revealed that most studies were

either of low or moderate quality, with only two studies rated as high. A lack of

representativeness of cases was a pervasive issue. All studies recruited transgender

participants through gender clinics. Given that individuals who seek formal

diagnosis/treatment of GI through gender clinics make up a small portion of a wider

population who identify as transgender or otherwise gender-diverse (Arcelus et al., 2015;

Meerwijk & Sevelius, 2017), this may contribute to selection bias. Some studies did not

adequately control for ethnicity (Foreman et al., 2019; Lombardo et al., 2013). This is

important, as genetic association studies require cases and controls to be ethnically

homogenous to control for differences in allele frequencies and linkage disequilibrium

between ancestral populations (Hellwege et al., 2017). Only three studies controlled for age

of onset of GI and the sexual preferences of the participants (Fernández, Delgado-Zayas, et

al., 2020; Fernández et al., 2016, 2018). It has been hypothesised that transgender individuals

are not a homogenous group, but rather may be subtyped based on age of onset and sexual

attraction (Blanchard, 1989), which may have support from neuroimaging studies (Guillamon

et al., 2016). Based on this hypothesis, it may therefore be necessary to control for these

factors in genetic studies to create as homogenous groups as possible, as different subtypes of

gender incongruence may have unique developmental pathways involving different genetic

factors.

Some limitations of this analysis should be acknowledged. Firstly, the power of the analysis

was limited by the paucity of studies in this area. This is especially true of the analyses in

transgender men, which only two studies reported data on. Secondly, the ability to interpret

heterogeneity in the individual study results with reference to study characteristics such as

ethnicity age of participants, clinical criteria used to define cases, age of onset of GI, or

sexuality was limited due to the lack of studies. Thirdly, the combined data came from

ethnically heterogeneous samples, making it difficult to determine how ethnicity may

influence the results, due to potential variability in the allele frequency between ethnic

populations (Zitzmann & Nieschlag, 2003). Lastly, we were unable to perform gene-gene

interaction analysis due to the unavailability of genotype data.

It would be remiss to not address the ethical implications of conducting genetic research in

transgender populations. Concerns have been raised that genetics research may negatively

impact transgender people in several ways. For example, a survey of transgender or otherwise

gender-diverse individuals found that people felt that genetics research could contribute to the

gatekeeping of transgender identities (i.e., by encouraging people to think only those with a

particular marker are 'truly' transgender), the gatekeeping of gender-affirming care (i.e., only

those with a particular marker can access treatment), the pathologisation of transgender

identities as mental disorders, the perpetuation of a binary conception of gender, or open the

door to eugenic practices (Rajkovic et al., 2022). Although these are valid concerns, they

should be weighed against the potential positives of conducting such research. Given the

nature of complex traits, it is extremely unlikely that a single marker will ever be able to

differentiate transgender and cisgender individuals; indeed, in this study, many cisgender

men possessed the marker that was overrepresented in transgender women. On the contrary,

demonstrating certain markers are overrepresented in people who experience GI could

contribute to the de-stigmatisation of transgender identities in the eyes of the public, as

several studies suggest that biological views of transgender identities are associated with

positive attitudes towards transgender people, whilst the view that transgender identities are

environmentally caused are associated with negative attitudes (Bowers & Whitley, 2020;

Brown et al., 2017; Campbell et al., 2019; Ching & Chen, 2022; Elischberger et al., 2016,

2018; Greenburg & Gaia, 2019; Landén & Innala, 2000; Rad et al., 2019; Woodford et al.,

2012). Additionally, studying the underlying genetics of gender identity may promote

understanding transgender identities as representing natural human variation, rather than

reinforcing a binary sense of gender. Such research may also have medical applications, such

as for personalised medicine for transgender individuals. The genetics of sex hormones which

may contribute to gender identity, as well as the use of exogeneous hormones for gender-

affirmation, may affect how medical conditions are treated in transgender individuals (Moyer

et al., 2019).

Conclusion

This study is the largest genetic analysis of transgender individuals conducted to date. A

significant association between a polymorphism in *ESR1* and gender incongruence amongst

natal males was identified. However, future studies exploring genetic contributions to gender

should employ genome-wide approaches. Candidate gene studies are prone to false-positives,

likely due to the small effect sizes of genes that contribute to complex traits requiring large

samples sizes to detect, and also due to ethnic heterogeneity between studies (Munafò et al.,

2004). Further still, genome-wide association studies have revealed that our *a priori*

assumptions as to what would constitute viable candidate genes are often wrong; significant

associations in candidate gene studies do not replicate when the entire genome is scrutinised

(Border et al., 2019; Hall et al., 2013; Johnson et al., 2017). Psychology is always highly

multigenic (Plomin et al., 2016), and therefore no account of a genetic contribution to gender

identity based on the examination of a few genes will be a complete one. So far, candidate

gene studies have focussed exclusively on sex steroidogenesis genes, motivated by the

hypothesis that the process of sexual differentiation of the brain is what is altered in

transgender individuals. If this is the case, it is also possible that sex chromosome genes may

have an influence, as brain differentiation has been shown to be influenced by sex

chromosome genes before sex steroidogenesis begins (Corre et al., 2016). In light of recent

evidence demonstrating the transgender individuals prior to receiving gender affirming

hormone therapy show global DNA methylation differences to cisgender people (Ramirez et

al., 2021), and that such therapy induces epigenetic changes that make their epigenetic

profiles more similar to the opposite sex (Shepherd et al., 2022), it is possible that genes

which control methylation in response to sex hormones could be implicated. Further still, it is

possible that mechanisms outside of the sex differentiation pathway are involved in the

development of GI which have not yet been considered.

Declarations

<u>Funding</u>

Full statement in title page submission to aid masking.

<u>Conflicts of Interest</u>

The authors declare that they have no competing interests.

<u>Availability of Data and Materials</u>

The datasets used and/or analysed during the current study are available from the

corresponding author on reasonable request.

<u>Code Availability</u>

The R code used in this analysis are available from the corresponding author on reasonable

request.

<u>Author Contributions</u>

BA extracted data, conducted quality assessment of studies, conducted all statistical analysis,

and wrote and edited the manuscript. VH conceived the project, extracted data, conducted

quality assessment, and contributed to editing of the manuscript.

<u>Ethics Approval</u>

Not applicable.

<u>Consent to Participate/Publish</u>

Not applicable.

<u>Acknowledgements</u>

We would like to thank Alice Anderson of Monash Health library for assisting in

constructing the database search strategy.

Compliance with Ethical Standards

<u>Disclosure of Potential Conflicts of Interest</u>

The authors have no conflicts of interest to declare.

<u>Research Involving Human Participants and/or Animals</u>

Not applicable.

<u>Informed Consent</u>

Not applicable.

References

Arcelus, J., Bouman, W. P., Van Den Noortgate, W., Claes, L., Witcomb, G., & Fernandez-

Aranda, F. (2015). Systematic review and meta-analysis of prevalence studies in

transsexualism. *European Psychiatry: The Journal of the Association of European*

Psychiatrists, *30*(6), 807–815. https://doi.org/10.1016/j.eurpsy.2015.04.005

Arnold, A. P. (2017). A general theory of sexual differentiation. *Journal of Neuroscience*

Research, *95*(1–2), 291–300. https://doi.org/10.1002/jnr.23884

Auger, A. P., Tetel, M. J., & McCarthy, M. M. (2000). Steroid receptor coactivator-1 (SRC-

1) mediates the development of sex-specific brain morphology and behavior.

Proceedings of the National Academy of Sciences, *97*(13), 7551–7555.

https://doi.org/10.1073/pnas.97.13.7551

Bakker, J., & Brock, O. (2010). Early Oestrogens in Shaping Reproductive Networks:

Evidence for a Potential Organisational Role of Oestradiol in Female Brain

Development. *Journal of Neuroendocrinology*, *22*(7), 728–735.

https://doi.org/10.1111/j.1365-2826.2010.02016.x

Bentz, E.-K., Hefler, L. A., Kaufmann, U., Huber, J. C., Kolbus, A., & Tempfer, C. B.

(2008). A polymorphism of the CYP17 gene related to sex steroid metabolism is

associated with female-to-male but not male-to-female transsexualism. *Fertility and*

Sterility, *90*(1), 56–59. https://doi.org/10.1016/j.fertnstert.2007.05.056

Blanchard, R. (1985). Typology of male-to-female transsexualism. *Archives of Sexual*

Behavior, *14*(3), 247–261. https://doi.org/10.1007/BF01542107

Blanchard, R. (1989). The classification and labeling of nonhomosexual gender dysphorias.

Archives of Sexual Behavior, *18*(4), 315–334. https://doi.org/10.1007/BF01541951

Border, R., Johnson, E. C., Evans, L. M., Smolen, A., Berley, N., Sullivan, P. F., & Keller,

M. C. (2019). No Support for Historical Candidate Gene or Candidate Gene-by-

Interaction Hypotheses for Major Depression Across Multiple Large Samples. *The*

American Journal of Psychiatry, *176*(5), 376–387.

https://doi.org/10.1176/appi.ajp.2018.18070881

Bowers, M. M., & Whitley, C. T. (2020). What Drives Support for Transgender Rights?

Assessing the Effects of Biological Attribution on U.S. Public Opinion of

Transgender Rights. *Sex Roles*, *83*(7), 399–411. https://doi.org/10.1007/s11199-019-

01118-9

Brooke, B. S., Schwartz, T. A., & Pawlik, T. M. (2021). MOOSE Reporting Guidelines for

Meta-analyses of Observational Studies. *JAMA Surgery*, *156*(8), 787–788.

https://doi.org/10.1001/jamasurg.2021.0522

Brown, C., Keller, C. J., Brownfield, J. M., & Lee, R. (2017). Predicting Trans-Inclusive

Attitudes of Undergraduate Nursing Students. *Journal of Nursing Education*, *56*(11),

660–669. https://doi.org/10.3928/01484834-20171020-05

Campbell, M., Hinton, J. D. X., & Anderson, J. R. (2019). A systematic review of the

relationship between religion and attitudes toward transgender and gender-variant

people. *International Journal of Transgenderism*, *20*(1), 21–38.

https://doi.org/10.1080/15552739.2018.1545149

Ching, B. H.-H., & Chen, T. T. (2022). Effects of Biological Determinism on Beliefs and

Attitudes About Transgender People: Psychological Essentialism and Biased

Assimilation. *Archives of Sexual Behavior*, *51*(4), 1927–1942.

https://doi.org/10.1007/s10508-021-02262-8

Cohen-Kettenis, P. T. (2005). Gender Change in 46,XY Persons with 5α-Reductase-2

Deficiency and 17β-Hydroxysteroid Dehydrogenase-3 Deficiency. *Archives of Sexual*

Behavior, *34*(4), 399–410. https://doi.org/10.1007/s10508-005-4339-4

Cordero, C. P., & Dans, A. L. (2021). Key concepts in clinical epidemiology: Detecting and

dealing with heterogeneity in meta-analyses. *Journal of Clinical Epidemiology*, *130*,

149–151. https://doi.org/10.1016/j.jclinepi.2020.09.045

Corre, C., Friedel, M., Vousden, D. A., Metcalf, A., Spring, S., Qiu, L. R., Lerch, J. P., &

Palmert, M. R. (2016). Separate effects of sex hormones and sex chromosomes on

brain structure and function revealed by high-resolution magnetic resonance imaging

and spatial navigation assessment of the Four Core Genotype mouse model. *Brain

Structure and Function*, *221*(2), 997–1016. https://doi.org/10.1007/s00429-014-0952-

Cortés-Cortés, J., Fernández, R., Teijeiro, N., Gómez-Gil, E., Esteva, I., Almaraz, M. C.,

Guillamón, A., & Pásaro, E. (2017). Genotypes and Haplotypes of the Estrogen

Receptor α Gene (ESR1) Are Associated With Female-to-Male Gender Dysphoria.

The Journal of Sexual Medicine, *14*(3), 464–472.

https://doi.org/10.1016/j.jsxm.2016.12.234

D'Andrea, S., Pallotti, F., Senofonte, G., Castellini, C., Paoli, D., Lombardo, F., Lenzi, A.,

Francavilla, S., Francavilla, F., & Barbonetti, A. (2020). Polymorphic Cytosine-

Adenine-Guanine Repeat Length of Androgen Receptor Gene and Gender

Incongruence in Trans Women: A Systematic Review and Meta-Analysis of Case-

Control Studies. *The Journal of Sexual Medicine*, *17*(3), 543–550.

https://doi.org/10.1016/j.jsxm.2019.12.010

Dessens, A. B., Slijper, F. M. E., & Drop, S. L. S. (2005). Gender dysphoria and gender

change in chromosomal females with congenital adrenal hyperplasia. *Archives of

Sexual Behavior*, *34*(4), 389–397. https://doi.org/10.1007/s10508-005-4338-5

Duval, S., & Tweedie, R. (2000). Trim and Fill: A Simple Funnel-Plot–Based Method of

Testing and Adjusting for Publication Bias in Meta-Analysis. *Biometrics*, *56*(2), 455–

463. https://doi.org/10.1111/j.0006-341X.2000.00455.x

Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected

by a simple, graphical test. *BMJ*, *315*(7109), 629–634.

https://doi.org/10.1136/bmj.315.7109.629

Elischberger, H. B., Glazier, J. J., Hill, E. D., & Verduzco-Baker, L. (2016). "Boys Don't

Cry"—or Do They? Adult Attitudes Toward and Beliefs About Transgender Youth.

Sex Roles, *75*(5), 197–214. https://doi.org/10.1007/s11199-016-0609-y

Elischberger, H. B., Glazier, J. J., Hill, E. D., & Verduzco-Baker, L. (2018). Attitudes

Toward and Beliefs about Transgender Youth: A Cross-Cultural Comparison

Between the United States and India. *Sex Roles*, *78*(1), 142–160.

https://doi.org/10.1007/s11199-017-0778-3

Fernández, R., Cortés-Cortés, J., Esteva, I., Gómez-Gil, E., Almaraz, M. C., Lema, E.,

Rumbo, T., Haro-Mora, J.-J., Roda, E., Guillamón, A., & Pásaro, E. (2015). The

CYP17 MspA1 Polymorphism and the Gender Dysphoria. *The Journal of Sexual*

Medicine, *12*(6), 1329–1333. https://doi.org/10.1111/jsm.12895

Fernández, R., Cortés-Cortés, J., Gómez-Gil, E., Esteva, I., Almaraz, M. C., Guillamón, A.,

& Pásaro, E. (2016). The CYP17-MspA1 rs743572 polymorphism is not associated

with gender dysphoria. *Genes & Genomics*, *38*(12), 1145–1150.

https://doi.org/10.1007/s13258-016-0456-9

Fernández, R., Delgado-Zayas, E., Ramírez, K., Cortés-Cortés, J., Gómez-Gil, E., Esteva, I.,

Almaraz, M. C., Guillamon, A., & Pásaro, E. (2020). Analysis of Four

Polymorphisms Located at the Promoter of the Estrogen Receptor Alpha ESR1 Gene

in a Population With Gender Incongruence. *Sexual Medicine*, *8*(3), 490–500.

https://doi.org/10.1016/j.esxm.2020.04.002

Fernández, R., Esteva, I., Gómez⬜Gil, E., Rumbo, T., Almaraz, M. C., Roda, E.,

Haro⬜Mora, J., Guillamón, A., & Pásaro, E. (2014). The (CA)n Polymorphism of

ERβ Gene is Associated with FtM Transsexualism. *The Journal of Sexual Medicine*,

11(3), 720–728. https://doi.org/10.1111/jsm.12398

Fernández, R., Esteva, I., Gómez-Gil, E., Rumbo, T., Almaraz, M. C., Roda, E., Haro-Mora,

J.-J., Guillamón, A., & Pásaro, E. (2014). Association study of ERβ, AR, and

CYP19A1 genes and MtF transsexualism. *The Journal of Sexual Medicine*, *11*(12),

2986–2994. https://doi.org/10.1111/jsm.12673

Fernández, R., Guillamon, A., Cortés-Cortés, J., Gómez-Gil, E., Jácome, A., Esteva, I.,

Almaraz, M., Mora, M., Aranda, G., & Pásaro, E. (2018). Molecular basis of Gender

Dysphoria: Androgen and estrogen receptor interaction. *Psychoneuroendocrinology*,

98, 161–167. https://doi.org/10.1016/j.psyneuen.2018.07.032

Fernández, R., Ramírez, K., Gómez-Gil, E., Cortés-Cortés, J., Mora, M., Aranda, G., Zayas,

E. D., Esteva, I., Almaraz, M. C., Guillamon, A., & Pásaro, E. (2020). Gender-

Affirming Hormone Therapy Modifies the CpG Methylation Pattern of the ESR1

Gene Promoter After Six Months of Treatment in Transmen. *The Journal of Sexual*

Medicine, *17*(9), 1795–1806. https://doi.org/10.1016/j.jsxm.2020.05.027

Foreman, M., Hare, L., York, K., Balakrishnan, K., Sánchez, F. J., Harte, F., Erasmus, J.,

Vilain, E., & Harley, V. R. (2019). Genetic Link Between Gender Dysphoria and Sex

Hormone Signaling. *The Journal of Clinical Endocrinology and Metabolism*, *104*(2),

390–396. https://doi.org/10.1210/jc.2018-01105

Garcia-Falgueras, A., & Swaab, D. F. (2008). A sex difference in the hypothalamic uncinate nucleus: Relationship to gender identity. *Brain*, *131*(12), 3132–3146. https://doi.org/10.1093/brain/awn276

Gennari, L., Masi, L., Merlotti, D., Picariello, L., Falchetti, A., Tanini, A., Mavilia, C., Del Monte, F., Gonnelli, S., Lucani, B., Gennari, C., & Brandi, M. L. (2004). A polymorphic CYP19 TTTA repeat influences aromatase activity and estrogen levels in elderly men: Effects on bone metabolism. *The Journal of Clinical Endocrinology and Metabolism*, *89*(6), 2803–2810. https://doi.org/10.1210/jc.2003-031342

Gennari, L., Merlotti, D., De Paola, V., Calabrò, A., Becherini, L., Martini, G., & Nuti, R. (2005). Estrogen Receptor Gene Polymorphisms and the Genetics of Osteoporosis: A HuGE Review. *American Journal of Epidemiology*, *161*(4), 307–320. https://doi.org/10.1093/aje/kwi055

Greenburg, J., & Gaia, A. (2019). Interpersonal Contact, Stereotype Acceptance, Gender Role Beliefs, Causal Attribution, and Religiosity as Predictors of Attitudes Toward Transgender Individuals. *Psi Chi Journal of Psychological Research*, *24*, 18–32. https://doi.org/10.24839/2325-7342.JN24.1.18

Guillamon, A., Junque, C., & Gómez-Gil, E. (2016). A Review of the Status of Brain Structure Research in Transsexualism. *Archives of Sexual Behavior*, *45*(7), 1615–1648. https://doi.org/10.1007/s10508-016-0768-5

Hall, F. S., Drgonova, J., Jain, S., & Uhl, G. R. (2013). Implications of genome wide association studies for addiction: Are our a priori assumptions all wrong? *Pharmacology & Therapeutics*, *140*(3), 267–279. https://doi.org/10.1016/j.pharmthera.2013.07.006

Hare, L., Bernard, P., Sánchez, F. J., Baird, P. N., Vilain, E., Kennedy, T., & Harley, V. R. (2009). Androgen receptor repeat length polymorphism associated with male-to-

658 female transsexualism. *Biological Psychiatry*, *65*(1), 93–96.

659 https://doi.org/10.1016/j.biopsych.2008.08.033

660 Hellwege, J. N., Keaton, J. M., Giri, A., Gao, X., Edwards, D. R. V., & Edwards, T. L.

661 (2017). Population Stratification in Genetic Association Studies. *Current Protocols in*

662 *Human Genetics*, *95*(1), 1.22.1-1.22.23. https://doi.org/10.1002/cphg.48

663 Henningsson, S., Westberg, L., Nilsson, S., Lundström, B., Ekselius, L., Bodlund, O.,

664 Lindström, E., Hellstrand, M., Rosmond, R., Eriksson, E., & Landén, M. (2005). Sex

665 steroid-related genes and male-to-female transsexualism. *Psychoneuroendocrinology*,

666 *30*(7), 657–664. https://doi.org/10.1016/j.psyneuen.2005.02.006

667 Heylens, G., Cuypere, G. D., Zucker, K. J., Schelfaut, C., Elaut, E., Bossche, H. V., Baere, E.

668 D., & T'Sjoen, G. (2012). Gender Identity Disorder in Twins: A Review of the Case

669 Report Literature. *The Journal of Sexual Medicine*, *9*(3), 751–757.

670 https://doi.org/10.1111/j.1743-6109.2011.02567.x

671 Hönekopp, J., & Watson, S. (2010). Meta-analysis of digit ratio 2D:4D shows greater sex

672 difference in the right hand. *American Journal of Human Biology*, *22*(5), 619–630.

673 https://doi.org/10.1002/ajhb.21054

674 Johnson, E. C., Border, R., Melroy-Greif, W. E., de Leeuw, C. A., Ehringer, M. A., & Keller,

675 M. C. (2017). No Evidence That Schizophrenia Candidate Genes Are More

676 Associated With Schizophrenia Than Noncandidate Genes. *Biological Psychiatry*,

677 *82*(10), 702–708. https://doi.org/10.1016/j.biopsych.2017.06.033

678 Laflamme, N., Nappi, R. E., Drolet, G., Labrie, C., & Rivest, S. (1998). Expression and

679 neuropeptidergic characterization of estrogen receptors (ERalpha and ERbeta)

680 throughout the rat brain: Anatomical evidence of distinct roles of each subtype.

681 *Journal of Neurobiology*, *36*(3), 357–378. https://doi.org/10.1002/(sici)1097-

682 4695(19980905)36:3<357::aid-neu5>3.0.co;2-v

Landén, M., & Innala, S. (2000). Attitudes Toward Transsexualism in a Swedish National

Survey. *Archives of Sexual Behavior*, *29*(4), 375–388.

https://doi.org/10.1023/A:1001970521182

Langdahl, B. L., Løkke, E., Carstens, M., Stenkjær, L. L., & Eriksen, E. F. (2000). A TA

Repeat Polymorphism in the Estrogen Receptor Gene Is Associated with Osteoporotic

Fractures but Polymorphisms in the First Exon and Intron Are Not. *Journal of Bone*

and Mineral Research, *15*(11), 2222–2230.

https://doi.org/10.1359/jbmr.2000.15.11.2222

Levin, R. N., Erickson-Schroth, L., Mak, K., & Edmiston, E. K. (2022). Biological studies of

transgender identity: A critical review. *Journal of Gay & Lesbian Mental Health*,

0(0), 1–30. https://doi.org/10.1080/19359705.2022.2127042

Lombardo, F., Toselli, L., Grassetti, D., Paoli, D., Masciandaro, P., Valentini, F., Lenzi, A.,

& Gandini, L. (2013). Hormone and genetic study in male to female transsexual

patients. *Journal of Endocrinological Investigation*, *36*(8), 550–557.

https://doi.org/10.3275/8813

Luoto, S., & Rantala, M. J. (2018). On estrogenic masculinization of the human brain and

behavior. *Hormones and Behavior*, *97*, 1–2.

https://doi.org/10.1016/j.yhbeh.2017.07.017

Mamikutty, R., Aly, A. S., & Marhazlinda, J. (2021). Selecting Risk of Bias Tools for

Observational Studies for a Systematic Review of Anthropometric Measurements and

Dental Caries among Children. *International Journal of Environmental Research and*

Public Health, *18*(16), Article 16. https://doi.org/10.3390/ijerph18168623

Mazur, T. (2005). Gender dysphoria and gender change in androgen insensitivity or

micropenis. *Archives of Sexual Behavior*, *34*(4), 411–421.

https://doi.org/10.1007/s10508-005-4341-x

McIntyre, M. (2006). The use of digit ratios as markers for perinatal androgen action. *Reproductive Biology and Endocrinology : RB&E*, *4*, 10. https://doi.org/10.1186/1477-7827-4-10

Meerwijk, E. L., & Sevelius, J. M. (2017). Transgender Population Size in the United States: A Meta-Regression of Population-Based Probability Samples. *American Journal of Public Health*, *107*(2), e1–e8. https://doi.org/10.2105/AJPH.2016.303578

Moyer, A. M., Matey, E. T., & Miller, V. M. (2019). Individualized medicine: Sex, hormones, genetics, and adverse drug reactions. *Pharmacology Research & Perspectives*, *7*(6), e00541. https://doi.org/10.1002/prp2.541

Mueller, S. C., Guillamon, A., Zubiaurre-Elorza, L., Junque, C., Gomez-Gil, E., Uribe, C., Khorashad, B. S., Khazai, B., Talaei, A., Habel, U., Votinov, M., Derntl, B., Lanzenberger, R., Seiger, R., Kranz, G. S., Kreukels, B. P. C., Kettenis, P. T. C., Burke, S. M., Lambalk, N. B., … Luders, E. (2021). The Neuroanatomy of Transgender Identity: Mega-Analytic Findings From the ENIGMA Transgender Persons Working Group. *The Journal of Sexual Medicine*, *18*(6), 1122–1129. https://doi.org/10.1016/j.jsxm.2021.03.079

Munafò, M. R., Clark, T. G., & Flint, J. (2004). Assessing publication bias in genetic association studies: Evidence from a recent meta-analysis. *Psychiatry Research*, *129*(1), 39–44. https://doi.org/10.1016/j.psychres.2004.06.011

Nugent, B. M., Wright, C. L., Shetty, A. C., Hodes, G. E., Lenz, K. M., Mahurkar, A., Russo, S. J., Devine, S. E., & McCarthy, M. M. (2015). Brain feminization requires active repression of masculinization via DNA methylation. *Nature Neuroscience*, *18*(5), Article 5. https://doi.org/10.1038/nn.3988

Ogawa, S., Hosoi, T., Shiraki, M., Orimo, H., Emi, M., Muramatsu, M., Ouchi, Y., & Inoue, S. (2000). Association of Estrogen Receptor β Gene Polymorphism with Bone

Mineral Density. *Biochemical and Biophysical Research Communications*, *269*(2),

537–541. https://doi.org/10.1006/bbrc.2000.2285

Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D.,

Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J.,

Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson,

E., McDonald, S., … Moher, D. (2021). The PRISMA 2020 statement: An updated

guideline for reporting systematic reviews. *PLOS Medicine*, *18*(3), e1003583.

https://doi.org/10.1371/journal.pmed.1003583

Pasterski, V., Zucker, K. J., Hindmarsh, P. C., Hughes, I. A., Acerini, C., Spencer, D.,

Neufeld, S., & Hines, M. (2015). Increased Cross-Gender Identification Independent

of Gender Role Behavior in Girls with Congenital Adrenal Hyperplasia: Results from

a Standardized Assessment of 4- to 11-Year-Old Children. *Archives of Sexual*

Behavior, *44*(5), 1363–1375. https://doi.org/10.1007/s10508-014-0385-0

Perrin, J. S., Hervé, P.-Y., Leonard, G., Perron, M., Pike, G. B., Pitiot, A., Richer, L.,

Veillette, S., Pausova, Z., & Paus, T. (2008). Growth of white matter in the adolescent

brain: Role of testosterone and androgen receptor. *The Journal of Neuroscience: The*

Official Journal of the Society for Neuroscience, *28*(38), 9519–9524.

https://doi.org/10.1523/JNEUROSCI.1212-08.2008

Petitti, D. B. (2001). Approaches to heterogeneity in meta-analysis. *Statistics in Medicine*,

20(23), 3625–3633. https://doi.org/10.1002/sim.1091

Plomin, R., DeFries, J. C., Knopik, V. S., & Neiderhiser, J. M. (2016). Top 10 Replicated

Findings from Behavioral Genetics. *Perspectives on Psychological Science: A*

Journal of the Association for Psychological Science, *11*(1), 3–23.

https://doi.org/10.1177/1745691615617439

Polderman, T. J. C., Kreukels, B. P. C., Irwig, M. S., Beach, L., Chan, Y.-M., Derks, E. M.,

Esteva, I., Ehrenfeld, J., Heijer, M. D., Posthuma, D., Raynor, L., Tishelman, A.,

Davis, L. K., & International Gender Diversity Genomics Consortium. (2018). The

Biological Contributions to Gender Identity and Gender Diversity: Bringing Data to

the Table. *Behavior Genetics*, *48*(2), 95–108. https://doi.org/10.1007/s10519-018-

9889-z

Puts, D., & Motta-Mena, N. V. (2018). Is human brain masculinization estrogen receptor-

mediated? Reply to Luoto and Rantala. *Hormones and Behavior*, *97*, 3–4.

https://doi.org/10.1016/j.yhbeh.2017.07.018

Rad, M. S., Shackleford, C., Lee, K. A., Jassin, K., & Ginges, J. (2019). Folk theories of

gender and anti-transgender attitudes: Gender differences and policy preferences.

PLOS ONE, *14*(12), e0226967. https://doi.org/10.1371/journal.pone.0226967

Rajkovic, A., Cirino, A. L., Berro, T., Koeller, D. R., & Zayhowski, K. (2022). Transgender

and gender-diverse (TGD) individuals' perspectives on research seeking genetic

variants associated with TGD identities: A qualitative study. *Journal of Community

Genetics*, *13*(1), 31–48. https://doi.org/10.1007/s12687-021-00554-z

Rametti, G., Carrillo, B., Gómez-Gil, E., Junque, C., Zubiarre-Elorza, L., Segovia, S.,

Gomez, Á., & Guillamon, A. (2011). The microstructure of white matter in male to

female transsexuals before cross-sex hormonal treatment. A DTI study. *Journal of

Psychiatric Research*, *45*(7), 949–954.

https://doi.org/10.1016/j.jpsychires.2010.11.007

Ramírez, K. del V., Fernández, R., Delgado-Zayas, E., Gómez-Gil, E., Esteva, I., Guillamon,

A., & Pásaro, E. (2021). Implications of the Estrogen Receptor Coactivators SRC1

and SRC2 in the Biological Basis of Gender Incongruence. *Sexual Medicine*, *9*(3).

https://doi.org/10.1016/j.esxm.2021.100368

Ramirez, K., Fernández, R., Collet, S., Kiyar, M., Delgado-Zayas, E., Gómez-Gil, E., Van

Den Eynde, T., T'Sjoen, G., Guillamon, A., Mueller, S. C., & Pásaro, E. (2021).

Epigenetics Is Implicated in the Basis of Gender Incongruence: An Epigenome-Wide

Association Analysis. *Frontiers in Neuroscience*, *15*, 701017.

https://doi.org/10.3389/fnins.2021.701017

Raznahan, A., Lee, Y., Stidd, R., Long, R., Greenstein, D., Clasen, L., Addington, A.,

Gogtay, N., Rapoport, J. L., & Giedd, J. N. (2010). Longitudinally mapping the

influence of sex and androgen signaling on the dynamics of human cortical

maturation in adolescence. *Proceedings of the National Academy of Sciences*,

107(39), 16988–16993. https://doi.org/10.1073/pnas.1006025107

Rosenthal, S. M. (2021). Challenges in the care of transgender and gender-diverse youth: An

endocrinologist's view. *Nature Reviews. Endocrinology*, *17*(10), 581–591.

https://doi.org/10.1038/s41574-021-00535-9

Scariano, J. K., Simplicio, S. G., Montoya, G. D., Garry, P. J., & Baumgartner, R. N. (2004).

Estrogen Receptor β Dinucleotide (CA) Repeat Polymorphism is Significantly

Associated with Bone Mineral Density in Postmenopausal Women. *Calcified Tissue

International*, *74*(6), 501–508. https://doi.org/10.1007/s00223-003-0170-x

Seneviratne, S. N., Jayarajah, U., Gunawardana, S., Samarasinghe, M., & de Silva, S. (2021).

Gender-role behaviour and gender identity in girls with classical congenital adrenal

hyperplasia. *BMC Pediatrics*, *21*(1), 262. https://doi.org/10.1186/s12887-021-02742-9

Sharp, L., Cardy, A. H., Cotton, S. C., & Little, J. (2004). CYP17 Gene Polymorphisms:

Prevalence and Associations with Hormone Levels and Related Factors. A HuGE

Review. *American Journal of Epidemiology*, *160*(8), 729–740.

https://doi.org/10.1093/aje/kwh287

Shepherd, R., Bretherton, I., Pang, K., Mansell, T., Czajko, A., Kim, B., Vlahos, A., Zajac, J.

D., Saffery, R., Cheung, A., & Novakovic, B. (2022). Gender-affirming hormone

therapy induces specific DNA methylation changes in blood. *Clinical Epigenetics*,

14(1), 24. https://doi.org/10.1186/s13148-022-01236-4

Siegmann, E.-M., Müller, T., Dziadeck, I., Mühle, C., Lenz, B., & Kornhuber, J. (2020).

Digit ratio (2D:4D) and transgender identity: New original data and a meta-analysis.

Scientific Reports, *10*(1), Article 1. https://doi.org/10.1038/s41598-020-72486-6

Song, F., Sheldon, T. A., Sutton, A. J., Abrams, K. R., & Jones, D. R. (2001). Methods for

Exploring Heterogeneity in Meta-Analysis. *Evaluation & the Health Professions*,

24(2), 126–151. https://doi.org/10.1177/016327870102400203

Tan, G. C.-Y., Chu, C., Lee, Y. T., Tan, C. C. K., Ashburner, J., Wood, N. W., &

Frackowiak, R. SJ. (2020). The influence of microsatellite polymorphisms in sex

steroid receptor genes ESR1, ESR2 and AR on sex differences in brain structure.

NeuroImage, *221*, 117087. https://doi.org/10.1016/j.neuroimage.2020.117087

Theisen, J. G., Sundaram, V., Filchak, M. S., Chorich, L. P., Sullivan, M. E., Knight, J., Kim,

H.-G., & Layman, L. C. (2019). The Use of Whole Exome Sequencing in a Cohort of

Transgender Individuals to Identify Rare Genetic Variants. *Scientific Reports*, *9*(1),

20099. https://doi.org/10.1038/s41598-019-53500-y

Ujike, H., Otani, K., Nakatsuka, M., Ishii, K., Sasaki, A., Oishi, T., Sato, T., Okahisa, Y.,

Matsumoto, Y., Namba, Y., Kimata, Y., & Kuroda, S. (2009). Association study of

gender identity disorder and sex hormone-related genes. *Progress in Neuro-*

Psychopharmacology and Biological Psychiatry, *33*(7), 1241–1244.

https://doi.org/10.1016/j.pnpbp.2009.07.008

Vaillancourt, K. L., Dinsdale, N. L., & Hurd, P. L. (2012). Estrogen receptor 1 promoter

polymorphism and digit ratio in men. *American Journal of Human Biology*, *24*(5),

682–689. https://doi.org/10.1002/ajhb.22297

Viechtbauer, W. (2010). Conducting Meta-Analyses in R with the metafor Package. *Journal*

of Statistical Software, *36*, 1–48. https://doi.org/10.18637/jss.v036.i03

Wan, X., Wang, W., Liu, J., & Tong, T. (2014). Estimating the sample mean and standard

deviation from the sample size, median, range and/or interquartile range. *BMC*

Medical Research Methodology, *14*(1), 135. https://doi.org/10.1186/1471-2288-14-

135

Wells, G., Shea, B., & O'Connell, J. (2014). The Newcastle-Ottawa Scale (NOS) for

Assessing The Quality of Nonrandomised Studies in Meta-analyses. *Ottawa Health*

Research Institute Web Site, *7*.

Weng, H., Li, S., Huang, J.-Y., He, Z.-Q., Meng, X.-Y., Cao, Y., Fang, C., & Zeng, X.-T.

(2017). Androgen receptor gene polymorphisms and risk of prostate cancer: A meta-

analysis. *Scientific Reports*, *7*(1), 40554. https://doi.org/10.1038/srep40554

Westberg, L., Baghaei, F., Rosmond, R., Hellstrand, M., Landén, M., Jansson, M., Holm, G.,

Björntorp, P., & Eriksson, E. (2001). Polymorphisms of the Androgen Receptor Gene

and the Estrogen Receptor β Gene Are Associated with Androgen Levels in Women1.

The Journal of Clinical Endocrinology & Metabolism, *86*(6), 2562–2568.

https://doi.org/10.1210/jcem.86.6.7614

Woodford, M. R., Silverschanz, P., Swank, E., Scherrer, K. S., & Raiz, L. (2012). Predictors

of Heterosexual College Students' Attitudes Toward LGBT People. *Journal of LGBT*

Youth, *9*(4), 297–320. https://doi.org/10.1080/19361653.2012.716697

Xiao, F., Lan, A., Lin, Z., Song, J., Zhang, Y., Li, J., Gu, K., Lv, B., Zhao, D., Zeng, S.,

Zhang, R., Zhao, W., Pan, Z., Deng, X., & Yang, X. (2016). Impact of CAG repeat

length in the androgen receptor gene on male infertility—A meta-analysis.

Reproductive Biomedicine Online, *33*(1), 39–49.

https://doi.org/10.1016/j.rbmo.2016.03.012

Yore, M. A., Im, D., Webb, L. K., Zhao, Y., Chadwick, J. G., Molenda-Figueira, H. A.,

Haidacher, S. J., Denner, L., & Tetel, M. J. (2010). Steroid receptor coactivator-2

expression in brain and physical associations with steroid receptors. *Neuroscience*,

169(3), 1017–1028. https://doi.org/10.1016/j.neuroscience.2010.05.053

Zhou, J.-N., Hofman, M. A., Gooren, L. J. G., & Swaab, D. F. (1995). A sex difference in the

human brain and its relation to transsexuality. *Nature*, *378*(6552), Article 6552.

https://doi.org/10.1038/378068a0

Zitzmann, M., & Nieschlag, E. (2003). The CAG repeat polymorphism within the androgen

receptor gene and maleness1. *International Journal of Andrology*, *26*(2), 76–83.

https://doi.org/10.1046/j.1365-2605.2003.00393.x

Zuloaga, D. G., Puts, D. A., Jordan, C. L., & Breedlove, S. M. (2008). The role of androgen

receptors in the masculinization of brain and behavior: What we've learned from the

testicular feminization mutation. *Hormones and Behavior*, *53*(5), 613–626.

https://doi.org/10.1016/j.yhbeh.2008.01.013

Records identified from
databases (n = 568)
Medline = 280
Embase = 288

Duplicate records
automatically removed with
EndNote (n = 129)

Records screened
(n = 439)

Records excluded with
Rayyan (n = 419)

Reports sought for retrieval
(n = 20)

Reports not retrieved
(n = 0)

Reports assessed for eligibility
(n = 20)

Reports excluded (n = 14):
- Duplicate study not identified previously (n = 1)
- Same cohort used in later study (n = 2)
- Previous cohort articles (n = 3)
- Included < 18 years (n = 1)
- Study design (n = 1)
- Wrong patient population (n = 1)
- Short duration (n = 1)
- Wrong intervention (n = 1)
- Not relevant outcomes (n = 1)
- Wrong dosing (n = 1)
- Wrong patient population (n = 1)

Studies included in review
(n = 6)

a

Study	Transgender Women	Cisgender Men	Odds Ratio [95% CI]

b

Study	Transgender Women			Cisgender Men			SMD [95% CI]
	Mean	SD	N	Mean	SD	N	