Mass-immigration shapes the antibiotic resistome of wastewater treatment plants

Lanping Zhang[1,2,3], Bob Adyari[1,2,3,4], Liyuan Hou[5,6], Xiaoyong Yang[7],

Mahmoud Gad[8], Yuwen Wang[1,9], Cong Ma[10], Qian Sun[1,2,3], Qiang Tang[11],

Yifeng Zhang[12], Chang-Ping Yu[1], Anyi Hu[1,2,3*]

1. CAS Key Laboratory of Urban pollutant Conversion, Institute of Urban

Environment, Chinese Academy of Sciences, Xiamen 361021, China

2. University of Chinese Academy of Sciences, Beijing 100049, China

3. Fujian Key Laboratory of Watershed Ecology, Institute of Urban Environment,

Chinese Academy of Sciences, Xiamen 361021, China

4. Department of Environmental Engineering, Universitas Pertamina, Jakarta 12220,

Indonesia

5. Department of Civil and Environmental Engineering, Utah state university, Utah

UT 84322, USA

6. Utah Water Research Laboratory, Utah State University, Logan, Utah 84322,

USA

7. School of Environmental and Material Engineering, Yantai University, 30

Qingquan Road, Yantai 264005, China

8. Water Pollution Research Department, National Research Centre, Giza 12622,

Egypt

9. Marine biology Institute, Shantou University, Shantou 515063, China

10. Xiamen Municipal Environmental Technology Co., Ltd., Xiamen 361001, China

11. Department of Environmental Science and Technology, University of Science and

Technology of China, Hefei, 230026, China

12. Department of Environmental and Resource Engineering, Technical University of

Denmark, DK-2800 Lyngby, Denmark

*Correspondence to:

Dr. Anyi Hu; Email: ayhu@iue.ac.cn; Tel: 86-592-6190582.

Abstract

Wastewater treatment plants (WWTPs) are the hotspots for the spread of

antibiotic resistance genes (ARGs) into the environment. Nevertheless, a

comprehensive assessment of the city-level and short-term daily variations of ARG

surveillance is still lacking in WWTPs. Here, 285 ARGs and ten mobile gene

elements (MGEs) were monitored in seven WWTPs in Xiamen via high-throughput

qPCR (HT-qPCR) for seven days. The average daily load of ARGs to WWTPs was

about 1.21×10^{20} copies/d, and a total of 1.44×10^{18} copies/d was discharged to the

environment across the entire city. Interestingly, no daily variations were observed in

ARG richness, abundance, and community composition. Stochastic processes were

the main force determining the assembly of ARG communities, with their relative

importance ranked in the order of influent (INF) > effluent (EFF) > activated sludge

(AS). Further analyses indicated that bacteria and ARGs from upstream treatment

units played an increasingly dominant role in shaping ARG communities in AS and

EFF, respectively, suggesting the importance of mass-immigration of bacteria and

ARGs from the source on ARG transport in wastewater treatment units. This

emphasizes the need to revise the way we mitigate ARG contamination but focus on

the source of ARGs in urban wastewater.

Keywords: Wastewater treatment plant; Antibiotic resistance genes; Mass

immigration; Assembly mechanisms; Daily variation

1. Introduction

Extensive consumption of antibiotics in humans and livestock exerts a major selective pressure on the environmental microbiome, contributing to the proliferation and spread of antibiotic resistance bacteria (ARB) and antibiotic resistance genes (ARGs) [1]. Therefore, ARB and ARGs have become major global health threats and concerns in environmental systems [2,3]. One Health framework, aiming to understand the mechanism of ARB movement across humans, animals, and the environment, can help guide antibiotic resistance management. Reliable surveillance data lays the foundation for the One Health framework [4]. Previous cross-regional investigations reveal that ARGs are widespread in both highly human-disturbed (e.g., urban city) and human-inaccessible areas (e.g., pristine plateau) [5–9]. Human activities can promote ARG transfer between various environmental interfaces.

Nowadays, WWTPs are widely recognized as the hotspots for ARG proliferation and dissemination [10]. Previous studies showed that the biological treatment processes (e.g., activated sludge) of WWTPs contained a high density of bacteria and various kinds of contaminants (e.g., heavy metals, antibiotics, etc.). These contaminants not only facilitated the proliferation of ARB [11] but also promoted the exchange of ARGs among bacteria [12–18]. Although WWTPs had a relatively high removal efficiency of ARGs for 1–2 log reduction, the effluent still harbored more than 10^7 copies/mL ARGs [19,20]. The release or recycling of effluents could lead to increasing ARGs abundance and shifting in ARGs community composition in the receiving environments [21–25]. In addition, numerous studies have demonstrated that ARG profiles in WWTP varied at different time scales. For example, at the seasonal scale, the diversity and composition of ARGs were significantly different between summer and winter in eleven WWTPs in China [20]. And the temporal changes of ARG types in effluents were mainly related to the prevalent antibiotics [19]. At the diurnal scale, the variations of ARGs were related to the dynamics of

wastewater flow [26,27]. However, the daily variation of ARGs in WWTPs at the city

level is still limited. This is important for setting suitable sampling frequencies and

monitoring targets to facilitate comparison within and among individual WWTPs. In

addition, obtaining this information will help understand the assembly mechanisms

underlying ARG spread within a short time, contributing to establishing suitable

wastewater-based surveillance approaches in cities.

Stochastic (i.e., dispersal and drift) and deterministic processes (i.e., bio-

interaction and environmental selection) collectively contribute to the community

assembly [28–30]. However, after decades of debate, the role of stochastic processes

in structuring communities has been increasingly appreciated. For instance, the

stochastic processes played a central role in structuring ARG communities in urban

lakes (e.g., Baiyang Lake and Tai Lake). However, the difference in the relative

importance of stochastic processes in Baiyang Lake and Tai Lake was correlated with

environmental selection [31]. In addition, Hou and colleagues showed that the

stochastic processes played a dominant role in shaping ARG community assembly in

urban water bodies under anthropogenic influences [32,33]. And they suggested that

the dominant role of stochastic processes may indicate the contribution of HGT to the

spread of ARGs in different urban environments. Although many studies explored the

influence of various factors (e.g., nutrients, chemical contaminants, operational

parameters, etc.) on ARG profiles in WWTPs or bioreactors [17,34–36], the

contribution of stochastic processes in ARG community assembly in the different

treatment processes of WWTPs is still limited. Understanding the assembly

mechanism of microbial communities including ARGs and their hosts is a promising

way to provide information for predicting their transport and spread.

Numerous studies reported that the combined effect of physiochemical parameters

and biological interaction shapes the ARG profiles [37–39]. Notably, several recent

studies have demonstrated that mass immigration via wastewater stream strongly

determined the bacterial communities in active sludge [40–42]. However, whether

mass immigration (i.e., stochastic processes) plays a role in the ARG community

assembly across the wastewater treatment processes is not known. Since influent

wastewater flows across different WWTP treatment units, the mass immigration of

microorganisms along the flow may shape the microbial and ARG communities in the

downstream processes such as the activated sludge and effluents [40].

In this study, we employed HT-qPCR to investigate the daily variation of 285

ARGs and 10 MGEs in INF, AS and EFF of seven WWTPs in Xiamen, China for

seven days. We aim to: 1) characterize the daily variations of the abundance, diversity,

and composition of ARGs in WWTPs across the whole city; 2) explore the

community assembly mechanisms of ARGs and identify the key biotic and abiotic

factors shaping the ARG profiles in different wastewater treatment units; and 3) assess

the daily ARG mass loads to WWTPs and ARG mass loads in receiving environment

at the city level, followed by potential health risk assessment. We assume that

stochastic processes will have a higher contribution to the ARG community assembly

in INF than in AS and EFF because the latter two will undergo specific environmental

pressure (i.e., operation parameters). Among various factors (i.e., biotic and abiotic

factors), the immigration of ARGs and microbial communities from upstream (e.g.,

INF) will affect the ARG profiles in the downstream processes (i.e., AS and EFF).

The result of this study will extend our understanding of ARG dynamics as well as the

underlying assembly mechanisms in WWTPs, providing fundamental information for

the surveillance and control of WWTP ARGs at a city-level perspective.

2. Methods and materials

2.1 Study area and sample collection

Xiamen is a typical subtropical monsoon city on the southeast coast of China

(117°4′E-118°21′E, 24°24′N-24°54′N) with a small range of annual temperature

variance (seasonal average temperature 13.7–27.9°C). It covers an area of 1699 km^2

and has a population of 3.92 million in 2016 [43]. Impressively, the urban land almost

doubled to 334 km^2 and the total amount of municipal wastewater increased from

around 210 million tons to 320 million tons during the past 10 years (Xiamen

Environmental Protection Bureau, 2007–2016). 92% of the municipal wastewater in

Xiamen was treated by seven WWTPs. Among these WWTPs, six WWTPs (listed as

W2~W7) employ the activated sludge technology, while one (i.e., W1) uses a

biological aerated filter treatment process. Detailed information on these WWTPs was

shown in Table S1. More information about Xiamen city and residents representing

the socioeconomic development such as the urbanization rate, per capita disposable

income (PCDI), and per capita consumption expenditure was obtained from the

Xiamen Bureau of Statistics [44].

The 24-hr composite samples of INF, AS, and EFF were collected from seven

WWTPs by using automatic samplers (Hachi, Sigma SD900) for 7 days (from Feb.

28th to March 6th, 2016). But there was no sampling event on March 5th, 2016 because

the staff had the day off. From all WWTPs, six were equipped with one AS tank each

and one (W5) was equipped with two tanks. For the first two days, we collected AS

samples from the two AS tanks of W5 separately, and one AS composite sample for

the subsequent days. Due to the limitations of sampling conditions, a total of 49 INF

samples, 49 EFF samples, and 48 AS samples were finally collected. Among these

samples, there were 46 pairs of INF-AS-EFF samples. All samples were kept on ice

and shipped to the lab within 4 h. For microbial community analysis, 100~300 mL of

INF and 800~1000 mL of EFF were filtered through the 0.2 μm pore size Sterivex-GP

filter (Millipore, Billerica, MA, USA), while AS samples were concentrated by

centrifuging at 8,000 rpm for 15 min. All microbial concentrated samples were stored

at -80°C before DNA extraction. The concentration of pharmaceuticals and personal

care products (PPCPs) was measured following the protocol of EPA 1694 with slight

modifications [45]. Here, we classified PPCPs into antibiotics, non-steroidal anti-

inflammatory drugs (NSAIDs), stimulants, and other PPCPs, and cited the mass loads

of information on each PPCP type from our previous study [45]. All detected PPCPs

were listed in **Table S2** and PPCP mass load per capita of each classification were

supplied in **Table S3**. The water quality parameters, including chemical oxygen

demand (COD), 5-day biological oxygen demand (BOD$_5$), total phosphorous (TP),

total nitrogen (TN), ammonia nitrogen (NH$_4^+$-N), and suspended solids (SS) of INF

and EFF, were provided by the wastewater treatment plants by using national standard

methods [46]. Activated sludge performance on specific removal rates of nutrients

(TN, TP, and NH$_4^+$-N) and organic matters (indicated as chemical oxygen demand;

COD) was calculated as the following equation [47]:

$$Removal\ rate = (C_{inf} - C_{eff})/C_{inf}$$

where C_{inf} and C_{eff} represent the concentration of each chemical in the INF and EFF

(mg/L), respectively.

2.2 DNA extraction, 16S rRNA gene amplicon sequencing, and data processing

The genomic DNA of the 146 samples was extracted by using FastDNA SPIN Kit

for Soil (Qbiogene-MP Biomedicals, Irvine, CA, USA) according to the

manufacturer's protocol with modifications [43]. The 16S rRNA amplicon sequencing

was conducted to characterize the prokaryotic bacterial communities. The 16S rRNA

genes in the hypervariable V4-V5 region were amplified with a universal primer set

515F and 907R [48]. The PCR products were then purified and sequenced on Illumina

HiSeq 2500 platform (Illumina Inc., San Diego, CA, USA) via a paired-end approach

(2 × 250 bp). The raw sequence data can be downloaded from the NCBI short reads

archive database under the BioProject number of PRJNA545043.

Raw 16S rRNA paired-end reads were trimmed and assembled using the LotuS

pipeline [43,49]. Only high-quality reads that meet the following criteria were

retained for subsequent analysis: (1) an average sequence quality > 27; (2) a sequence

length > 170 bp; (3) no ambiguous bases; (4) no mismatches to the barcodes or

primers; and (□) a homopolymer run < 8 bp. The reads were then checked for

chimera and clustered into operational taxonomic units (OTUs) with a threshold of 97%

similarity by using UPARSE [50].

2.3 HT-qPCR analysis of 285 ARGs and 10 MGEs

A total of 285 ARGs, 10 MGEs, and one 16S rRNA gene were determined on a

Warfergen SmartChip real-time PCR system (Wafergen Bio Inc., USA) with the

SYBR Green I$^{®}$ chemistry. The targeted ARGs involved the genes that were resistant

to all major classes of antibiotics, while those MGEs included 8 transposase and 2

integron genes. HT-qPCR was carried out in a volume of 100 nL reaction system and

was under the following thermal cycle condition: initial denaturing at 95°C for 10 min;

followed by 40 cycles of denaturation at 95°C for 30 s and annealing at 60°C for 30 s

[33,51,52]. The melting curve analyses were automatically generated by SmartChip

qPCR software (2.7.0.1). Only the results that meet the following conditions were

obtained for statistical analysis: (1) a threshold cycle (Ct) $\leq$ 31; (2) more than two

replicates having been amplified were retained as positive; (3) reaction with one peak;

(4) amplification efficiency within 90–110%. The relative copy number of each gene

can be calculated using:

$$\text{Relative gene copy number}_{gene} = 10^{(31-\text{Ct})/(10/3)}$$

208   Relative abundance (copies per 16S rRNA gene) and absolute abundance (copies

209 per mL) of ARGs and MGEs was calculated by the following equation:

$$\text{Relative abundance}_{gene} = \frac{\text{Relative gene copy number}_{gene}}{\text{Relative gene copy number}_{16S\ rRNA}}$$

$$\text{Absolute abundance}_{gene}$$
$$= \text{Relative abundance}_{gene}$$
$$\times \text{absolute abundance of 16S rRNA genes}$$

The absolute abundance of the 16S rRNA gene was measured on LightCycler®

Roche 480 Real-time PCR instrument (Roche Inc., Basel, Switzerland). A total of 20

µL reaction system consisted of 10 µL LightCycler® 480 SYBR Green I Master Mix

(2 ×) (Applied Bio-systems, CA, USA), 0.4 μM of 16S rRNA primer, 1 μg/L bovine

serum albumin (Sigma, Steinheim, Germany) and 20 ng of template DNA. The

standard curve was processed by 10-fold serially diluted 16S rRNA gene incorporated

plasmids with insert length 61 bp bases. The PCR reactions were performed as

follows: initial denaturation at 95°C for 3 min, followed by 40 cycles of denaturation

at 95°C for 15 s and annealing at 60°C for 1 min and extension at 72°C for 15 s. All

qPCR reactions were conducted in triplicate for each sample and negative controls

were included.

2.4 Daily ARG and MGE mass loads

Based on the total amount of wastewater and the served population, target gene

(i.e., ARGs and MGEs) daily mass loads in different WWTPs across the whole city

and target gene mass loads per inhabitant are calculated at the city level. ARG or

MGE mass loads in different treatment units were following the equations:

Daily target gene loads (copoies/d)

$$= \frac{\sum \text{absolute abundance of total target gene } \left(\frac{\text{copies}}{\text{mL}}\right) \text{in the target unit} \times \text{ Q (mL/d)}}{7 \text{ (d)}}$$

Daily target gene loads per capita (copies/(d · capita))

$$= \frac{\text{Daily target gene loads (copoies/d)}}{\text{population size}}$$

Where Q is the flow rate (mL/d) of the corresponding WWTP. Daily mass loads of

ARGs and MGEs per capita in INF and EFF represent the citizen's daily discharge of

ARGs and MGEs to WWTP and potential discharge loads to the environment or

release to humans, respectively.

2.5 Null model analysis

The null model was utilized to determine the relative contribution of stochastic

and deterministic processes to the assembly of ARG and bacterial communities [33].

The principle of the null model can be recognized as follows: the expected similarity

of null expected communities was calculated by 1000 random shuffles based on the

raw community data. The richness and proportion of species occupancy of the

expected community were consistent with the observed community in this step. Then,

we calculated the value of the relative importance of the stochastic processes by:

$$\text{Stochasticity } (\%) = 1 - (S_{obs} - E_{exp})/S_{obs}$$

where S_{obs} means the observed similarity of the original communities, and E_{exp} means

the average similarity of null expected communities.

2.6 SourceTracker analysis

The relative contribution of communities (ARGs or OTUs) from different source

habitats (i.e., INF, AS, EFF and unknown source) to sink habitats was assessed by

using SourceTracker v2.0 with default parameter settings based on previous studies

[53–55]. For the sink of AS, the source includes INF and the unknown source, while

the sources became INF, AS and the unknown source for the sink of EFF. The

unknown source represents the indigenous communities or those from uninvestigated

environments. For each sample, the predicted proportion for each of the potential

sources was run 5 times to calculate the average proportion. The accuracy of

SourceTracker prediction and corresponding standard deviation (SD) within each

habitat were evaluated by the consistency between the predicted source and the

original habitat. The ARGs and OTUs with an occurrence of less than 1% were

discharged during this analysis.

2.7 Statistical analysis

Daily variations of ARG richness and abundance for seven days were assessed by

the coefficient of variance (CV) and were calculated as the following formula: CV =

SD/the mean value of ARG richness or abundance. The time-dependent ARG and

bacterial community structures were analyzed by using a time decay model. The time

decay model takes the time interval matrices between samples as the independent

variable, and the β-diversity matrices based on the Bray-Curtis distance between

samples as the dependent variable. The variables were fitted with a linear model and

the obtained slope and R^2 were used to evaluate the collinear relationship.

The β-diversity patterns of ARG and bacterial communities were revealed by

nonmetric multidimensional scaling (NMDS) analysis based on Bray-Curtis distance

matrices. The significance of the differences within ARG and bacterial communities

among wastewater treatment units or WWTPs were performed by permutational

multivariate analysis of variance (Adonis) and analysis of similarity (ANOSIM).

VPA was used to distinguish the relative effects of biotic, abiotic factors, and

MGEs on the variance of β-diversity of ARG communities in each treatment process

[48]. The biotic factors included the first two NMDS axes of the bacterial

communities and the corresponding source proportions of ARG and bacterial

communities predicted by SourceTracker. Here, we only included the source

proportions in the analysis of ARG communities from AS and EFF. The abiotic

factors included physicochemical parameters of INF and EFF and the nutrient

removal rate (i.e., TN, TP, NH_4^+-N, and COD) of WWTPs. Here, MGEs were

included as an independent factor because they were commonly recognized as an

important driver causing the HGT of ARGs. Prior to VPA, the multicollinearity

between variables in each factor group was assessed by the variation inflation factor

(VIF), using "vifstep" function from the R package *usdm* [56]. The variables with VIF

< 10 were retained. To quantify the relative importance of an individual element of

factors, HP analysis was performed via the "rdacca.hp" function from the R package

rdacca.hp [57].

Statistical analysis and visualization were conducted using R 4.1.0 with the R

package *phyloseq* [58], *vegan* [59], and *ggplot2* [60].

3. Results

3.1 A full profile of ARGs in WWTPs across the whole city

A total of 208 ARG subtypes were detected in INF, AS, and EFF of seven WWTPs of Xiamen, corresponding to 8 major ARG types (i.e., aminoglycoside, beta-lactam, chloramphenicol, MLSB, multidrug, sulfonamide, tetracycline, and vancomycin resistance genes). The number of ARG subtypes in different treatment units decreased as follows: INF (94~141) > EFF (31~115) > AS (44~98) (**Fig. S1A**). 156 of 208 ARG subtypes (75%) were shared by three habitats (i.e., INF, EFF, and AS) (**Fig. S1B**). The shared ARGs mainly belonged to beta-lactam (43 ARG subtypes), multidrug (33), aminoglycoside (31), and MLSB (23) resistance genes.

The results indicated that the relative abundance of ARGs in INF was the highest (average 0.56 ± 0.22 copies/16S rRNA gene), and then decreased more than 50% in AS (0.20 ± 0.09 copies/16S rRNA gene). But in EFF this value increased again to 0.33 ± 0.16 copies/16S rRNA gene (**Fig. 1A**). Being inconsistent with this trend, AS had the highest absolute abundance of ARGs (average 4.14×10^9 copies/mL), followed by INF (1.34×10^8 copies/mL) and then EFF (2.40×10^6 copies/mL) (**Fig. 1B**). According to the absolute abundance of ARGs, aminoglycoside (ranging from 2.70×10^4 to 5.36×10^9 copies/mL), multidrug (3.49×10^4~4.82×10^9 copies/mL), MLSB (5.81×10^3~9.03×10^8 copies/mL) and beta-lactam (9.90×10^3~3.06×10^9 copies/mL) resistance genes were the top 4 dominant ARG types.

As for MGEs, the relative abundance in INF was the highest value (average 0.092 ± 0.05 copies/16S), followed by EFF (0.055 ± 0.07 copies/16S) and AS (0.016 ± 0.01 copies/16S) (**Fig. 1A**), while the absolute abundance is in the following order: AS (average 3.81×10^8 copies/mL) > INF (2.26×10^7 copies/mL) > EFF (4.10×10^5 copies/mL) (**Fig. 1B**). Notably, among MGE subtypes, the relative abundance of integrons in EFF increased to a level comparable to that in INF. Spearman correlation analysis indicated that the relative abundances of MGEs and ARGs were significantly

correlated with each other in both INF and EFF (r ≥ 0.50, $P < 0.001$), but not in AS

(**Fig. S2**). In addition, the relative abundance of transposases correlated significantly

with the majority of ARG types in both INF and EFF ($P < 0.05$). Interestingly, the

relationship between integrons and ARG types was stronger in INF than in EFF (**Fig.**

S3).

3.2 Daily variations of ARG abundance, diversity, and composition in different

treatment units

NMDS ordination analysis indicated that both ARGs and bacterial communities

tended to cluster based on the sample habitats (i.e., treatment units) (**Fig. 2A & C**).

The treatment unit ($R^2 = 0.21$, $P < 0.001$) and WWTPs ($R^2 = 0.25$, $P < 0.001$) showed

similar effects on bacterial community compositions of WWTPs based on Adonis

tests. However, the overall variance of the ARG communities driven by the treatment

unit ($R^2 = 0.35$, $P < 0.001$) was much higher than that of WWTPs ($R^2 = 0.14$, $P <$

0.001). Significance differences have been observed in both ARG and bacterial

communities between EFF and INF (ARGs: $R^2 = 0.19$, $P < 0.01$; bacteria: $R^2 = 0.14$,

$P < 0.01$), EFF and AS (ARGs: $R^2 = 0.20$, $P < 0.01$; bacteria: $R^2 = 0.11$, $P < 0.01$), as

well as INF and AS (ARGs: $R^2 = 0.43$, $P < 0.01$; bacteria: $R^2 = 0.26$, $P < 0.01$).

However, the differences in both ARG and bacterial communities between EFF and

INF, and EFF and AS were much smaller than those between INF and AS. This

indicated the significant shifts of ARG and bacterial communities from INF to AS. In

addition, bacteria and ARG communities showed pronounced spatial differences

across the treatment units, especially in AS (ARGs: $R^2 = 0.66$, $P < 0.001$; bacteria: R^2

$= 0.78$, $P < 0.001$) (**Fig. S4**). However, Procrustes analysis revealed a similar and

significant β-diversity pattern between ARG and bacterial communities within each

treatment process (**Fig. S5**).

The daily variation of the ARG communities among different treatment units

during the sampling period was presented in **Fig. 3**. The daily variation of ARG

richness showed statistically significant differences between INF and AS, and INF

and EFF (Wilcoxon test; $P < 0.05$). The relative abundance of ARGs exhibited much

smaller daily fluctuations compared to the absolute abundance based on the CV

analysis (**Fig. 3A**). Among different treatment units, AS had the greatest variation in

ARG richness over time but the smallest variation in ARG relative abundance

and OTU richness. Based on the time decay model, we found how the dissimilarity of

ARG community composition changes with time. No linear association between the

dissimilarity of ARG community composition and time has been observed. However,

the dissimilarity of the bacterial community composition showed a strong positive

correlation with time (**Fig. 3B**). This indicated that the daily variation of the ARG

community within the same treatment unit was stable.

3.3 Immigration of ARGs across different treatment units

SourceTracker analysis conducted the prediction of the source habitats (i.e., the

treatment units) for ARG communities in 78% (108/138) samples and for bacterial

communities in 91% (126/138) samples successfully (**Figs. S6A and S7A**).

Specifically, as for ARG communities, 89% samples (41/46), 89% samples (41/46),

and 57% samples (26/46) were predicted to be from INF, AS, and EFF with a

probability of 73%±15%, 79%±14%, and 62%±16%, respectively. This supported the

findings of NMDS ordination that the samples tended to cluster by their habitats (**Fig.**

2A & C), suggesting that each habitat had unique characteristics of ARG communities.

However, as the wastewater flows between different treatment units, the immigration

process of the upstream communities (i.e., INF) contributed to downstream

communities (e.g., AS and EFF). For instance, the contribution of INF to the ARG

community downstream (i.e., AS and EFF) increased greatly along with treatment

units from 25.5% in AS to 49.6% in EFF. AS also contributed to 24.3% of the ARG

community in EFF (**Fig. S6B**). As a comparison, the contribution of INF to the

bacterial community in AS was 56.1%, while INF and AS had similar contributions

(~33%) to the bacterial community in EFF (**Fig. S7B**).

3.4 Assembly mechanisms and controlling factors underlying ARGs in WWTPs

The null model analysis showed that stochastic processes played a dominant role

in shaping the ARG communities in WWTPs (the average stochastic ratio > 50%),

and the relative importance of stochastic processes ranked as follows: INF (average

83.7% ± 9.3%) > EFF (64.6% ± 13.9%) > AS (58.8% ± 13.9%) (**Fig. 2B**).

VPA results showed that biotic (i.e., local bacterial communities and immigration

of bacteria and ARGs), abiotic (i.e., physicochemical parameters) factors and MGEs

could explain 36.5%~40.5% of the total variance of ARG communities in different

treatment units (**Fig. 4A, C, & E**). Although the pure effects of abiotic factors and

MGEs are relatively low in three treatment units, the interaction effect of these factors

was high in INF. Notably, the pure effects of biotic factors on the total variance of

ARG communities increased greatly as wastewater flowed to the downstream

treatment units (5.5% in INF, 13.2% AS and 23.5% in EFF). The relative importance

of an individual factor to the variance of ARG communities was assessed by HP

analysis (**Fig. 4B, D, & F**). The results indicated that bacterial immigration from INF

is the most important to the total variance of the ARG communities in AS. And ARG

immigrations from INF and AS were the top two important factors affecting ARG

communities in EFF. Besides, the pure effects of MGEs on ARG communities

decreased along with treatment units.

3.5 Daily mass loads of ARGs to the urban environment at the city level

Daily mass loads received by WWTPs across the whole city were 1.21×10^{20}

copies/d ARGs and 2.01×10^{19} copies/d MGEs. And WWTPs discharged 1.44×10^{18}

copies/d ARGs and 2.43×10^{17} copies/d MGEs to the receiving environment. The

citizen's daily ARG and MGE mass loads to WWTPs can be calculated accordingly,

which are up to 3.13×10^{13} copies/d/capita and 5.20×10^{12} copies/d/capita,

respectively. And potential discharge loads to the environment or releasing to humans

from WWTPs were 3.72×10^{11} copies/d/capita, 6.31×10^{10} copies/d/capita,

respectively (**Fig. 5A & B**).

Spearman correlation analysis indicated that total PPCPs mass loads, other PPCPs

mass loads, and socioeconomic factors (i.e., urbanization rate and PCDI) were

significantly correlated with six major types of ARG mass loads and the MGE mass

loads, while antibiotics and NSAIDs had no significant correlation with any types of

ARGs (**Fig. 5C**). In addition, MGE loads had the highest correlation coefficient with

the total PPCPs mass loads.

4. Discussion

To facilitate the monitoring and control of ARG dissemination, setting up a

wastewater-based surveillance approach as well as understanding the ecological

processes and driving forces of the ARG communities along the wastewater treatment

processes are required. Unlike the previous studies that mainly focused on the fate of

ARGs in WWTPs, we targeted the variation and transport of ARGs in different

wastewater treatment units. Here, we found the stable daily variance of ARGs at the

city level through large-scale and continuous daily sampling. Furthermore, stochastic

processes were the dominant assembly mechanism of ARG communities in WWTPs,

although the importance in each treatment unit varied. Among the driving factors, the

immigration of bacteria and ARG in the upstream treatment unit significantly shaped

the downstream ARG community. At last, we estimated the daily mass loads of ARGs

to the WWTPs and the ones released from WWTPs at the city level.

4.1 Stable daily ARG profile and loads throughout WWTPs inspired ARG

surveillance strategies

WWTPs are recognized as the breeding reservoirs for ARG dissemination, which harbor ARGs with high diversity and abundance. However, there is limited information regarding the continuous ARG pollution at various WWTP units during wastewater treatment operations. A comprehensive understanding of the daily dynamics and profile of ARGs throughout WWTP units is highly needed to build wastewater-based surveillance approaches for One Health. Here, we found that the composition of the daily ARG communities tended to be stable throughout seven days from all WWTPs in the same city (**Fig. 3B**). Interestingly, the ARG richness and total ARG relative abundance in INF showed low daily variations (**Fig. 3A**). This indicated that the ARGs profiles introduced by sewage formed a relatively stable pattern across entire city, compared to the EFF, which is affected by daily treatment process variability [61]. Another reason for such abundant and stable ARGs is the plateau effect of microbes. That is, microbes have reached the highest carrying capacity of ARGs in special environment. The plateau effect was frequently observed in field irrigation experiments [24,62]. For instance, the relative abundance of *sul1* in fields irrigated with untreated wastewater for 1.5 years was similar to that in fields irrigated with wastewater for 10 years [62]. This phenomenon requires continuous stimulation of high ARGs to stabilize [24]. Furthermore, the high stability of ARG profiles in INF indicates the locational reproducibility, suggesting that applying a low sampling frequency of sewage would be useful for ARG surveillance in global WWTPs. Similar phenomenon, such as shifts and increases in variability of ARGs from INF to EFF, have been observed in other studies [61,63]. The relatively higher daily variation of ARGs in EFF also presents advantage in assessing WWTP performance and providing a measure of loads and potential associated exposures to the receiving environment. Additionally, it may be better than by INF surveillance alone for detecting and dealing with anomalies of concern [61]. Another landmark study in

Hong Kong WWTPs with similar climate and geographical location showed no
distinct seasonal variations over 9 years span. However, the ARG composition in AS
shifted every 2~3 years [64]. In this study, no such turnover period was captured, and
it is unclear to what extent the observed patterns are generalizable across all WWTPs.
Future research would focus long period sampling for explore the ARG turnover
pattern and dynamics across each treatment processes.

Our study revealed that the relative abundance of ARGs showed higher dynamic
robustness than the absolute abundances of ARGs, which is consistent with a previous
study [27]. This study suggested that applying the ARG relative abundance as the
index can avoid minor variations in sample processing. In addition, our hourly
composite sampling over a 24-hour period reduced the impact of accidental
residential water use and provided a closer representation of daily discharge of ARGs
into WWTPs [26,27]. Overall, the above results provided novel insights into sampling
and monitoring strategies for antibiotic resistance surveillance, particularly for
monitoring anomalous genes of clinical concern in WWTPs, that is, short-term
sampling for ARGs profiles can provide representative information for the distribution
of ARGs in WWTPs (at least a week).

4.2 Immigration process determined the assembly of the ARG communities

The ecological mechanisms of ARG community assembly have been investigated
mainly in urban semi-natural environments, such as urban ponds and reservoirs
[33,37,38,65]. However, little is known about the ARG community assembly
processes in engineered environments. Here, the central role of stochastic processes in
the ecological assembly processes of ARG communities in WWTPs was revealed. It
is consistent with our previous study with a small sample size [33]. The prevalence of
such high stochasticity in ARG communities throughout WWTPs in the current study
may be attributed to other factors beyond the effects of small-scale habitats and
homogeneous physicochemical conditions [65,66]. The high correlation between the

abundance of ARGs and MGEs in INF suggested that HGT may play an important

role in the spread of ARGs (**Fig. S3**). HP analysis found that MGEs were the second

most important factor in determining the ARG communities in INF (**Fig. 4B**). Indeed,

large amounts of nutrients provided beneficial conditions for nourishing microbes as

well as pathogens, which could facilitate cell contact and their HGT of ARGs there

[12,67].

In AS, we found that the immigration of bacteria from INF and the local bacterial

communities was more vital compared to the other two factors (i.e., MGEs and abiotic

factors) (**Fig. 4C & D**). In a previous study, the bacterial community was considered

to be the key factor that drove the ARG profiles in the high anthropogenic

environment [51]. The variation in the microbial community composition was

concurrent with stable metabolic functional structure and performance [68]. In the AS

bioreactor, to maintain high efficiency in nutrients removal, the growth of species

with desired function or high adaptability is facilitated via operation parameters such

as solids retention time and recycle ratio, which is a deterministic process that may

affect the proliferation of ARGs [36,40,69]. This indicated the importance of VGT for

ARG reproduction in AS. Recently, researchers confirmed that the VGT is

significantly involved in the ARG community of AS via single-cell microfluidics

[70,71]. Thus, the existing understanding suggested deterministic factors, such as the

design and operation of AS bioreactors, are suggested to contribute to the AS

microbial communities. However, we showed that microbial immigration from

upstream (stochastic processes) mainly controls the ARG communities in AS.

Similarly, Dottorini et al. found that the influent microbial community immigration

determined the assembly of the AS microbial communities [40]. Previously, INF

microbial communities were often considered as the seeding for the AS communities.

However, here, mass immigration is responsible for both the presence and abundance

of ARGs in the AS, especially for those species with ARGs that are not fit for the

function of AS.

In EFF, the low bacterial density and diversity, as well as the low concentration of nutrients and antibiotics limited the frequency of VGT occurrence, while the high potential for natural transformation of ARG via HGT was confirmed [72,73]. Interestingly, our analysis revealed that the immigration of upstream ARGs had the highest effects on shaping the ARG communities, followed by the upstream and local bacterial communities and then MGEs (**Fig. 4F**). Tong et al. think bacteria host with ARGs may survive better compared with the bacteria without ARGs during the treatment processes [18]. This explained why the immigration of ARGs played a more important role in controlling the ARG community than the bacterial community. This also tells us the source ARG community is essential for the ARG discharges from WWTPs to the environment, challenging us to revise our understanding to control ARGs in sewer systems before WWTPs. Besides, the contribution of the unknown source for the ARG community in AS and EFF decreased along with the treatment units, indicating the limitation of the general source library of ARGs in WWTPs. Future studies focusing on providing more source information into the analysis and therefore increasing the range of identifiable sources in WWTPs are needed.

Although researchers found that EFF largely reflected AS and INF by microbes escaping settling, it is hard to quantify the influence from upstream [61]. We reported the first result highlighting the importance of the immigration of bacteria and ARGs from the source/upstream in shaping the downstream ARGs communities in WWTPs. Our results revealed that the shift of the ARG community structure is mainly affected by the immigration from upstream treatment units, but the immigration of the bacterial and ARG communities contributed differently. WWTPs were continuously fed by upstream sewage sources and were open ecosystems [40,74]. The INF including bacteria was considered to be the main source of biomass in AS and may influence the diversity, abundance, and assembly of local microbial communities microbes via microbial immigration [40–42,75,76]. Although microbial immigration constantly contributes to community assemblage in natural and engineered microbial

ecosystems, the impact of microbial immigration varied due to different immigration

intensities [77,78]. ARGs are prone to persist in the receiving environment once

introduced [24]. For example, effluents of WWTPs rapidly stabilized the antibiotic

resistome in receiving freshwater, even if bacterial community composition was not

yet stable, implying the occurrence of ARGs and pathogens downstream was affected

by the exchange of ARGs from EFF [22,77,79]. However, no evidence was found of

taxa that could drive the stabilization of ARGs in receiving environment [77].

Recently, coupling a mass balance model with qPCR allows the calculation of the

effects of treatment performance and hydrologic characteristics on ARG communities

in receiving environment [21,23]. However, the relative importance of the

immigration of bacterial and ARG communities in shaping ARG communities

remains veiled in engineered ecosystems. Here, we found that the immigration of

bacteria and ARG community determined the ARG communities in AS and EFF,

respectively. The results fill up the gap by providing basic information on the

dissemination of ARGs throughout WWTP treatment units and novel insights for

policy maker to consider the role of mass immigration when developing strategies to

control ARGs.

4.3 WWTPs as ARG dissemination source under socioeconomic development at

the city level

With the rapid development of industry, antibiotics are widely used in animal

husbandry and disease treatment, thus making the resistance gradually become a

serious problem. In our study, we found that socioeconomic factors are closely related

to ARG pollution across the whole city from the perspective of ARG loads.

Population size is the key factor driving fluoroquinolones, macrolide antibiotics,

sulfamethoxazole, chloramphenicol, and daily loads of the corresponding ARGs in

INF [80]. Xiamen is a coastal city with rapid industrialization and economic

development. Two WWTPs with the highest ratio of domestic wastewater in INF

contained the highest relative abundance of total ARGs. This indicated that domestic

wastewater which is closely associated with population size is the main source of

ARGs. Our previous study found that socioeconomic factors (i.e., urbanization rates)

are significantly correlated with PPCP daily mass load per capita [45]. Here, a

positive correlation between two socioeconomic factors (i.e., urbanization rate and

PCDI) and daily ARGs per capita was verified. This suggested that higher human

activity facilitates ARG pollution in highly developed area.

A study found that two municipal WWTPs in South Korea covering < 40% of the

population in Xiamen received average ARG daily loads of around 10^{20} copies/d and

discharged 10^{16}~10^{18} copies/d to the environment [81]. The total ARG daily mass

loads in Xiamen WWTPs in China were in the same range, suggesting the burden of

ARG pollution in highly developed cities. ARG and MGE mass loads are another vital

indicators for ARG dissemination assessment [24]. Thus, despite the fact that two

orders of magnitude of ARGs were removed during the treatment processes of

WWTPs, a massive amount of ARGs and MGEs were released into the environment

across the whole city, leading to the subsequent wide-spread and high risk of exposure

of ARGs to humans.

5. Conclusion

After characterizing the ARG dynamics throughout the wastewater treatment units

in all WWTPs in a city for a week, the following conclusions were therefore drawn:

1. Short-term sampling for ARGs profiles can be representative enough for the

distribution of ARGs in WWTPs in the longer term (at least a week), such as ARG

richness, relative abundance, and absolute abundance of total ARGs, as well as the

compositional dissimilarities of the ARG communities in a week showed little daily

variation, especially in INF.

2. Stochastic processes in assembly ARG communities were dominant in three

treatment units, and the relative importance of stochastic processes in different

treatment units ranked as follows: INF > EFF > AS. The immigration of the bacterial communities upstream and the local bacterial communities played an important role in shaping the ARG communities in AS. The immigration of ARG communities from upstream in WWTPs combined with the local bacterial communities contributed to the ARG dissemination in EFF. This emphasizes the need to revise the way we mitigate ARG contamination but focus on the source in the urban environment.

3. We assessed the total ARG daily mass load received by and discharged from WWTPs at the city level, filling the knowledge gaps on the total amount of ARGs in the urban water cycle. We have linked the socioeconomic factors at the city level with the ARG pollution from WWTPs, providing an assessment of human exposure to ARGs for highly developed cities.

Overall, this study highlighted the daily pattern of ARGs throughout the wastewater treatment units and the importance of mass immigration of ARGs and bacteria from upstream on the formation of ARG communities downstream. This provides the fundamental information for establishing wastewater-based surveillance strategies and ARG ecological behavior in WWTPs.

Author contributions

Lanping Zhang: Conceptualization, Investigation, Methodology, Data Analysis, Software, Visualization, Writing - original draft. **Bob Adyari**: Validation, Software, Writing - review & editing. **Liyuan Hou**: Validation, Writing - review & editing. **Xiaoyong Yang**: Investigation, Writing - review & editing. **Mahmoud Gad**: Writing - review & editing. **Yuwen Wang**: Investigation, Methodology. **Cong Ma**: Investigation, Resources. **Qian Sun**: Methodology, Resources. **Qiang Tang**: Writing - review & editing, Funding acquisition. **Yifeng Zhang**: Writing - review & editing. **Chang-Ping Yu**: Resources, Funding acquisition. **Anyi Hu**: Conceptualization, Data

curation, Validation, Writing - review & editing, Supervision, Project administration,

Funding acquisition.

Acknowledgement

We thank Dr. Fuyi Huang and Miss. Xinyuan Zhou for their assistance with the

methodology. This work was supported by the National Key R&D Program of China

(2022YFE0120300), the National Natural Science Foundation of China (U1805244

and 31870475) and CAS Key Laboratory of Urban Pollutant Conversion Joint

Research Fund (KLUPC-2021-2). M. Gad was supported by the Science, Technology

& Innovation Funding Authority, Egypt (44201).

References

[1]   P. Buchy, S. Ascioglu, Y. Buisson, S. Datta, M. Nissen, P.A. Tambyah, S.

631   Vong, Impact of vaccines on antimicrobial resistance, Int. J. Infect. Dis. 90

632   (2020) 188–196. https://doi.org/10.1016/j.ijid.2019.10.005.

633   [2]   F. Prestinaci, P. Pezzotti, A. Pantosti, Antimicrobial resistance: a global

634   multifaceted phenomenon, Pathog. Glob. Health. 109 (2015) 309–318.

635   https://doi.org/10.1179/2047773215Y.0000000030.

636   [3]   J. Bengtsson-Palme, D.G.J.J. Larsson, Antibiotic resistance genes in the

637   environment: prioritizing risks, Nat. Rev. Microbiol. 13 (2015) 396–396.

638   https://doi.org/10.1038/nrmicro3399-c1.

639   [4]   J. Lebov, K. Grieger, D. Womack, D. Zaccaro, N. Whitehead, B. Kowalcyk,

640   P.D.M. MacDonald, A framework for One Health research, One Heal. 3 (2017)

641   44–50. https://doi.org/10.1016/j.onehlt.2017.03.004.

642   [5]   Y. Yao, Z. Liu, K.K. Yip, Y. Pu, W. Cheng, M. Li, O. Habimana, Cross-

643   regional scale pollution of freshwater biofilms unveiled by antibiotic resistance

644   genes, Sci. Total Environ. 818 (2022) 151835.

645   https://doi.org/10.1016/j.scitotenv.2021.151835.

646 [6] R.S. Hendriksen, P. Munk, P. Njage, B. van Bunnik, L. McNally, O.

647   Lukjancenko, T. Röder, D. Nieuwenhuijse, S.K. Pedersen, J. Kjeldgaard, R.S.

648   Kaas, P.T.L.C. Clausen, J.K. Vogt, P. Leekitcharoenphon, M.G.M.M. van de

649   Schans, T. Zuidema, A.M. de Roda Husman, S. Rasmussen, B. Petersen, A.

650   Bego, C. Rees, S. Cassar, K. Coventry, P. Collignon, F. Allerberger, T.O.

651   Rahube, G. Oliveira, I. Ivanov, Y. Vuthy, T. Sopheak, C.K. Yost, C. Ke, H.

652   Zheng, L. Baisheng, X. Jiao, P. Donado-Godoy, K.J. Coulibaly, M. Jergović, J.

653   Hrenovic, R. Karpíšková, J.E. Villacis, M. Legesse, T. Eguale, A. Heikinheimo,

654   L. Malania, A. Nitsche, A. Brinkmann, C.K.S. Saba, B. Kocsis, N. Solymosi,

655   T.R. Thorsteinsdottir, A.M. Hatha, M. Alebouyeh, D. Morris, M. Cormican, L.

656   O'Connor, J. Moran-Gilad, P. Alba, A. Battisti, Z. Shakenova, C. Kiiyukia, E.

657   Ng'eno, L. Raka, J. Avsejenko, A. Bērziņš, V. Bartkevics, C. Penny, H.

658   Rajandas, S. Parimannan, M.V. Haber, P. Pal, G.-J.J. Jeunen, N. Gemmell, K.

659   Fashae, R. Holmstad, R. Hasan, S. Shakoor, M.L.Z. Rojas, D. Wasyl, G.

660   Bosevska, M. Kochubovski, C. Radu, A. Gassama, V. Radosavljevic, S.

661   Wuertz, R. Zuniga-Montanez, M.Y.F.F. Tay, D. Gavačová, K. Pastuchova, P.

662   Truska, M. Trkov, K. Esterhuyse, K. Keddy, M. Cerdà-Cuéllar, S. Pathirage, L.

663   Norrgren, S. Örn, D.G.J.J. Larsson, T. Van der Heijden, H.H. Kumburu, B.

664   Sanneh, P. Bidjada, B.-M.M. Njanpop-Lafourcade, S.C. Nikiema-Pessinaba, B.

665   Levent, J.S. Meschke, N.K. Beck, C.D. Van, N. Do Phuc, D.M.N. Tran, G.

666   Kwenda, D. adjim Tabo, A.L. Wester, S. Cuadros-Orellana, C. Amid, G.

667   Cochrane, T. Sicheritz-Ponten, H. Schmitt, J.R.M. Alvarez, A. Aidara-Kane,

668   S.J. Pamp, O. Lund, T. Hald, M. Woolhouse, M.P. Koopmans, H. Vigre, T.N.

669   Petersen, F.M. Aarestrup, Global monitoring of antimicrobial resistance based

670   on metagenomics analyses of urban sewage, Nat. Commun. 10 (2019) 1124.

671   https://doi.org/10.1038/s41467-019-08853-3.

[7] S. Liu, P. Wang, C. Wang, X. Wang, J. Chen, Anthropogenic disturbances on

antibiotic resistome along the Yarlung Tsangpo River on the Tibetan Plateau:

Ecological dissemination mechanisms of antibiotic resistance genes to bacterial

pathogens, Water Res. 202 (2021) 117447.

https://doi.org/10.1016/j.watres.2021.117447.

[8] B. Chen, K. Yuan, X. Chen, Y. Yang, T. Zhang, Y. Wang, T. Luan, S. Zou, X.

Li, Metagenomic Analysis Revealing Antibiotic Resistance Genes (ARGs) and

Their Genetic Compartments in the Tibetan Environment, Environ. Sci.

Technol. 50 (2016) 6670–6679. https://doi.org/10.1021/acs.est.6b00619.

[9] J. Li, L. Zhang, Y. Li, K. Liu, Y. Liu, S. Huang, F. Li, C.-T.A. Chen, Y. Zhang,

A. Hu, A Comprehensive Profile of Antibiotic Resistance Genes in the Water

Column of a Shallow-Sea Hydrothermal Vent Ecosystem, Sustainability. 14

(2022) 1776. https://doi.org/10.3390/su14031776.

[10] A. Pruden, D.G.J. Larsson, A. Amézquita, P. Collignon, K.K. Brandt, D.W.

Graham, J.M. Lazorchak, S. Suzuki, P. Silley, J.R. Snape, E. Topp, T. Zhang,

Y.-G. Zhu, Management of Options for Reducing the Release of Antibiotics,

Environ. Health Perspect. 121 (2013) 878–885.

[11] J. Wang, L. Chu, L. Wojnárovits, E. Takács, Occurrence and fate of antibiotics,

antibiotic resistant genes (ARGs) and antibiotic resistant bacteria (ARB) in

municipal wastewater treatment plant: An overview, Sci. Total Environ. 744

(2020) 140997. https://doi.org/10.1016/j.scitotenv.2020.140997.

[12] Y. Zhao, T. Cocerva, S. Cox, S. Tardif, J.-Q. Su, Y.-G. Zhu, K.K. Brandt,

Evidence for co-selection of antibiotic resistance genes and mobile genetic

elements in metal polluted urban soils, Sci. Total Environ. 656 (2019) 512–520.

https://doi.org/10.1016/j.scitotenv.2018.11.372.

[13] Z.-H. Li, L. Yuan, L. Wang, Q.-H. Liu, G.-P. Sheng, Coexistence of silver ion

and tetracycline at environmentally relevant concentrations greatly enhanced

antibiotic resistance gene development in activated sludge bioreactor, J. Hazard.

Mater. 423 (2022) 127088. https://doi.org/10.1016/j.jhazmat.2021.127088.

[14] Y.-G.G. Zhu, T.A. Johnson, J.-Q.Q. Su, M. Qiao, G.-X.X. Guo, R.D. Stedtfeld,

S.A. Hashsham, J.M. Tiedje, Diverse and abundant antibiotic resistance genes

in Chinese swine farms, Proc. Natl. Acad. Sci. 110 (2013) 3435–3440.

https://doi.org/10.1073/pnas.1222743110.

[15] Z. Su, A. Li, J. Chen, B. Huang, Q. Mu, L. Chen, D. Wen, Wastewater

discharge drives ARGs spread in the coastal area: A case study in Hangzhou

Bay, China, Mar. Pollut. Bull. 151 (2020) 110856.

https://doi.org/10.1016/j.marpolbul.2019.110856.

[16] H. Lin, L. Jiang, B. Li, Y. Dong, Y. He, Y. Qiu, Screening and evaluation of

heavy metals facilitating antibiotic resistance gene transfer in a sludge bacterial

community, Sci. Total Environ. 695 (2019) 133862.

https://doi.org/10.1016/j.scitotenv.2019.133862.

[17] F. Ju, K. Beck, X. Yin, A. Maccagnan, C.S. McArdell, H.P. Singer, D.R.

Johnson, T. Zhang, H. Bürgmann, Wastewater treatment plant resistomes are

shaped by bacterial composition, genetic exchange, and upregulated expression

in the effluent microbiomes, ISME J. 13 (2019) 346–360.

https://doi.org/10.1038/s41396-018-0277-8.

[18] J. Tong, A. Tang, H. Wang, X. Liu, Z. Huang, Z. Wang, J. Zhang, Y. Wei, Y.

Su, Y. Zhang, Microbial community evolution and fate of antibiotic resistance

genes along six different full-scale municipal wastewater treatment processes,

Bioresour. Technol. 272 (2019) 489–500.

https://doi.org/10.1016/j.biortech.2018.10.079.

[19] X. Yin, Y. Yang, Y. Deng, Y. Huang, L. Li, L.Y.L. Chan, T. Zhang, An

assessment of resistome and mobilome in wastewater treatment plants through

temporal and spatial metagenomic analysis, Water Res. 209 (2022) 117885.

https://doi.org/10.1016/j.watres.2021.117885.

[20] X.-L.L. An, J.-Q.Q. Su, B. Li, W.-Y.Y. Ouyang, Y. Zhao, Q.-L.L. Chen, L. Cui,

H. Chen, M.R. Gillings, T. Zhang, Y.-G.G. Zhu, Tracking antibiotic resistome

during wastewater treatment using high throughput quantitative PCR, Environ.

Int. 117 (2018) 146–153. https://doi.org/10.1016/j.envint.2018.05.011.

[21] C.U. Schwermer, W. Uhl, Calculating expected effects of treatment effectivity

and river flow rates on the contribution of WWTP effluent to the ARG load of

a receiving river, J. Environ. Manage. 288 (2021) 112445.

https://doi.org/10.1016/j.jenvman.2021.112445.

[22] G. Reichert, S. Hilgert, J. Alexander, J.C. Rodrigues de Azevedo, T. Morck, S.

Fuchs, T. Schwartz, Determination of antibiotic resistance genes in a WWTP-

impacted river in surface water, sediment, and biofilm: Influence of seasonality

and water quality, Sci. Total Environ. 768 (2021) 144526.

https://doi.org/10.1016/j.scitotenv.2020.144526.

[23] J. Lee, F. Ju, A. Maile-Moskowitz, K. Beck, A. Maccagnan, C.S. McArdell, M.

Dal Molin, F. Fenicia, P.J. Vikesland, A. Pruden, C. Stamm, H. Bürgmann,

Unraveling the riverine antibiotic resistome: The downstream fate of

anthropogenic inputs, Water Res. 197 (2021) 117050.

https://doi.org/10.1016/j.watres.2021.117050.

[24] I.D. Kampouris, S. Agrawal, L. Orschler, D. Cacace, S. Kunze, T.U.

Berendonk, U. Klümper, Antibiotic resistance gene load and irrigation intensity

determine the impact of wastewater irrigation on antimicrobial resistance in the

soil microbiome, Water Res. 193 (2021).

https://doi.org/10.1016/j.watres.2021.116818.

[25] R.B.M. Marano, A. Zolti, E. Jurkevitch, E. Cytryn, Antibiotic resistance and

class 1 integron gene dynamics along effluent, reclaimed wastewater irrigated

soil, crop continua: elucidating potential risks and ecological constraints, Water

Res. 164 (2019) 114906. https://doi.org/10.1016/j.watres.2019.114906.

[26] M. Cai, Z. Wang, H. Gu, H. Dong, X. Zhang, N. Cui, L. Zhou, G. Chen, G.

Zou, Occurrence and temporal variation of antibiotics and antibiotic resistance

genes in hospital inpatient department wastewater: Impacts of daily schedule of

inpatients and wastewater treatment process, Chemosphere. 292 (2022) 133405.

https://doi.org/10.1016/j.chemosphere.2021.133405.

[27] S. Sun, J. Geng, B. Li, L. Ma, X. Sun, F. Meng, H. Qi, J. Shen, Temporal

variations of antibiotic resistance genes in influents and effluents of a WWTP

in cold regions, J. Clean. Prod. 328 (2021) 129632.

https://doi.org/10.1016/j.jclepro.2021.129632.

[28] M. Vellend, Conceptual Synthesis in Community Ecology, Q. Rev. Biol. 85

(2010) 183–206. https://doi.org/10.1086/652373.

[29] J. Zhou, D. Ning, Stochastic Community Assembly: Does It Matter in

Microbial Ecology?, Microbiol. Mol. Biol. Rev. 81 (2017).

https://doi.org/10.1128/MMBR.00002-17.

[30] N. Daliang, D. Ye, T.J. M., Z. Jizhong, A general framework for quantitatively

assessing ecological stochasticity, Proc. Natl. Acad. Sci. 116 (2019) 16892–

16898. https://doi.org/10.1073/pnas.1904623116.

[31] H. Chen, C. Liu, Y. Teng, Z. Zhang, Y. Chen, Y. Yang, Environmental risk

characterization and ecological process determination of bacterial antibiotic

resistome in lake sediments, Environ. Int. 147 (2021) 106345.

https://doi.org/10.1016/j.envint.2020.106345.

[32] L. Hou, H. Wang, Q. Chen, J.-Q.Q. Su, M. Gad, J. Li, S.I. Mulla, C.-P.P. Yu, A.

Hu, Fecal pollution mediates the dominance of stochastic assembly of

antibiotic resistome in an urban lagoon (Yundang lagoon), China, J. Hazard.

Mater. 417 (2021) 126083. https://doi.org/10.1016/j.jhazmat.2021.126083.

[33] L. Hou, L. Zhang, F. Li, S. Huang, J. Yang, C. Ma, D. Zhang, C.-P. Yu, A. Hu,

Urban ponds as hotspots of antibiotic resistome in the urban environment, J.

Hazard. Mater. 403 (2021) 124008.

https://doi.org/10.1016/j.jhazmat.2020.124008.

[34] R. Pallares-vega, H. Blaak, R. Van Der Plaats, A.M. De Roda, L. Hernandez,

M.C.M. Van Loosdrecht, D.G. Weissbrodt, H. Schmitt, A. Gene, Determinants

of presence and removal of antibiotic resistance genes during WWTP

treatment☐: A cross-sectional study, Water Res. 161 (2019) 319–328.

https://doi.org/10.1016/j.watres.2019.05.100.

[35] J. Chen, Z. Zhang, Z. Lei, K. Shimizu, P. Yao, Z. Su, D. Wen, Occurrence and

distribution of antibiotic resistance genes in the coastal sediments of effluent-

receiving areas of WWTPs, China, Bioresour. Technol. Reports. 11 (2020)

100511. https://doi.org/10.1016/j.biteb.2020.100511.

[36] D. Dai, C. Brown, H. Bürgmann, D.G.J.J. Larsson, I. Nambi, T. Zhang, C.-F.

Flach, A. Pruden, P.J. Vikesland, Long-read metagenomic sequencing reveals

shifts in associations of antibiotic resistance genes with mobile genetic

elements from sewage to activated sludge, Microbiome. 10 (2022) 20.

https://doi.org/10.1186/s40168-021-01216-5.

[37] Y. Guo, M. Liu, L. Liu, X. Liu, H. Chen, J. Yang, The antibiotic resistome of

free-living and particle-attached bacteria under a reservoir cyanobacterial

bloom, Environ. Int. 117 (2018) 107–115.

https://doi.org/10.1016/j.envint.2018.04.045.

[38] P. Fang, F. Peng, X. Gao, P. Xiao, J. Yang, Decoupling the dynamics of

bacterial taxonomy and antibiotic resistance function in a subtropical urban

reservoir as revealed by high-frequency sampling, Front. Microbiol. 10 (2019)

1–12. https://doi.org/10.3389/fmicb.2019.01448.

[39] Z. Zhao, J. Wang, Y. Han, J. Chen, G. Liu, H. Lu, B. Yan, S. Chen, Nutrients,

heavy metals and microbial communities co-driven distribution of antibiotic

resistance genes in adjacent environment of mariculture, Environ. Pollut. 220

(2017) 909–918. https://doi.org/10.1016/j.envpol.2016.10.075.

[40] G. Dottorini, T.Y. Michaelsen, S. Kucheryavskiy, K.S. Andersen, J.M.

Kristensen, M. Peces, D.S. Wagner, M. Nierychlo, P.H. Nielsen, Mass-

immigration determines the assembly of activated sludge microbial

communities, Proc. Natl. Acad. Sci. U. S. A. 118 (2021) 1–11.

https://doi.org/10.1073/pnas.2021589118.

[41] M. Ali, Z. Wang, K.W. Salam, A.R. Hari, M. Pronk, M.C.M. Van Loosdrecht,

P.E. Saikaly, Importance of species sorting and immigration on the bacterial

assembly of different-sized aggregates in a full-scale Aerobic granular sludge

plant, Environ. Sci. Technol. 53 (2019) 8291–8301.

https://doi.org/10.1021/acs.est.8b07303.

[42] R. Mei, W.-T.T. Liu, Quantifying the contribution of microbial immigration in

engineered water systems, Microbiome. 7 (2019) 1–8.

https://doi.org/10.1186/s40168-019-0760-0.

[43] A. Hu, S. Li, L. Zhang, H. Wang, J. Yang, Z. Luo, A. Rashid, S. Chen, W.

Huang, C.-P. Yu, Prokaryotic footprints in urban water ecosystems: A case

study of urban landscape ponds in a coastal city, China, Environ. Pollut. 242

(2018) 1729–1739. https://doi.org/10.1016/j.envpol.2018.07.097.

[44] Xiamen Bureau of Statistics, 2017 Yearbook of Xiamen Special Economic

Zone, China Statisitcs Press. (2017). https://doi.org/http://www.stats-

xm.gov.cn/2017/main0.htm.

[45] Y.Y. Wang, Y. Li, A. Hu, A. Rashid, M. Ashfaq, Y.Y. Wang, H. Wang, H. Luo,

C.P. Yu, Q. Sun, Monitoring, mass balance and fate of pharmaceuticals and

personal care products in seven wastewater treatment plants in Xiamen City,

China, J. Hazard. Mater. 354 (2018) 81–90.

https://doi.org/10.1016/j.jhazmat.2018.04.064.

[46] China's State Environment Protection Agency, Water and Waste Water

Monitoring and Analysis Methods, China Environ. Sci. Press. (2002).

[47] T. Han, J. Zheng, Y. Han, X. Xu, M. Li, C. Schwarz, L. Zhu, Comprehensive

insights into core microbial assemblages in activated sludge exposed to textile-

dyeing wastewater stress, Sci. Total Environ. 791 (2021) 148145.

https://doi.org/10.1016/j.scitotenv.2021.148145.

[48] A. Hu, F. Ju, L. Hou, J. Li, X. Yang, H. Wang, S.I. Mulla, Q. Sun, H.

Bürgmann, C.-P. Yu, Strong impact of anthropogenic contamination on the co-

occurrence patterns of a riverine microbial community, Environ. Microbiol. 19

(2017) 4993–5009. https://doi.org/10.1111/1462-2920.13942.

[49] F. Hildebrand, R. Tadeo, A.Y. Voigt, P. Bork, J. Raes, LotuS: an efficient and

user-friendly OTU processing pipeline, Microbiome. 2 (2014) 30.

https://doi.org/10.1186/2049-2618-2-30.

[50] R.C. Edgar, UPARSE: highly accurate OTU sequences from microbial

amplicon reads, Nat. Methods. 10 (2013) 996–998.

https://doi.org/10.1038/nmeth.2604.

[51] J. Su, B. Wei, W.-Y. Ou-Yang, F. Huang, Y. Zhao, H. Xu, Y. Zhu, Antibiotic

Resistome and Its Association with Bacterial Communities during Sewage

Sludge Composting, Environ. Sci. Technol. 49 (2015) 7356–7363.

https://doi.org/10.1021/acs.est.5b01012.

[52] Y.-G.G. Zhu, Y. Zhao, B. Li, C.-L.L. Huang, S.-Y.Y. Zhang, S. Yu, Y.-S.S.

Chen, T. Zhang, M.R. Gillings, J.-Q.Q. Su, Continental-scale pollution of

estuaries with antibiotic resistance genes, Nat. Microbiol. 2 (2017) 16270.

https://doi.org/10.1038/nmicrobiol.2016.270.

[53] D. Knights, J. Kuczynski, E.S. Charlson, J. Zaneveld, M.C. Mozer, R.G.

Collman, F.D. Bushman, R. Knight, S.T. Kelley, Bayesian community-wide

culture-independent microbial source tracking, Nat. Methods. 8 (2011) 761–

765. https://doi.org/10.1038/nmeth.1650.

[54] M. Gad, L. Hou, M. Cao, B. Adyari, L. Zhang, D. Qin, C.P. Yu, Q. Sun, A. Hu,

Tracking microeukaryotic footprint in a peri-urban watershed, China through

machine-learning approaches, Sci. Total Environ. 806 (2022) 150401.

https://doi.org/10.1016/j.scitotenv.2021.150401.

[55]   L.-G.G. Li, X. Yin, T. Zhang, Tracking antibiotic resistance gene pollution

from different sources using machine-learning classification, Microbiome. 6

(2018) 93. https://doi.org/10.1186/s40168-018-0480-x.

[56]   B. Naimi, N.A.S. Hamm, T.A. Groen, A.K. Skidmore, A.G. Toxopeus, Where

is positional uncertainty a problem for species distribution modelling?,

Ecography (Cop.). 37 (2014) 191–203. https://doi.org/10.1111/j.1600-

0587.2013.00205.x.

[57]   J. Lai, Y. Zou, J. Zhang, P.R. Peres-Neto, Generalizing hierarchical and

variation partitioning in multiple regression and canonical analyses using the

rdacca.hp R package, Methods Ecol. Evol. 2022 (2022) 782–788.

https://doi.org/10.1111/2041-210X.13800.

[58]   P.J. McMurdie, S. Holmes, phyloseq: An R Package for Reproducible

Interactive Analysis and Graphics of Microbiome Census Data, PLoS One. 8

(2013) e61217. https://doi.org/10.1371/journal.pone.0061217.

[59]   J. Oksanen, F.G. Blanchet, P. Kindt, R., Legendre, P.R. Minchin, R.B. O'Hara,

G.L. Simpson, P. Sólymos, M.H.H. Stevens, H. Wagner, Vegan: Community

Ecology Package, R Packag. Version 1.15-2. (2012).

https://doi.org/http://cran.r-project.org/web/ packages/vegan/index.html.

[60]   H. Wickham, A Layered Grammar of Graphics, J. Comput. Graph. Stat. 19

(2010) 3–28. https://doi.org/10.1198/jcgs.2009.07098.

[61]   H.J. Majeed, M. V. Riquelme, B.C. Davis, S. Gupta, L. Angeles, D.S. Aga, E.

Garner, A. Pruden, P.J. Vikesland, Evaluation of Metagenomic-Enabled

Antibiotic Resistance Surveillance at a Conventional Wastewater Treatment

Plant, Front. Microbiol. 12 (2021) 1–19.

https://doi.org/10.3389/fmicb.2021.657954.

[62] P. Dalkmann, M. Broszat, C. Siebe, E. Willaschek, T. Sakinc, J. Huebner, W.

Amelung, E. Grohmann, J. Siemens, Accumulation of pharmaceuticals,

enterococcus, and resistance genes in soils irrigated with wastewater for zero to

100 years in central Mexico, PLoS One. 7 (2012) e45397.

https://doi.org/10.1371/journal.pone.0045397.

[63] F. Lira, I. Vaz-Moreira, J. Tamames, C.M. Manaia, J.L. Martínez,

Metagenomic analysis of an urban resistome before and after wastewater

treatment, Sci. Rep. 10 (2020) 8174. https://doi.org/10.1038/s41598-020-

65031-y.

[64] X. Yin, Y. Deng, L. Ma, Y. Wang, L.Y.L. Chan, T. Zhang, Exploration of the

antibiotic resistome in a wastewater treatment plant by a nine-year longitudinal

metagenomic study, Environ. Int. 133 (2019) 105270.

https://doi.org/10.1016/j.envint.2019.105270.

[65] F. Peng, Y. Guo, A. Isabwe, H. Chen, Y. Wang, Y. Zhang, Z. Zhu, J. Yang,

Urbanization drives riverine bacterial antibiotic resistome more than taxonomic

community at watershed scale, Environ. Int. 137 (2020) 105524.

https://doi.org/10.1016/j.envint.2020.105524.

[66] W.T. Sloan, M. Lunn, S. Woodcock, I.M. Head, S. Nee, T.P. Curtis,

Quantifying the roles of immigration and chance in shaping prokaryote

community structure, Environ. Microbiol. 8 (2006) 732–740.

https://doi.org/10.1111/j.1462-2920.2005.00956.x.

[67] Y. He, M. Bai, Y. He, S. Wang, J. Zhang, S. Jiang, G. Wang, Suspended

particles are hotspots for pathogen-related bacteria and ARGs in coastal beach

waters of northern China, Sci. Total Environ. 817 (2022) 153004.

https://doi.org/10.1016/j.scitotenv.2022.153004.

[68] S. Louca, M. Doebeli, Taxonomic variability and functional stability in

microbial communities infected by phages, Environ. Microbiol. 19 (2017)

3863–3878. https://doi.org/10.1111/1462-2920.13743.

[69] C. Wang, Y. Yang, Y.Y. Wang, D. Wang, X. Xu, Y.Y. Wang, L. Li, C. Yang,

T. Zhang, Absolute quantification and genome-centric analyses elucidate the

dynamics of microbial populations in anaerobic digesters, Water Res. 224

(2022) 119049. https://doi.org/10.1016/j.watres.2022.119049.

[70] B. Li, Y. Qiu, Y. Song, H. Lin, H. Yin, Dissecting horizontal and vertical gene

transfer of antibiotic resistance plasmid in bacterial community using

microfluidics, Environ. Int. 131 (2019) 105007.

https://doi.org/10.1016/j.envint.2019.105007.

[71] Z. Wei, K. Feng, Z. Wang, Y. Zhang, M. Yang, Y.-G.G. Zhu, M.P.J.J. Virta, Y.

Deng, High-Throughput Single-Cell Technology Reveals the Contribution of

Horizontal Gene Transfer to Typical Antibiotic Resistance Gene Dissemination

in Wastewater Treatment Plants, Environ. Sci. Technol. 55 (2021) 11824–

11834. https://doi.org/10.1021/acs.est.1c01250.

[72] M. Jin, L. Liu, D. ning Wang, D. Yang, W. li Liu, J. Yin, Z. wei Yang, H. ran

Wang, Z. gang Qiu, Z. qiang Shen, D. yang Shi, H. bei Li, J. hua Guo, J. wen

Li, Chlorine disinfection promotes the exchange of antibiotic resistance genes

across bacterial genera by natural transformation, ISME J. 14 (2020) 1847–

1856. https://doi.org/10.1038/s41396-020-0656-9.

[73] P. Dong, H. Wang, T. Fang, Y. Wang, Q. Ye, Assessment of extracellular

antibiotic resistance genes (eARGs) in typical environmental samples and the

transforming ability of eARG, Environ. Int. 125 (2019) 90–96.

https://doi.org/10.1016/j.envint.2019.01.050.

[74] S.H. Lee, H.J. Kang, H.D. Park, Influence of influent wastewater communities

on temporal variation of activated sludge communities, Water Res. 73 (2015)

132–144. https://doi.org/10.1016/j.watres.2015.01.014.

[75] A.M. Saunders, M. Albertsen, J. Vollertsen, P.H. Nielsen, The activated sludge

ecosystem contains a core community of abundant organisms, ISME J. 10

(2016) 11–20. https://doi.org/10.1038/ismej.2015.117.

[76] S. Zhang, F. Meng, Core activated sludge communities are influenced little by

immigration: Case study of a membrane bioreactor plant, J. Environ. Sci. 102

(2021) 244–255. https://doi.org/10.1016/j.jes.2020.09.024.

[77] G. Corno, Y. Yang, E.M. Eckert, D. Fontaneto, A. Fiorentino, S. Galafassi, T.

Zhang, A. Di Cesare, Effluents of wastewater treatment plants promote the

rapid stabilization of the antibiotic resistome in receiving freshwater bodies,

Water Res. 158 (2019) 72–81. https://doi.org/10.1016/j.watres.2019.04.031.

[78] R. Mei, J. Kim, F.P. Wilson, B.T.W.W. Bocher, W.-T. Liu, Coupling growth

kinetics modeling with machine learning reveals microbial immigration

impacts and identifies key environmental parameters in a biological wastewater

treatment process, Microbiome. 7 (2019) 65. https://doi.org/10.1186/s40168-

019-0682-x.

[79] M. Harnisz, E. Kiedrzyńska, M. Kiedrzyński, E. Korzeniewska, M.

Czatzkowska, I. Koniuszewska, A. Jóźwik, S. Szklarek, S. Niestępski, M.

Zalewski, The impact of WWTP size and sampling season on the prevalence of

antibiotic resistance genes in wastewater and the river system, Sci. Total

Environ. 741 (2020) 140466. https://doi.org/10.1016/j.scitotenv.2020.140466.

[80] F.C.T. Elder, K. Proctor, R. Barden, W.H. Gaze, J. Snape, E.J. Feil, B.

Kasprzyk-Hordern, Spatiotemporal profiling of antibiotics and resistance genes

in a river catchment: Human population as the main driver of antibiotic and

antibiotic resistance gene presence in the environment, Water Res. 203 (2021)

117533. https://doi.org/10.1016/j.watres.2021.117533.

[81] J. Lee, J.H. Jeon, J. Shin, H.M. Jang, S. Kim, M.S. Song, Y.M. Kim,

Quantitative and qualitative changes in antibiotic resistance genes after passing

through treatment processes in municipal wastewater treatment plants, Sci.

Total Environ. 605–606 (2017) 906–914.

https://doi.org/10.1016/j.scitotenv.2017.06.250.

Figure 1. Relative abundance (A) and absolute abundance (B) of ARGs and MGEs in different wastewater treatment units. MLSB stands for Macrolide-Lincosamide-Streptogramin B.

Figure 2. Non-metric multidimensional scaling (NMDS) ordination based on Bray-Curtis distance showing the distribution pattern of ARG communities (A) and bacterial communities (C) in all WWTPs across the entire city. The ellipses were drawn at the 95% confidence level. Abundance-based β-null model was applied to assess the relative importance of the stochasticity process in governing the assembly of ARG (B) and bacterial communities (D) in each treatment unit during the sampling period (from day 1 to day 7). Significant differences were indicated by different letters among different units of the treatment process.

Figure 3. Daily variation of ARGs and bacteria at different wastewater treatment units. (A) Coefficient of variation (CV) of richness and abundance of overall ARG community and OTU richness, (B) time-decay relationships of the compositional dissimilarity of ARG and bacterial communities at different time intervals. Significant differences among different units of the wastewater treatment process were assessed by paired Wilcoxon test and indicated as follows: ***$P < 0.001$, **$P < 0.01$ and *$P < 0.05$. Lines represent the fitted linear regression curve. The shaded areas represent the 95% confidential interval of the fitted linear regression curve.

Figure 4. Variation partitioning analysis (VPA) of β-diversity variation of ARG communities among biotic, abiotic, and MGEs groups as well as their interactions (A, C, E). The absence of any values for the three interactions in Figure C & E implies that the interaction effect was 0. MGEs are considered to be a unique class of factors that contribute to the assembly of the ARG community, and therefore are separated from the biotic group as a single factor. The top five variables with the highest variance explained to β-diversity variation of ARG communities in each treatment unit are indicated by hierarchical partitioning (HP) analysis (B, D, F). The abbreviation of BAC NMDS1 and BAC NMDS2 represent the first two axes of NMDS results of the bacterial community in each treatment unit, and the abbreviation of BAC INF, BAC AS, ARG INF, and ARG AS represents the source of the bacterial community or ARG

community (upstream treatment units such as INF or AS). Significant differences were indicated as follows: ***$P < 0.001$, **$P < 0.01$ and *$P < 0.05$.

Figure 5. The daily ARG and MGE receiving loads per capita by WWTPs (A) and discharge loads from WWTPs (B) in Xiamen, China. Spearman correlation analysis was made between major types of the daily discharge ARG loads per capita and the daily PPCP loads in WWTPs and social development factors of the city (C). Significant differences were indicated as follows: ***P < 0.001, **P < 0.01 and *P < 0.05.