Comparative analysis of amphibian genomes: an emerging resource for basic and applied research

Running title: Comparative analysis of amphibian genomes

Tiffany A. Kosch[1], Andrew J. Crawford[2], Rachel Lockridge Mueller[3], Katharina C. Wollenberg Valero[4],

Megan L. Power[4], Ariel Rodríguez [5], Lauren A. O'Connell[6], Neil D. Young[1], and Lee F. Skerratt[1]

[1] Faculty of Science, University of Melbourne, Melbourne, Australia

[2] Departamento de Ciencias Biológicas, Universidad de los Andes, Bogotá, Colombia

[3] Department of Biology, Colorado State University, Colorado, USA

[4] School of Biology and Environmental Science, University College Dublin, Dublin, Ireland

[5] Institute of Zoology, University of Veterinary Medicine of Hannover, Hannover, Germany

[6] Department of Biology, Stanford University, California, USA

Corresponding author: tiffany.kosch@gmail.com

ABSTRACT

Amphibians are the most threatened group of vertebrates and are in dire need of conservation

intervention to ensure their continued survival. They exhibit unique features including a high

diversity of reproductive strategies, permeable and specialized skin capable of producing toxins and

antimicrobial compounds, multiple genetic mechanisms of sex determination, and in some lineages,

the ability to regenerate limbs and organs. Although genomics approaches would shed light on these

unique traits and aid conservation, sequencing and assembly of amphibian genomes has lagged

behind other taxa due to their comparatively large genome sizes. Fortunately, the development of

long-read sequencing technologies and initiatives has led to a recent burst of new amphibian

genome assemblies. Although growing, the field of amphibian genomics suffers from the lack of

annotation resources, tools for working with challenging genomes, and lack of high-quality

assemblies in multiple clades of amphibians. Here we analyze 51 publicly available amphibian

genomes to evaluate their usefulness for functional genomics research. We report considerable

variation in genome assembly quality and completeness, and report some of the highest

transposable element and repeat contents of any vertebrate. Additionally, we detected an

association between transposable element content and climatic variables. Our analysis provides

evidence of conserved genome synteny despite the long divergence times of this group, but we also

highlight inconsistencies in chromosome naming and orientation across genome assemblies. We

discuss sequencing gaps in the phylogeny and suggest key targets for future sequencing endeavors.

Finally, we propose increased investment in amphibian genomics research to promote their

conservation.

KEYWORDS

Amphibian genomes; comparative genomics; transposable elements; repeat expansion;

genome synteny

INTRODUCTION

Amphibians are an ancient lineage of vertebrates that predate amniotes by more than 100 million

years. Despite the considerable age of this lineage, amphibians are now the most threatened group

of vertebrates with more that 40% of species and are threatened by factors such as habitat change,

disease, and over-exploitation (IUCN, 2022; Scheele et al., 2019). Notably, many of these threats are

hard to reverse, suggesting that novel approaches that utilize genomic resources may lead to

improved management decisions for some of the most endangered taxa (Kosch et al., 2022; Scheele

et al., 2014).

We are only just beginning to understand the genetic basis of many of the unique features of

amphibians. Amphibians exhibit a high diversity of reproductive strategies including biphasic and

direct development, uniparental and biparental care, mouth and gastric brooding, and foam nesting

(Brown et al., 2010; Nunes-de-Almeida et al., 2021; Schulte et al., 2020). They also have specialized

skin capable of producing complex compounds of interest for drug discovery for the development of

antimicrobial drugs and analgesics (Daly et al., 2000; De Angelis et al., 2021; Liu et al., 2020).

Amphibians occur across habitat types from rainforests to deserts, freshwater streams to salt

marshes, and tropical to arctic climates (Duellman, 1999), but it is unclear how this ecological

diversity is reflected in genome composition. One potential way is the number of transposable

elements (TEs) present in the genome. TEs have a huge impact on the structure and function of

eukaryotic genomes, with amphibians having among the largest TE content among vertebrates.

There is increasing evidence that TE activity, and thus their relative proportion in genomes, is

influenced by abiotic factors (Pimpinelli & Piacentini, 2020). This in turn highlights their potential role

in the regulation of genetic mechanisms responsible for environmental adaptation (Casacuberta &

González, 2013; Pappalardo et al., 2021). Salamanders are an important resource for transplant and

regeneration research due to their ability to regenerate limbs and internal organs (Elewa et al., 2017;

Nowoshilow et al., 2018). Amphibians also have many of the same immune components of mammals

making them an important model resource for immunology (Paiola et al., 2023; Robert, 2020).

Despite the obvious value of amphibian genomes for research on ecology, evolution, medicine, and

improving their conservation, until recently, the generation of amphibian reference genomes has

been markedly slower than other vertebrates (Hotaling et al., 2021a; Womack et al., 2022). This lag

can be attributed to high costs and the computational challenges of assembling their often large and

complex genomes (Sun et al., 2020). Recent advances in sequencing technologies such as long read

sequencing and assembly algorithms that incorporate hybrid approaches have circumvented many of

these challenges leading to a surge of high quality, chromosome-level reference genomes. The next

challenge will be developing the tools for annotation and comparative analyses of these large

genomes.

In this study, we provide a synthesis of all available amphibian reference genome assemblies, 51 at

the time of our analysis, with the number growing every day. We evaluate assembly quality,

sequencing technology, gene completeness, transposable element and repeat content and its

ecological correlates, taxonomic representation, and synteny.

MATERIALS AND METHODS

Genomes

A search of the NCBI genome website using the search term "amphibians" conducted on August 25, 2023, revealed there were 90 amphibian genomes from 68 species. All genome files in fasta format were downloaded for assessment. Sixteen salamander genomes (Pyron et al., 2024) were excluded from our analyses due to their high degree of incompleteness (i.e., <10% of genome assembled). Of the remaining genomes, one genome was selected for each species for subsequent analysis. If there was more than one draft of a genome, the most recent draft and/or the primary haplotype was selected. In cases where there were multiple versions sequenced by different groups, the best genome was selected by lowest scaffold number. Entire genomes (including uncharacterized contigs but excluding mitochondrial genomes) were used for assessment unless indicated otherwise.

Genome databases NCBI Genomes, NCBI RefSeq (O'Leary et al., 2016), Ensembl (Cunningham et al., 2022), UCSC Genome Browser (Lee et al., 2022), and Genomes on a Tree (GoaT) (Sotero-Caio et al., 2021) were searched for information on the 51 amphibian species with reference genomes including chromosome number, annotation data, proteome availability, C-value, and sequencing technology. Sequencing strategy was classified as "short-single" for Illumina only sequencing, "long-single" for sequencing using long read technologies (e.g., PacBio and Oxford Nanopore), and "hybrid" for sequencing approaches using more than one approach (e.g., PacBio and Hi-C).

A search for amphibian proteome datasets on NCBI RefSeq (O'Leary et al., 2016), Ensembl (Cunningham et al., 2022), and UCSC Genome Browser (Lee et al., 2022) databases on June 24, 2022 revealed 11 proteomics datasets.

A search of the NCBI Organelle database on 15, February 2023 using search term "amphibian"

resulted in 353 mitochondrial genomes belonging to 345 species (Table S11). Seventeen

mitochondrial genomes overlapped with the amphibian nuclear genomes analyzed in this study.

Reference genome availability summary

The GoaT online database (Sotero-Caio et al., 2021) was searched on August 28, 2023 to summarize

genomes in progress or publicly available using the search terms "tax_tree(Amphibia) AND

tax_rank(species) AND sequencing_status=in_progress" or "tax_tree(Amphibia) AND

tax_rank(species) AND sequencing_status=insdc_open". The same search terms were used to

summarize publicly available genomes for mammals, birds, and non-avian reptiles with the

"tax_tree" search term replaced by appropriate Class.

Genome quality analyses

Genome quality assessment was performed with BBMap (v. 39.01) "statswrapper.sh" bash script

(https://github.com/BioInfoTools/BBMap). This tool generates metrics such as genome size, contig

N50, and scaffold count. Benchmarking Universal Single-Copy Orthologs (BUSCO) were summarized

with the BUSCO tool (v. 5.1.2) (Manni et al., 2021) using the OrthoDB Tetrapoda ortholog library (v.

odb10) (Kriventseva et al., 2018) (N=5310 orthologs) with the prompt "-m genome". Percentage of

the genome assembled to chromosomes was calculated with a custom bash script that computes the

genome length assigned to chromosomes and divides it by the "assembly length" value computed by

BBMap.

Phylogenetic tree

A species to family correspondence table was obtained from Jetz and Pyron (2018)

(https://vertlife.org/files_20170703/) and was filtered to include only the species with the longest

nucleotide sequence per family. This taxa subset was used to obtain a subset of 100 phylogenetic

trees from the posterior distribution of the Jetz and Pyron (2018) dataset, as available

from http://vertlife.org/phylosubsets. A consensus tree from these 100 trees was then obtained

using treeannotator (v2.7.5) (settings - target tree type: maximum clade credibility, node heights:

median burn-in percentage: 0, posterior probability limit: 0.0) (Drummond & Rambaut, 2007). The

species names of the tree tips were then substituted with the corresponding family names using the

"sub.taxa.label" function in the phylotools package (https://github.com/helixcn/phylotools) in R with

the aid of the species to family correspondence table, which was updated with the most recent

classification available in AmphibiaWeb (https://amphibiaweb.org) and the Amphibian Species of the

World database (https://amphibiansoftheworld.amnh.org/). In cases where these two references

disagreed, the AmphibiaWeb taxonomy was used.

Repeat modelling and annotation

Repeats were *de novo* modelled with RepeatModeler (Apptainer v. 1.2.3) (Flynn et al., 2020).

Genomes were then annotated using RepeatMasker (v. 4.1.2-p1) (Smit et al., 2013) with a

concatenated library of genome-specific repeats generated from RepeatModeler and the Dfam

amphibian repeat library (v. Dfam.h5) (Storer et al., 2021). Before annotation, any previous soft

masking of the genomes was reversed. The results were summarized using a custom bash and R

scripts.

Ecological correlates of transposable element content

Occurrence data for the 51 species were downloaded from the Global Biodiversity Information

Facility (GBIF) (https://www.gbif/org/; last accessed February 2024 (full DOI's for each occurrence

data set in Table S5). In addition, due to the putative involvement of temperature in TE activity,

BioClim variables associated with temperature (Bio1-Bio11) were obtained for the 51 amphibian

species (Table S6). As previous studies have explored the relationship between amphibian genome

size and environmental variables (Liedtke et al., 2022), here we focused on the relationship between

temperature variables, elevation, and amphibian transposable elements. Influence of these

bioclimatic variables (after removing highly collinear variables, see supplementary methods) on

transposable element content (summarized into three groups: proportion of total transposable

elements (TEs), proportion of retroelements, and proportion of DNA transposons) was modelled

using Bayesian mixed effect models (Hadfield, 2010). To correct for body size, log transformed body

size was included in the model structure with log transformed Bio2 (Mean diurnal range), Bio4

(temperature seasonality), Bio8 (mean temperature of wettest quarter), Bio10 (mean temperature of

warmest quarter) and elevation. Models were also corrected for phylogenetic non-independence

(Figure S5, see Supplementary Methods for further information) with phylogenetically independent

contrasts (Felsenstein, 1985; Garland Jr et al., 1992) plotted with and (Revell, 2024).

Synteny analysis

Synteny of BUSCO genes for chromosome level assemblies was analyzed with R Package GENESPACE

(v. 1.1.4) (Lovell et al., 2022), which uses OrthoFinder (v. 2.5.4) (Emms & Kelly, 2019) to infer

orthology. Synteny was analyzed using BUSCO "full_table.tsv" results files that were reformatted for

GENESPACE input using a custom bash script. Synteny plots were generated for all chromosome level

assemblies, all anuran chromosome level assemblies, for the two salamander genomes, and for the

three caecilian genomes using the GENESPACE plotting tool "plot_riparian". Chromosomes with

reversed orientation compared to the reference genome were inverted to improve visualization.

Quantification and statistical analysis

Regression analyses, ANOVAs, and Student's t-tests for comparing genome quality measurements

were conducted with the R statistics package (v. 4.1.2) (Team, 2013) in R Studio (v. 2022.02.3) (Team,

2022). Genome quality measures, contigN50, and scaffold count, were log transformed prior to

analysis. R-scripts for statistical analysis and plotting are available on GitHub at

https://doi.org/10.5281/zenodo.7679280.

RESULTS

Genome quality

A total of 51 nuclear amphibian genome assemblies were available for our study and were generated

with a variety of sequencing technologies, including Illumina (NextSeq, HiSeq), PacBio (RS11, Sequel),

and Oxford Nanopore. Sequenced genomes represented 25 of 73 amphibian families with reference

genomes distributed unevenly across the phylogeny (Fig. 1). For example, there are only two

salamander genomes representing the 798 extant species, no genomes representing anuran families

such as Leiopelmatidae or Hyperoliidae, yet there are seven Ranidae and six Pipidae genomes (Fig. 1).

Genome assembly length ranged from 0.48 Gb in *Scaphiopus couchii* to 28.21 Gb in *Ambystoma*

mexicanum and was strongly positively associated with c-value estimates of genome size (F_{49} = 330.5,

$p < 1 \times 10^{-15}$) (Table 1, Fig. S1). Twenty-eight of these genomes were assembled to the chromosome

level of which the percentage of the genome assigned to chromosomes ranged from 63.88 to 99.99%

(Table 1). Percentage of the genome assigned to chromosomes was positively associated with contig

N50 (F_{26} = 8.6, p = 0.007) and read length ($t_{29.2}$ = 3.07, p = 0.005) and negatively associated with the

number of scaffolds (F_{26} = 25.2, p < 0.00001). There are additionally mitochondrial genome

assemblies for 345 species of which 17 had nuclear reference genomes. Eleven of the species with

genomes had proteomics data (Table S1).

The quality of the amphibian genomes varied considerably (Table 1). Genomes generated with short-

read technologies were of lower quality than long-read or hybrid genome assemblies as indicated by

significantly lower contig N50s ($F_{2,48}$ = 26.91, p < 10^{-6}), percentage of complete Benchmarking

Universal Single-Copy Ortholog (BUSCO) genes (Fig. S3; $F_{2,48}$ = 10.52, p < 0.001), and higher scaffold

numbers ($F_{2,48}$ = 15.8, p < 10^{-5}).

Contig N50 ranged from 362 bp in *S. couchii* to 45.59 Mb in *Pleurodeles waltl* with a median of 611.23

Kb. Scaffold count varied considerably from 17 in *Spea bombifrons* to more than four million in

Bombina variegata with a median of 6.66 Kb (Table 1). Benchmarking Universal Single-Copy

Orthologs (BUSCO) scores ranged from 0.7 to 99.5% completeness (Tables 1, S1; Fig. 2) and were

positively associated with contig N50 (F_{49} = 82.6, p < 10^{-10}; Fig. S2) and scaffold count (F_{49} = 66.04, p <

10^{-8}). Most genomes had low percentages of duplicate BUSCO genes (< 6%), suggesting they may be

diploid except for *Ranitomeya imitator* and the known tetraploid species, *X. laevis and X. borealis*

(Fig. 2) (Tymowska & Fischberg, 1973).

234
Repeat content

Overall identified repeat percentage of the genomes ranged from 23% in *Platyplectrum ornatum* to

82% in *Oophaga sylvatica* and was positively associated with genome size (F_{49} = 13.24, p = 0.0006)

(Tables 1; Fig. S3). Repeat content varied across genomes with the anurans *Pseudophrne corroboree*,

Bombina bombina, and *O. sylvatica* dominated by Long Terminal Repeats (LTRs), the three caecilians

dominated by Long Interspersed Nuclear Elements (LINEs), and many of the ranid and bufonid

anurans dominated by DNA transposons (Fig. 3; Tables S2-S4). Salamander genomes *Ambystoma*

mexicanum and *Pleurodeles waltl* had fewer repeats than might be predicted given their large sizes

(Fig. S3).

The proportion of repeats that could be classified by RepeatMasker ranged from 7.4 % in *P. ornatum*

to 47.8 % in *P. corroboree* (Table S1) and was positively associated with genome quality measures

contigN50 (F_{49} = 23.49, p = 0.001), scaffold count (F_{49} = 8.71, p = 0.005), and percent BUSCO complete

(F_{49} = 10.27, p = 0.002). The ability to classify repeats was also positively associated with read length,

with longer reads resulting in better classification ($t_{35.622}$ = 4.73, p < 0.001).

Ecological correlates of transposable element content

A Bayesian mixed effect modelling approach was employed to examine the relationships between

proportion of transposable elements and environmental variables. Controlling for phylogenetic

relationships (by estimating Pagel's lambda, λ; de Villemereuil & Nakagawa, 2014), including body

size as a covariate (Spearman correlation with transposable element content ⍴= -0.772, p<0.001) and

excluding the three globally invasive species (*Rhinella marina*, *X. laevis*, and *Lithobates catesbeianus*)

our analysis revealed a significant influence (pMCMC = 0.014) of Bio8 (mean temperature of the

wettest quarter) on the proportion of total transposable elements (Figs 4, S5; Table S8). Inclusion of

these three invasive species did not change this relationship (Table S7). Further analysis indicated

that the relationship with Bio8 was not specific to a particular class of transposable elements, such as

retroelements or DNA transposons (Tables S9 and S10). Phylogenetic signal (Pagel's lambda, λ) was

moderate when considering total transposable elements and retroelements (0.555; Table S7) and

increased when we considered retroelements and DNA transposons alone (0.616 and 0.649; Table

S9).

Genome synteny

Genome synteny of BUSCO genes was highly conserved within amphibian orders (caecilians (Fig. S7),

caudates (Fig. S8), and anurans (Fig. S9); but was less conserved across the amphibian orders (Fig. 5,

S6). However, chromosome naming was inconsistent across all taxa (Figs 5, S6-S9). For example, *X.*

tropicalis chr1 is chr12 in *Leptobrachium ailaonicum* (but not *L. leishanense*), chr2 in *Bufo bufo* (but

not *Bufo gargarizans*) (Fig. S9), and most of the chromosomes for the two salamander genomes (Fig.

S8). Orientation of chromosomes was also inconsistent, including between species of the same genus

(e.g., *Bufo, Leptobrachium*) (Fig. S59) and among the three caecilians (Fig. S7). Multiple inversions

were evident including between chr3 of pipids *(Xenopus tropicalis* and *Hymenochirus boettgeri*) and

other anurans (chromosomes 1, 2, 3, 4, or 10), caecilians (chr3 and chr4/5/6), and even within

species of the same genus (chr7 *Bufo gargarizans*, chr 9 *B. bufo*; Figs 5, S7, S9). There was also

evidence of several chromosomal fissions including the separation of chr1 of *Leptobrachium*

leishanense into chr3 and chr6 in *Pyxicephalus adspersus* and into chr3 and chr7 in *Engystomops*

pustulosus; however, this chromosome remained mostly intact in the other anuran genomes (Fig.

S9).

DISCUSSION

In this study we analyzed 51 amphibian reference genomes from the public domain to evaluate their

content and usefulness for functional genetics research (Fig. 1, Table 1). There are considerably

fewer reference genomes for amphibians than exist for birds (N=754), mammals (N=406), and non-

avian reptiles (N=108). This scarcity of reference genomes results in many gaps in genome

representation across the amphibian tree of life including many entirely unrepresented groups and

with only two genomes representing the entire order Caudata (but see Myers & Pyron, 2024). The

unrepresented families include many of interest from a conservation perspective due to their high

number of IUCN RedList Critically Endangered species (e.g., Cryptobranchidae, Plethodontidae,

Strabomantidae, and Craugastoridae) (IUCN, 2022). However, our search of the Genomes on a Tree

(GoaT) database (Sotero-Caio et al., 2021) indicated that there are a further 20 amphibian genome

assemblies in progress (15 anurans, 5 caudates; Table S10) indicating that this resource will be

increasing by more than 40% in the next few years.

The quality and completeness of the genomes in our dataset varied considerably (e.g., Fig. 2). Much

of this variation can be attributed to the sequencing technology used to generate them, with short-

read sequencing approaches resulting in lower completeness and continuity (Fig. S2). These impacts

are a recognized limitation of short-read sequencing and have been reported to impact genome

quality in taxa from insects (Hotaling et al., 2021b) to other vertebrates (Rhie et al., 2021), but have

likely had a disproportionate impact in amphibian genomes due to the difficulty of assembling

genomes with high repeat content (Sun et al., 2020). Fortunately, most ongoing sequencing efforts

now use long-read or hybrid sequencing approaches (i.e., that incorporate scaffolding technologies

such as Hi-C sequencing), which along with improved sequencing algorithms, should result in higher

quality amphibian genomes (Hotaling et al., 2021a; Lawniczak et al., 2022; Rhie et al., 2021).

The variation we report here in genome quality, contiguity, and completeness may impact the value

of the genomes for functional genomics research. However, the improvements in all these measures

seen with the utilization of long read technologies or hybrid assemblies suggests that genome quality

will continue to improve as these approaches are used more frequently. Genome quality (i.e., high

continuity, contiguity, accuracy, completeness (Rhie et al., 2021)) are critical for applications such as

quantitative genetics where assembly errors can lead to incorrect inferences in genetic association or

genetic prediction. Quality also enhances the usefulness of genomes. For example, highly contiguous

chromosome-level assemblies decrease computational requirements for downstream analyses such

as mapping, variant calling, and alignment (Aganezov et al., 2022).

One of the most intriguing features of amphibian genomes is the huge range they exhibit in size

(Biscotti et al., 2019). This was exemplified in our dataset where assembly length ranged from 0.48

Gb in *Scaphiopus couchii* to 28 Gb in *Ambystoma mexicanum*. Why gigantic genomes exist in some

species, but not others, remains a key evolutionary question (Kapusta et al., 2017; Wang et al., 2021).

Explanations include differences in genome-level processes (e.g., insertion and deletion rates)

(Frahry et al., 2015; Sun et al., 2012b), development (e.g., developmental rate and complexity)

(Gregory, 2002; Liedtke et al., 2018), physiology (e.g., water loss) (Johnson et al., 2021), body size

(e.g. miniaturization) (Decena-Segarra et al., 2020), and demography (e.g., effective population size)

(Liedtke et al., 2018; Lynch & Walsh, 2007) (but see Mohlhenrich & Mueller, 2016). As more

amphibian genomes become available, these hypotheses can be more rigorously evaluated.

We report some of the largest estimates of repeat content of any vertebrate (82% in *Oophaga*

sylvatica and 77% in *Rana muscosa*), exceeded only by the Australian lungfish at 90% (Meyer et al.,

2021). As expected, genome size was correlated with repeat content affirming that much of the

variation in amphibian genome size is due to an excess of repeats and transposable elements rather

than coding regionds (Biscotti et al., 2019; Lamichhaney et al., 2021; Zuo et al., 2023).

In contrast to mammals, whose repeat landscape is mainly dominated by LTR retrotransposons (Platt

et al., 2018), amphibian repeat content varied considerably with some species dominated by DNA

transposons (as previously reported (Suda et al., 2022; Zuo et al., 2023), and others by non-LTR

retrotransposons including the three caecilian genomes which were dominated by LINEs. This agrees

with genomic data and transcriptomic data from the caecilian *Ichthyophis bannanicus*, where LINEs

were the second most abundant type of repeat (26% of the genome) behind Dictyostelium

intermediate repeat sequences (DIRS) (30%) (Wang et al., 2021); this is a similar percentage of LINES

to what we report in the three caecilian genomes in this study (19 to 26%) (Table S4).

These disparities in repeat percentage and content likely reflect differing evolutionary histories

among species, as indicated by three of the four congeneric species pairs in our dataset having

similar values (i.e., *Bufo*, *Leptobrachium,* and *Xenopus*; but not *Oophaga*). The differences we

observed in *Oophaga pumilio* and *O. sylvatica* are likely due to assembly quality rather than genome

content given that these two genomes were sequenced with different technologies and have

dramatically different genome qualities (e.g., contig N50s of 5.8 vs. 97.8 Kbp respectively).

A considerable proportion of the repeats could not be classified. This was likely due to incorrect

classification (e.g., genes categorized as repeats) and the lack of good amphibian-specific repeat

resources (Ou et al., 2019) for classification via nucleotide sequence homology. The majority of

amphibian curated repeat libraries are generated in reference to *Xenopus* species (e.g., Dfam); the

large divergence times of this genus from the other amphibian species suggests that it may be a

contributing factor to the lack of classification. However, we also report many unclassified repeats in

the two *Xenopus* genomes.

The largest genomes in our dataset from caudates, *A. mexicanum and P. waltl*, had fewer repeats

than predicted given their size (Fig S4) (Nowoshilow et al., 2018). This may be due, in part, to the

Dfam (Storer et al., 2021) library used for repeat annotation being anuran-based; however, we did

not observe this trend in the three caecilian genomes in our dataset. Also, we performed *de novo*

annotation of these genomes, which should have captured repetitive elements missing from Dfam.

More likely, this low number of repeats reflects low deletion rates and, thus, persistence of repeats

in the genome for extremely long periods of time, leading to their mutational decay into unique

sequences whose repetitive origin is obscured (Frahry et al., 2015; Keinath et al., 2015; Novák et al.,

2020; Sun et al., 2012a).

We also show that amphibian species that inhabit warm climates particularly during months with

high precipitation have a greater proportion of transposable elements. This observed trend does not

appear to be driven by a specific group of transposons suggesting it may be caused by climatic

factors. Recent studies indicate that transposable elements exhibit greater activity in hotter climates

(Baduel et al., 2021) with an increasing number of studies suggesting increased transposable element

activity contributes to genetic diversification and facilitates species adaptation (Li et al., 2018;

Schrader & Schmitz, 2019; Stapley et al., 2015). The pattern observed here likewise suggests the

potential for heightened transposable element activity and may help explain transposable element

accumulation and potentially the higher evolutionary rates observed in the genomes of tropical

amphibians (Pyron & Wiens, 2013).

Our study is the first to examine chromosomal synteny across all amphibian orders. We show that

overall synteny of amphibian genomes is relatively conserved, particularly within orders (Figs 5 and

S7). This aligns with previous results from anurans that reported conserved genome organization in

this group (Bredeson et al., 2021; Wu et al., 2022). However, chromosome content and number

varied across species, which seems to have been driven by multiple occurrences of chromosomal

fusions and fissions (e.g., Fig. 5). Chromosomal rearrangements have occurred throughout vertebrate

evolution, including the hypothesized fusion of microchromosomes in the ancestor of tetrapods to

create the larger macrochromosomes seen in amphibians and mammals and their subsequent fission

to create the microchromosomes of modern birds and non-avian reptiles (Waters et al., 2021).

Some of the structural rearrangements we detected may be due to assembly errors and should be

evaluated in future assemblies using long-read scaffolding approaches (e.g., Oxford nanopore

sequencing), chromosome conformation capture technologies (e.g., Hi-C), or chromosome mapping

approaches (e.g., FISH). We also identified incongruities with chromosome naming and orientation

caused by differences in assembly methods. These were apparent even within species of the same

genus (e.g., *Bufo*). We suggest potential revisions of existing genome annotations to improve

congruity and that future assemblies are curated consistently against high-quality reference genomes

(e.g., *Xenopus laevis*).

Conclusions

New sequencing technologies and assembly algorithms have resulted in a good number of genomes

for comparative analyses spanning the amphibian phylogeny. This has already begun to yield

important insights on the evolution (Lamichhaney et al., 2021; Wu et al., 2022), development

(Schloissnig et al., 2021; Stuckert et al., 2021), sex determination (Hime et al., 2019; Ma & Veltsos,

2021), and unique features (Fischer et al., 2019; Nowoshilow et al., 2018; Seidl et al., 2019) of this

interesting group of animals.

The increased availability of amphibian genomes can also aid conservation efforts in this highly

threatened group by facilitating research on genome-wide functional diversity, which can be used to

inform management decisions such as genetic rescue or targeted genetic intervention for species

threatened by habitat loss or chytridiomycosis (Chestnut et al., 2014; Kosch et al., 2022).

Additionally, well-annotated genomes can be used to create eDNA assays for population monitoring

(Breton et al., 2022; Saeed et al., 2022).

Future research efforts should focus on generating more reference genomes to fill the gaps in the

amphibian phylogeny and the identification of advantageous genetic traits against threats. Efforts

should also be made to increase the quality of genomes and expand transcriptome and annotation

databases. We suggest that these efforts strive to follow the recommendations of initiatives such as

the Earth BioGenome Project (Lawniczak et al., 2022), the Darwin Tree of Life Project (Blaxter et al.,

2022), and the Threatened Species Initiative (Hogg et al., 2022) to sequence at least one

representative from each family to ensure taxonomic coverage. Species selection should prioritize

species of interest for understanding valuable functional genetic traits; for example, for the purpose

of immunological research to understand disease resistance, or for conservation purposes to

enhance fitness.

ACKNOWLEDGEMENTS

T.A.K, N.D.Y, and L.F.S research were supported by The University of Melbourne's Research

Computing Services and the Petascale Campus Initiative. T.A.K. and L.F.S were supported by

Australian Research Council Grants (FT190100462, LP200301370). K.W.V and M.L.P were funded by

the European Union (ERC, MolStressH2O, 101044202). Views and opinions expressed are however

those of the author(s) only and do not necessarily reflect those of the European Union or the

European Research Council Executive Agency. Neither the European Union nor the granting authority

can be held responsible for them. We are grateful to A. Stuckert, N. Brajuka, J. Sproul, and E. Tescari

for their advice on repeat modelling, and J.T. Li for providing suggestions on synteny analyses. We

thank M. Moore for exploratory repeat analyses.

REFERENCES
Aganezov, S., Yan, S. M., Soto, D. C., Kirsche, M., Zarate, S., Avdeyev, P., Taylor, D. J., Shafin,
434   K., Shumate, A., Xiao, C., Wagner, J., McDaniel, J., Olson, N. D., Sauria, M. E. G.,
435   Vollger, M. R., Rhie, A., Meredith, M., Martin, S., Lee, J., Koren, S., Rosenfeld, J. A.,
436   Paten, B., Layer, R., Chin, C. S., Sedlazeck, F. J., Hansen, N. F., Miller, D. E., Phillippy, A.
437   M., Miga, K. H., McCoy, R. C., Dennis, M. Y., Zook, J. M., & Schatz, M. C. (2022). A
438   complete reference genome improves analysis of human genetic variation. *Science,*
439   *376*(6588), eabl3533. doi:10.1126/science.abl3533
440   Baduel, P., Leduque, B., Ignace, A., Gy, I., Gil Jr, J., Loudet, O., Colot, V., & Quadrana, L.
441   (2021). Genetic and environmental modulation of transposition shapes the
442   evolutionary potential of *Arabidopsis thaliana*. *Genome Biology, 22*(1), 138.
443   Biscotti, M. A., Carducci, F., Olmo, E., & Canapa, A. (2019). Vertebrate Genome Size and the
444   Impact of Transposable Elements in Genome Evolution. In P. Pontarotti (Ed.),
445   *Evolution, Origin of Life, Concepts and Methods* (pp. 233-251). Cham: Springer
446   International Publishing.
447   Blaxter, M., Archibald, J. M., Childers, A. K., Coddington, J. A., Crandall, K. A., Di Palma, F.,
448   Durbin, R., Edwards, S. V., Graves, J. A. M., Hackett, K. J., Hall, N., Jarvis, E. D.,
449   Johnson, R. N., Karlsson, E. K., Kress, W. J., Kuraku, S., Lawniczak, M. K. N., Lindblad-
450   Toh, K., Lopez, J. V., Moran, N. A., Robinson, G. E., Ryder, O. A., Shapiro, B., Soltis, P.
451   S., Warnow, T., Zhang, G., & Lewin, H. A. (2022). Why sequence all eukaryotes?

Proceedings of the National Academy of Sciences, 119(4), e2115636118.
doi:10.1073/pnas.2115636118

Bredeson, J. V., Mudd, A. B., Medina-Ruiz, S., Mitros, T., Smith, O. K., Miller, K. E., Lyons, J. B.,
Batra, S. S., Park, J., Berkoff, K. C., Plott, C., Grimwood, J., Schmutz, J., Aguirre-
Figueroa, G., Khokha, M. K., Lane, M., Philipp, I., Laslo, M., Hanken, J., Kerdivel, G.,
Buisine, N., Sachs, L. M., Buchholz, D. R., Kwon, T., Smith-Parker, H., Gridi-Papp, M.,
Ryan, M. J., Denton, R. D., Malone, J. H., Wallingford, J. B., Straight, A. F., Heald, R.,
Hockemeyer, D., Harland, R. M., & Rokhsar, D. S. (2021). Conserved chromatin and
repetitive patterns reveal slow genome evolution in frogs. *bioRxiv*,
2021.2010.2018.464293. doi:10.1101/2021.10.18.464293

Breton, B.-A. A., Beaty, L., Bennett, A. M., Kyle, C. J., Lesbarrères, D., Vilaça, S. T., Wikston,
M. J. H., Wilson, C. C., & Murray, D. L. (2022). Testing the effectiveness of
environmental DNA (eDNA) to quantify larval amphibian abundance. *Environmental*
DNA, 4(6), 1229-1240. doi:10.1002/edn3.332

Brown, J. L., Morales, V., & Summers, K. (2010). A key ecological trait drove the evolution of
biparental care and monogamy in an amphibian. *The American Naturalist, 175*(4),
436-446.

Casacuberta, E., & González, J. (2013). The impact of transposable elements in
environmental adaptation. *Molecular Ecology, 22*(6), 1503-1517.
doi:https://doi.org/10.1111/mec.12170

Chestnut, T., Anderson, C., Popa, R., Blaustein, A. R., Voytek, M., Olson, D. H., & Kirshtein, J.
(2014). Heterogeneous Occupancy and Density Estimates of the Pathogenic Fungus
Batrachochytrium dendrobatidis in Waters of North America. *PLOS ONE, 9*(9),
e106790. doi:10.1371/journal.pone.0106790

Cunningham, F., Allen, J. E., Allen, J., Alvarez-Jarreta, J., Amode, M. R., Armean, I. M.,
Austine-Orimoloye, O., Azov, A. G., Barnes, I., Bennett, R., Berry, A., Bhai, J., Bignell,
A., Billis, K., Boddu, S., Brooks, L., Charkhchi, M., Cummins, C., Da Rin Fioretto, L.,
Davidson, C., Dodiya, K., Donaldson, S., El Houdaigui, B., El Naboulsi, T., Fatima, R.,
Giron, C. G., Genez, T., Martinez, J. G., Guijarro-Clarke, C., Gymer, A., Hardy, M.,
Hollis, Z., Hourlier, T., Hunt, T., Juettemann, T., Kaikala, V., Kay, M., Lavidas, I., Le, T.,
Lemos, D., Marugán, J. C., Mohanan, S., Mushtaq, A., Naven, M., Ogeh, D. N., Parker,
A., Parton, A., Perry, M., Piližota, I., Prosovetskaia, I., Sakthivel, M. P., Salam, A. I. A.,
Schmitt, B. M., Schuilenburg, H., Sheppard, D., Pérez-Silva, J. G., Stark, W., Steed, E.,
Sutinen, K., Sukumaran, R., Sumathipala, D., Suner, M. M., Szpak, M., Thormann, A.,
Tricomi, F. F., Urbina-Gómez, D., Veidenberg, A., Walsh, T. A., Walts, B., Willhoft, N.,
Winterbottom, A., Wass, E., Chakiachvili, M., Flint, B., Frankish, A., Giorgetti, S.,
Haggerty, L., Hunt, S. E., GR, I. I., Loveland, J. E., Martin, F. J., Moore, B., Mudge, J. M.,
Muffato, M., Perry, E., Ruffier, M., Tate, J., Thybert, D., Trevanion, S. J., Dyer, S.,
Harrison, P. W., Howe, K. L., Yates, A. D., Zerbino, D. R., & Flicek, P. (2022). Ensembl
2022. *Nucleic Acids Research, 50*(D1), D988-d995. doi:10.1093/nar/gkab1049

Daly, J. W., Garraffo, H. M., Spande, T. F., Decker, M. W., Sullivan, J. P., & Williams, M.
(2000). Alkaloids from frog skin: the discovery of epibatidine and the potential for
developing novel non-opioid analgesics. *Natural product reports, 17*(2), 131-135.

De Angelis, M., Casciaro, B., Genovese, A., Brancaccio, D., Marcocci, M., Novellino, E.,
Carotenuto, A., Palamara, A., Mangoni, M., & Nencioni, L. (2021). Temporin G, an

amphibian antimicrobial peptide against influenza and parainfluenza respiratory
viruses: insights into biological activity and mechanism of action. *The FASEB Journal,*
35(2), e21358.
Decena-Segarra, L. P., Bizjak-Mali, L., Kladnik, A., Sessions, S. K., & Rovito, S. M. (2020).
Miniaturization, genome size, and biological size in a diverse clade of salamanders.
The American Naturalist, 196(5), 634-648. doi:10.5061/dryad.ht76hdrcg
Drummond, A. J., & Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling
trees. *BMC Evolutionary Biology, 7*, 214. doi:10.1186/1471-2148-7-214
Duellman, W. E. (1999). *Patterns of distribution of amphibians: a global perspective*: JHU
Press.
Elewa, A., Wang, H., Talavera-López, C., Joven, A., Brito, G., Kumar, A., Hameed, L. S.,
Penrad-Mobayed, M., Yao, Z., & Zamani, N. (2017). Reading and editing the
Pleurodeles waltl genome reveals novel features of tetrapod regeneration. *Nature*
Communications, 8(1), 1-9. doi:10.1038/s41467-017-01964-9
Emms, D. M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for
comparative genomics. *Genome Biology, 20*(1), 1-14.
Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap.
Evolution, 39(4), 783-791.
Fischer, E. K., Roland, A. B., Moskowitz, N. A., Tapia, E. E., Summers, K., Coloma, L. A., &
O'Connell, L. A. (2019). The neural basis of tadpole transport in poison frogs.
Proceedings of the Royal Society B: Biological Sciences, 286(1907), 20191084.
doi:10.1098/rspb.2019.1084
Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F. (2020).
RepeatModeler2 for automated genomic discovery of transposable element families.
Proceedings of the National Academy of Sciences, 117(17), 9451-9457.
doi:10.1073/pnas.1921046117
Frahry, M. B., Sun, C., Chong, R. A., & Mueller, R. L. (2015). Low levels of LTR retrotransposon
deletion by ectopic recombination in the gigantic genomes of salamanders. *Journal of*
Molecular Evolution, 80, 120-129. doi:10.1007/s00239-014-9663-7
Garland Jr, T., Harvey, P. H., & Ives, A. R. (1992). Procedures for the analysis of comparative
data using phylogenetically independent contrasts. *Systematic Biology, 41*(1), 18-32.
Gregory, T. R. (2002). Genome size and developmental complexity. *Genetica, 115*(1), 131-
146. doi:10.1023/A:1016032400147
Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models:
the MCMCglmm R package. *Journal of statistical software, 33*, 1-22.
Hime, P. M., Briggler, J. T., Reece, J. S., & Weisrock, D. W. (2019). Genomic Data Reveal
Conserved Female Heterogamety in Giant Salamanders with Gigantic Nuclear
Genomes. *G3: Genes|Genomes|Genetics*, g3.400556.402019.
doi:10.1534/g3.119.400556
Hogg, C. J., Ottewell, K., Latch, P., Rossetto, M., Biggs, J., Gilbert, A., Richmond, S., & Belov, K.
(2022). Threatened Species Initiative: Empowering conservation action using genomic
resources. *Proceedings of the National Academy of Sciences, 119*(4), e2115643118.
doi:10.1073/pnas.2115643118

Hotaling, S., Kelley, J. L., & Frandsen, P. B. (2021a). Toward a genome sequence for every
animal: Where are we now? *Proceedings of the National Academy of Sciences,*
118(52), e2109019118. doi:10.1073/pnas.2109019118
Hotaling, S., Sproul, J. S., Heckenhauer, J., Powell, A., Larracuente, A. M., Pauls, S. U., Kelley,
J. L., & Frandsen, P. B. (2021b). Long-reads are revolutionizing 20 years of insect
genome sequencing. *Genome Biology and Evolution*. doi:10.1093/gbe/evab138
IUCN. (2022). The IUCN Red List of Threatened Species. Version 2022-2. Accessed on [6
January 2023].   https://www.iucnredlist.org.
Jetz, W., & Pyron, R. A. (2018). The interplay of past diversification and evolutionary isolation
with present imperilment across the amphibian tree of life. *Nature Ecology &*
Evolution, 2(5), 850-858. doi:10.1038/s41559-018-0515-5
Johnson, B. B., Searle, J. B., & Sparks, J. P. (2021). Genome size influences adaptive plasticity
of water loss, but not metabolic rates, in lungless salamanders. *Journal of*
Experimental Biology, 224(8), jeb242196. doi:10.1242/jeb.242196
Kapusta, A., Suh, A., & Feschotte, C. (2017). Dynamics of genome size evolution in birds and
mammals. *Proceedings of the National Academy of Sciences, 114*(8), E1460-E1469.
doi:10.1073/pnas.1616702114
Keinath, M. C., Timoshevskiy, V. A., Timoshevskaya, N. Y., Tsonis, P. A., Voss, S. R., & Smith, J.
J. (2015). Initial characterization of the large genome of the salamander *Ambystoma*
mexicanum using shotgun and laser capture chromosome sequencing. *Scientific*
Reports, 5(1), 16413. doi:10.1038/srep16413
Kosch, T. A., Waddle, A. W., Cooper, C. A., Zenger, K. R., Garrick, D. J., Berger, L., & Skerratt,
L. F. (2022). Genetic approaches for increasing fitness in endangered species. *Trends*
in Ecology & Evolution, 37(4), 332-345. doi:10.1016/j.tree.2021.12.003
Kriventseva, E. V., Kuznetsov, D., Tegenfeldt, F., Manni, M., Dias, R., Simão, F. A., & Zdobnov,
E. M. (2018). OrthoDB v10: sampling the diversity of animal, plant, fungal, protist,
bacterial and viral genomes for evolutionary and functional annotations of orthologs.
Nucleic Acids Research, 47(D1), D807-D811. doi:10.1093/nar/gky1053
Lamichhaney, S., Catullo, R., Keogh, J. S., Clulow, S., Edwards, S. V., & Ezaz, T. (2021). A bird-
like genome from a frog: Mechanisms of genome size reduction in the ornate
burrowing frog, *Platyplectrum ornatum. Proceedings of the National Academy of*
Sciences, 118(11), e2011649118. doi:10.1073/pnas.2011649118
Lawniczak, M. K. N., Durbin, R., Flicek, P., Lindblad-Toh, K., Wei, X., Archibald, J. M., Baker,
W. J., Belov, K., Blaxter, M. L., Marques Bonet, T., Childers, A. K., Coddington, J. A.,
Crandall, K. A., Crawford, A. J., Davey, R. P., Di Palma, F., Fang, Q., Haerty, W., Hall, N.,
Hoff, K. J., Howe, K., Jarvis, E. D., Johnson, W. E., Johnson, R. N., Kersey, P. J., Liu, X.,
Lopez, J. V., Myers, E. W., Pettersson, O. V., Phillippy, A. M., Poelchau, M. F., Pruitt, K.
D., Rhie, A., Castilla-Rubio, J. C., Sahu, S. K., Salmon, N. A., Soltis, P. S., Swarbreck, D.,
Thibaud-Nissen, F., Wang, S., Wegrzyn, J. L., Zhang, G., Zhang, H., Lewin, H. A., &
Richards, S. (2022). Standards recommendations for the Earth BioGenome Project.
Proceedings of the National Academy of Sciences, 119(4), e2115639118.
doi:10.1073/pnas.2115639118
Lee, B. T., Barber, G. P., Benet-Pagès, A., Casper, J., Clawson, H., Diekhans, M., Fischer, C.,
Gonzalez, J. N., Hinrichs, A. S., Lee, C. M., Muthuraman, P., Nassar, L. R., Nguy, B.,
Pereira, T., Perez, G., Raney, B. J., Rosenbloom, K. R., Schmelter, D., Speir, M. L., Wick,

B. D., Zweig, A. S., Haussler, D., Kuhn, R. M., Haeussler, M., & Kent, W. J. (2022). The
UCSC Genome Browser database: 2022 update. *Nucleic Acids Research, 50*(D1),
D1115-d1122. doi:10.1093/nar/gkab959
Li, Z.-W., Hou, X.-H., Chen, J.-F., Xu, Y.-C., Wu, Q., González, J., & Guo, Y.-L. (2018).
Transposable elements contribute to the adaptation of *Arabidopsis thaliana*. *Genome
Biology and Evolution, 10*(8), 2140-2150. doi:10.1093/gbe/evy171
Liedtke, H. C., Cruz, F., Gómez-Garrido, J., Fuentes Palacios, D., Marcet-Houben, M., Gut, M.,
Alioto, T., Gabaldón, T., & Gomez-Mestre, I. (2022). Chromosome-level assembly,
annotation and phylome of *Pelobates cultripes*, the western spadefoot toad. *DNA
Research, 29*(3), dsac013.
Liedtke, H. C., Gower, D. J., Wilkinson, M., & Gomez-Mestre, I. (2018). Macroevolutionary
shift in the size of amphibian genomes and the role of life history and climate. *Nature
Ecology & Evolution, 2*(11), 1792-1799. doi:10.1038/s41559-018-0674-4
Liu, Y., Shi, D., Wang, J., Chen, X., Zhou, M., Xi, X., Cheng, J., Ma, C., Chen, T., & Shaw, C.
(2020). A novel amphibian antimicrobial peptide, phylloseptin-PV1, exhibits effective
anti-staphylococcal activity without inducing either hepatic or renal toxicity in mice.
Frontiers in Microbiology, 11, 565158.
Lovell, J. T., Sreedasyam, A., Schranz, M. E., Wilson, M., Carlson, J. W., Harkess, A., Emms, D.,
Goodstein, D. M., & Schmutz, J. (2022). GENESPACE tracks regions of interest and
gene copy number variation across multiple genomes. *eLife, 11*, e78526.
doi:10.7554/eLife.78526
Lynch, M., & Walsh, B. (2007). *The origins of genome architecture*. Sunderland: Sinauer
Associates.
Ma, W.-J., & Veltsos, P. (2021). The Diversity and Evolution of Sex Chromosomes in Frogs.
Genes, 12(4), 483. doi:10.3390/genes12040483
Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO
Update: Novel and Streamlined Workflows along with Broader and Deeper
Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes.
Molecular Biology and Evolution, 38(10), 4647-4654. doi:10.1093/molbev/msab199
Meyer, A., Schloissnig, S., Franchini, P., Du, K., Woltering, J. M., Irisarri, I., Wong, W. Y.,
Nowoshilow, S., Kneitz, S., Kawaguchi, A., Fabrizius, A., Xiong, P., Dechaud, C., Spaink,
H. P., Volff, J.-N., Simakov, O., Burmester, T., Tanaka, E. M., & Schartl, M. (2021).
Giant lungfish genome elucidates the conquest of land by vertebrates. *Nature,
590*(7845), 284-289. doi:10.1038/s41586-021-03198-8
Mohlhenrich, E. R., & Mueller, R. L. (2016). Genetic drift and mutational hazard in the
evolution of salamander genomic gigantism. *Evolution, 70*(12), 2865-2878.
Myers, E. A., & Pyron, R. A. (2024). The first complete assembly for a lungless urodelan with
a "miniaturized" genome, the Northern Dusky Salamander (Plethodontidae:
Desmognathus fuscus). *bioRxiv*, 2024.2004.2030.591895.
doi:10.1101/2024.04.30.591895
Novák, P., Guignard, M. S., Neumann, P., Kelly, L. J., Mlinarec, J., Koblížková, A., Dodsworth,
S., Kovařík, A., Pellicer, J., Wang, W., Macas, J., Leitch, I. J., & Leitch, A. R. (2020).
Repeat-sequence turnover shifts fundamentally in species with large genomes.
Nature Plants, 6(11), 1325-1329. doi:10.1038/s41477-020-00785-x

Nowoshilow, S., Schloissnig, S., Fei, J.-F., Dahl, A., Pang, A. W. C., Pippel, M., Winkler, S.,
Hastie, A. R., Young, G., Roscito, J. G., Falcon, F., Knapp, D., Powell, S., Cruz, A., Cao,
H., Habermann, B., Hiller, M., Tanaka, E. M., & Myers, E. W. (2018). The axolotl
genome and the evolution of key tissue formation regulators. *Nature*.
doi:10.1038/nature25458
https://www.nature.com/articles/nature25458#supplementary-information
Nunes-de-Almeida, C. H. L., Haddad, C. F. B., & Toledo, L. F. (2021). A revised classification of
the amphibian reproductive modes. *Salamandra, 57*(3), 413-427.
O'Leary, N. A., Wright, M. W., Brister, J. R., Ciufo, S., Haddad, D., McVeigh, R., Rajput, B.,
Robbertse, B., Smith-White, B., Ako-Adjei, D., Astashyn, A., Badretdin, A., Bao, Y.,
Blinkova, O., Brover, V., Chetvernin, V., Choi, J., Cox, E., Ermolaeva, O., Farrell, C. M.,
Goldfarb, T., Gupta, T., Haft, D., Hatcher, E., Hlavina, W., Joardar, V. S., Kodali, V. K.,
Li, W., Maglott, D., Masterson, P., McGarvey, K. M., Murphy, M. R., O'Neill, K., Pujar,
S., Rangwala, S. H., Rausch, D., Riddick, L. D., Schoch, C., Shkeda, A., Storz, S. S., Sun,
H., Thibaud-Nissen, F., Tolstoy, I., Tully, R. E., Vatsan, A. R., Wallin, C., Webb, D., Wu,
W., Landrum, M. J., Kimchi, A., Tatusova, T., DiCuccio, M., Kitts, P., Murphy, T. D., &
Pruitt, K. D. (2016). Reference sequence (RefSeq) database at NCBI: current status,
taxonomic expansion, and functional annotation. *Nucleic Acids Research, 44*(D1),
D733-745. doi:10.1093/nar/gkv1189
Ou, S., Su, W., Liao, Y., Chougule, K., Agda, J. R. A., Hellinga, A. J., Lugo, C. S. B., Elliott, T. A.,
Ware, D., Peterson, T., Jiang, N., Hirsch, C. N., & Hufford, M. B. (2019). Benchmarking
transposable element annotation methods for creation of a streamlined,
comprehensive pipeline. *Genome Biology, 20*(1), 275. doi:10.1186/s13059-019-1905-
y
Paiola, M., Dimitrakopoulou, D., Pavelka, M. S., & Robert, J. (2023). Amphibians as a model
to study the role of immune cell heterogeneity in host and mycobacterial
interactions. *Developmental & Comparative Immunology, 139*, 104594.
doi:https://doi.org/10.1016/j.dci.2022.104594
Pappalardo, A. M., Ferrito, V., Biscotti, M. A., Canapa, A., & Capriglione, T. (2021).
Transposable Elements and Stress in Vertebrates: An Overview. *Int J Mol Sci, 22*(4).
doi:10.3390/ijms22041970
Pimpinelli, S., & Piacentini, L. (2020). Environmental change and the evolution of genomes:
Transposable elements as translators of phenotypic plasticity into genotypic
variability. *Functional Ecology, 34*(2), 428-441.
Platt, R. N., 2nd, Vandewege, M. W., & Ray, D. A. (2018). Mammalian transposable elements
and their impacts on genome evolution. *Chromosome Res, 26*(1-2), 25-43.
doi:10.1007/s10577-017-9570-z
Pyron, R. A., Pirro, S., Hains, T., Colston, T. J., Myers, E. A., O'Connell, K. A., & Beamer, D. A.
(2024). The Draft Genome Sequences of 50 Salamander species (Caudata, Amphibia).
Biodiversity Genomes, 2024. doi:10.56179/001c.116891
Pyron, R. A., & Wiens, J. J. (2013). Large-scale phylogenetic analyses reveal the causes of high
tropical amphibian diversity. *Proceedings of the Royal Society B: Biological Sciences,*
280(1770), 20131622.
Revell, L. J. (2024). phytools 2.0: an updated R ecosystem for phylogenetic comparative
methods (and other things). *PeerJ, 12*, e16505. doi:10.7717/peerj.16505

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G. L., Cantin, L. J., Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M., Haase, B., Mountcastle, J., Winkler, S., Paez, S., Howard, J., Vernes, S. C., Lama, T. M., Grutzner, F., Warren, W. C., Balakrishnan, C. N., Burt, D., George, J. M., Biegler, M. T., Iorns, D., Digby, A., Eason, D., Robertson, B., Edwards, T., Wilkinson, M., Turner, G., Meyer, A., Kautt, A. F., Franchini, P., Detrich, H. W., Svardal, H., Wagner, M., Naylor, G. J. P., Pippel, M., Malinsky, M., Mooney, M., Simbirsky, M., Hannigan, B. T., Pesout, T., Houck, M., Misuraca, A., Kingan, S. B., Hall, R., Kronenberg, Z., Sović, I., Dunn, C., Ning, Z., Hastie, A., Lee, J., Selvaraj, S., Green, R. E., Putnam, N. H., Gut, I., Ghurye, J., Garrison, E., Sims, Y., Collins, J., Pelan, S., Torrance, J., Tracey, A., Wood, J., Dagnew, R. E., Guan, D., London, S. E., Clayton, D. F., Mello, C. V., Friedrich, S. R., Lovell, P. V., Osipova, E., Al-Ajli, F. O., Secomandi, S., Kim, H., Theofanopoulou, C., Hiller, M., Zhou, Y., Harris, R. S., Makova, K. D., Medvedev, P., Hoffman, J., Masterson, P., Clark, K., Martin, F., Howe, K., Flicek, P., Walenz, B. P., Kwak, W., Clawson, H., Diekhans, M., Nassar, L., Paten, B., Kraus, R. H. S., Crawford, A. J., Gilbert, M. T. P., Zhang, G., Venkatesh, B., Murphy, R. W., Koepfli, K.-P., Shapiro, B., Johnson, W. E., Di Palma, F., Marques-Bonet, T., Teeling, E. C., Warnow, T., Graves, J. M., Ryder, O. A., Haussler, D., O'Brien, S. J., Korlach, J., Lewin, H. A., Howe, K., Myers, E. W., Durbin, R., Phillippy, A. M., & Jarvis, E. D. (2021). Towards complete and error-free genome assemblies of all vertebrate species. *Nature, 592*(7856), 737-746. doi:10.1038/s41586-021-03451-0

Robert, J. (2020). Experimental platform using the amphibian *Xenopus laevis* for research in fundamental and medical immunology. *Cold Spring Harbor Protocols, 2020*(7), pdb. top106625.

Saeed, M., Rais, M., Akram, A., Williams, M. R., Kellner, K. F., Hashsham, S. A., & Davis, D. R. (2022). Development and validation of an eDNA protocol for monitoring endemic Asian spiny frogs in the Himalayan region of Pakistan. *Scientific Reports, 12*(1), 5624. doi:10.1038/s41598-022-09084-1

Scheele, B. C., Hunter, D. A., Grogan, L. F., Berger, L., Kolby, J. E., McFadden, M. S., Marantelli, G., Skerratt, L. F., & Driscoll, D. A. (2014). Interventions for Reducing Extinction Risk in Chytridiomycosis-Threatened Amphibians. *Conservation Biology, 28*(5), 1195-1205. doi:10.1111/cobi.12322

Scheele, B. C., Pasmans, F., Skerratt, L. F., Berger, L., Martel, A., Beukema, W., Acevedo, A. A., Burrowes, P. A., Carvalho, T., Catenazzi, A., De la Riva, I., Fisher, M. C., Flechas, S. V., Foster, C. N., Frías-Álvarez, P., Garner, T. W. J., Gratwicke, B., Guayasamin, J. M., Hirschfeld, M., Kolby, J. E., Kosch, T. A., La Marca, E., Lindenmayer, D. B., Lips, K. R., Longo, A. V., Maneyro, R., McDonald, C. A., Mendelson, J., Palacios-Rodriguez, P., Parra-Olea, G., Richards-Zawacki, C. L., Rödel, M.-O., Rovito, S. M., Soto-Azat, C., Toledo, L. F., Voyles, J., Weldon, C., Whitfield, S. M., Wilkinson, M., Zamudio, K. R., & Canessa, S. (2019). Amphibian fungal panzootic causes catastrophic and ongoing loss of biodiversity. *Science, 363*(6434), 1459-1463. doi:10.1126/science.aav0379

Schloissnig, S., Kawaguchi, A., Nowoshilow, S., Falcon, F., Otsuki, L., Tardivo, P., Timoshevskaya, N., Keinath, M. C., Smith, J. J., & Voss, S. R. (2021). The giant axolotl genome uncovers the evolution, scaling, and transcriptional control of complex gene loci. *Proceedings of the National Academy of Sciences, 118*(15).

Schrader, L., & Schmitz, J. (2019). The impact of transposable elements in adaptive
evolution. *Molecular Ecology, 28*(6), 1537-1549.
Schulte, L. M., Ringler, E., Rojas, B., & Stynoski, J. L. (2020). Developments in Amphibian
Parental Care Research: History, Present Advances, and Future Perspectives.
Herpetological Monographs, 34(1), 71-97. doi:10.1655/HERPMONOGRAPHS-D-19-
00002.1
Seidl, F., Levis, N. A., Schell, R., Pfennig, D. W., Pfennig, K. S., & Ehrenreich, I. M. (2019).
Genome of *Spea multiplicata*, a Rapidly Developing, Phenotypically Plastic, and
Desert-Adapted Spadefoot Toad. *G3: Genes|Genomes|Genetics, 9*(12), 3909-3919.
doi:10.1534/g3.119.400705
Smit, A., Hubley, R., & Green, P. (2013). RepeatMasker Open-4.0. 2013-2015. Retrieved from
<http://www.repeatmasker.org>
Sotero-Caio, C., Challis, R., Kumar, S., & Blaxter, M. (2021). Genomes on a Tree (GoaT): A
centralized resource for eukaryotic genome sequencing initiatives. *Biodiversity
Information Science and Standards*.
Stapley, J., Santure, A. W., & Dennis, S. R. (2015). Transposable elements as agents of rapid
adaptation may explain the genetic paradox of invasive species. *Molecular Ecology,
24*(9), 2241-2252.
Storer, J., Hubley, R., Rosen, J., Wheeler, T. J., & Smit, A. F. (2021). The Dfam community
resource of transposable element families, sequence models, and genome
annotations. *Mobile DNA, 12*(1), 1-14.
Stuckert, A. M., Chouteau, M., McClure, M., LaPolice, T. M., Linderoth, T., Nielsen, R.,
Summers, K., & MacManes, M. D. (2021). The genomics of mimicry: gene expression
throughout development provides insights into convergent and divergent
phenotypes in a Müllerian mimicry system. *Molecular Ecology, 30*(16), 4039-4061.
Suda, K., Hayashi, S. R., Tamura, K., Takamatsu, N., & Ito, M. (2022). Activation of DNA
Transposons and Evolution of piRNA Genes Through Interspecific Hybridization in
Xenopus Frogs. *Frontiers in Genetics, 13*. doi:10.3389/fgene.2022.766424
Sun, C., López Arriaza, J. R., & Mueller, R. L. (2012a). Slow DNA loss in the gigantic genomes
of salamanders. *Genome Biology and Evolution, 4*(12), 1340-1348.
Sun, C., Shepard, D. B., Chong, R. A., López Arriaza, J., Hall, K., Castoe, T. A., Feschotte, C.,
Pollock, D. D., & Mueller, R. L. (2012b). LTR retrotransposons contribute to genomic
gigantism in plethodontid salamanders. *Genome Biology and Evolution, 4*(2), 168-
183.
Sun, Y.-B., Zhang, Y., & Wang, K. (2020). Perspectives on studying molecular adaptations of
amphibians in the genomic era. *Zoological Research, 41*(4), 351.
Team, R. (2022). RStudio: integrated development for R. Boston, MA: RStudio. In: Inc.
Team, R. C. (2013). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. *http://www*. *R-project. org/*.
Tymowska, J., & Fischberg, M. (1973). Chromosome complements of the genus *Xenopus*.
Chromosoma, 44(3), 335-342. doi:10.1007/BF00291027
Wang, J., Itgen, M. W., Wang, H., Gong, Y., Jiang, J., Li, J., Sun, C., Sessions, S. K., & Mueller,
R. L. (2021). Gigantic Genomes Provide Empirical Tests of Transposable Element
Dynamics Models. *Genomics, Proteomics & Bioinformatics*.
doi:https://doi.org/10.1016/j.gpb.2020.11.005

Waters, P. D., Patel, H. R., Ruiz-Herrera, A., Álvarez-González, L., Lister, N. C., Simakov, O.,
Ezaz, T., Kaur, P., Frere, C., Grützner, F., Georges, A., & Graves, J. A. M. (2021).
Microchromosomes are building blocks of bird, reptile, and mammal chromosomes.
Proceedings of the National Academy of Sciences, 118(45), e2112494118.
doi:doi:10.1073/pnas.2112494118
Womack, M. C., Steigerwald, E., Blackburn, D. C., Cannatella, D. C., Catenazzi, A., Che, J., Koo,
M. S., McGuire, J. A., Ron, S. R., Spencer, C. L., Vredenburg, V. T., & Tarvin, R. D.
(2022). State of the Amphibia 2020: A Review of Five Years of Amphibian Research
and Existing Resources. *Ichthyology & Herpetology, 110*(4), 638-661, 624.
Wu, W., Gao, Y. D., Jiang, D. C., Lei, J., Ren, J. L., Liao, W. B., Deng, C., Wang, Z., Hillis, D. M.,
Zhang, Y. P., & Li, J. T. (2022). Genomic adaptations for arboreal locomotion in Asian
flying treefrogs. *Proc Natl Acad Sci U S A, 119*(13), e2116342119.
doi:10.1073/pnas.2116342119
Zuo, B., Nneji, L. M., & Sun, Y.-B. (2023). Comparative genomics reveals insights into anuran
genome size evolution. *BMC Genomics, 24*(1), 379. doi:10.1186/s12864-023-09499-8
779

DATA ACCESSIBILITY AND BENEFIT-SHARING

Data Accessibility Statement

Genetic data:

All the genomes used in this study are available on the NCBI Genomes database

(https://www.ncbi.nlm.nih.gov/genome/).

Code:

All original code has been deposited on GitHub and is publicly available at

(https://doi.org/10.5281/zenodo.7679280).

AUTHOR CONTRIBUTIONS

Conceptualization, T.A.K., A.J.C., L.A.O., A.R., and K.C.W.V; methodology, T.A.K, N.D.Y, R.L.M, and

A.R.; formal analysis, T.A.K., M.L.P.; investigation, T.A.K., N.D.Y, R.L.M., K.C.W.V, M.L.P.; resources,

L.A.O. and A.R.; writing – original draft, T.A.K.; writing – review & editing, all authors; project

administration, T.A.K; funding acquisition, T.A.K. and L.F.S.

TABLES AND FIGURES

796 **Table 1**. Genome quality measures.

Order	Family	N chrs	Assembly ID	N scaf	Genome size (Gb)	Scaf N50 (kb)	Cont N50 (kb)	% chrs	% assembled
Anura	Aromobatidae	12	in progress	6,658	5.32	625,512	879	0.00	76.60
Anura	Bombinatoridae	12	GCA_027579735.1	2,468	10.02	1,241,341	3,760	97.73	90.00
Anura	Bombinatoridae	12	GCA905336975.1	4,065,339	4.68	2	2	0.00	52.20
Anura	Bufonidae	11	GCA905171765.1	1,307	5.04	635,713	3,960	98.92	78.20
Anura	Bufonidae	11	GCA014858855.1	747	4.55	539,822	1,738	96.21	83.00
Anura	Hylidae	15	GCA_027789765.1	2,200	2.21	153,439	8,808	94.14	95.60
Anura	Alytidae	14	GCA_027410445.1	1,318	3.87	453,718	3,044	95.72	75.40
Anura	Eleutherodactylidae	13	GCA019857665.1	105,233	2.79	109,467	11	63.88	73.50
Anura	Leptodactylidae	11	GCA019512145.1	108,304	2.56	172,109	305	75.27	108.00
Anura	Microhylidae	11	GCA_027917425.1	1,002	4.34	489,283	4,963	98.02	91.00
Anura	Ranidae	13	GCA018402905.1	561,726	7.63	748	11	0.00	95.70
Anura	Centrolenidae	10	in progress	227,850	4.04	295,404	10	0.00	126.20
Anura	Hylidae	12	GCA_029499605.1	3,474	4.14	384,967	3,809	90.48	96.00
Anura	Pipidae	12	GCA019447015.1	25,835	3.21	293,321	784	82.53	135.10
Anura	Megophryidae	13	GCA018994145.1	5,370	3.54	412,425	821	98.22	129.60
Anura	Megophryidae	13	GCA009667805.1	5,302	3.55	394,693	1,946	93.34	130.10
Anura	Leptodactylidae	11	GCA_947044405.1	3,509	2.51	4,630	4,630	0.00	79.90
Anura	Limnodynastidae	11	GCA011038615.1	520,896	2.38	286	11	0.00	74.80
Anura	Ranidae	13	GCA002284835.2	1,544,635	6.25	39	6	0.00	85.20
Anura	Ranidae	13	GCA_028564925.1	4,861	5.15	611,110	2,499	98.80	90.30
Anura	Mantellidae	13	in progress	135,214	3.41	324,859	14	0.00	72.60
Anura	Dicroglossidae	13	GCA000935625.1	25,187	2.05	1,069	33	0.00	50.60
Anura	Dendrobatidae	10	GCA009801035.1	125,383	3.49	78	6	0.00	39.90
Anura	Dendrobatidae	10	in progress	15,287	5.19	571,703	98	0.00	59.30
Anura	Pelobatidae	13	GCA_933207985.1	3,574	3.09	330,124	130	98.65	75.10
Anura	Dicroglossidae	13	GCA_022657655.1	1,446,664	1.83	1	1	0.00	45.10
Anura	Hylidae	26	GCA_025379985.1	109,372	4.74	78	78	0.00	72.70
Anura	Pipidae	10	GCA_021901965.1	830,118	1.19	2	2	0.00	50.20
Anura	Pipidae	15	GCA019650415.1	283,592	1.37	7	6	0.00	57.70
Anura	Limnodynastidae	11	GCA_016617825.1	148,035	1.07	27	5	0.00	46.90
Anura	Myobatrachidae	12	GCA_028390025.1	3,127	8.87	846,895	6,819	92.31	103.70
Anura	Pyxicephalidae	13	GCA004786255.1	5,411	1.56	157,522	30	98.54	114.20
Anura	Ranidae	12	GCA_029574335.1	2,983	4.83	547,819	1,704	92.76	92.50
Anura	Ranidae	13	GCA_029206835.1	5,628	10.16	938,497	4,644	85.63	194.50
Anura	Ranidae	13	GCA905171775.1	555	4.11	481,763	6,265	98.26	93.60
Anura	Dendrobatidae	9	GCA_958301615.1	79,800	6.02	390	309	0.00	68.80
Anura	Bufonidae	11	GCA900303285.1	31,391	2.55	167	167	0.00	54.80
Anura	Scaphiopodidae	13	GCA009364435.1	1,378,905	0.48	0	0	0.00	46.30
Anura	Scaphiopodidae	13	GCA009364455.1	1,509,266	0.71	1	1	0.00	51.90
Anura	Pelobatidae	13	GCA_027358695.2	17	0.99	114,150	21,170	99.99	81.80
Anura	Pelobatidae	13	GCA_029215755.1	351	1.16	120,843	14,336	0.00	81.80
Anura	Pelobatidae	13	GCA009364415.1	49,736	1.08	72	31	0.00	82.10
Anura	Ranidae	13	GCA_951230385.1	22,068	3.98	628	611	0.00	63.80
Anura	Pipidae	18	GCA_024363595.1	23,435	2.75	143,989	21	96.13	78.90
Anura	Pipidae	18	GCA017654675.1	54	2.74	155,251	22,452	99.97	89.00
Anura	Pipidae	10	GCA000004195.4	166	1.45	153,961	14,634	99.86	85.30
Caudata	Ambystomatidae	14	GCA002915635.3	27,157	28.21	1,205,707	218	94.77	81.20
Caudata	Salamandridae	12	GCA_026652325.1	267	20.30	1,237,755	45,590	99.61	97.00
Gymnophiona	Dermophiidae	19	GCA902459505.2	164	3.78	272,612	20,657	99.85	82.40
Gymnophiona	Siphonopidae	12	GCA017765095.2	1,081	4.69	376,147	3,662	97.06	34.30
Gymnophiona	Rhinatrematidae	14	GCA901001135 2	1 330	5.32	486 875	3 216	97 36	59 90

798
799 **Figure 1.** Phylogenetic tree of amphibian families. Amphibian families with representative genomes
800 are highlighted and numbers indicate genome counts per family. (Green) anurans, (blue) caecilians,
801 and (orange) salamanders. *Engystomops pustulosus* (Family) image was taken by B. Gratwicke, other
802 amphibian images were licensed to T. Kosch by Adobe Stock and Shutterstock.

Figure 2. BUSCO (Benchmarking Universal Single-Copy Orthologs) assessment results for amphibian genomes.

815
816 **Figure 3**. Repeat content across the amphibian genomes. (LINEs) long interspersed nuclear elements,
817 (LTRs) long terminal repeats, and (SINEs) short interspersed nuclear elements.

827
828 **Figure 4.** Phylogenetic independent contrasts (PICs) between the proportion of transposable element
829 content relative to genome size and Bio8 (representing mean temperature of wettest quarter).

836
837 **Figure 5**. Synteny plot of BUSCOs (Benchmarking Universal Single-Copy Orthologs) for representative
838 amphibian chromosome-level genomes. The phylogenetic tree was created with Timetree.org. The
839 reference genome is *Ambystoma mexicanum*. *Indicate inverted chromosomes. Chromosomes
840 without BUSCOs were excluded from the plot.
841

Species	Order	Family	N chrs	Assembly ID	N scaf	Genome size (Gb)	Scaf N50 (kb)	Cont N50 (kb)	% chrs	% assembled	% repeats	BUSCO % complete
Allobates femoralis	Anura	Aromobatidae	12	in progress	6,658	5.32	625,512	879	0.00	76.60	73.52	81.37
Bombina bombina	Anura	Bombinatoridae	12	GCA_027579735.1	2,468	10.02	1,241,341	3,760	97.73	90.00	69.56	83.62
Bombina variegata	Anura	Bombinatoridae	12	GCA905336975.1	4,065,339	4.68	2	2	0.00	52.20	57.73	13.03
Bufo bufo	Anura	Bufonidae	11	GCA905171765.1	1,307	5.04	635,713	3,960	98.92	78.20	76.92	90.11
Bufo gargarizans	Anura	Bufonidae	11	GCA014858855.1	747	4.55	539,822	1,738	96.21	83.00	73.92	86.74
Dendropsophus ebraccatus	Anura	Hylidae	15	GCA_027789765.1	2,200	2.21	153,439	8,808	94.14	95.60	59.10	91.37
Discoglossus pictus	Anura	Alytidae	14	GCA_027410445.1	1,318	3.87	453,718	3,044	95.72	75.40	64.10	92.05
Eleutherodactylus coqui	Anura	Eleutherodactylidae	13	GCA019857665.1	105,233	2.79	109,467	11	63.88	73.50	52.96	79.11
Engystomops pustulosus	Anura	Leptodactylidae	11	GCA019512145.1	108,304	2.56	172,109	305	75.27	108.00	51.56	75.48
Gastrophryne carolinensis	Anura	Microhylidae	11	GCA_027917425.1	1,002	4.34	489,283	4,963	98.02	91.00	71.17	90.06
Glandirana rugosa	Anura	Ranidae	13	GCA018402905.1	561,726	7.63	748	11	0.00	95.70	62.54	79.62
Hyalinobatrachium fleischmanni	Anura	Centrolenidae	10	in progress	227,850	4.04	295,404	10	0.00	126.20	61.12	66.27
Hyla sarda	Anura	Hylidae	12	GCA_029499605.1	3,474	4.14	384,967	3,809	90.48	96.00	70.86	88.96
Hymenochirus boettgeri	Anura	Pipidae	12	GCA019447015.1	25,835	3.21	293,321	784	82.53	135.10	51.76	87.38
Leptobrachium ailaonicum	Anura	Megophryidae	13	GCA018994145.1	5,370	3.54	412,425	821	98.22	129.60	67.52	81.79
Leptobrachium leishanense	Anura	Megophryidae	13	GCA009667805.1	5,302	3.55	394,693	1,946	93.34	130.10	70.10	89.55
Leptodactylus fallax	Anura	Leptodactylidae	11	GCA_947044405.1	3,509	2.51	4,630	4,630	0.00	79.90	63.55	90.45
Limnodynastes dumerilii	Anura	Limnodynastidae	11	GCA011038615.1	520,896	2.38	286	11	0.00	74.80	46.17	79.10
Lithobates catesbeianus	Anura	Ranidae	13	GCA002284835.2	1,544,635	6.25	39	6	0.00	85.20	62.41	44.05
Lithobates sylvaticus	Anura	Ranidae	13	GCA_028564925.1	4,861	5.15	611,110	2,499	98.80	90.30	70.19	85.44
Mantella aurantiaca	Anura	Mantellidae	13	in progress	135,214	3.41	324,859	14	0.00	72.60	52.28	58.57
Nanorana parkeri	Anura	Dicroglossidae	13	GCA000935625.1	25,187	2.05	1,069	33	0.00	50.60	50.45	90.40
Oophaga pumilio	Anura	Dendrobatidae	10	GCA009801035.1	125,383	3.49	78	6	0.00	39.90	40.55	67.57
Oophaga sylvatica	Anura	Dendrobatidae	10	in progress	15,287	5.19	571,703	98	0.00	59.30	81.63	53.47
Pelobates cultripes	Anura	Pelobatidae	13	GCA_933207985.1	3,574	3.09	330,124	130	98.65	75.10	57.39	87.74
Phrynoglossus myanhessei	Anura	Dicroglossidae	13	GCA_022657655.1	1,446,664	1.83	1	1	0.00	45.10	47.65	8.96
Phyllomedusa bahiana	Anura	Hylidae	26	GCA_025379985.1	109,372	4.74	78	78	0.00	72.70	69.02	53.94
Pipa carvalhoi	Anura	Pipidae	10	GCA_021901965.1	830,118	1.19	2	2	0.00	50.20	45.27	11.81
Pipa parva	Anura	Pipidae	15	GCA019650415.1	283,592	1.37	7	6	0.00	57.70	42.92	26.72
Platyplectrum ornatum	Anura	Limnodynastidae	11	GCA016617825.1	148,035	1.07	27	5	0.00	46.90	23.42	43.45
Pseudophryne corroboree	Anura	Myobatrachidae	12	GCA_028390025.1	3,127	8.87	846,895	6,819	92.31	103.70	75.86	87.85
Pyxicephalus adspersus	Anura	Pyxicephalidae	13	GCA004786255.1	5,411	1.56	157,522	30	98.54	114.20	31.64	87.93
Rana kukunoris	Anura	Ranidae	12	GCA_029574335.1	2,983	4.83	547,819	1,704	92.76	92.50	71.31	89.15
Rana muscosa	Anura	Ranidae	13	GCA_029206835.1	5,628	10.16	938,497	4,644	85.63	194.50	77.28	89.19
Rana temporaria	Anura	Ranidae	13	GCA905171775.1	555	4.11	481,763	6,265	98.26	93.60	68.52	90.72
Ranitomeya imitator	Anura	Dendrobatidae	9	GCA_958301615.1	79,800	6.02	390	309	0.00	68.80	76.94	97.50
Rhinella marina	Anura	Bufonidae	11	GCA900303285.1	31,391	2.55	167	167	0.00	54.80	67.02	85.89
Scaphiopus couchii	Anura	Scaphiopodidae	13	GCA009364435.1	1,378,905	0.48	0	0	0.00	46.30	27.34	0.70
Scaphiopus holbrookii	Anura	Scaphiopodidae	13	GCA009364455.1	1,509,266	0.71	1	1	0.00	51.90	31.65	2.60
Spea bombifrons	Anura	Pelobatidae	13	GCA_027358695.2	17	0.99	114,150	21,170	99.99	81.80	39.55	91.96
Spea hammondii	Anura	Pelobatidae	13	GCA_029215755.1	351	1.16	120,843	14,336	0.00	81.80	46.92	91.51
Spea multiplicata	Anura	Pelobatidae	13	GCA009364415.1	49,736	1.08	72	31	0.00	82.10	33.15	76.59
Staurois parvus	Anura	Ranidae	13	GCA_951230385.1	22,068	3.98	628	611	0.00	63.80	65.85	84.39
Xenopus borealis	Anura	Pipidae	18	GCA_024363595.1	23,435	2.75	143,989	21	96.13	78.90	35.67	86.61
Xenopus laevis	Anura	Pipidae	18	GCA017654675.1	54	2.74	155,251	22,452	99.97	89.00	48.23	95.95
Xenopus tropicalis	Anura	Pipidae	10	GCA000004195.4	166	1.45	153,961	14,634	99.86	85.30	44.21	94.44
Ambystoma mexicanum	Caudata	Ambystomatidae	14	GCA002915635.3	27,157	28.21	1,205,707	218	94.77	81.20	65.91	66.06
Pleurodeles waltl	Caudata	Salamandridae	12	GCA_026652325.1	267	20.30	1,237,755	45,590	99.61	97.00	71.01	84.82
Geotrypetes seraphini	Gymnophiona	Dermophiidae	19	GCA902459505.2	164	3.78	272,612	20,657	99.85	82.40	71.22	91.39
Microcaecilia unicolor	Gymnophiona	Siphonopidae	12	GCA901765095.2	1,081	4.69	376,147	3,662	97.06	34.30	68.35	90.89
Rhinatrema bivittatum	Gymnophiona	Rhinatrematidae	14	GCA901001135.2	1,330	5.32	486,875	3,216	97.36	59.90	66.99	91.58