Comparative analysis of amphibian genomes: an emerging resource for basic and applied research

Running title: Comparative analysis of amphibian genomes

Tiffany A. Kosch[1], Andrew J. Crawford[2], Rachel Lockridge Mueller[3], Katharina C. Wollenberg Valero[4],

Megan L. Power[4], Ariel Rodríguez [5], Lauren A. O'Connell[6], Neil D. Young[1], and Lee F. Skerratt[1]

[1] Faculty of Science, University of Melbourne, Melbourne, Australia

[2] Departamento de Ciencias Biológicas, Universidad de los Andes, Bogotá, Colombia

[3] Department of Biology, Colorado State University, Colorado, USA

[4] School of Biology and Environmental Science, University College Dublin, Dublin, Ireland

[5] Institute of Zoology, University of Veterinary Medicine of Hannover, Hannover, Germany

[6] Department of Biology, Stanford University, California, USA

Corresponding author: tiffany.kosch@unimelb.edu.au

ABSTRACT

Amphibians are the most threatened group of vertebrates and are in dire need of conservation

intervention to ensure their continued survival. They have many unique features including a high

diversity of reproductive strategies, permeable and specialized skin capable of producing toxins and

antimicrobial compounds, multiple genetic mechanisms of sex determination, and in some lineages

even the ability to regenerate limbs and internal organs. Although genomics approaches would shed

light on these unique phenotypic traits and aid in conservation management, sequencing and

assembly of amphibian genomes has lagged far behind other taxa due to their comparatively large

genome sizes. Fortunately, the development of long-read sequencing technologies and initiatives has

led to a recent burst of new amphibian genome assemblies. Although growing, the field of amphibian

genomics suffers from the lack of annotation resources, tools for working with challenging genomes,

and lack of high-quality assemblies in multiple clades of amphibians. Here we analyze 32 publicly

available amphibian genomes to evaluate their usefulness for functional genomics analysis. We

report considerable variation in assembly quality and completeness, and report some of the highest

transposable element and repeat contents of any vertebrate, which is associated with climate. We

also provide evidence of conserved genome synteny despite the long divergence times of this group

but show that chromosome naming and orientation have been inconsistent across genome

assemblies. Additionally, we discuss sequencing gaps in the phylogeny and suggest key targets for

future sequencing endeavors. Lastly, we propose increased investment in amphibian genomics

research to promote their conservation.

KEYWORDS

Amphibian genomes; comparative genomics; transposable elements; repeat expansion;

genome synteny

INTRODUCTION

Amphibians are an ancient lineage of vertebrates that predate amniotes by more than 100 million years. Despite the considerable age of this lineage, amphibians are now the most threatened group of vertebrates with more that 40% of species threatened by factors such as habitat change, disease, and over-exploitation (IUCN, 2022; Scheele et al., 2019). Notably, many of these threats are hard to reverse, suggesting that novel approaches that utilize genomic resources may lead to improved management decisions for some of the most endangered taxa (Kosch et al., 2022; Scheele et al., 2014).

We are only just beginning to understand the genetic basis of many of the unique features of amphibians. Amphibians exhibit a high diversity of reproductive strategies including biphasic and direct development, uniparental and biparental care, mouth and gastric brooding, and foam nesting (Brown et al., 2010; Nunes-de-Almeida et al., 2021; Schulte et al., 2020). They also have specialized skin capable of producing complex compounds of interest for drug discovery for the development of antimicrobial drugs and analgesics (Daly et al., 2000; De Angelis et al., 2021; Liu et al., 2020). Amphibians occur across habitat types from rainforests to deserts, freshwater streams to salt marshes, and tropical to arctic climates (Duellman, 1999), but it is unclear how this ecological diversity is reflected in genome composition. Salamanders are an important resource for transplant and regeneration research due to their ability to regenerate limbs and internal organs (Elewa et al., 2017; Nowoshilow et al., 2018). Amphibians also have many of the same immune components of mammals making them an important model resource for immunology (Paiola et al., 2023; Robert, 2020).

Despite the obvious value of amphibian genomes for research on ecology, evolution, medicine, and improving their conservation, until recently, the generation of amphibian reference genomes has

been markedly slower than other vertebrates (Hotaling et al., 2021a). This lag can be attributed to

high costs and the computational challenges of assembling their often large and complex genomes

(Sun et al., 2020). Recent advances in sequencing technologies such as long read sequencing and

assembly algorithms that incorporate hybrid approaches have circumvented many of these

challenges leading to a surge of high quality, chromosome-level reference genomes. The next

challenge will be developing the tools for annotation and comparative analyses of these large

genomes.

In this study, we provide a state-of-the-field synthesis of amphibian reference genome assemblies by

analyzing 32 publicly available amphibian genomes. We evaluate assembly quality, sequencing

technology, gene completeness, transposable element and repeat content and its ecological

correlates, taxonomic representation, and synteny.

MATERIALS AND METHODS

Genomes

A search conducted on the NCBI genome website using search term "amphibians" conducted on

November 1, 2021 revealed there were 39 amphibian genomes from 30 species. All genome files in

fasta format were downloaded for assessment. One genome was selected for each species. If there

was more than one draft of a genome, the most recent draft and/or the primary haplotype was

selected. In cases where there were multiple versions sequenced by different groups, the best

genome was selected by lowest scaffold number. We excluded the *Ranitomeya imitator* genome as

the genome was later retracted from NCBI. Entire genomes (including uncharacterized contigs but

excluding mitochondrial genomes) were used for assessment unless indicated otherwise.

Genome databases NCBI Genomes, NCBI RefSeq (O'Leary et al., 2016), Ensembl (Cunningham et al.,

2022), UCSC Genome Browser (Lee et al., 2022), and Genomes on a Tree (GoaT) (Sotero-Caio et al.,

2021) were searched for information on the 34 amphibian species with reference genomes including

chromosome number, annotation data, proteome availability, C-value, and sequencing technology.

Read length was classified as "short" for Illumina sequencing and "long" for PacBio and Nanopore

technologies.

A search for amphibian proteome datasets on NCBI RefSeq (O'Leary et al., 2016), Ensembl

(Cunningham et al., 2022), and UCSC Genome Browser (Lee et al., 2022) databases on June 24, 2022

revealed 11 proteomics datasets.

A search of the NCBI Organelle database on 15, February 2023 using search term "amphibian"

resulted in 353 mitochondrial genomes belonging to 345 species (Table S11). Seventeen

mitochondrial genomes overlapped with the amphibian nuclear genomes analyzed in this study.

Reference genome availability summary

The GoaT online database (Sotero-Caio et al., 2021) was searched on January 6, 2023 to summarize

genomes in progress or publicly available using the search terms "tax_tree(Amphibia) AND

tax_rank(species) AND sequencing_status=in_progress" or "tax_tree(Amphibia) AND

tax_rank(species) AND sequencing_status=insdc_open". The same search terms were used to

summarize publicly available genomes for mammals, birds, and non-avian reptiles with the

"tax_tree" search term replaced by appropriate Class.

Genome quality analyses

Genome quality assessment was performed with BBMap (v. 38.76). Benchmarking Universal Single-

Copy Orthologs (BUSCO) were summarized with the BUSCO tool (v. 5.1.2) (Manni et al., 2021) using

the OrthoDB Tetrapoda ortholog library (v. odb10) (Kriventseva et al., 2018) (N=5310 orthologs).

Percentage of the genome assembled to chromosomes was calculated with a custom bash script.

Phylogenetic tree

A species to family correspondence table was obtained from Jetz and Pyron (2018)

(https://vertlife.org/files_20170703/) and was filtered to include only the species with the longest

nucleotide sequence per family. This taxa subset was used to obtain a subset of 100 phylogenetic

trees from the posterior distribution of the Jetz and Pyron (2018) dataset, as available

from http://vertlife.org/phylosubsets. A consensus tree from these 100 trees was then obtained

using treeannotator (Drummond & Rambaut, 2007). The species names of the tree tips were then

substituted with the corresponding family names using the "sub.taxa.label" function in the

phylotools package (https://github.com/helixcn/phylotools) in R with the aid of the species to family

correspondence table, which was updated with the most recent classification available in

AmphibiaWeb https://amphibiaweb.org and the Amphibian Species of the World

database https://amphibiansoftheworld.amnh.org/.

Repeat modelling and annotation

Repeats were *de novo* modelled with RepeatModeler (v. 2.0.3) (Flynn et al., 2020). Genomes were

then annotated using RepeatMasker (v. 4.1.2-p1) (Smit et al., 2013) with a concatenated library of

genome-specific repeats generated from RepeatModeler and the Dfam amphibian repeat library (v.

Dfam.h5) (Storer et al., 2021). Before annotation, any previous soft masking of the genomes was

reversed. The results were summarized using a custom bash script.

Ecological correlates of transposable element content

On the basis of Global Biodiversity Information Facility (GBIF) data (https://www.gbif/org/; accessed

May 2023), MaxEnt habitat suitability-weighted means of a range of bioclimatic variables were

obtained for the 32 amphibian species. As previous studies have explored the relationship between

amphibian genome size and environmental variables (Liedtke et al., 2022), here we focused on the

relationship between bioclimatic variables and amphibian transposable elements. Influence of these

bioclimatic variables on transposable element content was modelled using Bayesian mixed effect

models (Hadfield, 2010), corrected for body size and phylogenetic signal (see Supplementary

Methods for further information).

Synteny analysis

Synteny of BUSCO genes for chromosome level assemblies was analyzed with R Package GENESPACE

(v. 0.9.4) (Lovell et al., 2022), which uses OrthoFinder (v. 2.5.4) (Emms & Kelly, 2019) to infer

orthology. Synteny was analyzed using BUSCO "full_table.tsv" results files that were reformatted for

GENESPACE input using a custom bash script. Synteny plots were generated for all chromosome level

assemblies, all anuran chromosome level assemblies, and for the three caecilian genomes using the

GENESPACE plotting tool "plot_riparianHits". Chromosomes with reversed orientation compared to

the reference genome were inverted to improve visualisation.

Quantification and statistical analysis

Regression analyses and Student's t-tests for comparing genome quality measurements were

conducted with the R statistics package (v. 4.1.2) (Team, 2013) in R Studio (v. 2022.02.3) (Team,

2022). Genome quality measures, contigN50, and scaffold count, were log transformed prior to

analysis. R-scripts for statistical analysis and plotting are available on GitHub at

https://doi.org/10.5281/zenodo.7679280.

RESULTS

Genome quality

A total of 32 nuclear amphibian genome assemblies were available for our study and were generated

with a variety of sequencing technologies, including Illumina (NextSeq, HiSeq), PacBio (RS11, Sequel),

and Oxford Nanopore. Sequenced genomes represented 20 of 73 amphibian families with reference

genomes distributed unevenly across the phylogeny (Fig. 1). For example, there is only one

salamander genome representing the 798 extant species, no genomes representing the microhylids

or hyperoliids, yet there are five pipid and four pelobatid genomes (Fig. 1).

Genome assembly length ranged from 0.48 Gb in *Scaphiopus couchii* to 28.21 Gb in *Ambystoma*

mexicanum and was strongly positively associated with c-value estimates of genome size (F_{30} = 270.2,

$p < 1 \times 10^{-11}$) (Table 1, Fig. S2). Fifteen of these genomes were assembled to the chromosome level of

which the percentage of the genome assigned to chromosomes ranged from 63.88 to 99.96% (Table

1). Percentage of the genome assigned to chromosomes was positively associated with contig N50

(F_{13} = 10.6, p = 0.006), number of scaffolds (F_{13} = 21.35, p < 0.001) and read length ($t_{28.4}$ = 3.65, p =

0.001). There are additionally mitochondrial genome assemblies for 345 species of which 17 had

nuclear reference genomes. Eleven of the species with genomes had proteomics data (Table S1).

The quality of the amphibian genomes varied considerably (Table 1). Genomes generated with short-

read technologies were of lower quality than long-read or hybrid genome assemblies as indicated by

significantly lower contig N50s (t_{29} = 6.05, p < 0.00001), percentage of complete Benchmarking

Universal Single-Copy Ortholog (BUSCO) genes ($t_{13.4}$ = 2.85, p = 0.013), and higher scaffold numbers

(t_{29} = -4.86, p < 0.0001).

Contig N50 ranged from 362 bp in *S. couchii* to 22.45 Mb in *Xenopus laevis* with a median of 32.80

Kb. Scaffold count varied considerably from 54 in *X. laevis* to more than four million in *Bombina*

variegata with a median of 27.16 Kb (Table 1). Benchmarking Universal Single-Copy Orthologs

(BUSCO) scores ranged from 0.7 to 96% completeness (Tables 1, S1; Fig. 2) and were positively

associated with contig N50 (F_{29} = 55.09, p < 0.0000001; Fig. S3) and scaffold count (F_{29} = 42.98, p <

0.000001). All genomes had low percentages of duplicate BUSCO genes (< 6%), suggesting they may

be diploid except for the known tetraploid species, *X. laevis* (Fig. 2).

The variation we report here in genome quality, contiguity, and completeness may impact the value

of the genomes for functional genomics research. However, the improvements in all these measures

seen with the utilization of long read technologies or hybrid assemblies suggests that genome quality

will continue to improve as these approaches are used more frequently.

214
Repeat content

Overall identified repeat percentage of the genomes ranged from 23% in *Platyplectrum ornatum* to

82% in *Oophaga sylvatica* and was positively associated with genome size (F_{30} = 5.87, p = 0.022)

(Tables 1; S3). Repeat content varied across genomes with the anuran *O. sylvatica* and the

salamander *Ambystoma mexicanum* dominated by Long Terminal Repeats (LTRs), the three caecilians

dominated by Long Interspersed Nuclear Elements (LINEs), and the two bufonid anurans dominated

by DNA transposons (Fig. 3; Tables S2-S4). The *Ambystoma mexicanum* genome had fewer repeats

than might be predicted by its large size (Fig. 3).

The proportion of repeats that could be classified by RepeatMasker ranged from 6.6% in *P. ornatum*

to 48.2% in *O. sylvatica* (Table S4) and was positively associated with genome quality measures

contigN50 (F_{29} = 12.8, p = 0.001), scaffold count (F_{29} = 7.80, p = 0.009), and percent BUSCO complete

(F_{29} = 4.60, p = 0.041). The ability to classify repeats was also positively associated with read length,

with longer reads resulting in better classification ($t_{26.241}$ = 3.57, p = 0.001).

These disparities in repeat percentage and content likely reflect differing evolutionary histories

among species, as indicated by three of the four congeneric species pairs in our dataset having

similar values (i.e., *Bufo*, *Leptobrachium*, and *Xenopus*; but not *Oophaga*). The differences we

observed in *Oophaga pumilio* and *O. sylvatica* are likely due to assembly quality rather than genome

content given that these two genomes were sequenced with different technologies and have

dramatically different genome qualities (e.g., contig N50s of 5.8 vs. 97.8 Kbp respectively).

Ecological correlates of transposable element content

A Bayesian mixed effect modelling approach was employed to examine the relationships

between proportion of transposable elements and environmental variables. Controlling for

phylogenetic relationships (by estimating Pagel's lambda, λ) and body size (Spearman

correlation with TRANSPOSABLE ELEMENTcontent ρ= -0.772, p<0.001) and excluding the

three globally invasive species (*Rhinella marina, X. laevis*, and *Lithobates catesbeianus*) our

analysis revealed a significant influence (pMCMC = 0.004) of Bio18 (precipitation in the

warmest quarter) on proportion of total transposable elements (Figs 4, S6; Table S7).

Inclusion of these three invasives for a total of 32 species showed additional significant

relationships with Bio15 (precipitation seasonality) and elevation (Table S6). Further analysis

indicated that the relationship with Bio18 was not specific to a particular class of

transposable elements, such as retroelements or DNA transposons (Tables S7 and S8).

Phylogenetic signal (Pagel's lambda, λ) was moderate when considering total transposable

elements and retroelements (0.440; Tables S6 and S7) and increased when we considered

DNA transposons alone (0.603; Table S8).

Genome synteny

Genome synteny of BUSCO genes was highly conserved within anurans (Fig. 5) and within caecilians

(Fig. S8) but was less conserved across the amphibian orders (Fig. S7). However, chromosome

naming was inconsistent across all taxa (Fig. 5, S10-11). For example, *X. tropicalis* chr1 is chr12 in

Leptobrachium ailaonicum (but not *L. leishanense*) and chr2 in *Bufo bufo* (but not *Bufo gargarizans*)

(Fig. 5). Orientation of chromosomes was also inconsistent, including between species of the same

genus (e.g., *Bufo*, *Leptobrachium*) (Fig. 5) and among the three caecilians (Fig. S8). Multiple

inversions were evident including between chr3 of pipids *(Xenopus tropicalis* and *Hymenochirus*

boettgeri) and other anurans (chromosomes 1, 2, 3, 4, or 10), caecilians (chr3 and chr4/5/6), and

even within species of the same genus (chr7 *Bufo gargarizans*, chr 9 *B. bufo*; Figs 5, S10-11). There

was also evidence of several chromosomal fissions including the separation of chr1 of *Leptobrachium*

leishanense into chr3 and chr6 in *Pyxicephalus adspersus* and into chr3 and chr7 in *Engystomops*

pustulosus; however, this chromosome remained mostly intact in the other anuran genomes (Fig. 5).

Given that these genomes were generated using different technologies, it is unclear whether or not

the differences we observed in synteny are due to true differences or assembly errors.

DISCUSSION

In this study we analysed 32 amphibian reference genomes from the public domain to evaluate their

content and usefulness for functional genetics research (Fig. 1, Table 1). There are considerably

fewer reference genomes for amphibians than exist for birds (N=754), mammals (N=406), and non-

avian reptiles (N=108). This scarcity of reference genomes results in many gaps in genome

representation across the amphibian tree of life including many entirely unrepresented groups and

with only one genome representing the entire order Caudata. The unrepresented families include

many of interest from a conservation perspective due to their high number of IUCN RedList Critically

Endangered species (e.g., Cryptobranchidae, Plethodontidae, Strabomantidae, and Craugastoridae)

(IUCN, 2022). However, our search of the Genomes on a Tree (GoaT) database (Sotero-Caio et al.,

2021) indicated that there are a further 24 amphibian genome assemblies in progress (20 anurans, 4

caudatans) indicating that this resource will be increasing by nearly 70% in the next few years.

The quality and completeness of the genomes in our dataset varied considerably (e.g., Fig. 2). Much

of this variation can be attributed to the sequencing technology used to generate them, with short-

read sequencing approaches resulting in lower completeness and continuity (Fig. S3). These impacts

are a recognized limitation of short-read sequencing and have been reported to impact genome

quality in taxa from insects (Hotaling et al., 2021b) to other vertebrates (Rhie et al., 2021), but have

likely had a disproportionate impact in amphibian genomes due to the difficulty of assembling

genomes with high repeat content (Sun et al., 2020). Fortunately, most ongoing sequencing efforts

now use long-read or hybrid sequencing approaches, which along with improved sequencing

algorithms, should result in higher quality amphibian genomes (Hotaling et al., 2021a; Lawniczak et

al., 2022; Rhie et al., 2021).

Genome quality (i.e., high continuity, contiguity, accuracy, completeness (Rhie et al., 2021)) is critical

for applications such as quantitative genetics where assembly errors can lead to incorrect inferences

in genetic association or genetic prediction. Quality also enhances the usefulness of genomes. For

example, highly contiguous chromosome-level assemblies decrease computational requirements for

downstream analyses such as mapping, variant calling, and alignment (Aganezov et al., 2022).

One of the most intriguing features of amphibian genomes is the huge range they exhibit in genome

size (Biscotti et al., 2019). This was exemplified in our dataset where assembly length ranged from

0.48 Gb in *Scaphiopus couchii* to 28 Gb in *Ambystoma mexicanum*. Why gigantic genomes exist in

some species, but not others, remains a key evolutionary question (Kapusta et al., 2017; Wang et al.,

2021). Explanations include differences in genome-level processes (e.g., insertion and deletion rates)

(Frahry et al., 2015; Sun et al., 2012), development (e.g., developmental rate and complexity)

(Gregory, 2002; Liedtke et al., 2018), physiology (e.g., water loss) (Johnson et al., 2021), body size

(e.g. miniaturization) (Decena-Segarra et al., 2020), and demography (e.g., effective population size)

(Liedtke et al., 2018; Lynch & Walsh, 2007) (but see Mohlhenrich & Mueller, 2016). As more

amphibian genomes become available, these hypotheses can be more rigorously evaluated.

We report some of the largest estimates of repeat content of any vertebrate (82% in *Oophaga*

sylvatica and 77% in *Bufo bufo*), exceeded only by the Australian lungfish at 90% (Meyer et al., 2021).

As expected, genome size was correlated with repeat content affirming that much of the variation in

amphibian genome size is due to an excess of repeats and transposable elements (Biscotti et al.,

2019; Lamichhaney et al., 2021).

In contrast to mammals, whose repeat landscape is mainly dominated by LTR retrotransposons (Platt

et al., 2018), amphibian repeat content varied considerably with some species dominated by DNA

transposons (as previously reported (Suda et al., 2022), and others by non-LTR retrotransposons

including the three caecilian genomes which were dominated by LINEs. This somewhat agrees with

genomic data and transcriptomic data from the caecilian *Ichthyophis bannanicus*, where LINEs were

the second most abundant type of repeat (26% of the genome) behind DIRS (30%) (Wang et al.,

2021); this is a higher percentage of LINES than we report in the three caecilian genomes in this study

(7 to 14%) (Table S4).

A considerable proportion of the repeats could not be classified. This was likely due to incorrect

classification (e.g., genes categorized as repeats) and the lack of good amphibian-specific repeat

resources (Ou et al., 2019) for classification via nucleotide sequence homology. The majority of

amphibian curated repeat libraries are generated in reference to *Xenopus* species (e.g., Dfam); the

large divergence times of this genus from the other amphibian species suggests that it may be a

contributing factor to the lack of classification. However, we also report many unclassified repeats in

the two *Xenopus* genomes.

The largest genome in our dataset, *A. mexicanum*, had fewer repeats than predicted given its size

(Fig S3) (Nowoshilow et al., 2018). This may be due, in part, to the Dfam library used for repeat

annotation being anuran-based; however, we did not observe this trend in the three caecilian

genomes in our dataset. Also, we performed *de novo* annotation of this genome, which should have

captured repetitive elements missing from Dfam. More likely, this low number of repeats reflects low

deletion rates and, thus, persistence of repeats in the genome for extremely long periods of time,

leading to their mutational decay into unique sequences whose repetitive origin is obscured (Keinath

et al., 2015; Novák et al., 2020).

We also show that amphibian species that inhabit wet and warm climates particularly during the

summer months have a greater proportion of transposable elements. This observed trend does not

appear to be driven by a specific group of transposons suggesting it may be caused by climatic

factors. Recent studies indicate that transposable elements exhibit greater activity in hotter climates

(Baduel et al., 2021) with an increasing number of studies suggesting increased transposable element

activity contributes to genetic diversification and facilitates species adaptation (Li et al., 2018;

Schrader & Schmitz, 2019; Stapley et al., 2015). The pattern observed here likewise suggests the

potential for heightened transposable element activity, and may help explain transposable element

accumulation and potentially the higher evolutionary rates observed in the genomes of tropical

amphibians.

Our study is the first to examine chromosomal synteny across all amphibian orders. We show that

overall synteny of amphibian genomes is relatively conserved, particularly within orders (Figs 5 and

S8). This aligns with previous results from anurans that reported conserved genome organization in

this group (Bredeson et al., 2021; Wu et al., 2022). However, chromosome content and number

varied across species, which seems to have been driven by multiple occurrences of chromosomal

fusions and fissions (e.g., Fig. 5). Chromosomal rearrangements have occurred throughout vertebrate

evolution, including the hypothesized fusion of microchromosomes in the ancestor of tetrapods to

create the larger macrochromosomes seen in amphibians and mammals and their subsequent fission

to create the microchromosomes of modern birds and non-avian reptiles (Waters et al., 2021).

Some of the structural rearrangements we detected may be due to assembly errors and should be

evaluated in future assemblies using long-read scaffolding approaches (e.g., Oxford nanopore

sequencing), chromosome conformation capture technologies (e.g., Hi-C), or chromosome mapping

approaches (e.g., FISH). We also identified incongruities with chromosome naming and orientation

caused by differences in assembly methods. These were apparent even within species of the same

genus (e.g., *Bufo*). We suggest potential revisions of existing genome annotations to improve

congruity and that future assemblies are curated consistently against high-quality reference genomes

(e.g., *Xenopus laevis*).

Conclusions

New sequencing technologies and assembly algorithms have resulted in enough genomes for comparative analyses spanning the amphibian phylogeny. This has already begun to yield important insights on the evolution (Lamichhaney et al., 2021; Wu et al., 2022), development (Schloissnig et al., 2021; Stuckert et al., 2021), sex determination (Hime et al., 2019; Ma & Veltsos, 2021), and unique features (Fischer et al., 2019; Nowoshilow et al., 2018; Seidl et al., 2019) of this interesting group of animals.

The increased availability of amphibian genomes can also aid conservation efforts in this highly threatened group by facilitating research on genome-wide functional diversity, which can be used to inform management decisions such as genetic rescue or targeted genetic intervention for species threatened by habitat loss or chytridiomycosis (Chestnut et al., 2014; Kosch et al., 2022). Additionally, well-annotated genomes can be used to create eDNA assays for population monitoring (Breton et al., 2022; Saeed et al., 2022).

Future research efforts should focus on generating more reference genomes to fill the gaps in the amphibian phylogeny and the identification of advantageous genetic traits against threats. Efforts should also be made to increase the quality of genomes and expand transcriptome and annotation databases. We suggest that these efforts strive to follow the recommendations of initiatives such as the Earth BioGenome Project (Lawniczak et al., 2022), the Darwin Tree of Life Project (Blaxter et al., 2022), and the Threatened Species Initiative (Hogg et al., 2022) to sequence at least one representative from each family to ensure taxonomic coverage. Species selection should prioritize species of interest for understanding valuable functional genetics traits; for example, for the purpose of immunological research to understand disease resistance, or for conservation purposes to enhance fitness.

ACKNOWLEDGEMENTS

This research was supported by The University of Melbourne's Research Computing Services and the

Petascale Campus Initiative, and Australian Research Council Grants (FT190100462, LP200301370).

We are grateful to A. Stuckert, N. Brajuka and J. Sproul for their advice on repeat modelling, and J.T.

Li for providing suggestions on synteny analyses. We thank M. Moore for exploratory repeat

analyses.

REFERENCES
Aganezov, S., Yan, S. M., Soto, D. C., Kirsche, M., Zarate, S., Avdeyev, P., Taylor, D. J., Shafin,
K., Shumate, A., Xiao, C., Wagner, J., McDaniel, J., Olson, N. D., Sauria, M. E. G.,
Vollger, M. R., Rhie, A., Meredith, M., Martin, S., Lee, J., Koren, S., Rosenfeld, J. A.,
Paten, B., Layer, R., Chin, C. S., Sedlazeck, F. J., Hansen, N. F., Miller, D. E., Phillippy, A.
M., Miga, K. H., McCoy, R. C., Dennis, M. Y., Zook, J. M., & Schatz, M. C. (2022). A
complete reference genome improves analysis of human genetic variation. *Science,*
376(6588), eabl3533. doi:10.1126/science.abl3533
Baduel, P., Leduque, B., Ignace, A., Gy, I., Gil Jr, J., Loudet, O., Colot, V., & Quadrana, L.
(2021). Genetic and environmental modulation of transposition shapes the
evolutionary potential of Arabidopsis thaliana. *Genome Biology, 22*(1), 138.
Biscotti, M. A., Carducci, F., Olmo, E., & Canapa, A. (2019). Vertebrate Genome Size and the
Impact of Transposable Elements in Genome Evolution. In P. Pontarotti (Ed.),
Evolution, Origin of Life, Concepts and Methods (pp. 233-251). Cham: Springer
International Publishing.
Blaxter, M., Archibald, J. M., Childers, A. K., Coddington, J. A., Crandall, K. A., Di Palma, F.,
Durbin, R., Edwards, S. V., Graves, J. A. M., Hackett, K. J., Hall, N., Jarvis, E. D.,
Johnson, R. N., Karlsson, E. K., Kress, W. J., Kuraku, S., Lawniczak, M. K. N., Lindblad-
Toh, K., Lopez, J. V., Moran, N. A., Robinson, G. E., Ryder, O. A., Shapiro, B., Soltis, P.
S., Warnow, T., Zhang, G., & Lewin, H. A. (2022). Why sequence all eukaryotes?
Proceedings of the National Academy of Sciences, 119(4), e2115636118.
doi:10.1073/pnas.2115636118
Bredeson, J. V., Mudd, A. B., Medina-Ruiz, S., Mitros, T., Smith, O. K., Miller, K. E., Lyons, J. B.,
Batra, S. S., Park, J., Berkoff, K. C., Plott, C., Grimwood, J., Schmutz, J., Aguirre-
Figueroa, G., Khokha, M. K., Lane, M., Philipp, I., Laslo, M., Hanken, J., Kerdivel, G.,
Buisine, N., Sachs, L. M., Buchholz, D. R., Kwon, T., Smith-Parker, H., Gridi-Papp, M.,
Ryan, M. J., Denton, R. D., Malone, J. H., Wallingford, J. B., Straight, A. F., Heald, R.,
Hockemeyer, D., Harland, R. M., & Rokhsar, D. S. (2021). Conserved chromatin and
repetitive patterns reveal slow genome evolution in frogs. *bioRxiv*,
2021.2010.2018.464293. doi:10.1101/2021.10.18.464293
Breton, B.-A. A., Beaty, L., Bennett, A. M., Kyle, C. J., Lesbarrères, D., Vilaça, S. T., Wikston,
M. J. H., Wilson, C. C., & Murray, D. L. (2022). Testing the effectiveness of
environmental DNA (eDNA) to quantify larval amphibian abundance. *Environmental*
DNA, 4(6), 1229-1240. doi:10.1002/edn3.332

Brown, J. L., Morales, V., & Summers, K. (2010). A key ecological trait drove the evolution of biparental care and monogamy in an amphibian. *The American Naturalist, 175*(4), 436-446.

Chestnut, T., Anderson, C., Popa, R., Blaustein, A. R., Voytek, M., Olson, D. H., & Kirshtein, J. (2014). Heterogeneous Occupancy and Density Estimates of the Pathogenic Fungus *Batrachochytrium dendrobatidis* in Waters of North America. *PLOS ONE, 9*(9), e106790. doi:10.1371/journal.pone.0106790

Cunningham, F., Allen, J. E., Allen, J., Alvarez-Jarreta, J., Amode, M. R., Armean, I. M., Austine-Orimoloye, O., Azov, A. G., Barnes, I., Bennett, R., Berry, A., Bhai, J., Bignell, A., Billis, K., Boddu, S., Brooks, L., Charkhchi, M., Cummins, C., Da Rin Fioretto, L., Davidson, C., Dodiya, K., Donaldson, S., El Houdaigui, B., El Naboulsi, T., Fatima, R., Giron, C. G., Genez, T., Martinez, J. G., Guijarro-Clarke, C., Gymer, A., Hardy, M., Hollis, Z., Hourlier, T., Hunt, T., Juettemann, T., Kaikala, V., Kay, M., Lavidas, I., Le, T., Lemos, D., Marugán, J. C., Mohanan, S., Mushtaq, A., Naven, M., Ogeh, D. N., Parker, A., Parton, A., Perry, M., Piližota, I., Prosovetskaia, I., Sakthivel, M. P., Salam, A. I. A., Schmitt, B. M., Schuilenburg, H., Sheppard, D., Pérez-Silva, J. G., Stark, W., Steed, E., Sutinen, K., Sukumaran, R., Sumathipala, D., Suner, M. M., Szpak, M., Thormann, A., Tricomi, F. F., Urbina-Gómez, D., Veidenberg, A., Walsh, T. A., Walts, B., Willhoft, N., Winterbottom, A., Wass, E., Chakiachvili, M., Flint, B., Frankish, A., Giorgetti, S., Haggerty, L., Hunt, S. E., GR, I. I., Loveland, J. E., Martin, F. J., Moore, B., Mudge, J. M., Muffato, M., Perry, E., Ruffier, M., Tate, J., Thybert, D., Trevanion, S. J., Dyer, S., Harrison, P. W., Howe, K. L., Yates, A. D., Zerbino, D. R., & Flicek, P. (2022). Ensembl 2022. *Nucleic Acids Research, 50*(D1), D988-d995. doi:10.1093/nar/gkab1049

Daly, J. W., Garraffo, H. M., Spande, T. F., Decker, M. W., Sullivan, J. P., & Williams, M. (2000). Alkaloids from frog skin: the discovery of epibatidine and the potential for developing novel non-opioid analgesics. *Natural product reports, 17*(2), 131-135.

De Angelis, M., Casciaro, B., Genovese, A., Brancaccio, D., Marcocci, M., Novellino, E., Carotenuto, A., Palamara, A., Mangoni, M., & Nencioni, L. (2021). Temporin G, an amphibian antimicrobial peptide against influenza and parainfluenza respiratory viruses: insights into biological activity and mechanism of action. *The FASEB Journal, 35*(2), e21358.

Decena-Segarra, L. P., Bizjak-Mali, L., Kladnik, A., Sessions, S. K., & Rovito, S. M. (2020). Miniaturization, genome size, and biological size in a diverse clade of salamanders. *The American Naturalist, 196*(5), 634-648. doi:10.5061/dryad.ht76hdrcg

Drummond, A. J., & Rambaut, A. (2007). BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evolutionary Biology, 7*, 214. doi:10.1186/1471-2148-7-214

Duellman, W. E. (1999). *Patterns of distribution of amphibians: a global perspective*: JHU Press.

Elewa, A., Wang, H., Talavera-López, C., Joven, A., Brito, G., Kumar, A., Hameed, L. S., Penrad-Mobayed, M., Yao, Z., & Zamani, N. (2017). Reading and editing the *Pleurodeles waltl* genome reveals novel features of tetrapod regeneration. *Nature Communications, 8*(1), 1-9. doi:10.1038/s41467-017-01964-9

Emms, D. M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biology, 20*(1), 1-14.

Fischer, E. K., Roland, A. B., Moskowitz, N. A., Tapia, E. E., Summers, K., Coloma, L. A., & O'Connell, L. A. (2019). The neural basis of tadpole transport in poison frogs. *Proceedings of the Royal Society B: Biological Sciences, 286*(1907), 20191084. doi:10.1098/rspb.2019.1084

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F. (2020). RepeatModeler2 for automated genomic discovery of transposable element families. *Proceedings of the National Academy of Sciences, 117*(17), 9451-9457. doi:10.1073/pnas.1921046117

Frahry, M. B., Sun, C., Chong, R. A., & Mueller, R. L. (2015). Low levels of LTR retrotransposon deletion by ectopic recombination in the gigantic genomes of salamanders. *Journal of Molecular Evolution, 80*, 120-129. doi:10.1007/s00239-014-9663-7

Gregory, T. R. (2002). Genome size and developmental complexity. *Genetica, 115*(1), 131-146. doi:10.1023/A:1016032400147

Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. *Journal of statistical software, 33*, 1-22.

Hime, P. M., Briggler, J. T., Reece, J. S., & Weisrock, D. W. (2019). Genomic Data Reveal Conserved Female Heterogamety in Giant Salamanders with Gigantic Nuclear Genomes. *G3: Genes|Genomes|Genetics*, g3.400556.402019. doi:10.1534/g3.119.400556

Hogg, C. J., Ottewell, K., Latch, P., Rossetto, M., Biggs, J., Gilbert, A., Richmond, S., & Belov, K. (2022). Threatened Species Initiative: Empowering conservation action using genomic resources. *Proceedings of the National Academy of Sciences, 119*(4), e2115643118. doi:10.1073/pnas.2115643118

Hotaling, S., Kelley, J. L., & Frandsen, P. B. (2021a). Toward a genome sequence for every animal: Where are we now? *Proceedings of the National Academy of Sciences, 118*(52), e2109019118. doi:10.1073/pnas.2109019118

Hotaling, S., Sproul, J. S., Heckenhauer, J., Powell, A., Larracuente, A. M., Pauls, S. U., Kelley, J. L., & Frandsen, P. B. (2021b). Long-reads are revolutionizing 20 years of insect genome sequencing. *Genome Biology and Evolution*. doi:10.1093/gbe/evab138

IUCN. (2022). The IUCN Red List of Threatened Species. Version 2022-2. Accessed on [6 January 2023]. https://www.iucnredlist.org.

Jetz, W., & Pyron, R. A. (2018). The interplay of past diversification and evolutionary isolation with present imperilment across the amphibian tree of life. *Nature Ecology & Evolution, 2*(5), 850-858. doi:10.1038/s41559-018-0515-5

Johnson, B. B., Searle, J. B., & Sparks, J. P. (2021). Genome size influences adaptive plasticity of water loss, but not metabolic rates, in lungless salamanders. *Journal of Experimental Biology, 224*(8), jeb242196. doi:10.1242/jeb.242196

Kapusta, A., Suh, A., & Feschotte, C. (2017). Dynamics of genome size evolution in birds and mammals. *Proceedings of the National Academy of Sciences, 114*(8), E1460-E1469. doi:10.1073/pnas.1616702114

Keinath, M. C., Timoshevskiy, V. A., Timoshevskaya, N. Y., Tsonis, P. A., Voss, S. R., & Smith, J. J. (2015). Initial characterization of the large genome of the salamander *Ambystoma mexicanum* using shotgun and laser capture chromosome sequencing. *Scientific Reports, 5*(1), 16413. doi:10.1038/srep16413

Kosch, T. A., Waddle, A. W., Cooper, C. A., Zenger, K. R., Garrick, D. J., Berger, L., & Skerratt, L. F. (2022). Genetic approaches for increasing fitness in endangered species. *Trends in Ecology & Evolution, 37*(4), 332-345. doi:10.1016/j.tree.2021.12.003

Kriventseva, E. V., Kuznetsov, D., Tegenfeldt, F., Manni, M., Dias, R., Simão, F. A., & Zdobnov, E. M. (2018). OrthoDB v10: sampling the diversity of animal, plant, fungal, protist, bacterial and viral genomes for evolutionary and functional annotations of orthologs. *Nucleic Acids Research, 47*(D1), D807-D811. doi:10.1093/nar/gky1053

Lamichhaney, S., Catullo, R., Keogh, J. S., Clulow, S., Edwards, S. V., & Ezaz, T. (2021). A bird-like genome from a frog: Mechanisms of genome size reduction in the ornate burrowing frog, *Platyplectrum ornatum*. *Proceedings of the National Academy of Sciences, 118*(11), e2011649118. doi:10.1073/pnas.2011649118

Lawniczak, M. K. N., Durbin, R., Flicek, P., Lindblad-Toh, K., Wei, X., Archibald, J. M., Baker, W. J., Belov, K., Blaxter, M. L., Marques Bonet, T., Childers, A. K., Coddington, J. A., Crandall, K. A., Crawford, A. J., Davey, R. P., Di Palma, F., Fang, Q., Haerty, W., Hall, N., Hoff, K. J., Howe, K., Jarvis, E. D., Johnson, W. E., Johnson, R. N., Kersey, P. J., Liu, X., Lopez, J. V., Myers, E. W., Pettersson, O. V., Phillippy, A. M., Poelchau, M. F., Pruitt, K. D., Rhie, A., Castilla-Rubio, J. C., Sahu, S. K., Salmon, N. A., Soltis, P. S., Swarbreck, D., Thibaud-Nissen, F., Wang, S., Wegrzyn, J. L., Zhang, G., Zhang, H., Lewin, H. A., & Richards, S. (2022). Standards recommendations for the Earth BioGenome Project. *Proceedings of the National Academy of Sciences, 119*(4), e2115639118. doi:10.1073/pnas.2115639118

Lee, B. T., Barber, G. P., Benet-Pagès, A., Casper, J., Clawson, H., Diekhans, M., Fischer, C., Gonzalez, J. N., Hinrichs, A. S., Lee, C. M., Muthuraman, P., Nassar, L. R., Nguy, B., Pereira, T., Perez, G., Raney, B. J., Rosenbloom, K. R., Schmelter, D., Speir, M. L., Wick, B. D., Zweig, A. S., Haussler, D., Kuhn, R. M., Haeussler, M., & Kent, W. J. (2022). The UCSC Genome Browser database: 2022 update. *Nucleic Acids Research, 50*(D1), D1115-d1122. doi:10.1093/nar/gkab959

Li, Z.-W., Hou, X.-H., Chen, J.-F., Xu, Y.-C., Wu, Q., González, J., & Guo, Y.-L. (2018). Transposable elements contribute to the adaptation of *Arabidopsis thaliana*. *Genome Biology and Evolution, 10*(8), 2140-2150. doi:10.1093/gbe/evy171

Liedtke, H. C., Cruz, F., Gómez-Garrido, J., Fuentes Palacios, D., Marcet-Houben, M., Gut, M., Alioto, T., Gabaldón, T., & Gomez-Mestre, I. (2022). Chromosome-level assembly, annotation and phylome of Pelobates cultripes, the western spadefoot toad. *DNA Research, 29*(3), dsac013.

Liedtke, H. C., Gower, D. J., Wilkinson, M., & Gomez-Mestre, I. (2018). Macroevolutionary shift in the size of amphibian genomes and the role of life history and climate. *Nature Ecology & Evolution, 2*(11), 1792-1799. doi:10.1038/s41559-018-0674-4

Liu, Y., Shi, D., Wang, J., Chen, X., Zhou, M., Xi, X., Cheng, J., Ma, C., Chen, T., & Shaw, C. (2020). A novel amphibian antimicrobial peptide, phylloseptin-PV1, exhibits effective anti-staphylococcal activity without inducing either hepatic or renal toxicity in mice. *Frontiers in Microbiology, 11*, 565158.

Lovell, J. T., Sreedasyam, A., Schranz, M. E., Wilson, M., Carlson, J. W., Harkess, A., Emms, D., Goodstein, D. M., & Schmutz, J. (2022). GENESPACE tracks regions of interest and gene copy number variation across multiple genomes. *eLife, 11*, e78526. doi:10.7554/eLife.78526

569 Lynch, M., & Walsh, B. (2007). *The origins of genome architecture*. Sunderland: Sinauer
570   Associates.
571 Ma, W.-J., & Veltsos, P. (2021). The Diversity and Evolution of Sex Chromosomes in Frogs.
572   *Genes, 12*(4), 483. doi:10.3390/genes12040483
573 Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO
574   Update: Novel and Streamlined Workflows along with Broader and Deeper
575   Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes.
576   *Molecular Biology and Evolution, 38*(10), 4647-4654. doi:10.1093/molbev/msab199
577 Meyer, A., Schloissnig, S., Franchini, P., Du, K., Woltering, J. M., Irisarri, I., Wong, W. Y.,
578   Nowoshilow, S., Kneitz, S., Kawaguchi, A., Fabrizius, A., Xiong, P., Dechaud, C., Spaink,
579   H. P., Volff, J.-N., Simakov, O., Burmester, T., Tanaka, E. M., & Schartl, M. (2021).
580   Giant lungfish genome elucidates the conquest of land by vertebrates. *Nature,*
581   *590*(7845), 284-289. doi:10.1038/s41586-021-03198-8
582 Mohlhenrich, E. R., & Mueller, R. L. (2016). Genetic drift and mutational hazard in the
583   evolution of salamander genomic gigantism. *Evolution, 70*(12), 2865-2878.
584 Novák, P., Guignard, M. S., Neumann, P., Kelly, L. J., Mlinarec, J., Koblížková, A., Dodsworth,
585   S., Kovařík, A., Pellicer, J., Wang, W., Macas, J., Leitch, I. J., & Leitch, A. R. (2020).
586   Repeat-sequence turnover shifts fundamentally in species with large genomes.
587   *Nature Plants, 6*(11), 1325-1329. doi:10.1038/s41477-020-00785-x
588 Nowoshilow, S., Schloissnig, S., Fei, J.-F., Dahl, A., Pang, A. W. C., Pippel, M., Winkler, S.,
589   Hastie, A. R., Young, G., Roscito, J. G., Falcon, F., Knapp, D., Powell, S., Cruz, A., Cao,
590   H., Habermann, B., Hiller, M., Tanaka, E. M., & Myers, E. W. (2018). The axolotl
591   genome and the evolution of key tissue formation regulators. *Nature*.
592   doi:10.1038/nature25458
593 https://www.nature.com/articles/nature25458#supplementary-information
594 Nunes-de-Almeida, C. H. L., Haddad, C. F. B., & Toledo, L. F. (2021). A revised classification of
595   the amphibian reproductive modes. *Salamandra, 57*(3), 413-427.
596 O'Leary, N. A., Wright, M. W., Brister, J. R., Ciufo, S., Haddad, D., McVeigh, R., Rajput, B.,
597   Robbertse, B., Smith-White, B., Ako-Adjei, D., Astashyn, A., Badretdin, A., Bao, Y.,
598   Blinkova, O., Brover, V., Chetvernin, V., Choi, J., Cox, E., Ermolaeva, O., Farrell, C. M.,
599   Goldfarb, T., Gupta, T., Haft, D., Hatcher, E., Hlavina, W., Joardar, V. S., Kodali, V. K.,
600   Li, W., Maglott, D., Masterson, P., McGarvey, K. M., Murphy, M. R., O'Neill, K., Pujar,
601   S., Rangwala, S. H., Rausch, D., Riddick, L. D., Schoch, C., Shkeda, A., Storz, S. S., Sun,
602   H., Thibaud-Nissen, F., Tolstoy, I., Tully, R. E., Vatsan, A. R., Wallin, C., Webb, D., Wu,
603   W., Landrum, M. J., Kimchi, A., Tatusova, T., DiCuccio, M., Kitts, P., Murphy, T. D., &
604   Pruitt, K. D. (2016). Reference sequence (RefSeq) database at NCBI: current status,
605   taxonomic expansion, and functional annotation. *Nucleic Acids Research, 44*(D1),
606   D733-745. doi:10.1093/nar/gkv1189
607 Ou, S., Su, W., Liao, Y., Chougule, K., Agda, J. R. A., Hellinga, A. J., Lugo, C. S. B., Elliott, T. A.,
608   Ware, D., Peterson, T., Jiang, N., Hirsch, C. N., & Hufford, M. B. (2019). Benchmarking
609   transposable element annotation methods for creation of a streamlined,
610   comprehensive pipeline. *Genome Biology, 20*(1), 275. doi:10.1186/s13059-019-1905-
611   y
612 Paiola, M., Dimitrakopoulou, D., Pavelka, M. S., & Robert, J. (2023). Amphibians as a model
613   to study the role of immune cell heterogeneity in host and mycobacterial

interactions. *Developmental & Comparative Immunology, 139*, 104594. doi:https://doi.org/10.1016/j.dci.2022.104594

Platt, R. N., 2nd, Vandewege, M. W., & Ray, D. A. (2018). Mammalian transposable elements and their impacts on genome evolution. *Chromosome Res, 26*(1-2), 25-43. doi:10.1007/s10577-017-9570-z

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G. L., Cantin, L. J., Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M., Haase, B., Mountcastle, J., Winkler, S., Paez, S., Howard, J., Vernes, S. C., Lama, T. M., Grutzner, F., Warren, W. C., Balakrishnan, C. N., Burt, D., George, J. M., Biegler, M. T., Iorns, D., Digby, A., Eason, D., Robertson, B., Edwards, T., Wilkinson, M., Turner, G., Meyer, A., Kautt, A. F., Franchini, P., Detrich, H. W., Svardal, H., Wagner, M., Naylor, G. J. P., Pippel, M., Malinsky, M., Mooney, M., Simbirsky, M., Hannigan, B. T., Pesout, T., Houck, M., Misuraca, A., Kingan, S. B., Hall, R., Kronenberg, Z., Sović, I., Dunn, C., Ning, Z., Hastie, A., Lee, J., Selvaraj, S., Green, R. E., Putnam, N. H., Gut, I., Ghurye, J., Garrison, E., Sims, Y., Collins, J., Pelan, S., Torrance, J., Tracey, A., Wood, J., Dagnew, R. E., Guan, D., London, S. E., Clayton, D. F., Mello, C. V., Friedrich, S. R., Lovell, P. V., Osipova, E., Al-Ajli, F. O., Secomandi, S., Kim, H., Theofanopoulou, C., Hiller, M., Zhou, Y., Harris, R. S., Makova, K. D., Medvedev, P., Hoffman, J., Masterson, P., Clark, K., Martin, F., Howe, K., Flicek, P., Walenz, B. P., Kwak, W., Clawson, H., Diekhans, M., Nassar, L., Paten, B., Kraus, R. H. S., Crawford, A. J., Gilbert, M. T. P., Zhang, G., Venkatesh, B., Murphy, R. W., Koepfli, K.-P., Shapiro, B., Johnson, W. E., Di Palma, F., Marques-Bonet, T., Teeling, E. C., Warnow, T., Graves, J. M., Ryder, O. A., Haussler, D., O'Brien, S. J., Korlach, J., Lewin, H. A., Howe, K., Myers, E. W., Durbin, R., Phillippy, A. M., & Jarvis, E. D. (2021). Towards complete and error-free genome assemblies of all vertebrate species. *Nature, 592*(7856), 737-746. doi:10.1038/s41586-021-03451-0

Robert, J. (2020). Experimental platform using the amphibian *Xenopus laevis* for research in fundamental and medical immunology. *Cold Spring Harbor Protocols, 2020*(7), pdb. top106625.

Saeed, M., Rais, M., Akram, A., Williams, M. R., Kellner, K. F., Hashsham, S. A., & Davis, D. R. (2022). Development and validation of an eDNA protocol for monitoring endemic Asian spiny frogs in the Himalayan region of Pakistan. *Scientific Reports, 12*(1), 5624. doi:10.1038/s41598-022-09084-1

Scheele, B. C., Hunter, D. A., Grogan, L. F., Berger, L., Kolby, J. E., McFadden, M. S., Marantelli, G., Skerratt, L. F., & Driscoll, D. A. (2014). Interventions for Reducing Extinction Risk in Chytridiomycosis-Threatened Amphibians. *Conservation Biology, 28*(5), 1195-1205. doi:10.1111/cobi.12322

Scheele, B. C., Pasmans, F., Skerratt, L. F., Berger, L., Martel, A., Beukema, W., Acevedo, A. A., Burrowes, P. A., Carvalho, T., Catenazzi, A., De la Riva, I., Fisher, M. C., Flechas, S. V., Foster, C. N., Frías-Álvarez, P., Garner, T. W. J., Gratwicke, B., Guayasamin, J. M., Hirschfeld, M., Kolby, J. E., Kosch, T. A., La Marca, E., Lindenmayer, D. B., Lips, K. R., Longo, A. V., Maneyro, R., McDonald, C. A., Mendelson, J., Palacios-Rodriguez, P., Parra-Olea, G., Richards-Zawacki, C. L., Rödel, M.-O., Rovito, S. M., Soto-Azat, C., Toledo, L. F., Voyles, J., Weldon, C., Whitfield, S. M., Wilkinson, M., Zamudio, K. R., &

Canessa, S. (2019). Amphibian fungal panzootic causes catastrophic and ongoing loss
of biodiversity. *Science, 363*(6434), 1459-1463. doi:10.1126/science.aav0379
Schloissnig, S., Kawaguchi, A., Nowoshilow, S., Falcon, F., Otsuki, L., Tardivo, P.,
Timoshevskaya, N., Keinath, M. C., Smith, J. J., & Voss, S. R. (2021). The giant axolotl
genome uncovers the evolution, scaling, and transcriptional control of complex gene
loci. *Proceedings of the National Academy of Sciences, 118*(15).
Schrader, L., & Schmitz, J. (2019). The impact of transposable elements in adaptive
evolution. *Molecular Ecology, 28*(6), 1537-1549.
Schulte, L. M., Ringler, E., Rojas, B., & Stynoski, J. L. (2020). Developments in Amphibian
Parental Care Research: History, Present Advances, and Future Perspectives.
Herpetological Monographs, 34(1), 71-97. doi:10.1655/HERPMONOGRAPHS-D-19-
00002.1
Seidl, F., Levis, N. A., Schell, R., Pfennig, D. W., Pfennig, K. S., & Ehrenreich, I. M. (2019).
Genome of *Spea multiplicata*, a Rapidly Developing, Phenotypically Plastic, and
Desert-Adapted Spadefoot Toad. *G3: Genes|Genomes|Genetics, 9*(12), 3909-3919.
doi:10.1534/g3.119.400705
Smit, A., Hubley, R., & Green, P. (2013). RepeatMasker Open-4.0. 2013-2015. Retrieved from
<http://www.repeatmasker.org>
Sotero-Caio, C., Challis, R., Kumar, S., & Blaxter, M. (2021). Genomes on a Tree (GoaT): A
centralized resource for eukaryotic genome sequencing initiatives. *Biodiversity*
Information Science and Standards.
Stapley, J., Santure, A. W., & Dennis, S. R. (2015). Transposable elements as agents of rapid
adaptation may explain the genetic paradox of invasive species. *Molecular Ecology,*
24(9), 2241-2252.
Storer, J., Hubley, R., Rosen, J., Wheeler, T. J., & Smit, A. F. (2021). The Dfam community
resource of transposable element families, sequence models, and genome
annotations. *Mobile DNA, 12*(1), 1-14.
Stuckert, A. M., Chouteau, M., McClure, M., LaPolice, T. M., Linderoth, T., Nielsen, R.,
Summers, K., & MacManes, M. D. (2021). The genomics of mimicry: gene expression
throughout development provides insights into convergent and divergent
phenotypes in a Müllerian mimicry system. *Molecular Ecology*.
Suda, K., Hayashi, S. R., Tamura, K., Takamatsu, N., & Ito, M. (2022). Activation of DNA
Transposons and Evolution of piRNA Genes Through Interspecific Hybridization in
Xenopus Frogs. *Frontiers in Genetics, 13*. doi:10.3389/fgene.2022.766424
Sun, C., Shepard, D. B., Chong, R. A., López Arriaza, J., Hall, K., Castoe, T. A., Feschotte, C.,
Pollock, D. D., & Mueller, R. L. (2012). LTR retrotransposons contribute to genomic
gigantism in plethodontid salamanders. *Genome Biology and Evolution, 4*(2), 168-
183.
Sun, Y.-B., Zhang, Y., & Wang, K. (2020). Perspectives on studying molecular adaptations of
amphibians in the genomic era. *Zoological Research, 41*(4), 351.
Team, R. (2022). RStudio: integrated development for R. Boston, MA: RStudio. In: Inc.
Team, R. C. (2013). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. *http://www. R-project. org/*.
Wang, J., Itgen, M. W., Wang, H., Gong, Y., Jiang, J., Li, J., Sun, C., Sessions, S. K., & Mueller,
R. L. (2021). Gigantic Genomes Provide Empirical Tests of Transposable Element

Dynamics Models. *Genomics, Proteomics & Bioinformatics*.
doi:https://doi.org/10.1016/j.gpb.2020.11.005
Waters, P. D., Patel, H. R., Ruiz-Herrera, A., Álvarez-González, L., Lister, N. C., Simakov, O.,
Ezaz, T., Kaur, P., Frere, C., Grützner, F., Georges, A., & Graves, J. A. M. (2021).
Microchromosomes are building blocks of bird, reptile, and mammal chromosomes.
Proceedings of the National Academy of Sciences, 118(45), e2112494118.
doi:doi:10.1073/pnas.2112494118
Wu, W., Gao, Y. D., Jiang, D. C., Lei, J., Ren, J. L., Liao, W. B., Deng, C., Wang, Z., Hillis, D. M.,
Zhang, Y. P., & Li, J. T. (2022). Genomic adaptations for arboreal locomotion in Asian
flying treefrogs. *Proc Natl Acad Sci U S A, 119*(13), e2116342119.
doi:10.1073/pnas.2116342119
714

DATA ACCESSIBILITY AND BENEFIT-SHARING

Data Accessibility Statement

Genetic data:

All the genomes used in this study are available on the NCBI Genomes database

(https://www.ncbi.nlm.nih.gov/genome/).

Code:

All original code has been deposited on GitHub and is publicly available at

(https://doi.org/10.5281/zenodo.7679280).

AUTHOR CONTRIBUTIONS

Conceptualization, T.A.K., A.J.C., L.A.O., A.R., and K.C.W.V; methodology, T.A.K, N.D.Y, R.L.M, and

A.R.; formal analysis, T.A.K., M.L.P.; investigation, T.A.K., N.D.Y, R.L.M., K.C.W.V, M.L.P.; resources,

L.A.0. and A.R.; writing – original draft, T.A.K.; writing – review & editing, all authors; project

administration, T.A.K; funding acquisition, T.A.K. and L.F.S.

TABLES AND FIGURES

Table 1. Genome quality measures.

Species	Family	N chrs	Assembly ID	N scaffolds	Geno me size (Gb)	Gap %	Scaffold N50 (kb)	Contig N50 (kb)	% Chrs	% Repea ts	Busco % complet e
Ambystoma mexicanum	Ambystomatidae	14	GCA002915 635.3	27157	28.21	0.013	1205706.7	217.996	94.77	66.26	66.06
Allobates femoralis	Aromobatidae	0	unpublishe d	6658	5.32	0.041	625512.0	879.1	0.00	74.02	81.37
Bombina variegata	Bombinatoridae	0	GCA905336 975.1	4065339	4.68	0.564	2.0	1.9	0.00	56.99	13.03
Rhinella marina	Bufonidae	0	GCA900303 285.1	31391	2.55	0	167.5	167.5	0.00	67.71	85.89
Bufo gargarizans	Bufonidae	11	GCA014858 855.1	747	4.55	1.792	539822.1	1738.3	96.21	74.54	86.74
Bufo bufo	Bufonidae	11	GCA905171 765.1	1307	5.04	0.196	635713.4	3959.8	98.92	77.47	90.11
Hyalinobatrachium fleischmanni	Centrolenidae	0	unpublishe d	227850	4.04	11.337	295403.6	10.5	0.00	61.2	66.27
Oophaga pumilio	Dendrobatidae	0	GCA009801 035.1	125383	3.49	36.043	78.0	5.8	0.00	40.77	67.57
Oophaga sylvatica	Dendrobatidae	0	unpublishe d	15287	5.19	0.109	571702.6	97.8	0.00	81.8	53.47
Geotrypetes seraphini	Dermophiidae	19	GCA902459 505.2	164	3.78	0.146	272612.2	20656.6	99.85	72.51	91.39
Nanorana parkeri	Dicroglossidae	0	GCA000935 625.1	25187	2.05	3.704	1069.1	32.8	0.00	51.39	90.40
Eleutherodactylus coqui	Eleutherodactylida e	13	GCA019857 665.1	105233	2.79	15.115	109467.1	10.9	63.88	53.83	79.11
Engystomops pustulosus	Leptodactylidae	11	GCA019512 145.1	108304	2.56	0.38	172109.2	305.4	75.27	52.32	75.48
Platyplectrum ornatum	Limnodynastidae	0	GCA016617 825.1	148035	1.07	25.246	27.3	4.7	0.00	23.4	43.45
Limnodynastes dumerilii	Limnodynastidae	0	GCA011038 615.1	520896	2.38	10.162	286.0	10.6	0.00	46.25	79.10
Mantella aurantiaca	Mantellidae	0	unpublishe d	135214	3.41	14.489	324858.8	14.1	0.00	52.56	58.57
Leptobrachium ailaonicum	Megophryidae	13	GCA018994 145.1	5370	3.54	0.149	412424.8	821.1	98.22	67.92	81.79
Leptobrachium leishanense	Megophryidae	13	GCA009667 805.1	5302	3.55	0.009	394693.0	1946.3	93.34	70.63	89.55
Spea bombifrons	Pelobatidae	0	GCA009364 475.1	1629896	0.77	0	0.5	0.5	0.00	33.73	3.71
Spea multiplicata	Pelobatidae	0	GCA009364 415.1	49736	1.08	0.235	71.5	30.7	0.00	33.55	76.59
Pipa parva	Pipidae	0	GCA019650 415.1	283592	1.37	0.596	7.2	6.2	0.00	43.01	26.72
Xenopus tropicalis	Pipidae	10	GCA000004 195.4	166	1.45	0.196	153961.3	14634.3	99.86	49.09	94.44
Xenopus laevis	Pipidae	18	GCA017654 675.1	54	2.74	0.061	155250.6	22451.5	99.97	45.21	95.95

Hymenochirus boettgeri	Pipidae	9	GCA019447 015.1	25835	3.21	0.128	293320.9	783.8	82.53	52.72	87.38
Pyxicephalus adspersus	Pyxicephalidae	14	GCA004786 255.1	5411	1.56	10.517	157522.4	30.4	98.54	32.46	87.93
Rana temporaria	Ranidae	13	GCA905171 775.1	555	4.11	0.637	481763.2	6264.8	98.26	69.11	90.72
Lithobates catesbeianus	Ranidae	0	GCA002284 835.2	1544635	6.25	11.584	39.4	6.0	0.00	62.17	44.05
Glandirana rugosa	Ranidae	0	GCA018402 905.1	561726	7.63	10.864	747.6	11.0	0.00	62.38	79.62
Rhinatrema bivittatum	Rhinatrematidae	19	GCA901001 135.2	1330	5.32	0.638	486875.1	3216.3	97.36	67.86	91.58
Scaphiopus couchii	Scaphiopodidae	0	GCA009364 435.1	1378905	0.48	0	0.4	0.4	0.00	26.94	0.70
Scaphiopus holbrookii	Scaphiopodidae	0	GCA009364 455.1	1509266	0.71	0	0.5	0.5	0.00	31.42	2.60
Microcaecilia unicolor	Siphonopidae	14	GCA901765 095.2	1081	4.69	1.012	376147.1	3661.5	97.06	69.21	90.89

Figure 1. Phylogenetic tree of amphibian families. Amphibian families with representative genomes are highlighted and numbers indicate genome counts per family. (Green) anurans, (blue) caecilians, and (orange) salamanders. *Engystomops pustulosus* (Family) image was taken by B. Gratwicke, other amphibian images were licenced to T. Kosch by Adobe Stock and Shutterstock.

749
750 **Figure 2**. BUSCO (Benchmarking Universal Single-Copy Orthologs) assessment results for amphibian
751 genomes.

758
759 **Figure 3**. Repeat content across the amphibian genomes.

769
770 **Figure 4: Relationship between proportion of total transposable elements per genome (body size**
771 **corrected and log transformed) and Bio18 (precipitation of the warmest quarter, log transformed).**
772 **32** amphibian genomes have a higher proportion of transposable elements with increasing wet and
773 warm conditions. Colours range from purple - low to medium- green to orange - high values.

780
781
782 **Figure 5**. Synteny plot of BUSCOs (Benchmarking Universal Single-Copy Orthologs) for chromosome-
783 level anuran genomes. The phylogenetic tree was created with Timetree.org. The reference genome
784 is *Xenopus tropicalis*. Tetraploid anuran *Xenopus laevis* has been excluded. *Indicate inverted
785 chromosomes. Chromosomes without BUSCOs were excluded from the plot.
786