The function of ER-phagy receptors is regulated through phosphorylation-dependent
ubiquitination pathways.

Rayene Berkane[1,2], Hung Ho-Xuan[1,2, #], Marius Glogger[3, #], Pablo Sanz-Martinez[1,2, #], Lorène
Brunello[1,2], Tristan Glaesner[2], Santosh Kumar Kuncha[1,2], Katharina Holzhüter[4], Sara Cano-
Franco[1,2], Viviana Buonomo[5], Paloma Cabrerizo-Poveda[1,2], Ashwin Balakrishnan[3], Georg
Tascher[1], Koraljka Husnjak[1], Thomas Juretschke[7], Mohit Misra[1,2], Alexis González[1], Volker
Dötsch[4], Paolo Grumati[5,6], Mike Heilemann[3] and Alexandra Stolz[1,2,*]

Affiliations:
[1] Institute of Biochemistry II (IBC2), Faculty of Medicine, Goethe University, Frankfurt am
Main, Germany
[2] Buchmann Institute for Molecular Life Sciences (BMLS), Goethe University, Frankfurt am
Main, Germany
[3] Institute of Physical and Theoretical Chemistry, Goethe University, Frankfurt am Main,
Germany
[4] Institute of Biophysical Chemistry and Center for Biomolecular Magnetic Resonance,
Goethe University, Frankfurt am Main, Germany
[5] Telethon Institute of Genetics and Medicine (TIGEM), Pozzuoli, Italy
[6] Department of Clinical Medicine and Surgery, Federico II University, Naples, Italy
[7] Institute of Molecular Biology, Mainz, Germany
[#] Equal contribution
[*] Corresponding author

Keywords: endoplasmic reticulum, selective autophagy, FAM134B, FAM134C, lysosomal
degradation, ER remodeling, ER fragmentation, ER homeostasis, stress response, ATM, ATR,
CHEK1, CK2, mTOR, ubiquitination.

Abbreviations:
Akt: RAC-alpha serine/threonine-protein kinase
ATM: Ataxia-Telangiectasia Mutated
ATR: Ataxia-Telangiectasia and Rad3 related
BafA1: Bafilomycin A1
BiCAP: bimolecular complementation affinity purification

CAMKII: Ca²□/calmodulin-dependent protein kinase II

Chk1: Checkpoint kinase1

CK2: Casein Kinase 2

CTRL: control

DDR: DNA damage response

DMEM: Dulbecco's modified Eagle's medium

DOX: doxycycline

E1: ubiquitin-activating enzyme

ER: Endoplasmic Reticulum

FBS: fetal bovine serum

FCS: fetal calf serum

GABARAP: gamma aminobutyric acid receptor-associated protein

GFP: green fluorescent protein

HA: human infuenza hemmaglutinin

ITC: isothermal titration calorimetry

KD: knockdown

Ki: inhibition constant

KO: knockout

LC3: MAP1LC3 microtubule-associated proteins 1A/1B light chain 3

LIR: LC3 interaction motif

MS: mass spectrometry

MEF: mouse embryonic fibroblast

mTOR: mechanistic target of rapamycin

MALDI: matric-assisted laser desorption/ionization

NEM: N-ethylmaleimide

PI3KK: phosphatidylinositol 3-kinase-related kinase

PTM: post-translational modification

RFP: red fluorescent protein

RHD: reticulon homology domain

SMLM: single-molecule localization microscopy

STED: stimulated emission depletion

TUBE: tandem ubiquitin binding entity

WT: Wild-type

Abstract

Selective autophagy of the endoplasmic reticulum (ER), known as ER-phagy, is an important regulator of ER remodeling and essential to maintain cellular homeostasis during environmental changes. We recently showed that members of the FAM134 family play a critical role during stress-induced ER-phagy. However, the mechanisms on how they are activated remain largely unknown. In this study, we analyzed phosphorylation of FAM134 as a trigger of FAM134-driven ER-phagy upon mTOR (mechanistic target of rapamycin) inhibition. An unbiased screen of kinase inhibitors revealed that CK2 is essential for FAM134B- and FAM134C-driven ER-phagy after mTOR inhibition. Furthermore, we provide evidence that ER-phagy receptors are regulated by ubiquitination events and that treatment with E1 inhibitor suppresses Torin1-induced ER-phagy flux. Using super-resolution microscopy, we show that CK2 activity is essential for the formation of high-density FAM134B and FAM134C clusters. In addition, dense clustering of FAM134B and FAM134C requires phosphorylation-dependent ubiquitination of FAM134B and FAM134C. Treatment with the CK2 inhibitor SGC-CK2-1 or mutation of FAM134B and FAM134C phosphosites prevents ubiquitination of FAM134 proteins, formation of high-density clusters, as well as Torin1-induced ER-phagy flux. Therefore, we propose that CK2-dependent phosphorylation of ER-phagy receptors precedes ubiquitin-dependent activation of ER-phagy flux.

Introduction

The endoplasmic reticulum (ER) is a continuum of membrane structures that dominate a considerable fraction of cell volume and form a central hub for many molecular events such as protein folding, lipid and steroid synthesis, as well as calcium storage (1). The multifunctional nature of this organelle requires the protection of its integrity under multiple stress conditions to achieve optimal function of individual proteins and the entire organelle (2).

Selective turnover of the ER by autophagy (ER-phagy) has emerged as a major quality control pathway ensuring programmed renovation of the ER through elimination of discrete fragments sequestered into autophagosomes and degraded through the lysosomal machinery (macroautophagy) (3). Receptor-mediated macroautophagy (in the following called autophagy) relies on autophagy receptors selectively recognizing cargo, linking it to the autophagic membrane, and thereby promoting cargo sequestration in autophagosomes. To facilitate this function, most ER-phagy receptors are linked to the ER through a transmembrane domain (cargo binding) and also have an LC3 interaction region (LIR) that allows them to interact with members of the LC3 / GABARAP (MAP1LC3 microtubule-associated proteins 1A/1B light chain 3 (LC3) / gamma aminobutyric acid receptor-associated protein) protein family on the autophagic membrane (4). So far, eight membrane-embedded ER-phagy receptors have been identified: FAM134B/RETREG1 (5), SEC62 (6), RTN3L (7), CCPG1 (8), ATL3 (9), TEX264 (10), FAM134A/RETREG2 and FAM134C/RETREG3 (11), which participate in basal ER turnover, ER reshaping after stress-related expansion, as well as lysosomal degradation of selected ER proteins (2).

Members of the FAM134 family carry a conserved LIR motif located at their C-terminus, and their membrane domain consists of a reticulon homology domain (RHD), comprised of two hydrophobic regions embedded in the membrane that are bridged by a flexible cytoplasmic linker (12). We recently showed that the three FAM134 proteins have shared and distinct functions and differ in the dynamics of their RHDs (11). All FAM134 proteins provide critical functions in maintaining ER homeostasis and their individual absence results in massive deformation of the ER network and swelling of the ER. Although quite similar, the three paralogues may operate differently: while FAM134B is constitutively active, FAM134A and FAM134C predominantly exist in an inactive state under basal conditions (11). Upon ER stress, all three receptors can induce significant fragmentation and degradation of the ER via the lysosome. We also demonstrated that FAM134C can act in concert with FAM134B and

enhances its activity in degrading misfolded pro-collagen (11). The cooperative performance

of these two paralogues is likely to be tightly regulated through processes including post-

translational modifications (PTMs) and structural organization. Along these lines, it was

recently shown that phosphorylation of FAM134B-RHD, within the flexible cytoplasmic

linker by CAMKII (Ca^{2+}/calmodulin-dependent protein kinase II), drives the oligomerization

of FAM134B and promotes ER fragmentation prior to ER-phagy (13). The question of how

phosphorylation mechanistically drives oligomerization and if other kinases are involved in

this regulatory pathway remains unknown.

Here, we show that the activity of both FAM134B and FAM134C can be modulated by

inhibiting mTOR (the master kinase of autophagy) with Torin1 and identify several

downstream kinases using a chemical screening approach. In this unbiased screen, we identify

CK2 downstream of Torin1-induced and FAM134B-, as well as FAM134C-driven ER-phagy.

Phosphorylation sites within the RHD of FAM134 proteins were important for ER-phagy

induction by Torin1 treatment. Phosphorylation of FAM134B and FAM134C was also a

prerequisite for subsequent ubiquitination of FAM134B and FAM134C and the formation of

high-density receptor clusters. This suggests phosphorylation-dependent ubiquitination of

both FAM134 proteins as the driving force behind high-density clustering of FAM134B and

FAM134C at active ER-phagy sites.

Results

ER-phagy driven by FAM134C and FAM134B is activated upon inhibition of mTOR

Our recent findings showed that FAM134C is an ER-phagy receptor that works in concert

with FAM134B and that overexpressed FAM134C is inactive under basal conditions and

activated upon starvation (11). We hypothesized that phosphorylation events activate ER-

phagy and tested the impact of mTOR inhibition on ER-phagy flux driven by FAM134C and

FAM134B using the ER-phagy reporter ssRFP-GFP-KDEL (10) (later referred to as the

KDEL reporter). GFP is rapidly quenched within the acidic environment of the lysosomes,

while the fluorescence of RFP is preserved. Therefore, changes in the RFP/GFP ratio (total

integrated fluorescence) can be used to compare the ER-phagy flux between conditions

independent of the monitored cell number (Fig. 1A). Several ER-phagy reporters have been

identified so far (5–11) and activation of any of these receptors can, in principle, affect KDEL

monitored flux. To specifically measure FAM134-driven ER-phagy flux and minimize the

impact of other ER-phagy receptors, we used U2OS cell lines constitutively expressing the

KDEL reporter (CTRL) and stably overexpressing HA-FLAG-FAM134B (FAM134B) and
HA-FLAG-FAM134C (FAM134C) in a doxycycline (DOX)-inducible manner (11). Under
basal conditions (DMSO), overexpression of FAM134B or FAM134C slightly increased ER-
phagy flux compared to control cells (Fig. 1B). Upon treatment with 250 nM Torin1, the flux
of ER-phagy increased strongly over time in all cell lines tested and specifically upon
overexpression of FAM134B or FAM134C. Induction of ER-phagy flux was approximately
3-fold higher in FAM134B and FAM134C overexpressing cells compared to CTRL cells,
indicating that under these conditions the vast majority of measured ER-phagy flux results
from FAM134-driven ER-phagy (Fig. 1B). We then investigated whether loss of endogenous
Fam134b or Fam134c decreases the amount of Torin1-induced ER-phagy (Fig. 1C, D). For
that, we used mouse embryonic fibroblasts (MEFs) isolated from *Fam134b* and *Fam134c*
single knockout (KO) mice complemented with the ssRFP-GFP-KDEL reporter (11). Our
data confirms that the contribution of FAM134 proteins to mTOR-mediated ER-phagy is an
endogenous function of FAM134. The difference between ER-phagy flux in cells
overexpressing FAM134B or FAM134C and the CTRL cell line represents Torin1-mediated,
FAM134-driven ER-phagy flux and can serve as a screening window to identify factors
important for FAM134 activation (Fig. 1B). Notably, the mTORC1 inhibitor Rapamycin has a
similar, but weaker impact on ER-phagy flux (Fig. S1A). All together, these findings suggest
that FAM134s function is regulated through mTOR inhibition.
189
Kinases involved in mTOR-mediated and FAM134-driven ER-phagy
To understand which downstream kinases regulate FAM134B and FAM134C-driven ER-
phagy upon mTOR inhibition, we screened a library of approximatively 100 highly selective
kinase inhibitors (chemogenomic set of EUbOPEN; see EUbOPEN.org) and available
negative controls for their potency to prevent ER-phagy induction. These target-optimized
inhibitors have a K_i in the nanomolar range and the screening was performed uniformly with
1 µM final compound concentration. After 24 hours of DOX treatment to induce FAM134B
and FAM134C overexpression, respectively, cells were preincubated for 1 hour with kinase
inhibitors to block target kinase activity (-1h) and subsequently treated with 250 nM Torin1 to
induce ER-phagy (0h). Time-point measurements of the ratio of total integrated red (RFP) and
green (GFP) fluorescence, as well as cell confluence (phase), were taken every 2 hours over a
total period of 48 hours. Heatmaps summarizing the time-resolved screening results on ER-
phagy flux and cell growth are shown in Fig. 1E and Fig. S1B, respectively. Inhibition of
several kinases prevented Torin1-induced, FAM134-driven ER-phagy, indicated by the loss

of red color toward higher time points in the heatmap (Fig. 1E). To present the screening

results in a visually receptive format, the data were transformed and normalized: the

difference in the RFP/GFP ratio between the untreated and Torin1-treated cells was defined as

100% (Fig. S1C). Early time points (<12h) were excluded due to the small screening window

and subsequent misinterpretation of the early data points. The impact of individual

compounds was represented as % of the relative reduction of FAM134-mediated ER-phagy

per time point (Fig. S1C). Consequently, ER-phagy relevant compounds (HITs) were defined

as compounds causing >20% relative reduction in ER-phagy flux over a minimum time

period of 12 hours (6 time points). We identified 10 kinase inhibitors with impact on

FAM134B and 15 kinase inhibitors with impact on FAM134C activation, with 8 overlapping

HITs (Fig. S1D and Fig. S2A). From identified HITs, we were specifically interested in those

that target kinases closely related to known mTOR signaling pathways and not inducing

obvious cell death. Five candidate compounds corresponded to these criteria: VE-822, KU-

60019, CHIR-124, MK-2206 2HCL and SGC-CK2-1 targeting ATR, ATM, Chk1, Akt, and

CK2, respectively (Fig. S2A, B). These kinases were reported to be related to or part of the

mTOR signaling axis. For instance, ATM and ATR, which orchestrate the DNA damage

response (DDR), are members of the phosphatidylinositol 3-kinase (PI3K)-related family

(PI3KKs) that also comprises mTOR (13,14). CK2 was previously associated with mTOR, as

its inhibition was shown to have an anticancer effect through downregulation of the

AKT/mTOR signaling pathway (15). For validation, selected HITs were retested for dose-

response. Five selected HIT compounds, VE-822, KU-60019, CHIR-124, MK-2206 2HCL

and SGC-CK2-1 were confirmed and blocked FAM134B, as well as FAM134C-mediated ER-

phagy in a dose-dependent manner (Fig. 2A, B). We also tested the impact of selected HIT

compounds on Torin1-induced ER-phagy in U2OS CTRL cells. Here, the screening window

is relatively narrow. However, HIT compounds prevented Torin1-induced ER-phagy to some

extent, indicating endogenous regulatory function of the target kinases (Fig. S2C).

mTOR inhibition triggers the phosphorylation of FAM134B and FAM134C by CK2

Parallel to the phenotypic chemical screening, the phosphorylation status of FAM134B and

FAM134C was analyzed using a mass spectrometry (MS) approach (Fig. 3A, B). Several

phosphorylation sites on highly conserved residues were identified under basal or Torin1-

induced conditions (Fig. 3B, Fig. S3A) including previously reported functional sites within

FAM134B (16). While this in principle supports the robustness of our MS approach, we were

surprised that an inhibitor of the reported CAMKII kinase was absent from the list of our HIT

compounds (Fig. 1E, Fig. S2A). The previous report tested KN-93, a CaMKII inhibitor with a

K_i of 2.58 µM at 10 µM (16). Although KN-93 is relatively selective for CaMKII, it has been

found to have several off-target effects at higher concentrations that can indirectly affect

FAM134-dependent ER-phagy flux, including effects on Ca^{2+} channels, Ca^{2+} release, and

calmodulin activity (17). Our screen included a more potent inhibitor of the CaMKII kinase

with a K_i of 0.37 µM (KN-93 phosphate). We reanalyzed both inhibitors in dose-response and

found doses of >4 µM KN-93 to impact ER-phagy flux over time. KN-93 phosphate showed

no effect (defined as >20% relative reduction over min. 12h) up to 2 µM (Fig. S2D).

Identified phosphorylation sites were also analyzed using the motif-match analysis tool

PHOSIDA (18) to predict kinases. Casein kinase 2 (CK2) was the most frequently predicted

kinase for our data set, suggesting that CK2 can directly phosphorylate FAM134 proteins

(Fig. S3B). The prediction nicely correlated with our screening results, where the CK2

inhibitor SGC-CK2-1 reduced up to 60% and 70% of Torin1-induced ER-phagy in FAM134B

and FAM134C overexpressing cells, respectively (Fig. 2A, B). To further validate our

hypothesis that CK2 directly phosphorylates FAM134 proteins, we analyzed the ability of

purified CK2 to phosphorylate FAM134B *in vitro*. A reaction of FAM134B and CK2

(subunits CK2A1 and CK2A2) purified from bacteria showed a shift of FAM134B on a phos-

tag gel, indicating CK2-dependent phosphorylation (Fig. 3C). To validate the *in cellulo*

phosphorylation sites within the RHD domain of FAM134B, the assay was repeated with

purified FAM134B-RHD and analyzed by MS (Fig. S3C). We could identify one of the

potentially three phosphorylation sites *in vitro*, indicating that CK2 is indeed capable of

directly phosphorylating FAM134B (Fig. 3D, E, Fig. S3D).

CK2 activity positively impacts FAM134-driven ER-phagy

For validation of the functional impact of the identified phosphorylation sites, stable inducible

cell lines expressing mutated FAM134B S149,151,153A or FAM134C S258A (phospho-

mutants) were generated and subjected to our ER-phagy flux assay (Fig. 3E, F). The

FAM134C S258A mutation did not impact ER-phagy flux under basal conditions, however,

had a strong negative impact on Torin1-induced ER-phagy flux. Mutation of the FAM134B

phosphorylation sites also negatively affected the ability of FAM134B to increase the ER-

phagy flux after Torin1-induction. At the same time, the FAM134B phospho-mutant

increased basal ER-phagy flux – visible by a slightly higher steady-state red/green ratio at 0h

– indicating stress / a dominant negative effect upon expression of the mutant protein. In line

with these data, loss of endogenous CK2 prevented induction of ER-phagy flux after Torin1

treatment in cells overexpressing FAM134B and FAM134C, respectively (Fig. S3E,F).

We previously showed that FAM134 proteins are involved in the degradation of misfolded

collagen (11). To test our hypothesis that phosphorylation of FAM134 proteins is necessary

for the activation of receptors and the maintenance of a high ER-phagy flux capacity in times

of stress, we reconstituted MEFs lacking Fam134b (*Fam134b* KO) and Fam134c (*Fam134c*

KO), with respective phospho-mutants. In contrast to the wild-type (WT) protein, FAM134

proteins lacking identified phospho-sites were unable to recover the endogenous function of

the WT protein and showed elevated levels of collagen (Fig. 3G-J). Our findings stress the

importance of the identified phosphorylation sites in the activation process of FAM134

proteins.

Phosphorylation by CK2 triggers ubiquitination of FAM134

When immunoprecipitating FAM134C for MS experiments, we noticed dominant bands

detected by anti-HA-antibody (FAM134C) >50kDa on control western blots. This indicated

PTMs beyond phosphorylation that can shift the protein to higher molecular weights. One

possibility was multi- or poly-ubiquitination. To test this hypothesis, we used the tandem

ubiquitin binding entity 2 (TUBE 2) system, which is capable of pulling down mono- and

poly-ubiquitinated proteins (Fig. S4A). Using lysates of cells overexpressing HA-FAM134B

and HA-FAM134C, we were able to identify ubiquitinated species of FAM134B and

FAM134C, respectively (Fig. S4B). The amount of ubiquitinated FAM134 species increased

after treatment with Bafilomycin A1 (BafA1), indicating that ubiquitinated FAM134 proteins

indeed undergo ER-phagy and therefore accumulate after blocking lysosomal degradation

(Fig. S4B).

To test whether ubiquitination is functionally important for FAM134-driven ER-phagy, we

used our ER-phagy flux assay and co-treated cells with Torin1 and different concentrations of

an E1 inhibitor (MLN7243) blocking the first step of ubiquitination. Inhibition of the

ubiquitin system had a negative impact on ER-phagy flux, and higher concentrations of the E1

inhibitor suppressed FAM134-driven ER-phagy to more than 80% (Fig. 4A, B). Along the

hypothesis that FAM134 ubiquitination promotes ER-phagy, a higher amount of ubiquitinated

species was found for endogenous FAM134B and FAM134C (Fig. 4C), as well as in the

overexpression system (Fig. 4D, E) upon Torin1-induction. With a ~3.4-fold increase, the

difference in ubiquitination levels between basal and induced conditions was much more

pronounced for FAM134B than for FAM134C (~1.4-fold) (Fig. 4E).

We then asked the question of whether there is a cross-talk between phosphorylation and

ubiquitination. Therefore, the ubiquitination status of FAM134B and FAM134C after Torin1

treatment was analyzed in the presence and absence of the CK2 inhibitor. High molecular

species disappeared after co-treatment with Torin1 and CK2 inhibitor or co-treatment with the

E1 inhibitor (control) and CK2 inhibitor, indicating that CK2-mediated phosphorylation is

indeed a prerequisite for the ubiquitination of FAM134 proteins (Fig. 4C-E). Likewise,

genetic depletion of CK2 prevented ubiquitination of FAM134 proteins under Torin1-induced

conditions (Fig. 4F, G). To exclude the possibility that other CK2-mediated phosphorylation

events are the cause of ubiquitination of FAM134, we also analyzed the ubiquitination status

of our FAM134 phospho-mutants. Indeed, FAM134B and FAM134C phospho-mutants had

no ubiquitin modifications both under basal and Torin1-induced conditions (Fig. S4C).

Oligomeric FAM134B and FAM134C is ubiquitinated in a CK2-dependent manner

Receptor oligomerization is believed to be important for vesiculation / fragmentation of the

ER and subsequently delivery to the lysosome (19,20). Therefore, we investigated the

ubiquitination status of oligomerized FAM134 proteins by performing immunoprecipitation

experiments using the split Venus system (21), which drives and preserves dimerization (Fig.

S4D). Two constructs carrying half of the GFP were generated for FAM134B and FAM134C,

respectively. After transient co-transfection of both constructs, FAM134 proteins oligomerize

and split GFP molecules can fuse to form a complete and stable GFP molecule (Fig. S4D, E).

We immunoprecipitated FAM134B or FAM134C homo-dimers using GPF-Trap beads and

detected for ubiquitin. Again, we were able to detect high-molecular weight species of

FAM134B and FAM134C under basal and induced conditions (Fig. 4H). These may represent

SDS-resistant dimers as suggested previously (16), as well as multi/poly-ubiquitinated

FAM134 proteins. The detected ubiquitination signal completely vanished upon co-treatment

with the CK2 inhibitor, further supporting our claim that CK2-dependent phosphorylation

events are a prerequisite for the ubiquitination of FAM134B and FAM134C.

Phosphorylation increases the size of the FAM134B nanoscale cluster

To refine the impact of FAM134-RHD phosphorylation on clustering we used single-

molecule localization microscopy (SMLM), an optical super-resolution method that achieves

near-molecular resolution (22) and enables a quantitative readout (23) to analyze nanoscale

clustering (50-200 nm) of FAM134 proteins. Optimized sample preparation conditions

consisted of mild HA-FAM134 overexpression (12h of DOX induction), followed by 6h of

treatment with DMSO, Torin1, and Torin1+CK2 inhibitor, respectively. Using antibodies

targeting the HA epitope in combination with accumulation of DNA points in nanoscale

topography (DNA-PAINT) (23), we visualized FAM134 clusters with sub-diffraction spatial

resolution (Fig. 5A, B). A quantitative analysis of FAM134B nanoscale clustering (see

Methods) revealed a unimodal distribution of cluster diameters under basal conditions (mode

= 66 nm, σ = 0.23 nm) (Fig. 5C). Upon Torin1 treatment, a second population emerged at a

larger cluster diameter (mode = 104 nm, σ = 0.24 nm), while the first population was

maintained. Cotreatment with the CK2 inhibitor prevented the formation of the larger

nanoscale cluster population, while retaining the first population of small clusters (mode = 69

nm, σ = 0.23 nm) (Fig. 5C), indicating that the size of FAM134B clusters is mediated and

positively regulated by CK2 activity. A quantitative analysis of the diameters of FAM134C

nanoscale clusters also showed a unimodal, but broader distribution under basal conditions

(mode = 52 nm, σ = 0.52 nm) (Fig. 5D). Unlike FAM134B, Torin1 treatment did not lead to

the emergence of a second population of FAM134C clusters with larger diameter (mode = 53

nm, σ = 0.51 nm) (Fig. 5D). However, statistical analysis revealed a small, yet significant

change in the diameter of the overall cluster, which was reversed by co-treatment with the

CK2 inhibitor (Fig. S5A). Because of the relatively large size of the HA-tag and HA-

antibodies that may hinder efficient labeling of dense protein assemblies, we additionally

generated a stable inducible cell lines expressing FAM134 tagged with the recently developed

ALFA-tag (Fig. S5D) (24). Speed optimized DNA-PAINT microscopy (25) confirmed our

findings that Torin1 treatment increased the average size of FAM134B clusters, while the size

of the FAM134C cluster remained unchanged (Fig. S5E). Using shorter induction times, we

observed a unimodal population of FAM134B cluster diameters for Torin1 treated cells under

these settings (Fig. S5E), indicating that the majority of visualized clusters change from an

inactive to an active ER-phagy site. In addition, these experiments revealed that inhibition of

CK2 not only reverts the size of the FAM134B cluster, but also caused a significant decrease

compared to the basal conditions (DMSO) (Fig. S5E). This presumably represents the small

contribution of endogenous ER-phagy receptors to the visualized clusters, which is only

detected by the advanced ALFA-tag system. The impact of inhibition of CK2 on the

diameters of the FAM134B cluster can be caused by direct (phosphorylation of FAM134B) or

indirect actions (phosphorylation of other regulatory or functional proteins within the ER-

phagy process). To support a direct regulation of FAM134 by CK2, we analyzed the

established phospho-mutants of FAM134B and FAM134C. The cluster diameter distribution

of the phospho-mutant of FAM134B was similar to that of FAM134B WT under basal

conditions (mode = 63 nm, σ = 0.17 nm), however, did not increase after Torin1 treatment

(mode = 65 nm, σ = 0.19 nm) (Fig. 5C). This indicates that mutated phosphorylation sites

have a direct impact on the organization of FAM134B and that CK2-mediated

phosphorylation drives/boosts the formation of larger FAM134B nanoscale clusters after ER-

phagy induction by Torin1. Regarding the wild-type form of FAM134C, we did not observe

an impact of inhibition of CK2 on the diameters of the FAM134C S258A cluster diameters

(Fig 5D). Our data therefore suggests that serine 258 phosphorylation has regulatory functions

that are not impacting the size of FAM134C clusters in our overexpression system.

CK2-mediated phosphorylation is essential for the formation of high-density clusters of

FAM134B and FAM134C

The number of FAM134 proteins within nanoscale clusters may vary between clusters of the

same size, depending on the space between individual FAM134 molecules (Fig. 5A, Fig. 6A).

Close packing of FAM134 proteins (high density) may be required for their ER-phagy

function, e.g., bending of the ER membrane to facilitate efficient fragmentation (12). To

further refine the characteristics of nanoscale clusters, we analyzed the relative protein density

within individual FAM134 nanoclusters by extracting the number of single-molecule binding

events (localizations) for different conditions and phosphorylation states (phospho-mutants)

(Fig. 5E-J). A direct comparison of the WT and FAM134 phospho-mutants showed that

clusters with high numbers of localizations (=high density) were largely absent for the

inactive phospho-mutants (colored dots versus colored triangles). We then defined wild-type

high-density FAM134 clusters by thresholding (>100 localizations/cluster, cluster diameter >

90 nm; grey regions). For FAM134B, the number of high-density clusters (as percentage of

all analyzed clusters) increased after Torin1 treatment (5.6 %) compared to baseline

conditions (2 %) and inhibition of CK2 (3.7 %) (Fig. 5E-G). Interestingly, we measured only

a minor increase in high-density clusters for FAM134C after Torin1 treatment (6.6 % for

DMSO and 7.3%) but a strong decline after CK2 inhibition (2.6 %) (Fig. 5H-J). This

correlates with the ubiquitination status of FAM134C and indicates that FAM134C is

regulated by CK2 also under basal conditions. Subdividing nanoscale clusters into small (30-

90 nm), medium (90-150 nm) and large clusters (> 150 nm) and plotting the relative

frequency distribution of localizations per cluster refined the distribution of high-density

within clusters of the same size (Fig. S5B, C). All findings could be reproduced using ALFA-

tagged FAM134 proteins (Fig. S5F-I).

Together, our data reflect higher FAM134B and FAM134C cluster densities after Torin1-
induced phosphorylation and hint at a dynamic process, where FAM134 clusters reach a
certain size (>90 nm) and then get further activated (by phosphorylation-dependent
ubiquitination) to increase the density of included molecules. Consistent with this hypothesis,
nanoscale clusters formed by phospho-mutants of FAM134B and FAM134C were able to
reach a size of >150 nm, but were unable to further mature to respective high-density clusters
(Fig. 5E-J). Combining this with our findings that inhibition of CK2 after Torin1 treatment
suppresses ER-phagy flux (Fig. 2A, B), we predict that only high-density clusters are
functional active ER-phagy sites that subsequently support transport of ER-fragments to the
lysosome (Fig. 6).
416
Discussion
One of the challenging questions in the field of ER-phagy is how the pathway is fine-tuned
and controlled, starting from the initial trigger to bind cargo and initiate ER curvature towards
ER fragmentation up to the final delivery to the lysosome. It is obvious that ER-phagy
receptors play a critical role along this path, but the molecular mechanisms that drive their
activation and functions remain poorly understood. In this manuscript, we provide evidence
that CK2-mediated phosphorylation within the RHD domain and subsequent phosphorylation-
mediated ubiquitination of FAM134B and FAM134C are essential for the formation of high-
density clusters of FAM134 proteins on the ER membrane and FAM134-driven ER-phagy.
These findings originate from our previous report that FAM134C is in an inactive state under
basal conditions and can serve as an enhancer of the FAM134B-dependent ER-phagy flux
after stress (11). We hypothesized that a phosphorylation cascade upon mTOR inhibition can
serve as an activation signal for FAM134-driven ER-phagy. An unbiased screen against a
chemical library of kinase inhibitors modulating ER-phagy flux revealed ATR, ATM, Chk1,
Akt, and CK2 to be downstream of and essential for induction of FAM134B- and FAM134C-
driven ER-phagy upon mTOR inhibition. Furthermore, we identified potentially dynamically
regulated phosphorylation sites after Torin1 treatment within the RHD domain and adjacent
regions of FAM134B and FAM134C. In line with our chemical screening, the prediction of
the target site suggests that CK2 can phosphorylate FAM134 proteins directly, which we
further showed *in vitro* using purified proteins. CK2 is known to be ubiquitously expressed
and to be a constitutively active kinase that regulates several biological processes (26).
Additionally, CK2 has been linked to autophagy-related diseases such as Alzheimer,
Parkinson and Huntington disease (27). In this study we attributed CK2 a role in orchestrating

FAM134B- and FAM134C-mediated ER-phagy by phosphorylating specific sites within the
RHD domain that drive the clustering of ER-phagy receptors and promote ER-phagy flux.
The phosphorylation of FAM134B and subsequent induction of its oligomerization were
previously reported and at that time linked to CaMKII activity (16). Within our settings,
CaMKII inhibition performed poorly in terms of preventing FAM134B-driven ER-phagy,
while chemical inhibition or knockdown (KD) of CK2 had a strong effect. However, our
findings do not rule out possible cooperation between CK2 and CaMKII kinases in fine-
tuning these events, specifically since CAMKII was previously shown to form a complex with
CK2 (28). Mechanistically, phosphorylation could alter the structure and create new binding
interfaces both towards the membrane and the cytosolic side of FAM134 proteins and recruit
downstream regulatory factors.
Very recently another study has highlighted the role of CK2 in ER-phagy (29). In contrast to
the above results, the study proposes CK2 as a negative regulator of ER-phagy. More
precisely, CK2 was shown to phosphorylate a site close to the LIR motif of FAM134C and it
was suggested that this phosphorylation could prevent its binding to LC3 (29). The presented
in vivo and *in silico* data supported the idea, however, the impact of phosphorylation on the
binding ability of FAM134C to LC3 was not fully characterized in detail *in vitro*. We closed
this gap by performing isothermal titration calorimetry (ITC) measurements with synthetic
FAM134C-LIR peptides in their phosphorylated and non-phosphorylated form (Fig. S6A).
Additionally, we purified GFP-FAM134C-LIR from bacteria, phosphorylated it with purified
CK2 and performed pulldown experiments with purified GST-LC3B (Fig. S6B-E). Both
experimental setups suggest that the FAM134C-LIR behaves as a classical phospho-LIR (30),
where phosphorylation close to the LIR motif enhances binding to LC3 and therefore
promotes ER-phagy. These contrasting evidences suggest a dynamic role of phosphorylation
in ER-phagy and hence needs to be looked at in greater detail. We propose that the
discrepancy between the *in vivo* and *in vitro* data arises from an additional, yet unknown,
regulatory factor with a high affinity for phosphorylated FAM134-LIR that is able to
outcompete LC3 in binding. This hypothesis is specifically intriguing as it would allow
organisms to selectively regulate FAM134-driven ER-phagy in different cell types and
tissues, while using the commonly expressed and constitutively active CK2 kinase as a
mediator.
In addition of being phosphorylated, we show that FAM134B and FAM134C are also
ubiquitinated in an mTOR- and CK2-dependent manner. Phosphorylation of FAM134
proteins may therefore create a new cytosolic binding surface important for the recruitment

and/or activation of a ubiquitin ligase. The accumulation of ubiquitinated FAM134 proteins

upon autophagy inhibition (BafA1 treatment) and the lack of ER-phagy capacity upon Torin1-

induction in an overexpression system as well as at the endogenous level indicate that the

ubiquitination of FAM134 proteins has functional relevance for FAM134-driven ER-phagy.

Therefore, we hypothesized that ubiquitination is needed for the regulation and/or activation

of FAM134-driven ER-phagy. In agreement with this hypothesis, inhibition of the

ubiquitination system (treatment with E1 inhibitor) suppressed FAM134-driven ER-phagy

flux upon mTOR inhibition. Treatment with the CK2 inhibitor abolished ubiquitination of

FAM134B and FAM134C, indicating that ubiquitination is downstream of and dependent on

CK2-mediated phosphorylation of FAM134.

Previous studies suggested that super-clustering of RHD-containing receptors, e.g., Atg40 in

yeast and FAM134B in mammals, generates high-curved regions that subsequently can pinch-

off the ER and get delivered to the lysosome (12,31). Our data implies an interesting chain of

reactions where phosphorylation and subsequent ubiquitination are involved in the regulation

and formation of such super-receptor-clusters. Our data are further supported by a manuscript

published during the revision of this study and reporting on the role of ubiquitination in ER-

phagy regulation (32).

To dissect the consequences of PTMs on oligomerization, we analyzed two parameters of

nanoscale clusters (50-200nm) of either WT or phospho-mutants of FAM134 proteins by

super-resolution microscopy: changes in cluster diameter (size) and differences in

localizations within a cluster of a specific size (density). The nanoscale clusters of FAM134B

WT proteins changed from unimodal (DMSO/basal) to bimodal distribution after induction of

ER-phagy (Torin1). The emerging population after induction clustered around 104 nm (σ =

0.24), indicating that a cluster size of > 90nm is necessary for FAM134B-driven ER-phagy.

Since the appearance of the larger population of FAM134B clusters was suppressed by CK2

inhibition as well as mutation of phosphorylation sites, phosphorylation and/or subsequent

ubiquitination are essential for its formation. On the contrary, the size distribution of the

clusters built by FAM134C WT or phospho-mutants was similar under all tested conditions.

We also found that the density of FAM134 molecules in clusters >90 nm started to vary

dramatically, both in FAM134B and FAM134C-formed clusters in a CK2-dependent manner.

This finding hints at a mechanism where clusters first grow to a certain size and then further

mature by increasing the density of FAM134 molecules within a cluster. An increase in the

local concentration of FAM134 proteins may be necessary to sufficiently bend the ER

membrane to initiate vesiculation. Along this line, the maturation to high-density FAM134B

clusters increased approximately 3 times after ER-phagy induction (2% vs 5.6% relative

abundance) and was greatly suppressed by treatment with CK2 inhibitor (3.7%). Interestingly,

FAM134C was already forming high-density clusters under basal conditions (6.6%) and their

relative abundance only slightly increased upon induction of ER-phagy (7.3%). At the same

time, co-treatment with the CK2 inhibitor considerably reduced its abundance to levels below

basal conditions (2.6%), indicating that FAM134C oligomerization is also regulated by CK2-

dependent phosphorylation. The appearance of high-density clusters mirrors the

ubiquitination status of both FAM134 proteins: the ubiquitination level of FAM134B is low

under basal conditions and strongly increases upon Torin1 treatment, while FAM134C is

already ubiquitinated under basal conditions and its ubiquitination level only increases

slightly upon inhibition of mTOR.

Together, our data show that phosphorylation by CK2 upon mTOR inhibition, as well as

phosphorylation-dependent ubiquitination of FAM134 proteins play essential roles in the

regulation and activation of FAM134B- and FAM134C-driven ER-phagy. In this context, we

suggest that only high-density nanoscale clusters >90nm and >100 locations are functionally

active during ER-phagy. For future studies, we hypothesize that phosphorylation-dependent

ubiquitination facilitates the formation of high-density FAM134 clusters, presumably by

introducing conformational changes between adjacent FAM134 molecules and/or by

recruiting cytosolic factors with oligomerizing properties to the forming ER vesicle. As

FAM134C serves as an enhancer of FAM134B (11), we further hypothesize that pre-existing,

ubiquitinated FAM134C nanoscale clusters may serve as collection points for ER-phagy

substrates and subsequently as nucleation points for FAM134B and that this is the reason

behind the rapid occurrence of a second, larger FAM134B cluster population upon Torin1

treatment. It is also interesting to note that in our system overexpression of FAM134C has a

greater impact on ER-phagy flux than FAM134B, which may imply that FAM134C is also an

enhancer for other, endogenously expressed ER-phagy receptors.

Material and methods

Reagents and antibodies:

Kinase inhibitor library was provided by the SGC Frankfurt donated probes program and

complemented by additional kinase inhibitors. Torin1 was purchased from Bio-techne, 4247

and BafA1 from Cayman Chemical Corporation, 11038. Following primary antibodies were

used for immunoblotting and immunofluorescent staining: rabbit anti-HA tag (Cell signaling

Technology, 3724T), rat anti-HA tag (Roche, 11867432001), rabbit anti-FAM134B (Sigma
Prestige, HPA012077), rabbit anti-FAM134C (Sigma Prestige, HPA016492), mouse anti-
REEP5 (Santa Cruz Biotechnology, sc-393508), rabbit anti-GST (Cell Signaling Technology,
2625S), mouse anti-GFP (Santa Cruz Biotechnology, sc-9996), rabbit anti-CK2 (Cell
signaling Technology, 2656S), mouse anti-ubiquitin P4D1 (abcam, ab7254), rabbit anti-
GAPDH (Cell Signaling Technology, 2118L). Following secondary antibodies were used for
immunoblotting: anti-mouse (Thermo Fisher Scientific, 31326) anti-rabbit (Thermo Fisher
Scientific, 32460) anti-rat (Cell Signaling Technology, 70775). Following secondary
antibodies were used for immunofluorescent staining and super-resolution microscopy: anti-
mouse Alexa Fluor® 532 (Life technologies, A11002), anti-rabbit (AffiniPure, 711-005-152)
and anti-mouse (AffiniPure, 115-005-003), covalently labeled with short oligonucleotide
strands anti-P1 (5´-ATCTACATATT-3´) and anti-P5 (5´-TATGTAACTTT-3´), respectively.
554

Plasmid
pCW57-CMV-ssRFP-GFP-KDEL was a gift from Noboru Mizushima (University of Tokyo,
Tokyo, Japan), (Addgene plasmid # 128257; RRID: Addgene_128257). pLenti CMV GFP
Hygro (656-4) was a gift from Eric Campeau and Paul Kaufman (Addgene plasmid # 17446;
http://n2t.net/addgene:17446; RRID: Addgene_17446). pLenti CMV GFP Hygro (656-4) was
used as the backbone to create pLenti-CMV-Hygro- ssRFP-GFP-KDEL for consecutively
expressing ssRFP-GFP-KDEL. Briefly, pLenti CMV GFP Hygro (656-4) was digested with
two restriction enzymes (BamHI and SalI) to remove the GFP cassette. ssRFP-GFP-KDEL
was then amplified by PCR and cloned into pLenti CMV GFP Hygro (656-4) plasmids using
Sequence and Ligation Independent Cloning (SLIC) with T4 DNA polymerase. iTAP-FLAG-
HA-FAM134B (also referred to as pMCSV-puro-FLAG-HA-FAM134B) and iTAP-FLAG-
HA-FAM134B (also referred to as pMCSV-puro-FLAG-HA-FAM134C) were gifts from Ivan
Dikic (Goethe University Frankfurt am Main, Germany). iTAP-FLAG-HA-FAM134B and
iTAP-FLAG-HA-FAM134C were used as the backbone to create different phosphorylation
mutants, as well as the ALFA-tag expressing cells using quick-change PCR mutagenesis. The
FLAG sequence from the iTAP-FLAG-HA-FAM134B and iTAP-FLAG-HA-FAM134C
vectors was replaced by the sequence of ALFA-tag to create vectors expressing ALFA-
tagged- FAM134B and FAM134C. The amino acid sequence of the ALFA-tag
(SRLEEELRRRLTE) was used based on the previous publication (33).
574

Cell culture

U2OS T-REx and HEK293T cells were maintained in a 37°C incubator with humidified 5%

CO_2 atmosphere and cultured in Dulbecco's modified Eagle's medium (DMEM) (Thermo

Fisher Scientific) supplemented with 10% fetal bovine serum (FBS) (Thermo Fisher

Scientific) and 100 U/ml (1%) penicillin/streptomycin (PS) (Thermo Fisher Scientific).

U2OS T-REx cells stably expressing ssRFP-GFP-KDEL and inducibly individual FAM134

proteins were described previously (11). These cells were treated with 1 µg/ml DOX (Sigma-

Aldrich, D9891) to induce FAM134 protein expression. Prior to each treatment, cells were

plated the day before in order to perform the experiments when cells had a final confluency of

70-80%. HEK293T was purchased from ATCC and was used for transient transfections. For

transient expression, DNA plasmids were transfected with JetPEI (Polyplus, 101000020)

according to manufacturer instructions.

Cell line generation

U2OS T-REx cells were a gift from Stephen Blacklow (Brigham and Women's Hospital and

Harvard Medical School) (34) and were culture using DMEM supplemented with 10% FBS,

1% PS and 15 µg/ml blasticidin S (Thermo Fisher scientific, A1113902). Lentiviral particles

were produced in HEK293T cells using pLenti-CMV-Hygro-ssRFP-GFP-KDEL and

pMD2.G and psPAX2. Retroviral particles were produced in HEK293T using iTAP-FLAG-

HA or or iTAP-ALFA-HA plasmids expressing FAM134B, FAM134C or different

phosphorylation mutants, packaging plasmid (pCMV-Gag-Pol) and the envelope plasmid

(pCMV-VSV-G). The U2OS T-REx cells were plated and propagated in 6-well plates

overnight. On the next day, transduction with viruses was performed. Two days later,

transduced cells with iTAP-FLAG-HA or iTAP-ALFA-HA plasmids were selected using 15

µg/ml blasticidin S and 2 µg/mL puromycin (Invitrogen, Thermo Fisher Scientific, Waltham,

Massachusetts, USA) for 3 passages. U2OS T-REx cells expressing FAM134B, FAM134C or

different mutants were used for transduction with ssRFP-GFP-KDEL retroviruses. The

transduced cells were selected using 15 µg/ml blasticidin S, 2 µg/mL puromycin and 150

µg/ml hygromycin B. Afterward, the cells were sorted for different populations of RFP- and

GFP-expressing cells using SH800S Cell Sorter (Sony Biotechnology, San Jose, CA, USA).

Stable clones were maintained in the same medium as used for selection.

Fam134b and *Fam134c* knockout MEFs were provided by Prof. Christian Huebner (Jena

University) and were culture using DMEM supplemented with 10% FBS. These cells were

obtained from *Fam134b* or *Fam134c* single knockout mice and have been reconstituted with

cDNAs cloned in a retroviral pBABE plasmid. Retrovirus were generated in HEK293T cells

co-transfecting the retroviral vector, containing the selected cDNA, with the packaging plasmid (viral Gag-Pol) and the envelope plasmid (VSV-G). Lentivirus were also generated co-transfecting HEK293T cells with the pCW57-CMV-ssRFP-GFP-KDEL (Addgene #128257) vector together withpPAX2 packaging plasmid and pMD2.G envelope plasmid.

Protein purification

The recombinant GFP-FAM134C-LIR protein was over-expressed in *E. coli* strain BL21(DE3). Briefly, primary cultures were grown overnight followed by secondary cultures in LB media containing appropriate antibiotics at 37°C. The secondary cultures of *E.coli* were grown up to 0.6 $OD_{(600)}$ and induced the cells with 0.25 mM IPTG for 18 hours at 18°C. The cultures were harvested by centrifugation at 8000 rpm for 8 minutes. Resulting cell pellets were lysed using sonicator. The 6X-Histidine tagged proteins were purified using the Ni-NTA affinity chromatography method using solution containing 200 mM NaCl and 100 mM Tris pH 8.0. An imidazole gradient of 20 mM to 250 mM was used to elute the bound protein. Eluted fractions containing protein were pooled and concentrated to reach an appropriate concentration and stored at -20°C in buffer containing 50% glycerol. Purification of the FAM134B-RHD domain has been published before (35).

Endogenous pulldown of FAM134 proteins

HEK293T cells were plated in 15cm dishes and subjected to following treatments (DMSO (Sigma-Aldrich), and 250 nM Torin1 (Bio-techne) in combination with 1 µM CK2 inhibitor (SGC donated probes). 3 copies of each individual condition were harvested and pulled to be then lysed in 1ml of ice-cold RIPA buffer: 10mM Tris-HCl, (pH 8.0), 1mM EDTA, 0.5mM EGTA, 1% (v/v) Triton X-100, 0.1% SDS, 140 mM NaCl, cOmplete protease inhibitor cocktail (Roche), phosphatase inhibitor cocktail 1 and 2 (Sigma-Aldrich). Lysates were cleared by centrifugation at 20000 x g for 15 min at 4°C. 3 mg of proteins were subsequently incubated for 1h with 10 µl of FAM134B or FAM134C antibodies (Sigma Prestige-HPA012077 and -HPA016492, respectively) and then overnight at 4°C with A/G PLUS agarose beads (Santa Cruz Biotechnology) with constant mixing. Beads were then washed 3 times in wash buffer (10 mM Tris-HCl (pH 8.0), 1 mM EDTA, 0.5 mM EGTA, 1% (v/v) Triton X-100, 0.1% SDS, 250 mM NaCl, cOmplete protease inhibitor cocktail (Roche), phosphatase inhibitor cocktail 1 and 2 (Sigma-Aldrich)), resuspended in SDS sample buffer and boiled at 95°C for 10 min. Beads were collected by centrifugation at $2500 \times g$ for 2 min

643 and supernatants were loaded on SDS-PAGE and western blot was performed using the
644 indicated antibodies.
645

In vitro and in cellulo GFP-Trap pulldown
For *in cellulo* experiment, HEK293T cells were transiently co-transfected with two constructs
carrying either V1-FAM134B and V2-FAM134B or V1-FAM134C and V2-FAM134C each
independently tagged with half GFP at the N-terminus. Following 6h treatment with the
indicated conditions, cells were washed with 1x PBS and lysed in ice-cold RIPA buffer: 10
mM Tris-HCl, (pH 8.0), 1mM EDTA, 0.5mM EGTA, 1% (v/v) Triton X-100, 0.1% SDS,
140mM NaCl, cOmplete protease inhibitor cocktail (Roche), phosphatase inhibitor cocktail 1
and 2 (Sigma-Aldrich) and deubiquitinase inhibitor (N-ethylmaleimide, NEM). Samples were
incubated on ice for 15 min and centrifuged at $20000 \times g$ for 15 min at 4°C. GFP-Trap beads
(Chromotek) were equilibrated and washed 3 times in 500 µL of ice-cold RIPA buffer and
were spun down at $2500 \times g$ for 2 min at 4°C. In total, 1 mg of lysates was added to 5 µl of
equilibrated GFP-Trap beads and incubated overnight 4°C with constant mixing. Tubes were
spun at $2500 \times g$ for 3 min at 4°C. GFP-Trap beads were washed three times with 500 µl ice-
cold lysis buffer. 50 µL of 1X SDS sample buffer was then added to the GFP-Trap beads and
were boiled for 10 min at 95°C. Beads were collected by centrifugation at $2500 \times g$ for 2 min
and SDS-PAGE was performed with the supernatant.
For *in vitro* experiment, purified proteins (1µM GST-LC3B, 1 µM GFP-FAM134C-LIR or 1
µM of phosphorylated GFP-FAM134C-LIR) were mixed with 100 mM NaCl, 100 mM Tris
pH 7.5 and 5 µl of equilibrated GFP-Trap beads (Chromotek) and incubated overnight at 4°C
with constant mixing. Samples were then washed three times with 500 µl ice-cold wash buffer
(300 mM NaCl, 100 mM Tris pH 7.5) and processed as described in the *in cellulo* experiment.
667

Tandem Ubiquitin Binding Entities 2 (TUBE 2) assay
U2OS cells expression FAM134 proteins priorly subjected or not to KD experiments were
induced with 24h DOX (1 µg/ml) treatment. Following treatments (DMSO, Torin1, CK2
inhibitor, BafA1), confluent 10 cm dishes of U2OS cells were harvested in 500 µL of ice-cold
lysis buffer: 50 mM Tris-HCl (pH 8.0), 120 mM NaCl, 1% (v/v) NP-40, cOmplete protease
inhibitor cocktail, phosphatase inhibitor cocktail 1 and 2. Lysates were cleared by
centrifugation at 20000 x g for 15 min at 4°C and 1mg of proteins was incubated overnight at
4°C with 10 µL of agarose-TUBE 2 resin. Reactions were rotated overnight at 4°C, washed

three times with NP-40 buffer and analysed by SDS-PAGE. Anti-HA antibody was used to

detect HA-tagged FAM134 proteins and ubiquitinated species.

Immunoblotting of total cell lysates

Following KD experiment, U2OS cells were harvested after removing the culture medium and

washing with 1X PBS and lysed in lysis buffer (150 mM NaCl, 50 mM Tris-HCl pH 7.5, 10

mM NaF, 1 mM EGTA, 5 mM MgCl$_2$, 0.1% SDS) supplemented with protease and

phosphatase inhibitors. Insoluble cell components were separated at $20000 \times$ g for 15 min at

4°C and total protein concentration was assessed using BCA assay (Thermo Fisher

Scientific). Protein denaturation was performed in SDS sample buffer at 95°C for 10 min.

Protein lysates were resolved in SDS-PAGE gels and transferred to PVDF (Millipore, 0.2 µm)

membrane. Membranes were saturated in PBST 0.1% Tween-20 containing 5% BSA and

incubated overnight at 4°C with the specific primary antibody.

In Cellulo Mass Spectometry Identification of Phospho-Serines

U2OS cell lines stably expressing FAM134 proteins were cultured in 15 cm dishes up to 70%

confluency in the presence of 1 µg/ml DOX prior to any treatment. Each replicate from every

condition consisted of 2x 15 cm dishes lysates (50 mM Tris PH 7.4, 150 mM NaCl, 1% NP-

40, fresh protease and phosphatase inhibitors) pooled together. The purification of FAM134

proteins from each condition was performed via anti-HA antibody-conjugated beads. To

remove as many impurities and weak interactors as possible, 6 washing steps were performed.

For the first two washes, 50 mM Tris PH 7.4, 500 mM NaCl, 1% NP-40 containing

phosphatase inhibitor and Roche protease inhibitor tablets was used. The following two

washes were performed with 50 mM Tris PH 7.4, 150 mM NaCl and all inhibitors. The last

two washes were performed with 1X PBS to remove all traces of detergent prior to MS

processing. Beads were incubated with SDC buffer (2% SDC, 1m M TCEP, 4 mM CAA, 50

mM Tris pH 8.5) at 65°C for 30 minutes. Proteins were digested with 500 ng of trypsin

(Promega) and LysC (Wako Chemicals). Digestion was stopped with 1% t rifluoroacetic acid

(TFA) in isopropanol. Following digestion, SDB-RPS columns with a capacity of 10 µg/14G

plug were utilized for sample clean up. Samples were analyzed on a Q Exactive HF coupled

to an easy nLC 1200 (Thermo Fisher Scientific) using a 35 cm long, 75 µm ID fused-silica

column packed in house with 1.9 µm C18 particles (Reprosil pur, Dr. Maisch), and kept at

50°C using an integrated column oven (Sonation). Peptides were eluted by a non-linear

gradient from 4-28% acetonitrile over 45 minutes and directly sprayed into the mass-

spectrometer equipped with a nanoFlex ion source (Thermo Fisher Scientific). Full scan MS

spectra (350-1650 m/z) were acquired in Profile mode at a resolution of 60,000 at m/z 200, a

maximum injection time of 20 ms and an AGC target value of 3×10^6 charges. Up to 10 most

intense peptides per full scan were isolated using a 1.4 Th window and fragmented using

higher energy collisional dissociation (normalized collision energy of 27). MS/MS spectra

were acquired in centroid mode with a resolution of 30,000, a maximum injection time of 110

ms and an AGC target value of 1×10^5. Single charged ions, ions with a charge state above 5

and ions with unassigned charge states were not considered for fragmentation and dynamic

exclusion was set to 20 s.

In Vitro Mass Spectrometry Identification of Phospho-Serines

Three technical replicates containing each 1 µM of the purified RHD region of FAM134B

with 1 µM of each purified subunit of CK2 (CSNK2A1 and CSNK2A2) were incubated for

2h at 37°C. ATP was added at a final concentration of 10 mM and mixed with CK2 buffer (20

mM Tris-HCl pH 7.5, 50 nM KCl, 10 mM $MgCl_2$). The negative controls had no CK2 added

to them. Samples were processed and analyzed for MS as previously discussed with the

exception of a lower amount of trypsine/LysC (200 ng).

Mass spectrometry data processing and analysis

MS raw data processing was performed with MaxQuant (v 2.1.0.0) and its in-build label-free

quantification algorithm MaxLFQ applying default parameters (36). Acquired spectra were

searched against the human reference proteome (Taxonomy ID 9606) downloaded from

UniProt (21-05-2022; 82.492 protein count), self-made fasta file containing the precise

sequence of the RHD construct used for the *in vitro* experiment, and a collection of common

contaminants (244 entries) using the Andromeda search engine integrated in MaxQuant (37).

Identifications were filtered to obtain false discovery rates (FDR) below 5% for both peptide

spectrum matches (PSM; minimum length of 7 amino acids) and proteins using a target-decoy

strategy (38).

The MS/MS spectra for the modified and best-localized peptides were taken from the

visualization option of MaxQuant software. All the tables in Fig S3, show the different

confidence scores, posterior error probability (PEP), and localization probability. It also

shows the positions they hold within the WT full length FAM134B protein and in which

samples the peptides were found.

ER-phagy flux assay

U2OS T-REx FLAG-HA-FAM134s stable and inducible cell lines were infected with lentivirus carrying the pCW57-CMV-ssRFP-GFP-KDEL (Addgene #128257). We previously deleted the tetracycline response element, via PCR, from ssRFP-GFP-KDEL in order to have constitutive expression. Wild-type and *Fam134*s KO MEFs were infected with pCW57-CMV-ssRFP-GFP-KDEL (Addgene #128257) in order to generate ER-phagy reporter inducible cell lines. Fluorescence of GFP and RFP, as well as cell confluence (phase), were monitored over time via the IncuCyte® S3 (Sartorius, Germany) in 384-well format. 1500 cells (U2OS) per well were seeded in 50 µl DMEM media, supplemented with 10% FBS, 1% PS, 2 µg/ml puromycin, 15 µg/ml blasticidin S, 1 µg/ml DOX and incubated for 24h before treatment. For indicated treatments, 50 µl of media containing indicated treatments were added. Between the actual treatment and the first scan point (0hrs) appx. 5-10min passed for technical reasons. Screens in all three channels were taken at indicated time points following the treatment. ER-phagy flux was monitored through changes in the ratio of the total fluorescence intensity of RFP/GFP. Each point represents the averaged ratio of data obtained from three individual wells (technical replicates).

Antibody-oligonucleotide conjugation for DNA-PAINT

Secondary antibodies donkey anti-rabbit (AffiniPure, 711-005-152) and goat anti-mouse (AffiniPure, 115-005-003) were covalently labeled with short oligonucleotide strands anti-P1 (5´-ATCTACATATT-3´) and anti-P5 (5´-TATGTAACTTT-3´), respectively, via DBCO-sulfo-NHS ester chemistry. In brief, concentrated secondary antibodies were incubated with a 20-fold molar excess of DBCO-sulfo-NHS ester (Jena Bioscience, CLK-A124-10) for 90 min at 4°C. Excess reagent was removed using spin filters (Merck, Germany) and docking strand labeled antibodies were incubated with a tenfold molar excess of azide-functionalized DNA-docking strands overnight at 4°C. Samples were then filtered through Amicon centrifugal filters (100.000 MWCO) to remove unbound oligonucleotides and change storage buffer to PBS. Finally, antibody-DNA complexes were concentrated to 5 mg/mL and stored at 4°C.

Sample preparation for DNA-PAINT

For DNA-PAINT experiments, U2OS cells were seeded in fibronectin-coated Ibidi µ-slide VI chambers at a density/confluence of 60% and FAM134s protein expression was induced with 1 µg/ml DOX overnight. Cells were chemically fixed with prewarmed (37°C) 4% methanol-free formaldehyde (Sigma-Aldrich, Germany) in PBS for 20 min followed by three washing

steps with PBS. Primary antibodies rabbit anti-HA (1:500) was subsequently added in
permeabilization/blocking buffer (10% fetal calf serum (FCS) (Gibco), 0.1% saponin (Sigma-
Aldrich)) and incubated for 60 min at room temperature. Excess primary antibodies were
removed from the chambers by three washing steps with PBS and custom docking-strand
labeled secondary antibodies (1:100 in permeabilization /blocking buffer) were added to the
chambers. After incubation for 60 min at room temperature, samples were washed thrice to
remove excess secondary antibodies. Finally, samples were post-fixed using 4% methanol-
free formaldehyde in PBS for 10 min at room temperature followed by three washing steps
with PBS. For colocalization experiments, fiducial markers were used for image registration
(125 nm gold beads (Nanopartz, USA). Gold beads were sonicated for 10 min and diluted
1:30 in PBS. Diluted samples were again sonicated for 10 min and 100 µL of the solutions
were added to the microscopy chambers. After settlement of the gold beads (5 min), the
samples were washed thrice with PBS. Cell chambers were finally assembled to a
microfluidic device (Bruker, USA) for automated buffer exchange. Prior to image acquisition,
ATTO-655 labeled imager Strands P1 or P5 (400 pM, 0.5 M NaCl, PBS, pH 8.3) were
injected to the flow chamber at a flow rate of 600 µl/min. For colocalization experiments,
imager strands were exchanged by washing the samples with 1 mL PBS followed by injection
of new imager strands.

Microscopy setup and data acquisition
DNA-PAINT based super-resolution microscopy was performed at the N-STORM super-
resolution microscopy system (Nikon, Japan), equipped with an oil immersion objective
(Apo, 100x, NA 1.49) and an EMCCD camera (DU-897U-CS0-#BV, Andor Technology,
Ireland) as described previously (39). Fluorophore-conjugated oligonucleotides were excited
with a collimated 647 nm laser beam at an intensity of 2 kW/cm^2 (measured at the objective)
at highly inclined and laminated optical sheet (HILO) mode. 40.000 and 20.000 consecutive
frames were acquired for imager strands P1 (5´-TAGATGTAT-3´-ATTO655) and P5 (5´-
CATACATTGA-3´-ATTO655), respectively. Image acquisition was performed at 20 Hz (P1)
and 10 Hz (P5) in active frame transfer mode with an EMCCD gain of 200, a pre amp gain of
1 and at an effective pixel size of 158 nm. Software tools NIS Elements (Nikon, Japan),
LCControl (Agilent, USA), and Micro-Manager 1.4.22 (40) were used for control of the
optical setup and image acquisition. FAM134B and FAM134C were imaged sequentially
using a microfluidic assisted exchange of imager strands. The spatial resolution of our super-

resolution images was determined to 22 ± 2 nm (mean ± s.d., (41), allowing the detection of

FAM134 nanoscale cluster.

Image processing

Localization of single molecules and reconstruction of super-resolution images were

performed with modular software package Picasso (42). Integrated Gaussian maximum

likelihood estimation was used for detection of single molecules using the following

parameter: Min. net. gradient of 15000, a baseline of 205, a sensitivity of 4.78, a quantum

efficiency of 0.95. Sample drift correction and image alignment were performed via

redundant cross-correlation algorithm using fiducial marker. For FAM134B (WT and

S149,151,153A) and FAM134C (WT and S258A), localizations were subsequently filtered

based on ATTO-655 single-molecule footprint (PSF symmetry

$0.6 < FWHM(x)/FWHM(y) < 1.4$). Signals from the same origin detected in consecutive

images were linked within a radius of five times the nearest-neighbor based analysis (NeNa)

(43) localization precision and a maximum dark time of 8 consecutive frames. Remaining

signals from fiducial marker were removed by excluding traces from the same origin with a

length larger than 20 consecutive frames and a min. net gradient < 100.000. To account for

the high expression level in FAM134C WT expressing cells, nanoscale cluster were first

segmented from ER-located background signal via Density-based spatial clustering and

application with noise algorithm (44) analysis using a radius of 16 nm and a minimum density

of 30 localizations followed by data filtering as described previously.

Cluster analysis

DBSCAN algorithm was used to identify protein nanocluster in DNA-PAINT super-

resolution images. A radius of 32 nm (FAM134B WT and S149,151,153A and FAM134C

S258A) and a minimum density of 10 localizations were used as parameter. For FAM134C

WT, a radius of 16 nm and a minimum density of 5 localizations were used for DBSCAN

analysis. For the determination of nanoscale cluster density in U2OS cells, protein clusters

were normalized with respect to the cell area. Therefore, cell contours were drawn in super-

resolution images using ImageJ and the cell area was determined.

Statistical analysis

FAM134B and FAM134C nanocluster diameter were tested for normal distribution via

Shapiro-Wilk normality test and statistical significance via non-parametric Mann-Whitney U-

test. The relative frequency distribution of nanocluster diameter was fitted with a Log-normal

distribution and the mode of distribution was calculated as $e^{(\mu-\sigma^2)}$. All statistical analysis

were performed in OriginPro (Origin Lab) or GraphPad Prism for quantified blots.

Material and methods: supplementary information

Sample preparation for STED imaging

Sample preparation for stimulated emission depletion (STED) microscopy was as described

for exchange DNA-PAINT in material and methods with the exception of the antibody used.

ALFA-FAM134s were stained directly using a nanobody against ALFA tag coupled to a

Abberior Star 635P fluorophore (FluoTag®-X2 anti-ALFA - AbberiorStar635P, NanoTag

Biotechnologies GmbH) diluted 1: 500 in permeabilization/blocking buffer (10% FCS

(Gibco), 0.1% saponin (Sigma-Aldrich)) and incubated for 60 min at room temperature.

Sample preparation for DNA-PAINT

Sample preparation for DNA-PAINT was the same as described for exchange DNA-PAINT

in material and methods with the exception of the antibody used. ALFA-FAM134s were

targeted directly using an anti-ALFA nanobody carrying a 5xR1 docking strand (5-

TCCTCCTCCTCCTCCTCCT-3´) (Massive-TAG-Q Anti-ALFA Custom, Massive Photonics,

GmbH) which was diluted in permeabilization/blocking buffer (10% FCS (Gibco), 0.1%

saponin (Sigma-Aldrich)) at a dilution of 1:500 and incubated for 60 min at room

temperature.

Microscopy setup and data acquisition

DNA-PAINT based super-resolution microscopy was performed using an N-STORM super-

resolution microscopy system as described in material and methods. 25,000 consecutives

frames were acquired for the imager strand R1 (5´-AGGAGGA-3´-ATTO655) at 10 Hz using

settings as described in material and methods.

Image processing

Localization of single molecules and reconstruction of super-resolution images were

performed with modular software package Picasso as described in material and methods with

no additional filtering for FAM134C.

Cluster analysis

DBSCAN algorithm was used to identify protein nanocluster in DNA-PAINT super-resolution images using a radius of 32 nm and a minimum density of 5 localizations (ALFA-FAM134B) or 10 localizations (ALFA-FAM134C). Only clusters that contain more than 20 localizations were used for further analysis. All cluster with a mean frame (mean number of frames in which localizations occurred) in the first or last 30% of the total frame number and a standard deviation $20\% < \sigma < 35\%$ were excluded from the analysis.

Isothermal titration calorimetry

For ITC experiments, LC3B was obtained by Ub-fusion technology based on protocols described previously (45). Synthetic peptides were purchased from the Tufts University Core Facility (0.1 mmol scale, HPLC-purified). Before experiments, proteins and peptides were equilibrated in buffer containing 25 mM Hepes pH 7.5, 100 mM NaCl. All titration experiments were performed in triplicates at 25°C using a VP-ITC microcalorimeter (Malvern Instruments Ltd., UK). ITC data were analyzed using NITPIC for baseline correction and SEDPHAT for fitting. The peptides at a concentration of 250 μM were titrated into 20 μM LC3B in 25 steps. Dilution heat from titrating peptide into buffer or buffer into protein was subtracted. Concentrations of proteins and peptides were calculated from the UV absorption at 280 nm using a Nanodrop spectrophotometer (Thermo Fisher Scientific, DE; USA).

Data availability

The mass spectrometry proteomics data have been deposited to the proteomeXchange Consortium via the PRIDE (46–48) partner repository with the dataset identifier PXD043003.

Acknowledgements

We thank all members of the Quantitative Proteomics Unit at IBC2 (Goethe University, Frankfurt), in particular, Rajeshwari Ratore and Marina Hoffmann for support and expertise in Proteomics methodology and data analysis, Martin Adrian-Allgood for technical support, Kristina Wagner for producing LC columns, and David Krause for help in (bio)informatics. The authors thank Stefan Knapp and Andreas Krämer for providing tools and reagents including purified CK2 and Ivan Dikic for critical comments on the text. We thank Julia Pomirska for support in the purification of the RHD domain and Sergio Alejandro Poveda Cuevas for help with the RHD domain scheme. We thank To Dinh Nguyen Ngu and Ivana Peluso for technical support.

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation) – grant numbers 259130777 (SFB1177); 515275293 (FuGG; cell sorter);
403765277 (FuGG; mass spectrometer); and 512574446 (FuGG; spinning disc confocal), the
Dr. Rolf M. Schwiete Stiftung (13/2017), the EU/EFPIA/OICR/McGill/KTH/Diamond
Innovative Medicines Initiative 2 Joint Undertaking (EUbOPEN grant n° 875510), the
Telethon Foundation (grant TMPGMFU22TT), and AIRC (grant MFAG 2020 ID 24856).
Images were created in part using BioRender.

Discloser statement

The authors declare no conflict of interest.

Authors contribution

RB: conceptualization, investigation, formal analysis, visualization, writing – original draft
preparation; HHX: investigation, validation, writing – review and editing; MG: methodology,
investigation, formal analysis, visualization, writing – original draft preparation; PSM:
investigation, formal analysis, data curation, writing – review and editing; LB: investigation;
TG: investigation, validation; SKK: investigation; KH: investigation, formal analysis; SCF:
investigation, validation; VB: investigation; PCP: investigation; AB: methodology,
investigation; KH: methodology; GT: methodology, formal analysis; TJ: investigation, formal
analysis; MM: resources; AG: resources; VD: methodology, supervision; PG: validation,
resources, supervision, writing – review and editing; MH: methodology, supervision, writing
– review and editing, funding acquisition; AS: conceptualization, methodology, formal
analysis, validation, data curation, visualization, writing – original draft preparation,
supervision, project administration, funding acquisition.

References

Schwarz DS, Blower MD. The endoplasmic reticulum: Structure, function and response to
cellular signaling. Vol. 73, Cellular and Molecular Life Sciences. Birkhauser Verlag AG;
2016. p. 79–94.

Chino H, Mizushima N. ER-Phagy: Quality Control and Turnover of Endoplasmic Reticulum.
Vol. 30, Trends in Cell Biology. Elsevier Ltd; 2020. p. 384–98.

Stolz A, Grumati P. The various shades of ER-phagy. Vol. 286, FEBS Journal. Blackwell
Publishing Ltd; 2019. p. 4642–9.

Hübner CA, Dikic I. ER-phagy and human diseases. Vol. 27, Cell Death and Differentiation.

Springer Nature; 2020. p. 833–42.

Khaminets A, Heinrich T, Mari M, Grumati P, Huebner AK, Akutsu M, et al. Regulation of

endoplasmic reticulum turnover by selective autophagy. Nature. 2015 Jun 18;522(7556):354–

8.

Fumagalli F, Noack J, Bergmann TJ, Presmanes EC, Pisoni GB, Fasana E, et al. Translocon

component Sec62 acts in endoplasmic reticulum turnover during stress recovery. Nat Cell

Biol. 2016 Oct 27;18(11):1173–84.

Zhang H, Grumati P, Morozzi G, Hö Lper S, Mari M, Harwardt MLI, et al. Full length RTN3

regulates turnover of tubular endoplasmic reticulum via selective autophagy.

Smith MD, Harley ME, Kemp AJ, Wills J, Lee M, Arends M, et al. CCPG1 Is a Non-

canonical Autophagy Cargo Receptor Essential for ER-Phagy and Pancreatic ER Proteostasis.

Dev Cell. 2018 Jan 22;44(2):217-232.e11.

Chen Q, Xiao Y, Chai P, Zheng P, Teng J, Chen J. ATL3 Is a Tubular ER-Phagy Receptor for

GABARAP-Mediated Selective Autophagy. Current Biology. 2019 Mar 4;29(5):846-855.e6.

Chino H, Hatta T, Natsume T, Mizushima N. Intrinsically Disordered Protein TEX264

Mediates ER-phagy. Mol Cell. 2019 Jun 6;74(5):909-921.e6.

Reggio A, Buonomo V, Berkane R, Bhaskara RM, Tellechea M, Peluso I, et al. Role of

FAM134 paralogues in endoplasmic reticulum remodeling, ER phagy, and Collagen quality

control. EMBO Rep. 2021 Sep 6;22(9).

Bhaskara RM, Grumati P, Garcia-Pardo J, Kalayil S, Covarrubias-Pinto A, Chen W, et al.

Curvature induction and membrane remodeling by FAM134B reticulon homology domain

assist selective ER-phagy. Nat Commun. 2019 Dec 1;10(1).

Alao JP, Legon L, Rallis C. Crosstalk between the mtor and dna damage response pathways

in fission yeast. Vol. 10, Cells. MDPI; 2021. p. 1–11.

Maréchal A, Zou L. DNA damage sensing by the ATM and ATR kinases. Cold Spring Harb

Perspect Biol. 2013 Sep;5(9).

So KS, Rho JK, Choi YJ, Kim SY, Choi CM, Chun YJ, et al. AKT/mTOR Down-regulation

by CX-4945, a CK2 Inhibitor, Promotes Apoptosis in Chemorefractory Non-small Cell Lung

Cancer Cells. Anticancer Res [Internet]. 2015 [cited 2023 Jun 5];35(3):1537–42. Available

from: https://ar.iiarjournals.org/content/anticanres/35/3/1537.full.pdf

Jiang X, Wang X, Ding X, Du M, Li B, Weng X, et al. FAM 134B oligomerization drives

endoplasmic reticulum membrane scission for ER phagy . EMBO J. 2020 Mar 2;39(5).

979. Nassal D, Gratz D, Hund TJ. Challenges and opportunities for therapeutic targeting of calmodulin kinase II in heart. Vol. 11, Frontiers in Pharmacology. Frontiers Media S.A.; 2020.

982. Gnad F, Gunawardena J, Mann M. PHOSIDA 2011: The posttranslational modification database. Nucleic Acids Res. 2011 Jan;39(SUPPL. 1).

984. Mochida K, Yamasaki A, Matoba K, Kirisako H, Noda NN, Nakatogawa H. Super-assembly of ER-phagy receptor Atg40 induces local ER remodeling at contacts with forming autophagosomal membranes. Nat Commun. 2020 Dec 1;11(1).

987. Bhaskara RM, Grumati P, Garcia-Pardo J, Kalayil S, Covarrubias-Pinto A, Chen W, et al. Curvature induction and membrane remodeling by FAM134B reticulon homology domain assist selective ER-phagy. Nat Commun. 2019 Dec 1;10(1).

990. Croucher DR, Iconomou M, Hastings JF, Kennedy SP, Han JZR, Shearer RF, et al. Bimolecular complementation affinity purification (BiCAP) reveals dimer-specific protein interactions for ERBB2 dimers [Internet]. Available from: www.SCIENCESIGNALING.org

993. Sauer M, Heilemann M. Single-Molecule Localization Microscopy in Eukaryotes. Vol. 117, Chemical Reviews. American Chemical Society; 2017. p. 7478–509.

995. Jungmann R, Steinhauer C, Scheible M, Kuzyk A, Tinnefeld P, Simmel FC. Single-molecule kinetics and super-resolution microscopy by fluorescence imaging of transient binding on DNA origami. Nano Lett. 2010 Nov 10;10(11):4756–61.

998. Götzke H, Kilisch M, Martínez-Carranza M, Sograte-Idrissi S, Rajavel A, Schlichthaerle T, et al. The ALFA-tag is a highly versatile tool for nanobody-based bioscience applications. Nat Commun. 2019 Dec 1;10(1).

1001. Strauss S, Jungmann R. Up to 100-fold speed-up and multiplexing in optimized DNA-PAINT. Nat Methods. 2020 Aug 1;17(8):789–91.

1003. Roffey SE, Litchfield DW. CK2 regulation: Perspectives in 2021. Vol. 9, Biomedicines. MDPI; 2021.

1005. Borgo C, D'Amore C, Sarno S, Salvi M, Ruzzene M. Protein kinase CK2: a potential therapeutic target for diverse human diseases. Vol. 6, Signal Transduction and Targeted Therapy. Springer Nature; 2021.

1008. Sanz-Clemente A, Gray JA, Ogilvie KA, Nicoll RA, Roche KW. Activated CaMKII Couples GluN2B and Casein Kinase 2 to Control Synaptic NMDA Receptors. Cell Rep. 2013;3(3):607–14.

11. Di Lorenzo G, Iavarone F, Maddaluno M, Belén Plata-Gómez A, Aureli S, Paz Quezada Meza C, et al. Phosphorylation of FAM134C by CK2 controls starvation-induced ER-phagy [Internet]. Vol. 8, Sci. Adv. 2022. Available from: https://www.science.org

14. Chino H, Yamasaki A, Ode KL, Ueda HR, N 4 Noda N, Mizushima N. Phosphorylation by casein kinase 2 ensures ER-phagy receptor TEX264 binding to ATG8 proteins 2 3. 4259. Available from: https://doi.org/10.1101/2022.02.11.480038

17. Mochida K, Yamasaki A, Matoba K, Kirisako H, Noda NN, Nakatogawa H. Super-assembly of ER-phagy receptor Atg40 induces local ER remodeling at contacts with forming autophagosomal membranes. Nat Commun. 2020 Dec 1;11(1).

20. González A, Covarrubias-Pinto A, Bhaskara RM, Glogger M, Kuncha SK, Xavier A, et al. Ubiquitination regulates ER-phagy and remodelling of endoplasmic reticulum. Nature [Internet]. 2023 Jun 8;618(7964):394–401. Available from: https://www.nature.com/articles/s41586-023-06089-2

23. Götzke H, Kilisch M, Martínez-Carranza M, Sograte-Idrissi S, Rajavel A, Schlichthaerle T, et al. The ALFA-tag is a highly versatile tool for nanobody-based bioscience applications. Nat Commun. 2019 Dec 1;10(1).

27. Gordon WR, Vardar-Ulu D, L'Heureux S, Ashworth T, Malecki MJ, Sanchez-Irizarry C, et al. Effects of S1 cleavage on the structure, surface export, and signaling activity of human Notch1 and Notch2. PLoS One. 2009 Aug 24;4(8).

35. González A, Covarrubias-Pinto A, Bhaskara RM, Glogger M, Kuncha SK, Xavier A, et al. Ubiquitination regulates ER-phagy and remodelling of endoplasmic reticulum. Nature [Internet]. 2023 May 24; Available from: https://www.nature.com/articles/s41586-023-06089-2

36. Tyanova S, Temu T, Cox J. The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. Nat Protoc. 2016 Dec 1;11(12):2301–19.

36. Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen J V., Mann M. Andromeda: A peptide search engine integrated into the MaxQuant environment. J Proteome Res. 2011 Apr 1;10(4):1794–805.

38. Elias JE, Gygi SP. Target-decoy search strategy for increased confidence in large-scale protein identifications by mass spectrometry. Nat Methods. 2007 Mar;4(3):207–14.

39. Edelstein AD, Tsuchida MA, Amodaj N, Pinkard H, Vale RD, Stuurman N. Advanced methods of microscope control using µManager software. J Biol Methods. 2014 Nov 7;1(2):e10.

44. Schnitzbauer J, Strauss MT, Schlichthaerle T, Schueder F, Jungmann R. Super-resolution microscopy with DNA-PAINT. Nat Protoc. 2017 Jun 1;12(6):1198–228.

45. Descloux A, Grußmayer KS, Radenovic A. Parameter-free image resolution estimation based on decorrelation analysis. Nat Methods. 2019 Sep 1;16(9):918–24.

46. Malkusch S, Heilemann M. Extracting quantitative information from single-molecule super-resolution imaging data with LAMA - LocAlization Microscopy Analyzer. Sci Rep. 2016 Oct 5;6.

47. Ester M, Kriegel HP, Sander J, Xu X. A Density-Based Algorithm for Discovering Clusters in Large Spatial Databases with Noise [Internet]. 1996. Available from: www.aaai.org

48. el Beheiry M, Dahan M. ViSP: Representing single-particle localizations in three dimensions. Vol. 10, Nature Methods. 2013. p. 689–90.

49. Rozenknop A, Rogov V V., Rogova NY, Löhr F, Güntert P, Dikic I, et al. Characterization of the interaction of GABARAPL-1 with the LIR motif of NBR1. J Mol Biol. 2011 Jul 15;410(3):477–87.

50. Perez-Riverol Y, Xu QW, Wang R, Uszkoreit J, Griss J, Sanchez A, et al. PRIDE inspector toolsuite: Moving toward a universal visualization tool for proteomics data standard formats and quality assessment of proteomexchange datasets. Molecular and Cellular Proteomics. 2016 Jan 1;15(1):305–17.

51. Deutsch EW, Bandeira N, Perez-Riverol Y, Sharma V, Carver JJ, Mendoza L, et al. The ProteomeXchange consortium at 10 years: 2023 update. Nucleic Acids Res. 2023 Jan 6;51(D1):D1539–48.

52. Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S, et al. The PRIDE database resources in 2022: A hub for mass spectrometry-based proteomics evidences. Nucleic Acids Res. 2022 Jan 7;50(D1):D543–52.

Figure legends

Figure 1. Chemical screen to identify kinases involved FAM134B- and FAM134C-driven ER-phagy

(**A**) Schematic representation of the ER-phagy flux assay using U2OS cells constitutively expressing the ssRFP-GFP-KDEL reporter and in addition overexpressing FAM134B (FAM134B) or FAM134C (FAM134C) in DOX-inducible manner. Cells carrying the empty overexpression vector are used as control cells (CTRL). ER-phagy flux is represented by the red/green (RFP/GFP) ratio of the total integrated fluorescence intensities. Higher values

indicate higher ER-phagy flux. (**B**) ER-phay flux analysis under basal (DMSO) and Torin1-

induced conditions in the absence and presence of FAM134B/C overexpression. Differences

in the ER-phagy flux of CTRL and FAM134B/C overexpressing cells serve as the screening

window for (**E**) to identify kinases involved in FAM134B/C-driven ER-phagy upon Torin1

treatment. n=3, error bars=±STDV, [au]: arbitrary unit. (**C**) Representative blots image of

Fam134 proteins in indicated MEFs isolated from *Fam134b* and *Fam134c* KO mice,

expressing ER-phagy reporter ssRFP-GFP-KDEL. (**D**) ER-phagy flux in ssRFP-GFP-KDEL

Fam134 WT and KO MEFs, represented by the RFP/GFP ratio of total integrated intensities

in basal (DMSO) or induced (Torin1) conditions. n=3, error bars=±STDV. [au]= arbitrary

unit. **E**) Heatmap of an ER-phagy flux screen of appx. 100 probe and probe-like kinase

inhibitors and available negative control compounds (1µM final concentration) in the

presence of 250nM Torin1, using U2OS cells described in (**A**). Total integrated fluorescent

intensities (RFP/red and GFP/green) were monitored in the IncuCyte® S3 over a time course

of 48 h. Data points represent the averaged red/green ratio of three images comprising >150

cells and taken from three independent wells.

1093

Figure 2. mTOR inhibition triggers phosphorylation of FAM134B and FAM134C by CK2

(**A-B**) Validation of kinase screening HITs counteracting the effect of Torin1 in FAM134B-

(**A**) or FAM134C- (**B**) overexpressing cells. Left panels: Analytic representation of

FAM134B or FAM134C - induced ER-phagy changes upon treatment with a dilution series of

selected compounds (SGC-CK2-1 (CK2i), VE-822 (ATRi), KU-60019/KU-55933 (ATMi),

CHIR-124 (Chk1i) and MK-2206 2HCL (AKTi)) ranging from 0.01 µM to 4 µM combined

with Torin1 treatment. DMSO was set to represent 100% inhibition of FAM134B- and

FAM134C-induced ER-phagy while Torin1 represents 0%. Averaged RFP/GFP ratio

measured in response to HIT compound treatments was normalized at each time point to the

averaged RFP/GFP ratio of DMSO and Torin1. Right panels: Growth of U2OS cells

overexpressing FAM134B or FAM134C treated with indicated compounds and

concentrations and recorded over a time period of 48h (data from 2 time points are shown in

the bar graphs). n=3, error bars=±STDV, [au]: arbitrary unit.

1107

Figure 3. CK2 activity positively impacts FAM134-driven ER-phagy

(**A**) Schematic representation of the mass spectrometry experimental procedure to identify

FAM134s phosphorylation sites under basal (DMSO) or induced (Torin1) conditions. (**B**)

Schematic representation of FAM134B or FAM134C proteins highlighting the reticulon

homology domain (RHD; white) comprising two transmembrane domains (TM1/2 and

TM3/4; blue) flanked by membrane associated linker regions (green), the conserved LC3

interacting motif (LIR), as well as sites of phosphorylation events. (**C**) Representative blot

image of *in vitro* phosphorylation of full-length GST-FAM134B by purified CK2 and

corresponding Coomassie-stained gel (lower panel). Anti-GST antibody was used to detect

GST-FAM134B (upper panel). (**D**) Schematic three-dimensional representation of

FAM134B-RHD highlighting serine 153 phosphorylated *in vitro* by CK2. (**E,F**) Left panels:

representative blot images validating the stable expression of FAM134 WT or phospho-

mutant proteins under DOX induction in the background of U2OS cells expressing ER-phagy

reporter ssRFP-GFP-KDEL. Right panels: ER-phagy flux in U2OS cells expressing

FAM134s WT or phospho-mutants and ER-phagy reporter ssRFP-GFP-KDEL, represented

by the RFP/GFP ratio of total integrated intensities in basal (DMSO) or induced (Torin1)

conditions. n=3, error bars=±STDV. (**G,H**) Representative blot image (**G**) and the relative bar

plot (**H**) showing Collagen I accumulation in WT MEFs, Fam134b KO or cells reconstituted

with FAM134B WT and phospho-mutant (S149,151,153A). (**I,J**) Representative blot image

(**I**) and the relative bar plot (**J**) showing Collagen I accumulation in WT MEFs, Fam134c KO

or cells reconstituted with FAM134C WT and phospho-mutant (S258A).

Data information: GAPDH was used as a reference for ratio calculation. The statistical

significance was estimated by one-way ANOVA. n=4 biological experiments for (G,H) and

n=3 biological experiments for (I,J), error bars=±STDV, [au]: arbitrary unit.

1132

Figure 4. FAM134B and FAM134C phosphorylation by CK2 is prerequisite for their

ubiquitination and Torin1-dependant ER-phagy induction

(**A**) Exemplary pictures of U2OS cells overexpressing FAM134B or FAM134C and stably

expressing KDEL reporter and exposed to indicated conditions. Pictures were acquired using

the IncuCyte® S3 (10x) and show the overlaid GFP and RFP signal from DMSO, Torin1 and

Torin1 + E1 inhibitor treated cells at time point 24h. (**B**) Quantitative representation of ER-

phagy flux essay performed in (**A**) analyzing changes over time in FAM134B- (left panel) and

FAM134C-induced ER-phagy (right panel) upon treatment with E1 inhibitor dilution series in

combination with 250 nM Torin1. (**C**) Representative blot image of endogenously

immunoprecipitated FAM134 proteins from total cellular extracts collected from U2OS cells

previously treated for 6h with the indicated treatments. Anti-FAM134B or anti-FAM134C

antibodies were used to detect endogenous proteins while anti-Ub (P4D1) was used to detect

ubiquitination. (**D,E**) Representative blot image (**D**) and corresponding quantification (**E**) of

TUBE 2 assay in U2OS cells overexpressing FAM134B (left panel) or FAM134C (right

panel) subjected for 6h to the indicated conditions prior to cell lysis. anti-HA antibody was

used to detect HA-tagged FAM134B and FAM134C and ubiquitinated species. (**F,G**)

Representative blot image (**F**) and relative bar plot (**G**) of TUBE 2 assay in U2OS cells

overexpressing FAM134B (left panel) or FAM134C (right panel) transfected with control

siRNA (siCTRL) or siPOOL targeting CSNK2A1 and CSNK2A3 (siCK2). After 48h siRNA

transfection, cells were DOX-induced overnight and treated for 6h with indicated conditions

prior to cell lysis. anti-HA antibody was used to detect HA-tagged FAM134B and FAM134C

and ubiquitinated species while anti-CK2 antibody was used to detect endogenous CK2

levels. (**H**) Representative western blot of the bimolecular complementation affinity

purification assay (BiCAP) showing ubiquitination status of FAM134B (left panel) and

FAM134C (right panel) dimers. GFP-Trap pulldown was performed to immunoprecipitate

reconstituted GFP representing FAM134B or FAM134C dimers. Anti-FAM134B or anti-

FAM134C antibodies were used to detect endogenous proteins, while anti-Ub (P4D1) was

used to detect ubiquitination.

Data information: for TUBE 2 assays, DMSO condition was used as reference for ratio

calculation. The statistical significance was estimated by one-way ANOVA. n=3 biological

experiments for (D,E) and n=5 biological experiments for (F,G), error bars=±STDV, [au]:

arbitrary unit.

1165

Figure 5. Phosphorylation of FAM134B and FAM134C proteins impacts their clustering

(A) Schematic representation of sample preparation for exchange DNA-PAINT experiment

performed on U2OS overexpressing FAM134B and FAM134C upon 12h DOX induction.

Cells were grown under basal, 250 nM Torin1 or 250 nM Torin1 + 1 µM CK2 inhibitor

conditions. (**B**) Super-resolved DNA-PAINT images showing HA-FAM134B (left panel) and

HA-FAM134C (right panel) labeling in U2OS cells treated as described in Fig. 5A.

Localization of single molecules and reconstruction of clusters are annotated in the region of

interest. Scale bars: 10 and 1 µm. (**C-D**) Quantitative analysis of FAM134B (**C**) and

FAM134C (**D**) nanocluster diameters in indicated conditions (250 nM Torin1 or 250 nM

Torin1 + 1 µM CK2 inhibitor). Quantification was performed via DBSCAN clustering

algorithm 3 and distribution of clusters was assessed in WT FAM134B- and FAM134C-

overexpressing U2OS cells and their respective phospho-mutants (FAM134B S149,151,153A

and FAM134C S258A) (**E-J**). Comparative analysis of the relative protein density for

FAM134B or mutant (**E-G**) and FAM134C or mutant (**H-J**). Frequency of transient binding

events (localization) per nanoscale cluster was analyzed under basal, 250 nM Torin1 or 250

nM Torin1 + 1 µM CK2 inhibitor conditions.

1182

Figure 6. FAM134B- and FAM134C- dependent ER-phagy induction is regulated by early

interdependent posttranslational modification (PTMs)

Schematic representation of the model of the study: induction of ER stress via mTOR

inhibition triggers phosphorylation of FAM134B and FAM134C, which primes for their

ubiquitination. Both PTM are essential for the oligomerization process of both FAM134s and

control the size and density of the clustering events leading to ER budding and ER-phagy

induction.

1190

1191 **Supplementary figure legends**

1192

Supplementary figure 1. Identification of kinases involved in FAM134B/C and mTOR-

regulated ER-phagy

(**A**) ER-phay flux in ssRFP-GFP-KDEL U2OS cells overexpressing FAM134B or FAM134C

in basal (DMSO), autophagy-induced (40nM Rapamycin) condition in monotreatment or in

combination with CK2i. Total integrated fluorescent intensities (RFP/red and GFP/green)

were monitored in the IncuCyte® S3 over a time course of 48h. Data points represent the

averaged red/green ratio of three images comprising >150 cells and taken from three

independent wells. n=3, error bars=±STDV, [au]: arbitrary unit. (**B**) Heatmap indicating the

cell confluence over time (% of covered area) corresponding to Fig. 1E. (**C**) Schematic

workflow for data transformation to calculate the relative reduction of Torin1-induced ER-

phagy flux. Cpd: compounds. (**D**) Relative reduction of ER-phagy flux caused by HIT

compounds counteracting Torin1-induced FAM134B- and FAM134C-driven ER-phagy,

respectively. HITs were defined as compounds causing >20% relative reduction in ER-phagy

flux over a minimum time period of 12h.

1207

Supplementary figure 2. Validation of chemical kinase screen and identification of CK2 as

kinase regulating FAM134B- and FAM134C-dependant ER-phagy

(**A**) List of selected HITs from the ER-phagy screening performed in Fig. 1E showing

compounds, and their targets, with an effect on FAM134B (left panel) or FAM134C (right

panel) activities. (**B**) Sample pictures of U2OS cells overexpressing FAM134B or FAM134C

and stably expressing KDEL reporter. Pictures were acquired by the IncuCyte® S3 (10x) and

show the overlaid GFP and RFP signal from annotated conditions at time point 48h. (**C**) Impact of selected HITs on ER-phagy flux in U2OS control cells. U2OS cells stably expressing KDEL reporter were subjected to a dilution series of selected compound HITs (SGC-CK2-1 (CK2i), VE-822 (ATRi), KU-60019 (ATMi), CHIR-124 (Chk1i) and MK-2206 2HCL (AKTi)) ranging from 0.01 µM to 4 µM combined with Torin1 treatment. Outcome of ER-phagy flux was recorded over 48h. (**D**) Relative reduction of FAM134B-driven ER-phagy flux upon CaMKII inhibition.

Supplementary figure 3. CK2 mediates FAM134s phosphorylation

(**A**) Multiple sequence alignment of FAM134B and FAM134C showing the conservation level of phosphorylation sites presented in Fig. 3B. (**B**) Predicted kinases phosphorylating identified serines of FAM134B and FAM134C proteins using motif matcher tool PHOSIDA. (**C**) Schematic representation of the workflow of *in vitro* mass spectrometry performed using purified FAM134B RHD domain. (**D**) Tables showing *in vitro* phosphorylation mass spectrometry results. (**E**) Representative western blot image showing KD efficiency of CK2 performed in U2OS cells transfected with control siRNA (siCTRL) or siPOOL targeting CSNK2A1 and CSNK2A3 (siCK2). (**F**) ER-phagy flux in ssRFP-GFP-KDEL U2OS cells overexpressing FAM134B or FAM134C with CK2 present or depleted in the background. ER-phagy flux was assessed either in basal (DMSO) or autophagy-induced (Torin1) conditions. Total integrated fluorescent intensities (RFP/red and GFP/green) were monitored in the IncuCyte® S3 over a time course of 48 h. Data points represent the averaged red/green ratio of three images comprising >150 cells and taken from three independent wells. n=3, error bars=±STDV, [au]: arbitrary unit.

Supplementary figure 4. FAM134B and FAM134C phosphorylation is prerequisite for their ubiquitination and oligomerization in Torin1-dependent manner

(**A**) Schematic representation of Tandem Ubiquitin Binding Entities pulldown assay (TUBE 2) performed in U2OS cells overexpressing FAM134B or FAM134C in DOX-dependent manner which were subjected to different conditions to then purify mono- and poly-ubiquitinated FAM134s proteins. (**B**) Representative western blot image showing enrichment for poly ubiquitinated FAM134 proteins using TUBE 2 assay. U2OS cells overexpressing FAM134B or FAM134C proteins were treated or not with 200 ng/ml BafA1 3 h prior to cell lysis and pulldown experiment. Anti-HA antibody was used to detect HA-tagged FAM134B and FAM134C, as well as ubiquitinated species. (**C**) Representative western blot showing

enrichment for poly ubiquitinated FAM134 proteins using TUBE 2 assay. U2OS cells overexpressing FAM134B WT or S149,151,153A, FAM134C WT or S258A proteins were treated for 6 h with indicated conditions prior to cell lysis. Pulldown of ubiquitinated entities was performed as in Fig. S3A. anti-HA antibody was used to detect HA-tagged FAM134B and FAM134C and ubiquitinated species. **(D)** Representative scheme outlining the principle behind BiCAP assay using the Venus split system coupled with GFP-Trap pulldown assay. **(E)** Confocal fluorescence microscopy imaging of the BiCAP assay validating the *in cellulo* interaction between V1–FAM134B and V2–FAM134B or V1–FAM134C and V2–FAM134C. Fixed cells expressing the indicated plasmids were stained with REEP5 to mark the ER. Scale bar =10 µm. 10 cells were imaged for control purpose.

Supplementary figure 5. Phosphorylation of FAM134B and FAM134C regulates cluster size and density

(A) Cluster diameter distributions (boxplots) for FAM134B (upper panel) and FAM134C (lower panel) under basal (DMSO), autophagy induced (Torin1) and Torin1 + CK2i condition. Nanoscale cluster diameters were tested for normal distribution via Shapiro-Wilk normality test and statistical significance via non-parametric Mann-Whitney U-test. For further details see material and methods. **(B-C)** Comparative analysis of FAM134B (A) and FAM134C (B) nanoscale cluster diameters classified into small (30-90 nm), medium (90-150 nm) and large clusters (> 150 nm). **(D)** Super-resolved STED images showing DOX-induced expression (0.1 µg/ml) of ALFA-FAM134B (left panel) and ALFA-FAM134C (right panel) in U2OS cells. Fixed cells were stained with ALFA-tag nanobody to visualize FAM134B/C expression. Both images complement the results of SMLM, where FAM134B is sparsely populated in the ER and FAM134C is densely populated in the ER. **(E)** Quantitative analysis of DNA-PAINT imaging data reports cluster diameters of ALFA-FAM134B (left panel) and ALFA-FAM134C (right panel) for DMSO, 250 nM Torin1 and 250 nM Torin1 + 1 µM CK2 inhibitor. Cluster analysis was performed with the DBSCAN clustering algorithm in WT FAM134B- and FAM134C-overexpressing U2OS cells **(F-I).** Comparative analysis of nanoscale clusters densities in ALFA-FAM134B **(F, G)** or ALFA-FAM134C **(H, I)**. Fig. S5E-H represent an exact replicate of the experiment conducted in Fig. 5 using ALFA-tag system and show reproducibility in these settings.

Supplementary figure 6. Phosphorylation of FAM134C increases its affinity for LC3B

(A) ITC profiles for the titration of purchased FAM134-LIR (left) or FAM13C-pLIR (right)

peptide into LC3B. The upper panel shows the raw measurements after baseline correction,

the bottom diagram displays the integrated heat per titration step (black circles) with best-fit

curves (line). Each measurement was performed in triplicates, representative results are

shown. Dissociation constant (K_d) values are indicated, errors represent standard deviations.

(B) SDS-PAGE image showing Coomassie-stained gel of purified GFP-FAM134C-LIR

peptide. E1-3 (elution fraction 1-3). **(C)** Phosphorylation level of GFP-FAM134C-LIR in the

presence or absence of purified CK2. Peptide phosphorylation was assessed using MALDI

(Matric-assisted laser desorption/ionization). **(D,E)** Representative western blot image of

GFP-Trap pulldown assay (**D**) and the relative bar plot (**E**) showing binding intensities of

phosphorylated or non-phosphorylated GFP-FAM134C-LIR peptide (bait) to GST-LC3B

(prey). Anti-GFP was used to detect GFP-FAM134C-LIR while anti-GST antibody was used

to detect LC3B. GFP was used as a reference for ratio calculation. Two-sided unpaired

student's *t*-test was used to estimate statistical significance. n=3, error bars=±STDV, [au]:

arbitrary unit.

1296

1297

1298

1299

1300

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6