Challenges of Whole Genome Sequencing based Molecular Identification of Zoonotic

Tuberculosis caused by *Mycobacterium orygis*

Md Rashedul Islam[a], Meenu K. Sharma[a,b], Rupinder KhunKhun[c], Cary Shandro[d], Inna Sekirov[c],

Gregory J. Tyrell[d], and Hafid Soualhine[a,b*]

[a]National Reference Centre for Mycobacteriology, National Microbiology Laboratory, Public

Health Agency of Canada, Winnipeg, Manitoba, Canada

[b]Department of Medical Microbiology and Infectious Diseases, Max Rady College of Medicine,

University of Manitoba, Winnipeg, Manitoba, Canada

[c]BC Center for Disease Control Public Health Laboratory, Vancouver, British Columbia, Canada

[d]Provincial Laboratory for Public Health, Alberta Precision Labs. Edmonton, Alberta, Canada

18 [*]Corresponding author's contact information:

19 E-mail: hafid.soualhine@phac-aspc.gc.ca; Phone: 204.789.6051

20 National Reference Centre for Mycobacteriology, National Microbiology Laboratory, Public

21 Health Agency of Canada, Winnipeg, Manitoba, Canada

ABSTRACT A recently described member of the *Mycobacterium tuberculosis* complex (MTBC) is *Mycobacterium orygis,* which can cause disease in both animals and humans. Although *M. orygis* has been reported from different geographic regions around the world, due to a lack of proper identification techniques, the contribution of this emerging pathogen to the global burden of zoonotic tuberculosis is not fully understood. In the present work, we report single nucleotide polymorphisms (SNPs) analysis using whole genome sequencing (WGS) that can accurately identify *M. orygis* and differentiate it from other members of MTBC species. WGS-based SNPs analysis was performed for 61 isolates from different Provinces in Canada that were identified as *M. orygis*. A total of 56 *M. orygis* sequences from the public databases were also included in the analysis. Several unique SNPs in *gyrB, PPE55, Rv2042c, leuS, mmpL6,* and *mmpS6* genes were used to determine their effectiveness as genetic markers for the identification of *M. orygis*. To the best of our knowledge, five of these SNPs, *viz., gyrB*^{520} (A→G), *gyrB*^{1721} (T→C), *leuS*^{1064} (A→T), *mmpL6*^{486} (T→C), and *mmpS6*^{334} (C→G) are reported for the first time in this study. Our results also revealed several SNPs specific to other species within MTBC. The phylogenetic analysis shows that studied genomes were genetically diverse and clustered with *M. orygis* sequences of human and animal origin reported from different geographic locations. Therefore, the present study provides a new insight into the high confidence identification of *M. orygis* from MTBC species based on WGS data, which can be useful for reference and diagnostic laboratories.

KEYWORDS *Mycobacterium orygis*, *Mycobacterium tuberculosis* complex, Zoonotic tuberculosis, Whole genome sequencing, Single nucleotide polymorphism, Phylogeny

Tuberculosis is a severe and complex infectious disease caused by the *Mycobacterium*

tuberculosis complex (MTBC) and remains a public health concern leading to the death of 1·6

million people per year worldwide, predominantly in low-income and middle income countries

(1). *Mycobacterium orygis,* a species belonging to MTBC, was first described by van Ingen et al.

(2). *M. orygis* is capable of causing infection in both animal and human hosts. This species has

received considerable interest in recent years, and has been reported to be isolated from dairy cattle

and captive monkeys (3), captive wild animals (4), deer (5), rhinoceros (6), black buck (5), and

humans (2, 7, 8). *M. orygis* infection has been recognized as a zoonotic source of human

tuberculosis (9). Moreover, in New Zealand, a presumptive transmission of *M. orygis* from human

to animal was reported, with the original infection being mapped out from contacts with domestic

animals in India (10). *M. orygis* is endemic in Southeast Asian countries including Bangladesh,

India, Nepal, and Pakistan (3, 7, 10). Although tuberculosis incidence in Canada is relatively low

(11), zoonotic tuberculosis is progressively being recognized as a significant menace to the public

health. Hence, zoonotic tuberculosis could pose a significant challenge in controlling this disease

and meeting global tuberculosis elimination goals (12).

Along with *M. orygis*, other phylogenetically related MTBC species are *M. tuberculosis*,

M. bovis and its variant BCG, *M. africanum*, *M. caprae*, *M. pinnipedii*, *M. microti*, *M. canettii*,

and members of animal-adapted clade A1 such as "Dassie" bacillus, Chimpanzee bacillus, *M.*

suricattae, and *M. mungi* (13-15). To improve human and animal health surveillance, it is

important to implement proper identification methods and analysis tools for quickly and accurately

discriminating MTBC species. Probes hybridization-based assays have been used to differentiate

the causative agents of tuberculosis, however, studies have reported a limitation of this method in

differentiating *M. orygis* from *M. africanum* (16). The mutations in genes including *gyrB* and

Rv2042c, and the regions of difference (RDs) deletion have been used to discriminate *M. orygis*

from MTBC species, using PCR-based approaches (2, 3, 17). Since *M. orygis* strains share *gyrB*^{1450}

(G→T) mutation with MTBC members including *M. africanum* and *M. pinnipedii*, they may have

previously been mislabeled as *M. africanum* (3, 17, 18).

Whole genome sequencing (WGS) technologies are now increasingly being used in clinical

and research laboratories to investigate tuberculosis surveillance, outbreak detection, antimicrobial

resistance prediction, characterization and diversity of MTBC species (14, 19). One of the key

challenges for adopting WGS for these applications is data analysis that requires bioinformatics

support and data interpretation (20). Consequently, a number of analytical tools have been

developed to detect pathogenic bacterial strains using WGS data, with single nucleotide

polymorphisms (SNP)-based methods being the common ones used in public health laboratories

(21-23). BioHansel, for example, performs high-resolution genotyping by detecting

phylogenetically informative SNPs in WGS data (24).

In order to improve our ability to accurately identify *M. orygis* and other species within

MTBC, whole genome analysis was performed in this investigation. We report the identification

of 61 new *M. orygis* from Canada by WGS-based SNP analysis, and to validate the results, publicly

available 56 *M. orygis* genomes were added to the analysis. We performed molecular markers

characterization on these genomes that demonstrate a clear differentiation of *M. orygis* from

members of other MTBC species. Furthermore, the newly sequenced *M. orygis* isolates were

phylogenetically analyzed to determine their diversity and global distribution. This study may

improve our understanding of this emerging zoonotic pathogen poorly monitored and address it

burden in Canada and globally.

MATERIALS AND METHODS

Sample collection, and project background. Cultures and clinical samples from across Canada were received at the National Microbiology Laboratory, Public Health Agency of Canada, Winnipeg (NRCM) for *M. tuberculosis* testing between 2009 and 2022. The cultures were grown on mycobacteria growth indicator tube (MGIT) media and Middlebrook 7H10 plates using standard and aerobic growth conditions. At NRCM, the presence of the insertion sequence IS*6110* and region of difference RD9 was confirmed by RT-PCR analysis to detect MTBC and *M. tuberculosis*, respectively. Classical methods such as *gyrB*, RD1, RD4, RD7 and mycobacterial interspersed repetitive unit-variable number of tandem repeat (MIRU-VNTR) were also used for identification and genotyping purpose. These methods can distinguish between mostly isolated MTBC species, *M. tuberculosis, M. bovis* and its variant BCG, *M. caprae* and *M. africanum*, but they are unable to identify some rare species of the complex such as *M. orygis* and *M. pinnipedii*. Since 2018, our group started using routinely WGS technologies and BioHansel (24) program for identification and differentiation between MTBC species. BioHansel performs high-resolution genotyping of MTBC by detecting phylogenetically informative SNPs in WGS data. However, the SNPs in the *gyrB* gene (at positions 675, 756, 1113, 1311, 1410, and 1450) currently incorporated in BioHansel cannot differentiate *M. orygis*, and the pipeline eventually identifies this species, with low confidence, as a member of an animal lineage of the MTBC (probable *M. tuberculosis* var. *orygis*) based on SNP typing (2). We then separated all isolates identified as animal lineage "probable *M. orygis*",all isolates with same MIRU-VNTR profile and *M. africanum* or *M. pinnipedii* from the inventory and investigated in this study. Therefore, a total of 137 isolates were interrogated, from which 61 were identified to be *M. orygis*. Most of the isolates were from British Columbia 'BC' (n = 38), followed by Alberta 'AB' (n = 19), Manitoba 'MB' (n = 2), Nova Scotia

'NS' (n = 1), and Saskatchewan 'SK' (n = 1). The detailed information (*e.g.,* source Province,

specimen type, collection year, gender, and identification) on *M. orygis* isolates from this study is

depicted in Table 1.

Whole genome sequencing (WGS). (i) DNA extraction, sequencing and data

processing. Genomic DNA was extracted from the isolates using the InstaGene[TM] Matrix

comprising 6% (*w/v*) Chelex resin for PCR-ready DNA purification (Bio-Rad #7326030;

California, United States). DNA quality was checked by measuring the *A260/280* ratio using

NanoDrop, and quantified fluorometrically using the Qubit 3.0 (Thermofisher, Massachusetts,

United States). The sequencing library was prepared using the Illumina DNA Prep Kit (Illumina,

California, United States), according to the manufacturer's protocol. Paired-end sequencing was

conducted on an Illumina MiSeq platform (Illumina Inc.) using a 600-cycle MiSeq Reagent Kit v3

(Illumina, California, United States), creating 2×300 bp paired-end reads.

Raw sequencing reads were processed in the IRIDA platform (25) (http://irida.ca) using

Shovill for assembly (26), Prokka for annotation (27), and QUAST for assembly assessment (28).

Sequencing   Report   workflow   v.2.3,   where   FastQC   (29)   and   SMALT

(https://www.sanger.ac.uk/tool/smalt-0/) are the two main tools, was implemented in the Galaxy

platform (30) for all studied isolates by aligning the reads against the reference genome of *M.*

tuberculosis H37Rv (accession # NC_000962.3). Only sequences with at least 60% of mapped

reads and a >28x average coverage were filtered for further analysis. While 54 out of 61 studied

M. orygis sequences had mapped reads of over 80%, the remaining seven sequences were found

to have 60-80% of mapped reads, and only one sequence (2100725) had average coverage below

30x (28.85).

(ii) Genomic comparison of the new sequences with public database. The newly

sequenced genomes were compared with 56 publicly available *M. orygis* genomes reported from

around the world, downloaded from the National Center for Biotechnology Information (NCBI)

GenBank (31), the NCBI Sequence Read Archive (32), and the European Nucleotide Archive (33).

The list of reference *M. orygis* genomes with their accession numbers, collection centres, country

of origin, host species and specimen source, is presented in Table S1A. Aside from *M. orygis*, a

total of 190 previously published selected genomes sequences (complete/draft/SRA) of MTBC

species and lineages were also downloaded from the above repositories and added to the analysis

(Table S1B).

Gene selection and R analysis: Unique SNPs for *M. orygis* and other MTBC

species/lineages. A set of six genes, *viz., gyrB, PPE55, Rv2042c, leuS, mmpL6,* and *mmpS6* were

selected for SNPs analysis, and the first five genes were chosen based on the previous publications

(2, 17, 34). The sequence for *gyrB, PPE55, Rv2042c* and *leuS* genes were obtained from H37Rv

(accession # NC_00962.3), and *mmpL6* and *mmpS6* gene sequences were recovered from Mtb-

specific deletion 1 region 'TbD1' (accession # AJ426486.1). The SNPs positions in the *gyrB* gene

are described according to previous nomenclature (accession # L27512.1, (35), while the positions

within *Rv2042c, PPE55,* and *leuS* genes are relative to *M. tuberculosis* H37Rv reference genome

(accession # NC_000962.3, (36). The numerical positions for the new and known SNPs within

mmpL6 and *mmpS6* genes are provided following TbD1 region (AJ426486.1, (37)). All identified

SNPs within the six genes are numbered according to gene position (5′ to 3′), and the

complementary sequences were used for genes *Rv2042c* and *PPE55* (Table 2).

To extract the above-mentioned six gene sequences from *M. orygis* and MTBC

species/lineage genomes, a high-throughput analysis was performed using our custom

in-house-developed R-scripts in RStudio v.2021.9.2.382 (38), which used Basic Local Alignment

Search Tool (BLAST) (39), with the e-value cutoff option set to 10e-100. The scripts interrogated

the assembled genomes using a reference gene to identify and extract the sequence corresponding

to the SNPs of interest. NCBI BLAST search was also done for all genes, with any unique

sequences investigated. In order to identify *M. orygis* or other MTBC species-specific new and

known SNPs, the extracted gene sequences were aligned using MUSCLE (40), and sequence

variations of the unique SNPs were detected using MEGA-X v.10.0.5 (41) and Geneious Prime

2022 (Geneious Prime 2022.0.2, https://www. geneious.com).

The SNPs-based phylogenetic reconstruction of *M. orygis*. To compare *M. orygis*

sequences from the present study with those available in the public repositories, a maximum

likelihood tree was constructed using the single nucleotide variant phylogenomics (SNVPhyl)

pipeline v.1.2.3 in IRIDA (23). *M. tuberculosis* H37Rv (NC_00962.3) was chosen as a reference

genome with parameters: minimum SNV coverage of 15, SNV abundance ratio of 0.75, and

minimum mapping quality of 30. The SNV density filtering was enabled to remove high-density

SNV regions that could suggest possible recombination. The identified SNVs were used in the

construction of a phylogeny by calculating genetic distance between isolates by a generalized time

reversible (GTR) model with PhyML 3.0 (42).

Data availability. The whole genome sequences of *M. orygis* isolates from this project

have been deposited in GenBank and are available through the SRA under BioProject no.

PRJNA934340 with the submission ID SUB12537725. Accession numbers for all publicly

available genomes used in this study are listed in Table S1.

RESULTS

Data collection and genome analysis. In the present study, a total of 61 *M. orygis* isolates were identified from cultures submitted by five different Provinces in Canada over a period of 14 years (2009-2022), with more than 90% from the Provinces of British Columbia and Alberta. The first isolates of this zoonotic pathogen were misidentified since 2009 as *M. africanum*. The cultures were recovered from pulmonary (65%) and extra-pulmonary samples. Forty-five *M. orygis* cases were female patients (74%), 14 were male patients (30%) and two cases were with unknown gender patient. The PCR, MIRU-VNTR or BioHansel and Mykrobe Predictor v.0.7.0 (43) analyses identified the isolates as *M. africanum* or *M. pinnipedii* or *M. africanum* with MTBC animal lineages (Table 1). We then performed WGS for all 61 isolates and determined their identification as *M. orygis* using SNPs analysis. The sequencing data were augmented with a total of 56 publicly available *M. orygis* genome sequences to validate the identification and their discrimination from other MTBC species. Of these 56 sequences, 35 were reported from India, followed by eight from The United States, five from Norway, four from Switzerland, two from The United Kingdom, one from The Netherlands, and one from Canada (Table S1A).

Whole genome-based SNPs analysis. We performed the WGS-based SNPs analysis for identification and genetic differentiation of *M. orygis* from other MTBC members. Several new and known SNPs in *gyrB* (2028 bp)*, PPE55* (2405 bp)*, Rv2042c* (798 bp)*, leuS* (2910 bp)*, mmpL6* (2904 bp)*,* and *mmpS6* (447 bp) genes were evaluated to ascertain their effectiveness as molecular markers for *M. orygis*. By gene sequence analysis of multiple sequence alignment from 117 *M. orygis* genomes (61 studied and 56 public repositories) and 190 sequences of MTBC species and lineages, we identified 10 unique SNPs across six selected genes that can identify and discriminate

M. orygis from all members of MTBC. These *M. orygis* specific SNPs along with their

corresponding genomic positions, mutation type, locus, and function are listed in Table 2.

A multiple alignment of the *gyrB* gene showed the presence of two novel nonsynonymous

mutations at positions 520 (A→G) and 1721 (T→C), along with a known synonymous mutation

at position 1113 (G→A) (Table 2, Table 3). Within *PPE55* gene, our study identified two *M. orygis*

specific SNPs at positions 2162 (C→T) and 2163 (T→G). The SNPs analysis also identified a

unique SNP in *Rv2042c* gene at position 113 (T→G). In addition, we found two *M. orygis* specific

SNPs within *leuS* gene; one of which was a novel SNP (*leuS*^{1064} A→T) and associated with

synonymous mutation, and another one (*leuS*^{1251} G→T) was previously reported. The sequence of

the TbD1 region (AJ426486.1) from the ancestral *M. tuberculosis* contains genes *mmpS6* and

mmpL6. Two more novel SNPs were detected in this region and were found to be associated with

a synonymous (*mmpL6*^{486} T→C) and a nonsynonymous (*mmpS6*^{334} C→G) mutations (Table 2,

Table 3). The genes *mmpL6* and *mmpS6* encode membrane proteins belonging to a large and a

small family, respectively. The above 10 unique SNPs situated within a set of six genes were

identified in all 117 *M. orygis* strains screened in the present study.

In order to define the discriminatory power of the WGS-based SNPs analysis in identifying

all MTBC species and lineages, a small MTBC dataset consisting of 190 genome sequences was

also included in this investigation, and results of representative sequence from each species and

lineage are shown in Table 3. Apart from *M. orygis* specific SNPs, we identified some new and

previously reported SNPs that are unique for all MTBC members and lineages. These include 11

SNPs in *gyrB*, 10 SNPs in *leuS*, eight SNPs in *PPE55*, six SNPs in *mmpL6*, three SNPs in *Rv2042c*,

and two SNPs in *mmpS6* genes.

In our study, we detected a series of SNPs specific for *M. canettii* in all six selected genes,

including *gyrB* (456G→A, 735C→T, 1011C→T, 1458C→T and 2097T→C), *PPE55*

(1445G→A), *Rv2042c* (134A→G), *mmpL6* (1879G→C), and *leuS* (321C→A and 2199T→C).

Species-specific polymorphisms were also found for *M. caprae* (*gyrB*^{1311} T→G and *leuS*^{1262}

A→C), *M. microti* (*gyrB*^{675} C→T) and *M. mungi* (*leuS*^{2343} G→A). Interestingly, one SNP

identified in *leuS* gene (564A→G) of *M. microti* was overlapped in *M. pinnipedii* strains. The gene

sequence analysis identified a SNP (*mmpL6*^{306} C→T) that can differentiate *M. africanum* lineage

6 from other MTBC members and lineage. While the SNPs in *gyrB* gene at position 1410 (C→T)

and in *leuS* at position 1632 (G→A) were found to discriminate *M. bovis* and *M. bovis* BCG from

all others, the SNP *PPE55*^{1701} A→G could even separate *M. bovis* from BCG. Furthermore, *M.*

bovis and BCG shared SNPs *gyrB*^{756} G→A and *PPE55*^{556} C→A with *M. caprae* strains (Table 3).

In addition to the above, we detected SNP markers in the selected six genes that differentiated the

strains belonging to different lineages from other MTBC members examined in this study.

Single nucleotide polymorphisms-based phylogenetic inference of *M. orygis*. A whole

genome SNP-based phylogenetic tree was built to determine the genetic similarity of the newly

sequenced *M. orygis* isolates in Canada with *M. orygis* sequences from the public repositories (Fig.

1). The SNV distance matrix generated from the whole genome alignments lists the pairwise SNV

distances between every sequence (Table S2). The genome sequences from this study clustered

with *M. orygis* sequences reported from all five geographic locations.

M. orygis sequences from different samples of the same patient clustered together in the

SNVPhyl tree, as expected. For example, BC isolate 1801225 clustered with 1801535, 1900059

with 1900097, 2001127 with 2001229, 2001439 with 2001480, and 2200196 with 2200263, and

appeared to have a SNVs distance of 0-5. The latter two isolates also closely clustered (29-52

SNVs distance) with two *M. orygis* sequences from Norway (ERR5336158) and The United States

(SRR5642712), and one previously reported *M. orygis* sequence 51145 (SRR16643349) from

Quebec Canada. Similarly, 11 *M. orygis* sequences from the present study formed a large cluster

with three sequences (ERR2659153, ERR2659154, ERR2659156) from Switzerland, one

sequence (ERR5336157) from Norway, and another sequence (ERR015582) reported from The

United Kingdom, with two sequences (1900228 from AB and 1800965 from BC) showing a

distance of 92-96 SNVs with the *M. orygis* sequences from Switzerland (Fig. 1, Table S2).

The studied isolates 1901282 and 2100725 from AB clustered with 11 *M. orygis* sequences

of animal origin from India and one sequence of human origin from The United States with a

pairwise SNV distance of 34-88. Interestingly, the isolate 2100725 closely clustered with seven of

these animal originated sequences (SRR10251185, SRR10251189-92, SRR10251197, and

SRR10251200) by 34 SNV distances, highlighting the zoonotic potential of *M. orygis*.

DISCUSSION

The present study is the first to report a greater number of *M. orygis* from a single geographic area.

M. orygis as a causal agent of tuberculosis in both animal and human hosts was first described in

2012 (2), and thereafter this organism has been reported by several groups (3-5, 7, 8). The MTBC

species *M. bovis* has long been believed to be the only agent that causes zoonotic tuberculosis,

however, recovering a larger number of *M. orygis* from both animals and humans in recent years

from different areas of the world highlights the need for considering this bacterium as a zoonotic

pathogen (44).

Since tuberculosis cases caused by *M. orygis* are often identified as MTBC or misidentified

and published as *M. africanum* or *M. bovis*, the actual number of infections associated with this

bacterium may have been underreported (3, 16, 45). Our investigation revealed that *M. orygis* was

misidentified as *M. africanum* since 2009 in human TB patients. Rahim et al. (18) initially reported

M. africanum from four dairy cows that were later identified to be *M. orygis* by refined analysis

(3). A part of the confusion towards this misidentification is that *M. orygis* shares the $gyrB^{1450}$

(G→T) mutations with *M. africanum*, *M. bovis, M. microti, M. caprae, M. mungi*, and *M.*

pinnipedii ((17), Table 3). Moreover, MTBC is genetically highly clonal, and thus without proper

identification tools and analysis approaches the species differentiation could be challenging.

Since currently available bioinformatics tools are struggling to differentiate animal

lineages, in the present investigation we used WGS-based SNPs analysis targeting *gyrB, PPE55,*

Rv2042c, leuS, *mmpL6,* and *mmpS6* genes that can accurately identify *M. orygis* and

unambiguously differentiate it from all members of MTBC species. The *gyrB* gene encodes for the

β-subunit of the DNA gyrase and has been used as a molecular marker for several bacterial genera

including members of MTBC. The discriminatory power of polymorphisms in *gyrB* gene in

identifying *M. orygis* has been evaluated in previous studies (3, 17). While these authors described

only one unique SNP ($gyrB^{1113}$ C→T), we detected two more novel and useful genetic markers

within *gyrB* gene ($gyrB^{520}$ A→G, and $gyrB^{1721}$ T→C) in the screened *M. orygis* genomes. The

PPE55 (Rv3347c) gene is specific to MTBC species and play a major role in host-pathogen

interactions. Huard et al. (17) earlier reported two *M. orygis* specific SNPs in *PPE55* gene

($PPE55^{2162}$ C→T, and $PPE55^{2163}$ T→G), that was also revealed in our study. The results suggest

that these SNPs could be used as genetic markers for the identification of *M. orygis*. We also

evaluated *Rv2042c* gene for possible unique SNPs to be used as a molecular marker to identify *M.*

orygis. In accordance with van Ingen et al. (2), our SNPs analysis identified a nonsynonymous

mutation in the 38th codon of *Rv2042c* gene at position 113 (T→G). In mycobacterial species,

leuS gene encodes for L-leucyl-tRNA synthetase, which is involved in translation. In this study, in

addition to the SNP *leuS*[1251] (G→T) described by Napier et al. (34), we showed that another novel

SNP *leuS*[1064] (A→T) associated with nonsynonymous mutation is also present in *M. orygis* strains

(Table 2, Table 3). Mtb-specific deletion 1 region 'TbD1' region comprises the *mmpL6* and

mmpS6 genes, which encode for mycobacterial membrane protein families. In modern *M.*

tuberculosis strains, the *mmpS6* gene is fully deleted and the *mmpL6* gene is trimmed (46). Our

result shows two SNP *mmpL6*[486] T→C, and *mmpS6*[334] C→G, as two novel and distinct genetic

markers for the identification of *M. orygis*.

We included previously described *M. orygis* specific polymorphisms in the WGS-based

SNPs analysis, and they were reconfirmed in a larger number of *M. orygis* genomes tested in this

work. Thus, the results suggest WGS-based SNPs analysis is a useful tool to rapidly identify *M.*

orygis, and to clearly differentiate this emerging pathogen from all MTBC species and lineages.

To the best of our knowledge, the unique SNPs *gyrB*[520] (A→G), *gyrB*[1721] (T→C), *leuS*[1064] (A→T),

mmpL6[486] (T→C), and *mmpS6*[334] (C→G) are the first to be reported in the present study.

WGS-based SNPs analysis has been employed for the identification of MTBC species (9),

and this approach has also become pertinent to MTBC genotyping (37, 47). In this study, we

further identified the species-specific polymorphisms for members of MTBC species (excluding

M. orygis here, as discussed above) and lineages, and results from *gyrB* gene confirm the

discrimination of several MTBC members including *M. tuberculosis*, *M. bovis, M. canettii*, *M.*

microti, and *M. caprae*, (Table 3). These results are in agreement with Huard et al. (17), who

reported the identification of the above species (along with *M. orygis*) using a PCR-based SNPs

analysis. Our study supports that many genomic characteristics could be shared between *M. bovis*

and *M. caprae* strains, for example, the *gyrB* mutation at position 756, and *PPE55* mutation at

position 556 (17).

Furthermore, we detected unique polymorphisms for lineage 1 in *gyrB*^{1116} gene (G→C);

for lineage 2 in *Rv2042c*^{597} (C→G) gene; for lineage 3 in *PPE55* gene at positions 1173 (G→C),

1177-1178 (A→G), 1179 (C→T) and 1182 (C→G); for lineage 5 in *PPE55*^{923} (G→T), *mmpS6*^{360}

(C→T) and *leuS*^{2736} (T→C) genes; for lineage 7 in *mmpL6*^{1780} (G→C) and *mmpS6*^{339} (C→T)

genes; and for lineage 9 in *leuS*^{1363} (C→T) gene. The reason for not being able to extract some

SNPs in *mmpL6* and *mmpS6* genes for *M. tuberculosis* H37Rv and lineages 1-4 in Table 3 is that

these genes are partly or fully deleted from the modern *M. tuberculosis*, as discussed above. Our

results indicate that WGS-based SNPs analysis could be successfully used to distinguish all

members of MTBC species and lineages.

The use of whole genome SNPs-based phylogenetic tree allowed us to inspect the genetic

relation of *M. orygis* recovered from Canada and those reported from other geographic regions.

The *M. orygis* isolates of human origin from this study were found to be distributed across the

phylogeny (Fig. 1). Phylogenetic analysis of *M. orygis* genomes from the same patient on the same

episode (two isolates from each of five patients) showed a difference of 0-5 SNV.

We sought to determine whether *M. orygis* isolated from human clinical samples in this

study showed a close phylogenetic relation with animal strains. The result shows that the isolate

2100725 from an AB female clustered (34 SNVs apart) with several *M. orygis* sequences of animal

origin reported from India (Fig. 1, Table S2). This result may be indicative of the adaptation of

animal origin *M. orygis* strains to a human host. The zoonotic potentials of *M. orygis* has been

discussed in a previous work (44).

Furthermore, two isolates from BC were related (5-39 SNVs apart) to *M. orygis* sequences

of human origin reported from Norway (ERR5336158, (8) and The United States (SRR5642712,

(7), and segregated by 49-52 SNVs from previously published Canadian strain 51145 (48). Thus,

the newly sequenced *M. orygis* from Canada phylogenetically clustered with *M. orygis* sequences

reported from different geographic locations, placing it within the global context.

In conclusion, the WGS analysis in the present study evaluated 10 novel and known unique

SNPs within a set of six genes that could be used as molecular genetic markers to accurately

identify *M. orygis* and unambiguously discriminate it from all members of MTBC species and

lineages. As WGS technologies are increasingly being used by the healthcare systems, our

approach will be helpful to diagnosis and surveillance of *M. orygis* associated tuberculosis, and

optimizing the clinical management of this disease. The analysis of *M. orygis* sequences will

improve our understanding of molecular characteristics and phylogenetic diversity of this

emerging pathogen and its implication as a zoonosis. The ever-increasing evidence of *M. orygis*

linked endemicity, and the identification of a greater number of *M. orygis* from animals and

humans around the world highlight the urgency for a multi-sectoral collaboration linking the

clinical and veterinary sectors towards One Health approach. The origin, epidemiology and

transmission dynamics of *M. orygis* within Canada is currently under investigation.

ACKNOWLEDGMENTS

The authors would like to thank Darrell Johnstone, Debra Janella and members of the

National Reference Centre for Mycobacteriology (NRCM) and DNA Sequencing Core Services

for their technical assistance with this project. The authors also acknowledge Aaron Petkau of

National Microbiology Laboratory (NML) for helpful suggestions with phylogenetic analysis.

This work is supported under GRDI program funded by Public Health Agency of Canada.

CONFLICTS OF INTEREST

The authors declare that there are no conflicts of interest.

REFERENCES

1. WHO. 2021. Global tuberculosis report 2021. World Health Organization, Geneva.

2. van Ingen J, Rahim Z, Mulder A, Boeree MJ, Simeone R, Brosch R, van Soolingen D. 2012. Characterization of *Mycobacterium orygis* as *M. tuberculosis* complex subspecies. *Emerg Infect Dis* 18:653-5.

3. Rahim Z, Thapa J, Fukushima Y, van der Zanden AGM, Gordon SV, Suzuki Y, Nakajima C. 2017. Tuberculosis Caused by *Mycobacterium orygis* in Dairy Cattle and Captured Monkeys in Bangladesh: a New Scenario of Tuberculosis in South Asia. *Transbound Emerg Dis* 64:1965-1969.

4. Thapa J, Nakajima C, Maharjan B, Poudell A, Suzuki Y. 2015. Molecular characterization of *Mycobacterium orygis* isolates from wild animals of Nepal. *Jpn J Vet Res* 63:151-8.

5. Refaya AK, Ramanujam H, Ramalingam M, Rao GVS, Ravikumar D, Sangamithrai D, Shanmugam S, Palaniyandi K. 2022. Tuberculosis caused by *Mycobacterium orygis* in wild ungulates in Chennai, South India. *Transbound Emerg Dis* 69:e3327-e3333.

6. Thapa J, Paudel S, Sadaula A, Shah Y, Maharjan B, Kaufman GE, McCauley D, Gairhe KP, Tsubota T, Suzuki Y, Nakajima C. 2016. *Mycobacterium orygis*-Associated Tuberculosis in Free-Ranging Rhinoceros, Nepal, 2015. *Emerg Infect Dis* 22:570-2.

7. Marcos LA, Spitzer ED, Mahapatra R, Ma Y, Halse TA, Shea J, Isabelle M, Lapierre P, Escuyer VE. 2017. *Mycobacterium orygis* Lymphadenitis in New York, USA. *Emerg Infect Dis* 23:1749-1751.

8. Eldholm V, Rønning JO, Mengshoel AT, Arnesen T. 2021. Import and transmission of *Mycobacterium orygis* and *Mycobacterium africanum*, Norway. *BMC Infect Dis* 21:562.

9.   Lipworth S, Jajou R, de Neeling A, Bradley P, van der Hoek W, Maphalala G, Bonnet M,

Sanchez-Padilla E, Diel R, Niemann S, Iqbal Z, Smith G, Peto T, Crook D, Walker T, van

Soolingen D. 2019. SNP-IT Tool for Identifying Subspecies and Associated Lineages of

Mycobacterium tuberculosis Complex. *Emerg Infect Dis* 25:482-488.

10.   Dawson KL, Bell A, Kawakami RP, Coley K, Yates G, Collins Desmond M. 2012.

Transmission of *Mycobacterium orygis* (*M. tuberculosis* Complex Species) from a

Tuberculosis Patient to a Dairy Cow in New Zealand. *J Clin Microbiol* 50:3136-3138.

11.   Mounchili A, Perera R, Lee RS, Njoo H, Brooks J. 2022. Chapter 1: Epidemiology of

tuberculosis in Canada. *Can J Respir Crit Care Sleep Med* 6:8-21.

12.   WHO. 2016. Towards TB elimination in low-incidence countries: TB treatment and care

2016.

13.   Boritsch EC, Khanna V, Pawlik A, Honoré N, Navas VH, Ma L, Bouchier C, Seemann T,

Supply P, Stinear TP, Brosch R. 2016. Key experimental evidence of chromosomal DNA

transfer among selected tuberculosis-causing mycobacteria. *Proc Natl Acad Sci U S A*

113:9876-9881.

14.   Brites D, Loiseau C, Menardo F, Borrell S, Boniotti MB, Warren R, Dippenaar A, Parsons

SDC, Beisel C, Behr MA, Fyfe JA, Coscolla M, Gagneux S. 2018. A New Phylogenetic

Framework for the Animal-Adapted *Mycobacterium tuberculosis* Complex. *Front*

Microbiol 9.

15.   Landolt P, Stephan R, Stevens MJA, Scherrer S. 2019. Three-reaction high-resolution

melting assay for rapid differentiation of *Mycobacterium tuberculosis* complex members.

Microbiologyopen 8:e919.

16. Loiseau C, Brites D, Moser I, Coll F, Pourcel C, Robbe-Austerman S, Escuyer V, Musser

Kimberlee A, Peacock Sharon J, Feuerriegel S, Kohl Thomas A, Niemann S, Gagneux S,

Köser Claudio U. 2019. Revised Interpretation of the Hain Lifescience GenoType MTBC

To Differentiate *Mycobacterium canettii* and Members of the *Mycobacterium tuberculosis*

Complex. *Antimicrob Agents Chemother* 63:e00159-19.

17. Huard Richard C, Fabre M, de Haas P, Claudio Oliveira Lazzarini L, van Soolingen D,

Cousins D, Ho John L. 2006. Novel Genetic Polymorphisms That Further Delineate the

Phylogeny of the *Mycobacterium tuberculosis* Complex. *J Bacteriol* 188:4271-4287.

18. Rahim Z, Möllers M, te Koppele-Vije A, de Beer J, Zaman K, Matin MA, Kamal M,

Raquib R, van Soolingen D, Baqi MA, Heilmann FG, van der Zanden AG. 2007.

Characterization of *Mycobacterium africanum* subtype I among cows in a dairy farm in

Bangladesh using spoligotyping. *Southeast Asian J Trop Med Public Health* 38:706-13.

19. Meehan CJ, Goig GA, Kohl TA, Verboven L, Dippenaar A, Ezewudo M, Farhat MR,

Guthrie JL, Laukens K, Miotto P, Ofori-Anyinam B, Dreyer V, Supply P, Suresh A, Utpatel

C, van Soolingen D, Zhou Y, Ashton PM, Brites D, Cabibbe AM, de Jong BC, de Vos M,

Menardo F, Gagneux S, Gao Q, Heupink TH, Liu Q, Loiseau C, Rigouts L, Rodwell TC,

Tagliani E, Walker TM, Warren RM, Zhao Y, Zignol M, Schito M, Gardy J, Cirillo DM,

Niemann S, Comas I, Van Rie A. 2019. Whole genome sequencing of *Mycobacterium*

tuberculosis: current standards and open issues. *Nat Rev Microbiol* 17:533-545.

20. Deurenberg RH, Bathoorn E, Chlebowicz MA, Couto N, Ferdous M, García-Cobos S,

Kooistra-Smid AMD, Raangs EC, Rosema S, Veloo ACM, Zhou K, Friedrich AW, Rossen

JWA. 2017. Application of next generation sequencing in clinical microbiology and

infection prevention. *J Biotechnol* 243:16-24.

21. Nadon C, Van Walle I, Gerner-Smidt P, Campos J, Chinen I, Concepcion-Acevedo J,

Gilpin B, Smith AM, Man Kam K, Perez E, Trees E, Kubota K, Takkinen J, Nielsen EM,

Carleton H. 2017. PulseNet International: Vision for the implementation of whole genome

sequencing (WGS) for global food-borne disease surveillance. *Euro Surveill* 22.

22. Dallman T, Ashton P, Schafer U, Jironkin A, Painset A, Shaaban S, Hartman H, Myers R,

Underwood A, Jenkins C, Grant K. 2018. SnapperDB: a database solution for routine

sequencing analysis of bacterial isolates. *Bioinformatics* 34:3028-3029.

23. Petkau A, Mabon P, Sieffert C, Knox NC, Cabral J, Iskander M, Iskander M, Weedmark

K, Zaheer R, Katz LS, Nadon C, Reimer A, Taboada E, Beiko RG, Hsiao W, Brinkman F,

Graham M, Van Domselaar G. 2017. SNVPhyl: a single nucleotide variant phylogenomics

pipeline for microbial genomic epidemiology. *Microb Genom* 3:e000116.

24. Labbé G, Kruczkiewicz P, Robertson J, Mabon P, Schonfeld J, Kein D, Rankin MA, Gopez

M, Hole D, Son D, Knox N, Laing CR, Bessonov K, Taboada EN, Yoshida C, Ziebell K,

Nichani A, Johnson RP, Van Domselaar G, Nash JHE. 2021. Rapid and accurate SNP

genotyping of clonal bacterial pathogens with BioHansel. *Microb Genom* 7:000651.

25. Matthews TC, Bristow FR, Griffiths EJ, Petkau A, Adam J, Dooley D, Kruczkiewicz P,

Curatcha J, Cabral J, Fornika D, Winsor GL, Courtot M, Bertelli C, Roudgar A, Feijao P,

Mabon P, Enns E, Thiessen J, Keddy A, Isaac-Renton J, Gardy JL, Tang P, João A Carriço

TIC, Chindelevitch L, Chauve C, Graham MR, McArthur AG, Taboada EN, Beiko RG,

Brinkman FSL, Hsiao WWL, Domselaar GV. 2018. The Integrated Rapid Infectious

Disease Analysis (IRIDA) Platform. *bioRxiv* doi:10.1101/381830:381830.

26. Seemann T. 2019. Shovill. https://github.com/tseemann/shovill.

27.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068-

457   2069.

458   28.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for

459   genome assemblies. *Bioinformatics* 29:1072-1075.

460   29.   Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.

461   Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

462   30.   The Galaxy C. 2022. The Galaxy platform for accessible, reproducible and collaborative

463   biomedical analyses: 2022 update. *Nucleic Acids Res* 50:W345-W351.

464   31.   NCBI. 2022a. https://www.ncbi.nlm.nih.gov/genbank/. Accessed December 9, 2022.

465   32.   NCBI. 2022b. https://www.ncbi.nlm.nih.gov/sra. Accessed December 9, 2022.

466   33.   ENA. 2022. https://www.ebi.ac.uk/ena. Accessed December 9, 2022.

467   34.   Napier G, Campino S, Merid Y, Abebe M, Woldeamanuel Y, Aseffa A, Hibberd ML,

468   Phelan J, Clark TG. 2020. Robust barcoding and identification of *Mycobacterium*

469   *tuberculosis* lineages for epidemiological and clinical studies. *Genome Med* 12:114.

470   35.   Kasai H, Ezaki T, Harayama S. 2000. Differentiation of phylogenetically related slowly

471   growing mycobacteria by their gyrB sequences. *J Clin Microbiol* 38:301-8.

472   36.   Cole ST, Brosch R, Parkhill J, Garnier T, Churcher C, Harris D, Gordon SV, Eiglmeier K,

473   Gas S, Barry CE, Tekaia F, Badcock K, Basham D, Brown D, Chillingworth T, Connor R,

474   Davies R, Devlin K, Feltwell T, Gentles S, Hamlin N, Holroyd S, Hornsby T, Jagels K,

475   Krogh A, McLean J, Moule S, Murphy L, Oliver K, Osborne J, Quail MA, Rajandream

476   MA, Rogers J, Rutter S, Seeger K, Skelton J, Squares R, Squares S, Sulston JE, Taylor K,

477   Whitehead S, Barrell BG. 1998. Deciphering the biology of *Mycobacterium tuberculosis*

478   from the complete genome sequence. *Nature* 396:190-190.

37.   Brosch R, Gordon SV, Marmiesse M, Brodin P, Buchrieser C, Eiglmeier K, Garnier T,

Gutierrez C, Hewinson G, Kremer K, Parsons LM, Pym AS, Samper S, van Soolingen D,

Cole ST. 2002. A new evolutionary scenario for the *Mycobacterium tuberculosis* complex.

Proc Natl Acad Sci U S A 99:3684-3689.

38.   RStudio T. 2020. RStudio: Integrated Development for R. RStudio, PBC, Boston, MA.

URL http://www.rstudio.com/.

39.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search

tool. *J Mol Biol* 215:403-410.

40.   Edgar RC. 2004. Local homology recognition and distance measures in linear time using

compressed amino acid alphabets. *Nucleic Acids Res* 32:380-385.

41.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms. *Mol Biol Evol* 35:1547-1549.

42.   Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New

Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing the

Performance of PhyML 3.0. *Syst Biol* 59:307-321.

43.   Hunt M, Bradley P, Lapierre SG, Heys S, Thomsit M, Hall MB, Malone KM, Wintringer

P, Walker TM, Cirillo DM, Comas I, Farhat MR, Fowler P, Gardy J, Ismail N, Kohl TA,

Mathys V, Merker M, Niemann S, Omar SV, Sintchenko V, Smith G, van Soolingen D,

Supply P, Tahseen S, Wilcox M, Arandjelovic I, Peto TEA, Crook DW, Iqbal Z. 2019.

Antibiotic resistance prediction for *Mycobacterium tuberculosis* from genome sequence

data with Mykrobe. *Wellcome Open Res* 4:191.

44.   Duffy SC, Srinivasan S, Schilling MA, Stuber T, Danchuk SN, Michael JS, Venkatesan

M, Bansal N, Maan S, Jindal N, Chaudhary D, Dandapat P, Katani R, Chothe S, Veerasami

M, Robbe-Austerman S, Juleff N, Kapur V, Behr MA. 2020. Reconsidering

Mycobacterium bovis as a proxy for zoonotic tuberculosis: a molecular epidemiological

surveillance study. *Lancet Microbe* 1:e66-e73.

45. Tortoli E. 2014. Microbiological features and clinical relevance of new species of the genus

Mycobacterium. *Clin Microbiol Rev* 27:727-52.

46. Bottai D, Frigui W, Sayes F, Di Luca M, Spadoni D, Pawlik A, Zoppo M, Orgeur M,

Khanna V, Hardy D, Mangenot S, Barbe V, Medigue C, Ma L, Bouchier C, Tavanti A,

Larrouy-Maumus G, Brosch R. 2020. TbD1 deletion as a driver of the evolutionary success

of modern epidemic *Mycobacterium tuberculosis* lineages. *Nat Commun* 11:684.

47. Coll F, McNerney R, Guerra-Assunção JA, Glynn JR, Perdigão J, Viveiros M, Portugal I,

Pain A, Martin N, Clark TG. 2014. A robust SNP barcode for typing *Mycobacterium*

tuberculosis complex strains. *Nat Commun* 5:4812.

48. Rufai Syed B, McIntosh F, Poojary I, Chothe S, Sebastian A, Albert I, Praul C, Venkatesan

M, Mahata G, Maity H, Dandapat P, Michael Joy S, Katani R, Kapur V, Behr Marcel A.

2021. Complete Genome Sequence of *Mycobacterium orygis* Strain 51145. *Microbiol*

Resour Announc 10:e01279-20.

 TABLE 1 Newly identified 61 *M. orygis* isolates from this study

Isolate	Province	Specimen	Collection year	Gender	Identification (PCR/MIRU-VNTR or BioHansel)
900123	BC	Bronchoalveolar lavage LLL	2009	Female	*M. africanum* by PCR
900124	BC	Bronchoalveolar lavage LUL	2009	Female	*M. africanum* by PCR
900125	BC	Bronchoalveolar lavage RUL	2009	Female	*M. africanum* by PCR
900942	SK	Bronchial wash	2009	Unknown	*M. africanum* by PCR
1501282	BC	Sputum	2015	Female	*M. africanum* by PCR/MIRU-VNTR, MTBC by BioHansel
1600127	BC	Bronchial wash	2016	Female	*M. africanum* by PCR/MIRU-VNTR, MTBC by BioHansel
1601131	MB	Sputum	2016	Female	*M. africanum* by PCR/MIRU-VNTR, MTBC by BioHansel
1601221	AB	Sputum	2016	Female	*M. africanum* by PCR/MIRU-VNTR, MTBC by BioHansel
1601335	MB	Endotracheal	2016	Male	*M. africanum* by PCR/MIRU-VNTR, MTBC by BioHansel
1601614	BC	Bronchial wash	2016	Female	*M. africanum* by PCR/MIRU-VNTR, MTBC by BioHansel
1700074	AB	Stool	2017	Female	*M. africanum* by PCR/MIRU-VNTR, MTBC by BioHansel
1701554	AB	Sputum	2017	Male	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1701642	AB	Sputum	2017	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1701797	BC	Urine	2017	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1701798	BC	Bronchial wash	2017	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1800159	BC	Ascitic fluid	2018	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1800300	AB	Sputum	2018	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1800446	BC	Pericardial fluid	2018	Male	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1800448	BC	Bronchial wash	2018	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel

1800722	AB	Sputum	2018	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1800965	BC	Sputum	2018	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1801127	BC	Sputum	2018	Male	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1801225	BC	Bronchial wash RUL	2018	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1801256	BC	Bronchial wash	2018	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1801535	BC	Sputum	2018	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1900059	BC	Thoracentesis fluid	2019	Male	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1900097	BC	Sputum	2019	Male	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1900228	AB	Right chest wall	2019	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1900323	BC	Bronchial wash	2019	Female	*M. africanum/M. pinnipedii* by PCR/MIRU-VNTR, MTBC by BioHansel
1900674	BC	Abscess L fallopian tube	2019	Female	MTBC by BioHansel
1900698	AB	Sputum	2019	Female	MTBC by BioHansel
1900817	BC	Bronchial wash RUL	2019	Female	MTBC by BioHansel
1901041	BC	Bronchial wash	2019	Male	MTBC by BioHansel
1901282	AB	Sputum	2019	Female	MTBC by BioHansel
1901423	BC	Abscess deep R psoas	2019	Female	MTBC by BioHansel
1901424	BC	Bronchial washing LUL	2019	Female	MTBC by BioHansel
2000368	BC	Bronchial wash RUL	2020	Female	MTBC by BioHansel
2000728	AB	Lymph node fluid	2020	Female	MTBC by BioHansel
2001127	BC	Pleural fluid	2020	Female	MTBC by BioHansel
2001229	BC	Tissue - diaphramatic nodule	2020	Female	MTBC by BioHansel

2001439	BC	Pleural fluid	2020	Male	MTBC by BioHansel
2001463	AB	Lymph node FNA	2020	Female	MTBC by BioHansel
2001464	AB	Sputum	2020	Female	MTBC by BioHansel
2001465	AB	Blood	2020	Male	MTBC by BioHansel
2001479	BC	Bronchial wash R lung	2020	Female	MTBC by BioHansel
2001480	BC	Sputum	2020	Male	MTBC by BioHansel
2100119	AB	Sputum	2021	Female	MTBC by BioHansel
2100187	BC	Bronchial wash	2021	Female	MTBC by BioHansel
2100376	BC	Pleural fluid	2021	Male	MTBC by BioHansel
2100430	AB	Sputum	2021	Female	MTBC by BioHansel
2100725	AB	Lymph node	2021	Female	MTBC by BioHansel
2100976	BC	Sputum	2021	Male	MTBC by BioHansel
2100977	BC	Bronchial wash	2021	Female	MTBC by BioHansel
2101097	BC	Tissue - lymph node mediastinal	2021	Male	MTBC by BioHansel
2101205	AB	Sputum	2021	Female	MTBC by BioHansel
2101352	BC	Sputum	2021	Female	MTBC by BioHansel
2101375	AB	Urine	2021	Female	MTBC by BioHansel
2101452	NS	Lymph node	2021	Unknown	MTBC by BioHansel
2200196	BC	Urine	2022	Female	MTBC by BioHansel
2200263	BC	Sputum	2022	Female	MTBC by BioHansel
2201129	AB	Pleural fluid	2022	Male	MTBC by BioHansel

PCR, Polymerase chain reaction.

MIRU-VNTR, Mycobacterial interspersed repetitive unit - Variable number tandem repeat.

LLL, Left lower lobe; LUL, Left upper lobe; RUL, Right upper lobe; FNA, F*ine needle aspiration.*

AB, Alberta; BC, British Columbia; MB, Manitoba; NS, Nova Scotia; SK, Saskatchewan.

MTBC, Mycobacterium tuberculosis complex.

TABLE 2 Signature single nucleotide polymorphisms (SNPs) in *M. orygis*, gene/genomic position, mutation type, locus, transcript

biotype/function as referenced to H37Rv (NC_000962.3) and Mtb-specific *deletion* 1 region 'TbD1' (AJ426486.1*)

Gene	Gene position	Genomic position of mutation	Nucleotide change	Mutation type	Locus	Transcript biotype/Function	Reference
gyrB	5240-7267	5516	A>G	Nonsynonymous	Rv0005	Protein coding	This study
gyrB	5240-7267	6109	G>A	Synonymous	Rv0005	Protein coding	Huard et al. (17)
gyrB	5240-7267	6717	T>C	Nonsynonymous	Rv0005	Protein coding	This study
Rv2042c	c2287884-2288681	2288569	T>G	Nonsynonymous	Rv2042c	Hypothetical protein	van Ingen et al. (2)
PPE55	c3750780-3753184	3751022	T>G	Synonymous	Rv3347c	PPE family protein	Huard et al. (17)
PPE55	c3750780-3753184	3751023	C>T	Nonsynonymous	Rv3347c	PPE family protein	Huard et al. (17)
leuS	43562-46471	44625	A>T	Nonsynonymous	Rv0041	Protein coding	This study
leuS	43562-46471	44812	G>T	Synonymous	Rv0041	Protein coding	Napier et al. (34)
mmpL6*	735-3638	1220	T>C	Synonymous	TbD1 flanking region	Putative transporter	This study
mmpS6*	292-738	625	C>G	Nonsynonymous	TbD1 flanking region	Putative small membrane protein	This study

'c' in 'Gene position' column for genes *Rv2042c* and *PPE55* denotes complementary sequence.

The numerical positions for the SNPs within *gyrB* gene are provided according to previous nomenclature (accession # L27512.1,

Kasai et al. (35)), while the positions within *Rv2042c, PPE55* and *leuS* genes are relative to *M. tuberculosis* H37Rv (accession #

NC_000962.3).

*The positions for the new SNPs within *mmpL6* and *mmpS6* genes are given as per TbD1 region (accession # AJ426486.1).

TABLE 3 Summary of single nucleotide polymorphisms (SNPs) analysis for *M. orygis* differentiation from other MTBC species and

lineages. Fourteen SNPs on *gyrB,* eleven SNPs on *PPE55*, four SNPs on *Rv2042c*, seven SNPs on *mmpL6*, three SNPs on *mmpS6*, and

twelve SNPs on *leuS* genes are shown

MTBC species/lineage	Gene name and reference position																							
	gyrB														*PPE55*									
																		1177-1178						
	456	520	675	735	756	1011	1113	1116	1311	1410	1450	1458	1721	2097	556	923	1173	1177-1178	1179	1182	1445	1701	2162	2163
M. tuberculosis H37Rv / TbD1	G	A	C	C	G	C	G	G	T	C	G	C	T	T	C	G	G	A	C	C	G	A	G	A
M. bovis AF2122/97	G	A	C	C	A	C	G	G	T	T	T	C	T	T	A	G	G	A	C	C	G	A	G	A
M. bovis BCG 1173P2	G	A	C	C	A	C	G	G	T	T	T	C	T	T	A	G	G	A	C	C	G	G	G	A
M. africanum GM041182	G	A	C	C	G	C	G	G	T	C	T	C	T	T	C	G	G	A	C	C	G	A	G	A
M. orygis 51145	G	G	C	C	G	C	A	G	T	C	T	C	C	T	C	G	G	A	C	C	G	A	A	C
M. canettii CIPT140010059	A	A	C	T	G	T	G	G	T	C	G	T	T	C	C	G	G	A	C	C	A	A	G	A
M. caprae ATCC BAA-824	G	A	C	C	A	C	G	G	G	C	T	C	T	T	A	G	G	A	C	C	G	A	G	A
M. microti ATCC 35782	G	A	T	C	G	C	G	G	T	C	T	C	T	T	-	-	-	-	-	-	-	-	-	-
M. mungi BM22813	G	A	C	C	G	C	G	G	T	C	T	C	T	T	C	G	G	A	C	C	G	A	G	A
M. pinnipedii MP1	G	A	C	C	G	C	G	G	T	C	T	C	T	T	-	-	-	-	-	-	-	-	-	-
Lineage 1 (NC_021740.1)	G	A	C	C	G	C	G	C	T	C	G	C	T	T	C	G	G	A	C	C	G	A	G	A
Lineage 2 (CP007803.1)	G	A	C	C	G	C	G	G	T	C	G	C	T	T	C	G	G	A	C	C	G	A	G	A
Lineage 3 (CP010340.1)	G	A	C	C	G	C	G	G	T	C	G	C	T	T	C	G	C	G	T	G	G	A	G	A
Lineage 4 (NC_020559.1)	G	A	C	C	G	C	G	G	T	C	G	C	T	T	C	G	G	A	C	C	G	A	G	A
Lineage 5 (SRR998618)	G	A	C	C	G	C	G	G	T	C	T	C	T	T	C	T	G	A	C	C	G	A	G	A
Lineage 6 (SRR1577831)	G	A	C	C	G	C	G	G	T	C	T	C	T	T	C	G	G	A	C	C	G	A	G	A
Lineage 7 (ERR1971865)	G	A	C	C	G	C	G	G	T	C	G	C	T	T	C	G	G	A	C	C	G	A	G	A
Lineage 8 (SRR10828835)	G	A	C	C	G	C	G	G	T	C	G	C	T	T	C	G	G	A	C	C	G	A	G	A
Lineage 9 (ERR4162024)	G	A	C	C	G	C	G	G	T	C	T	C	T	T	C	G	G	A	C	C	G	A	G	A

539
540

TABLE 3 Continued …

MTBC species/lineage	Rv2042c				mmpL6							mmpS6			leuS											
	113	134	399	597	306	486	969	1653	1780	1879	2633	334	339	360	321	384	564	1064	1251	1262	1363	1632	2199	2343	2682	2736
M. tuberculosis H37Rv / TbD1	T	A	C	C	C	T	T	C	G	G	T	C	C	C	C	A	A	A	G	A	C	G	T	G	C	T
M. bovis AF2122/97	T	A	C	C	C	T	T	G	G	G	T	C	C	C	C	G	A	A	G	A	C	A	T	G	C	T
M. bovis BCG 1173P2	T	A	C	C	C	T	T	G	G	G	T	C	C	C	C	G	A	A	G	A	C	A	T	G	C	T
M. africanum GM041182	T	A	C	C	T	T	T	C	G	G	T	C	C	C	C	G	A	A	G	A	C	G	T	G	C	T
M. orygis 51145	G	A	C	C	C	C	T	G	G	G	T	G	C	C	C	G	A	T	T	A	C	G	T	G	C	T
M. canettii CIPT140010059	T	G	T	C	C	T	C	C	G	C	C	C	C	C	A	G	A	A	G	A	C	G	C	G	T	T
M. caprae ATCC BAA-824	T	A	C	C	C	T	T	G	G	G	T	C	C	C	C	G	A	A	G	C	C	G	T	G	C	T
M. microti ATCC 35782	T	A	C	C	C	T	T	G	G	G	T	C	C	C	C	G	G	A	G	A	C	G	T	G	C	T
M. mungi BM22813	T	A	C	C	C	T	T	C	G	G	T	C	C	C	C	G	A	A	G	A	C	G	T	A	C	T
M. pinnipedii MP1	T	A	C	C	C	T	T	G	G	G	T	C	C	C	C	G	G	A	G	A	C	G	T	G	C	T
Lineage 1 (NC_021740.1)	T	A	C	C	-	-	-	-	G	G	T	-	-	-	C	G	A	A	G	A	C	G	T	G	C	T
Lineage 2 (CP007803.1)	T	A	C	G	-	-	-	-	G	G	T	-	-	-	C	A	A	A	G	A	C	G	T	G	C	T
Lineage 3 (CP010340.1)	T	A	C	C	-	-	-	-	G	G	T	-	-	-	C	A	A	A	G	A	C	G	T	G	C	T
Lineage 4 (NC_020559.1)	T	A	C	C	-	-	-	-	G	G	T	-	-	-	C	A	A	A	G	A	C	G	T	G	C	T
Lineage 5 (SRR998618)	T	A	C	C	C	T	T	C	G	G	T	C	C	T	C	G	A	A	G	A	C	G	T	G	C	C
Lineage 6 (SRR1577831)	T	A	C	C	T	T	T	C	G	G	T	C	C	C	C	G	A	A	G	A	C	G	T	G	C	T
Lineage 7 (ERR1971865)	T	A	C	C	C	T	T	C	C	G	T	C	T	C	C	A	A	A	G	A	C	G	T	G	C	T
Lineage 8 (SRR10828835)	T	A	C	C	-	-	T	C	G	G	T	C	C	C	C	G	A	A	G	A	C	G	T	G	C	T
Lineage 9 (ERR4162024)	T	A	C	C	C	T	T	C	G	G	T	C	C	C	C	G	A	A	G	A	T	G	T	G	C	T

The nucleotides in red color show unique SNPs for a particular species/lineage, represented in a different colored block.

'-' indicates that the nucleotides for those positions were not extracted in the analysis.

The reference nucleotide positions for *gyrB* gene are given according to gene sequence described by Kasai et al. (35) (accession #

L27512.1).

The nucleotide positions within *PPE55*, *Rv2042c* and *leuS* genes are relative to *M. tuberculosis* H37Rv (accession # NC_000962.3, Cole

et al. (36)).

The nucleotide positions for *mmpL6* and *mmpS6* genes are provided according TbD1 region (accession # AJ426486.1, Brosch et al.

(37)).

FIG 1 Phylogeny of *M. orygis* based on single nucleotide variants (SNVs) analysis using SNVPhyl pipeline v.1.2.3. A maximum

likelihood tree of 61 newly sequenced *M. orygis* isolates from this study and the *M. orygis* genomes collected from public repositories.

The outer ring shows the geographic location of *M. orygis* (Canada in green, India in blue, Norway in pink, Switzerland in cyan, UK in

red, and USA in yellow), and the inner ring denotes the source of *M. orygis* (animal in orange, and human in purple). *M. tuberculosis*

H37Rv (NC_000962.3) used as a reference in this analysis. Scale bar indicates average number of substitutions per site. The studied

isolates mostly clustered with *M. orygis* of human origin recovered from different geographic regions. The isolate 2100725 showed a

good phylogenetic relation (pairwise distances of 34 SNVs) with seven *M. orygis* of animal origin from India.