Sing to me, baby:

Infants show neural tracking and rhythmic movements to live and dynamic maternal singing

Trinh Nguyen[1,2]*, Susanne Reisner[1], Anja Lueger[1], Samuel V. Wass[3], Stefanie Hoehl[1], & Gabriela Markova[1,4]*

[1]Faculty of Psychology, University of Vienna, Liebiggasse 5, 1010 Vienna, Austria
susanne.reisner@univie.ac.at, anja.lueger@univie.ac.at, stefanie.hoehl@univie.ac.at, gabriela.markova@univie.ac.at

[2]Neuroscience of Perception and Action Lab, Italian Institute of Technology, Viale Regina Elena 291, 00161 Rome, Italy.
trinh.nguyen@iit.it

[3]Department of Psychology, University of East London, University Way, London, E16 2RD, United Kingdom
s.v.wass@uel.ac.uk

[4]Institute for Early Life Care, Paracelsus Medical University, Strubergasse 13, 5020 Salzburg, Austria

*Corresponding authors: Trinh Nguyen, Viale Regina Elena, 291, 00161 Rome, Italy, trinh.nguyen@iit.it; Gabriela Markova, Liebiggasse 5, 1010 Vienna, Austria, +431427747478, gabriela.markova@univie.ac.at.

Abstract

Infant-directed singing has unique acoustic characteristics that may allow even very young infants to respond to the rhythms carried through the caregiver's voice. The goal of this study was to examine neural and movement responses to live and dynamic maternal singing in 7-month-old infants and their relation to linguistic development. In total, 60 mother-infant dyads were observed during two singing conditions (playsong and lullaby). In Study 1 ($n = 30$), we measured infant EEG and used an encoding approach utilizing ridge regressions to measure neural tracking. In Study 2 ($n = 30$), we coded infant rhythmic movements during the mother's singing. In both studies, we assessed children's vocabulary when they were 20 months old. In Study 1, we found above-threshold neural tracking of maternal singing, with superior tracking of lullabies than playsongs, and the acoustic features of infant-directed singing modulated tracking. In Study 2, infants showed more rhythmic movement to playsongs than lullabies. Importantly, neural coordination and rhythmic movement to playsongs were positively related to infants' expressive vocabulary at 20 months. These results highlight the importance of infants' brain and movement coordination to their caregiver's musical presentations, potentially as a function of musical variability.

Keywords: infant-directed singing, social interaction, EEG, neural tracking, rhythmic movement

Sing to me, baby:

Infants show neural tracking and rhythmic movement to live and dynamic maternal singing

Musicality, the ability to perceive, process, and produce music, is a universal human ability with deep ontogenetic roots (e.g., Mehr et al., 2021; Savage et al., 2021). Fetuses at 35 weeks of gestational age already show more movement to music in comparison to speech (Kisilevsky et al., 2004). After birth, infants are exposed to a variety of musical stimuli throughout the day. Most prominently, caregivers sing to them to capture their attention, regulate their arousal, and share emotions with or entertain them (Trehub, 2019). While infants seem to show neural tracking of rhythmic beats (Cirelli et al., 2016) as well as meter (Lenc et al., 2022) and move their bodies more to music than to other auditory stimuli (Ilari, 2015; Zentner & Eerola, 2010), no research to date has examined whether young infants also show neural tracking of and movement changes in response to their caregivers' singing and what the developmental outcomes of such coordination processes might be. Consequently, the goal of the present study was to examine 7-month-old infants' neural and sensorimotor responses to live maternal infant-directed (ID) singing and its linguistic developmental consequences. Examining the sensorimotor processes underlying infants' engagement with ID singing is key to understanding the communicative functions of early musicality.

From early in life, children have sophisticated perceptual capabilities when listening to music. Infants are sensitive to tempo- and pitch-related organization of musical sequences, such as changes in melodic patterns, harmony, grouping, beat, and meter (e.g., Ferland & Mendelson, 1989; Háden et al., 2015; Krumhansl & Jusczyk, 1990; Trainor & Heinmiller, 1998; Winkler et al., 2009). This early sensitivity to musical information allows even very young children to coordinate with the music they hear. Such coordination, in particular to the rhythmic structure of the musical input, modifies brain activity and behavior in such a way that they eventually lock onto a common periodicity (Vuilleumier & Trost, 2015). For example, Cirelli et al. (2016) showed that 7-month-old infants' brains track the beat and meter of rhythmic patterns. At the behavioral level, infants coordinate their gaze with the beat of ID singing as early as 2 months of age (Lense et al., 2022), and adjust their movements to the tempo of the music they hear by 6 months (Ilari, 2015; Zentner & Eerola, 2010). This evidence suggests that young infants are not only receptive to but often also coordinate with music and other rhythmic sounds within their abilities.

Research examining infant musicality used predominantly pre-recorded or digitalized stimuli that are easier to control and manipulate but have a phenomenologically different quality than live musical performances (Trehub, 2017). In contrast, caregivers sing to infants on a daily basis (Steinberg et al., 2021), and ID singing is part of intimate interactions between caregivers and infants with the goal of engaging or soothing (Trainor et al., 1997). ID singing, compared to non-ID singing, is characterized by high pitch, slow tempo, enhanced temporal regularity, and greater dynamic range (Nakata & Trehub, 2011; Trainor et al., 1997). ID songs are often accompanied by idiosyncratic gestures (Eckerdal & Merker, 2009), energetic movement, and caregivers' positive affect (Cirelli et al., 2020; Trehub et al., 2016), all of which are used by caregivers to attract infants' attention, activate, and engage with them (Cirelli et al., 2020; Trehub & Trainor, 1998). Moreover, caregivers adjust their facial expressions and gaze to the timing of the ID songs (Lense et al., 2022). Still, it remains unclear how infants perceive and act upon *live* and *dynamic* ID singing.

The acoustic features of ID songs can be further specified by the context in which they are used, namely, to play or to soothe. Playsongs are characterized by higher rhythmicity, tempo, and pitch level, as well as more dynamic variability and thus increased complexity in comparison to lullabies (Cirelli et al., 2020; Rock et al., 1999; Trehub & Trainor, 1998). On the other hand, lullabies are characterized by lower tempo, pitch, and less variability, intending to soothe and calm an infant. Accordingly, the different acoustic features might affect how infants coordinate their neural and movement responses. Recent research in adults shows that neural tracking was enhanced in slower music (Weineck et al., 2022) and was guided by melodic expectations (Di Liberto et al., 2020). We thus expected enhanced neural tracking and coherence in slower and "simpler" ID songs (i.e., lullabies), while we hypothesized that the complexity of playsongs might attenuate infants' neural tracking. Importantly, the specific acoustic features of lullabies and playsongs might have different effects on infants' rhythmic movement than on neural tracking. On the behavioral level, increased rhythmic complexity was related to more movement and reported groove (i.e., the inclination to move) in children and adults (Cameron et al., 2022; Kragness et al., 2022). The enhanced complexity of playsongs might thus be more engaging to children.

Infants' underlying cognitive resources and neural substrates for communicative musicality are shared with other domains, such as the processing of speech (Brandt et al., 2012). Brandt and colleagues (2012) propose that the musical aspects of language (i.e.,

prosody, rhythm, timbral contrast) scaffold the later development of semantic and syntactic aspects of language. In fact, available evidence suggests that an enriched musical environment (i.e., music classes, music at home) during infancy and preschool can promote language development (Papadimitriou et al., 2021; Politimou et al., 2019; Putkinen et al., 2013; Williams et al., 2015; Zhao & Kuhl, 2016). Moreover, rhythmic motor abilities provide relevant practice of rhythmic, closely timed actions, as they are required for speech (Iverson, 2010; Iverson et al., 2007). There is also, albeit limited, evidence suggesting a link between ID singing and language development. Brain responses of newborns to sung as compared to spoken streams of syllables predict expressive vocabulary at 18 months (François et al., 2017), and ID singing has been found to facilitate aspects of phonetic perception and word learning in infants in the second half of the first year of life (Lebedeva & Kuhl, 2010; Thiessen & Saffran, 2009). In addition, recent evidence showed that neural tracking of nursery rhymes is predictive of infants' later language outcomes (Attaheri et al., preprint, Menn et al., 2022b). Yet, it remains unknown whether infants' coordination with live ID singing is also related to their language development.

Overall, we still know very little about infants' neural coordination and rhythmic movement to live and dynamic maternal singing. Closing this gap will help us understand how we might be able to facilitate infants' physical, social, and cognitive development through musical communication. In the present study, we used the naturally occurring variance in acoustic features of playsongs and lullabies during live and dynamic ID singing to gain a deeper understanding of the tracking effects on infants' bodies and brains. In Study 1, we measured 7-month-old infants' neural tracking (using EEG) to maternal singing. In order to avoid motion artifacts, we observed an additional sample of 7-month-old infants' rhythmic movements to maternal singing in Study 2. We hypothesized that infants' neural responses are more coordinated with lullabies than playsongs. In contrast, their rhythmic movements were expected to be more frequent with playsongs than lullabies. At 20 months, we assessed infants' vocabulary to examine whether early musical engagement relates to infants' language development. We hypothesized that higher neural tracking of ID songs, as well as more rhythmic movements, predict children's vocabulary.

Methods

Participants

In Study 1, 30 7-month-old infants (13 females; age: 228.80 ± 8.20 days [*M* ± *SD*]) and their mothers (age: 33.96 ± 4.88 years) were included in the final sample. An additional 29 infants were tested but could not be included due to fussiness (*n* = 4), technical problems during the recording (*n* = 9), and bad signal quality due to the naturalistic setup (*n* = 16). In Study 2, 30 7-month-old infants (13 female; age: 226.67 ± 6.1 days) and their mothers (age: 35.10 ± 4.00 years) were included in the final sample. We excluded 16 additional mother-infant dyads from the final analysis due to infant fussiness (*n* = 4), equipment failure (*n* = 11), or because the mother did not follow instructions (*n* = 1).

Participants were recruited from a database of families who expressed interest in taking part in developmental research and have consented to be contacted by our staff. These families were recruited in neonatal units at local hospitals, in mother-child activity classes, and through social media. All infants were born full-term (38-42 weeks gestational age), had a birth weight of > 2500 g and were without any known developmental delays or neurological or hearing impairments. Infants grew up in predominantly German-speaking households. Mothers were highly educated, with 93.1% (Study 1) and 86.7% (Study 2) of mothers holding a university degree. Fifty-five percent of the mothers in both samples reported playing an instrument, and 20% were singing in a choir or a band. The study was conducted according to the declaration of Helsinki, approved by the university's ethics committee, and parents provided written informed consent.

Procedure

After arrival in the laboratory, parents and infants were familiarized with the environment. Parents were informed about the study and signed a consent form. In Study 1, the EEG setup was prepared while the infant was sitting on the parent's lap. Infants were seated in an infant car seat, and their mothers faced them (*Figure 1A*). Study 2 followed the same procedure as Study 1, except that infants sat upright in an infant highchair that allowed more movement. In Study 2, we additionally measured infants' electro-cardiac rhythms using a two-lead derivation. These results will be reported elsewhere.

Each dyad was observed during two experimental singing blocks (*Figure 1B*), the order of which was randomized between participants. Each singing block was preceded and

followed by a baseline (i.e., three in total), during which infants watched videos of slowly moving shapes for 60 s without further communication. During each singing block, mothers continued holding the tablet showing a calm aquarium video to help with infant fussiness. Mothers were instructed to sing one of two types of songs: a playsong or a lullaby. Songs were selected according to their level of familiarity during pilot testing. All mothers sang the same songs and were informed about and provided recordings of the songs at the time of recruitment. The majority (87%) of mothers were familiar with the playsong, and 97% of mothers were familiar with the lullaby. All mothers sang four verses of each song and repeated each song, resulting in eight verses for each song. Mothers were prompted with a metronome before the commencement of each song (i.e., playsong = 170 bpm; lullaby = 100 bpm). Three cameras (synchronized in VideoSyncPro by Mangold International) were used to record the experiment.

Figure 1. (A) Illustrated experimental setup. An experimenter held the car seat with the infant facing their mother in Study 1. Video cameras were used to record the mother-infant dyad from two angles. (B) Procedure. Mothers sang two repetitions of four verses for each lullaby and playsong. During both conditions, a tablet held by the mother played a video of fish swimming in an aquarium. A 60-second baseline, during which the mother did not

interact with the infant, preceded, and followed each song, during which the tablet showed moving geometric shapes.

Measures

Maternal Singing

Audio of maternal singing was recorded using a portable microphone (Mangold International) at a 44,100 Hz sampling rate. Audio data were pre-processed using the software Audacity (https://www.audacityteam.org/). Excerpts containing infant vocalizations and environmental noise were manually removed. To examine whether mothers differentiated between the two singing conditions, we analyzed maternal audio data in terms of mean and standard deviation in tempo (beats per minute), pitch (Hz), root mean square (RMS; perceived loudness), and pulse clarity using MIRtoolbox (Lartillot et al., 2008).

In order to extract the sound envelopes for further EEG analyses (Jessen et al., 2021), the audio data were then submitted to the NSL toolbox (http://nsl.isr.umd.edu/downloads.html). The resulting matrix contained band-specific envelopes of 128 frequency bands of uniform width on a logarithmic scale with center frequencies logarithmically spaced between 0.1 and 4 kHz. We obtained the broadband temporal envelope of the audio soundtrack by summing up band-specific envelopes across all frequencies to obtain one temporal envelope. The spectrogram of the envelope is included in the supplements (Figure S1). Lastly, the amplitude envelope data were down sampled to match the 500 Hz sampling rate of the EEG data.

Infant EEG (Study 1)

We recorded infant EEG using a BrainAmp DC 64-Channel System and a 32 channels ActiCap system (Brain Products, Germany) containing active Ag/AgCl electrodes, mounted according to scalp locations of the 10-20 system. Horizontal and vertical electrooculograms were recorded bipolarly to control for eye movements. EEG data were sampled at 500 Hz. Impedances were controlled at the beginning of the experiment and accepted when below 20 kΩ.

EEG data were processed using the *eeglab* (Delorme & Makeig, 2004) and *fieldtrip* toolboxes (Oostenveld et al., 2010) as well as custom MATLAB code. EEG data were band-

pass filtered with a noncausal finite impulse response filter (0.3 Hz - 100 Hz, -6 dB cutoff: 0.15 Hz - 100.15 Hz). Next, we reduced line noise (50/100 Hz) by using ZapLine from NoiseTools (de Cheveigné, 2020). Noisy channels were identified by assessing the normed joint probability of the average log power from 1 – 100 Hz and rejected if exceeding a threshold of 3 SD from the mean (number of removed channels: $M = 4.61$; *range* $= 0 – 10$). This step was repeated twice. Next, we used wavelet thresholding to identify and remove artifacts (threshold = hard, wavelet = coif4, level-dependent thresholding; Lopez et al., 2022). The previously rejected channels were then interpolated using spherical splines. Afterward, all channels were re-referenced to the average of all scalp electrodes (F7, F8, F9, F3, Fz, F4, FT7, FC3, FCz, FC4, FT8, T8 C4, Cz, C3, T7, CP3, CP4, TP10, P8, P4, Pz, P3, P7, PO9, PO10, O1, Oz, O2) and segmented into non-overlapping 1-s-epochs. In preparation for further analysis, epochs were automatically examined to see if they were still contaminated by artifacts. The standard deviation was computed in a sliding window of 200 ms. If the standard deviation exceeded 100 µV☐ at any electrode, the entire epoch was discarded. Participants were excluded from further analysis if less than 90 artifact-free epochs could be retained. Infants' processed EEG data comprised on average 144 epochs from the playsong condition and 126 epochs from the lullaby condition. The spectrogram of the data is provided in Figure S2. EEG data were then FIR-type bandpass filtered between 1 – 10 Hz (Jessen et al., 2019).

Neural tracking analyses. To quantify the degree to which the 7-month-old infants showed neural tracking of maternal singing, we used encoding models (i.e., temporal response functions [TRF]) to predict the infants' EEG from the amplitude envelope of mothers' singing. The temporal response function regresses the neural response onto the auditory signal, dividing out the autocovariance structure of the stimulus from the model. The response model is defined as:

$$r(t, n) = \sum_l \quad w(l, n)s(t - l) + \varepsilon(t, n)$$

where $r(t, n)$ is the neural response, across each channel included in the model *(n)*, across all time points *(t)*; *w* is the TRF that describes the linear transformation of the stimulus *(s)* to the ongoing neural response *(t)*, and ε is the residual error at each channel not explained by the model.

The TRF (w) is estimated using the following equation:

$$w = (S^T S + \lambda I)^{-1} S^T r$$

where $S^T r$ is the result of convolution between the zero-lagged neural response *(r)*, and the speech stimulus at different time lags (S), $S^T S$ is the autocovariance matrix of the stimulus at each time lag which is divided out from the linear relationship modeled in $S^T r$. I is the identity matrix, and λ is the ridge parameter, which enforces a smoothness constraint on the output of the model (w). The output (w) is a time-windows by EEG channels matrix, containing the resulting TRF weights for each channel at each time window (see Crosse et al., 2016 for further explanations). A constant term is included in the model by concatenating a column of ones to the left of *S*.

A method of ridge regression is used to improve the reliability of the estimated coefficients and prevent over-fitting. Ridge regression works by introducing a bias term to the model, which penalizes TRF values as a function of their distance from 0 (Crosse et al., 2016). This reduces over-estimation problems in *w* and reduces the likelihood of high frequency artifacts in the estimated TRF model (Fiedler et al., 2017; Haufe et al., 2014) .

Following Jessen et al. (2019), we estimated the predictive accuracy of the model by using individual models (computation of individual response function for each infant). We assumed individual models to be suitable for our purposes as ID songs were highly different between dyads, therefore leading to vastly different regressor weights between infants. To that end, 80% of the available data for a given participant was used to train the model. The resulting response function was then correlated with the response observed in the remaining 20% of the data. To accommodate a time lag between changes in the EEG signal and changes in the stimulus signal, predictions were computed over a range of time lags (i.e., 2 ms) between -100 and 500 ms later than the stimulus signal (see Jessen et al., 2021, for further details).

We obtained the optimal regularization parameter λ by training the respective model on the concatenated EEG and audio data of each infant using a variety of λ values between 10^{-7} and 10^7. We increased the exponent in steps of 1 and used the resulting models to predict the EEG signal for each participant. By doing so, we obtained a total of fourteen different

models (and predictions) based on the different λ parameters. For each of these 14 different models, we computed the mean response function across n-1 participants and used this response function to predict the EEG of the n[th] fold (i.e., n-fold leave-one-out cross-validation). One fold comprised 1/5 of the data. Finally, we computed the predictive accuracy (i.e., Pearson's correlation coefficient *r* between the predicted EEG and the actual EEG) for each infant and each electrode, resulting in 31 accuracy values per infant and stimulus parameters for each λ value. For each infant, stimulus parameter, and electrode set, we chose the λ value for which the model yielded the highest correlation between the predicted and the actual EEG time-series (Jessen et al., 2021).

For statistical evaluation, we computed two different predictive accuracies per infant and song conditions. First, we computed the correlation between the predicted response generated on a model trained on 80% of the data on the other 20% of the data ("individual model"). Second, a permuted or null predictive accuracy ("circular shift control") was obtained. Before calculating accuracy this way, we reversed and applied a random circular shift to the true EEG time series for participant *n* (to ensure exceeding the potential autoregressive structure of the EEG) and computed the correlation between the shifted EEG and the predicted response from the individual model (for $n = 400$ permutations). The permutations were then averaged, resulting in one so-called "randomized" predictive accuracy value per infant.

To evaluate the resulting response functions, we computed a cluster-based permutation test with 1000 randomizations, testing the obtained response functions against zero. A cluster was defined along with the dimensions of time (lags between the sound envelope and the EEG) and electrode position, with the constraint that a cluster had to extend over at least two adjacent electrodes. A type-1 error probability of less than 0.05 was ensured at the cluster level (Meyer et al., 2021). The cluster-based permutation analysis revealed no significant clusters in the TRF potentially due to the large individual differences between the infants' evoked responses as well the acoustic features of the ID songs. The detailed results of this evaluation are reported in the supplements (see Figure S3).

Infant Rhythmic Movement (Study 2)

We coded the duration of infant rhythmic movement during the two singing conditions. Infant rhythmic movement was defined as at least three repetitions of the same movement in a body part or the entire body at regular short intervals (1 second or less; Thelen, 1979) We calculated proportionate movement values to compensate for differences in recording length. To ensure inter-rater reliability, two independent observers coded 30% of all videos yielding high inter-rater reliability with κ = .83.

Infant Gaze

To control for infant visual attention in both studies, we coded the following categories of infant gaze from video recordings using INTERACT by Mangold: (1) *gaze at mother's face*; (2) *gaze at mother's body; (3) gaze at tablet*; (4) *gaze away* as infant gaze directed away from their mother and the tablet at something else in their surrounding (Markova & Legerstee, 2006). Coding was conducted without sound, frame-by-frame, and each dwell time had to last at least 1 s. Inter-rater reliability was assessed on 30% of data coded by three pairs of independent observers, yielding moderate to excellent inter-rater reliability within a range of κ = .70-.99 for all categories. Analyses regarding infant gaze are reported in the supplements (Table S1).

Questionnaires

To assess language development, we used the Austrian Communicative Development Inventory - Level 2SF (ACDI-2 SF; Marschik et al., 2007) that mothers filled out when children were 20 months of age (age: 20.23 ±1.35 months [$M \pm SD$]). Further self-reports on mothers' depression symptoms, anxiety levels after singing, and familiarity of infants with the songs are included in the supplements.

Statistical Analysis

All statistical analyses were conducted in RStudio (RStudio Team, 2022).

Study 1

First, we ran a linear mixed-effects model comparing the prediction accuracy values between song type (playsong vs lullabies), data type (true: predicted infant EEG from original sound envelopes correlated with the original infant EEG vs shifted: the correlation of the

predicted infant EEG from the circular shifted sound envelope with the original infant EEG) in each channel. The prediction accuracy values were Fisher's z transformed.

Prediction accuracy ~ song type * data type * channel + (1+ song type + data type | ID)

Next, we included the acoustic features of the playsong and lullaby (i.e., tempo, pitch, pulse clarity, and RMS) as fixed and interaction effects to examine whether differences in these features were related to differences in infants' neural tracking of ID singing.

Prediction accuracy ~ song type * (tempo+pitch+pulse clarity +RMS) + channel + (1 | ID)

Prediction accuracy ~ song type * (tempo.sd+pitch.sd+pulse clarity.sd +RMS.sd) + channel + (1 | ID)

Finally, we included infants' expressive vocabularies at 20 months as fixed and interaction effects to examine whether variation in their vocabulary was related to differences in infants' neural tracking of ID singing.

Prediction accuracy ~ song type * vocabulary + channel + (1 | ID)

Study 2

We ran a generalized linear mixed-effects model comparing the frequency of rhythmic movements in both lullaby and playsong conditions.

Rhythmic movement ~ song type + (1 + song type | ID)

Next, we included the acoustic features of the playsong and lullaby (i.e., tempo, pitch, pulse clarity, and RMS) as fixed and interaction effects to examine whether differences in these features were related to differences in infants' rhythmic movement during ID singing.

Rhythmic movement ~ song type * (tempo+pitch+pulse clarity +RMS) + (1 | ID)

Rhythmic movement ~ song type * (tempo.sd+pitch.sd+pulse clarity.sd+RMS.sd) + (1 | ID)

Finally, we included infants' expressive vocabularies at 20 months as fixed and interaction effects to examine whether variation in their vocabulary was related to differences in infants' rhythmic movement during ID singing.

*Rhythmic movement ~ song type * vocabulary + (1 | ID)*

Results

Acoustic Features of Maternal Singing (Study 1 & 2)

First, we compared the acoustic features of mothers' singing the playsong and the lullaby (see Table 1 for descriptive statistics). Multiple comparisons were corrected for using the false discovery rate, which is indicated by the q-value (Benjamini & Hochberg, 1995). Playsongs were sung louder ($\chi^2(1) = 34.15$, $q < .001$), faster ($\chi^2(1) = 40.77$, $q < .001$), and in a higher pitch ($\chi^2(1) = 8.83$, $q = .004$) than lullabies. Playsongs and lullabies did not differ in pulse clarity ($q = .248$). Playsong varied more in perceived loudness ($\chi^2(1) = 35.97$, $q < .001$), tempo ($\chi^2(1) = 6.79$, $q = .012$), and pitch ($\chi^2(1) = 57.50$, $q < .001$) compared to lullabies. Lullabies varied more in pulse clarity compared to playsongs ($\chi^2(1) = 4.505$, $q = .034$).

Table 1. Descriptive statistics on acoustic features of live-sung lullabies and playsongs

	Lullaby				Playsong			
Variables	*M*	*SD*	*min*	*max*	*M*	*SD*	*min*	*max*
RMS (perceived loudness)	0.045	0.035	0.009	0.184	0.056	0.033	0.009	0.210
RMS SD	0.028	0.020	0.006	0.117	0.037	0.021	0.005	0.136
Tempo (BPM)	137.55	26.18	95.97	199.63	163.63	23.51	109.95	199.81
Tempo SD (BPM)	33.143	6.174	21.808	44.532	35.842	6.770	20.869	48.351
Pitch (Fundamental frequency f0, Hz)	260.51	36.29	167.39	368.70	268.64	32.71	168.04	322.56
Pitch SD (Hz)	34.138	6.385	23.297	60.007	42.222	6.229	0.011	0.009
Pulse clarity	0.059	0.030	0.010	0.147	0.065	0.026	0.008	0.145
Pulse clarity SD	0.064	0.011	0.047	0.104	0.061	0.009	0.039	0.081

Neural Tracking of Maternal Singing (Study 1)

First, we compared the predictive accuracy obtained by the individual models of both playsong and lullaby. The model results displayed a significant fixed effect of ID song type ($\chi^2(1) = 42.135$, $p < .001$), a fixed effect of data type ($\chi^2(1) = 2430.361$, $p < .001$), and an interaction effect between ID song type and data type, $\chi^2(1) = 40.881$, $p < .001$. The set of effects was further tested in post-hoc contrasts (corrected for multiple comparison) and showed that infants' neural tracking of ID singing (data type true) was more accurate (estimates = 0.065, SE = 0.002, 95% CI = [0.062 0.068]) in comparison to the randomized control (data type shifted; estimates = 0.001, SE = 0.002, 95% CI = [0.004 0.004]). In addition, neural tracking of lullabies (estimates = 0.074, SE = 0.002, 95% CI = [0.070 0.077]) was more accurate than that of playsongs (estimates = 0.057, SE = 0.002, 95% CI = [0.053 0.060], Figure 2A) in data type true, while neural tracking of the two song types did not differ in data type shifted, $p = .999$. Other effects, such as the fixed effect of channel, the interaction effect between channel and type of song, the interaction effect between channel and data type, and the three-way interaction between channel, type of song, and data type were not significant ($p > .086$). In sum, we found above-threshold neural tracking of lullabies and more accurate tracking for lullabies than playsongs across the infant brain.

Figure 2. (A) Graph depicting the interaction effect between true vs. shifted data (x Axis) in the different types of ID songs (columns) on prediction accuracy (y Axis). The difference in prediction accuracy (y Axis) between song types (columns) is modulated by (B) the

perceived loudness of playsongs (x Axis), (C) the tempo of lullabies (x Axis), and (D) the pulse clarity of playsongs (x Axis).

Next, we examined the acoustic features, namely tempo, pitch, pulse clarity, RMS, and their variability on the predictive accuracies of mTRF models tested on the original data. First, the mean of acoustic features was included as fixed and interaction effects in the linear mixed-effects model. The model output model showed that the fixed effect of ID song type remained when controlling for the acoustic features, $\chi^2(1) = 16.164$, $p < .001$. There were, however, significant interaction effects between ID song type and RMS, $\chi^2(1) = 14.264$, $p < .001$ (Figure 2B), ID song type and tempo, $\chi^2(1) = 4.925$, $p = .026$ (Figure 2C), as well as ID song type and pulse clarity, $\chi^2(1) = 9.796$, $p = .002$ (Figure 2D). Post-hoc analyses of the interaction effects showed that lower RMS (perceived loudness) was related to more accurate neural tracking of the playsong (*trend* = 0.015, *SE* = 0.004, *95% CI* = [-0.024 -0.008]), while RMS was not related to neural tracking accuracy of lullabies (*trend* = -0.003, *SE* = 0.003, *95% CI* = [-0.009 0.002]). Moreover, slower-sung lullabies were related to more accurate neural tracking (*trend* = -0.008, *SE* = 0.004, *95% CI* = [-0.016 -0.001]), while the tempo of playsongs did not relate to neural tracking (*trend* = 0.000, *SE* = 0.002, *95% CI* = [-0.004 0.005]). Higher pulse clarity in playsongs was further related to enhanced neural tracking (*trend* = 0.010, *SE* = 0.003, *95% CI* = [0.004 0.015]), but did not affect neural tracking of lullabies (*trend* = -0.003, *SE* = 0.002, *95% CI* = [-0.008 0.002]).

In the next step, we tested the variability of the acoustic features as fixed and interaction effects on neural tracking. We found significant fixed effects of ID song type ($\chi^2(1) = 8.866$, $p = .003$) and variability of tempo ($\chi^2(1) = 5.972$, $p = .015$), indicating that neural tracking was enhanced when the tempo of ID songs varied less, estimate = -0.008, SE = 0.003, *95% CI* = [-0.013 -0.002]. There were also significant interaction effects between ID song type and RMS variability, $\chi^2(1) = 11.762$, $p = .001$, as well as ID song type and pitch variability, $\chi^2(1) = 11.329$, $p = .008$. Post-hoc analyses showed that lower RMS variability was related to more accurate neural tracking of playsongs (*trend* = -0.011, *SE* = 0.004, *95% CI* = [-0.019 -0.004]), while RMS variability was not related to neural tracking of lullabies (*trend* = -0.000, *SE* = 0.003, *95% CI* = [-0.006 0.005]). Lower pitch variability was related to enhanced neural tracking of lullabies (*trend* = -0.006, *SE* = 0.003, *95% CI* = [-0.012 -0.001]), while it was not significantly associated with neural tracking of playsongs (*trend* = 0.006, *SE*

= 0.004, *95% CI* = [-0.001 0.013]). All other acoustic features were not related to predictive accuracy (*p* > .063).

Rhythmic Movement to Maternal Singing (Study 2)

We tested whether infants showed more rhythmic movement during the lullaby or the playsong condition. The generalized linear mixed-effects model output displayed a significant condition effect, $\chi^2(1) = 9.514$, *p* = .004 (Figure 3A). Model estimates revealed that infants showed less rhythmic movement in the lullaby condition (*estimate* = 3.56, *SE* = 0.09, *95% CI* = [3.39 3.74]) than in the playsong condition (*estimate* = 3.72, *SE* = 0.07, *95% CI* = [3.58 3.85]).

Figure 3. (A) Graph depicting the fixed effect of the type of ID song (x Axis) on rhythmic movement frequency (y Axis). (B) Infants generally showed more rhythmic movements (y Axis) when mothers varied the perceived loudness (x Axis) during both song types (columns).

Next, we tested the role of the acoustic features of maternal ID singing on infants' rhythmic movement and found only one significant association between the variability in RMS and the amount of rhythmic movement, *estimate* = 0.157, *SE* = 0.095, *95% CI* = [3.39 3.74], $X^2(1)$ = 5.673, *p* = .017 (Figure 3B). All other associations with mean values and standard deviations of acoustic features were not significant (*p* > .113).

Relations Between Neural Tracking (Study 1), Rhythmic Movement (Study 2), and Children's Language Development

Finally, we examined whether infants' neural tracking and rhythmic movement in response to maternal ID singing are related to their expressive vocabulary at 20 months. The model outputs for Study 1 revealed a significant correlation between vocabulary size at 20 months and neural tracking of ID songs, $\chi^2(1)$ = 5.236, *p* = .022. The post-hoc analysis (see Figure 4A) revealed a significant, but not robust, trend for enhanced neural tracking of playsongs in relation to larger vocabulary size at 20 months (*trend* = 0.002, *SE* = 0.002, *95% CI* = [-0.002 0.006]), and no significant relation between neural tracking of lullabies and vocabulary size (*trend* = -0.002, *SE* = 0.002, *95% CI* = [-0.006 0.002]). Results for Study 2 showed that infants' rhythmic movement was positively related to infants' expressive vocabulary depending on the song type, $\chi^2(1)$ = 4.933, *p* = .026. Specifically, post-hoc trend

analysis showed that infants' rhythmic movement during the playsong, but not the lullaby, was positively related to their vocabulary (see Figure 4B). The trend was, however, not robust (*trend* = 0.072, *SE* = 0.068, *95% CI* = [-0.062 0.205]).

Figure 4. (A) Graph depicting the relation between neural tracking (x Axis) and (B) rhythmic movement at 7 months (x Axis) on vocabulary size at 20 months (y Axis).

General Discussion

In the two studies reported here, we investigated 7-month-old infants' neural tracking of and rhythmic movement to live and dynamic maternal ID singing. In addition, we tested whether and how these forms of coordination are related to children's language development. We found that 7-month-old infants coordinate both at the neural and behavioral levels with their mothers' live ID singing. In line with our hypotheses, infants showed better neural tracking of lullabies than playsongs, while they displayed more rhythmic movements during playsongs than during lullabies. Interestingly, only infants' neural and behavioral coordination with their mothers' playsongs, but not lullabies, predicted their vocabulary size at 20 months.

Infants showed above-threshold neural tracking of live-sung ID songs in comparison to a shuffled control. This finding highlights that infants are able to neurally track live musical communication beyond rhythmic pure tones (Cirelli et al., 2016), recordings of nursery rhymes (Attaheri et al., 2022), or infant-directed speech (Menn, Michel, et al., 2022). Interestingly, our results indicate that infants were more accurate at tracking lullabies than playsongs but showed fewer rhythmic movements during lullabies than playsongs. Lullabies are sung with a regular and repetitive pattern (Cirelli et al., 2020; Trainor et al., 1997) that

could help infants neurally track this type of song. This interpretation is supported by acoustic analyses of maternal singing in both studies showing that lullabies, compared to playsongs, were sung slower, lower in pitch, and less variable, possibly making them more predictable. These individual acoustic features of lullabies were further related to enhanced neural tracking, highlighting the relation between a clear structure in tempo and pitch and potential ease of neural coordination. This increased clarity in tempo- and pitch-related structures in lullabies might also be particularly associated with their regulatory functions, such as modulating emotions (Trehub et al., 2015) and maintaining infant attention and composure (Shenfield et al., 2003). There is some evidence showing that infants are more relaxed when they listen to lullabies compared to other types of music (Bainbridge et al., 2021; Cirelli et al., 2020; Cirelli & Trehub, 2020), corroborating our finding of infants' decreased rhythmic movements when compared to playsongs. Overall, infants showed higher neural but lower behavioral coordination with lullabies, in line with the soothing function associated with the acoustic features of this type of ID song.

Despite their more variable and faster structure, infants were also able to neurally track playsongs and actually showed more rhythmic movement during playsongs than lullabies. In line with our argumentation, neural coordination was enhanced when caregivers sang with higher pulse clarity, quieter, and less variability in loudness. We, therefore, suggest that caregivers might counterbalance the complexity of playsongs by enhancing their rhythmic structure (Zentner & Eerola, 2010) as well as using loudness to guide infants' attention (Calignano et al., 2021). In previous research, pulse clarity was also related to infants' amounts of rhythmic movement (Zentner & Eerola, 2010). Generally, our results suggest that variability in acoustic features, potentially up to a certain threshold, does not hinder infants' neural tracking of ID songs. Instead, increased rhythmic variability might increase engagement (Cameron et al., 2022), as indicated by more frequent rhythmic movement. The discrepancy in rhythmic movement frequency between playsongs and lullabies is likely further pronounced by infants' familiarity with the songs (see Supplements) and potentially their expectation to engage in an activating social context. Alternatively, rhythmic movement might arise to help infants process these acoustically more variable ID songs. In fact, the perception of rhythm while listening to music often involves spontaneous movement on these periodicities (Hurley et al., 2014; Toiviainen et al., 2010). Music stimulates movement and activates motor pathways (Grahn & Brett, 2007; Vuilleumier &

Trost, 2015), and already 5-month-olds move rhythmically significantly more to music than speech, suggesting that there may be a predisposition for rhythmic movement in response to music (Zentner & Eerola, 2010). Interestingly, passive body movement was also found to influence 7-month-old infants' encoding of musical rhythms (Phillips-Silver & Trainor, 2005), indicating that their own motor experiences may contribute to young infants' processing of more variable acoustic input.

Importantly, we found that infants' diverging coordination to live and dynamic lullaby and playsong was also reflected in the association with language outcomes at 20 months. Results showed that infants' neural tracking of and rhythmic movement to playsongs, but not lullabies, were positively related to their later vocabulary size, even though the relation was not robust. This finding supports the proposition that musicality evolved as a common ability for infants to decipher music and language (Brandt et al., 2012). Preverbal infants, who are not yet familiar with the rules of language, classify speech according to pitch, melody, and rhythm in order to understand its meaning (Goswami, 2012). It is precisely prosodic information (i.e., slow speech rhythms) that is crucial for language acquisition (Goswami, 2012; Langus et al., 2017). Neural tracking of the more variable playsong might, for example, relate to infants' ability to extract prosodic information from a vocal stream and thus possibly support their word segmentation abilities (Menn, Ward, et al., 2022), marked by larger expressive vocabulary size at age 20 months. ID singing may be particularly helpful for language acquisition, in addition to ID speech, due to its metric structure that makes prosodic stress more pronounced (Nakata & Trehub, 2011; Suppanen et al., 2019; Trainor et al., 1997). In addition, moving rhythmically could help infants process the prosodic information from the song, a notion supported by arguments that rhythmic abilities are a central part of language acquisition (Thomson & Goswami, 2008). An alternative explanation we cannot rule out is that infants' movements affected the EEG response; therefore, movement-related artifacts in the EEG might have driven the relation between infant neural tracking in the playsong and infants' later language development. Nonetheless, our study provides the first promising evidence for the connection between early musical experiences and language development.

Despite the study's interactional and naturalistic setting, the paradigm was still an observational situation in a laboratory and thus has its limitations. Both mothers and infants were restricted in their movement and choice of song. We further need to consider the fit of

the musical piece to the particular situation in which we observed infants. The context of the laboratory setting likely afforded the playsong more than the lullaby, as the mothers and the experimenter intended to keep infants attentive and entertained. Thus, the congruence between the observational situation and the playsong may have contributed to our pattern of results. Moreover, it is possible that mothers themselves moved differently during the two types of songs, and infants simply tracked and imitated their mothers' movement type and frequency. Playsongs, in particular, are not only expressed through voice but also the body and are often intrinsically multimodal (Eckerdal & Merker, 2009; Lense et al., 2022). In fact, our supplementary analyses showed that infants looked significantly longer at their mothers during the playsong than during the lullaby. However, infant gaze behavior during the playsong was neither associated with their neural tracking nor rhythmic movements in that condition. Instead, we found that neural tracking of lullabies was weaker when infants looked away from mothers' faces and bodies, suggesting that neural tracking is related to infants' attention towards the face (Lense et al., 2022). To disentangle factors underlying infants' neural tracking during ID songs, future studies thus need to include more in-depth analyses of the accompanying sensorimotor synchronization processes, including caregivers' movements (e.g., rocking, playful gestures, etc.).

Conclusion

Singing represents a flexible tool for caregivers to meet the various needs of their infants. However, it remains a question of great theoretical controversy why ID singing is such a powerful communication method with very young infants (Mehr et al., 2021; Savage et al., 2021). Markova et al. (2019) argued that ID singing has inherent acoustic characteristics that support infants' coordination to a particular interactive rhythm purported by the caregiver, and this process could have important developmental implications (Markova et al., 2020). Findings of the present study corroborate these hypotheses by showing that lullabies may be used to help infants track a slower and regular rhythmic pattern, while playsongs, containing more variability, could be perceived as more engaging and thus make infants move rhythmically. The acoustic variability of playsongs also seems to promote language acquisition in that rhythmic variability may draw attention to the songs' prosodic information (Hannon & Johnson, 2005). These findings highlight potential avenues for future studies to test the neural and behavioral basis of the communicative function of mother-infant musical interaction by experimentally manipulating the relevant acoustic features. With its

semi-naturalistic design, the present study provides a better understanding of the dynamics, mechanisms, and developmental outcomes of early musical interactions.

Acknowledgements

Declaration of Interest. We declare we have no competing interests.

Funding. This research was funded by the University of Vienna and by a doctoral stipend from the Studienstiftung des Deutschen Volkes granted to T.N.

Acknowledgments. We are grateful to Liesbeth Forsthuber, Christina Schaetz, Martina de Eccher, Sarah Paul, Marlies Schermann, Anna Matyk, Rahel Loos, Jakob Weickmann and Elisa Roberti, for their support in data acquisition, processing, and video coding. Additionally, we thank all families who participated in the study and the Department of Obstetrics and Gynecology of the Vienna General Hospital for supporting our participant recruitment.

CRediT author statement. *Trinh Nguyen*: Conceptualization, Methodology, Formal analysis, Data curation, Writing – Original Draft, Writing – Review & Editing, Visualization, Project administration *Susanne Reisner*: Data curation, Writing – Original Draft, Writing – Review & Editing, Visualization *Anja Lueger*: Investigation, Data – Curation, Writing – Original Draft *Sam V. Wass*: Methodology, Formal analysis, Writing – Review & Editing *Stefanie Hoehl:* Conceptualization, Methodology, Resources, Writing – Review & Editing, Supervision, Funding acquisition *Gabriela Markova:* Conceptualization, Methodology, Investigation, Data curation, Writing – Original Draft, Writing – Review & Editing, Supervision, Project administration

References

Attaheri, A., Choisdealbha, Á. N., Di Liberto, G. M., Rocha, S., Brusini, P., Mead, N., Olawole-Scott, H., Boutris, P., Gibbon, S., Williams, I., Grey, C., Flanagan, S., & Goswami, U. (2022). Delta- and theta-band cortical tracking and phase-amplitude coupling to sung speech by infants. *NeuroImage*, *247*, 118698. https://doi.org/10.1016/j.neuroimage.2021.118698

Bainbridge, C. M., Bertolo, M., Youngers, J., Atwood, S., Yurdum, L., Simson, J., Lopez, K., Xing, F., Martin, A., & Mehr, S. A. (2021). Infants relax in response to unfamiliar foreign lullabies. *Nature Human Behaviour*, *5*(2), Article 2. https://doi.org/10.1038/s41562-020-00963-z

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Brandt, A., Slevc, R., & Gebrian, M. (2012). Music and Early Language Acquisition. *Frontiers in Psychology*, *3*. https://www.frontiersin.org/articles/10.3389/fpsyg.2012.00327

Calignano, G., Dispaldro, M., Russo, S., & Valenza, E. (2021). Attentional engagement during syllable discrimination: The role of salient prosodic cues in 6- to 8-month-old infants. *Infant Behavior and Development*, *62*, 101504. https://doi.org/10.1016/j.infbeh.2020.101504

Cameron, D. J., Caldarone, N., Psaris, M., Carrillo, C., & Trainor, L. J. (2022). The complexity-aesthetics relationship for musical rhythm is more fixed than flexible: Evidence from children and expert dancers. *Developmental Science*, e13360. https://doi.org/10.1111/desc.13360

Cirelli, L. K., Jurewicz, Z. B., & Trehub, S. E. (2020). Effects of Maternal Singing Style on Mother–Infant Arousal and Behavior. *Journal of Cognitive Neuroscience*, *32*(7), 1213–1220. https://doi.org/10.1162/jocn_a_01402

Cirelli, L. K., Spinelli, C., Nozaradan, S., & Trainor, L. J. (2016). Measuring Neural Entrainment to Beat and Meter in Infants: Effects of Music Background. *Frontiers in Neuroscience*, *10*. https://doi.org/10.3389/fnins.2016.00229

Cirelli, L. K., & Trehub, S. E. (2020). Familiar songs reduce infant distress. *Developmental Psychology*, *56*(5), 861–868. https://doi.org/10.1037/dev0000917

Crosse, M. J., Di Liberto, G. M., Bednar, A., & Lalor, E. C. (2016). The Multivariate Temporal Response Function (mTRF) Toolbox: A MATLAB Toolbox for Relating Neural Signals to Continuous Stimuli. *Frontiers in Human Neuroscience*, *10*. https://www.frontiersin.org/articles/10.3389/fnhum.2016.00604

de Cheveigné, A. (2020). ZapLine: A simple and effective method to remove power line artifacts. *NeuroImage*, *207*, 116356. https://doi.org/10.1016/j.neuroimage.2019.116356

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Di Liberto, G. M., Pelofi, C., Bianco, R., Patel, P., Mehta, A. D., Herrero, J. L., de Cheveigné, A., Shamma, S., & Mesgarani, N. (2020). Cortical encoding of melodic expectations in human temporal cortex. *ELife*, *9*, e51784. https://doi.org/10.7554/eLife.51784

Eckerdal, P., & Merker, B. (2009). "Music" and the "action song" in infant development: An interpretation. In *Communicative musicality: Exploring the basis of human companionship* (pp. 241–262). Oxford University Press.

Ferland, M. B., & Mendelson, M. J. (1989). Infants' categorization of melodic contour. *Infant Behavior and Development*, *12*(3), 341–355. https://doi.org/10.1016/0163-6383(89)90043-X

Fiedler, L., Wöstmann, M., Graversen, C., Brandmeyer, A., Lunner, T., & Obleser, J. (2017). Single-channel in-ear-EEG detects the focus of auditory attention to concurrent tone streams and mixed speech. *Journal of Neural Engineering*, *14*(3), 036020. https://doi.org/10.1088/1741-2552/aa66dd

François, C., Teixidó, M., Takerkart, S., Agut, T., Bosch, L., & Rodriguez-Fornells, A. (2017). Enhanced Neonatal Brain Responses To Sung Streams Predict Vocabulary Outcomes By Age 18 Months. *Scientific Reports*, *7*(1), 12451. https://doi.org/10.1038/s41598-017-12798-2

Goswami, U. (2012). *Entraining the Brain: Applications to Language Research and Links to Musical Entrainment*. https://doi.org/10.18061/1811/52980

Grahn, J. A., & Brett, M. (2007). Rhythm and Beat Perception in Motor Areas of the Brain. *Journal of Cognitive Neuroscience*, *19*(5), 893–906. https://doi.org/10.1162/jocn.2007.19.5.893

Háden, G. P., Honing, H., Török, M., & Winkler, I. (2015). Detecting the temporal structure of sound sequences in newborn infants. *International Journal of Psychophysiology*, *96*(1), 23–28. https://doi.org/10.1016/j.ijpsycho.2015.02.024

Hannon, E. E., & Johnson, S. P. (2005). Infants use meter to categorize rhythms and melodies: Implications for musical structure learning. *Cognitive Psychology*, *50*(4), 354–377. https://doi.org/10.1016/j.cogpsych.2004.09.003

Haufe, S., Meinecke, F., Görgen, K., Dähne, S., Haynes, J.-D., Blankertz, B., & Bießmann, F. (2014). On the interpretation of weight vectors of linear models in multivariate neuroimaging. *NeuroImage*, *87*, 96–110. https://doi.org/10.1016/j.neuroimage.2013.10.067

Hurley, B. K., Martens, P. A., & Janata, P. (2014). Spontaneous sensorimotor coupling with multipart music. *Journal of Experimental Psychology: Human Perception and Performance*, *40*, 1679–1696. https://doi.org/10.1037/a0037154

Ilari, B. (2015). Rhythmic Engagement With Music in Early Childhood: A Replication and Extension. *Journal of Research in Music Education*, *62*(4), 332–343. https://doi.org/10.1177/0022429414555984

Iverson, J. M. (2010). Developing language in a developing body: The relationship between motor development and language development. *Journal of Child Language*, *37*(2), 229. https://doi.org/10.1017/S0305000909990432

Iverson, J. M., Hall, A. J., Nickel, L., & Wozniak, R. H. (2007). The Relationship between Reduplicated Babble Onset and Laterality Biases in Infant Rhythmic Arm Movements. *Brain and Language*, *101*(3), 198. https://doi.org/10.1016/j.bandl.2006.11.004

Jessen, S., Fiedler, L., Münte, T. F., & Obleser, J. (2019). Quantifying the individual auditory and visual brain response in 7-month-old infants watching a brief cartoon movie. *NeuroImage*, *202*, 116060. https://doi.org/10.1016/j.neuroimage.2019.116060

Jessen, S., Obleser, J., & Tune, S. (2021). Neural tracking in infants – An analytical tool for multisensory social processing in development. *Developmental Cognitive Neuroscience*, *52*, 101034. https://doi.org/10.1016/j.dcn.2021.101034

Kisilevsky, B. s., Hains, S. m. j., Jacquet, A.-Y., Granier-Deferre, C., & Lecanuet, J. p. (2004). Maturation of fetal responses to music. *Developmental Science*, *7*(5), 550–559. https://doi.org/10.1111/j.1467-7687.2004.00379.x

Kragness, H. E., Anderson, L., Chow, E., Schmuckler, M., & Cirelli, L. K. (2022). Musical groove shapes children's free dancing. *Developmental Science*. https://doi.org/10.1111/desc.13249

Krumhansl, C. L., & Jusczyk, P. W. (1990). Infants' Perception of Phrase Structure in Music. *Psychological Science*, *1*(1), 70–73. https://doi.org/10.1111/j.1467-9280.1990.tb00070.x

Langus, A., Mehler, J., & Nespor, M. (2017). Rhythm in language acquisition. *Neuroscience & Biobehavioral Reviews*, *81*, 158–166. https://doi.org/10.1016/j.neubiorev.2016.12.012

Lartillot, O., Toiviainen, P., & Eerola, T. (2008). A Matlab Toolbox for Music Information Retrieval. In C. Preisach, H. Burkhardt, L. Schmidt-Thieme, & R. Decker (Eds.), *Data Analysis, Machine Learning and Applications* (pp. 261–268). Springer. https://doi.org/10.1007/978-3-540-78246-9_31

Lebedeva, G. C., & Kuhl, P. K. (2010). Sing that tune: Infants' perception of melody and lyrics and the facilitation of phonetic recognition in songs. *Infant Behavior & Development*, *33*, 419–430. https://doi.org/10.1016/j.infbeh.2010.04.006

Lenc, T., Peter, V., Hooper, C., Keller, P. E., Burnham, D., & Nozaradan, S. (2022). Infants show enhanced neural responses to musical meter frequencies beyond low-level features. *Developmental Science*, *n/a*(n/a), e13353. https://doi.org/10.1111/desc.13353

Lense, M. D., Shultz, S., Astésano, C., & Jones, W. (2022). Music of infant-directed singing entrains infants' social visual behavior. *Proceedings of the National Academy of Sciences*, *119*(45), e2116967119. https://doi.org/10.1073/pnas.2116967119

Lopez, K. L., Monachino, A. D., Morales, S., Leach, S. C., Bowers, M. E., & Gabard-Durnam, L. J. (2022). HAPPILEE: HAPPE In Low Electrode Electroencephalography, a standardized pre-processing software for lower density recordings. *NeuroImage*, *260*, 119390. https://doi.org/10.1016/j.neuroimage.2022.119390

Markova, G., & Legerstee, M. (2006). Contingency, imitation, and affect sharing: Foundations of infants' social awareness. *Developmental Psychology*, *42*(1), 132–141. https://doi.org/10.1037/0012-1649.42.1.132

Markova, G., Nguyen, T., & Hoehl, S. (2019). Neurobehavioral Interpersonal Synchrony in Early Development: The Role of Interactional Rhythms. *Frontiers in Psychology*, *10*, 2078. https://doi.org/10.3389/fpsyg.2019.02078

Markova, G., Nguyen, T., Schätz, C., & de Eccher, M. (2020). Singing in Tune – Being in Tune: Relationship Between Maternal Playful Singing and Interpersonal Synchrony. *Enfance*, *1*(1), 89–107. Cairn.info. https://doi.org/10.3917/enf2.201.0089

Marschik, P. B., Einspieler, C., Garzarolli, B., & Prechtl, H. F. R. (2007). Events at early development: Are they associated with early word production and neurodevelopmental abilities at the preschool age? *Early Human Development*, *83*(2), 107–114. https://doi.org/10.1016/j.earlhumdev.2006.05.009

Mehr, S. A., Krasnow, M. M., Bryant, G. A., & Hagen, E. H. (2021). Origins of music in credible signaling. *Behavioral and Brain Sciences*, *44*, e60. https://doi.org/10.1017/S0140525X20000345

Menn, K. H., Michel, C., Meyer, L., Hoehl, S., & Männel, C. (2022). Natural infant-directed speech facilitates neural tracking of prosody. *NeuroImage*, *251*, 118991. https://doi.org/10.1016/j.neuroimage.2022.118991

Menn, K. H., Ward, E. K., Braukmann, R., van den Boomen, C., Buitelaar, J., Hunnius, S., & Snijders, T. M. (2022). Neural Tracking in Infancy Predicts Language Development in Children With and Without Family History of Autism. *Neurobiology of Language*, *3*(3), 495–514. https://doi.org/10.1162/nol_a_00074

Meyer, M., Lamers, D., Kayhan, E., Hunnius, S., & Oostenveld, R. (2021). Enhancing reproducibility in developmental EEG research: BIDS, cluster-based permutation tests, and effect sizes. *Developmental Cognitive Neuroscience*, *52*, 101036. https://doi.org/10.1016/j.dcn.2021.101036

Nakata, T., & Trehub, S. E. (2011). Expressive timing and dynamics in infant-directed and non-infant-directed singing. *Psychomusicology: Music, Mind and Brain*, *21*(1–2), 45–53. https://doi.org/10.1037/h0094003

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2010). FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational Intelligence and Neuroscience*, *2011*, e156869. https://doi.org/10.1155/2011/156869

Papadimitriou, A., Smyth, C., Politimou, N., Franco, F., & Stewart, L. (2021). The impact of the home musical environment on infants' language development. *Infant Behavior and Development*, *65*, 101651. https://doi.org/10.1016/j.infbeh.2021.101651

Phillips-Silver, J., & Trainor, L. J. (2005). Feeling the Beat: Movement Influences Infant Rhythm Perception. *Science*, *308*(5727), 1430–1430. https://doi.org/10.1126/science.1110922

Politimou, N., Dalla Bella, S., Farrugia, N., & Franco, F. (2019). Born to Speak and Sing: Musical Predictors of Language Development in Pre-schoolers. *Frontiers in Psychology*, *10*. https://doi.org/10.3389/fpsyg.2019.00948

Putkinen, V., Saarikivi, K., & Tervaniemi, M. (2013). Do informal musical activities shape auditory skill development in preschool-age children? *Frontiers in Psychology*, *4*. https://www.frontiersin.org/articles/10.3389/fpsyg.2013.00572

Rock, A. M. L., Trainor, L. J., & Addison, T. L. (1999). Distinctive messages in infant-directed lullabies and play songs. *Developmental Psychology*, *35*(2), 527–534. https://doi.org/10.1037/0012-1649.35.2.527

Savage, P. E., Loui, P., Tarr, B., Schachner, A., Glowacki, L., Mithen, S., & Fitch, W. T. (2021). Music as a coevolved system for social bonding. *Behavioral and Brain Sciences*, *44*, e59. https://doi.org/10.1017/S0140525X20000333

Shenfield, T., Trehub, S. E., & Nakata, T. (2003). Maternal Singing Modulates Infant Arousal. *Psychology of Music*, *31*(4), 365–375. https://doi.org/10.1177/03057356030314002

Steinberg, S., Liu, T., & Lense, M. D. (2021). Musical Engagement and Parent-Child Attachment in Families With Young Children During the Covid-19 Pandemic. *Frontiers in Psychology*, *12*, 641733. https://doi.org/10.3389/fpsyg.2021.641733

Suppanen, E., Huotilainen, M., & Ylinen, S. (2019). Rhythmic structure facilitates learning from auditory input in newborn infants. *Infant Behavior and Development*, *57*, 101346. https://doi.org/10.1016/j.infbeh.2019.101346

Thelen, E. (1979). Rhythmical stereotypies in normal human infants. *Animal Behaviour*, *27*, 699–715. https://doi.org/10.1016/0003-3472(79)90006-X

Thiessen, E. D., & Saffran, J. R. (2009). How the Melody Facilitates the Message and Vice Versa in Infant Learning and Memory. *Annals of the New York Academy of Sciences*, *1169*(1), 225–233. https://doi.org/10.1111/j.1749-6632.2009.04547.x

Thomson, J. M., & Goswami, U. (2008). Rhythmic processing in children with developmental dyslexia: Auditory and motor rhythms link to reading and spelling. *Journal of Physiology-Paris*, *102*(1), 120–129. https://doi.org/10.1016/j.jphysparis.2008.03.007

Toiviainen, P., Luck, G., & Thompson, M. R. (2010). Embodied Meter: Hierarchical Eigenmodes in Music-Induced Movement. *Music Perception*, *28*(1), 59–70. https://doi.org/10.1525/mp.2010.28.1.59

Trainor, L. J., Clark, E. D., Huntley, A., & Adams, B. A. (1997). The acoustic basis of preferences for infant-directed singing. *Infant Behavior and Development*, *20*(3), 383–396. https://doi.org/10.1016/S0163-6383(97)90009-6

Trainor, L. J., & Heinmiller, B. M. (1998). The development of evaluative responses to music: *Infant Behavior and Development*, *21*(1), 77–88. https://doi.org/10.1016/S0163-6383(98)90055-8

Trehub, S. E. (2017). The Maternal Voice as a Special Signal for Infants. In M. Filippa, P. Kuhn, & B. Westrup (Eds.), *Early Vocal Contact and Preterm Infant Brain Development: Bridging the Gaps Between Research and Practice* (pp. 39–54). Springer International Publishing. https://doi.org/10.1007/978-3-319-65077-7_3

Trehub, S. E. (2019). Nurturing infants with music. *International Journal of Music in Early Childhood*, *14*(1), 9–15. https://doi.org/10.1386/ijmec.14.1.9_1

Trehub, S. E., Ghazban, N., & Corbeil, M. (2015). Musical affect regulation in infancy. *Annals of the New York Academy of Sciences*, *1337*(1), 186–192. https://doi.org/10.1111/nyas.12622

Trehub, S. E., Plantinga, J., & Russo, F. A. (2016). Maternal Vocal Interactions with Infants: Reciprocal Visual Influences: Maternal Speech and Singing. *Social Development*, *25*(3), 665–683. https://doi.org/10.1111/sode.12164

Trehub, S., & Trainor, L. (1998). Singing to infants: Lullabies and play songs. *Advances in Infancy Research*, *12*, 43–77.

Vuilleumier, P., & Trost, W. (2015). Music and emotions: From enchantment to entrainment. *Annals of the New York Academy of Sciences*, *1337*(1), 212–222. https://doi.org/10.1111/nyas.12676

Weineck, K., Wen, O. X., & Henry, M. J. (2022). Neural synchronization is strongest to the spectral flux of slow music and depends on familiarity and beat salience. *ELife*, *11*, e75515. https://doi.org/10.7554/eLife.75515

Williams, K. E., Barrett, M. S., Welch, G. F., Abad, V., & Broughton, M. (2015). Associations between early shared music activities in the home and later child outcomes: Findings from the Longitudinal Study of Australian Children. *Early Childhood Research Quarterly*, *31*, 113–124. https://doi.org/10.1016/j.ecresq.2015.01.004

Winkler, I., Háden, G. P., Ladinig, O., Sziller, I., & Honing, H. (2009). Newborn infants detect the beat in music. *Proceedings of the National Academy of Sciences*, *106*(7), 2468–2471. https://doi.org/10.1073/pnas.0809035106

Zentner, M. R., & Eerola, T. (2010). Rhythmic engagement with music in infancy. *Proceedings of the National Academy of Sciences*, *107*(13), 5768–5773. https://doi.org/10.1073/pnas.1000121107

Zhao, T. C., & Kuhl, P. K. (2016). Musical intervention enhances infants' neural processing of temporal structure in music and speech. *Proceedings of the National Academy of Sciences*, *113*(19), 5212–5217. https://doi.org/10.1073/pnas.1603984113