1

The acetylase activity of Cdu1 protects *Chlamydia* effectors from degradation to regulate

bacterial exit from infected cells

Robert J. Bastidas[1][*], Mateusz Kędzior[1], Lee Dolat[1], Barbara S. Sixt[2,3,4], Jonathan N. Pruneda[5],

Raphael H. Valdivia[1,6][*]

Author affiliations: [1]Department of Immunology, Duke University, Durham, N.C 27708, USA.

[2]Deparment of Molecular Biology, Umeå University, Umeå, Sweden. [3]The Laboratory for

Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå, Sweden. [4]Umeå Centre

for Microbial Research (UCMR), Umeå University, Umeå, Sweden. [5]Department of Molecular

Microbiology & Immunology, Oregon Health & Science University, Portland, OR 97239, USA.

[6]Department of Molecular Genetics and Microbiology, Duke University, Durham, N.C 27708,

USA.

[*] Correspondence: robert.bastidas@duke.edu; raphael.valdivia@duke.edu

Abstract

Many cellular processes are regulated by ubiquitin-mediated proteasomal degradation. Bacterial pathogens can regulate eukaryotic proteolysis through the delivery of proteins with de-ubiquitinating (DUB) activities. The obligate intracellular pathogen *Chlamydia trachomatis* secretes Cdu1 (ChlaDUB1), a dual deubiquitinase and Lys-acetyltransferase, that promotes Golgi remodeling and survival of infected host cells presumably by regulating the ubiquitination of host and bacterial proteins. Here we determined that Cdu1's acetylase but not its DUB activity is important to protect Cdu1 from ubiquitin-mediated degradation. We further identified three *C. trachomatis* proteins on the pathogen-containing vacuole (InaC, IpaM, and CTL0480) that required Cdu1's acetylase activity for protection from degradation and determined that Cdu1 and these Cdu1-protected proteins are required for optimal egress of *Chlamydia* from host cells. These findings highlight a non-canonical mechanism of pathogen-mediated protection of virulence factors from degradation after their delivery into host cells and the coordinated regulation of secreted effector proteins.

Introduction

Ubiquitination is a conserved and ubiquitous post-translational modification (PTM) of proteins involving the conjugation of the carboxy-terminal glycine residue of ubiquitin (Ub) to lysine residues of target proteins. Poly-ubiquitination of substrates involves further conjugation of a Ub internal lysine residue or amino-terminal methionine (M1) with a second Ub molecule. Seven internal lysines in Ub (K6, K11, K27, K29, K33, K48, K63) and M1 are utilized by Ub conjugating enzymes to form homogeneous, branched, or mixed poly-ubiquitin (polyUb) chains (Komander and Rape., 2012). PolyUb chains with different linkage types exhibit distinct structures and functions. For example, K48- and K11-linked polyUb chains exhibit a compact conformation and are substrates for 26S proteasome-mediated degradation (Varadan et al., 2004; Tenno et al., 2004; Eddins et al., 2007; Bremm et al., 2010; Saeki., 2017). In contrast, K63-linked polyUb

conjugates adopt more open conformations that enable the recruitment of multiprotein

complexes that regulate the function of the target protein by proteolytic independent events

(Komander et al., 2009b; Weeks et al., 2009; Datta et al., 2009; Komander and Rape., 2012).

Mixed and branched polyUb chains are also emerging as important regulators of physiological

functions (Swatek and Komander., 2016; Ohtake and Tsuchiya., 2017).

Protein ubiquitination regulates numerous eukaryotic cell processes including protein

degradation, signal transduction, cell cycle regulation, selective autophagy, the DNA damage

response, and programmed cell death. Ub also plays key roles in modulating host innate

immune responses to bacterial infection (Li et al., 2016), bacterial proteins, pathogen-containing

vacuoles, and bacteria themselves by targeting them for Ub-mediated degradation by

proteasomal or autophagic machineries (Li et al., 2016). Because the Ub system is critical for

pathogen containment, many pathogens have evolved mechanisms to counteract the impact of

this PTM (Vozandychova et al., 2021). For instance, bacterial deubiquitinases (DUBs) can

remove Ub from ubiquitinated substrates thereby dampening inflammatory and cell-autonomous

defense mechanisms (Kubori et al., 2019). Many DUBs are cysteine proteases with a catalytic

Cys, a nearby His and an Asn/Asp (Komander et al., 2009a). DUBs are typically dedicated to

the removal of Ub moieties and are unable to hydrolyze other Ub-like (Ubl) modifications such

as SUMO or NEDD8. However, the CE clan of Ubl proteases (ULPs) can catalyze the removal

of both SUMO and NEDD8 (Ronau et al., 2016). Bacterial pathogens also encode CE clan

enzymes that function as DUBs, ULPs or both. For instance, *Salmonella* Typhimurium SseL*,*

Escherichia coli ElaD, and *Shigella flexneri* ShiCE function as Ub specific proteases (Rytkönen

et al., 2007; Catic et al., 2007; Pruneda et al., 2016) while RickCE from *Rickettsia belli* functions

as a protease directed towards both Ub and NEDD8 as does SidE from *Legionella pneumophila*

which displays mixed activities towards Ub, NEDD8, and ISG15 (Sheedlo et al., 2015; Pruneda

et al., 2016). Similarly, XopD from *Xanthamonas campestris* and LotB from *L. pneumophila* are

isopeptidases exhibiting cross reactivity towards both Ub and SUMO (Pruneda et al., 2016;

Schubert et al., 2020). Some CE clan bacterial effectors display acetyltransferase activity. *L.*

pneumophila LegCE, *S.* Typhimurium AvrA, and YopJ from *Yersinia pestis* function exclusively

as acetyltransferases (Mittal et al., 2006; Mukherjee et al., 2006; Jones et al., 2008; Pruneda et

al., 2016). In contrast, the *Chlamydia trachomatis* (Ct) effector Cdu1/*Chla*DUB1 is a CE clan

protein that exhibits both acetyltransferase and deubiquitinating activities (Misaghi et al., 2006;

Pruneda et al., 2016; Fischer et al., 2017; Pruneda et al., 2018).

Ct is an obligate intracellular bacterial pathogen responsible for human diseases of significant

clinical and public health importance (Haggerty et al., 2010). Ct has a biphasic developmental

cycle in which the Ct infectious propagule or elementary body (EB) invades the target host cell.

Upon internalization the EB transitions to the reticulate body (RB). RBs replicate by binary

fission within a pathogenic vacuole ("inclusion") and asynchronously differentiate back to EBs.

In cell culture, starting at around 48 hours-post infection (hpi) Ct will egress after lysis of the

host cell or by a process termed extrusion, wherein the entire inclusion is exocytosed from the

infected cell (Moulder., 1991; Abdelrahman and Belland., 2005; Hybiske and Stephens., 2007;

Lee et al., 2018). The effector Cdu1 was originally identified as a deneddylating and

deubiquitinating enzyme and subsequently shown to exhibit *in-vitro* isopeptidase activity

towards both Lys48 and Lys63 linked di-Ub substrates (Misaghi et al., 2006; Claessen et al.,

2013; Pruneda et al., 2016; Fischer et al., 2017). Cdu1 is unique among CE clan enzymes in

that it also functions as a bona fide lysine acetylase with both is acetylase and DUB activities

catalyzed by the same catalytic active site (Pruneda et al., 2018). Intriguingly, Cdu1

autoacetylation is directed towards lysines unlike other CE clan acetylases that predominantly

target serine and threonine residues (Pruneda et al., 2018). In infected cells, Cdu1 localizes to

the inclusion membrane where it functions to stabilize the anti-apoptotic protein Mcl-1 and to

promote the repositioning of Golgi ministacks around the Ct inclusion (Fischer et al., 2017;

Wang et al., 2018; Pruneda et al., 2018; Kunz et al., 2019; Auer et al., 2020). However, the

mechanism by which Cdu1 promotes redeployment of Golgi ministacks and any additional roles

that Cdu1 may play during Ct infection of epithelial cells remains unknown.

In this study, we show that Cdu1 protects itself and three secreted Ct effectors, InaC, IpaM, and

CTL0480 from targeted ubiquitination and proteasomal degradation. InaC, IpaM, and CTL0480

are members of a larger family of bacterial proteins embedded within the inclusion membrane

(Inc proteins) (Bannantine et al., 2000; Rockey et al., 2002; Chen et al., 2006; Li et al., 2008;

Alzhanov et al., 2009; Dehoux et al., 2011; Lutter et al., 2012; Lutter et al., 2013; Kokes et al.,

2015; Weber et al., 2015). We show that Cdu1-mediated protection from degradation is

independent from its DUB activity but relies upon its Lys acetylase activity. We show that Cdu1

protects InaC to promote repositioning of Golgi ministacks and formation of actin scaffolds

around the Ct inclusion, and CTL0480 to promote recruitment of myosin phosphatase target

subunit 1 (MYPT1) to the inclusion. In addition, we determined that Cdu1 and Cdu1-protected

Incs are required for optimal extrusion of inclusions from host cells at the late stages of

infection.

Results

The *C. trachomatis* inclusion membrane proteins InaC, IpaM, and CTL0480 are

differentially ubiquitinated in the absence of Cdu1

Cdu1 is required for Golgi repositioning around the Ct inclusion (Pruneda et al., 2018; Auer et

al., 2020). To understand how Cdu1 promotes Golgi redistribution, we first generated a *cdu1*

null strain in a Ct L2 (LGV L2 434 Bu) background by TargeTron mediated insertional

mutagenesis (pDFTT3-*aadA*) (Lowden et al., 2015) (S. Figure 1). Loss of Cdu1 expression in

the resulting L2 *cdu1*::GII *aadA* (*cdu1*::GII) strain was verified by western blot analysis and by

indirect immunofluorescence with antibodies raised against Cdu1 (S. Figures 2A and 2B).

Because *cdu2* resides directly downstream of the *cdu1* locus and encodes a second Ct DUB

(Cdu2/*Chla*DUB2) (Misaghi et al., 2006), we first determined whether the disruption of *cdu1*

impacted the expression of *cdu2*. We detected *cdu1* and *cdu2* transcripts in HeLa cells infected

with Ct L2 but not for the juncture between *cdu1* and *cdu2* (S. Figure 2C). In cells infected with

Ct *cdu1*::GII we only detected *cdu2* transcripts (S. Figure 2C) confirming that *cdu1* and *cdu2* are

not co-expressed as part of an operon in accordance with previous observations (Albrecht et al.,

2010).

We hypothesized that Cdu1s' DUB activity promoted Golgi redistribution around inclusions and

that we could identify potential targets by comparing the protein ubiquitination profile of cells

infected with WT or *cdu1*::GII strains by quantitative mass spectrometry (MS). HeLa cell were

mock infected or infected with either WT L2 or *cdu1*::GII strains. At 24 hpi, poly-ubiquitinated

proteins were enriched from lysed cells using Tandem Ubiquitin Binding Entities (TUBEs)

(LifeSensors). TUBEs consist of concatenated Ub binding associated domains (UBAs) that bind

to polyUb-modified proteins with nanomolar affinities. Poly-ubiquitinated proteins of both human

and Ct origin were enriched and identified by quantitative LC-MS/MS analysis.

Over 2,000 non-ubiquitinated proteins co-precipitated with TUBE 1 bound proteins across all

three conditions (mock, L2, and *cdu1*::GII infected HeLa cells) and 3 biological replicates (S.

Table 1). Among these, 47 human proteins were significantly enriched in mock infected HeLa

cells and 50 human proteins were significantly enriched during Ct infection (L2 and *cdu1*::GII)

(S. Table 3, S. Figure 3) Pathway enrichment analysis revealed that proteins involved in RNA

metabolism were overrepresented among co-precipitating proteins from mock infected cells (S.

Table 4, S. Figure 4) while no biological pathways or processes were overrepresented in

proteins enriched from infected cells (S. Figure 4). We also identified 8 TUBE1 co-precipitating

Ct proteins in HeLa cells infected with L2 and *cdu1*::GII (S. Table 5, S. Figure 3).

TUBE 1 affinity capture lead to the identification of 43 ubiquitinated proteins (35 human proteins

and 8 Ct proteins across all 3 conditions and replicates) based on the presence of peptides

containing a di-glycine remnant motif (Peng et al., 2003) (S. Tables 6-8). The lack of widespread

poly-ubiquitination of either human or Ct proteins in response to Ct infection (Figure 1) was

surprising given that wholesale changes in protein ubiquitination has been reported during

infection of HeLa cells by intracellular pathogens like *S.* Typhimurium (Fiskin et al., 2016). Only

two human ubiquitinated proteins (ZC3H7A and DDIT4) were found to be significantly enriched

in response to WT L2 infection (Figures 1A and 1C, S. Table 8) while only one human protein

(MGC3121) was preferentially ubiquitinated in HeLa cells infected with the *cdu1*::GII mutant

strain (Figures 1B and 1C, S. Table 8). In contrast three Ct proteins, InaC (K104, K107, and

K149), IpaM (K29), and CTL0480 (K115) were ubiquitinated at Lys residues in the absence of

Cdu1 (Figures 1B, 1C, and 1D, S. Table 8). InaC, IpaM, and CTL0480 are Ct effector proteins

that localize to the inclusion membrane (Chen et al., 2006; Alzhanov et al., 2009; Lutter et al.,

2013; Kokes et al., 2015). These Type 3 secretion substrates belong to a family of over 36

inclusion membrane proteins that contain a signature bi-lobal hydrophobic transmembrane

domain (Bannantine et al., 2000; Rockey et al., 2002; Li et al., 2008; Dehoux et al., 2011; Lutter

et al., 2012). Because Cdu1 also localizes to the inclusion membrane (Fischer et al., 2017;

Wang et al., 2018; Pruneda et al., 2018; Kunz et al., 2019) we postulated that Cdu1 directly

protects InaC, IpaM, and CTL0480 from ubiquitination.

Cdu1 associates with InaC, IpaM, and CTL0480

We first determined if Cdu1 co-localized with InaC, IpaM, and CTL0480 at the inclusion

membrane. HeLa cells were infected for 24 hours with WT L2 or L2 expressing CTL0480-Flag

from its endogenous promoter and immunostained with antibodies against Cdu1, InaC, IpaM, or

the Flag epitope. Both InaC and CTL0480-Flag localized throughout the inclusion membrane

while IpaM was restricted to discrete microdomains as previously reported (Alzhanov et al.,

2009; Dumoux et al., 2015) (Figure 2A). Cdu1 co-localized with InaC and CTL0480-Flag and

with IpaM at microdomains (Figures 2A and 2B). All four antibodies specifically recognized their

corresponding antigens since immunostaining for Cdu1, InaC, IpaM or the Flag epitope was not

observed in Ct strains lacking Cdu1 (*cdu1*::GII), InaC (M407, Kokes et al., 2015) IpaM

(*ipaM*::GII, Meier et al., 2022), or a strain that does not express CTL0480-Flag (Figure 2A).

We next determined if Cdu1 can interact with InaC, IpaM, and CTL0480 by co-transfecting HEK

293 cells with vectors expressing either full length Cdu1-GFP or truncated versions of Cdu1-

GFP lacking transmembrane or catalytic domains (Figure 2C), and vectors expressing Flag-

InaC, V5-IpaM, and V5-CTL0480. Transfected cells were lysed and Cdu1-GFP was

immunoprecipitated with antibodies against GFP. Western blot analysis of the

immunoprecipitates showed that Flag-InaC, V5-IpaM, and V5-CTL0480 co-precipitated with

Cdu1-GFP (Figures 2D-F). Moreover, the transmembrane domain of Cdu1 was necessary for

Cdu1-GFP to interact with all three Incs (Figures 2D-F). The interaction between Cdu1-GFP and

the three tagged Incs was specific since we did not detect interactions between Cdu1-GFP and

a V5-tagged version of the inclusion membrane protein CpoS (Sixt et al., 2017) (Figure 2G). We

conclude from these results that Cdu1 can interact with all three Incs even in the absence of

infection and that interactions are facilitated by the transmembrane domain of Cdu1.

Cdu1 protects InaC, IpaM, and CTL0480 proteins from degradation during infection

We next assessed if Cdu1 was required to stabilize endogenous InaC, IpaM, and CTL0480 in

infected cells. HeLa cells were infected with either WT L2 or *cdu1*::GII strains and at various

time points in the Ct infectious cycle crude cell lysates were analyzed by western blot to assess

the relative abundance of Inc proteins. At 24 hpi, we observed a decrease in InaC protein levels

in cells infected with *cdu1*::GII relative to cells infected with WT L2 (Figure 3A). In contrast, IpaM

protein levels did not decrease until 36 hpi in *cdu1*::GII infected cells (Figure 3B). For CTL0480,

we were not able to detect CTL0480 by western blot but were successful in following CTL0480

expression by indirect immunofluorescence (Figure 3C). The relative abundance of CTL0480 at

inclusion membranes, like IpaM, was not affected in *cdu1*::GII inclusions at 24 hpi. However, at

36 hpi, a subpopulation of cells lost CTL0480 immunoreactivity and by 48 hpi, inclusion

membranes of the *cdu1*::GII strain were devoid of CTL0480 while CTL0480 was prominently

detected at the inclusion membranes of WT L2 (Figures 3C and 3D). As controls for the

specificity of antibodies used for western blots and for indirect immunofluorescence, we

included cells infected with Ct lacking *ipaM (ipaM*::GII, Meier et al., 2022), with an *inaC*

nonsense mutant (M407, Kokes et al., 2015), and with Ct lacking *ctl0480* (*ctl0480*::GII, Shaw

et al., 2018). Overall, our results indicate that steady state protein levels of InaC and IpaM and

CTL0480 localization at the inclusion membrane, especially at late stages of infection, are

dependent on Cdu1, and that Cdu1 acts at different stages in the infection cycle.

The acetylase activity of Cdu1 is required for Cdu1 to protect itself, InaC, and IpaM from

polyubiquitination and proteasomal degradation

The crystal structures of Cdu1 bound to Ub or Coenzyme A indicated that the adenosine and

phosphate groups of Coenzyme A make contact with a helix in variable region 3 (VR-3) of Cdu1

while the Ile36-patch of Ub binds to the opposite face of the same helix (Pruneda et al., 2018).

Although Cdu1 catalyzes both of its DUB and acetylase (Act) activities with the same active site

(Pruneda et al., 2018) the two activities of Cdu1 can be uncoupled by the amino acid

substitution K268E in VR-3 which disrupts Coenzyme A binding required for Act activity and by

the amino acid substitution I225A in the Ub-binding region of VR-3 required for DUB activity

(Pruneda et al., 2018). These substitutions allowed us to test which of Cdu1's enzymatic

functions are required for the observed effects on protein stability. We generated Ct shuttle

plasmids (pBOMB4-MCI backbone), expressing WT Cdu1, a catalytically inactive variant of

Cdu1 lacking both DUB and Act activities (Cdu1^{C345A}) (Pruneda et al., 2018), a Cdu1 DUB-

deficient variant (Cdu1^{I225A}), and a Cdu1 Act-deficient variant (Cdu1^{K268E}). All Cdu1 constructs

were expressed as 3X Flag epitope-tagged versions and from the Cdu1 endogenous promoter.

Plasmids expressing each Cdu1 variant were transformed into the *cdu*1::GII mutant and the

resulting strains used to infect HeLa cells for 36 and 48 hours. The levels of endogenous InaC

and IpaM in cell extracts of infected cells were monitored by western blot analysis. At 36 hpi,

InaC protein levels drastically decreased in cells infected with *cdu1* null strains transformed with

empty vector or expressing the catalytic inactive variant of Cdu1 (Cdu1^{C345A}-Flag) (Figure 4A).

Likewise, IpaM protein levels diminished at 48 hpi during infection with the same strains (Figure

4B). Both InaC and IpaM protein levels were restored to wild type levels in *cdu1* null strains

complemented with wild type Cdu1-Flag (Figures 4A and 4B). Unexpectedly, cells infected with

a *cdu1* null strain ectopically expressing the DUB deficient Cdu1 variant (Cdu1^{I225A}-Flag)

displayed wild type levels of InaC and IpaM while the Act deficient variant (Cdu1^{K268E}-Flag) did

not (Figures 4A and 4B). These results suggest that the acetylase activity of Cdu1 rather than

its DUB activity is required for Cdu1's ability to stabilize InaC and IpaM proteins.

When we monitored the stability of each Flag-tagged Cdu1 variant, we observed that the

catalytically inactive variant of Cdu1 (C345A-Flag) was destabilized (Figures 4A and 4B- Flag

WB). We reasoned that Cdu1 also protects itself from being targeted for degradation in infected

cells. As with InaC and IpaM, the acetylase but not the DUB activity of Cdu1 was required for

Cdu1's stability (Figures 4A and 4B). Although the C345A (DUB-, Act-) and K268E (Act-) amino

acid substitutions in Cdu1 do not destabilize Cdu1 expressed in *E. coli* (Pruneda et al., 2018), it

was possible that these substitutions impacted the expression and/or folding of Cdu1 in

Chlamydia. To determine if these Cdu1 mutants were inherently unstable, we expressed

Cdu1^{C345A}-Flag and Cdu1^{K268E}-Flag in WT L2 in the presence of endogenous Cdu1. We found

that each Flag-tagged variant was stabilized (Figure 4C), indicating that endogenous Cdu1

protected the catalytically-deficient Cdu1 variants *in trans.* In addition, western blot analysis of

immunoprecipitated Cdu1-Flag and Cdu1^{C345A}-Flag expressed in a *cdu1*::GII mutant showed

that while wild-type Cdu1-Flag was not modified by Lys48-linked poly-ubiquitination, Cdu1^{C345A}-

Flag was robustly modified by Lys48-linked polyUb in the presence of the proteasome inhibitor

MG132 (Figure 4D). Moreover, endogenous Cdu1 protected Cdu1^{C345A}-Flag from Lys48-linked

polyUb when Cdu1^{C345A}-Flag was expressed in a wild type L2 background (Figure 4D). These

results indicate that the loss of Cdu1 activity likely leads to its Lys48-linked poly-ubiquitination

and subsequent proteasome-dependent degradation.

Because Cdu1 autoacetylates itself and its Act activity is directed towards lysines (Pruneda et

al., 2018) we postulated that Cdu1 may stabilize proteins from degradation by acetylating lysine

residues that are potential targets of ubiquitination. We tested this hypothesis by assessing

whether Cdu1, InaC, CTL0480, and IpaM are acetylated at lysines during infection. Fractions

enriched for inclusion membranes were isolated by sub-cellular fractionation from HeLa cells

infected with wild-type L2 (24 hpi), and proteins acetylated at lysines were immunoprecipitated.

Western blot analysis of acetyl-lysine immunoprecipitates indicated that InaC and IpaM, but not

the Inc protein IncA, were acetylated (Figure 4E). Western blot analysis of anti-acetyl-lysine

immunoprecipitates of inclusion membrane-enriched membrane fractions of HeLa cells infected

with L2 expressing Cdu1-Flag (24 hpi), CTL0480-Flag (40 hpi), and IpaM-Flag (40 hpi) also

showed that all three Flag tagged effectors were acetylated at lysines (Figure 4F). We also

determined that Flag-tagged InaC expressed in an M407 (*inaC* null) background was acetylated

at lysines as determined by western blot analysis of anti-acetyl-lysine immunoprecipitates

(Figure 4G). In addition, we identified acetylated forms of Cdu1 from mass spectrometric

analysis of Flag immunoprecipitates derived from extracts of HeLa cells infected with L2

expressing Cdu1-Flag, (24 hpi) (Figure 4H).

We reasoned that the lysine acetylase activity of Cdu1 is a prominent mechanism by which

Cdu1 protects client proteins (at 24 hpi), since loss of Cdu1 or expression of the acetylase

deficient variant of Cdu1 (Cdu1^{K268E}-Flag) leads to a marked increase in Ub immunostaining at

or near the periphery of *cdu1*::GII inclusions (> 80% of inclusions) compared to HeLa cells

infected with *cdu1*::GII strains complemented with wild type or DUB deficient (I225A-Flag) Cdu1

strains (Figures 5A and 5B). Given that Cdu1 appears to localize exclusively at inclusion

membranes we predicted that its activity would be spatially restricted to the inclusion periphery.

We tested this premise by co-infecting HeLa cells with an *incA* null strain (M923 (IncA R197*),

Kokes et al., 2015) and the *cdu1*::GII strain. IncA mediates homotypic fusion of inclusion

membranes and loss of IncA results in the accumulation of multiple unfused inclusions in cells

infected at high MOIs (Hackstadt et al., 1999; Suchland et al., 2000; Pannekoek et al., 2005)

(Figure 5C). As expected, *incA* mutants which retain Cdu1 activity did not accumulate Ub at or

near the periphery of inclusion membranes. In HeLa cells bearing both *cdu1*::GII and M923

inclusions, *cdu1::GII* Cdu1^{C345A} and M923 inclusions, or *cdu1*::GII Cdu1^{K268E} and M923

inclusions, only the M923 inclusions were protected from ubiquitination (Figures 5C and 5D).

Based on these observations we conclude that the acetylase activity of Cdu1 protects proteins

in cis and that this activity is constrained to the membrane of the pathogenic vacuole consistent

with previous reports (Auer et al., 2020).

Cdu1 is required for F-actin assembly and Golgi ministack repositioning around the Ct

inclusion, and for MYPT1 recruitment to Ct inclusions

InaC is required for Ct to assemble F-actin scaffolds and to reposition Golgi mini stacks

around the peripheries of inclusion membranes (Kokes et al., 2015; Wesolowski et al., 2017;

Haines et al., 2021). Because Cdu1 regulates InaC levels, we predicted that *cdu1* mutants

would phenocopy *inaC* mutants. We quantified the number of inclusions surrounded by F-actin

cages at 40 hpi. In cells infected with Rif-R L2 (wild type parental strain of M407, Nguyen and

Valdivia., 2012; Kokes et al., 2015), 25% of inclusions were surrounded by F-actin, consistent

with previous observations (Chin et al., 2012; Kokes et al., 2015) (Figures 6A and 6E). The

number of inclusions surrounded by F-actin decreased to 7% in cells infected with M407 (*inaC*

null) and increased to 49% in HeLa cells infected with M407 complemented with wild type InaC

(Figures 6A and 6E). A significant increase in inclusions surrounded by F-actin was also

observed in host cells infected with *cdu1*::GII mutants expressing Cdu1-Flag and Cdu1^{I225A}-

Flag (DUB-) (52% and 46% respectively) (Figures 6A and 6E) while *cdu1*::GII mutants

transformed with an empty plasmid or expressing Cdu1^{C345A}-Flag (DUB- Act-) and Cdu1^{K268E}-

Flag (Act-) resulted in only 8%,13%, and 10% of inclusions enveloped by F-actin respectively

(Figures 6A and 6E). From these observations we conclude that the acetylase activity of Cdu1

is required for Ct to promote assembly of F-actin around the Ct inclusion likely through the

stabilization of InaC.

We next quantified Golgi dispersal in infected HeLa cells at 24 hpi, a process that is also

dependent on InaC (Kokes et al., 2015; Wesolowski et al., 2017). In Hela cells infected with an

inaC null strain (M407) Golgi dispersal was limited to only 26% of the Ct inclusion perimeter. In

contrast, cells infected with either L2 Rif-R or with M407 complemented with wild type InaC, the

Golgi is dispersed around 45% of the inclusion perimeter (Figures 6B and 6D). Similarly, Golgi

dispersal around inclusions during infection with WT Ct L2 and in *cdu1*::GII mutants expressing

wild type Cdu1-Flag or Cdu1^{I225A}-Flag (DUB-) was 43%, 41%, and 43% respectively (Figures

6B and 6D). In HeLa cells infected with *cdu1*::GII and *cdu1*::GII strains expressing Cdu1^{C345A}-

Flag (DUB- Act-), and Cdu1^{K268E}-Flag (Act-), Golgi repositioning was restricted to only 24%,

23%, and 23% of inclusion perimeters respectively (Figures 6B and 6D). These results confirm

that both InaC and Cdu1 are required for efficient repositioning of the Golgi around the Ct

inclusion as previously reported (Kokes et al., 2015; Wesolowski et al., 2017; Pruneda et al.,

2018; Auer et al., 2020) and that this process is independent of Cdu1's DUB activity but requires

its acetylase activity. Moreover, our results suggest that Cdu1 promotes Golgi repositioning by

protecting InaC-mediated redistribution of the Golgi around the Ct inclusion.

CTL0480 promotes recruitment of the myosin phosphatase subunit MYPT1 to the inclusion

membrane where it regulates the extrusion of intact inclusions from host cells (Lutter et al.,

2013; Shaw et al., 2018). Consistent with the gradual loss of CTL0480 from inclusions in cells

infected with the *cdu1*::GII strain starting at 36 hpi (Figures 3C and 3D) we also observed a

complete loss of MYPT1 recruitment to inclusions by 48 hpi (Figures 6C and 6F).

Cdu1, InaC, IpaM, and CTL0480 are required for optimal extrusion of Ct from host cells

In the absence of Cdu1, the levels of InaC, CTL0480, and IpaM decreased late in infection (36

hpi and 48 hpi, Figures 3 and 4) suggesting that a prominent role of Cdu1 is to protect these

Incs from degradation late in infection. At the end of its developmental cycle, *Chlamydia* exits

host cells by promoting cellular lysis or by extrusion of intact inclusions (Hybiske and Stephens.,

2007). Ct host cell exit by extrusion is an active process requiring a remodeling of the actin

cytoskeleton and the function of Inc proteins (Hybiske and Stephens., 2007; Chin et al., 2012;

Lutter et al., 2013; Shaw et al., 2018; Nguyen et al., 2018). CTL0480 recruits MYPT1 (an

inhibitor of Myosin II motor complexes) to the inclusion membrane which prevents premature

extrusion of Ct inclusions and loss of CTL0480 leads to increased rates of extrusion by Ct from

infected HeLa cells (Lutter et al., 2013; Shaw et al., 2018) (Figure 7). Actin polymerization is

also required for Ct extrusion (Hybiske and Stephens., 2007; Chin et al., 2012) suggesting that

InaC dependent recruitment of F-actin to the inclusion may also contribute to optimal Ct

extrusion. IpaM localizes to microdomains in the inclusion membrane that are proposed to

function as foci for extrusion (Nguyen et al., 2018). Based on these observations, we postulated

that Cdu1-mediated protection of CTL0480, InaC, and IpaM regulates the extrusion of Ct

inclusions. We quantified the number of extrusions released from infected HeLa cells at 52 hpi

and observed a 60% reduction in the number of extrusions in HeLa cells infected with the

cdu1::GII strain relative to cells infected with WT L2 (Figures 7A and 7B). Complementation of

cdu1::GII with either wild type Cdu1-Flag or Cdu1^{I225A}-Flag (DUB-) restored extrusion

production to near wild type levels. Infection of HeLa cells with *inaC* (*inaC*::GII, Wesolowski et

al., 2017), or *ipaM* (*ipaM*::GII, Meier et al., 2020) null strains led to a 75% and 58% reduction in

extrusion production respectively (Figures 7A and 7B). Impaired extrusion of inclusions by

cdu1::GII, *inaC*::GII, and *ipaM*::GII strains was not due to defects in inclusion biogenesis, since

all three strains produced similar number of inclusions at 48 hpi relative to cells infected with WT

L2 (Supp. Figure 5). We conclude that Cdu1, InaC, and IpaM function to promote optimal

extrusion of Ct inclusions from host cells.

In contrast, infection of HeLa cells with a *ctl0480::GII* mutant strain led to an increase in the

number of extruded inclusions as previously observed (Shaw et al., 2018) (Figures 7A and 7B).

Therefore, even though the Cdu1-mediated protection of InaC and IpaM is important for the

extrusion of inclusions and *cdu1* mutants phenocopy the loss of InaC and IpaM, the phenotypic

similarities do not extend to the increased number of extruded inclusions observed in cells

infected with the *ctl0480*::GII mutant strain (Figures 7A and 7B). We infer from these

observations that functions for both InaC and IpaM in the extrusion of inclusions are epistatic to

CTL0480. Extruded inclusions produced during infection of HeLa cells also varied in size with

an average diameter of 40 µm (Figures 7A and 7C). Interestingly, the loss of IpaM and over

expression of Cdu1-Flag and Cdu1^{I225A}-Flag (DUB-) shifted the size distribution of extrusions

toward larger extrusions (Figures 7A and 7C) suggesting that Ct regulates the size of extruded

inclusions through Cdu1.

Discussion

Several *Chlamydia* effector proteins localize to the inclusion membrane to regulate interactions

between the *Chlamydia* pathogenic vacuole and the host cytoskeleton, organelles, and

vesicular trafficking pathways. They also modulate host cell death programs and promote

Chlamydia exit from host cells (reviewed in Bugalhão and Mota., 2019)**.** Given the central roles

that Incs play in promoting *Chlamydia* intracellular infection, it is not surprising that they are

targeted for inactivation by host cellular defenses. In response, *Chlamydia* has evolved

mechanisms to protect Incs. In this study we show that the acetylase activity of the effector

Cdu1 protects itself as well as three independent Incs; InaC, IpaM, and CTL0480, from

ubiquitination and degradation (Figure 8). Interestingly, all three Incs play prominent roles in

regulating the extrusion of *Chlamydia* inclusions from host cells (Figure 7). Observations that

the encapsulation of *Chlamydia* within an extruded inclusion enhances survival of *Chlamydia*

within macrophages (Zuck et al., 2017) together with the broad conservation of extrusion as an

exit strategy among *Chlamydia* (Zuck et al., 2016) suggests that this mechanism is important for

Chlamydia pathogenesis. Thus, targeting *Chlamydia* Incs that regulate extrusion for Ub-

mediated destruction may be advantageous for the host. In this context, we hypothesize that

InaC is targeted for degradation due to its role in promoting F-actin assembly around the

inclusion, since F-actin is essential for extrusion (Hybiske and Stephens., 2013; Chin et al.,

2012). However, other roles of InaC like its regulation of microtubule scaffolds around the

inclusion cannot be ruled out (Wesolowski et al., 2017; Haines et al., 2021). CTL0480 was

previously shown to function as an inhibitor of extrusions through its role in modulating the

activity of myosin light chain 2 (MLC_2) (Lutter et al., 2013; Shaw et al., 2018). IpaM localizes to

specialized microdomains in the inclusion membrane which are also sites of enrichment for over

9 inclusion membrane proteins including CTL0480 and MrcA, both of which are required for

Chlamydia extrusion (Mital et al., 2010; Lutter et al., 2013; Nguyen et al., 2018). IpaM is a third

inclusion membrane protein localizing to microdomains that is required for the optimal extrusion

of inclusions and lends support to a potential role for microdomains as foci for extrusion

(Nguyen et al., 2018). We also find that the loss of IpaM shifted the size distribution of

extrusions towards larger inclusions indicating that Ct constraints the size of extruded inclusions

(Figure 7). We speculate that heterogeneity in the size of extrusions might facilitate uptake of

some extrusions by innate immune cells at infected mucosal sites. Uptake of *Chlamydia* by host

phagocytes has been proposed to promote *Chlamydia* LGV dissemination to distal sites in the

genital tract to avoid clearance of *Chlamydia* by other immune cells (Zuck et al., 2017).

Protection of IpaM by Cdu1 could therefore be advantageous for *Chlamydia* EBs within

extrusions to escape to distal sites and avoid exposure to localized inflammatory responses.

Effectors that modulate the activity of other translocated effectors are referred to as

"metaeffectors", a term coined by Kubori and colleagues after observing that the *L. pneumophila*

effector LubX which functions as an E3 ligase, ubiquitinates the translocated effector SidH

leading to its degradation (Kubori et al., 2010). Several other effector-metaeffector interactions

have been described in *L. pneumophila*, *Salmonella enterica*, and *Brucella abortus* which

regulate the activity of other effectors either directly or indirectly by modifying the same host

target or cellular process (Kubori et al., 2010; Neunuebel et al., 2011; Jeong et al., 2015;

Urbanus et al., 2016; Smith et al., 2020; Iyer and Das., 2021). In this context Cdu1 functions as

an atypical metaeffector since it targets multiple effectors and unlike other effector-metaeffector

pairs it is not encoded in the same locus as its targets. We also observed that Cdu1 interactions

with InaC, IpaM, and CTL0480 likely occur independently from each other and that the

stoichiometry and kinetics of degradation varies for each Inc during infection of host cells. For

example, InaC protein levels decrease as early as 24 hpi while IpaM and CTL0480 protein

levels decrease at later stages of infection (36 and 48 hpi) (Figure 3). Furthermore, in the

absence of Cdu1, we observe almost complete loss of InaC at 36 hpi (Figure 4A) and of

CTL0480 at 48 hpi (Figures 3C and 3D), while only a subpopulation of IpaM is destabilized

throughout infection (Figures 3B and 4B).

The observation that the DUB activity of Cdu1 was not required to protect InaC, IpaM, and

CTL0480 from ubiquitination was unexpected and pointed to a potential role for its acetylation

activity instead. Because Cdu1 can interact with InaC, IpaM, and CTL0480 we hypothesized

that Cdu1 may protect these Incs through acetylation of lysine residues that are otherwise

targeted for ubiquitination. Indeed, we found that all three Incs and Cdu1 are acetylated at

lysines in infected cells, however, we were unable to determine if lysine acetylation in all four

proteins was dependent on Cdu1s' Act activity or if these PTMs are protective. Why the DUB

activity of Cdu1 is unable to compensate for loss of its Act activity remains unknown. It is

possible that Cdu1, like other DUBs, is regulated by PTMs (Komander et al., 2009a). For

instance, phosphorylation of human CYLD inhibits its DUB activity towards TRAF2 while

phosphorylation of human USP8 inhibits its DUB activity toward EGFR (Reiley et al., 2005;

Mizuno et al., 2007). Mass spectrometry analysis of immunoprecipitated Flag tagged Cdu1

expressed in Ct revealed that Cdu1 is phosphorylated at multiple serine and threonine residues

(Figure 4H) as previously suggested (Zadora et al., 2019). We identified three PX(S/T)P MAPK

phosphorylation consensus sequence motifs in the proline rich domain (PRD) of Cdu1,

suggesting that MAPKs may regulate the DUB activity of Cdu1.

Cdu1 homologs are found in multiple *Chlamydia* species including *C. trachomatis*, *C.*

muridarum, *C. suis*, *C. psitacci*, *C. abortus*, *C. caviae*, and *C. felis* but is notably absent in the

genomes of *C. pneumoniae* and *C. pecorum*. The acquisition of a second deubiquitinase

paralog (Cdu2) has also occurred in *C. trachomatis*, *C. muridarum*, and *C. suis*. In the genomes

of all three species, *cdu2* resides directly adjacent to *cdu1*; an arrangement that presumptively

arose from a gene duplication event. Cdu2 is a dedicated ULP with deubiquitinating and

deneddylating activities (Misaghi et al., 2006; Pruneda et al., 2016). New evidence suggest that

both paralogues might not be functionally redundant. The crystal structure of Cdu2 has revealed

differences in residues involved in substrate recognition between Cdu1 and Cdu2 and that each

paralog might recognize polyUb chains differently (Hausman et al., 2020). The processivity

rates for removal of terminal Ub from polyUb chains also differs between both isopeptidases

with Cdu2 exhibiting limited trimming of polyUb as compared to Cdu1 (Hausman et al., 2020).

Moreover, Cdu2 lacks the proline rich domain found in Cdu1 which might be important for

regulation of Cdu1 enzymatic activity. The presence of Cdu2 might also explain the low

incidence of human and Ct proteins that were differentially ubiquitinated in the absence of Cdu1

(Figure 1). Whereas several *Chlamydia* species have acquired either one or two deubiquitinase

paralogs, both *C. pneumoniae* and *C. pecorum* have not. Instead, both species have acquired

an unrelated deubiquitinase (*Chla*OTU) belonging to the OTU family of proteases (Makarova et

al., 2000; Furtado et al., 2013). Curiously, *Chla*OTU is also found in *C. psitacci*, *C. abortus*, *C.*

caviae, and *C. felis* all of which encode only Cdu1 and is absent in *C. trachomatis*, *C.*

muridarum, and *C. suis*, all of which encode Cdu1 and Cdu2. It is noteworthy that *Chlamydia*

species have independently acquired deubiquitinases multiple times (Cdu1, Cdu2, *Chla*OTU)

and that some of these deubiquitinases have evolved into moonlighting enzymes reflecting the

diverse strategies adopted by pathogenic *Chlamydia* as they adapt to their particular niche.

Acknowledgements

We thank LifeSensors and the Duke Proteomics and Metabolomics Shared Resource Center for

their proteomics services. We thank the Duke Light Microscopy Core Facility for microscopy

services. We also thank Marcela Kokes for generating the IncA-Flag constructs used in this

study. This work was supported by NIH grants AI140019 to R.J.B and AI134891 to R.H.V.

Author contributions

R.J.B and R.H.V designed the study. R.J.B wrote the manuscript with input from all listed

authors. L.D generated the *cdu1*::GII strain. M.K verified generation of the *cdu1*::GII mutant

strain and showed interaction of Cdu1-GFP variants with InaC, IpaM, and CTL0480 in

transfected HEK cells. R.J.B performed all other experiments. B.S.S and J.N.P shared reagents.

R.J.B and R.H.V proofed the manuscript.

Declaration of interests

R.H.V is a founder of Bloom Sciences (San Diego, CA), which is a microbiome therapeutics

company. Findings reported in this study are unrelated to the work being performed with Bloom

Sciences.

Material and Methods

Key Resource Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		

See S. Table 9		
Bacterial and virus strains		
Chlamydia trachomatis LGV biovar L2 434 Bu (L2)	Richard Stephens (UC Berkeley)	N/A
L2 pBOMB-MCI (Parent: LGV L2 434 Bu)	This paper	N/A
L2 pBOMB-MCI_CTL0480-3XFlag (Parent: LGV L2 434 Bu)	This paper	N/A
L2 pBOMB-MCI_IpaM-3XFlag (Parent: LGV L2 434 Bu)	This paper	N/A
L2 pBOMB-MCI_Cdu1-3XFlag (Parent: LGV L2 434 Bu)	This paper	N/A
L2 pBOMB-MCI_Cdu1 ^{C345A}-3XFlag (Parent: LGV L2 434 Bu)	This paper	N/A
L2 pBOMB-MCI_Cdu1 ^{K268E}-3XFlag (Parent: LGV L2 434 Bu)	This paper	N/A
L2 *cdu1*::GII *aadA* (Parent: LGV L2 434 Bu)	This paper	N/A
L2 *cdu1*::GII *aadA* pBOMB-MCI (Parent: L2 *cdu1*::GII *aadA*)	This paper	N/A
L2 *cdu1*::GII *aadA* pBOMB-MCI_Cdu1-3XFlag (Parent: L2 *cdu1*::GII *aadA*)	This paper	N/A
L2 *cdu1*::GII *aadA* pBOMB-MCI_Cdu1 ^{C345A}-3XFlag (Parent: L2 *cdu1*::GII *aadA*)	This paper	N/A
L2 *cdu1*::GII *aadA* pBOMB-MCI_Cdu1 ^{I225A}-3XFlag (Parent: L2 *cdu1*::GII *aadA*)	This paper	N/A
L2 *cdu1*::GII *aadA* pBOMB-MCI_Cdu1 ^{K268E}-3XFlag (Parent: L2 *cdu1*::GII *aadA*)	This paper	N/A
L2 Rif-R (Parent: L2 434 Bu)	Nguyen et al., 2012	N/A
M407 (*inaC C307T*, InaC Q103*) (Parent: L2 Rif-R)	Kokes et al., 2015	N/A
M407 p2TK2 (Parent: M407)	Kokes et al., 2015	N/A
M407 p2TK2_InaC (Parent: M407)	Kokes et al., 2015	N/A
M407 p2TK2_InaC-3X Flag (Parent: M407)	This paper	N/A
inaC::GII *bla* (Parent: L2 434 Bu)	Wesolowski et al., 2017	N/A
ipaM::GII *cat* (Parent: L2 434 Bu)	Meier et al., 2022	N/A
ctl0480::GII *aadA* (Parent: L2 434 Bu)	Shaw et al., 2018	N/A
M923 (*incA C589T*, IncA R197*) (Parent: L2 Rif-R)	Kokes et al., 2015	N/A
M923 pBOMB-MCI (Parent: M923)	Sixt et al., 2017	N/A
Biological samples		
N/A		
Chemicals, peptides, and recombinant proteins		
PR-619 (DUB inhibitor)	Sigma-Aldrich	Cat# SML0430

MG132 (proteasome inhibitor)	Sigma-Aldrich	Cat# 474791
TUBE-1 (pUb affinity capture reagent)	LifeSensors	Cat# UM401M
Acti-stain™ 488 (Phalloidin 488)	Cytoskeleton Inc.	Cat# PHDG1
ChromoTek GFP-Trap® Agarose	Proteintech	Cat# gta
Protein A/G PLUS-Agarose	Santa Cruz Biotechnology	Cat# sc-2203
3xFLAG peptide	APExBIO	Cat# A6001
LGV L2 434 Bu Cdu1 recombinant protein	Jonathan Pruneda (Oregon Health and Science University)	N/A
Critical commercial assays		
TargeTron™ gene knockout system	Sigma-Aldrich	Cat# TA0100
Deposited data		
C. trachomatis LGV L2 434 Bu proteome	NCBI	NCBI:txid47472
Homo sapiens (Human) proteome	UniProt	Proteome ID: UP000005640
TUBE-1 affinity capture proteomics data	PRIDE	To be deposited
Cdu1 lysine acetylation proteomic data	PRIDE	To be deposited
Experimental models: Cell lines		
Hela cells	ATCC	Cat# CCL-2 RRID:CVCL_0030
Vero cells	ATCC	Cat# CCL-81 RRID:CVCL_0059
HEK 293T cells	ATCC	Cat# CRL-3216 RRID:CVCL_0063
Experimental models: Organisms/strains		
Chlamydia trachomatis LGV L2 434 Bu	Richard Stephens (UC Berkeley)	N/A
Oligonucleotides		
See S. Table 10		
Recombinant DNA		
Plasmid: pDFTT3-aadA_Cdu1 635/636	This paper	N/A
Plasmid: pOPIN-GFP_Cdu1 FL (aa 1-401)	Jonathan Pruneda (Oregon Health and Science University)	N/A
Plasmid: pOPIN-GFP_Cdu1 TMD- (aa 71-401)	Jonathan Pruneda (Oregon Health and Science University)	N/A
Plasmid: pOPIN-GFP_Cdu1 CD- (aa 1-130)	Jonathan Pruneda (Oregon Health and Science University)	N/A
Plasmid: pCDNA-DEST53 (w/o GFP)_InaC (CT813)-3XFLAG	This paper	N/A
Plasmid: pcDNA3.1/nV5-DEST_IpaM	This paper	N/A
Plasmid: pcDNA3.1/nV5-DEST_CTL0480 p	This paper	N/A
Plasmid: pcDNA3.1/nV5-DEST_CpoS	This paper	N/A
Plasmid: pBOMB4-MCI_CTL0480-3X Flag	This paper	N/A
Plasmid: pBOMB4-MCI_IpaM-3X Flag	This paper	N/A
Plasmid: pBOMB4-MCI_Cdu1-3XFlag	This paper	N/A
Plasmid: pBOMB4-MCI_Cdu1 C345A-3XFlag	This paper	N/A

Plasmid: pBOMB4-MCI_Cdu1 I225A-3XFlag	This paper	N/A
Plasmid: pBOMB4-MCI_Cdu1 K263E-3XFlag	This paper	N/A
Plasmid: p2TK2_SW2_InaC-3XFlag	This paper	N/A
Software and algorithms		
Targetronics	Targetronics, LLC	www.targetrons.com
Proteome Discoverer 2.3	Thermo Fisher Scientific	https://www.thermofisher.com/us/en/home/industrial/mass-spectrometry/liquid-chromatography-mass-spectrometry-lc-ms/lc-ms-software/multi-omics-data-analysis/proteome-discoverer-software.html
Mascot software	Matrix Science	https://www.matrixscience.com
MaxQuant 1.6.2.3		https://www.maxquant.org
Scaffold PTM	Proteome Software	https://www.proteomesoftware.com/products/scaffold-ptm
VolcaNoseR	Goedhart, and Luijsterburg., 2020	https://huygens.science.uva.nl/VolcaNoseR/
Metascape	Zhou et al., 2019	https://metascape.org/gp/index.html#/main/step1
DAVID Bioinformatic Resources	Huang et al., 2009a., Huang et al., 2009b	https://david.ncifcrf.gov/tools.jsp
Image J	Schneider et al., 2012	https://imagej.nih.gov/ij/
NEBaseChanger	New England Biolabs	https://nebasechanger.neb.com
Prism 9	GraphPad	https://www.graphpad.com/updates/prism-900-release-notes
Other		
N/A		

505
506

507 **Resource Availability**

508 **Materials availability**

509 • All newly generated materials associated with this study will be freely available upon

510   request.

Data and code availability

- Proteomics data will be deposited at PRIDE (Proteomics Identification Database-EMBL-EBI) and publicly available at the time of publication. Accession numbers will be listed in the key resources table.

- Original data used for microscopy and western blots in this study will be deposited online (Mendeley data) and made available at the time of publication.

- Any additional information required to reanalyze the data reported in this paper is available upon request.

- This paper does not report original code.

Experimental Model And Subject Details

Cell lines

Vero (CCL-81; RRID:CVCL_0030), HeLa (CCL-2; RRID:CVCL_0059), and HEK293T (CRL-3216; CVCL_0063) cells were purchased from ATCC and cultured in High Glucose Dulbecco's Modified Eagle's Medium supplemented with L-glutamine, sodium pyruvate (DMEM; Gibco) and 10% fetal bovine serum (FBS; Sigma-Aldrich). Cells were grown at 37°C in a 5% CO2 humidified incubator. Vero, Hela, and HEK293T cells were derived from females. All three cell lines have been authenticated by the Duke Cell Culture and DNA analysis facility.

Chlamydia **strains and propagation**

Chlamydia strains used in this study are listed in the Key Resources Table. Ct strains were

propagated in Vero cells and harvested by osmotic lysis at 48 hours post infection. Following

lysis extracts were sonicated and bacteria pelleted by centrifugation at 21,000 x g. Bacteria

were resuspended in SPG storage buffer (75g/L sucrose, 0.5 g/L KH_4HPO_4, 1.2 g/L Na_2HPO_4,

0.72 g/L glutamic acid, pH 7.5) and stored as single use aliquots at -80°C.

Method Details

***Chlamydia* infections**

Chlamydia infections were synchronized by centrifugation (2,500 x *g* for 30 minutes at 10°C)

onto HeLa cell monolayers and incubated for the indicated times. Co-infections were performed

by infecting HeLa cell monolayers at a 1:1 ratio using MOIs of 2 for each co-infecting strain.

Insertional mutagenesis of CTL0247 (*cdu1*)

Primer sequences for TargeTron™ mediated mutagenesis of the LGV L2 434 Bu *cdu1*

(CTL0247) ORF were designed at the TARGETRONICS, LLC web portal

(www.targetrons.com). IBS1/2, EBS1/delta, and EBS2 primers (primer sequences are listed in

S. Table 10) were used in a PCR reaction to generate homing sequences for TargeTron™

integration between nucleotides 635 and 636 of the *cdu1* ORF using a TargeTron™ gene

knockout system (Sigma-Aldrich; TA0100) according to the manufacturer instructions. Homing

sequences were gel purified, digested with HindIII and BsrGI, and ligated into HindIII and BsrGI

digested pDFTT3-*aadA* (Lowden et al., 2015). Ligations were transformed into *E. coli* DH5α,

clones isolated, and *cdu1* redirected pDFTT3-aadA plasmids identified by restriction digest and

verified by Sanger sequencing (Eton Bioscience) using a T7-promoter specific primer. The

resulting plasmid was transformed into a *C. trachomatis* LGV L2 434 Bu strain and

transformants selected with 150 μg/mL spectinomycin and plaque purified as previously

described (Kędzior and Bastidas., 2019). Insertion of the GII *aadA* intron at the *cdu1* locus was

verified by PCR analysis (S. Figure1) using primers that amplify amplicons spanning the

cdu1::GII 5' (RBP409 and RBP436) and 3' (RBP468 and RBP118) junctions, the *cdu1* CDS

(RBP409 and RBP118), and the *aadA* CDS (RBP512 and RBP513). Primer sequences are

listed in S. Table 10. Loss of Cdu1 protein was verified by western blot and indirect

immunofluorescence analysis (S. Figures 2A and 2B).

Analysis of *cdu1* and *cdu2* transcription by RT-PCR

Confluent HeLa cell monolayers (2.9 x 10^6 cells/infection) were infected with wild type L2 434

Bu or L2 *cdu1*::GII *aadA* strains. At 24 hpi, total RNA was isolated with a Qiagen RNeasy kit

(Qiagen; 74004) according to the manufacturer instructions. Total RNA was treated twice with

DNAse I (NEB; M0303S) and used for cDNA synthesis using a SuperScript IV Reverse

Transcriptase kit (Thermo Fisher Scientific; 18090010). cDNAs synthesized with and without

reverse transcriptase were used as templates for PCR analysis (S. Figure 2C) using primers

that amplify amplicons spanning the *cdu*1 (CTL0247_F and CTL0247_R) and *cdu2* ORFs

(CTL0246_F and CTL0247_R), and the intergenic junction between the *cdu1* and *cdu2* ORFs

(CTL0246-0247_F and CTL0246-0247_R). Primer sequences can be found in the S. Table 10.

TUBE1 based global ubiquitin profiling

Confluent HeLa cell monolayers (5.04 x 10^6 cells/infection) were mock infected or separately

infected with WT LGV L2 434 Bu or a L2 *cdu1*::GII *aadA* strain at MOIs of 3. At 24 hpi cells were

collected and spun down (700 x *g* for 10 minutes), frozen at -80°C and shipped on dry ice to

LifeSensors (Malvern, PA) for quantitative TUBE1-based Mass Spectrometry Analysis. Cell

pellets from three independent biological replicates were sent to LifeSensors for analysis. Cell

were subsequently lysed in lysis buffer (50 mM Tris-HCL, pH 7.5, 150 mM NaCl, 2 mM EDTA,

1% NP-40, 10% glycerol, 1% Sodium Deoxycholate) supplemented with a protease inhibitor

cocktail, the DUB inhibitor PR-619 (Sigma-Aldrich; SML0430), and the proteasomal inhibitor

MG-132 (Sigma-Aldrich; 474791). Lysates were clarified by high-speed centrifugation (14,000 x

g, 10 minutes, 4°C) and supernatants containing 5 mg of protein were equilibrated with

magnetic TUBE-1 (LifeSensors; UM401M) and incubated overnight at 4°C under rotation.

TUBEs were isolated with a magnetic stand and washed sequentially with PBST and TUBE

wash buffer. Poly-ubiquitinated and associated proteins were eluted with TUBE elution buffer.

Eluted supernatants were neutralized with neutralization buffer and loaded onto SDS-gels and

run until SDS buffer reached 0.5 cm into the gel. Gels were stained with Coomassie Blue and

lanes excised, reduced with TCEP, alkylated with iodoacetamide, and digested with Trypsin

(Trypsin Gold, Mass Spectrometry Grade) (Promega; V5280). Tryptic digests were analyzed

using a 150 min LC run on a Thermo Scientific™ Q Exactive HF Orbitrap™ LC-MS/MS system.

MS data was searched against the UniProt human database (UniProt; Proteome ID:

UP000005640) and the *Chlamydia trachomatis* L2 434 Bu reference database (NCBI:txid47472)

using MaxQuant 1.6.2.3 (https://www.maxquant.org). Proteins, peptides, and site identification

was set to a false discovery rate of 1%. N-terminal acetylation, Met oxidation, and diGly remnant

on lysine residues was also identified. All peptides and proteins identified can be found in S.

Table 1. The intensities (sum of all peptide MS peak areas for a protein or ubiquitinated peptide)

for each protein and ubiquitinated peptide across all three biological replicates were used to

determine mean intensities and to calculate *p*-values based on one-way student t-tests. Volcano

plots of mean intensities vs. *p*-values were generated with VolcaNoseR (Goedhart and

Lujsterburg., 2020) (https://huygens.science.uva.nl/VolcaNoseR/) and used to identify

significantly enriched proteins and ubiquitinated peptides. Data used to generate each Volcano

plot can be found in S. Table 2. Pathway enrichment analysis was performed with Metascape

(Zhou et al., 2019) (https://metascape.org/gp/index.html#/main/step1) and DAVID bioinformatic

resources (Huang et al., 2009a; Huang et al., 2009b) (https://david.ncifcrf.gov/tools.jsp).

Inhibitors, antibodies, and western blots

MG-132 (25 μM) (Sigma-Aldrich; 474791) was added to infected cell monolayers 5 hours prior

to extract preparations. Recombinant Cdu1 protein (LGV L2 434 Bu, amino acids 71-401) was

generated as previously described (Pruneda et al., 2018) and kindly provided by Jonathan

Pruneda (Oregon Health and Science University, OR). Recombinant Cdu1 protein was used to

generate antibodies in immunized New Zealand White rabbits. Cdu1 antisera was pre-adsorbed

with crude cell extracts from HeLa cells infected with a *cdu1*::GII *aadA* strain. Pre-adsorbed

antisera was used for western blot analysis at a 1:500 dilution in a solution containing 5% BSA

supplemented with crude extracts from HeLa cells infected with a *cdu1*::GII *aadA* strain

(0.1mg/mL total protein). Antibodies, antibody dilutions, and antibody diluents used in this study

are listed in S. Table 9. For western blot analysis, lysates from infected HeLa cell monolayers

(2.4 x 10^6 cells) were prepared by incubating cell monolayers with boiling hot 1% SDS lysis

buffer (1% SDS, 100 mM NaCl, 50 mM Tris, pH 7.5). Lysates were collected, briefly sonicated,

and total protein concentration measured with a DC^{TM} Protein Assay Kit (BIO-RAD; 5000111). 8

μg (Slc1 and alpha Tubulin blots) and 25 μg of total protein lysates (all other blots) were loaded

onto 4-15% Mini-PROTEAN and TGX Stain Free^{TM} Protein Gels (BIO-RAD; 4568084),

transferred to PVDF membranes (BIO-RAD; 1620177), blocked with 5% Milk/TBSt, and

incubated with primary antibodies overnight at 4$°$C. Protein signals were detected with Goat anti

mouse (H+L) IgG (ThermoFisher scientific; 31430) or Goat anti rabbit (H+L) IgG HRP

(ThermoFisher scientific; 31460) conjugated secondary antibodies (1:1000 in 5% Milk/TBSt) and

SuperSignal West Femto HRP substrate (ThermoFisher scientific; 34096). Antibody bound

membranes were imaged with a LI-COR imaging system (LI-COR, Inc.). Varying amounts of

protein extracts were used to determine the linear range of detection for InaC, IpaM, and Slc1

antibodies prior to quantification of western blot images with LI-COR imaging software (data not

shown).

Immunofluorescence microscopy

HeLa cells were grown on coverslips to 50% confluency (0.1×10^5 cells) and infected at MOIs of

0.6. At indicated times, infected cells were separately fixed with ice cold Methanol or with warm

PBS containing 4% formaldehyde for 20 minutes. After fixative removal, cells were washed with

PBS and formaldehyde fixed cells were incubated either in 5% BSA/PBS supplemented with

0.1% Triton X-100 or in 5% BSA/PBS supplemented with 0.05% Saponin for 30 minutes with

gentle rocking. Following washing with PBS, Methanol fixed cells were incubated with primary

antibodies diluted in 5% BSA/PBS and formaldehyde fixed cells were incubated with primary

antibodies diluted in 5% BSA/PBS supplemented with 0.1% Triton X-100 or 0.05% Saponin for

1 hour with gentle rocking. Dilutions for each antibody used can be found in S. Table 9.

Methanol fixed cells were washed with PBS and incubated with secondary antibodies diluted in

5% BSA/PBS and supplemented with Hoechst 33342 (2 µg/mL) (ThermoFisher Scientific;

H3570). Formaldehyde fixed cells were washed and incubated with 5% BSA/PBS supplemented

with 0.1% Triton X-100 and Hoechst or 0.05% Saponin and Hoechst for 1 hour protected from

light and with gentle rocking. For detection of F-actin, Phalloidin conjugated to Alexa Fluor™

488 (1:5000) (Act-Stain 488 Phalloidin; Cytoskeleton Inc.; PHDG1) was added for the last 20

minutes of incubation with the secondary antibodies. Coverslips were transferred to glass slides,

mounted with 10 µL of Vectashield (Vector Labs; H-1000) and incubated over night at room

temperature prior to imaging. Secondary antibodies used were goat anti-mouse (H+L) IgG

(ThermoFisher scientific; A-11001 and A-21235) and goat anti-rabbit (H+L) IgG (ThermoFisher

scientific; A-11008 and A-21244) conjugated to Alexa Fluor™ 488 and Alexa Fluor™ 647. All of

the antibodies used for indirect immunofluorescence analysis were analyzed under all three

staining conditions (Methanol, Formaldehyde/Triton X-100, and Formaldehyde/Saponin).

Representative images were acquired with an inverted confocal laser scanning microscope

(Zeiss 880) equipped with an Airyscan detector (Hamamatsu) and with diode (405 nm), argon

ion (488 nm), double solid-state (561 nm) and helium-neon (633) lasers. Images were acquired

with a 63x C-Apochromatic NA 1.2 oil-objective (Zeiss). Images acquired in Airyscan mode were

deconvoluted using automatic Airyscan processing in Zen software (Zeiss). Image acquisition

was performed at the Light Microscopy Core Facility at Duke University. Images used for

quantification were captured in an inverted microscope (Ti2-Nikon instruments) equipped with

an ORCA Flash 4.0 V3 sCMOS camera (Hamamatsu) and a SOLA solid-state white light

illuminator (Lumencro). Images were acquired using a 60x Plan Apochromatic NA 1.40 oil

objective. All images were opened with ImageJ (Schneider et al., 2012) and only linear

adjustments were made to fluorescence intensity for the entire image. Images were exported as

TIFFs and compiled with Adobe suite software (Illustrator).

Vector construction and *C. trachomatis* transformation

Constructs used in co-transfection experiments: Mammalian vectors expressing Cdu1-GFP

constructs were kindly provided by Jonathan Pruneda (Oregon Health and Science University,

OR). Briefly, geneblocks encoding full length Cdu1 (LGV L2 434 Bu, CTL0247) (amino acids 1-

401), Cdu1 lacking its transmembrane domain (amino acids 71-401), and Cdu1 lacking its

catalytic domain (amino acids 1-130) were generated and inserted into the pOPIN-GFP vector

(Berrow et al., 2007) by In-Fusion[TM] cloning (Takara Bio; 638947), resulting in Cdu1 constructs

with a C-terminal eGFP-His tag preceded by a 3C protease cleavage site. The Flag-InaC

mammalian expression vector was derived from a Gateway[TM] entry clone containing the *C.*

trachomatis Serovar D/UW-3/CX CT813 (*inaC*) ORF (amino acids 41-264) obtained from a *C.*

trachomatis ORFeome library (Roan et al., 2006). The entry vector was used as a donor

plasmid for Gateway[TM] based transfer into a modified pcDNA[TM] DEST53 (ThermoFisher

Scientific; 12288015) vector in which the cycle 3 *GFP* ORF was removed. A NEB Q5[®]-Site

Directed Mutagenesis Kit (New England Biolabs; E0554S) was used to introduce a 3X Flag

epitope tag at the N-terminus of the CT813 ORF and a stop codon at the end of the CT813

ORF. L2 *ipaM* (CTL0476), L2 CTL0480, and L2 *cpoS* (CTL0481) ORFs were PCR amplified

from cell lysates derived from Vero cells infected with wild type L2 LGV 434 Bu with primers

containing attB sequences (primers *ipaM* forward, *ipaM* reverse, CTL0480 forward, CTL0480

reverse, *cpoS* forward, and *cpoS* reverse). Primer sequences can be found in the S. Table 10.

PCR amplicons were used as donors for Gateway[TM] BP Clonase[TM] based transfers into the

donor vector pDONR[TM]221 (ThermoFisher Scientific; 12536017) to generate entry plasmids.

Entry plasmids were used to transfer *ipaM*, CTL0480, and *cpoS* into the Gateway[TM] destination

vector pcDNA[TM]3.1/nV5-DEST (ThermoFisher Scientific; 12290010) by Gateway[TM] LR

Clonase[TM] based reactions. The resulting mammalian expression vectors express IpaM,

CTL0480, and CpoS with V5-epitopes fused to their N-terminus.

pBOMB4-MCl based plasmids. CTL0480 and *ipaM* ORFs were amplified by PCR from cell

extracts derived from Vero cells infected with wild type LGV L2 434 Bu. The CTL0480 ORF, 149

b.p of upstream sequence, and a 3X FLAG epitope was amplified by PCR using primers

RBP628 and RBP629. The CTL0476 (*ipaM*) ORF, 400 b.p of upstream sequence, and a 3X

FLAG epitope was amplified with primers RBP623 and RBP624. CTL0480 and *ipaM* amplicons

were digested with Not1 and Pst1 and cloned into Not1 and Pst1 digested pBOMB4-MCl

(Bauler and Hackstadt., 2014) to generate pBOMB4-MCl_CTL0480-3X Flag and pBOMB4-

MCl_IpaM-3X Flag plasmids respectively. pBOMB4-MCl_Cdu1-3XFlag plasmids were

generated by PCR amplification of 175 b.p of genomic sequence directly upstream of the L2

434 Bu CTL0247 (*cdu1*) ORF and the entire *cdu1* ORF tagged with a C-terminal 3X Flag

epitope tag (primers RBP460 and RBP461). PCR amplicons were generated from gradient

purified LGV L2 434 Bu EBs and cloned into a pCR™-Blunt II TOPO® vector using a Zero

Blunt™ TOPO™ PCR cloning Kit (ThermoFisher Scientific; K2800J10) according to the

manufacturer instructions. Cdu1 catalytic variants were generated with a NEB Q5®-Site Directed

Mutagenesis Kit (New England Biolabs; E0554S) using the *cdu1*p-*cdu1*-3X Flag construct

cloned into pCR™-Blunt II TOPO® as a template and following the manufacturer instructions.

Primers for introducing base pair changes were designed on the NEBaseChanger website

(https://nebasechanger.neb.com). The Cdu1^{C345A} variant was generated by changing the *TGC*

codon located at positions 1033-1035 in the *cdu1* ORF to *GCT* (primers RBP525 and RBP526).

The Cdu1^{I225A} variant was generated by substituting the *ATC* codon located at positions 673-

675 for *GCT* (primers RBP527and RBP528). The Cdu1^{K268E} variant was generated by

introducing an *A802G* base pair substitution (primers RBP529 and RBP530). Wild type *cdu1*p-

cdu1-3XFLAG and all three *cdu1* variants were digested with Not1 and Pst1 and ligated into

Not1 and Pst1 digested pBOMB4-MCI (Bauler and Hackstadt., 2014). Primer sequences can be

found in the S. Table 10.

p2TK2_SW2-inaC-3XFlag. The CTL0184 (*inaC*) ORF and 250 b.p of upstream sequence was

amplified by PCR from cell extracts derived from Vero cells infected with L2 434 Bu and cloned

into the p2TK2_SW2 vector (Agaisse and Derré., 2013). A NEB Q5®-Site Directed Mutagenesis

Kit (New England Biolabs; E0554S) was used to insert a 3X FLAG epitope sequence at the C-

terminus (stop codon removed) of the CTL0184 ORF to generate the p2TK2_SW2-InaC-3XFlag

plasmid.

pBOMB4-MCI based plasmids and p2TK2_SW2-InaC-3X Flag plasmids were transformed into

corresponding *Chlamydia* strains, and transformants were selected with 10 U/mL Penicillin G

and plaque purified as previously described (Kędzior and Bastidas., 2019). All primer

sequences and plasmids generated in this study are listed in S. Table 10 and Key Resources

Table.

Subcellular fractionation

HeLa cells (2.16×10^7 cells/strain) seeded in six well plates were mock infected or infected with

Chlamydia L2 434 Bu strains. At indicated time points cells were washed with ice-cold PBS,

collected in ice-cold PBS with a cell scraper, and transferred to 15 mL conical tubes. Cell

suspensions were centrifuged at 500 x g for 5 minutes at 4°C and cell pellets were resuspended

in 400 μL of ice-cold subcellular fractionation buffer (20 mM HEPES (pH 7.4), 10 mM KCl, 2mM

$MgCl_2$, 1mM EDTA, 1mM EGTA) supplemented with 1mM DTT and a 1x cOmplete Mini-EDTA

free protease inhibitor cocktail (Sigma-Aldrich; 11836170001). Cells were incubated on ice for

20 minutes and lysed with 30 strokes of a Dounce homogenizer. Cell lysates were sequentially

centrifuged twice at 720 x g for 5 minutes at 4°C to remove intact nuclei. Supernatants were

centrifuged at 10,000 x g for 5 minutes at 4°C and the heavy membrane (inclusion) fraction was

recovered and resuspended in IP lysis buffer (25 mM Tris-Hcl, 150 mM NaCl, 1% NP-40, 5%

Glycerol) supplemented with 1mM PMSF and a 1x cOmplete Mini-EDTA free protease inhibitor

cocktail (Sigma-Aldrich; 11836170001).

Immunoprecipitations (IPs)

Transfections and GFP-immunoprecipitations: HEK 293T cell monolayers (1.32×10^7

cells/transfection) seeded in 10 cm cell culture dishes pre coated with poly-L-Lysine (Sigma-

Aldrich; P4707) were grown to 50% confluency and transfected with 10 μg of each plasmid used

per co-transfection in a 1.5 to 1 ratio of jetOPTIMUS® (Polyplus; 101000051) transfection

reagent to total plasmid DNA according to the manufacturer instructions. At 24 hpi, transfected

cells were lysed in IP lysis buffer (described above) supplemented with 1mM PMSF and 1x

cOmplete Mini-EDTA free protease inhibitor cocktail (Sigma-Aldrich; 11836170001). Lysates

were transferred to Eppendorf tubes, sonicated, and cleared by centrifugation (21,000 x g, 15

minutes, 4°C). Supernatants containing 2 mg of total protein were incubated with magnetic

GFP-Trap® agarose (Proteintech; gta) for 1 hour at 4°C with rotation. Beads were washed

according to manufacturer instructions and immunoprecipitated proteins eluted with 2X Laemmli

sample buffer.

Flag immunoprecipitations: Mock and infected HeLa cell monolayers (1.44 x 10^7 cells) grown in

six well plates were lysed in IP lysis buffer (described above) and transferred to Eppendorf

tubes. Lysates were sonicated and cleared by centrifugation (21,000 x g, 15 minutes, 4°C).

Supernatants containing 2 mg of total protein were pre-cleared by incubating with Protein A/G

PLUS-Agarose (Santa Cruz Biotechnology; sc-2203) for 30 minutes at 4°C followed by

sedimentation of agarose resins by centrifugation. Supernatants were incubated with M2-anti

Flag mouse mAb (1:400) (Sigma-Aldrich; F1804) overnight at 4°C with rotation followed by

incubation with Protein A/G PLUS-Agarose for 3 hours at 4°C with rotation. Agarose resins were

sedimented and washed according to manufacturer instructions and immunoprecipitated

proteins eluted with 50 μL of 100 μg/mL 3xFLAG peptides (APExBIO; A6001).

Acetylated lysine immunoprecipitations. Heavy membrane (inclusion) subcellular fractions

isolated from infected HeLa cells and containing 1 mg of total protein were pre-cleared by

incubating with Protein A/G PLUS-Agarose (Santa Cruz Biotechnology; sc-2203) for 30 minutes

at 4°C followed by sedimentation of agarose resins by centrifugation. Supernatants were

incubated with an anti-acetylated lysine rabbit antibody (Cell signaling; #9441) (1:100) overnight

at 4°C with rotation followed by incubation with Protein A/G PLUS-Agarose for 3 hours at 4°C

with rotation. Agarose resins were sedimented and washed according to manufacturer

instructions and immunoprecipitated proteins eluted with 50 μL of 2X Laemmli sample buffer.

Input (8 μg of total protein for Slc1 and alpha Tubulin blots, and 25 μg for all other blots) and

immunoprecipitates (GFP, Flag, proteins acetylated at lysines) were loaded onto 4-15% Mini-

PROTEAN and TGX Stain Free™ Protein Gels (BIO-RAD; 4568084), transferred to

PVDF membranes (BIO-RAD; 1620177), blocked with 5% Milk/TBSt, and incubated with

primary antibodies overnight at 4°C. Protein signals were detected with Goat anti mouse (H+L)

IgG (ThermoFisher scientific; 31430) or Goat anti rabbit (H+L) IgG HRP (ThermoFisher

scientific; 31460) conjugated secondary antibodies (1:1000 in 5% Milk/TBSt) and SuperSignal

West Femto HRP substrate (ThermoFisher scientific; 34096). Antibody bound membranes were

imaged with a LI-COR imaging system (LI-COR, Inc.).

Identification Cdu1 lysine acetylation and phosphorylation sites by LC-MS/MS

HeLa cell monolayers (8.64 x10^7 cells/strain) grown in six well plates were infected with a wild

type L2 strain transformed with empty pBOMB4-MCI (Bauler and Hackstadt., 2014) plasmid or

with a wild type L2 strain transformed with a pBOMB4-MCI_*cdu1*-3X Flag plasmid. At 24 hpi

infected cells were lysed and Flag tagged Cdu1 was immunoprecipitated as described above.

Flag eluates from 3 independent biological replicates were sent to the Proteomics and

Metabolomics Shared Resource Facility at Duke University for quantitative LC-MS/MS analysis.

Samples were spiked with undigested casein, reduced with 10 mM dithiothreitol, and alkylated

with 20 mM iodoacetamide. Eluates were then supplemented with 1.2% phosphoric acid and S-

Trap (Protifi) binding buffer (90% Methanol, 100 mM TEAB). Proteins were trapped on the S-

Trap, digested with 20 ng/μL Trypsin (Trypsin Gold, Mass Spectrometry Grade) (Promega;

V5280), and eluted with 50 mM TEAB, 0.2% FA, and 50% ACN/0.2% FA. Samples were

lyophilized and resuspended in 1% TFA/2% acetonitrile containing 12.5 fmol/μL yeast alcohol

dehydrogenase. Quantitative LC/MS/MS was performed using a nanoAcquitiy UPLC system

(Waters Corp) coupled to Thermo Scientific[TM] Orbitrap[TM] Fusion Lumos high resolution accurate

mass tandem mass spectrometer via a nanoelectrospray ionization source. Data was analyzed

with Proteome Discoverer 2.3 (Thermo Fisher Scientific[TM].) and MS/MS data searched against

the *Chlamydia trachomatis* LGV L2 434 Bu reference database (NCBI:txid47472). Cdu1-Flag

MS/MS data was analyzed with Mascot software (Matrix Science) using Trypsin/P specificity for

N-terminal acetylation, lysine acetylation, lysine Ub, and S/T/Y phosphorylation identification.

Analysis identified multiple acetylated and phosphorylated Cdu1 peptides and no Cdu1

ubiquitinated peptides. Data was viewed in Scaffold with Scaffold PTM (Scaffold Software).

Isolation and imaging of *Chlamydia* inclusion extrusions

HeLa cell monolayers (1.2×10^6 cells) were infected with Ct strains at MOIs of 0.8. At 48 hpi,

infected monolayers were imaged using an EVOS FL Cell Imaging System (ThermoFisher

Scientific) equipped with a 20x/0.4 NA objective and a CCD camera. Following imaging, growth

media was removed, cell monolayers washed with fresh growth media, and monolayers

incubated for an additional 4 hours at 37°C. At 52 hpi growth supernatants were collected and

transferred to Eppendorf tubes. Extrusions were enriched by centrifugation (1,500 rpm, 5 min)

and pellets (not always visible) containing extrusions were resuspended in 30 μL of 4%

Formaldehyde/PBS supplemented with Hoechst (2 μg/mL) and 0.2% Trypan Blue Solution

(Gibco, 0.4%). Extrusions were analyzed by plating 10 μL drops on a glass side (without

coverslips) and immediately imaged using an EVOS FL Cell Imaging System (ThermoFisher

Scientific) equipped with a 20x/0.4 NA objective and a CCD camera. Intact extrusions were

identified based on morphology, lacking nuclei, and being impermeable to trypan blue. Images

were opened in ImageJ (Schneider et al., 2012) and enumeration of inclusions and extrusions

was performed manually. The sizes of individual extrusions and inclusions were determined by

manually tracing a line around the perimeter of each extrusion and inclusion in ImageJ and

measuring perimeter length. All measurements were exported to Microsoft Excel. Data plots and

statistical analyses were done with Prism 9 (GraphPad) software. Datasets were analyzed for

significance using a paired student t-test.

Image analysis

Line scan profiles of Cdu1 co-localization with inclusion membrane proteins was performed with

ImageJ (Schneider et al., 2012) by tracing a line through regions of interest and plotting

fluorescent signal intensities with the Plot Profile function. Localization of CTL0480, recruitment

of MYPT1, and association of Ub with Ct inclusions was performed manually from maximum

projections in ImageJ. Assessment of F-actin recruitment to Ct inclusions was performed

manually in ImageJ by projecting four to five sections in order to capture the entire inclusion.

Redistribution of Golgi around the Ct inclusion was measured in ImageJ from maximum

projections. The perimeters of individual inclusions were manually traced, and lengths

measured. The length of dispersed Golgi was measured by tracing and measuring the length of

the GM130 signal directly adjacent to each inclusion. Dispersed Golgi length was divided by

inclusion perimeter length. All measurements were exported to Microsoft Excel for

quantification. Data plots and statistical analyses were done with Prism 9 (GraphPad).

Quantification And Statistical Analysis

Quantifications were generated from three independent experiments and measurements

derived from blinded images. Data plots and statistical analyses were done with Prism 9

(GraphPad) software. Datasets were analyzed for significance using a paired student t-test or

one-way ANOVAs with a Student-Newman-Keuls post hoc test. Data graphs show means and

error bars represent standard error. *p*-values less than 0.05 are defined as statistically

significant. The indicated statistical test for each experiment can be found in the figure legends.

References

1. Abdelrahman YM, Belland RJ. 2005. The chlamydial developmental cycle. *FEMS Microbiol*

Rev **29**:949–959. doi:10.1016/j.femsre.2005.03.002

2. Agaisse H, Derré I. 2013. A *C. trachomatis* cloning vector and the generation of *C.*

trachomatis strains expressing fluorescent proteins under the control of a *C. trachomatis*

promoter. *PLoS ONE* **8**:e57090. doi:10.1371/journal.pone.0057090

3. Albrecht M, Sharma CM, Reinhardt R, Vogel J, Rudel T. 2010. Deep sequencing-based

discovery of the *Chlamydia trachomatis* transcriptome. *Nucleic Acids Res* **38**:868–877.

doi:10.1093/nar/gkp1032

4. Alzhanov DT, Weeks SK, Burnett JR, Rockey DD. 2009. Cytokinesis is blocked in

mammalian cells transfected with *Chlamydia trachomatis* gene CT223. *BMC Microbiol* **9**:2.

doi:10.1186/1471-2180-9-2

5. Auer D, Hügelschäffer SD, Fischer AB, Rudel T. 2020. The chlamydial deubiquitinase Cdu1

supports recruitment of Golgi vesicles to the inclusion. *Cell Microbiol* **22**:e13136.

doi:10.1111/cmi.13136

6. Bannantine JP, Griffiths RS, Viratyosin W, Brown WJ, Rockey DD. 2000. A secondary

structure motif predictive of protein localization to the chlamydial inclusion membrane. *Cell*

Microbiol **2**:35–47. doi:10.1046/j.1462-5822.2000.00029.x

7. Bauler LD, Hackstadt T. 2014. Expression and targeting of secreted proteins from

Chlamydia trachomatis. *J Bacteriol* **196**:1325–1334. doi:10.1128/JB.01290-13

8. Berrow NS, Alderton D, Sainsbury S, Nettleship J, Assenberg R, Rahman N, Stuart DI,

Owens RJ. 2007. A versatile ligation-independent cloning method suitable for high-

throughput expression screening applications. *Nucleic Acids Res* **35**:e45.

doi:10.1093/nar/gkm047

9. Bremm A, Freund SMV, Komander D. 2010. Lys11-linked ubiquitin chains adopt compact

conformations and are preferentially hydrolyzed by the deubiquitinase Cezanne. *Nat Struct*

Mol Biol **17**:939–947. doi:10.1038/nsmb.1873

10. Bugalhão JN, Mota LJ. 2019. The multiple functions of the numerous *Chlamydia trachomatis*

secreted proteins: the tip of the iceberg. *Microb Cell* **6**:414–449.

doi:10.15698/mic2019.09.691

11. Catic A, Misaghi S, Korbel GA, Ploegh HL. 2007. ElaD, a Deubiquitinating protease

expressed by *E. coli*. *PLoS ONE* **2**:e381. doi:10.1371/journal.pone.0000381

12. Chen C, Chen D, Sharma J, Cheng W, Zhong Y, Liu K, Jensen J, Shain R, Arulanandam B,

Zhong G. 2006. The hypothetical protein CT813 is localized in the *Chlamydia trachomatis*

inclusion membrane and is immunogenic in women urogenitally infected with *C.*

trachomatis. Infect Immun **74**:4826–4840. doi:10.1128/IAI.00081-06

13. Chen Y-S, Bastidas RJ, Saka HA, Carpenter VK, Richards KL, Plano GV, Valdivia RH.

2014. The *Chlamydia trachomatis* type III secretion chaperone Slc1 engages multiple early

effectors, including TepP, a tyrosine-phosphorylated protein required for the recruitment of

CrkI-II to nascent inclusions and innate immune signaling. *PLoS Pathog* **10**:e1003954.

doi:10.1371/journal.ppat.1003954

14. Chin E, Kirker K, Zuck M, James G, Hybiske K. 2012. Actin recruitment to the *Chlamydia*

inclusion is spatiotemporally regulated by a mechanism that requires host and bacterial

factors. *PLoS ONE* **7**:e46949. doi:10.1371/journal.pone.0046949

15. Claessen JHL, Witte MD, Yoder NC, Zhu AY, Spooner E, Ploegh HL. 2013. Catch-and-

release probes applied to semi-intact cells reveal ubiquitin-specific protease expression in

Chlamydia trachomatis infection. *Chembiochem* **14**:343–352. doi:10.1002/cbic.201200701

16. Datta AB, Hura GL, Wolberger C. 2009. The structure and conformation of Lys63-linked

tetraubiquitin. *J Mol Biol* **392**:1117–1124. doi:10.1016/j.jmb.2009.07.090

17. Dehoux P, Flores R, Dauga C, Zhong G, Subtil A. 2011. Multi-genome identification and

characterization of chlamydiae-specific type III secretion substrates: the Inc proteins. *BMC*

Genomics **12**:109. doi:10.1186/1471-2164-12-109

18. Dumoux M, Menny A, Delacour D, Hayward RD. 2015. A *Chlamydia* effector recruits

CEP170 to reprogram host microtubule organization. *J Cell Sci* **128**:3420–3434.

doi:10.1242/jcs.169318

19. Eddins MJ, Varadan R, Fushman D, Pickart CM, Wolberger C. 2007. Crystal structure and

solution NMR studies of Lys48-linked tetraubiquitin at neutral pH. *J Mol Biol* **367**:204–211.

doi:10.1016/j.jmb.2006.12.065

20. Fischer A, Harrison KS, Ramirez Y, Auer D, Chowdhury SR, Prusty BK, Sauer F, Dimond Z,

Kisker C, Hefty PS, Rudel T. 2017. *Chlamydia trachomatis*-containing vacuole serves as

deubiquitination platform to stabilize Mcl-1 and to interfere with host defense. *eLife* **6**.

doi:10.7554/eLife.21465

21. Fiskin E, Bionda T, Dikic I, Behrends C. 2016. Global analysis of host and bacterial

ubiquitinome in response to *Salmonella* Typhimurium infection. *Mol Cell* **62**:967–981.

doi:10.1016/j.molcel.2016.04.015

22. Furtado AR, Essid M, Perrinet S, Balañá ME, Yoder N, Dehoux P, Subtil A. 2013. The

chlamydial OTU domain-containing protein *Chla*OTU is an early type III secretion effector

targeting ubiquitin and NDP52. *Cell Microbiol* **15**:2064–2079. doi:10.1111/cmi.12171

23. Gehre L, Gorgette O, Perrinet S, Prevost M-C, Ducatez M, Giebel AM, Nelson DE, Ball SG,

Subtil A. 2016. Sequestration of host metabolism by an intracellular

pathogen. *eLife* **5**:e12552. doi:10.7554/eLife.12552

24. Goedhart J, Luijsterburg MS. 2020. VolcaNoseR is a web app for creating, exploring,

labeling and sharing volcano plots. *Sci Rep* **10**:20560. doi:10.1038/s41598-020-76603-3

25. Hackstadt T, Scidmore-Carlson MA, Shaw EI, Fischer ER. 1999. The *Chlamydia*

trachomatis IncA protein is required for homotypic vesicle fusion. *Cell Microbiol* **1**:119–130.

doi:10.1046/j.1462-5822.1999.00012.x

26. Haggerty CL, Gottlieb SL, Taylor BD, Low N, Xu F, Ness RB. 2010. Risk of sequelae after

Chlamydia trachomatis genital infection in women. *J Infect Dis* **201 Suppl 2**:S134-55.

doi:10.1086/652395

27. Haines A, Wesolowski J, Ryan NM, Monteiro-Brás T, Paumet F. 2021. Cross Talk between

ARF1 and RhoA Coordinates the Formation of Cytoskeletal Scaffolds during *Chlamydia*

Infection. MBio **12**:e0239721. doi:10.1128/mBio.02397-21

28. Hausman JM, Kenny S, Iyer S, Babar A, Qiu J, Fu J, Luo Z-Q, Das C. 2020. The Two

Deubiquitinating Enzymes from *Chlamydia trachomatis* Have Distinct Ubiquitin Recognition

Properties. *Biochemistry* **59**:1604–1617. doi:10.1021/acs.biochem.9b01107

29. Huang DW, Sherman BT, Lempicki RA. 2009a. Bioinformatics enrichment tools: paths

toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**:1–

13. doi:10.1093/nar/gkn923

30. Huang DW, Sherman BT, Lempicki RA. 2009b. Systematic and integrative analysis of large

gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**:44–57.

doi:10.1038/nprot.2008.211

31. Hybiske K, Stephens RS. 2007. Mechanisms of host cell exit by the intracellular bacterium

Chlamydia. Proc Natl Acad Sci USA **104**:11430–11435. doi:10.1073/pnas.0703218104

32. Iyer S, Das C. 2021. The unity of opposites: Strategic interplay between bacterial effectors

to regulate cellular homeostasis. *J Biol Chem* **297**:101340. doi:10.1016/j.jbc.2021.101340

33. Jeong KC, Sexton JA, Vogel JP. 2015. Spatiotemporal regulation of a *Legionella*

pneumophila T4SS substrate by the metaeffector SidJ. *PLoS Pathog* **11**:e1004695.

doi:10.1371/journal.ppat.1004695

34. Jones RM, Wu H, Wentworth C, Luo L, Collier-Hyams L, Neish AS. 2008. *Salmonella avra*

coordinates suppression of host immune and apoptotic defenses via JNK pathway

blockade. *Cell Host Microbe* **3**:233–244. doi:10.1016/j.chom.2008.02.016

35. Kędzior M, Bastidas RJ. 2019. Forward and reverse genetic analysis of *Chlamydia. Methods*

Mol Biol **2042**:185–204. doi:10.1007/978-1-4939-9694-0_13

1001

1002 36. Kokes M, Dunn JD, Granek JA, Nguyen BD, Barker JR, Valdivia RH, Bastidas RJ. 2015.

1003 Integrating chemical mutagenesis and whole-genome sequencing as a platform for forward

1004 and reverse genetic analysis of *Chlamydia. Cell Host Microbe* **17**:716–725.

1005 doi:10.1016/j.chom.2015.03.014

1006

1007 37. Komander D, Clague MJ, Urbé S. 2009a. Breaking the chains: structure and function of the

1008 deubiquitinases. *Nat Rev Mol Cell Biol* **10**:550–563. doi:10.1038/nrm2731

1009

1010 38. Komander D, Reyes-Turcu F, Licchesi JDF, Odenwaelder P, Wilkinson KD, Barford D.

1011   2009b. Molecular discrimination of structurally equivalent Lys 63-linked and linear

1012   polyubiquitin chains. *EMBO Rep* **10**:466–473. doi:10.1038/embor.2009.55

1013

1014 39. Komander D, Rape M. 2012. The ubiquitin code. *Annu Rev Biochem* **81**:203–229.

1015   doi:10.1146/annurev-biochem-060310-170328

1016

1017 40. Kubori T, Shinzawa N, Kanuka H, Nagai H. 2010. *Legionella* metaeffector exploits host

1018   proteasome to temporally regulate cognate effector. *PLoS Pathog* **6**:e1001216.

1019   doi:10.1371/journal.ppat.1001216

1020

1021 41. Kubori T, Kitao T, Nagai H. 2019. Emerging insights into bacterial deubiquitinases. *Curr*

1022   *Opin Microbiol* **47**:14–19. doi:10.1016/j.mib.2018.10.001

1023

1024 42. Kunz TC, Götz R, Sauer M, Rudel T. 2019. Detection of *Chlamydia* developmental forms

1025   and secreted effectors by expansion microscopy. *Front Cell Infect Microbiol* **9**:276.

1026   doi:10.3389/fcimb.2019.00276

1027

1028 43. Lee JK, Enciso GA, Boassa D, Chander CN, Lou TH, Pairawan SS, Guo MC, Wan FYM,

1029   Ellisman MH, Sütterlin C, Tan M. 2018. Replication-dependent size reduction precedes

1030   differentiation in *Chlamydia trachomatis*. *Nat Commun* **9**:45. doi:10.1038/s41467-017-

1031   02432-0

1032

1033 44. Li J, Chai Q-Y, Liu CH. 2016. The ubiquitin system: a critical regulator of innate immunity

1034   and pathogen-host interactions. *Cell Mol Immunol* **13**:560–576. doi:10.1038/cmi.2016.40

1035

1036   45. Li Z, Chen C, Chen D, Wu Y, Zhong Y, Zhong G. 2008. Characterization of fifty putative

1037   inclusion membrane proteins encoded in the *Chlamydia trachomatis* genome. *Infect*

1038   *Immun* **76**:2746–2757. doi:10.1128/IAI.00010-08

1039

1040   46. Lowden NM, Yeruva L, Johnson CM, Bowlin AK, Fisher DJ. 2015. Use of aminoglycoside 3'

1041   adenyltransferase as a selection marker for *Chlamydia trachomatis* intron-mutagenesis and

1042   in vivo intron stability. *BMC Res Notes* **8**:570. doi:10.1186/s13104-015-1542-9

1043

1044   47. Lutter EI, Martens C, Hackstadt T. 2012. Evolution and conservation of predicted inclusion

1045   membrane proteins in *Chlamydiae. Comp Funct Genomics* **2012**:362104.

1046   doi:10.1155/2012/362104

1047

1048   48. Lutter EI, Barger AC, Nair V, Hackstadt T. 2013. *Chlamydia trachomatis* inclusion

1049   membrane protein CT228 recruits elements of the myosin phosphatase pathway to regulate

1050   release mechanisms. *Cell Rep* **3**:1921–1931. doi:10.1016/j.celrep.2013.04.027

1051

1052   49. Makarova KS, Aravind L, Koonin EV. 2000. A novel superfamily of predicted cysteine

1053   proteases from eukaryotes, viruses and *Chlamydia pneumoniae. Trends Biochem*

1054   *Sci* **25**:50–52. doi:10.1016/s0968-0004(99)01530-3

1055

1056   50. Meier K, Jachmann LH, Pérez L, Kepp O, Valdivia RH, Kroemer G, Sixt BS. 2022.

1057   The *Chlamydia* protein CpoS modulates the inclusion microenvironment and restricts the

1058   interferon response by acting on Rab35. *BioRxiv*. doi:10.1101/2022.02.18.481055

1059

1060   51. Misaghi S, Balsara ZR, Catic A, Spooner E, Ploegh HL, Starnbach MN. 2006. *Chlamydia*

1061   *trachomatis*-derived deubiquitinating enzymes in mammalian cells during infection. *Mol*

1062   *Microbiol* **61**:142–150. doi:10.1111/j.1365-2958.2006.05199.x

1063

1064   52. Mittal R, Peak-Chew S-Y, McMahon HT. 2006. Acetylation of MEK2 and IκB kinase (IKK)

1065   activation loop residues by YopJ inhibits signaling. *Proc Natl Acad Sci USA* **103**:18574–

1066   18579. doi:10.1073/pnas.0608995103

1067

1068   53. Mital J, Miller NJ, Fischer ER, Hackstadt T. 2010. Specific chlamydial inclusion membrane

1069   proteins associate with active Src family kinases in microdomains that interact with the host

1070   microtubule network. *Cell Microbiol***12**:1235–1249. doi:10.1111/j.1462-5822.2010.01465.x

1071

1072   54. Mizuno E, Kitamura N, Komada M. 2007. 14-3-3-dependent inhibition of the deubiquitinating

1073   activity of UBPY and its cancellation in the M phase. *Exp Cell Res* **313**:3624–3634.

1074   doi:10.1016/j.yexcr.2007.07.028

1075

1076   55. Moulder JW. 1991. Interaction of *Chlamydiae* and host cells in vitro. *Microbiol Rev* **55**:143–

1077   190.

1078

1079   56. Mukherjee S, Keitany G, Li Y, Wang Y, Ball HL, Goldsmith EJ, Orth K. 2006. *Yersinia* YopJ

1080   acetylates and inhibits kinase activation by blocking phosphorylation. *Science* **312**:1211–

1081   1214. doi:10.1126/science.1126867

1082

1083   57. Neunuebel MR, Chen Y, Gaspar AH, Backlund PS, Yergey A, Machner MP. 2011. De-

1084   AMPylation of the small GTPase Rab1 by the pathogen *Legionella*

1085   *pneumophila. Science* **333**:453–456. doi:10.1126/science.1207193

1086

1087   58. Nguyen BD, Valdivia RH. 2012. Virulence determinants in the obligate intracellular pathogen

1088   *Chlamydia trachomatis* revealed by forward genetic approaches. *Proc Natl Acad Sci*

1089   *USA* **109**:1263–1268. doi:10.1073/pnas.1117884109

1090

1091   59. Nguyen PH, Lutter EI, Hackstadt T. 2018. *Chlamydia trachomatis* inclusion membrane

1092   protein MrcA interacts with the inositol 1,4,5-trisphosphate receptor type 3 (ITPR3) to

1093   regulate extrusion formation. *PLoS Pathog***14**:e1006911. doi:10.1371/journal.ppat.1006911

1094

1095   60. Ohtake F, Tsuchiya H. 2017. The emerging complexity of ubiquitin architecture. *J*

1096   *Biochem* **161**:125–133. doi:10.1093/jb/mvw088

1097

1098   61. Pannekoek Y, Spaargaren J, Langerak AAJ, Merks J, Morré SA, van der Ende A. 2005.

1099   Interrelationship between polymorphisms of *incA*, fusogenic properties of *Chlamydia*

1100   *trachomatis* strains, and clinical manifestations in patients in The Netherlands. *J Clin*

1101   *Microbiol* **43**:2441–2443. doi:10.1128/JCM.43.5.2441-2443.2005

1102

1103   62. Peng J, Schwartz D, Elias JE, Thoreen CC, Cheng D, Marsischky G, Roelofs J, Finley D,

1104   Gygi SP. 2003. A proteomics approach to understanding protein ubiquitination. *Nat*

1105   *Biotechnol* **21**:921–926. doi:10.1038/nbt849

1106

1107   63. Pruneda JN, Durkin CH, Geurink PP, Ovaa H, Santhanam B, Holden DW, Komander D.

1108   2016. The Molecular Basis for Ubiquitin and Ubiquitin-like Specificities in Bacterial Effector

1109   Proteases. *Mol Cell* **63**:261–276. doi:10.1016/j.molcel.2016.06.015

1110

1111 64. Pruneda JN, Bastidas RJ, Bertsoulaki E, Swatek KN, Santhanam B, Clague MJ, Valdivia

1112   RH, Urbé S, Komander D. 2018. A *Chlamydia* effector combining deubiquitination and

1113   acetylation activities induces Golgi fragmentation. *Nat Microbiol* **3**:1377–1384.

1114   doi:10.1038/s41564-018-0271-y

1115

1116 65. Reiley W, Zhang M, Wu X, Granger E, Sun S-C. 2005. Regulation of the deubiquitinating

1117   enzyme CYLD by IκB kinase gamma-dependent phosphorylation. *Mol Cell Biol* **25**:3886–

1118   3895. doi:10.1128/MCB.25.10.3886-3895.2005

1119

1120 66. Roan NR, Gierahn TM, Higgins DE, Starnbach MN. 2006. Monitoring the T cell response to

1121   genital tract infection. *Proc Natl Acad Sci USA* **103**:12069–12074.

1122   doi:10.1073/pnas.0603866103

1123

1124 67. Rockey DD, Scidmore MA, Bannantine JP, Brown WJ. 2002. Proteins in the chlamydial

1125   inclusion membrane. *Microbes Infect* **4**:333–340. doi:10.1016/s1286-4579(02)01546-0

1126

1127 68. Ronau JA, Beckmann JF, Hochstrasser M. 2016. Substrate specificity of the ubiquitin and

1128   Ubl proteases. *Cell Res* **26**:441–456. doi:10.1038/cr.2016.38

1129

1130 69. Rytkönen A, Poh J, Garmendia J, Boyle C, Thompson A, Liu M, Freemont P, Hinton JCD,

1131   Holden DW. 2007. SseL, a *Salmonella* deubiquitinase required for macrophage killing and

1132   virulence. *Proc Natl Acad Sci USA***104**:3502–3507. doi:10.1073/pnas.0610095104

1133

1134 70. Saeki Y. 2017. Ubiquitin recognition by the proteasome. *J Biochem* **161**:113–124.

1135   doi:10.1093/jb/mvw091

1136

1137   71. Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image

1138   analysis. *Nat Methods* **9**:671–675. doi:10.1038/nmeth.2089

1139

1140   72. Schubert AF, Nguyen JV, Franklin TG, Geurink PP, Roberts CG, Sanderson DJ, Miller LN,

1141   Ovaa H, Hofmann K, Pruneda JN, Komander D. 2020. Identification and characterization of

1142   diverse OTU deubiquitinases in bacteria. *EMBO J* **39**:e105127.

1143   doi:10.15252/embj.2020105127

1144

1145   73. Shaw JH, Key CE, Snider TA, Sah P, Shaw EI, Fisher DJ, Lutter EI. 2018. Genetic

1146   Inactivation of *Chlamydia trachomatis* Inclusion Membrane Protein CT228 Alters MYPT1

1147   Recruitment, Extrusion Production, and Longevity of Infection. *Front Cell Infect*

1148   *Microbiol* **8**:415. doi:10.3389/fcimb.2018.00415

1149

1150   74. Sheedlo MJ, Qiu J, Tan Y, Paul LN, Luo Z-Q, Das C. 2015. Structural basis of substrate

1151   recognition by a bacterial deubiquitinase important for dynamics of phagosome

1152   ubiquitination. *Proc Natl Acad Sci USA* **112**:15090–15095. doi:10.1073/pnas.1514568112

1153

1154   75. Sixt BS, Bastidas RJ, Finethy R, Baxter RM, Carpenter VK, Kroemer G, Coers J, Valdivia

1155   RH. 2017. The *Chlamydia trachomatis* Inclusion Membrane Protein CpoS Counteracts

1156   STING-Mediated Cellular Surveillance and Suicide Programs. *Cell Host Microbe* **21**:113–

1157   121. doi:10.1016/j.chom.2016.12.002

1158

1159   76. Smith EP, Cotto-Rosario A, Borghesan E, Held K, Miller CN, Celli J. 2020. Epistatic Interplay

1160   between Type IV Secretion Effectors Engages the Small GTPase Rab2 in the *Brucella*

1161   Intracellular Cycle. *MBio* **11**. doi:10.1128/mBio.03350-19

1162

1163   77. Suchland RJ, Rockey DD, Bannantine JP, Stamm WE. 2000. Isolates of *Chlamydia*

1164   *trachomatis* that occupy nonfusogenic inclusions lack IncA, a protein localized to the

1165   inclusion membrane. *Infect Immun* **68**:360–367. doi:10.1128/IAI.68.1.360-367.2000

1166

1167   78. Swatek KN, Komander D. 2016. Ubiquitin modifications. *Cell Res* **26**:399–422.

1168   doi:10.1038/cr.2016.39

1169

1170   79. Tenno T, Fujiwara K, Tochio H, Iwai K, Morita EH, Hayashi H, Murata S, Hiroaki H, Sato M,

1171   Tanaka K, Shirakawa M. 2004. Structural basis for distinct roles of Lys63- and Lys48-linked

1172   polyubiquitin chains. *Genes Cells***9**:865–875. doi:10.1111/j.1365-2443.2004.00780.x

1173

1174   80. Urbanus ML, Quaile AT, Stogios PJ, Morar M, Rao C, Di Leo R, Evdokimova E, Lam M,

1175   Oatway C, Cuff ME, Osipiuk J, Michalska K, Nocek BP, Taipale M, Savchenko A, Ensminger

1176   AW. 2016. Diverse mechanisms of metaeffector activity in an intracellular bacterial

1177   pathogen, *Legionella pneumophila. Mol Syst Biol* **12**:893. doi:10.15252/msb.20167381

1178

1179   81. Varadan R, Walker O, Pickart C, Fushman D. 2002. Structural properties of polyubiquitin

1180   chains in solution. *J Mol Biol* **324**:637–647. doi:10.1016/s0022-2836(02)01198-1

1181

1182   82. Vozandychova V, Stojkova P, Hercik K, Rehulka P, Stulik J. 2021. The Ubiquitination

1183   System within Bacterial Host-Pathogen Interactions. *Microorganisms* **9**.

1184   doi:10.3390/microorganisms9030638

1185

1186   83. Wang X, Hybiske K, Stephens RS. 2018. Direct visualization of the expression and

1187   localization of chlamydial effector proteins within infected host cells. *Pathog Dis* **76**.

1188   doi:10.1093/femspd/fty011

1189

1190   84. Weber MM, Bauler LD, Lam J, Hackstadt T. 2015. Expression and localization of predicted

1191   inclusion membrane proteins in *Chlamydia trachomatis*. *Infect Immun* **83**:4710–4718.

1192   doi:10.1128/IAI.01075-15

1193

1194   85. Weeks SD, Grasty KC, Hernandez-Cuebas L, Loll PJ. 2009. Crystal structures of Lys-63-

1195   linked tri- and di-ubiquitin reveal a highly extended chain architecture. *Proteins* **77**:753–759.

1196   doi:10.1002/prot.22568

1197

1198   86. Wesolowski J, Weber MM, Nawrotek A, Dooley CA, Calderon M, St Croix CM, Hackstadt T,

1199   Cherfils J, Paumet F. 2017. *Chlamydia* hijacks ARF gtpases to coordinate microtubule

1200   posttranslational modifications and golgi complex positioning. *MBio* **8**.

1201   doi:10.1128/mBio.02280-16

1202

1203   87. Zadora PK, Chumduri C, Imami K, Berger H, Mi Y, Selbach M, Meyer TF, Gurumurthy RK.

1204   2019. Integrated Phosphoproteome and Transcriptome Analysis Reveals *Chlamydia*-

1205   Induced Epithelial-to-Mesenchymal Transition in Host Cells. *Cell Rep* **26**:1286-1302.e8.

1206   doi:10.1016/j.celrep.2019.01.006

1207

1208   88. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C, Chanda

1209   SK. 2019. Metascape provides a biologist-oriented resource for the analysis of systems-

1210   level datasets. *Nat Commun* **10**:1523. doi:10.1038/s41467-019-09234-6

1211

1212   89. Zuck M, Ellis T, Venida A, Hybiske K. 2017. Extrusions are phagocytosed and promote

1213   *Chlamydia* survival within macrophages. *Cell Microbiol* **19**. doi:10.1111/cmi.12683

1214

1215   90. Zuck M, Sherrid A, Suchland R, Ellis T, Hybiske K. 2016. Conservation of extrusion as an

1216   exit mechanism for *Chlamydia*. *Pathog Dis* **74**. doi:10.1093/femspd/ftw093

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238   **Figures**

1239
1240
Figure 1
The *C. trachomatis* inclusion membrane proteins InaC, IpaM, and CTL0480 are
ubiquitinated in the absence of Cdu1. (A-C) Volcano plots (pairwise comparisons) of the
relative abundance of human and Ct ubiquitinated peptides between mock infected HeLa cells
versus HeLa cells infected with L2 (24 hpi) **(A),** mock infected HeLa cells versus HeLa cells
infected with a *cdu1* null strain (*cdu1*::GII) (24 hpi) **(B),** and HeLa cells infected with L2 (24 hpi)
versus HeLa cells infected with a *cdu1* null strain (24 hpi) **(C)**. All infections were performed for
24 hours. Significance values were interpolated from 3 independent biological replicates.
Ubiquitinated proteins were enriched with magnetic TUBE 1 beads (binds to polyubiquitinated
proteins) and peptides identified by quantitative LC MS/MS analysis. Three Ct proteins, InaC,
IpaM, and CTL0480 were differentially ubiquitinated in the absence of Cdu1. **(D)**
InaC was ubiquitinated at K104, K107, and K149, IpaM at K290, and CTL0480 at K115 in the
absence of Cdu1. TMD: Transmembrane domain. Numbering corresponds to amino acids in the
protein sequence of each respective inclusion membrane protein.
1255

 Figure 2
Cdu1 associates with InaC, IpaM, and CTL0480. (A) Co-localization of Cdu1(magenta) with endogenous InaC (green), IpaM (green), and ectopically expressed CTL0480-Flag (green) at the L2 inclusion membrane during infection of HeLa cells at 24 hpi. HeLa cells infected with an *inaC* null strain (M407), an *ipaM* null strain (*ipaM*::GII), and L2 pBOMB were used as antibody specificity controls. DNA stained with Hoechst is shown in blue. Scale bar: 10μm. Images are representative of multiple images captured across three independent replicates. **(B)** Line scan profiles of fluorescent signal intensities displayed in (A) showing co-localization of fluorescence intensities for endogenous Cdu1 with endogenous InaC, and IpaM and with CTL0480-Flag

along the L2 inclusion membrane. **(C)** Schematic of Cdu1-GFP(C) (L2) fusion (Cdu1-GFP) and
Cdu1-GFP variants used in co-transfections of HEK 293 cells. GFP: Green fluorescent protein.
TMD: Transmembrane domain. PRD: Proline rich domain. CD: Catalytic domain. FL: Full length.
TMD-: Cdu1-GFP variant lacking TMD domain. CD-: Cdu1-GFP variant lacking CD domain. **(D-**
G) Western blot analysis of GFP immunoprecipitates from HEK 293 cells co-transfected with
mammalian plasmids expressing: Cdu1-GFP variants and **(D)** truncated 3XFlag(N)-InaC
(D/UW-3/CX CT813, amino acids 96-264), **(E)** V5(N)-IpaM (L2, full length), **(F)** V5(N)-CTL0480
(L2, full length), and **(G)** V5(N)-CpoS (L2, full length). Vec1: Empty pOPINN-GFP vector. Vec2:
Empty pDEST53 vector. Vec3: Empty pcDNA^{TM}3.1/nV5-DEST vector. Western blot images are
representative from two independent experiments.

1290
1291
Figure 3
Cdu1 stabilizes InaC, IpaM, and CTL0480. (A) Western blot analysis of InaC in HeLa cells
infected for 24 hours with L2, *inaC* null, and *cdu1* null strains. Ct Slc1 and human alpha tubulin
were used to determine Ct burdens and equal loading of protein extracts respectively. Western
blot images are representative of 3 independent experiments. **(B)** Western blot analysis of
endogenous IpaM in HeLa cells infected with L2, *cdu1* null, and *ipaM* null strains for 24, 36, and
48 hours. Western blot images are representative of 2 independent experiments. **(C)**
Localization of CTL0480 during Ct infection of HeLa cells at 24, 36, and 48 hpi. CTL0480 signal
(green) co-localizes with the inclusion membrane protein IncA (magenta) at the L2 inclusion
membrane. Arrowheads highlight *cdu1* null inclusions lacking CTL0480 at 36 hpi. DNA stained
with Hoechst is shown in blue. Scale bar: 10μm. **(D)** Quantification of CTL0480 localization at Ct
inclusion membranes as shown in (C). CTL0480 fluorescent signal was used to determine the
percent of inclusions displaying CTL0480 at the inclusion membranes. Representative images
and quantification of CTL0480 positive inclusions are derived from Ct inclusions imaged in 10
fields across 3 independent biological replicates. *p* values were determined by a student paired
t-test.
1308

1309
1310
Figure 4
The acetylase activity of Cdu1 is required to stabilize Cdu1, InaC, and IpaM. (A) Western
blot analysis of endogenous InaC and Cdu1-Flag catalytic variants expressed from a plasmid.
HeLa cells were infected for 36 hours with L2, a *cdu1* null strain, and *cdu1* null strains
expressing wild type Cdu1-Flag and the Cdu1 variants C345A-Flag (catalytic inactive), I225A-

Flag (DUB deficient), and K268E-Flag (Act deficient). Protein lysates from HeLa cells infected
with M407 (*inaC* null) were used to control for the specificity of anti-InaC antibodies. Western
blot images are representative of two independent experiments. **(B)** Western blot analysis of
endogenous IpaM and Cdu1-FLAG variants in crude extracts of HeLa cells infected for 48 hours
with the same strains as describe in (A). Infection of HeLa cells with *ipaM*::GII was used to test
for the specificity of the anti-IpaM antibody. Western blot images are representative of two
independent experiments. **(C)** Western blot analysis of Cdu1^{C345A}-Flag (catalytic inactive) and
Cdu1^{K268E}-Flag (Act deficient) expressed in a *cdu1* null strain or wild type L2 background after
infection of HeLa cells for 24 hours. Both Cdu1 variants are stabilized by endogenous Cdu1. **(D)**
Western blot analysis of Cdu1-Flag and Cdu1^{C345A}-Flag expressed in a *cdu1* null strain and
Cdu1^{C345A}-Flag expressed in wild type L2 following immunoprecipitation (anti-Flag) from HeLa
cell extracts after infection for 24 hours and treatment with MG132 (25 μM, 5 hours). Western
blot image is a representative blot from at least three independent experiments. **(E)** Western
blot analysis of endogenous InaC and IpaM following immunoprecipitation of proteins acetylated
at lysines from inclusion membrane subcellular fractions derived from HeLa cells infected with
WT L2 for 24 hours. Western blot image is representative of two independent experiments. **(F)**
Western blot analysis (Flag WB) of acetylated lysine immunoprecipitates generated from
inclusion membrane subcellular fractions derived from HeLa cells infected with WT L2 strains
expressing Cdu1-Flag, CTL0480-Flag, and IpaM-Flag at 40 hpi. Western blot image is
representative of two independent experiments. **(G)** Flag WB of acetylated lysine
immunoprecipitates generated from inclusion membrane subcellular fractions derived from
HeLa cells infected with Rif-R L2 (M407 parental strain) and with M407 (*inaC* null) expressing
InaC-Flag at 24 hpi. Western blot image is representative of two independent experiments. **(H)**
The initiator methionine, Lys10, Lys126, and Lys263 of Cdu1 are acetylated during Ct infection
of HeLa cells (24 hpi). One tyrosine (Y) residue and multiple serine (S) and threonine (T)
residues in Cdu1 are also phosphorylated during Ct infection of HeLa cells (24 hpi). Three
PX(S/T)P MAPK phosphorylation consensus sequence motifs were identified in the proline rich
domain of Cdu1. Modified residues were identified by quantitative LC MS/MS analysis of
immunoprecipitated Cdu1-Flag across 3 independent biological replicates. TMD:
Transmembrane domain. PR: Proline rich. PSTP: PX(S/T)P motifs. Ac: Acetylation. P:
Phosphorylation. Numbering corresponds to amino acids in Cdu1 protein sequence.
1347

Figure 5

The DUB activity of Cdu1 is not required for blocking the ubiquitination of inclusion

membranes. (A) Loss of Cdu1's acetylase activity results in the accumulation of Ub (magenta

signal) at the periphery of the inclusion. HeLa cells were infected for 24 hours. Antisera against

the membrane protein Cap1 (green) was used to mark inclusion membranes. DNA stained with

Hoechst is shown in blue. Scale bar: 10μm. **(B)** Quantification of ubiquitinated inclusions as

shown in (A). The Ub fluorescent signal was used to determine the number of infected cells with

Ub decorated inclusions. The total number of ubiquitinated inclusions was divided by the total
number of inclusions analyzed (defined by Cap1 staining). 87%, 86%, and 91% of inclusions
were decorated with Ub in HeLa cells infected with a *cdu1* null strain and *cdu1* null strains
expressing Cdu1^{C345A} (DUB-, Act-), and Cdu1^{K268E} (Act-) variants respectively. Representative
images in (A) and quantification of ubiquitinated inclusions in (B) were obtained from inclusions
imaged in 10 fields across 3 independent biological replicates for each strain. *p* values were
determined by a student paired t-test. **(C)** HeLa cells co-infected with *cdu1*::GII (*cdu1Δ*) and
incA null (*incAΔ*, M923) strains at 24 hpi. IncA-deficient inclusions do not fuse with other
inclusions. In co-infected cells, Cdu1 present on the inclusion membranes of *incAΔ* strains did
not block ubiquitination events at or near the inclusion membranes of neighboring *cdu1Δ* strains
(mCherry signal from pBOMB4-MCI plasmid). DNA stained with Hoechst is shown in blue. Scale
bar: 10μm. **(D)** Quantification of *cdu1Δ* inclusions as shown in (C) in which ubiquitination events
are observed in regions of *cdu1Δ* inclusion membranes that are in direct apposition to *incAΔ*
inclusion membranes (junctions). The total number of *cdu1Δ* inclusions ubiquitinated at inclusion
junctions was divided by the total number of inclusions analyzed (Cap1 staining). 66% and 61%
of *cdu1Δ* inclusions were decorated with Ub at junctions in HeLa cells co-infected with *incAΔ*
and *cdu1Δ* strains or with *incAΔ* and a *cdu1Δ* strain ectopically expressing Cdu1^{K268E} (Act-)
respectively. Representative images in (C) and quantification of *cdu1Δ* inclusions ubiquitinated
at junctions in (D) are derived from inclusions imaged in 6 fields across 3 independent biological
replicates for each condition. *p* values were determined by a student paired t-test.
1377

1378
1379
1380 **Figure 6**

The DUB activity of Cdu1 is not required for assembly of F-actin or Golgi ministack repositioning around the Ct inclusion while Cdu1 is required for MYPT1 recruitment to the inclusion.
Representative images of F-actin (green, Alexa Fluor™ Phalloidin) assembled around the Ct inclusion (magenta, anti Cdu1 and Cap1 staining) in HeLa cells infected for 40 hours. **(B)** Representative images of Golgi (anti GM130 staining, magenta) around Ct inclusions (arrowheads) in HeLa cells infected for 24 hours. HeLa cells were infected with an *inaC* null strain transformed with empty p2TK2 vector (M407 p2TK2), the wild type strain in which M407 was derived from (Rif-R L2) and M407 ectopically expressing full length InaC (M407 p2TK2-InaC). HeLa cells were also infected with L2 pBOMB, *cdu1*::GII pBOMB, and *cdu1* null strains ectopically expressing Cdu1-Flag (Cdu1), and the Cdu1 variants C345A-Flag (DUB-, Act-), I225A-Flag (DUB-), and K268E-Flag (Act-). **(C)** Representative images of MYPT1 (magenta) at Ct inclusions (green, anti-IncA staining). Arrowheads represent *cdu1* null inclusions with low MYPT1 signal. DNA stained with Hoechst is shown in blue in panels A-C. Scale bar: 10μm in panels A-C. **(D)** Quantification of Golgi dispersal around the Ct inclusion as shown in (B). The length of Golgi dispersed around each Ct inclusion imaged was measured and normalized to the perimeter length of each inclusion (% Golgi dispersion around the inclusion). Representative images in (B) and Golgi dispersal quantified in (D) were performed from inclusions imaged in 6 fields across 3 independent biological replicates. *p* values were determined by a student paired t-test. **(E)** Quantification of Ct inclusion surrounded by F-actin as shown in (A) normalized to total inclusions analyzed. Representative images in (A) and quantification of F-actin surrounding Ct inclusions in (E) were obtained from inclusions imaged in 6 fields across 3 independent biological replicates. *p* values were determined by one-way ANOVAs with a Student-Newman-Keuls post hoc test. **(F)** Quantification of MYPT1 at Ct inclusions as shown in (C). Representative images in (C) and quantification of MYPT1 recruitment in (F) were obtained from inclusions imaged in 6 fields across 3 independent replicates. *p* values were determined by a student paired t-test.

1409
Figure 7
Cdu1, InaC, and IpaM are required for optimal extrusion of Ct inclusions from HeLa cells.
(A) Representative images of extrusions isolated from HeLa cell monolayers infected with Ct
strains for 52 hours. Scale bar: 200 μm **(B)** Quantification of the number of extruded inclusions
produced by infected HeLa cell monolayers. *p* values were determined by a student paired t-
test. **(C)** Quantification of the size of extruded inclusions quantified in (B). Extruded inclusions
varied in size among cells infected with wild type L2 (average :43 μm), *ipaM* mutants (average:
56 μm) and *cdu1* mutants complemented wild type Cdu1 (average: 52 μm) and Cdu1^{I225A}
(DUB-) (average: 60 μm). *p* values were determined by one-way ANOVAs with a Student-
Newman-Keuls post hoc test. Representative images in (A) and quantification of extrusion
number in (B) and extrusion size in (C) are based on images obtained from 6 fields across 3
independent biological replicates.
1422

1423
1424
Figure 8.
A model for acetylation mediated protection of the translocated Ct effectors InaC, IpaM,
and CTL0480 from degradation.
The human cellular Ub machinery targets *C. trachomatis* effectors, including the inclusion
membrane proteins InaC, IpaM, and CTL0480 for ubiquitination and subsequent protein
degradation. *C. trachomatis* has evolved strategies to counter such defense mechanisms by
translocating the effector Cdu1. Cdu1 utilizes its acetylase activity rather than its deubiquitinase
activity to protect itself and all three inclusion membrane proteins from being targeted for
ubiquitination and degradation. Cdu1 protects InaC in order to promote the recruitment of F-
actin scaffolds and Golgi ministacks to the inclusion perimeter and CTL0480 to recruit the
Myosin II regulator MYPT1. All three inclusion proteins and Cdu1 also participate in promoting
extrusion of *C. trachomatis* inclusions late in infection. Image created with BioRender.com.
1437

A

WT L2

cdu1 (1.3 kb)

cdu1 ORF

1   750 b.p

cdu1*::GII *aadA

cdu1 (3 kb)

5' junction (1 k.b)   aadA (0.4 k.b)   3' junction (0.3 k.b)

cdu1 ORF   aadA

1   635   2,831
                                                                                b.p

GII (aadA) intron

B

5' junction	3' junction	cdu1	aadA	+ control

WT / cdu1::GII aadA (repeated across lanes)

Base pairs

3,000
2,500
2,000
1,500
1,000
750
500
250

*

1438

Supplementary Figure 1
TargeTron mediated disruption of the L2 *cdu1* ORF. (A) Depiction of the *cdu1* ORFs in WT
L2 and *cdu1*::GII strains with corresponding diagnostic PCR amplicons. (**B**) PCR analysis of the
disrupted *cdu1* ORF in the *cdu1*::GII strain. *Amplification of *cdu1* ORF in *cdu1*::GII strain was
unsuccessful due to size of amplicon.

1445

1446
Supplementary Figure 2
Generation of a *cdu1* null strain in *C. trachomatis* (L2). (**A**) Western blot analysis of protein
lysates derived from HeLa cells infected with a Ct L2 434 Bu parental strain (L2) and an L2
cdu1::GII *aadA* (*cdu1*::GII) strain for 24 hours. Blots were probed with antibodies raised against
recombinant Cdu1 (amino acids 71-401), Ct RpoB (bacterial RNA polymerase), and human
α tubulin. (**B**) HeLa cells infected with L2 transformed with pBOMB4-MCI empty vector (L2
pBOMB) and *cdu1*::GII transformed with empty pBOMB4-MCI (*cdu1*::GII pBOMB) were fixed at
24 hpi and stained with Cdu1 antisera. Cdu1 signal is depicted in green, Ct cells expressing
mCherry (encoded in pBOMB4-MCI vector) are shown in red, and DNA stained with Hoechst in
blue. Scale bar: 10μm. (**C**) Total RNA isolated from HeLa cells infected with L2 and *cdu1*::GII
strains for 24 hours was used to synthesize cDNAs which served as templates for PCR analysis
of *cdu1*, *cdu2*, and *cdu1-cdu2* bicistronic expression. RT+: Total RNA + reverse transcriptase.
RT-: Total RNA without reverse transcriptase. gDNA: Genomic DNA.
1460

Supplementary Figure 3
Proteins co-precipitating (non ubiquitinated) with human and Ct proteins enriched by
TUBE 1 pulldowns. (A) Volcano plots (pairwise comparisons) of human proteins (non-
ubiquitinated) that co-precipitate with TUBE1 bound proteins in mock infected HeLa cells and
HeLa cells infected with L2 or *cdu1* null strains (24 hpi). **(B)** Volcano plots (pairwise

1467   comparisons) of Ct TUBE1 co-precipitating proteins (non-ubiquitinated) identified in mock
1468   infected HeLa cells and HeLa cells infected with L2 or *cdu1* null strains (24 hpi).

A

Metascape pathway enrichment analysis: Human proteins (non-ubiquitinated) that co-precipitate with TUBE1 bound proteins in mock infected Hela cells

B

Metascape pathway enrichment analysis: Human proteins (non-ubiquitinated) that co-precipitate with TUBE1 bound proteins in *cdu1*::GII infected Hela cells

GO:0008610:lipid biosynthetic process

C

DAVID pathway enrichment analysis: *Chlamydia trachomatis* (L2) proteins (non-ubiquitinated) that co-precipitate with TUBE1 bound proteins in infected Hela cells

UP_KEYWORDs: Ribonucleoprotein
UP_KEYWORDS: Ribosomal protein
GOTERM_MF_DIRECT: structurual component of ribosomes
GOTERM_BP_DIRECT: translation
KEGG_PATHWAY: Ribosome

1482
Supplementary Figure 4
Pathway enrichment analysis of human and Ct proteins that co-precipitate (non
ubiquitinated) with TUBE1 bound proteins. (A) Metascape functional enrichment analysis of
human co-precipitating proteins in mock infected HeLa cells. **(B)** Metascape functional
enrichment analysis of human co-precipitating proteins in *cdu1* null infected HeLa cells. **(C)**
DAVID pathway enrichment analysis of Ct co-precipitating proteins enriched in infected (L2 and
cdu1 null) HeLa cells.
1490

1491
1492 **Supplementary Figure 5**
1493 **The number of inclusions and the size of inclusions in *cdu1* null, *inaC* null, *ipaM* null, and**
1494 ***ctl0480* null strains are similar across each strain (A)** Representative images of inclusions in
1495 infected HeLa cell monolayers at 48 hpi. Scale bar: 200 μm **(B)** Quantification of the number of
1496 inclusions in infected HeLa cell monolayers. **(C)** Quantification of inclusion size.
1497
1498