1

The acetylase activity of Cdu1 regulates bacterial exit from infected cells by protecting

***Chlamydia* effectors from degradation**

Robert J. Bastidas[1]*, Mateusz Kędzior[1], Robert K. Davidson[2,#], Stephen C. Walsh[2,#], Lee

Dolat[1], Barbara S. Sixt[3,4,5], Jonathan N. Pruneda[6] , Jörn Coers[1,2], Raphael H. Valdivia[1,2]*

Author affiliations: [1]Department of Integrative Immunobiology, Duke University, Durham, N.C

27708, USA. [2]Department of Molecular Genetics and Microbiology, Duke University, Durham,

N.C 27708, USA. [3]Deparment of Molecular Biology, Umeå University, Umeå, Sweden. [4]The

Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå, Sweden.

[5]Umeå Centre for Microbial Research (UCMR), Umeå University, Umeå, Sweden. [6]Department

of Molecular Microbiology & Immunology, Oregon Health & Science University, Portland, OR

97239, USA. [#]Authors contributed equally.

* Correspondence: robert.bastidas@duke.edu; raphael.valdivia@duke.edu

Abstract

Many cellular processes are regulated by ubiquitin-mediated proteasomal degradation. Pathogens can regulate eukaryotic proteolysis through the delivery of proteins with de-ubiquitinating (DUB) activities. The obligate intracellular pathogen *Chlamydia trachomatis* secretes Cdu1 (ChlaDUB1), a dual deubiquitinase and Lys-acetyltransferase, that promotes Golgi remodeling and survival of infected host cells presumably by regulating the ubiquitination of host and bacterial proteins. Here we determined that Cdu1's acetylase but not its DUB activity is important to protect Cdu1 from ubiquitin-mediated degradation. We further identified three *C. trachomatis* proteins on the pathogen-containing vacuole (InaC, IpaM, and CTL0480) that required Cdu1's acetylase activity for protection from degradation and determined that Cdu1 and these Cdu1-protected proteins are required for optimal egress of *Chlamydia* from host cells. These findings highlight a non-canonical mechanism of pathogen-mediated protection of virulence factors from degradation after their delivery into host cells and the coordinated regulation of secreted effector proteins.

Introduction

Ubiquitination is a conserved and ubiquitous post-translational modification (PTM) of proteins involving the conjugation of the carboxy-terminal glycine residue of ubiquitin (Ub) to lysine residues of target proteins. Poly-ubiquitination of substrates involves further conjugation of a Ub internal lysine residue or amino-terminal methionine (M1) with a second Ub molecule. Seven internal lysines in Ub (K6, K11, K27, K29, K33, K48, K63) and M1 are utilized by Ub conjugating enzymes to form homogeneous, branched, or mixed poly-ubiquitin (polyUb) chains (Komander and Rape., 2012). PolyUb chains with different linkage types exhibit distinct structures and functions. For example, K48- and K11-linked polyUb chains exhibit a compact conformation and are substrates for 26S proteasome-mediated degradation (Varadan et al., 2004; Tenno et al., 2004; Eddins et al., 2007; Bremm et al., 2010; Saeki., 2017). In contrast, K63-linked polyUb

conjugates adopt more open conformations that enable the recruitment of multiprotein

complexes that regulate the function of the target protein by proteolytic independent events

(Komander et al., 2009b; Weeks et al., 2009; Datta et al., 2009; Komander and Rape., 2012).

Mixed and branched polyUb chains are also emerging as important regulators of physiological

functions (Swatek and Komander., 2016; Ohtake and Tsuchiya., 2017).

Protein ubiquitination regulates numerous eukaryotic cell processes including protein

degradation, signal transduction, cell cycle regulation, selective autophagy, the DNA damage

response, and programmed cell death. Ub also plays key roles in modulating host innate

immune responses to bacterial infection (Li et al., 2016), bacterial proteins, pathogen-containing

vacuoles, and bacteria themselves by targeting them for Ub-mediated degradation by

proteasomal or autophagic machineries (Li et al., 2016). Because the Ub system is critical for

pathogen containment, many pathogens have evolved mechanisms to counteract the impact of

this PTM (Vozandychova et al., 2021). For instance, bacterial deubiquitinases (DUBs) can

remove Ub from ubiquitinated substrates thereby dampening inflammatory and cell-autonomous

defense mechanisms (Kubori et al., 2019). Many DUBs are cysteine proteases with a catalytic

Cys, a nearby His and an Asn/Asp (Komander et al., 2009a). DUBs are typically dedicated to

the removal of Ub moieties and are unable to hydrolyze other Ub-like (Ubl) modifications such

as SUMO or NEDD8. However, the CE clan of Ubl proteases (ULPs) can catalyze the removal

of both SUMO and NEDD8 (Ronau et al., 2016). Bacterial pathogens also encode CE clan

enzymes that function as DUBs, ULPs or both. For instance, *Salmonella* Typhimurium SseL*,*

Escherichia coli ElaD, and *Shigella flexneri* ShiCE function as Ub specific proteases (Rytkönen

et al., 2007; Catic et al., 2007; Pruneda et al., 2016) while RickCE from *Rickettsia belli* functions

as a protease directed towards both Ub and NEDD8 as does SidE from *Legionella pneumophila*

which displays mixed activities towards Ub, NEDD8, and ISG15 (Sheedlo et al., 2015; Pruneda

et al., 2016). Similarly, XopD from *Xanthamonas campestris* and LotB from *L. pneumophila* are

isopeptidases exhibiting cross reactivity towards both Ub and SUMO (Pruneda et al., 2016;

Schubert et al., 2020). Some CE clan bacterial effectors display acetyltransferase activity. *L.*

pneumophila LegCE, *S.* Typhimurium AvrA, and YopJ from *Yersinia pestis* function exclusively

as acetyltransferases (Mittal et al., 2006; Mukherjee et al., 2006; Jones et al., 2008; Pruneda et

al., 2016). In contrast, the *Chlamydia trachomatis* (Ct) effector Cdu1/*Chla*DUB1 is a CE clan

protein that exhibits both acetyltransferase and deubiquitinating activities (Misaghi et al., 2006;

Pruneda et al., 2016; Fischer et al., 2017; Pruneda et al., 2018).

Ct is an obligate intracellular bacterial pathogen responsible for human diseases of significant

clinical and public health importance (Haggerty et al., 2010). Ct has a biphasic developmental

cycle in which the Ct infectious propagule or elementary body (EB) invades the target host cell.

Upon internalization the EB transitions to the reticulate body (RB). RBs replicate by binary

fission within a pathogenic vacuole ("inclusion") and asynchronously differentiate back to EBs.

In cell culture, starting at around 48 hours-post infection (hpi) Ct will egress after lysis of the

host cell or by a process termed extrusion, wherein the intact inclusion exits from the infected

cell (Moulder., 1991; Abdelrahman and Belland., 2005; Hybiske and Stephens., 2007; Lee et al.,

2018). The effector Cdu1 was originally identified as a deneddylating and deubiquitinating

enzyme and subsequently shown to exhibit *in-vitro* isopeptidase activity towards both Lys48 and

Lys63 linked di-Ub substrates (Misaghi et al., 2006; Claessen et al., 2013; Pruneda et al., 2016;

Fischer et al., 2017). Cdu1 is unique among CE clan enzymes in that it also functions as a bona

fide lysine acetylase with both is acetylase and DUB activities catalyzed by the same catalytic

active site (Pruneda et al., 2018). Intriguingly, Cdu1 autoacetylation is directed towards lysines

unlike other CE clan acetylases that predominantly target serine and threonine residues

(Pruneda et al., 2018). In transfected cells, Cdu1 protects the NFκB cytoplasmic retention factor

IκBα from ubiquitination and proteasomal degradation (Le Negrate et al., 2008). In infected

cells, Cdu1 localizes to the inclusion membrane where it functions to stabilize the anti-apoptotic

protein Mcl-1 and to promote the repositioning of Golgi ministacks around the Ct inclusion

(Fischer et al., 2017; Wang et al., 2018; Pruneda et al., 2018; Kunz et al., 2019; Auer et al.,

2020). However, the mechanism by which Cdu1 promotes redeployment of Golgi ministacks

and any additional roles that Cdu1 may play during Ct infection of epithelial cells remains

unknown.

In this study, we show that Cdu1 protects itself and three secreted Ct effectors, InaC, IpaM, and

CTL0480 from targeted ubiquitination and proteasomal degradation. InaC, IpaM, and CTL0480

are members of a larger family of bacterial proteins embedded within the inclusion membrane

(Inc proteins) (Bannantine et al., 2000; Rockey et al., 2002; Chen et al., 2006; Li et al., 2008;

Alzhanov et al., 2009; Dehoux et al., 2011; Lutter et al., 2012; Lutter et al., 2013; Kokes et al.,

2015; Weber et al., 2015). We show that Cdu1-mediated protection from degradation is

independent from its DUB activity but relies upon its Lys acetylase activity. We show that Cdu1

protects InaC to promote repositioning of Golgi ministacks and formation of actin scaffolds

around the Ct inclusion, and CTL0480 to promote recruitment of myosin phosphatase target

subunit 1 (MYPT1) to the inclusion. In addition, we determined that Cdu1 and Cdu1-protected

Incs are required for optimal extrusion of inclusions from host cells at the late stages of

infection.

Results

The *C. trachomatis* inclusion membrane proteins InaC, IpaM, and CTL0480 are

differentially ubiquitinated in the absence of Cdu1

Cdu1 is required for Golgi repositioning around the Ct inclusion (Pruneda et al., 2018; Auer et

al., 2020). To understand how Cdu1 promotes Golgi redistribution, we first generated a *cdu1*

null strain in a Ct L2 background by TargeTron mediated insertional mutagenesis (pDFTT3-

aadA) (Lowden et al., 2015) (S. Figure 1). Loss of Cdu1 expression in the resulting L2 *cdu1*::GII

aadA (*cdu1*::GII) strain was verified by western blot analysis and by indirect

immunofluorescence with antibodies raised against Cdu1 (S. Figures 2A and 2B). Because

cdu2 resides directly downstream of the *cdu1* locus and encodes a second Ct DUB

(Cdu2/*Chla*DUB2) (Misaghi et al., 2006), we first determined whether the disruption of *cdu1*

impacted the expression of *cdu2*. We detected *cdu1* and *cdu2* transcripts in HeLa cells infected

with Ct L2 but not for the juncture between *cdu1* and *cdu2* (S. Figure 2C). In cells infected with

Ct *cdu1*::GII we only detected *cdu2* transcripts (S. Figure 2C) confirming that *cdu1* and *cdu2* are

not co-expressed as part of an operon in accordance with previous observations (Albrecht et al.,

2010).

In transfected Hela cells, Cdu1's DUB activity has been linked to fragmentation of the Golgi

apparatus (Pruneda et al., 2018). We therefore hypothesized that Cdu1s' DUB activity in

infected cells promoted Golgi redistribution around inclusions and that we could identify

potential targets by comparing the protein ubiquitination profile of cells infected with WT or

cdu1::GII strains by quantitative mass spectrometry (MS). HeLa cell were mock infected or

infected with either WT L2 or *cdu1*::GII strains. At 24 hpi, poly-ubiquitinated proteins were

enriched from lysed cells using Tandem Ubiquitin Binding Entities (TUBEs) (LifeSensors).

TUBEs consist of concatenated Ub binding associated domains (UBAs) that bind to polyUb-

modified proteins with nanomolar affinities. Poly-ubiquitinated proteins of both human and Ct

origin were enriched and identified by quantitative LC-MS/MS analysis.

Over 2,000 non-ubiquitinated proteins co-precipitated with TUBE 1 bound proteins across all

three conditions (mock, L2, and *cdu1*::GII infected HeLa cells) and 3 biological replicates (S.

Table 1). Among these, 47 human proteins were significantly enriched in mock infected HeLa

cells and 50 human proteins were significantly enriched during Ct infection (L2 and *cdu1*::GII)

(S. Table 3, S. Figure 3) Pathway enrichment analysis revealed that proteins involved in RNA

metabolism were overrepresented among co-precipitating proteins from mock infected cells (S.

Table 4, S. Figure 4) while no biological pathways or processes were overrepresented in

proteins enriched from infected cells (S. Figure 4). We also identified 8 TUBE1 co-precipitating

Ct proteins in HeLa cells infected with L2 and *cdu1*::GII (S. Table 5, S. Figure 3).

TUBE 1 affinity capture lead to the identification of 43 ubiquitinated proteins (35 human proteins

and 8 Ct proteins across all 3 conditions and replicates) based on the presence of peptides

containing a di-glycine remnant motif (Peng et al., 2003) (S. Tables 6-8). The lack of widespread

poly-ubiquitination of either human or Ct proteins in response to Ct infection (Figure 1) was

surprising given that wholesale changes in protein ubiquitination has been reported during

infection of HeLa cells by intracellular pathogens like *S.* Typhimurium (Fiskin et al., 2016). It is

also possible that had we conducted our analysis at different time points post-infection (hpi), we

might have identified additional Cdu1 targets, such as Mcl1 and IκBα (Le Negrate et al., 2008;

Fischer et al., 2017) which were not identified in our analysis. However, given that we observed

Cdu1 at the inclusion membrane as early as 1 hpi (data not shown), we opted to focus on an

earlier stage of the infection cycle. Only two human ubiquitinated proteins (ZC3H7A and DDIT4)

were found to be significantly enriched in response to WT L2 infection at 24 hpi (Figures 1A and

1C, S. Table 8) while only one human protein (MGC3121) was preferentially ubiquitinated in

HeLa cells infected with the *cdu1*::GII mutant strain (Figures 1B and 1C, S. Table 8). In contrast

three Ct proteins, InaC (K104, K107, and K149), IpaM (K29), and CTL0480 (K115) were

ubiquitinated at Lys residues in the absence of Cdu1 (Figures 1B, 1C, and 1D, S. Table 8).

InaC, IpaM, and CTL0480 are Ct effector proteins that localize to the inclusion membrane (Chen

et al., 2006; Alzhanov et al., 2009; Lutter et al., 2013; Kokes et al., 2015). These Type 3

secretion substrates belong to a family of over 36 inclusion membrane proteins (Incs) that

contain a signature bi-lobal hydrophobic transmembrane domain (Bannantine et al., 2000;

Rockey et al., 2002; Li et al., 2008; Dehoux et al., 2011; Lutter et al., 2012). Incs provide many

functions important for Ct intracellular replication ranging from providing structural integrity to

the inclusion membrane, to regulating membrane trafficking, to mediating interactions with host

organelles and cytoskeletal structures (reviewed in Bugalhão et al., 2019). InaC facilitates the

activation of the small GTPase RhoA, a crucial step for the assembly of actin scaffolds around

the inclusion (Haines et al., 2021; Kumar and Valdivia., 2008). Additionally, InaC plays a pivotal

role in the activation of Arf GTPases. This activation subsequently induces post-translational

modifications of microtubules in close proximity to the inclusion membrane, which are essential

for Ct to initiate the repositioning of Golgi ministacks around the inclusion (Wesolowski et al.,

2017). IpaM exhibits localization to discrete patches in the inclusion termed microdomains

(Alzhanov et al., 2009; Dumoux et al., 2015). Upon ectopic expression, IpaM induces alterations

in microtubule organization (Dumoux et al., 2015). CTL0480, facilitates the recruitment of

MYPT1 (myosin phosphatase target subunit 1) to the inclusion membrane. Recrutiment of

MYPT1 is required for the efficient exit of Ct from host cells (Lutter et al., 2013; Shaw et al.,

2018). Because Cdu1 also localizes to the inclusion membrane (Fischer et al., 2017; Wang et

al., 2018; Pruneda et al., 2018; Kunz et al., 2019) we postulated that Cdu1 directly protects

InaC, IpaM, and CTL0480 from ubiquitination.

Cdu1 associates with InaC, IpaM, and CTL0480

We first determined if Cdu1 co-localized with InaC, IpaM, and CTL0480 at the inclusion

membrane. HeLa cells were infected for 24 hours with WT L2 or L2 expressing CTL0480-Flag

from its endogenous promoter and immunostained with antibodies against Cdu1, InaC, IpaM, or

the Flag epitope. Both InaC and CTL0480-Flag localized throughout the inclusion membrane

while IpaM was restricted to discrete microdomains as previously reported (Alzhanov et al.,

2009; Dumoux et al., 2015) (Figure 2A). Cdu1 co-localized with InaC and CTL0480-Flag and

with IpaM at microdomains (Figures 2A and 2B). All four antibodies specifically recognized their

corresponding antigens since immunostaining for Cdu1, InaC, IpaM or the Flag epitope was not

observed in Ct strains lacking Cdu1 (*cdu1*::GII), InaC (M407, Kokes et al., 2015) IpaM

(*ipaM*::GII, Meier et al., 2022), or a strain that does not express CTL0480-Flag (Figure 2A).

We next determined if Cdu1 can interact with InaC, IpaM, and CTL0480 by co-transfecting HEK

293 cells with vectors expressing either full length Cdu1-GFP or truncated versions of Cdu1-

GFP lacking transmembrane or catalytic domains (Figure 2C), and vectors expressing Flag-

InaC, V5-IpaM, and V5-CTL0480. Transfected cells were lysed and Cdu1-GFP was

immunoprecipitated with antibodies against GFP. Western blot analysis of the

immunoprecipitates showed that Flag-InaC, V5-IpaM, and V5-CTL0480 co-precipitated with

Cdu1-GFP (Figures 2D-F). Moreover, the transmembrane domain of Cdu1 was necessary for

Cdu1-GFP to interact with all three Incs (Figures 2D-F). The interaction between Cdu1-GFP and

the three tagged Incs was specific since we did not detect interactions between Cdu1-GFP and

a V5-tagged version of the inclusion membrane protein CpoS (Sixt et al., 2017) (Figure 2G). We

expected these interactions to be transient at the inclusion membrane as the engagement of

Cdu1 with its target(s) should mimic that of most enzymes with their substrates. Therefore, while

we could capture these complexes in co-immunoprecipitations in the context of overexpression,

this assay was not sensitive enough to reliably document the formation of complexes among

low abundance endogenous Ct proteins. Nevertheless, our findings from transfection

experiments lead us to conclude that Cdu1 can selectively interact with all three Incs even in the

absence of infection, and these interactions are facilitated by the transmembrane domain of

Cdu1.

Cdu1 protects InaC, IpaM, and CTL0480 proteins from degradation during infection

We next assessed if Cdu1 was required to stabilize endogenous InaC, IpaM, and CTL0480 in

infected cells. HeLa cells were infected with either WT L2 or *cdu1*::GII strains and at various

time points in the Ct infectious cycle crude cell lysates were analyzed by western blot to assess

the relative abundance of Inc proteins. At 36 and 48 hpi, the levels of InaC protein were found to

be undetectable in cells infected with *cdu1*::GII in comparison to cells infected with the WT L2

strain (Figures 3A and 3B). Conversely, IpaM protein levels exhibited a decrease at 36 and 48

hpi in cells infected with *cdu1*::GII (Figure 3C and 3D). We were not able to detect CTL0480

levels by western blot but were successful in following CTL0480 expression by indirect

immunofluorescence (Figure 3E). The relative abundance of CTL0480 at inclusion membranes

was not affected in *cdu1*::GII inclusions at 24 hpi. However, at 36 hpi, a subpopulation of cells

lost CTL0480 immunoreactivity and by 48 hpi, inclusion membranes of the *cdu1*::GII strain were

devoid of CTL0480 while CTL0480 was prominently detected at the inclusion membranes of WT

L2 (Figures 3E and Supplemental Figure 5). As controls for the specificity of antibodies used for

western blots and for indirect immunofluorescence, we included cells infected with Ct lacking

ipaM (ipaM::GII, Meier et al., 2022), with an *inaC* nonsense mutant (M407, Kokes et al.,

2015), and with Ct lacking *ctl0480* (*ctl0480*::GII, Shaw et al., 2018). Overall, our results

indicate that steady state protein levels of InaC and IpaM and CTL0480 localization at the

inclusion membrane, especially at late stages of infection, are dependent on Cdu1, and that

Cdu1 acts at different stages in the infection cycle.

The acetylase activity of Cdu1 is required for Cdu1 to protect itself, InaC, and IpaM from

polyubiquitination and proteasomal degradation

The crystal structures of Cdu1 bound to Ub or Coenzyme A indicated that the adenosine and

phosphate groups of Coenzyme A make contact with a helix in variable region 3 (VR-3) of Cdu1

while the Ile36-patch of Ub binds to the opposite face of the same helix (Pruneda et al., 2018).

Although Cdu1 catalyzes both of its DUB and acetylase (Act) activities with the same active site

(Pruneda et al., 2018) the two activities of Cdu1 can be uncoupled by the amino acid

substitution K268E in VR-3 which disrupts Coenzyme A binding required for Act activity and by

the amino acid substitution I225A in the Ub-binding region of VR-3 required for DUB activity

(Pruneda et al., 2018). These substitutions allowed us to test which of Cdu1's enzymatic

functions are required for the observed effects on protein stability. We generated Ct shuttle

plasmids, expressing WT Cdu1, a catalytically inactive variant of Cdu1 lacking both DUB and

Act activities (Cdu1^{C345A}) (Pruneda et al., 2018), a Cdu1 DUB-deficient variant (Cdu1^{I225A}), and

a Cdu1 Act-deficient variant (Cdu1^{K268E}). All Cdu1 constructs were expressed from the *cdu1*

endogenous promoter as 3X Flag epitope-tagged proteins.

Plasmids expressing each Cdu1 variant were transformed into the *cdu*1::GII mutant and the

resulting strains used to infect HeLa cells for 36 and 48 hours. The levels of endogenous InaC

and IpaM in cell extracts of infected cells were monitored by western blot analysis. At 36 hpi,

InaC protein levels drastically decreased in cells infected with *cdu1* null strains transformed with

empty vector or expressing the catalytic inactive variant of Cdu1 (Cdu1^{C345A}-Flag) (Figure 4A).

Likewise, IpaM protein levels diminished at 48 hpi during infection with the same strains (Figure

4B). Both InaC and IpaM protein levels were restored to wild type levels in *cdu1* null strains

complemented with wild type Cdu1-Flag (Figures 4A and 4B). Unexpectedly, cells infected with

a *cdu1* null strain ectopically expressing the DUB deficient Cdu1 variant (Cdu1^{I225A}-Flag)

displayed wild type levels of InaC and IpaM while the Act deficient variant (Cdu1^{K268E}-Flag) did

not (Figures 4A and 4B). These results suggest that the acetylase activity of Cdu1 rather than

its DUB activity is required for Cdu1's ability to stabilize InaC and IpaM proteins.

When we monitored the stability of each Flag-tagged Cdu1 variant, we observed that the

catalytically inactive variant of Cdu1 (C345A-Flag) was destabilized (Figures 4A and 4B- Flag

WB). We reasoned that Cdu1 also protects itself from being targeted for degradation in infected

cells. As with InaC and IpaM, the acetylase but not the DUB activity of Cdu1 was required for

Cdu1's stability (Figures 4A and 4B). Although the C345A (DUB-, Act-) and K268E (Act-) amino

acid substitutions in Cdu1 do not destabilize Cdu1 expressed in *E. coli* (Pruneda et al., 2018), it

was possible that these substitutions impacted the expression and/or folding of Cdu1 in

Chlamydia. To determine if these Cdu1 mutants were inherently unstable, we expressed

Cdu1^{C345A}-Flag and Cdu1^{K268E}-Flag in WT L2 in the presence of endogenous Cdu1. We found

that each Flag-tagged variant was stabilized (Figure 4C), indicating that endogenous Cdu1

protected the catalytically-deficient Cdu1 variants *in trans*. In addition, western blot analysis of

immunoprecipitated Cdu1-Flag and Cdu1^{C345A}-Flag expressed in a *cdu1*::GII mutant showed

that while WT Cdu1-Flag was not modified by Lys48-linked poly-ubiquitination, Cdu1^{C345A}-Flag

was robustly modified by Lys48-linked polyUb in the presence of the proteasome inhibitor

MG132 (Figure 4D). Moreover, endogenous Cdu1 protected Cdu1^{C345A}-Flag from Lys48-linked

polyUb when Cdu1^{C345A}-Flag was expressed in a wild type L2 background (Figure 4D). These

results indicate that the loss of Cdu1 activity likely leads to its Lys48-linked poly-ubiquitination

and subsequent proteasome-dependent degradation.

Because Cdu1 autoacetylates itself and its Act activity is directed towards lysines (Pruneda et

al., 2018) we postulated that Cdu1 may stabilize proteins from degradation by acetylating lysine

residues that are potential targets of ubiquitination. We tested this hypothesis by assessing

whether Cdu1, InaC, CTL0480, and IpaM are acetylated at lysines during infection. Fractions

enriched for inclusion membranes were isolated by sub-cellular fractionation from HeLa cells

infected with wild-type L2 (24 hpi), and proteins acetylated at lysines were immunoprecipitated.

Western blot analysis of acetyl-lysine immunoprecipitates indicated that InaC and IpaM, but not

the Inc protein IncA, were acetylated (Figure 4E). Western blot analysis of anti-acetyl-lysine

immunoprecipitates of inclusion membrane-enriched membrane fractions of HeLa cells infected

with L2 expressing Cdu1-Flag (24 hpi), CTL0480-Flag (40 hpi), and IpaM-Flag (40 hpi) also

showed that all three Flag tagged effectors were acetylated at lysines (Figure 4F). We also

determined that Flag-tagged InaC expressed in an *inaC* null (M407) background was acetylated

at lysines as determined by western blot analysis of anti-acetyl-lysine immunoprecipitates

(Figure 4G). In addition, we identified acetylated forms of Cdu1 from mass spectrometric

analysis of Flag immunoprecipitates derived from extracts of HeLa cells infected with L2

expressing Cdu1-Flag, (24 hpi) (Figure 4H).

Cdu1's acetylase activity shields inclusions from ubiquitination but is not sufficient to

protect against IFNγ mediated antimicrobial activity.

We reasoned that the lysine acetylase activity of Cdu1 is a prominent mechanism by which

Cdu1 protects client proteins (at 24 hpi), since loss of Cdu1 or expression of the acetylase

deficient variant of Cdu1 (Cdu1^{K268E}-Flag) leads to a marked increase in Ub immunostaining at

or near the periphery of *cdu1*::GII inclusions (> 80% of inclusions) compared to HeLa cells

infected with *cdu1*::GII strains complemented with wild type or DUB deficient (I225A-Flag) Cdu1

strains (Figures 5A, 5B, and Supplemental Figure 6A). Given that Cdu1 appears to localize

exclusively at inclusion membranes we predicted that its activity would be spatially restricted to

the inclusion periphery. We tested this premise by co-infecting HeLa cells with an *incA* null

strain (M923 (IncA^{R197*}), Kokes et al., 2015) and the *cdu1*::GII strain. IncA mediates

homotypic fusion of inclusion membranes and loss of IncA results in the accumulation of

multiple unfused inclusions in cells infected at high MOIs (Hackstadt et al., 1999; Suchland et

al., 2000; Pannekoek et al., 2005) (Figure 5C). As expected, *incA* mutants which retain Cdu1

activity did not accumulate Ub at or near the periphery of inclusion membranes. In HeLa cells

coinfected with both *cdu1*::GII and M923 (IncA^{R197*}), *cdu1::GII* Cdu1^{C345A} and M923, or

cdu1::GII Cdu1^{K268E} and M923, only the *incA* null inclusions were protected from ubiquitination

(Figures 5C, 5D, and Supplemental Figure 6B). Based on these observations we conclude that

the acetylase activity of Cdu1 protects proteins *in cis* and that this activity is constrained to the

membrane of the pathogenic vacuole consistent with previous reports (Auer et al., 2020).

Recently, the Ct inclusion membrane protein GarD was identified as a Ct effector that shields Ct

from γ-interferon mediated ubiquitination by the IFNγ-inducible human ubiquitin E3 ligase

RNF213 (Walsh et al., 2022). Because Cdu1 also protects the Ct inclusion from ubiquitination,

we tested if Cdu1 also plays a role in protecting Ct from IFNγ-induced cell immunity. A549 cells

were pretreated with IFNγ (100 U/mL) and infected with WT Ct, *cdu1*::GII, or *garD*::GII strains.

Infections with a *garD*::GII strain led to an approximate 90% decrease in the number of

inclusions formed relative to infections with its parental WT Ct strain (Figure 5E) while infections

with a *cdu1*::GII or its parental WT Ct strain showed a modest reduction in inclusion formation

(approximately 26% and 12% respectively) (Figure 5E). These results suggest that Cdu1 likely

does not play a role in protecting Ct from IFNγ-mediated cellular immunity.

We also tested whether RNF213 localizes to inclusions that lack Cdu1, as observed in IFNγ-

primed A549 cells infected with *garD::*GII strains (Walsh et al., 2022). RNF213 did not localize

to Ct inclusions when cells were infected with either WT Ct or *cdu1*::GII strains, regardless of

whether or not the A549 cells were treated with IFNγ (Figures 5F and 5G). In contrast, RNF213

localized to approximately 37% of inclusions in cells infected with *garD*::GII mutants of unprimed

A549 cells, and 81% in IFNγ treated cells (Figures 5F and 5G). Based on these results, we

conclude that Cdu1 does not play a role in protecting Ct from IFNγ-induced antimicrobial activity.

Cdu1 is required for F-actin assembly and Golgi ministack repositioning around the Ct

inclusion, and for MYPT1 recruitment to Ct inclusions

InaC is required for Ct to assemble F-actin scaffolds and to reposition Golgi mini stacks

around the periphery of the inclusion membrane (Kokes et al., 2015; Wesolowski et al., 2017;

Haines et al., 2021). Because Cdu1 regulates InaC levels, we predicted that *cdu1* mutants

would phenocopy *inaC* mutants. We quantified the number of inclusions surrounded by F-actin

cages at 40 hpi. In cells infected with WT L2 (parental strain of M407 (*inaC* null), Nguyen and

Valdivia., 2012; Kokes et al., 2015), approximately 25% of inclusions were surrounded by F-

actin, consistent with previous observations (Chin et al., 2012; Kokes et al., 2015) (Figures 6A

and 6B, and Supplemental Figure 7). The number of inclusions surrounded by F-actin

decreased to approximately 7% in cells infected with an *inaC* null strain (M407) and increased

to approximately 49% in HeLa cells infected with an *inaC* null strain (M407) complemented with

wild type InaC (Figures 6A and 6B, and Supplemental Figure 7). Cells infected with *cdu1*::GII

mutants transformed with an empty plasmid or expressing Cdu1^{C345A}-Flag (DUB- Act-) and

Cdu1^{K268E}-Flag (Act-) resulted in approximately 8%,13%, and 10% of of F-actin positive

inclusions respectively (Figures 6A and 6B, and Supplemental Figure 7). In contrast, cells

infected with *cdu1*::GII mutants expressing Cdu1-Flag and Cdu1^{I225A}-Flag (DUB-) led to a

marked increase in F-actin inclusions (approximately 52% and 46% respectively) (Figures 6A

and 6B, and Supplemental Figure 7). From these observations we conclude that the acetylase

activity of Cdu1 is required for Ct to promote assembly of F-actin around the Ct inclusion likely

through the stabilization of InaC.

We next quantified Golgi dispersal in infected HeLa cells at 24 hpi, a process that is also

dependent on InaC (Kokes et al., 2015; Wesolowski et al., 2017). In Hela cells infected with an

inaC null strain (M407) Golgi dispersal was limited to approximately 26% of the Ct inclusion

perimeter. In contrast, cells infected with either its parental WT L2 or with an *inaC* null strain

(M407) complemented with wild type InaC, the Golgi is dispersed around 45% of the inclusion

perimeter (Figures 6C and 6D, and Supplmental Figure 8). Similarly, Golgi dispersal around

inclusions during infection with WT L2 and in *cdu1*::GII mutants expressing wild type Cdu1-Flag

or Cdu1^{I225A}-Flag (DUB-) was approximately 43%, 41%, and 43% respectively (Figures 6C and

6D, and Supplmental Figure 8). In HeLa cells infected with *cdu1*::GII and *cdu1*::GII strains

expressing Cdu1^{C345A}-Flag (DUB- Act-), and Cdu1^{K268E}-Flag (Act-), Golgi repositioning was

restricted to approximately 24%, 23%, and 23% of inclusion perimeters respectively (Figures 6C

and 6D, and Supplemental Figure 8). These results confirm that both InaC and Cdu1 are

required for efficient repositioning of the Golgi around the Ct inclusion as previously reported

(Kokes et al., 2015; Wesolowski et al., 2017; Pruneda et al., 2018; Auer et al., 2020) and that

this process is independent of Cdu1's DUB activity but requires its acetylase activity. Moreover,

our results suggest that Cdu1 promotes Golgi repositioning by protecting InaC-mediated

redistribution of the Golgi around the Ct inclusion.

CTL0480 promotes recruitment of the myosin phosphatase subunit MYPT1 to the inclusion

membrane where it regulates the extrusion of intact inclusions from host cells (Lutter et al.,

2013; Shaw et al., 2018). Consistent with the gradual loss of CTL0480 from inclusions in cells

infected with the *cdu1*::GII strain starting at 36 hpi (Figures 3E and Supplemental Figure 5) we

also observed a complete loss of MYPT1 recruitment to inclusions by 48 hpi (Figures 6E and

6F).

Cdu1, InaC, IpaM, and CTL0480 are required for optimal extrusion of Ct from host cells

In the absence of Cdu1, the levels of InaC, CTL0480, and IpaM decreased late in infection (36

hpi and 48 hpi, Figures 3 and 4) suggesting that a prominent role of Cdu1 is to protect these

Incs from degradation late in infection. At the end of its developmental cycle, *Chlamydia* exits

host cells by promoting cellular lysis or by extrusion of intact inclusions (Hybiske and Stephens.,

2007). Ct host cell exit by extrusion is an active process requiring a remodeling of the actin

cytoskeleton and the function of Inc proteins (Hybiske and Stephens., 2007; Chin et al., 2012;

Lutter et al., 2013; Shaw et al., 2018; Nguyen et al., 2018). CTL0480 recruits MYPT1 (an

inhibitor of Myosin II motor complexes) to the inclusion membrane which prevents premature

extrusion of Ct inclusions and loss of CTL0480 leads to increased rates of extrusion by Ct from

infected HeLa cells (Lutter et al., 2013; Shaw et al., 2018) (Figure 7). Actin polymerization is

also required for Ct extrusion (Hybiske and Stephens., 2007; Chin et al., 2012) suggesting that

InaC dependent recruitment of F-actin to the inclusion may also contribute to optimal Ct

extrusion. IpaM localizes to microdomains in the inclusion membrane that are proposed to

function as foci for extrusion (Nguyen et al., 2018). Based on these observations, we postulated

that Cdu1-mediated protection of CTL0480, InaC, and IpaM regulates the extrusion of Ct

inclusions. We quantified the number of extrusions released from infected HeLa cells at 52 hpi

and observed a 60% reduction in the number of extrusions in HeLa cells infected with the

cdu1::GII strain relative to cells infected with WT L2 (Figures 7A and 7B). Complementation of

cdu1::GII with either wild type Cdu1-Flag or Cdu1^{I225A}-Flag (DUB-) restored extrusion

production to near wild type levels. In contrast, HeLa cells infected with *cdu1*::GII mutants

expressing Cdu1^{K268E}-Flag (Act-), or *inaC* (*inaC*::GII, Wesolowski et al., 2017) and *ipaM*

(*ipaM*::GII, Meier et al., 2020) null strains led to a 42%, 75%, and 58% reduction in extrusion

production respectively (Figures 7A and 7B). The decrease in the number of extruded inclusions

by these strains was not attributed to defects in inclusion biogenesis as they produced a

comparable number of inclusions at 48 hpi relative to cells infected with WT L2 (Supplemental

Figure 9). Consequently, we infer that InaC, IpaM, and Cdu1 collectively contribute to the

promotion of optimal extrusion of Ct inclusions from host cells, with Cdu1 playing a central

regulatory role by protecting these effectors from degradation.

In contrast, infection of HeLa cells with a *ctl0480::GII* mutant strain led to an increase in the

number of extruded inclusions as previously observed (Shaw et al., 2018) (Figures 7A and 7B).

Therefore, even though the Cdu1-mediated protection of InaC and IpaM is important for the

extrusion of inclusions and *cdu1* mutants phenocopy the loss of InaC and IpaM, the phenotypic

similarities do not extend to the increased number of extruded inclusions observed in cells

infected with the *ctl0480*::GII mutant strain (Figures 7A and 7B). We infer from these

observations that functions for both InaC and IpaM in the extrusion of inclusions are epistatic to

CTL0480. Extruded inclusions produced during infection of HeLa cells also varied in size with

an average diameter of 40 μm (Figures 7A and 7C). Interestingly, the loss of IpaM and over

expression of Cdu1-Flag and Cdu1^{I225A}-Flag (DUB-) shifted the size distribution of extrusions

toward larger extrusions (Figures 7A and 7C) suggesting that Ct regulates the size of extruded

inclusions through Cdu1.

Discussion

Several *Chlamydia* Inc proteins regulate interactions between the pathogenic vacuole and the

host cytoskeleton, organelles, and vesicular trafficking pathways. These Inc proteins also

modulate host cell death programs and promote *Chlamydia* exit from host cells (reviewed in

Bugalhão and Mota., 2019)**.** Given the central roles that Incs play in promoting *Chlamydia*

intracellular infection, it is not surprising that they are targeted for inactivation by host cellular

defenses. In response, *Chlamydia* has evolved mechanisms to protect Incs. In this study we

show that the acetylase activity of the effector Cdu1 protects itself and three Inc proteins; InaC,

IpaM, and CTL0480, from ubiquitination and degradation (Figure 8). Interestingly, all three Inc

proteins play prominent roles in regulating the extrusion of inclusions from host cells (Figure 7).

Observations that the encapsulation of *Chlamydia* within an extruded inclusion enhances

survival of *Chlamydia* within macrophages (Zuck et al., 2017) together with the broad

conservation of extrusion as an exit strategy among *Chlamydia* (Zuck et al., 2016) suggests that

this mechanism is important for *Chlamydia* pathogenesis. Notably, a *cdu1* mutant strain

(*cdu1*::Tn, Fischer et al., 2017) displays reduced bacterial loads in a murine model of upper

genital tract infections (Fischer et al., 2017). While neither this strain nor our *cdu1*::GII mutant

strain shows evident growth impairments during infection of Hela cells (data not shown), the

observed reduction in bacterial load in the absence of Cdu1 in animal models of infection could

potentially stem from defects in extrusion production or from perturbations in Cdu1-dependent

regulation of extrusion size. Thus, targeting Inc proteins that regulate extrusion for Ub-mediated

destruction may be advantageous for the host. For instance, targeting InaC for degradation

would limit F-actin dependent extrusions (Hybiske and Stephens., 2013; Chin et al., 2012) and

InaC-dependent microtubule scaffolds around the inclusion (Wesolowski et al., 2017; Haines et

al., 2021). CTL0480 functions as an inhibitor of extrusions through its role in modulating the

activity of myosin light chain 2 (MLC$_2$) (Lutter et al., 2013; Shaw et al., 2018). IpaM localizes to

specialized microdomains in the inclusion membrane which are also sites of enrichment for over

9 inclusion membrane proteins including Ctl0480 and MrcA, both of which are required for

Chlamydia extrusion (Mital et al., 2010; Lutter et al., 2013; Nguyen et al., 2018). We also find

that the loss of IpaM shifted the size distribution of extrusions towards larger inclusions (Figure

7). We speculate that heterogeneity in the size of extrusions might facilitate uptake of some

extrusions by innate immune cells at infected mucosal sites to promote *Chlamydia* LGV

dissemination to distal sites in the genital tract and avoid clearance of *Chlamydia* by other

immune cells (Zuck et al., 2017).

Effectors that modulate the activity of other translocated effectors are referred to as

"metaeffectors", a term coined by Kubori and colleagues after observing that the *L. pneumophila*

effector LubX which functions as an E3 ligase, ubiquitinates the translocated effector SidH

leading to its degradation (Kubori et al., 2010). Several other effector-metaeffector interactions

have been described in *L. pneumophila*, *Salmonella enterica*, and *Brucella abortus* which

regulate the activity of other effectors either directly or indirectly by modifying the same host

target or cellular process (Kubori et al., 2010; Neunuebel et al., 2011; Jeong et al., 2015;

Urbanus et al., 2016; Smith et al., 2020; Iyer and Das., 2021). In this context we propose that

Cdu1 functions as a metaeffector in Ct to protect multiple effectors. We also observed that Cdu1

interactions with InaC, IpaM, and CTL0480 likely occur independently from each other and that

the kinetics of degradation in the absence of Cdu1 varies for each Inc (Figure 3).

Our findings indicate that the DUB activity of Cdu1 was not required to protect InaC, IpaM, and

CTL0480 from ubiquitination. Instead we find that Cdu1's lysine acetylase activity is required to

protect these Inc proteins and Cdu1 itself from ubiquitination. Indeed, we found that all three

Incs and Cdu1 are acetylated at lysines in infected cells. However, we were unable to determine

if lysine acetylation in all four proteins was dependent on Cdu1s' Act activity or if these PTMs

are protective. Why the DUB activity of Cdu1 is unable to compensate for loss of its Act activity

remains unknown. It is possible that Cdu1, like other DUBs, is regulated by PTMs (Komander et

al., 2009a). For instance, phosphorylation of human CYLD inhibits its DUB activity towards

TRAF2 while phosphorylation of human USP8 inhibits its DUB activity toward EGFR (Reiley et

al., 2005; Mizuno et al., 2007). Mass spectrometry analysis of immunoprecipitated Flag tagged

Cdu1 expressed in Ct revealed that Cdu1 is phosphorylated at multiple serine and threonine

residues (Figure 4H) as previously suggested (Zadora et al., 2019). We identified three

PX(S/T)P MAPK phosphorylation consensus sequence motifs in the proline rich domain (PRD)

of Cdu1, suggesting that MAPKs may regulate the DUB activity of Cdu1.

Cdu1 homologs are found in multiple *Chlamydia* species including *C. trachomatis*, *C.*

muridarum, *C. suis*, *C. psitacci*, *C. abortus*, *C. caviae*, and *C. felis* but is notably absent in the

genomes of *C. pneumoniae* and *C. pecorum*. The acquisition of a second deubiquitinase

paralog (Cdu2) has also occurred in *C. trachomatis*, *C. muridarum*, and *C. suis*. In the genomes

of all three species, *cdu2* resides directly adjacent to *cdu1*; an arrangement that presumptively

arose from a gene duplication event. Cdu2 is a dedicated ULP with deubiquitinating and

deneddylating activities (Misaghi et al., 2006; Pruneda et al., 2016). New evidence suggest that

both paralogues might not be functionally redundant. The crystal structure of Cdu2 has revealed

differences in residues involved in substrate recognition between Cdu1 and Cdu2 and that each

paralog might recognize polyUb chains differently (Hausman et al., 2020). The processivity

rates for removal of terminal Ub from polyUb chains also differs between both isopeptidases

with Cdu2 exhibiting limited trimming of polyUb as compared to Cdu1 (Hausman et al., 2020).

Moreover, Cdu2 lacks the proline rich domain found in Cdu1 which might be important for

regulation of Cdu1 enzymatic activity. The presence of Cdu2 might also explain the low

incidence of human and Ct proteins that were differentially ubiquitinated in the absence of Cdu1

(Figure 1). Whereas several *Chlamydia* species have acquired either one or two deubiquitinase

paralogs, both *C. pneumoniae* and *C. pecorum* have not. Instead, both species have acquired

an unrelated deubiquitinase (*Chla*OTU) belonging to the OTU family of proteases (Makarova et

al., 2000; Furtado et al., 2013). Curiously, *Chla*OTU is also found in *C. psitacci*, *C. abortus*, *C.*

caviae, and *C. felis* all of which encode only Cdu1 and is absent in *C. trachomatis*, *C.*

muridarum, and *C. suis*, all of which encode Cdu1 and Cdu2. It is noteworthy that *Chlamydia*

species have independently acquired deubiquitinases multiple times (Cdu1, Cdu2, *Chla*OTU)

and that some of these deubiquitinases have evolved into moonlighting enzymes reflecting the

diverse strategies adopted by pathogenic *Chlamydia* as they adapt to their particular niche.

Acknowledgements

We thank LifeSensors and the Duke Proteomics and Metabolomics Shared Resource Center for

their proteomics services. We thank the Duke Light Microscopy Core Facility for microscopy

services. We also thank Marcela Kokes for generating the IncA-Flag constructs used in this

study. This work was supported by NIH grants GM142486 to J.N.P, AI103197 to J.C, AI140019

to R.J.B and AI134891 to R.H.V.

Author contributions

R.J.B and R.H.V designed the study. R.J.B wrote the manuscript with input from all listed

authors. L.D generated the *cdu1*::GII strain. M.K verified generation of the *cdu1*::GII mutant

strain and showed interaction of Cdu1-GFP variants with InaC, IpaM, and CTL0480 in

transfected HEK cells. R.K.D and S.C.W contributed equally to this work and quantified

inclusion production and RNF213 localization in unprimed and IFNγ-primed A549 cells. R.J.B

performed all other experiments. J.C, B.S.S, and J.N.P shared reagents. R.J.B and R.H.V

proofed the manuscript.

Declaration of interests

R.H.V is a founder of Bloom Sciences (San Diego, CA), which is a microbiome therapeutics

company. Findings reported in this study are unrelated to the work being performed with Bloom

Sciences.

Material and Methods

Key Resource Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
See S. Table 9		
Bacterial and virus strains		
Chlamydia trachomatis LGV biovar L2 434 Bu (L2)	Richard Stephens (UC Berkeley)	N/A
L2 pBOMB-MCI (Parent: LGV L2 434 Bu)	This paper	N/A
L2 pBOMB-MCI_CTL0480-3XFlag (Parent: LGV L2 434 Bu)	This paper	N/A
L2 pBOMB-MCI_IpaM-3XFlag (Parent: LGV L2 434 Bu)	This paper	N/A
L2 pBOMB-MCI_Cdu1-3XFlag (Parent: LGV L2 434 Bu)	This paper	N/A
L2 pBOMB-MCI_Cdu1 ^{C345A}-3XFlag (Parent: LGV L2 434 Bu)	This paper	N/A
L2 pBOMB-MCI_Cdu1 ^{K268E}-3XFlag (Parent: LGV L2 434 Bu)	This paper	N/A
L2 *cdu1*::GII *aadA* (Parent: LGV L2 434 Bu)	This paper	N/A
L2 *cdu1*::GII *aadA* pBOMB-MCI (Parent: L2 *cdu1*::GII *aadA*)	This paper	N/A
L2 *cdu1*::GII *aadA* pBOMB-MCI_Cdu1-3XFlag (Parent: L2 *cdu1*::GII *aadA*)	This paper	N/A
L2 *cdu1*::GII *aadA* pBOMB-MCI_Cdu1 ^{C345A}-3XFlag (Parent: L2 *cdu1*::GII *aadA*)	This paper	N/A
L2 *cdu1*::GII *aadA* pBOMB-MCI_Cdu1 ^{I225A}-3XFlag (Parent: L2 *cdu1*::GII *aadA*)	This paper	N/A
L2 *cdu1*::GII *aadA* pBOMB-MCI_Cdu1 ^{K268E}-3XFlag (Parent: L2 *cdu1*::GII *aadA*)	This paper	N/A
L2 Rif-R (Parent: L2 434 Bu)	Nguyen et al., 2012	N/A
M407 (*inaC C307T*, InaC Q103*) (Parent: L2 Rif-R)	Kokes et al., 2015	N/A
M407 p2TK2 (Parent: M407)	Kokes et al., 2015	N/A
M407 p2TK2_InaC (Parent: M407)	Kokes et al., 2015	N/A
M407 p2TK2_InaC-3X Flag (Parent: M407)	This paper	N/A
inaC::GII *bla* (Parent: L2 434 Bu)	Wesolowski et al., 2017	N/A
ipaM::GII *cat* (Parent: L2 434 Bu)	Meier et al., 2022	N/A
ctl0480::GII *aadA* (Parent: L2 434 Bu)	Shaw et al., 2018	N/A
M923 (*incA C589T*, IncA R197*) (Parent: L2 Rif-R)	Kokes et al., 2015	N/A
M923 pBOMB-MCI (Parent: L2 M923)	Sixt et al., 2017	N/A
L2 434 Bu (Parent of L2 *garD*::GII)	Walsh et al., 2022	N/A

L2 *ctl0390*::GII *aadA* (*garD*::GII)	Walsh et al., 2022	N/A
Biological samples		
N/A		
Chemicals, peptides, and recombinant proteins		
PR-619 (DUB inhibitor)	Sigma-Aldrich	Cat# SML0430
MG132 (proteasome inhibitor)	Sigma-Aldrich	Cat# 474791
TUBE-1 (pUb affinity capture reagent)	LifeSensors	Cat# UM401M
Acti-stain[TM] 488 (Phalloidin 488)	Cytoskeleton Inc.	Cat# PHDG1
ChromoTek GFP-Trap® Agarose	Proteintech	Cat# gta
Protein A/G PLUS-Agarose	Santa Cruz Biotechnology	Cat# sc-2203
3xFLAG peptide	APExBIO	Cat# A6001
Recombinant human interferon gamma (IFNγ)	Millipore	Cat# IF005
LGV L2 434 Bu Cdu1 recombinant protein	Jonathan Pruneda (Oregon Health and Science University)	N/A
Critical commercial assays		
TargeTron[TM] gene knockout system	Sigma-Aldrich	Cat# TA0100
Deposited data		
C. trachomatis LGV L2 434 Bu proteome	NCBI	NCBI:txid47472
Homo sapiens (Human) proteome	UniProt	Proteome ID: UP000005640
TUBE-1 affinity capture proteomics data	LifeSensors	Supplemental Table 1
Experimental models: Cell lines		
Hela cells	ATCC	Cat# CCL-2 RRID:CVCL_0030
Vero cells	ATCC	Cat# CCL-81 RRID:CVCL_0059
HEK 293T cells	ATCC	Cat# CRL-3216 RRID:CVCL_0063
A549 cells	ATCC	Cat# CCL-185 RRID:CVCL_0023
Experimental models: Organisms/strains		
Chlamydia trachomatis LGV L2 434 Bu	Richard Stephens (UC Berkeley)	N/A
Oligonucleotides		
See S. Table 10		
Recombinant DNA		
Plasmid: pDFTT3-aadA_Cdu1 635/636	This paper	N/A
Plasmid: pOPIN-GFP_Cdu1 FL (aa 1-401)	Jonathan Pruneda (Oregon Health and Science University)	N/A
Plasmid: pOPIN-GFP_Cdu1 TMD- (aa 71-401)	Jonathan Pruneda (Oregon Health and Science University)	N/A
Plasmid: pOPIN-GFP_Cdu1 CD- (aa 1-130)	Jonathan Pruneda (Oregon Health and Science University)	N/A

Plasmid: pCDNA-DEST53 (w/o GFP)_InaC (CT813)-3XFLAG	This paper	N/A
Plasmid: pcDNA3.1/nV5-DEST_IpaM	This paper	N/A
Plasmid: pcDNA3.1/nV5-DEST_CTL0480 p	This paper	N/A
Plasmid: pcDNA3.1/nV5-DEST_CpoS	This paper	N/A
Plasmid: pBOMB4-MCI_CTL0480-3X Flag	This paper	N/A
Plasmid: pBOMB4-MCI_IpaM-3X Flag	This paper	N/A
Plasmid: pBOMB4-MCI_Cdu1-3XFlag	This paper	N/A
Plasmid: pBOMB4-MCI_Cdu1 C345A-3XFlag	This paper	N/A
Plasmid: pBOMB4-MCI_Cdu1 I225A-3XFlag	This paper	N/A
Plasmid: pBOMB4-MCI_Cdu1 K263E-3XFlag	This paper	N/A
Plasmid: p2TK2_SW2_InaC-3XFlag	This paper	N/A
Software and algorithms		
Targetronics	Targetronics, LLC	www.targetrons.com
Proteome Discoverer 2.3	Thermo Fisher Scientific	https://www.thermofisher.com/us/en/home/industrial/mass-spectrometry/liquid-chromatography-mass-spectrometry-lc-ms/lc-ms-software/multi-omics-data-analysis/proteome-discoverer-software.html
Mascot software	Matrix Science	https://www.matrixscience.com
MaxQuant 1.6.2.3		https://www.maxquant.org
Scaffold PTM	Proteome Software	https://www.proteomesoftware.com/products/scaffold-ptm
VolcaNoseR	Goedhart, and Luijsterburg., 2020	https://huygens.science.uva.nl/VolcaNoseR/
Metascape	Zhou et al., 2019	https://metascape.org/gp/index.html#/main/step1
DAVID Bioinformatic Resources	Huang et al., 2009a., Huang et al., 2009b	https://david.ncifcrf.gov/tools.jsp
Image J	Schneider et al., 2012	https://imagej.nih.gov/ij/
NEBaseChanger	New England Biolabs	https://nebasechanger.neb.com
Prism 9	GraphPad	https://www.graphpad.com/updates/prism-900-release-notes
HCS Studio Cell Analysis Software	Thermo Fischer Scientific	Cat# CX51110
Other		
N/A		

Resource Availability

Materials availability

- All newly generated materials associated with this study will be freely available upon request.

Data and code availability

- Unprocessed (raw) proteomics data received from LifeSensors can be found in Supplemental Table 1.

- Original data used for microscopy and western blots in this study can be found at Mendeley data repository (data.mendeley.com):

Bastidas, Robert; Kędzior, Mateusz; Davidson, Robert; Walsh, Stephen; Dolat, Lee; Sixt, Barbara; Pruneda, Jonathan; Coers, Jörn; Valdivia, Raphael (2023), "The acetylase activity of Cdu1 regulates bacterial exit from infected cells by protecting *Chlamydia* effectors from degradation", Mendeley Data, V1, doi: 10.17632/xt3nmkm375.1

- Any additional information required to reanalyze the data reported in this paper is available upon request.

- This paper does not report original code.

Experimental Model And Subject Details

Cell lines

Vero (CCL-81; RRID:CVCL_0030), HeLa (CCL-2; RRID:CVCL_0059), HEK293T (CRL-3216;

RRID:CVCL_0063), and A549 (CCL-185; RRID:CVCL_0023) cells were purchased from ATCC

and cultured in High Glucose Dulbecco's Modified Eagle's Medium supplemented with L-

glutamine, sodium pyruvate (DMEM; Gibco) and 10% fetal bovine serum (FBS; Sigma-Aldrich).

Cells were grown at 37°C in a 5% CO2 humidified incubator. Vero, Hela, and HEK293T cells

were derived from females while A549 cells were derived from a male. All four cell lines have

been authenticated by the Duke Cell Culture and DNA analysis facility.

Chlamydia strains and propagation

Chlamydia strains used in this study are listed in the Key Resources Table. Ct strains were

propagated in Vero cells and harvested by osmotic lysis at 48 hours post infection. Following

lysis extracts were sonicated and bacteria pelleted by centrifugation at 21,000 x g. Bacteria

were resuspended in SPG storage buffer (75g/L sucrose, 0.5 g/L KH_4HPO_4, 1.2 g/L Na_2HPO_4,

0.72 g/L glutamic acid, pH 7.5) and stored as single use aliquots at -80°C.

Method Details

Chlamydia infections

Chlamydia infections were synchronized by centrifugation (2,500 x _g_ for 30 minutes at 10°C)

onto HeLa cell monolayers and incubated for the indicated times. Co-infections were performed

by infecting HeLa cell monolayers at a 1:1 ratio using MOIs of 2 for each co-infecting strain.

Insertional mutagenesis of CTL0247 (_cdu1_)

Primer sequences for TargeTron[TM] mediated mutagenesis of the LGV L2 434 Bu _cdu1_

(CTL0247) ORF were designed at the TARGETRONICS, LLC web portal

(www.targetrons.com). IBS1/2, EBS1/delta, and EBS2 primers (primer sequences are listed in

S. Table 10) were used in a PCR reaction to generate homing sequences for TargeTron^{TM}

integration between nucleotides 635 and 636 of the *cdu1* ORF using a TargeTron^{TM} gene

knockout system (Sigma-Aldrich; TA0100) according to the manufacturer instructions. Homing

sequences were gel purified, digested with HindIII and BsrGI, and ligated into HindIII and BsrGI

digested pDFTT3-*aadA* (Lowden et al., 2015). Ligations were transformed into *E. coli* DH5α,

clones isolated, and *cdu1* redirected pDFTT3-aadA plasmids identified by restriction digest and

verified by Sanger sequencing (Eton Bioscience) using a T7-promoter specific primer. The

resulting plasmid was transformed into a *C. trachomatis* LGV L2 434 Bu strain and

transformants selected with 150 µg/mL spectinomycin and plaque purified as previously

described (Kędzior and Bastidas., 2019). Insertion of the GII *aadA* intron at the *cdu1* locus was

verified by PCR analysis (S. Figure1) using primers that amplify amplicons spanning the

cdu1::GII 5' (RBP409 and RBP436) and 3' (RBP468 and RBP118) junctions, the *cdu1* CDS

(RBP409 and RBP118), and the *aadA* CDS (RBP512 and RBP513). Primer sequences are

listed in S. Table 10. Loss of Cdu1 protein was verified by western blot and indirect

immunofluorescence analysis (S. Figures 2A and 2B).

Analysis of *cdu1* and *cdu2* transcription by RT-PCR

Confluent HeLa cell monolayers (2.9 x 10^6 cells/infection) were infected with wild type L2 434

Bu or L2 *cdu1*::GII *aadA* strains. At 24 hpi, total RNA was isolated with a Qiagen RNeasy kit

(Qiagen; 74004) according to the manufacturer instructions. Total RNA was treated twice with

DNAse I (NEB; M0303S) and used for cDNA synthesis using a SuperScript IV Reverse

Transcriptase kit (Thermo Fisher Scientific; 18090010). cDNAs synthesized with and without

reverse transcriptase were used as templates for PCR analysis (S. Figure 2C) using primers

that amplify amplicons spanning the *cdu*1 (CTL0247_F and CTL0247_R) and *cdu2* ORFs

(CTL0246_F and CTL0247_R), and the intergenic junction between the *cdu1* and *cdu2* ORFs

(CTL0246-0247_F and CTL0246-0247_R). Primer sequences can be found in the S. Table 10.

TUBE1 based global ubiquitin profiling

Confluent HeLa cell monolayers (5.04×10^6 cells/infection) were mock infected or separately

infected with WT LGV L2 434 Bu or a L2 *cdu1*::GII *aadA* strain at MOIs of 3. At 24 hpi cells were

collected and spun down (700 x *g* for 10 minutes), frozen at -80°C and shipped on dry ice to

LifeSensors (Malvern, PA) for quantitative TUBE1-based Mass Spectrometry Analysis. Cell

pellets from three independent biological replicates were sent to LifeSensors for analysis. Cell

were subsequently lysed in lysis buffer (50 mM Tris-HCL, pH 7.5, 150 mM NaCl, 2 mM EDTA,

1% NP-40, 10% glycerol, 1% Sodium Deoxycholate) supplemented with a protease inhibitor

cocktail, the DUB inhibitor PR-619 (Sigma-Aldrich; SML0430), and the proteasomal inhibitor

MG-132 (Sigma-Aldrich; 474791). Lysates were clarified by high-speed centrifugation (14,000 x

g, 10 minutes, 4°C) and supernatants containing 5 mg of protein were equilibrated with

magnetic TUBE-1 (LifeSensors; UM401M) and incubated overnight at 4°C under rotation.

TUBEs were isolated with a magnetic stand and washed sequentially with PBST and TUBE

wash buffer. Poly-ubiquitinated and associated proteins were eluted with TUBE elution buffer.

Eluted supernatants were neutralized with neutralization buffer and loaded onto SDS-gels and

run until SDS buffer reached 0.5 cm into the gel. Gels were stained with Coomassie Blue and

lanes excised, reduced with TCEP, alkylated with iodoacetamide, and digested with Trypsin

(Trypsin Gold, Mass Spectrometry Grade) (Promega; V5280). Tryptic digests were analyzed

using a 150 min LC run on a Thermo Scientific™ Q Exactive HF Orbitrap™ LC-MS/MS system.

MS data was searched against the UniProt human database (UniProt; Proteome ID:

UP000005640) and the *Chlamydia trachomatis* L2 434 Bu reference database (NCBI:txid47472)

using MaxQuant 1.6.2.3 (https://www.maxquant.org). Proteins, peptides, and site identification

was set to a false discovery rate of 1%. N-terminal acetylation, Met oxidation, and diGly remnant

on lysine residues was also identified. All peptides and proteins identified can be found in S.

Table 1. The intensities (sum of all peptide MS peak areas for a protein or ubiquitinated peptide)

for each protein and ubiquitinated peptide across all three biological replicates were used to

determine mean intensities and to calculate *p*-values based on one-way student t-tests. Volcano

plots of mean intensities vs. *p*-values were generated with VolcaNoseR (Goedhart and

Lujsterburg., 2020) (https://huygens.science.uva.nl/VolcaNoseR/) and used to identify

significantly enriched proteins and ubiquitinated peptides. Data used to generate each Volcano

plot can be found in S. Table 2. Pathway enrichment analysis was performed with Metascape

(Zhou et al., 2019) (https://metascape.org/gp/index.html#/main/step1) and DAVID bioinformatic

resources (Huang et al., 2009a; Huang et al., 2009b) (https://david.ncifcrf.gov/tools.jsp).

Inhibitors, antibodies, western blots, and densitometry analysis

MG-132 (25 μM) (Sigma-Aldrich; 474791) was added to infected cell monolayers 5 hours prior

to extract preparations. Recombinant Cdu1 protein (LGV L2 434 Bu, amino acids 71-401) was

generated as previously described (Pruneda et al., 2018) and kindly provided by Jonathan

Pruneda (Oregon Health and Science University, OR). Recombinant Cdu1 protein was used to

generate antibodies in immunized New Zealand White rabbits. Cdu1 antisera was pre-adsorbed

with crude cell extracts from HeLa cells infected with a *cdu1*::GII *aadA* strain. Pre-adsorbed

antisera was used for western blot analysis at a 1:500 dilution in a solution containing 5% BSA

supplemented with crude extracts from HeLa cells infected with a *cdu1*::GII *aadA* strain

(0.1mg/mL total protein). Antibodies, antibody dilutions, and antibody diluents used in this study

are listed in S. Table 9. For western blot analysis, lysates from infected HeLa cell monolayers

(2.4 x 10^6 cells) were prepared by incubating cell monolayers with boiling hot 1% SDS lysis

buffer (1% SDS, 100 mM NaCl, 50 mM Tris, pH 7.5). Lysates were collected, briefly sonicated,

and total protein concentration measured with a DC^{TM} Protein Assay Kit (BIO-RAD; 5000111). 8

μg (Slc1 and alpha Tubulin blots) and 25 μg of total protein lysates (all other blots) were loaded

onto 4-15% Mini-PROTEAN and TGX Stain Free™ Protein Gels (BIO-RAD; 4568084),

transferred to PVDF membranes (BIO-RAD; 1620177), blocked with 5% Milk/TBSt, and

incubated with primary antibodies overnight at 4°C. Protein signals were detected with Goat anti

mouse (H+L) IgG (ThermoFisher scientific; 31430) or Goat anti rabbit (H+L) IgG HRP

(ThermoFisher scientific; 31460) conjugated secondary antibodies (1:1000 in 5% Milk/TBSt) and

SuperSignal West Femto HRP substrate (ThermoFisher scientific; 34096). Antibody bound

membranes were imaged with a LI-COR Odyssey Fc Imager (LI-COR, Inc.). Varying amounts of

protein extracts were used to determine the linear range of detection for InaC, IpaM, and Slc1

antibodies prior to quantification of western blot images (data not shown). Protein bands were

quantified using western blot densitometry analysis with LI-COR Image Studio Software (LI-

COR, Inc.). InaC and IpaM densitometry measurments were normalized to corresponding Slc1

densitometry measurments.

Immunofluorescence microscopy

HeLa cells were grown on coverslips to 50% confluency (0.1 x 10^5 cells) and infected at MOIs of

0.6. At indicated times, infected cells were separately fixed with ice cold Methanol or with warm

PBS containing 4% formaldehyde for 20 minutes. After fixative removal, cells were washed with

PBS and formaldehyde fixed cells were incubated either in 5% BSA/PBS supplemented with

0.1% Triton X-100 or in 5% BSA/PBS supplemented with 0.05% Saponin for 30 minutes with

gentle rocking. Following washing with PBS, Methanol fixed cells were incubated with primary

antibodies diluted in 5% BSA/PBS and formaldehyde fixed cells were incubated with primary

antibodies diluted in 5% BSA/PBS supplemented with 0.1% Triton X-100 or 0.05% Saponin for

1 hour with gentle rocking. Dilutions for each antibody used can be found in S. Table 9.

Methanol fixed cells were washed with PBS and incubated with secondary antibodies diluted in

5% BSA/PBS and supplemented with Hoechst 33342 (2 µg/mL) (ThermoFisher Scientific;

H3570). Formaldehyde fixed cells were washed and incubated with 5% BSA/PBS supplemented

with 0.1% Triton X-100 and Hoechst or 0.05% Saponin and Hoechst for 1 hour protected from

light and with gentle rocking. For detection of F-actin, Phalloidin conjugated to Alexa Fluor^{TM}

488 (1:5000) (Act-Stain 488 Phalloidin; Cytoskeleton Inc.; PHDG1) was added for the last 20

minutes of incubation with the secondary antibodies. Coverslips were transferred to glass slides,

mounted with 10 µL of Vectashield (Vector Labs; H-1000) and incubated over night at room

temperature prior to imaging. Secondary antibodies used were goat anti-mouse (H+L) IgG

(ThermoFisher scientific; A-11001 and A-21235) and goat anti-rabbit (H+L) IgG (ThermoFisher

scientific; A-11008 and A-21244) conjugated to Alexa Fluor^{TM} 488 and Alexa Fluor^{TM} 647. All of

the antibodies used for indirect immunofluorescence analysis were analyzed under all three

staining conditions (Methanol, Formaldehyde/Triton X-100, and Formaldehyde/Saponin).

Quantitative immunofluorescent microscopy for RNF213 was performed as previously described

(Walsh et. al., 2022). Briefly, A549 cells were grown on coverslips in 24-well plates to full

confluency (~2 x 10^5 cells). Cells were infected with indicated *C. trachomatis* strains at an MOI

of 2. At 3 hours post-infection, all cells were given fresh DMEM supplemented with L-tryptophan

(100 µg/mL) with half of the wells given interferon-gamma (100U/mL; Millipore, IF005). At 24

hours post-infection, cells were fixed with cold, 4% PFA in PBS for 20 minutes. Cells were

permeabilized with ice-cold methanol for 1 minute and blocked in PBS containing 5% BSA and

2.2% glycine for 30 minutes. Antibody incubations and microscope slide mounting was

performed as described. Samples were blinded using tape and imaged on a Zeiss Axio

Observer.Z1 epifluorescent microscope. For each sample, at least 6 separate fields of view and

100 *Chlamydia* inclusions were captured, saved and further blinded using the ImageJ Blind

Analysis Tool plugin (https://imagej.net/plugins/blind-analysis-tools). Quantification of the

number of inclusions with RNF213 targeted to the inclusion membrane was performed using

ImageJ (Schneider et al., 2012). Targeted inclusions were scored as having the indicated

protein signal colocalize with >50% of the inclusion membrane signal (incA positive antibody

staining).

Representative images were acquired with an inverted confocal laser scanning microscope

(Zeiss 880) equipped with an Airyscan detector (Hamamatsu) and with diode (405 nm), argon

ion (488 nm), double solid-state (561 nm) and helium-neon (633) lasers. Images were acquired

with a 63x C-Apochromatic NA 1.2 oil-objective (Zeiss). Images acquired in Airyscan mode were

deconvoluted using automatic Airyscan processing in Zen software (Zeiss). Image acquisition

was performed at the Light Microscopy Core Facility at Duke University. Images used for

quantification were captured in an inverted microscope (Ti2-Nikon instruments) equipped with

an ORCA Flash 4.0 V3 sCMOS camera (Hamamatsu) and a SOLA solid-state white light

illuminator (Lumencro). Images were acquired using a 60x Plan Apochromatic NA 1.40 oil

objective. All images were opened with ImageJ (Schneider et al., 2012) and only linear

adjustments were made to fluorescence intensity for the entire image. Images were exported as

TIFFs and compiled with Adobe suite software (Illustrator).

Vector construction and *C. trachomatis* transformation

Constructs used in co-transfection experiments: Mammalian vectors expressing Cdu1-GFP

constructs were kindly provided by Jonathan Pruneda (Oregon Health and Science University,

OR). Briefly, geneblocks encoding full length Cdu1 (LGV L2 434 Bu, CTL0247) (amino acids 1-

401), Cdu1 lacking its transmembrane domain (amino acids 71-401), and Cdu1 lacking its

catalytic domain (amino acids 1-130) were generated and inserted into the pOPIN-GFP vector

(Berrow et al., 2007) by In-Fusion^TM cloning (Takara Bio; 638947), resulting in Cdu1 constructs

with a C-terminal eGFP-His tag preceded by a 3C protease cleavage site. The Flag-InaC

mammalian expression vector was derived from a Gateway^{TM} entry clone containing the *C.*

trachomatis Serovar D/UW-3/CX CT813 (*inaC*) ORF (amino acids 41-264) obtained from a *C.*

trachomatis ORFeome library (Roan et al., 2006). The entry vector was used as a donor

plasmid for Gateway^{TM} based transfer into a modified pcDNA^{TM} DEST53 (ThermoFisher

Scientific; 12288015) vector in which the cycle 3 *GFP* ORF was removed. A NEB Q5®-Site

Directed Mutagenesis Kit (New England Biolabs; E0554S) was used to introduce a 3X Flag

epitope tag at the N-terminus of the CT813 ORF and a stop codon at the end of the CT813

ORF. L2 *ipaM* (CTL0476), L2 CTL0480, and L2 *cpoS* (CTL0481) ORFs were PCR amplified

from cell lysates derived from Vero cells infected with wild type L2 LGV 434 Bu with primers

containing attB sequences (primers *ipaM* forward, *ipaM* reverse, CTL0480 forward, CTL0480

reverse, *cpoS* forward, and *cpoS* reverse). Primer sequences can be found in the S. Table 10.

PCR amplicons were used as donors for Gateway^{TM} BP Clonase^{TM} based transfers into the

donor vector pDONR^{TM}221 (ThermoFisher Scientific; 12536017) to generate entry plasmids.

Entry plasmids were used to transfer *ipaM*, CTL0480, and *cpoS* into the Gateway^{TM} destination

vector pcDNA^{TM}3.1/nV5-DEST (ThermoFisher Scientific; 12290010) by Gateway^{TM} LR

Clonase^{TM} based reactions. The resulting mammalian expression vectors express IpaM,

CTL0480, and CpoS with V5-epitopes fused to their N-terminus.

pBOMB4-MCI based plasmids. CTL0480 and *ipaM* ORFs were amplified by PCR from cell

extracts derived from Vero cells infected with wild type LGV L2 434 Bu. The CTL0480 ORF, 149

b.p of upstream sequence, and a 3X FLAG epitope was amplified by PCR using primers

RBP628 and RBP629. The CTL0476 (*ipaM*) ORF, 400 b.p of upstream sequence, and a 3X

FLAG epitope was amplified with primers RBP623 and RBP624. CTL0480 and *ipaM* amplicons

were digested with Not1 and Pst1 and cloned into Not1 and Pst1 digested pBOMB4-MCI

(Bauler and Hackstadt., 2014) to generate pBOMB4-MCI_CTL0480-3X Flag and pBOMB4-

MCI_IpaM-3X Flag plasmids respectively. pBOMB4-MCI_Cdu1-3XFlag plasmids were

generated by PCR amplification of 175 b.p of genomic sequence directly upstream of the L2

434 Bu CTL0247 (*cdu1*) ORF and the entire *cdu1* ORF tagged with a C-terminal 3X Flag

epitope tag (primers RBP460 and RBP461). PCR amplicons were generated from gradient

purified LGV L2 434 Bu EBs and cloned into a pCR™-Blunt II TOPO® vector using a Zero

Blunt™ TOPO™ PCR cloning Kit (ThermoFisher Scientific; K2800J10) according to the

manufacturer instructions. Cdu1 catalytic variants were generated with a NEB Q5®-Site Directed

Mutagenesis Kit (New England Biolabs; E0554S) using the *cdu1*p-*cdu1*-3X Flag construct

cloned into pCR™-Blunt II TOPO® as a template and following the manufacturer instructions.

Primers for introducing base pair changes were designed on the NEBaseChanger website

(https://nebasechanger.neb.com). The Cdu1^{C345A} variant was generated by changing the *TGC*

codon located at positions 1033-1035 in the *cdu1* ORF to *GCT* (primers RBP525 and RBP526).

The Cdu1^{I225A} variant was generated by substituting the *ATC* codon located at positions 673-

675 for *GCT* (primers RBP527and RBP528). The Cdu1^{K268E} variant was generated by

introducing an *A802G* base pair substitution (primers RBP529 and RBP530). Wild type *cdu1*p-

cdu1-3XFLAG and all three *cdu1* variants were digested with Not1 and Pst1 and ligated into

Not1 and Pst1 digested pBOMB4-MCI (Bauler and Hackstadt., 2014). Primer sequences can be

found in the S. Table 10.

p2TK2_SW2-inaC-3XFlag. The CTL0184 (*inaC*) ORF and 250 b.p of upstream sequence was

amplified by PCR from cell extracts derived from Vero cells infected with L2 434 Bu and cloned

into the p2TK2_SW2 vector (Agaisse and Derré., 2013). A NEB Q5®-Site Directed Mutagenesis

Kit (New England Biolabs; E0554S) was used to insert a 3X FLAG epitope sequence at the C-

terminus (stop codon removed) of the CTL0184 ORF to generate the p2TK2_SW2-InaC-3XFlag

plasmid.

pBOMB4-MCI based plasmids and p2TK2_SW2-InaC-3X Flag plasmids were transformed into

corresponding *Chlamydia* strains, and transformants were selected with 10 U/mL Penicillin G

and plaque purified as previously described (Kędzior and Bastidas., 2019). All primer

sequences and plasmids generated in this study are listed in S. Table 10 and Key Resources

Table.

Subcellular fractionation

HeLa cells (2.16×10^7 cells/strain) seeded in six well plates were mock infected or infected with

Chlamydia L2 434 Bu strains. At indicated time points cells were washed with ice-cold PBS,

collected in ice-cold PBS with a cell scraper, and transferred to 15 mL conical tubes. Cell

suspensions were centrifuged at 500 x g for 5 minutes at 4°C and cell pellets were resuspended

in 400 μL of ice-cold subcellular fractionation buffer (20 mM HEPES (pH 7.4), 10 mM KCl, 2mM

$MgCl_2$, 1mM EDTA, 1mM EGTA) supplemented with 1mM DTT and a 1x cOmplete Mini-EDTA

free protease inhibitor cocktail (Sigma-Aldrich; 11836170001). Cells were incubated on ice for

20 minutes and lysed with 30 strokes of a Dounce homogenizer. Cell lysates were sequentially

centrifuged twice at 720 x g for 5 minutes at 4°C to remove intact nuclei. Supernatants were

centrifuged at 10,000 x g for 5 minutes at 4°C and the heavy membrane (inclusion) fraction was

recovered and resuspended in IP lysis buffer (25 mM Tris-Hcl, 150 mM NaCl, 1% NP-40, 5%

Glycerol) supplemented with 1mM PMSF and a 1x cOmplete Mini-EDTA free protease inhibitor

cocktail (Sigma-Aldrich; 11836170001).

Immunoprecipitations (IPs)

Transfections and GFP-immunoprecipitations: HEK 293T cell monolayers (1.32×10^7

cells/transfection) seeded in 10 cm cell culture dishes pre coated with poly-L-Lysine (Sigma-

Aldrich; P4707) were grown to 50% confluency and transfected with 10 μg of each plasmid used

per co-transfection in a 1.5 to 1 ratio of jetOPTIMUS® (Polyplus; 101000051) transfection

reagent to total plasmid DNA according to the manufacturer instructions. At 24 hpi, transfected

cells were lysed in IP lysis buffer (described above) supplemented with 1mM PMSF and 1x

cOmplete Mini-EDTA free protease inhibitor cocktail (Sigma-Aldrich; 11836170001). Lysates

were transferred to Eppendorf tubes, sonicated, and cleared by centrifugation (21,000 x g, 15

minutes, 4°C). Supernatants containing 2 mg of total protein were incubated with magnetic

GFP-Trap® agarose (Proteintech; gta) for 1 hour at 4°C with rotation. Beads were washed

according to manufacturer instructions and immunoprecipitated proteins eluted with 2X Laemmli

sample buffer.

Flag immunoprecipitations: Mock and infected HeLa cell monolayers (1.44 x 10^7 cells) grown in

six well plates were lysed in IP lysis buffer (described above) and transferred to Eppendorf

tubes. Lysates were sonicated and cleared by centrifugation (21,000 x g, 15 minutes, 4°C).

Supernatants containing 2 mg of total protein were pre-cleared by incubating with Protein A/G

PLUS-Agarose (Santa Cruz Biotechnology; sc-2203) for 30 minutes at 4°C followed by

sedimentation of agarose resins by centrifugation. Supernatants were incubated with M2-anti

Flag mouse mAb (1:400) (Sigma-Aldrich; F1804) overnight at 4°C with rotation followed by

incubation with Protein A/G PLUS-Agarose for 3 hours at 4°C with rotation. Agarose resins were

sedimented and washed according to manufacturer instructions and immunoprecipitated

proteins eluted with 50 µL of 100 µg/mL 3xFLAG peptides (APExBIO; A6001).

Acetylated lysine immunoprecipitations. Heavy membrane (inclusion) subcellular fractions

isolated from infected HeLa cells and containing 1 mg of total protein were pre-cleared by

incubating with Protein A/G PLUS-Agarose (Santa Cruz Biotechnology; sc-2203) for 30 minutes

at 4°C followed by sedimentation of agarose resins by centrifugation. Supernatants were

incubated with an anti-acetylated lysine rabbit antibody (Cell signaling; #9441) (1:100) overnight

at 4°C with rotation followed by incubation with Protein A/G PLUS-Agarose for 3 hours at 4°C

with rotation. Agarose resins were sedimented and washed according to manufacturer

instructions and immunoprecipitated proteins eluted with 50 μL of 2X Laemmli sample buffer.

Input (8 μg of total protein for Slc1 and alpha Tubulin blots, and 25 μg for all other blots) and

immunoprecipitates (GFP, Flag, proteins acetylated at lysines) were loaded onto 4-15% Mini-

PROTEAN and TGX Stain Free™ Protein Gels (BIO-RAD; 4568084), transferred to

PVDF membranes (BIO-RAD; 1620177), blocked with 5% Milk/TBSt, and incubated with

primary antibodies overnight at 4°C. Protein signals were detected with Goat anti mouse (H+L)

IgG (ThermoFisher scientific; 31430) or Goat anti rabbit (H+L) IgG HRP (ThermoFisher

scientific; 31460) conjugated secondary antibodies (1:1000 in 5% Milk/TBSt) and SuperSignal

West Femto HRP substrate (ThermoFisher scientific; 34096). Antibody bound membranes were

imaged with a LI-COR imaging system (LI-COR, Inc.).

Identification Cdu1 lysine acetylation and phosphorylation sites by LC-MS/MS

HeLa cell monolayers (8.64 x10^7 cells/strain) grown in six well plates were infected with a wild

type L2 strain transformed with empty pBOMB4-MCI (Bauler and Hackstadt., 2014) plasmid or

with a wild type L2 strain transformed with a pBOMB4-MCI_*cdu1*-3X Flag plasmid. At 24 hpi

infected cells were lysed and Flag tagged Cdu1 was immunoprecipitated as described above.

Flag eluates from 3 independent biological replicates were sent to the Proteomics and

Metabolomics Shared Resource Facility at Duke University for quantitative LC-MS/MS analysis.

Samples were spiked with undigested casein, reduced with 10 mM dithiothreitol, and alkylated

with 20 mM iodoacetamide. Eluates were then supplemented with 1.2% phosphoric acid and S-

Trap (Protifi) binding buffer (90% Methanol, 100 mM TEAB). Proteins were trapped on the S-

Trap, digested with 20 ng/μL Trypsin (Trypsin Gold, Mass Spectrometry Grade) (Promega;

V5280), and eluted with 50 mM TEAB, 0.2% FA, and 50% ACN/0.2% FA. Samples were

lyophilized and resuspended in 1% TFA/2% acetonitrile containing 12.5 fmol/μL yeast alcohol

dehydrogenase. Quantitative LC/MS/MS was performed using a nanoAcquitiy UPLC system

(Waters Corp) coupled to Thermo Scientific™ Orbitrap™ Fusion Lumos high resolution accurate

mass tandem mass spectrometer via a nanoelectrospray ionization source. Data was analyzed

with Proteome Discoverer 2.3 (Thermo Fisher Scientific™.) and MS/MS data searched against

the *Chlamydia trachomatis* LGV L2 434 Bu reference database (NCBI:txid47472). Cdu1-Flag

MS/MS data was analyzed with Mascot software (Matrix Science) using Trypsin/P specificity for

N-terminal acetylation, lysine acetylation, lysine Ub, and S/T/Y phosphorylation identification.

Analysis identified multiple acetylated and phosphorylated Cdu1 peptides and no Cdu1

ubiquitinated peptides. Data was viewed in Scaffold with Scaffold PTM (Scaffold Software).

Interferon gamma sensitivity assays

Ct sensitivities to interferon-gamma was assayed as previously described (Walsh et. al., 2022).

Briefly, A549 cells were seeded in black 96-well clear-bottomed plates (Corning). The next day,

cells were stimulated with 0 U/mL or 100 U/mL interferon gamma (IFNγ; Millipore, IF005) in

DMEM supplemented with L-tryptophan (100 μg/mL). After 20 hours, cells were infected in

technical duplicate with indicated *Chlamydia* strains at an MOI of 2. At 24 hours post-infection,

plates were fixed with cold 4% PFA in PBS for 20 minutes. Samples were nuclear stained with

Hoechst in PBS for 10 minutes and sealed using an aluminum adhesive (Thermo). Inclusions

and host cell nuclei were imaged and quantified using the CellInsight CX5 High Content

Screening platform (Thermo; CX51110). Relative bacterial infectivities were calculated as the

number of inclusions divided by the total number of host nuclei for each sample. Interferon

sensitivity was calculated by normalizing the infectivities of each strain to it's "untreated" (-IFNγ)

control and expressed as a percentage.

Isolation and imaging of *Chlamydia* inclusion extrusions

HeLa cell monolayers (1.2 x 10^6 cells) were infected with Ct strains at MOIs of 0.8. At 48 hpi,

infected monolayers were imaged using an EVOS FL Cell Imaging System (ThermoFisher

Scientific) equipped with a 20x/0.4 NA objective and a CCD camera. Following imaging, growth

media was removed, cell monolayers washed with fresh growth media, and monolayers

incubated for an additional 4 hours at 37°C. At 52 hpi growth supernatants were collected and

transferred to Eppendorf tubes. Extrusions were enriched by centrifugation (1,500 rpm, 5 min)

and pellets (not always visible) containing extrusions were resuspended in 30 μL of 4%

Formaldehyde/PBS supplemented with Hoechst (2 μg/mL) and 0.2% Trypan Blue Solution

(Gibco, 0.4%). Extrusions were analyzed by plating 10 μL drops on a glass side (without

coverslips) and immediately imaged using an EVOS FL Cell Imaging System (ThermoFisher

Scientific) equipped with a 20x/0.4 NA objective and a CCD camera. Intact extrusions were

identified based on morphology, lacking nuclei, and being impermeable to trypan blue. Images

were opened in ImageJ (Schneider et al., 2012) and enumeration of inclusions and extrusions

was performed manually. The sizes of individual extrusions and inclusions were determined by

manually tracing a line around the perimeter of each extrusion and inclusion in ImageJ and

measuring perimeter length. All measurements were exported to Microsoft Excel. Data plots and

statistical analyses were done with Prism 9 (GraphPad) software. Datasets were analyzed for

significance using a paired student t-test.

Image analysis

Line scan profiles of Cdu1 co-localization with inclusion membrane proteins was performed with

ImageJ (Schneider et al., 2012) by tracing a line through regions of interest and plotting

fluorescent signal intensities with the Plot Profile function. Localization of CTL0480, recruitment

of MYPT1, and association of Ub with Ct inclusions was performed manually from maximum

projections in ImageJ. Assessment of F-actin recruitment to Ct inclusions was performed

manually in ImageJ by projecting four to five sections in order to capture the entire inclusion.

Redistribution of Golgi around the Ct inclusion was measured in ImageJ from maximum

projections. The perimeters of individual inclusions were manually traced, and lengths

measured. The length of dispersed Golgi was measured by tracing and measuring the length of

the GM130 signal directly adjacent to each inclusion. Dispersed Golgi length was divided by

inclusion perimeter length. All measurements were exported to Microsoft Excel for

quantification. Data plots and statistical analyses were done with Prism 9 (GraphPad).

Quantification And Statistical Analysis

Quantifications were generated from three independent experiments and measurements

derived from blinded images. Data plots and statistical analyses were done with Prism 9

(GraphPad) software. Datasets were analyzed for significance using a paired student t-test,

one-way ANOVAs with a Student-Newman-Keuls post hoc test, or two-way ANOVAS with a

Turkey post hoc test. Data graphs show means and error bars represent standard error. *p*-

values less than 0.05 are defined as statistically significant. The indicated statistical test for

each experiment can be found in the figure legends.

References

1. Abdelrahman YM, Belland RJ. 2005. The chlamydial developmental cycle. *FEMS Microbiol*

Rev **29**:949–959. doi:10.1016/j.femsre.2005.03.002

2. Agaisse H, Derré I. 2013. A *C. trachomatis* cloning vector and the generation of *C.*

trachomatis strains expressing fluorescent proteins under the control of a *C. trachomatis*

promoter. *PLoS ONE* **8**:e57090. doi:10.1371/journal.pone.0057090

3. Albrecht M, Sharma CM, Reinhardt R, Vogel J, Rudel T. 2010. Deep sequencing-based

discovery of the *Chlamydia trachomatis* transcriptome. *Nucleic Acids Res* **38**:868–877.

doi:10.1093/nar/gkp1032

4. Alzhanov DT, Weeks SK, Burnett JR, Rockey DD. 2009. Cytokinesis is blocked in

mammalian cells transfected with *Chlamydia trachomatis* gene CT223. *BMC Microbiol* **9**:2.

doi:10.1186/1471-2180-9-2

5. Auer D, Hügelschäffer SD, Fischer AB, Rudel T. 2020. The chlamydial deubiquitinase Cdu1

supports recruitment of Golgi vesicles to the inclusion. *Cell Microbiol* **22**:e13136.

doi:10.1111/cmi.13136

6. Bannantine JP, Griffiths RS, Viratyosin W, Brown WJ, Rockey DD. 2000. A secondary

structure motif predictive of protein localization to the chlamydial inclusion membrane. *Cell*

Microbiol **2**:35–47. doi:10.1046/j.1462-5822.2000.00029.x

7. Bauler LD, Hackstadt T. 2014. Expression and targeting of secreted proteins from

Chlamydia trachomatis. J Bacteriol **196**:1325–1334. doi:10.1128/JB.01290-13

8. Berrow NS, Alderton D, Sainsbury S, Nettleship J, Assenberg R, Rahman N, Stuart DI,

Owens RJ. 2007. A versatile ligation-independent cloning method suitable for high-

throughput expression screening applications. *Nucleic Acids Res* **35**:e45.

doi:10.1093/nar/gkm047

9. Bremm A, Freund SMV, Komander D. 2010. Lys11-linked ubiquitin chains adopt compact

conformations and are preferentially hydrolyzed by the deubiquitinase Cezanne. *Nat Struct*

Mol Biol **17**:939–947. doi:10.1038/nsmb.1873

10. Bugalhão JN, Mota LJ. 2019. The multiple functions of the numerous *Chlamydia trachomatis*

secreted proteins: the tip of the iceberg. *Microb Cell* **6**:414–449.

doi:10.15698/mic2019.09.691

11. Catic A, Misaghi S, Korbel GA, Ploegh HL. 2007. ElaD, a Deubiquitinating protease

expressed by *E. coli. PLoS ONE* **2**:e381. doi:10.1371/journal.pone.0000381

12. Chen C, Chen D, Sharma J, Cheng W, Zhong Y, Liu K, Jensen J, Shain R, Arulanandam B,

Zhong G. 2006. The hypothetical protein CT813 is localized in the *Chlamydia trachomatis*

inclusion membrane and is immunogenic in women urogenitally infected with *C.*

trachomatis. Infect Immun **74**:4826–4840. doi:10.1128/IAI.00081-06

1000

1001 13. Chen Y-S, Bastidas RJ, Saka HA, Carpenter VK, Richards KL, Plano GV, Valdivia RH.

1002 2014. The *Chlamydia trachomatis* type III secretion chaperone Slc1 engages multiple early

1003 effectors, including TepP, a tyrosine-phosphorylated protein required for the recruitment of

1004 CrkI-II to nascent inclusions and innate immune signaling. *PLoS Pathog* **10**:e1003954.

1005 doi:10.1371/journal.ppat.1003954

1006

1007   14. Chin E, Kirker K, Zuck M, James G, Hybiske K. 2012. Actin recruitment to the *Chlamydia*

1008   inclusion is spatiotemporally regulated by a mechanism that requires host and bacterial

1009   factors. *PLoS ONE* **7**:e46949. doi:10.1371/journal.pone.0046949

1010

1011   15. Claessen JHL, Witte MD, Yoder NC, Zhu AY, Spooner E, Ploegh HL. 2013. Catch-and-

1012   release probes applied to semi-intact cells reveal ubiquitin-specific protease expression in

1013   *Chlamydia trachomatis* infection. *Chembiochem* **14**:343–352. doi:10.1002/cbic.201200701

1014

1015   16. Datta AB, Hura GL, Wolberger C. 2009. The structure and conformation of Lys63-linked

1016   tetraubiquitin. *J Mol Biol* **392**:1117–1124. doi:10.1016/j.jmb.2009.07.090

1017

1018   17. Dehoux P, Flores R, Dauga C, Zhong G, Subtil A. 2011. Multi-genome identification and

1019   characterization of chlamydiae-specific type III secretion substrates: the Inc proteins. *BMC*

1020   *Genomics* **12**:109. doi:10.1186/1471-2164-12-109

1021

1022   18. Dumoux M, Menny A, Delacour D, Hayward RD. 2015. A *Chlamydia* effector recruits

1023   CEP170 to reprogram host microtubule organization. *J Cell Sci* **128**:3420–3434.

1024   doi:10.1242/jcs.169318

1025

1026   19. Eddins MJ, Varadan R, Fushman D, Pickart CM, Wolberger C. 2007. Crystal structure and

1027   solution NMR studies of Lys48-linked tetraubiquitin at neutral pH. *J Mol Biol* **367**:204–211.

1028   doi:10.1016/j.jmb.2006.12.065

1029

1030   20. Fischer A, Harrison KS, Ramirez Y, Auer D, Chowdhury SR, Prusty BK, Sauer F, Dimond Z,

1031   Kisker C, Hefty PS, Rudel T. 2017. *Chlamydia trachomatis*-containing vacuole serves as

1032   deubiquitination platform to stabilize Mcl-1 and to interfere with host defense. *eLife* **6**.

1033   doi:10.7554/eLife.21465

1034

1035   21. Fiskin E, Bionda T, Dikic I, Behrends C. 2016. Global analysis of host and bacterial

1036   ubiquitinome in response to *Salmonella* Typhimurium infection. *Mol Cell* **62**:967–981.

1037   doi:10.1016/j.molcel.2016.04.015

1038

1039   22. Furtado AR, Essid M, Perrinet S, Balañá ME, Yoder N, Dehoux P, Subtil A. 2013. The

1040   chlamydial OTU domain-containing protein *Chla*OTU is an early type III secretion effector

1041   targeting ubiquitin and NDP52. *Cell Microbiol* **15**:2064–2079. doi:10.1111/cmi.12171

1042

1043   23. Gehre L, Gorgette O, Perrinet S, Prevost M-C, Ducatez M, Giebel AM, Nelson DE, Ball SG,

1044   Subtil A. 2016. Sequestration of host metabolism by an intracellular

1045   pathogen. *eLife* **5**:e12552. doi:10.7554/eLife.12552

1046

1047   24. Goedhart J, Luijsterburg MS. 2020. VolcaNoseR is a web app for creating, exploring,

1048   labeling and sharing volcano plots. *Sci Rep* **10**:20560. doi:10.1038/s41598-020-76603-3

1049

1050   25. Hackstadt T, Scidmore-Carlson MA, Shaw EI, Fischer ER. 1999. The *Chlamydia*

1051   *trachomatis* IncA protein is required for homotypic vesicle fusion. *Cell Microbiol* **1**:119–130.

1052   doi:10.1046/j.1462-5822.1999.00012.x

1053

1054   26. Haggerty CL, Gottlieb SL, Taylor BD, Low N, Xu F, Ness RB. 2010. Risk of sequelae after

1055   *Chlamydia trachomatis* genital infection in women. *J Infect Dis* **201 Suppl 2**:S134-55.

1056   doi:10.1086/652395

1057

1058   27. Haines A, Wesolowski J, Ryan NM, Monteiro-Brás T, Paumet F. 2021. Cross Talk between

1059   ARF1 and RhoA Coordinates the Formation of Cytoskeletal Scaffolds during *Chlamydia*

1060   *Infection. MBio* **12**:e0239721. doi:10.1128/mBio.02397-21

1061

1062   28. Hausman JM, Kenny S, Iyer S, Babar A, Qiu J, Fu J, Luo Z-Q, Das C. 2020. The Two

1063   Deubiquitinating Enzymes from *Chlamydia trachomatis* Have Distinct Ubiquitin Recognition

1064   Properties. *Biochemistry* **59**:1604–1617. doi:10.1021/acs.biochem.9b01107

1065

1066   29. Huang DW, Sherman BT, Lempicki RA. 2009a. Bioinformatics enrichment tools: paths

1067   toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**:1–

1068   13. doi:10.1093/nar/gkn923

1069

1070   30. Huang DW, Sherman BT, Lempicki RA. 2009b. Systematic and integrative analysis of large

1071   gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**:44–57.

1072   doi:10.1038/nprot.2008.211

1073

1074   31. Hybiske K, Stephens RS. 2007. Mechanisms of host cell exit by the intracellular bacterium

1075   *Chlamydia. Proc Natl Acad Sci USA* **104**:11430–11435. doi:10.1073/pnas.0703218104

1076

1077   32. Iyer S, Das C. 2021. The unity of opposites: Strategic interplay between bacterial effectors

1078   to regulate cellular homeostasis. *J Biol Chem* **297**:101340. doi:10.1016/j.jbc.2021.101340

1079

1080   33. Jeong KC, Sexton JA, Vogel JP. 2015. Spatiotemporal regulation of a *Legionella*

1081   *pneumophila* T4SS substrate by the metaeffector SidJ. *PLoS Pathog* **11**:e1004695.

1082   doi:10.1371/journal.ppat.1004695

1083

1084   34. Jones RM, Wu H, Wentworth C, Luo L, Collier-Hyams L, Neish AS. 2008. *Salmonella avra*

1085   coordinates suppression of host immune and apoptotic defenses via JNK pathway

1086   blockade. *Cell Host Microbe* **3**:233–244. doi:10.1016/j.chom.2008.02.016

1087

1088   35. Kędzior M, Bastidas RJ. 2019. Forward and reverse genetic analysis of *Chlamydia. Methods*

1089   *Mol Biol* **2042**:185–204. doi:10.1007/978-1-4939-9694-0_13

1090

1091   36. Kokes M, Dunn JD, Granek JA, Nguyen BD, Barker JR, Valdivia RH, Bastidas RJ. 2015.

1092   Integrating chemical mutagenesis and whole-genome sequencing as a platform for forward

1093   and reverse genetic analysis of *Chlamydia. Cell Host Microbe* **17**:716–725.

1094   doi:10.1016/j.chom.2015.03.014

1095

1096   37. Komander D, Clague MJ, Urbé S. 2009a. Breaking the chains: structure and function of the

1097   deubiquitinases. *Nat Rev Mol Cell Biol* **10**:550–563. doi:10.1038/nrm2731

1098

1099   38. Komander D, Reyes-Turcu F, Licchesi JDF, Odenwaelder P, Wilkinson KD, Barford D.

1100   2009b. Molecular discrimination of structurally equivalent Lys 63-linked and linear

1101   polyubiquitin chains. *EMBO Rep* **10**:466–473. doi:10.1038/embor.2009.55

1102

1103   39. Komander D, Rape M. 2012. The ubiquitin code. *Annu Rev Biochem* **81**:203–229.

1104   doi:10.1146/annurev-biochem-060310-170328

1105

1106   40. Kubori T, Shinzawa N, Kanuka H, Nagai H. 2010. *Legionella* metaeffector exploits host

1107   proteasome to temporally regulate cognate effector. *PLoS Pathog* **6**:e1001216.

1108   doi:10.1371/journal.ppat.1001216

1109

1110   41. Kubori T, Kitao T, Nagai H. 2019. Emerging insights into bacterial deubiquitinases. *Curr*

1111   *Opin Microbiol* **47**:14–19. doi:10.1016/j.mib.2018.10.001

1112

1113   42. Kumar Y, Valdivia RH. 2008. Actin and intermediate filaments stabilize the *Chlamydia*

1114   *trachomatis* vacuole by forming dynamic structural scaffolds. *Cell Host Microbe* **4**:159–169.

1115   doi:10.1016/j.chom.2008.05.018

1116
1117   43. Kunz TC, Götz R, Sauer M, Rudel T. 2019. Detection of *Chlamydia* developmental forms

1118   and secreted effectors by expansion microscopy. *Front Cell Infect Microbiol* **9**:276.

1119   doi:10.3389/fcimb.2019.00276

1120

1121   44. Lee JK, Enciso GA, Boassa D, Chander CN, Lou TH, Pairawan SS, Guo MC, Wan FYM,

1122   Ellisman MH, Sütterlin C, Tan M. 2018. Replication-dependent size reduction precedes

1123   differentiation in *Chlamydia trachomatis*. *Nat Commun* **9**:45. doi:10.1038/s41467-017-

1124   02432-0

1125

1126   45. Le Negrate G, Krieg A, Faustin B, Loeffler M, Godzik A, Krajewski S, Reed JC. 2008.

1127   ChlaDub1 of *Chlamydia trachomatis* suppresses NF-κB activation and inhibits IκBα

1128   ubiquitination and degradation. *Cell Microbiol* **10**:1879–1892. doi:10.1111/j.1462-

1129   5822.2008.01178.x

1130
1131   46. Li J, Chai Q-Y, Liu CH. 2016. The ubiquitin system: a critical regulator of innate immunity

1132   and pathogen-host interactions. *Cell Mol Immunol* **13**:560–576. doi:10.1038/cmi.2016.40

1133

1134   47. Li Z, Chen C, Chen D, Wu Y, Zhong Y, Zhong G. 2008. Characterization of fifty putative

1135   inclusion membrane proteins encoded in the *Chlamydia trachomatis* genome. *Infect*

1136   *Immun* **76**:2746–2757. doi:10.1128/IAI.00010-08

1137

48. Lowden NM, Yeruva L, Johnson CM, Bowlin AK, Fisher DJ. 2015. Use of aminoglycoside 3'

adenyltransferase as a selection marker for *Chlamydia trachomatis* intron-mutagenesis and

in vivo intron stability. *BMC Res Notes* **8**:570. doi:10.1186/s13104-015-1542-9

1141

49. Lutter EI, Martens C, Hackstadt T. 2012. Evolution and conservation of predicted inclusion

membrane proteins in *Chlamydiae. Comp Funct Genomics* **2012**:362104.

doi:10.1155/2012/362104

1145

50. Lutter EI, Barger AC, Nair V, Hackstadt T. 2013. *Chlamydia trachomatis* inclusion

membrane protein CT228 recruits elements of the myosin phosphatase pathway to regulate

release mechanisms. *Cell Rep* **3**:1921–1931. doi:10.1016/j.celrep.2013.04.027

1149

51. Makarova KS, Aravind L, Koonin EV. 2000. A novel superfamily of predicted cysteine

proteases from eukaryotes, viruses and *Chlamydia pneumoniae. Trends Biochem

Sci* **25**:50–52. doi:10.1016/s0968-0004(99)01530-3

1153

52. Meier K, Jachmann LH, Türköz G, Babu Sait MR, Pérez L, Kepp O, Valdivia RH, Kroemer

G, Sixt BS. 2023. The *Chlamydia* effector CpoS modulates the inclusion microenvironment

and restricts the interferon response by acting on Rab35. *MBio* **14**:e0319022.

doi:10.1128/mbio.03190-22

1158
53. Meier K, Jachmann LH, Pérez L, Kepp O, Valdivia RH, Kroemer G, Sixt BS. 2022.

The *Chlamydia* protein CpoS modulates the inclusion microenvironment and restricts the

interferon response by acting on Rab35. *BioRxiv*. doi:10.1101/2022.02.18.481055

1162

1163   54. Misaghi S, Balsara ZR, Catic A, Spooner E, Ploegh HL, Starnbach MN. 2006. *Chlamydia*

1164   *trachomatis*-derived deubiquitinating enzymes in mammalian cells during infection. *Mol*

1165   *Microbiol* **61**:142–150. doi:10.1111/j.1365-2958.2006.05199.x

1166

1167   55. Mittal R, Peak-Chew S-Y, McMahon HT. 2006. Acetylation of MEK2 and IκB kinase (IKK)

1168   activation loop residues by YopJ inhibits signaling. *Proc Natl Acad Sci USA* **103**:18574–

1169   18579. doi:10.1073/pnas.0608995103

1170

1171   56. Mital J, Miller NJ, Fischer ER, Hackstadt T. 2010. Specific chlamydial inclusion membrane

1172   proteins associate with active Src family kinases in microdomains that interact with the host

1173   microtubule network. *Cell Microbiol* **12**:1235–1249. doi:10.1111/j.1462-5822.2010.01465.x

1174

1175   57. Mizuno E, Kitamura N, Komada M. 2007. 14-3-3-dependent inhibition of the deubiquitinating

1176   activity of UBPY and its cancellation in the M phase. *Exp Cell Res* **313**:3624–3634.

1177   doi:10.1016/j.yexcr.2007.07.028

1178

1179   58. Moulder JW. 1991. Interaction of *Chlamydiae* and host cells in vitro. *Microbiol Rev* **55**:143–

1180   190.

1181

1182   59. Mukherjee S, Keitany G, Li Y, Wang Y, Ball HL, Goldsmith EJ, Orth K. 2006. *Yersinia* YopJ

1183   acetylates and inhibits kinase activation by blocking phosphorylation. *Science* **312**:1211–

1184   1214. doi:10.1126/science.1126867

1185

1186   60. Neunuebel MR, Chen Y, Gaspar AH, Backlund PS, Yergey A, Machner MP. 2011. De-

1187   AMPylation of the small GTPase Rab1 by the pathogen *Legionella*

1188   *pneumophila. Science* **333**:453–456. doi:10.1126/science.1207193

1189

61. Nguyen BD, Valdivia RH. 2012. Virulence determinants in the obligate intracellular pathogen

    Chlamydia trachomatis revealed by forward genetic approaches. *Proc Natl Acad Sci*

    USA **109**:1263–1268. doi:10.1073/pnas.1117884109

1193

62. Nguyen PH, Lutter EI, Hackstadt T. 2018. *Chlamydia trachomatis* inclusion membrane

    protein MrcA interacts with the inositol 1,4,5-trisphosphate receptor type 3 (ITPR3) to

    regulate extrusion formation. *PLoS Pathog***14**:e1006911. doi:10.1371/journal.ppat.1006911

1197

63. Ohtake F, Tsuchiya H. 2017. The emerging complexity of ubiquitin architecture. *J*

    Biochem **161**:125–133. doi:10.1093/jb/mvw088

1200

64. Pannekoek Y, Spaargaren J, Langerak AAJ, Merks J, Morré SA, van der Ende A. 2005.

    Interrelationship between polymorphisms of *incA*, fusogenic properties of *Chlamydia*

    trachomatis strains, and clinical manifestations in patients in The Netherlands. *J Clin*

    Microbiol **43**:2441–2443. doi:10.1128/JCM.43.5.2441-2443.2005

1205

65. Peng J, Schwartz D, Elias JE, Thoreen CC, Cheng D, Marsischky G, Roelofs J, Finley D,

    Gygi SP. 2003. A proteomics approach to understanding protein ubiquitination. *Nat*

    Biotechnol **21**:921–926. doi:10.1038/nbt849

1209

66. Pruneda JN, Durkin CH, Geurink PP, Ovaa H, Santhanam B, Holden DW, Komander D.

    2016. The Molecular Basis for Ubiquitin and Ubiquitin-like Specificities in Bacterial Effector

    Proteases. *Mol Cell* **63**:261–276. doi:10.1016/j.molcel.2016.06.015

1213

1214   67. Pruneda JN, Bastidas RJ, Bertsoulaki E, Swatek KN, Santhanam B, Clague MJ, Valdivia

1215   RH, Urbé S, Komander D. 2018. A *Chlamydia* effector combining deubiquitination and

1216   acetylation activities induces Golgi fragmentation. *Nat Microbiol* **3**:1377–1384.

1217   doi:10.1038/s41564-018-0271-y

1218

1219   68. Reiley W, Zhang M, Wu X, Granger E, Sun S-C. 2005. Regulation of the deubiquitinating

1220   enzyme CYLD by IκB kinase gamma-dependent phosphorylation. *Mol Cell Biol* **25**:3886–

1221   3895. doi:10.1128/MCB.25.10.3886-3895.2005

1222

1223   69. Roan NR, Gierahn TM, Higgins DE, Starnbach MN. 2006. Monitoring the T cell response to

1224   genital tract infection. *Proc Natl Acad Sci USA* **103**:12069–12074.

1225   doi:10.1073/pnas.0603866103

1226

1227   70. Rockey DD, Scidmore MA, Bannantine JP, Brown WJ. 2002. Proteins in the chlamydial

1228   inclusion membrane. *Microbes Infect* **4**:333–340. doi:10.1016/s1286-4579(02)01546-0

1229

1230   71. Ronau JA, Beckmann JF, Hochstrasser M. 2016. Substrate specificity of the ubiquitin and

1231   Ubl proteases. *Cell Res* **26**:441–456. doi:10.1038/cr.2016.38

1232

1233   72. Rytkönen A, Poh J, Garmendia J, Boyle C, Thompson A, Liu M, Freemont P, Hinton JCD,

1234   Holden DW. 2007. SseL, a *Salmonella* deubiquitinase required for macrophage killing and

1235   virulence. *Proc Natl Acad Sci USA***104**:3502–3507. doi:10.1073/pnas.0610095104

1236

1237   73. Saeki Y. 2017. Ubiquitin recognition by the proteasome. *J Biochem* **161**:113–124.

1238   doi:10.1093/jb/mvw091

1239

1240   74. Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image

1241   analysis. *Nat Methods* **9**:671–675. doi:10.1038/nmeth.2089

1242

1243   75. Schubert AF, Nguyen JV, Franklin TG, Geurink PP, Roberts CG, Sanderson DJ, Miller LN,

1244   Ovaa H, Hofmann K, Pruneda JN, Komander D. 2020. Identification and characterization of

1245   diverse OTU deubiquitinases in bacteria. *EMBO J* **39**:e105127.

1246   doi:10.15252/embj.2020105127

1247

1248   76. Shaw JH, Key CE, Snider TA, Sah P, Shaw EI, Fisher DJ, Lutter EI. 2018. Genetic

1249   Inactivation of *Chlamydia trachomatis* Inclusion Membrane Protein CT228 Alters MYPT1

1250   Recruitment, Extrusion Production, and Longevity of Infection. *Front Cell Infect*

1251   *Microbiol* **8**:415. doi:10.3389/fcimb.2018.00415

1252

1253   77. Sheedlo MJ, Qiu J, Tan Y, Paul LN, Luo Z-Q, Das C. 2015. Structural basis of substrate

1254   recognition by a bacterial deubiquitinase important for dynamics of phagosome

1255   ubiquitination. *Proc Natl Acad Sci USA* **112**:15090–15095. doi:10.1073/pnas.1514568112

1256

1257   78. Sixt BS, Bastidas RJ, Finethy R, Baxter RM, Carpenter VK, Kroemer G, Coers J, Valdivia

1258   RH. 2017. The *Chlamydia trachomatis* Inclusion Membrane Protein CpoS Counteracts

1259   STING-Mediated Cellular Surveillance and Suicide Programs. *Cell Host Microbe* **21**:113–

1260   121. doi:10.1016/j.chom.2016.12.002

1261

1262   79. Smith EP, Cotto-Rosario A, Borghesan E, Held K, Miller CN, Celli J. 2020. Epistatic Interplay

1263   between Type IV Secretion Effectors Engages the Small GTPase Rab2 in the *Brucella*

1264   Intracellular Cycle. *MBio* **11**. doi:10.1128/mBio.03350-19

1265

1266 80. Suchland RJ, Rockey DD, Bannantine JP, Stamm WE. 2000. Isolates of *Chlamydia*

1267 *trachomatis* that occupy nonfusogenic inclusions lack IncA, a protein localized to the

1268 inclusion membrane. *Infect Immun* **68**:360–367. doi:10.1128/IAI.68.1.360-367.2000

1269

1270 81. Swatek KN, Komander D. 2016. Ubiquitin modifications. *Cell Res* **26**:399–422.

1271 doi:10.1038/cr.2016.39

1272

1273 82. Tenno T, Fujiwara K, Tochio H, Iwai K, Morita EH, Hayashi H, Murata S, Hiroaki H, Sato M,

1274 Tanaka K, Shirakawa M. 2004. Structural basis for distinct roles of Lys63- and Lys48-linked

1275 polyubiquitin chains. *Genes Cells***9**:865–875. doi:10.1111/j.1365-2443.2004.00780.x

1276

1277 83. Urbanus ML, Quaile AT, Stogios PJ, Morar M, Rao C, Di Leo R, Evdokimova E, Lam M,

1278 Oatway C, Cuff ME, Osipiuk J, Michalska K, Nocek BP, Taipale M, Savchenko A, Ensminger

1279 AW. 2016. Diverse mechanisms of metaeffector activity in an intracellular bacterial

1280 pathogen, *Legionella pneumophila. Mol Syst Biol* **12**:893. doi:10.15252/msb.20167381

1281

1282 84. Varadan R, Walker O, Pickart C, Fushman D. 2002. Structural properties of polyubiquitin

1283 chains in solution. *J Mol Biol* **324**:637–647. doi:10.1016/s0022-2836(02)01198-1

1284

1285 85. Vozandychova V, Stojkova P, Hercik K, Rehulka P, Stulik J. 2021. The Ubiquitination

1286 System within Bacterial Host-Pathogen Interactions. *Microorganisms* **9**.

1287 doi:10.3390/microorganisms9030638

1288

1289 86. Walsh SC, Reitano JR, Dickinson MS, Kutsch M, Hernandez D, Barnes AB, Schott BH,

1290 Wang L, Ko DC, Kim SY, Valdivia RH, Bastidas RJ, Coers J. 2022. The bacterial effector

1291   GarD shields *Chlamydia trachomatis* inclusions from RNF213-mediated ubiquitylation and

1292   destruction. *Cell Host Microbe*. doi:10.1016/j.chom.2022.08.008

1293
1294 87. Wang X, Hybiske K, Stephens RS. 2018. Direct visualization of the expression and

1295   localization of chlamydial effector proteins within infected host cells. *Pathog Dis* **76**.

1296   doi:10.1093/femspd/fty011

1297

1298 88. Weber MM, Bauler LD, Lam J, Hackstadt T. 2015. Expression and localization of predicted

1299   inclusion membrane proteins in *Chlamydia trachomatis*. *Infect Immun* **83**:4710–4718.

1300   doi:10.1128/IAI.01075-15

1301

1302 89. Weeks SD, Grasty KC, Hernandez-Cuebas L, Loll PJ. 2009. Crystal structures of Lys-63-

1303   linked tri- and di-ubiquitin reveal a highly extended chain architecture. *Proteins* **77**:753–759.

1304   doi:10.1002/prot.22568

1305

1306 90. Wesolowski J, Weber MM, Nawrotek A, Dooley CA, Calderon M, St Croix CM, Hackstadt T,

1307   Cherfils J, Paumet F. 2017. *Chlamydia* hijacks ARF gtpases to coordinate microtubule

1308   posttranslational modifications and golgi complex positioning. *MBio* **8**.

1309   doi:10.1128/mBio.02280-16

1310

1311 91. Zadora PK, Chumduri C, Imami K, Berger H, Mi Y, Selbach M, Meyer TF, Gurumurthy RK.

1312   2019. Integrated Phosphoproteome and Transcriptome Analysis Reveals *Chlamydia*-

1313   Induced Epithelial-to-Mesenchymal Transition in Host Cells. *Cell Rep* **26**:1286-1302.e8.

1314   doi:10.1016/j.celrep.2019.01.006

1315

1316   92. Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C, Chanda

1317   SK. 2019. Metascape provides a biologist-oriented resource for the analysis of systems-

1318   level datasets. *Nat Commun* **10**:1523. doi:10.1038/s41467-019-09234-6

1319

1320   93. Zuck M, Ellis T, Venida A, Hybiske K. 2017. Extrusions are phagocytosed and promote

1321   *Chlamydia* survival within macrophages. *Cell Microbiol* **19**. doi:10.1111/cmi.12683

1322

1323   94. Zuck M, Sherrid A, Suchland R, Ellis T, Hybiske K. 2016. Conservation of extrusion as an

1324   exit mechanism for *Chlamydia*. *Pathog Dis* **74**. doi:10.1093/femspd/ftw093

1325

1326   **Figures**

1327
1328
Figure 1
The *C. trachomatis* inclusion membrane proteins InaC, IpaM, and CTL0480 are
ubiquitinated in the absence of Cdu1. (A-C) Volcano plots (pairwise comparisons) of the
relative abundance of human and Ct ubiquitinated peptides.**(A)** Mock infected HeLa cells versus
HeLa cells infected with WT Ct (L2 434 Bu pBOMB) (24 hpi). **(B)** Mock infected HeLa cells
versus HeLa cells infected with a *cdu1* null strain (*cdu1*::GII pBOMB) (24 hpi) **(C)** HeLa cells
infected with WT Ct (24 hpi) versus HeLa cells infected with a *cdu1* null strain (24 hpi).
Significance values were interpolated from 3 independent biological replicates. Ubiquitinated
proteins were enriched with magnetic TUBE 1 beads (binds to polyubiquitinated proteins) and
peptides identified by quantitative LC MS/MS analysis. Three Ct inclusion membrane proteins,
InaC, IpaM, and CTL0480 were differentially ubiquitinated in the absence of Cdu1. **(D)** InaC was
ubiquitinated at K104, K107, and K149, IpaM at K290, and CTL0480 at K115 in the absence of
Cdu1. TMD: Transmembrane domain. Numbering corresponds to amino acids in the protein
sequence of each respective inclusion membrane protein.
1343
1344

1345
Figure 2
Cdu1 associates with InaC, IpaM, and CTL0480. (A) Co-localization of Cdu1(magenta) with
endogenous InaC (green), IpaM (green), and ectopically expressed CTL0480-Flag (green) at
the Ct (L2) inclusion membrane of HeLa cells infected for 24 h. HeLa cells infected with an *inaC*
null strain (M407), an *ipaM* null strain (*ipaM*::GII), and WT Ct (L2 434 Bu pBOMB) were used as
controls for antibody specificity. DNA stained with Hoechst is shown in blue. Scale bar: 10μm.
Images are representative of multiple images captured across three independent replicates. **(B)**
Line scan profiles of fluorescent signal intensities displayed in (A) showing co-localization of
fluorescence intensities for endogenous Cdu1 with endogenous InaC and IpaM, and with
CTL0480-Flag along the L2 inclusion membrane. **(C)** Schematic of Cdu1-GFP(C) (L2) fusion
(Cdu1-GFP) and Cdu1-GFP variants used in co-transfections of HEK 293 cells. GFP: Green
fluorescent protein. TMD: Transmembrane domain. PRD: Proline rich domain. CD: Catalytic

domain. FL: Full length. TMD-: Cdu1-GFP variant lacking TMD domain. CD-: Cdu1-GFP variant
lacking CD domain. **(D-G)** Western blot analysis of GFP immunoprecipitates from HEK 293 cells
co-transfected with mammalian plasmids expressing: Cdu1-GFP variants and **(D)** truncated
3XFlag(N)-InaC (D/UW-3/CX CT813, amino acids 96-264), **(E)** V5(N)-IpaM (L2, full length), **(F)**
V5(N)-CTL0480 (L2, full length), and **(G)** V5(N)-CpoS (L2, full length). Vec1: Empty pOPINN-
GFP vector. Vec2: Empty pDEST53 vector. Vec3: Empty pcDNA™3.1/nV5-DEST vector.
Western blot images are representative from two independent experiments.

1370
Figure 3
Cdu1 stabilizes InaC, IpaM, and CTL0480. (A) Western blot analysis of endogenous InaC in
HeLa cells infected for 24, 36, and 48 hours with Wt Ct (L2 pBOMB), *cdu1* null (*cdu1*::GII
pBOMB), and *inaC* null (M407) strains. Ct Slc1 and human alpha tubulin were used to
determine Ct burdens and equal loading of protein extracts respectively. Western blot images
are representative of 2 independent experiments. **(B)** Quantification of InaC abundance (InaC
western blot signal from (A)) normalized to Slc1 western blot signal (from panel A) in Hela cells
infectedwith a *cdu1* null strain, relative to normalized InaC abundance in Hela cells infected with
Wt Ct. **(C)** Western blot analysis of endogenous IpaM in HeLa cells infected for 24, 36, and 48
hours with Wt Ct, *cdu1* null, and *ipam* null (*ipam*::GII) strains. Western blot images are
representative of 2 independent experiments. **(D)** Quantification of normalized IpaM abundance

(from (C) in Hela cells infectedwith a *cdu1* null strain, relative to normalized IpaM abundance in
Hela cells infected with Wt Ct. **(E)** Localization of CTL0480 during Ct infection of HeLa cells at
24, 36, and 48 hpi. CTL0480 signal (green) co-localizes with the inclusion membrane protein
IncA (magenta) at the Ct inclusion membrane. Arrowheads highlight *cdu1* null inclusions lacking
CTL0480 at 36 hpi. DNA stained with Hoechst is shown in blue. Scale bar: 10µm. Quantification
of MYPT1 localization at inclusion membranes can be found in Supplemental Figure 5

1388
1389
1390

1391
Figure 4
The acetylase activity of Cdu1 is required to stabilize Cdu1, InaC, and IpaM. (A) Western
blot analysis of endogenous InaC and Cdu1-Flag catalytic variants expressed from a plasmid
(pBOMB). HeLa cells were infected for 36 hours with WT Ct (L2 434 Bu pBOMB), a *cdu1* null
strain (*cdu1*::GII pBOMB), and *cdu1* null strains expressing wild type Cdu1-Flag and the Cdu1
variants C345A-Flag (catalytic inactive), I225A-Flag (DUB deficient), and K268E-Flag (Act

deficient). Cdu1-Flag variants were expressed from a pBOMB shuttle plasmid. Protein lysates
from HeLa cells infected with an *inaC* null (M407) strain were used to control for the specificity
of anti-InaC antibodies. Western blot images are representative of two independent
experiments. **(B)** Western blot analysis of endogenous IpaM and Cdu1-FLAG variants in crude
extracts of HeLa cells infected for 48 hours with the same strains as describe in (A). Infection of
HeLa cells with *ipaM*::GII was used to test for the specificity of the anti-IpaM antibody. Western
blot images are representative of two independent experiments. **(C)** Western blot analysis of
Cdu1^{C345A}-Flag (catalytic inactive) and Cdu1^{K268E}-Flag (Act deficient) expressed in a *cdu1* null
strain or WT Ct (L2 434 Bu) background after infection of HeLa cells for 24 hours. Both Cdu1
variants are stabilized by Cdu1 expressed in WT Ct. **(D)** Western blot analysis of Cdu1-Flag and
Cdu1^{C345A}-Flag expressed in a *cdu1* null strain and Cdu1^{C345A}-Flag expressed in WT Ct (L2
434 Bu) following immunoprecipitation (anti-Flag) from HeLa cell extracts after infection for 24
hours and treatment with MG132 (25 μM, 5 hours). Western blot image is a representative blot
from at least three independent experiments. **(E)** Western blot analysis of endogenous InaC and
IpaM following immunoprecipitation of inclusion membrane enriched subcellular fractions
(24hpi) with anti acetylated lysine antibodies. Western blot image is representative of two
independent experiments. **(F)** Western blot analysis (Flag WB) of acetylated lysine
immunoprecipitates generated from inclusion membrane enriched subcellular fractions (40 hpi)
derived from HeLa cells infected with WT Ct strains expressing Cdu1-Flag, CTL0480-Flag, or
IpaM-Flag. Western blot image is representative of two independent experiments. **(G)** WB of
acetylated lysine immunoprecipitates of inclusion membrane enriched fractions (24 hpi) of HeLa
cells infected with WT Ct and with an *inaC* null strain (M407) expressing InaC-Flag. Western
blot image is representative of two independent experiments. **(H)** The initiator methionine,
Lys10, Lys126, and Lys263 of Cdu1 are acetylated by 24 hpi. One tyrosine (Y) residue and
multiple serine (S) and threonine (T) residues in Cdu1 are also phosphorylated during Ct
infection of HeLa cells (24 hpi). Three PX(S/T)P MAPK phosphorylation consensus sequence
motifs were identified in the proline rich domain of Cdu1. Modified residues were identified by
quantitative LC MS/MS analysis of immunoprecipitated Cdu1-Flag across 3 independent
biological replicates. TMD: Transmembrane domain. PR: Proline rich. PSTP: PX(S/T)P motifs.
Ac: Acetylation. P: Phosphorylation. Numbering corresponds to amino acids in Cdu1 protein
sequence.
1429
1430

1431
1432
Figure 5
The DUB activity of Cdu1 is not required for blocking the ubiquitination of inclusion
membranes and Cdu1 is not required for protection against IFNγ mediated cellular
immunity. (A) Representative images of Ct inclusions decorated with ubiquitin during infection
of HeLa cells with a *cdu1*::GII strain for 24 hours. Representative images of infected Hela cells
used for quantification of ubiquitin decorated inclusions in (B) are shown in Supplemental Figure
6A. Antisera against the membrane protein Cap1 (green) was used to mark inclusion

membranes. DNA stained with Hoechst is shown in blue. Scale bar: 10μm. **(B)** Quantification of
ubiquitinated inclusions as shown in (A). The Ub fluorescent signal was used to determine the
number of infected cells with Ub decorated inclusions. The total number of ubiquitinated
inclusions was divided by the total number of inclusions analyzed (defined by Cap1 staining).
87%, 86%, and 91% of inclusions were decorated with Ub in HeLa cells infected with a *cdu1*
null strain and *cdu1* null strains expressing Cdu1^{C345A} (DUB-, Act-), and Cdu1^{K268E} (Act-)
variants respectively. Representative images (panel (A) and Supplemental Figure 6A) and
quantification of ubiquitinated inclusions were obtained from inclusions imaged in 10 fields
across 3 independent biological replicates for each strain. *p* values were determined by a
student paired t-test. **(C)** Representative images of HeLa cells co-infected with *cdu1*::GII
(*cdu1Δ*) and *incA* null (*incAΔ*, M923) strains at 24 hpi. IncA-deficient inclusions do not fuse with
other inclusions. In co-infected cells, Cdu1 present on the inclusion membranes of *incAΔ* strains
did not block ubiquitination events at or near the inclusion membranes of neighboring *cdu1Δ*
strains (mCherry signal from pBOMB4-MCI plasmid). Representative images of Hela cells
infected with strains quantified in (D) are shown in Supplemental Figure 6B. DNA stained with
Hoechst is shown in blue. Scale bar: 10μm. **(D)** Quantification of *cdu1Δ* inclusions as shown in
(C) in which ubiquitination events are observed in regions of *cdu1Δ* inclusion membranes that
are in direct apposition to *incAΔ* inclusion membranes (junctions). The total number of *cdu1Δ*
inclusions ubiquitinated at inclusion junctions was divided by the total number of inclusions
analyzed (Cap1 staining). 66% and 61% of *cdu1Δ* inclusions were decorated with Ub at
junctions in HeLa cells co-infected with *incAΔ* and *cdu1Δ* strains or with *incAΔ* and a *cdu1Δ*
strain ectopically expressing Cdu1^{K268E} (Act-) respectively. Representative images (panel (C)
and Supplemental Figure 6B) and quantification of *cdu1Δ* inclusions ubiquitinated at junctions
are derived from inclusions imaged in 6 fields across 3 independent biological replicates for
each condition. *p* values were determined by a student paired t-test. **(E)** Quantification of Ct
inclusion production during infection of unprimed and IFNγ-primed (100 U/mL) A549 cells at 24
hours post infection. Inclusions were quantified by high-content imaging analysis. Plot reflects
inclusion counts across 9 fields of view and 3 independent biological replicates. Inclusion counts
by each strain in unprimed A549 cells were set to 100%. Inclusion counts resulting from
cdu1::GII and *garD*::GII strains were normalized to corresponding parental Ct inclusion (100%)
counts in unprimed cells. P-values were calculated by 2 way ANOVA analysis. **(F)**
Representative images of RNF213 localizing to inclusions of WT Ct, *garD*::GII, and *cdu1*::GII
strains during infection of A549 cells primed with IFNγ (100 U/mL). **(G)** Quantification of RNF213
localizing to Ct inclusions during infection of unprimed and IFNγ-primed (100 U/mL) A549 cells
at 24 hours post infection. Plot reflects inclusion counts across 6 fields and 3 independent
biological replicates.
1476
1477

Figure 6
Cdu1 is required for assembly of F-actin, Golgi ministack repositioning, and MYPT1
recruitment to the inclusion.
(A) Examples of representative images of F-actin (arrowheads) (green, Alexa Fluor^{TM}
Phalloidin) assembled around the Ct inclusion (magenta, anti Cdu1 and Cap1 staining) in HeLa
cells infected for 40 hours. Representative images for each strain analyzed can be found in
Supplemental Figure 7. **(B)** Quantification of Ct inclusion surrounded by F-actin normalized to
the total number of inclusions analyzed during infection of Hela cells at 40 hpi. Quantification of
surrounding F-actin were obtained from inclusions imaged in 6 fields across 3 independent
biological replicates. *p* values were determined by one-way ANOVAs with a Student-Newman-
Keuls post hoc test. Strains used: WT L2 (Rif-R 434 Bu, parent of M407), M407 (*inaC* null
strain) p2TK2, M407 p2TK2-InaC, WT L2 (434 Bu) pBOMB, *cdu1*::GII pBOMB, *cdu1*::GII
pBOMB-Cdu1 Flag, *cdu1*::GII pBOMB-Cdu1^{C345A} Flag, *cdu1*::GII pBOMB-Cdu1^{I225A} Flag, and
cdu1::GII pBOMB-Cdu1^{K268E} Flag. **(C)** Sample representative images of Golgi (anti GM130
staining, magenta) around Ct inclusions in HeLa cells infected for 24 hours. Representative
images for each strain analyzed can be found in Supplemental Figure 8. **(D)** Quantification of
Golgi dispersal around the Ct inclusion during infection of Hela cells for 24 hpi. The length of
Golgi dispersed around each Ct inclusion imaged was measured and normalized to the
perimeter length of each inclusion (% Golgi dispersion around the inclusion). Golgi dispersal
around Ct inclusions was quantified from inclusions imaged in 6 fields across 3 independent
biological replicates. *p* values were determined by a student paired t-test. Strains analyzed were
the same ones as mentioned in (B). **(E)** Representative images of MYPT1 (magenta) at Ct
inclusions (green, anti-IncA staining). Arrowheads represent *cdu1* null inclusions with low
MYPT1 signal. DNA stained with Hoechst is shown in blue in panels C and E. Scale bar: 10μm.
(F) Quantification of MYPT1 at Ct inclusions as shown in (E). Representative images in (E) and
quantification of MYPT1 recruitment in (F) were obtained from inclusions imaged in 6 fields
across 3 independent replicates. *p* values were determined by a student paired t-test.
1509
1510

1511
Figure 7
Cdu1, InaC, and IpaM are required for optimal extrusion of Ct inclusions from HeLa cells.
(A) Representative images of extrusions isolated from HeLa cell monolayers infected with Ct
strains for 52 hours. Scale bar: 200 µm **(B)** Quantification of the number of extruded inclusions
produced by infected HeLa cell monolayers. *p* values were determined by a student paired t-
test. **(C)** Quantification of the size of extruded inclusions quantified in (B). Extruded inclusions
varied in size among cells infected with wild type L2 (average :43 µm), *ipaM* mutants (average:
56 µm) and *cdu1* mutants complemented wild type Cdu1 (average: 52 µm) and Cdu1^{I225A}
(DUB-) (average: 60 µm). *p* values were determined by one-way ANOVAs with a Student-
Newman-Keuls post hoc test. Representative images in (A) and quantification of extrusion
number in (B) and extrusion size in (C) are based on images obtained from 6 fields across 3
independent biological replicates.
1524

1525
1526
Figure 8.
A model for acetylation mediated protection of the Inc proteins InaC, IpaM, and Ctl0480
from degradation.
The cellular Ub machinery targets *C. trachomatis* effectors, including the Inc proteins InaC,
IpaM, and CTL0480 for ubiquitination and subsequent protein degradation. *C. trachomatis*
counters such defense mechanisms by translocating Cdu1 which protect itself and all three Inc
proteins from being targeted for ubiquitination and degradation through its acetylase activity.
Cdu1 protects InaC and enables the recruitment of F-actin scaffolds and Golgi ministacks to the
inclusion perimeter and CTL0480 to recruit the Myosin II regulator MYPT1. All three inclusion
proteins and Cdu1 promote extrusion and dissemination of *C. trachomatis* inclusions late in
infection. Image generated with BioRender.com.
1538
1539

A

WT L2

cdu1 (1.3 kb)

cdu1 ORF

1   750 bp

cdu1*::GII *aadA

cdu1 (3 kb)

5' junction (1 kb)   *aadA* (0.4 kb)   3' junction (0.3 kb)

cdu1 ORF   *aadA*

1   635   2,831
                    GII (*aadA*) intron   bp

B

1540

Supplemental Figure 1
TargeTron mediated disruption of the L2 *cdu1* ORF. (A) Depiction of the *cdu1* ORFs in WT
L2 and *cdu1*::GII strains with corresponding diagnostic PCR amplicons. (**B**) PCR analysis of the
disrupted *cdu1* ORF in the *cdu1*::GII strain. *Amplification of *cdu1* ORF in *cdu1*::GII strain was
unsuccessful due to size of amplicon. + control: WT *cdu2*.

1546

1547

1548
Supplemental Figure 2
Generation of a *cdu1* null strain in *C. trachomatis* (L2). (**A**) Western blot analysis of protein
lysates derived from HeLa cells infected with a Ct L2 434 Bu parental strain (L2) and an L2
cdu1::GII *aadA* (*cdu1*::GII) strain for 24 hours. Blots were probed with antibodies raised against
recombinant Cdu1 (amino acids 71-401), Ct RpoB (bacterial RNA polymerase), and human
α tubulin. (**B**) HeLa cells infected with L2 transformed with pBOMB4-MCI empty vector (L2
pBOMB) and *cdu1*::GII transformed with empty pBOMB4-MCI (*cdu1*::GII pBOMB) were fixed at
24 hpi and stained with Cdu1 antisera. Cdu1 signal is depicted in green, Ct cells expressing
mCherry (encoded in pBOMB4-MCI vector) are shown in red, and DNA stained with Hoechst in
blue. Scale bar: 10μm. (**C**) Total RNA isolated from HeLa cells infected with L2 and *cdu1*::GII
strains for 24 hours was used to synthesize cDNAs which served as templates for PCR analysis
of *cdu1*, *cdu2*, and *cdu1-cdu2* bicistronic expression. RT+: Total RNA + reverse transcriptase.
RT-: Total RNA without reverse transcriptase. gDNA: Genomic DNA.
1562

1563
Supplemental Figure 3
Proteins co-precipitating (non ubiquitinated) with human and Ct proteins enriched by
TUBE 1 pulldowns. (A) Volcano plots (pairwise comparisons) of human proteins (non-
ubiquitinated) that co-precipitate with TUBE1 bound proteins in mock infected HeLa cells and
HeLa cells infected with L2 or *cdu1* null strains (24 hpi). **(B)** Volcano plots (pairwise

comparisons) of Ct TUBE1 co-precipitating proteins (non-ubiquitinated) identified in mock
infected HeLa cells and HeLa cells infected with L2 or *cdu1* null strains (24 hpi).

A

B

C

1585
Supplemental Figure 4
Pathway enrichment analysis of human and Ct proteins that co-precipitate (non
ubiquitinated) with TUBE1 bound proteins. (A) Metascape functional enrichment analysis of
human co-precipitating proteins in mock infected HeLa cells. **(B)** Metascape functional
enrichment analysis of human co-precipitating proteins in *cdu1* null infected HeLa cells. **(C)**
DAVID pathway enrichment analysis of Ct co-precipitating proteins enriched in infected (L2 and
cdu1 null) HeLa cells.

1601
Supplemental Figure 5
Quantification of CTL0480 at Ct inclusion membranes.
Quantification of CTL0480 localization at Ct inclusion membranes as shown in Figure 3E.
CTL0480 fluorescent signal was used to determine the percent of inclusions displaying
CTL0480 at the inclusion membranes. Representative images and quantification of CTL0480
positive inclusions are derived from Ct inclusions imaged in 10 fields across 3 independent
biological replicates. *p* values were determined by a student paired t-test.
1609
1610
1611
1612

1613
Supplemental Figure 6.
The Acetylase activity of Cdu1 is the predominant activity of Cdu1 responsible for
protecting Ct inclusions from ubiquitination.
(A) Representative images of Ct inclusions (green, Cap1) decorated with Ubiquitin (magenta,
FK2 antibody) in HeLa cells infected for 24 hours. Strains used: WT Ct (L2 434 Bu), cdu1::GII
pBOMB MCI, *cdu1*::GII pBOMB MCI-Cdu1 Flag, *cdu1*::GII pBOMB MCI-Cdu1^{C345A} Flag,
cdu1::GII pBOMB MCI-Cdu1^{I225A} Flag, and *cdu1*::GII pBOMB MCI-Cdu1^{K268E} Flag. **(B)**
Representative images of Hela cells co-infected with an *incA* null strain and *cdu1*::GII strains
expressing Cdu1 variants for 24 hours. Strains used: *cdu1*::GII pBOMB MCI, *inaC* null (M923),
cdu1::GII pBOMB MCI-Cdu1 Flag, *cdu1*::GII pBOMB MCI-Cdu1^{C345A} Flag, *cdu1*::GII pBOMB
MCI-Cdu1^{I225A} Flag, and *cdu1*::GII pBOMB MCI-Cdu1^{K268E} Flag. DNA stained with Hoechst is
shown in blue. Scale bar: 10µm. MCI= mCherry.
1626

1627

Supplemental Figure 7
The DUB activity of Cdu1 is not required for assembly of F-actin around the Ct inclusion.
Representative images of F-actin (arrowheads) (green, Alexa Fluor™ Phalloidin) assembled
around the Ct inclusion (magenta, anti Cdu1 and Cap1 staining) in HeLa cells infected for 40
hours. Strains used: WT L2 (Rif-R 434 Bu, parent of M407), M407 (*inaC* null strain) p2TK2,
M407 p2TK2-InaC, WT L2 (434 Bu) pBOMB, *cdu1*::GII pBOMB, *cdu1*::GII pBOMB-Cdu1 Flag,
cdu1::GII pBOMB-Cdu1^{C345A} Flag, *cdu1*::GII pBOMB-Cdu1^{I225A} Flag, and *cdu1*::GII pBOMB-
Cdu1^{K268E} Flag. DNA stained with Hoechst is shown in blue. Scale bar: 10µm.

1649
1650
Supplemental Figure 8
The DUB activity of Cdu1 is not required for Golgi ministack repositioning around the Ct
inclusion.
Representative images of Golgi (anti GM130 staining, magenta) around Ct inclusions in HeLa
cells infected for 24 hours. Strains used: WT L2 (Rif-R 434 Bu, parent of M407), M407 (*inaC* null
strain) p2TK2, M407 p2TK2-InaC, WT L2 (434 Bu) pBOMB, *cdu1*::GII pBOMB, *cdu1*::GII
pBOMB-Cdu1 Flag, *cdu1*::GII pBOMB-Cdu1[C345A] Flag, *cdu1*::GII pBOMB-Cdu1[I225A] Flag, and
cdu1::GII pBOMB-Cdu1[K268E] Flag. DNA stained with Hoechst is shown in blue. Scale bar: 10μm.
1659

1660
1661 **Supplemental Figure 9**
1662 **The number of inclusions and the size of inclusions in *cdu1* null, *inaC* null, *ipaM* null, and**
1663 ***ctl0480* null strains are similar across each strain (A)** Representative images of inclusions in
1664 infected HeLa cell monolayers at 48 hpi. Scale bar: 200 μm **(B)** Quantification of the number of
1665 inclusions in infected HeLa cell monolayers. **(C)** Quantification of inclusion size.
1666
1667