1  **Photoreceptor genes in a trechine beetle, *Trechiama kuznetsovi*, living in the upper**
2  **hypogean zone**
3

4  Takuma Niida[1*], Yuto Terashima[1], Hitoshi Aonuma[2] and Shigeyuki Koshikawa[1,3*]
5
6  [1]Graduate School of Environmental Science, Hokkaido University, Sapporo, Japan
7  [2]Department of Biology, Graduate School of Science, Kobe University, Kobe, Japan
8  [3]Faculty of Environmental Earth Science, Hokkaido University, Sapporo, Japan
9
*authors for correspondence
Takuma Niida: ymbnw8bwgmxikb91sqck@gmail.com
Shigeyuki Koshikawa (submitting author): koshi@ees.hokudai.ac.jp

**Abstract**

To address how organisms adapt to a new environment, subterranean organisms whose

ancestors colonized subterranean habitats from surface habitats have been studied.

Photoreception abilities have been shown to have degenerated in organisms living in

caves and calcrete aquifers. Meanwhile, the organisms living in a shallow subterranean

environment, which are inferred to reflect an intermediate stage in an evolutionary

pathway to colonization of a deeper subterranean environment, have not been studied

well. In the present study, we examined the photoreception ability in a trechine beetle,

*Trechiama kuznetsovi*, which inhabits the upper hypogean zone and has a vestigial

compound eye. By *de novo* assembly of genome and transcript sequences, we were able

to identify photoreceptor genes and phototransduction genes. Specifically, we focused

on *opsin* genes, where one *long wavelength opsin* gene and one *ultraviolet opsin* gene

were identified. The encoded amino acid sequences had neither a premature stop codon

nor a frameshift mutation, and appeared to be subject to purifying selection.

Subsequently, we examined the internal structure of the compound eye and nerve tissue

in the adult head, and found potential photoreceptor cells in the compound eye and

nerve bundle connected to the brain. The present findings suggest that *T. kuznetsovi* has

retained the ability of photoreception. This species represents a transitional stage of

vision, in which the compound eye regresses, but it may retains the ability of

photoreception using the vestigial eye.

Key words: blind ground beetle, adaptation, evolution, Carabidae, mesovoid shallow

substratum

**Background**

How organisms adapt to a new environment is one of the fundamental research questions in evolutionary biology [1]. Subterranean organisms whose ancestors originally lived in a surface environment are ideal for investigating this issue [2, 3]. Subterranean habitats are not continuously exposed to light, and can be categorized into cave habitats, interstitial habitats and superficial subterranean habitats [4, 5]. Degeneration of eyes is generally observed in various taxa colonizing these subterranean environments [6, 7].

Do the organisms having a regressed eye also have decreased ability of photoreception? Previous studies focused on various aspects of subterranean adaptation, including signatures in photoreceptor proteins [8, 9]. For example, one previous study described the expression of visual opsin genes which encode seven-transmembrane photoreceptor proteins in the binocular eye of the Mexican blind cavefish, *Astyanax mexicanus* [8]. Transcripts of visual opsin genes were underrepresented in the cavefish as compared with a conspecific surface population, and this could be attributed to reduction of photoreceptor cells in the cavefish [10].

In Insecta, subterranean diving beetles (Dytiscidae), which have highly regressed or no eyes and inhabit a calcrete aquifer located 10 m underground in Western Australia, were subjected to a similar analysis [11]. Transcripts were not detected for *long wavelength opsin* or *ultraviolet opsin* at the adult stage of the diving beetles [12], and pseudogenization of *long wavelength opsin*, *ultraviolet opsin* and some phototransduction genes (i.e. *arrestin*, *neither inactivation nor afterpotential C*, *transient receptor potential* and *transient receptor potential channel-like*) was observed [13, 14].

Besides calcrete aquifers, insects have also been found to colonize superficial subterranean habitats, such as rock fissures near the surface [15]. Insects living in a superficial subterranean habitat can be exposed to light due to unexpected environmental fluctuation and migration of insects themselves toward the surface, unlike insects living in deep subterranean habitats. The colonization of a superficial subterranean habitat is inferred to reflect an intermediate stage in an evolutionary pathway to colonization of deeper and extreme environments [4]. Despite this importance, the biological features of species living in a superficial subterranean habitat remain unexplored.

The present study focused on one trechine beetle species, *Trechiama kuznetsovi* (Coleoptera: Carabidae: Trechinae). This species has a vestigial compound eye and inhabits the upper hypogean zone, which is a similar environment to mesovoid shallow

74 substratum (MSS) [16]. We aimed to reveal the photoreception ability in this species.
75 We obtained genome and transcript sequences to examine photoreceptor genes and
76 phototransduction genes and estimate selective pressure on visual opsin genes. Also,
77 histological investigations were performed to observe the internal structure of the
78 vestigial compound eye and a nerve bundle connecting it to the brain.
79
80

**Methods**
Sample collection
*Trechiama kuznetsovi* samples were collected at Yûbari City, Hokkaido, Japan.
We collected *T. kuznetsovi* adults from the upper hypogean zone consisting of small
rocks and clay, by digging in the slope of a v-shaped valley to the depth of a few to
some dozen centimeters (Fig. 1).
87

DNA and RNA sequencing
We used one adult male of *T. kuznetsovi* stored in 99.5% ethanol for genome
sequencing. Before DNA extraction, mites adhering to the body surface were removed
and the male genitalia was preserved in 99.5% ethanol for identification. Genomic DNA
was extracted using a Wizard Genomic DNA Purification Kit (Promega, Madison, WI,
USA). A library was constructed using a TruSeq Nano DNA Library Prep Kit (Illumina,
San Diego, CA, USA) and sequenced on the NovaSeq 6000 platform (Illumina) by
Macrogen Service (Macrogen, Seoul, South Korea). $2 \times 151$ bp paired-end reads were
generated (Table S1).
We used one live adult male of *T. kuznetsovi* for transcript sequencing. Before
RNA extraction, the beetle was washed with 99.5% ethanol, mites adhering on its body
surface were removed and the male genitalia was preserved in 99.5% ethanol for
identification. Total RNA was immediately extracted using an RNeasy Micro Kit
(Qiagen, Hilden, Germany). A library was constructed using a SMARTer Stranded
RNA-Seq Kit (Illumina) and sequenced on the NovaSeq 6000 platform (Illumina) by
Macrogen Service. $2 \times 101$ bp paired-end reads were generated (Table S1).
104

Assembly and mapping
Summary statistics of raw reads and adapter contamination were checked using
FastQC (v0.11.9; Babraham Institute). Quality control was performed using fastp
v0.20.1 [17] and Trimmomatic v0.39 [18] to trim off one base from the 3' end, low
quality sequences and adapter sequences. Then, summary statistics were rechecked

using FastQC. The kmer content of reads from genome sequencing and the genome size
were calculated using KmerGenie v1.7051 [19]. For the reads that passed the quality
control, genome and transcript assembly were conducted with Platanus v1.2.4 [20] and
Trinity v2.8.4 [21, 22]. In the scaffolding step of genome assembly, contigs smaller than
500 bp were excluded [23]. Completeness of the assembled genome and transcript was
assessed using BUSCO_v5 for insecta core gene sets and CEGMA for invertebrate core
gene sets in gVolante web server [24]. Summary statistics of the assembled sequences
were calculated using SeqKit v.0.16.1 [25].

RNA reads were mapped to the assembled genome scaffolds using HISAT
v2.1.0 [26]. The result of the mapping was visualized using IGV v2.11.9 [27].

Homology search

We conducted a search for genes encoding three photoreceptor proteins and 15
phototransduction proteins from the assembled genome and transcripts with tblastn
program using the following queries (unless noted, protein sequences of *Tribolium*
*castaneum* [Tenebrionidae] were used): Long wavelength opsin (Lw opsin) of *Pogonus*
*chalceus* (Carabidae: Trechinae), Ultraviolet opsin (Uv opsin) of *P. chalceus*, C-opsin,
Arrestin 1 (Arr1), Arrestin 2 (Arr2), G protein α-subunit 49B (Gα49B), G protein
β-subunit 76C (Gβ76C), G protein γ-subunit 30A (Gγ30A), G protein-coupled receptor
kinase 1 (Gprk1), Inactivation no afterpotential D (InaD), Neither inactivation nor
afterpotential A (NinaA), Neither inactivation nor afterpotential C (NinaC), No receptor
potential A (NorpA), Protein C kinase 53E (Pkc53E) of *Drosophila melanogaster*,
Rab-protein 6 (Rab6), Retinal degeneration C (RdgC), Transient receptor potential (Trp),
and Trp-like (Trpl). Selection of these proteins was based on previous studies [28, 29].
Blast-hit sequences with an e-value $< 1 \times e^{-20}$ were treated as having high similarity [30].
If no sequence matched this criterion, Blast-hit sequences were examined in order from
the best-hit sequence. In BLAST-search for the transcripts, the presence of premature
stop codons and frameshift mutations was examined.

We subsequently conducted a further analysis of *lw opsin* and *uv opsin*, which
are visual photoreceptor genes in Coleoptera, while some opsin genes are known to
have light-independent roles in *D. melanogaster* [31-35]. We checked whether the
blast-hit transcripts matched exon regions with mapped short reads of transcripts by
HISAT. Matched transcripts were used for subsequent comparative analyses with a
related surface species: *P. chalceus*, whose opsin amino acid sequences were already
registered in NCBI protein database [32].

Identification of *lw opsin* gene

A part of the genome sequence that has high similarity score to the Lw opsin amino acid sequence of *P. chalceus* was divided into three scaffolds (Table S2). To join these scaffolds together, primers were designed on each scaffold using Primer3Plus (https://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi) (Table S3) and PCR was performed using PrimeSTAR Max DNA Polymerase (Takara Bio, Shiga, Japan). The sequences of the PCR products, which were determined by Sanger sequencing, were overlapped on the scaffolds. The transcript sequence that showed similarity to the Lw opsin amino acid sequence of *P. chalceus* using tblastn was divided into two contigs (Table S2). These contigs were joined together by Sanger sequencing in the same method as above. Primers were designed on each contig (Table S3) and RT-PCR was performed using a 3′ RACE CORE Set (Takara Bio).

To specify exon and intron regions, the acquired transcript sequence of the *lw opsin* gene was aligned to the acquired genome sequence of the *lw opsin* gene with Exonerate v2.4.0 [36]. The exon and intron regions were illustrated with GenePalette [37].

Identification of *uv opsin* gene

There was one genome scaffold that had high similarity score to the Uv opsin amino acid sequence of *P. chalceus* using tblastn (Table S2), but no transcript contig was found. However, short read sequences originated from RNA were mapped to the scaffold of the *uv opsin* gene. We determined the transcript sequence of the *uv opsin* gene by Sanger sequencing. Firstly, the exon regions were predicted with Exonerate using the Uv opsin amino acid sequence of *P. chalceus*. Then primers were designed on the predicted exon regions with Primer3Plus (Table S3) and RT-PCR was performed using a 3′ RACE CORE Set.

To specify exon and intron regions, the acquired transcript sequence of the *uv opsin* gene was aligned to the acquired genome sequence of the *uv opsin* gene with Exonerate. The exon and intron regions were illustrated with GenePalette.

Opsin phylogeny and tests of selection

Blastp was performed using the acquired amino acid sequences of opsins in *T. kuznetsovi* as queries for non-redundant protein sequences in the NCBI database (Table S4). Amino acid sequences of opsins in *T. kuznetsovi*, four beetle species (*P. chalceus*, *Gyrinus marinus*, *Thermonectus marmoratus* and *T. castaneum*) and a honeybee (*Apis mellifera*) were aligned with MUSCLE in MEGA v 11 [38]. Based on the maximum

likelihood method, a phylogenetic tree of nucleotide sequences was reconstructed under
the best-fit GTR + G + I model with 1000 bootstrap generations.
The ancestral sequences of opsin gene sequences between subterranean *T.*
*kuznetsovi* and surface *P. chalceus* were estimated, based on above five beetle's
phylogenetic relationship using MEGA. Based on the maximum likelihood method [39],
the ratios of nonsynonymous (Ka) to synonymous (Ks) nucleotide substitution rates
were calculated between an ancestral sequence and a sequence of *T. kuznetsovi* and
between the ancestor sequence and a sequence of *P. chalceus* using KaKs_Calculator v
3.0 [40]. Fisher's exact test on a $2 \times 2$ contingency table was conducted using the
number of synonymous and nonsynonymous sites and synonymous and
nonsynonymous substitutions.
The Ka/Ks analysis is able to suggest that observed changes in a sequence have
been influenced by positive selection (Ka/Ks > 1), neutral evolution (Ka/Ks = 1), or
negative (purifying) selection (Ka/Ks < 1). In our study, the apparent result is expected
to be that opsin genes of *T. kuznetsovi* were under purifying selection, because along the
evolutionary branch from an ancestor to *T. kuznetsovi*, opsins will have been selected
under surface habitat before this lineage colonized subterranean habitat. To resolve this
problem, we also compared the degrees of purifying selection between opsin genes of *T.*
*kuznetsovi* (test) and *P. chalceus* (reference), carrying out branch-by-branch analyses
with RELAX in Hyphy [41]. As the result of this model, $k < 1$ is indicative of relaxed
selection, while $k > 1$ is indicative of purifying selection.
203
Histological study
The internal structure of a vestigial compound eye in *T. kuznetsovi* adults was
observed using paraffin sections. Adult heads were fixed in 50% alcohol Bouin solution
(ethanol:Bouin solution [for pathology, Fujifilm Wako Pure Chemicals] = 1:1) at room
temperature overnight or longer. The fixed samples were rinsed in 70% ethanol,
dehydrated in increasing concentrations of ethanol (90%, 95% and 100%) [42, 43], and
then cleared in xylene. Next, the samples were embedded in paraffin (Paraplast Plus;
Sigma Aldrich, MO, USA), and transverse sections (6 µm) were serially cut with a
microtome (OSK 97LF506; Ogawa Seiki, Tokyo, Japan). Sections were stained with
hematoxylin, observed with a microscope (CX-43; OLYMPUS, Tokyo, Japan) and
photographed with a mounted camera (EOS Kiss X9; Canon, Tokyo, Japan).
Because we could not obtain the complete series of cross sections due to their
friability, the nerve bundle between a compound eye and a brain in *T. kuznetsovi* adults
was observed with dissection. Adult heads were dissected in phosphate-buffered saline

and stained with 0.5 % methylene blue solution (22409-32; NAKALAI TESQUE,
Kyoto, Japan) for 1 hour. The samples were washed in phosphate-buffered saline,
observed with a stereo microscope (SZX16; OLYMPUS) and photographed with a
mounted camera (EOS Kiss X9; Canon).
222
223

**Results**

DNA and RNA sequencing

The genome size was estimated to be 554,652,206 bp based on the k-mer
frequency distribution of genome reads with KmerGenie. The assembled genome
contained 55,616 scaffolds with a total length of 456,726,283 bp (N50: 16,592 bp) and
96.20 % BUSCO completeness (Table 1). The assembled transcripts contained 71,303
contigs with a total length of 67,456,903 bp (N50: 1,883 bp) and 89.10 % BUSCO
completeness (Table 1). 81.90% of RNA reads were mapped to the assembled genome
with HISAT. The information of paired-end reads is summarized in Table S1.

Expression of photoreceptor genes and phototransduction genes

In the assembled genome, BLAST search found *lw opsin* gene, *uv opsin* gene
and non-visual *c-opsin* gene (Table 2). *lw opsin* and *c-opsin* were found in the
assembled transcripts, but *uv opsin* was not. Transcripts for 14 of the 15
BLAST-searched phototransduction genes were detected, namely, *Arr1*, *Arr2*, *Gα49B*,
*Gβ76C*, *Gγ30A*, *Gprk1*, *inaD, ninaA*, *ninaC*, *norpA*, *Pkc53E*, *Rab6*, *rdgC* and *trp*. Most
of these transcript sequences had neither a premature stop codon nor a frameshift
mutation, but transcript sequences of *Gα49B*, *Gβ76C*, *Gprk1*, *norpA* and *rdgC* had
premature stop codons or frameshift mutations in some transcript isoforms. These
transcripts might include primary transcripts before splicing. One phototransduction
gene, *trpl* was not detected in the transcripts, but was found in the genome.

Opsin genes

One *lw opsin* gene of *T. kuznetsovi* was identified with BLAST search. The
gene was divided into three genome scaffolds and two transcript contigs (Table S2). The
scaffolds and contigs were joined together using PCR and RT-PCR, and then exon and
intron regions of the *lw opsin* gene were determined (Fig. 2A). The conceptually
translated Lw opsin amino acid sequence was 379 residues and consisted of six exons.
There was neither a premature stop codon nor a frameshift mutation in the coding
sequence.

One *uv opsin* gene of *T. kuznetsovi* was identified by performing BLAST

search. The gene was present within one scaffold in the genome and no contig was

found in the transcripts (Table S2). The cDNA sequence of the *uv opsin* amino acid

sequence was determined using RT-PCR, and then exon and intron regions of the *uv*

*opsin* gene were determined (Fig. 2B). The conceptually translated Uv opsin amino acid

sequence was 373 residues and consisted of six exons. There was neither a premature

stop codon nor a frameshift mutation in the coding sequences.

Opsin phylogeny and selective pressure

A molecular phylogenetic tree of opsin genes of *T. kuznetsovi*, *P. chalceus*, *G.*

*marinus*, *T. marmoratus* and *T. castaneum* and *A. mellifera* was reconstructed (Fig. 3,

Table S4). The *lw opsin* and *uv opsin* of *T. kuznetsovi* were clustered with those of *P.*

*chalceus*, in accordance with their taxonomic relationship.

In *lw opsin* genes, Ka/Ks ratio was 0.158131 between the sequences of the

ancestor and *T. kuznetsovi* ($p = 1.12862e^{-013}$, Fisher's exact test), and Ka/Ks ratio was

0.0275072 between the sequences of the ancestor and *P. chalceus* ($p = 2.70467e^{-109}$,

Fisher's exact test) (Table 3). In *uv opsin*, Ka/Ks ratio was 0.138044 between the

sequences of the ancestor and *T. kuznetsovi* ($p = 1.34285e^{-024}$, Fisher's exact test), and

Ka/Ks ratio was 0.0695552 between the sequences of the ancestor and *P. chalceus* ($p =$

$7.58434e^{-092}$, Fisher's exact test). In all of these cases, Ka/Ks ratios were far below 1.0,

indicating that opsin genes have been under negative (purifying) selection in both the

lineage leading to *T. kuznetsovi* and that leading to *P. chalceus.*

Subsequently, the difference in the degree of the purifying selection between

the lineages from the ancestor to *T. kuznetsovi* and to *P. chalceus* was tested. As the

result of Relax analysis in Hyphy, *k* value was 0.61 ($p = 0.225$) in *lw opsin* and 2.14 ($p =$

0.539) in *uv opsin* (Table 3). Because this analysis did not show a statistical significance,

we were unable to conclude whether the selection on opsin genes was relaxed or

intensified in the lineage leading to *T. kuznetsovi* compared to the control lineage.

Internal structure of a compound eye

Putative photoreceptor cells stained by hematoxylin were observed in the

internal structure of a vestigial compound eye in a *T. kuznetsovi* adult (Fig. 4A). The

surface was covered by a transparent cuticle, a cornea. By observation from the outer

surface of the head, we could see a transparent cornea and ocular ridge with black

pigmentation (Fig. 1D). There was no pigmentation in cells within the eye structure,

unlike compound eyes of other carabid beetles [44]. No crystalline cones or any similar

structure were found [45]. An optic stalk, which is a nerve bundle connecting a
compound eye and a brain, was observed [46] (Fig. 4B).
292
293

**Discussion**
To understand the process of subterranean colonization of organisms, the
question of whether shallow subterranean habitats are a gateway to colonizing deep
zones has been featured in subterranean biology [3, 4]. In previous studies, organisms
inhabiting a cave or a calcrete aquifer and having extremely regressed structure and
function of vision were examined [8, 11-14]. To understand the transitional stage in
subterranean evolution, we focused on a trechine beetle, *Trechima kuznetsovi*, which
inhabits the upper hypogean zone and has a vestigial compound eye [16]. We evaluated
the ability of photoreception in *T. kuznetsovi* by genomics and histological observation.
We identified one *lw opsin* gene and one *uv opsin* gene in the genome and in
the transcripts in the adult. No frameshift mutation or premature stop codon was found
in these exon regions, Ka/Ks ratios were less than 1.0, and there was no significant
difference in the selective pressure between evolutionary lineages of subterranean *T.*
*kuznetsovi* and surface *P. chalceus*. These analyses implied that Lw opsin and Uv opsin
are under functional constraint, which could be one piece of evidence supporting the
possibility that *T. kuznetsovi* has photoreception ability for long wavelength and
ultraviolet light. Transcripts of 15 phototransduction genes without deleterious
mutations (premature stop codons or frameshift mutations) were detected in the
assembled transcripts. One phototransduction gene, *trpl,* was found in the assembled
genome. These results suggested the ability of phototransduction of *T. kuznetsovi*.
In subterranean diving beetles in Western Australia, the *lw opsin* gene became a
pseudogene due to frameshift mutations, and neither *lw opsin* nor *uv opsin* transcripts
were observed [12-14]. Frameshift mutations and premature stop codons occurred in
some phototransduction genes: *Arr1*, *Arr2*, *ninaC*, *trp* and *trpl* [14]. There are two
possible causes for these differences between *T. kuznetsovi* and the subterranean diving
beetles. The first possibility is the difference in their ecological niches. The calcrete
aquifer, in which subterranean diving beetles live, is located at a depth of 10–30 m
underground [11, 47]. In contrast, the upper hypogean zone, in which *T. kuznetsovi*
adults live, is a few or some dozen centimeters below the slope surface of a v-shaped
valley. *Trechiama kuznetsovi* adults would occasionally be exposed to the surface due to
landslides occurring as a result of precipitation or earthquakes [48-50]. This temporary
light stimulus could function as a selective pressure to retain the function of opsin genes

and other phototransduction genes. Friedrich et al. [29] also suggested that the retention
of photoreceptor genes and phototransduction genes in *Ptomaphagus hirtus* (Leiodidae)
inhabiting the Frozen Niagara cave entrance of Mammoth Cave in Kentucky, USA.
Beetles in shallow subterranean environments (the upper hypogean zone and cave
entrance) could tend to maintain the photoreception. The second possibility is that opsin
genes and other phototransduction genes of *T. kuznetsovi* have not spent enough time to
become pseudogenes. Niemiller et al. [51] showed that some cave lineages in
Amblyopsidae still possess functional rhodopsin, although they inhabit an aphotic
environment. This retained functionality is thought to be due to insufficient
accumulation of mutations during recent subterranean colonization. As in these cave
lineages, pseudogenization of opsin genes in *T. kuznetsovi* could not be observed
because divergence between *T. kuznetsovi* and related terrestrial species occurred
recently. To further examine this possibility, the divergence time of *Trechiama* species
needs to be studied.

By performing paraffin sectioning and dissection, we observed the cells inside
the compound eye and the optic stalk connecting the compound eye and the brain in *T.*
*kuznetsovi*. These observations suggested that photoreception is structurally possible
even with the vestigial compound eyes of this species. Complete loss of compound eyes
and optic lobes was observed in *Sinaphaenops wangorum* (Trechinae) inhabiting the
deep area of a cave in Guangxi Autonomous Region, China [52]. Thus, the existence of
the optic stalk is thought to be due to the retention of photoreception ability by *T.*
*kuznetsovi*, not to an inability of the visual system to degenerate any further.

Collectively, the results of genomics and histology performed here suggested
the ability of photoreception in *T. kuznetsovi*. This species is thought to possess both a
surface trait (photoreception) and some subterranean traits (vestigial compound eye,
underdeveloped body pigmentation and other morphological adaptations) [53, 54].
These characteristics would reflect an intermediate phase toward colonizing a deeper
subterranean niche. Further understanding of the visual degeneration process will be
obtained by clarifying the phylogenetic relationship between subterranean species and
surface species of trechine beetles.

**Conclusions**

By *de novo* assembly of genome and transcript sequences, we identified
photoreceptor genes and phototransduction genes of a trechine beetle, *Trechiama*
*kuznetsovi*, which inhabits the upper hypogean zone. The encoded amino acid sequences
of *lw opsin* and *uv opsin* had neither a premature stop codon nor a frameshift mutation,

and appeared to be subject to purifying selection. We identified potential photoreceptor
cells in the compound eye and nerve bundle connected to the brain. The present findings
suggest that *T. kuznetsovi* has retained the ability of photoreception.
365

**Abbreviations**
Lw opsin: Long wavelength opsin
Uv opsin: Ultraviolet opsin
Arr1: Arrestin 1
Arr2: Arrestin 2
Gα49B: G protein α-subunit 49B
Gβ76C: G protein β-subunit 76C
Gγ30A: G protein γ-subunit 30A
Gprk1: G protein-coupled receptor kinase 1
InaD: Inactivation no afterpotential D
NinaA: Neither inactivation nor afterpotential A
NinaC: Neither inactivation nor afterpotential C
NorpA: No receptor potential A
Pkc53E: Protein C kinase 53E
Rab6: Rab-protein 6
RdgC: Retinal degeneration C
Trp: Transient receptor potential
Trpl: Trp-like
CO: Cornea
R: Putative photoreceptor cells
OR: Ocular ridge
CE: Compound eye
NB: Nerve bundle
BR: Brain
390

**Declarations**
Ethical approval and consent to participate: Not applicable.
Consent for publication: Not applicable.
Availability of data and material: Sequence reads and scaffolds of the genome and
transcripts of *Trechiama kuznetsovi* were deposited to DDBJ/EMBL/GenBank
(DRR415324, DRR415325).
Competing interests: The authors declare no competing interests.

Funding: This study was supported by KAKENHI (22J12200), the Sasakawa Scientific Research Grant from The Japan Science Society, Asahi Glass Foundation, Kurita Water and Environment Foundation, and the research expense fund of a Hokkaido University Ambitious Doctoral Fellowship.

Authors contributions: Conceptualization, T.N. and S.K.; Data acquisition, T.N., Y.T., and H.A.; Data analysis, T.N., Y.T, H.A. and S.K.; Writing, T.N. and S.K.; All authors approved the final manuscript.

Acknowledgements: We thank Takashi Hayakawa and Juno Shimada for technical help; Cédric Finet for technical advice; Masahiro Ôhara for collection site information; Elizabeth Nakajima for English editing; and members of the Koshikawa lab for sample collection and discussion. The super-computing resource was provided by Human Genome Center (the Univ. of Tokyo).

Table 1

The summary statistics for assembly of the genome and transcripts in *T. kuznetsovi*.

|  | Genome | Transcripts |
|---|---|---|
| Total assembled length (bp) | 456,726,283 | 67,456,903 |
| Scaffolds (n) | 55,616 | 71,303 |
| The longest scaffold (bp) | 306,214 | 52,805 |
| The shortest scaffold (bp) | 386 | 181 |
| Scaffold N50 (bp) | 16,592 | 1,883 |
| BUSCO pipeline analysis (% complete / partial) | 96.20 / 98.83 | 89.10 / 95.32 |
| CEGMA pipeline analysis (% complete / partial) | 93.55 / 100 | Not Available |

Table 2

Photoreceptor and phototransduction genes detected in the assembled genome and transcripts.

| Query | | | *T. kuznetsovi* | |
|---|---|---|---|---|
| Protein name | Species | Accession no. | Genome | Transcripts |
| Visual opsin proteins | | | | |
| Lw opsin | Pcha | APY20654 | yes | yes (+) |
| Uv opsin | Pcha | APY20653 | yes | no |
| Non-visual opsin protein | | | | |
| C-opsin | Tcas | NP_001138950 | yes | yes (+) |
| Phototransduction proteins | | | | |
| Arr1 | Tcas | XP_966595 | yes | yes (+) |
| Arr2 | Tcas | NP_001164084 | yes | yes (+) |
| Gα49B | Tcas | XP_966311 | yes | yes (+/-) |
| Gβ76C | Tcas | XP_973851 | yes | yes (+/-) |
| Gγ30A | Tcas | XP_972123 | yes* | yes (+) |
| Gprk1 | Tcas | XP_966480 | yes | yes (+/-) |
| InaD | Tcas | XP_015837295 | yes | yes (+) |
| NinaA | Tcas | XP_973192 | yes | yes (+) |
| NinaC | Tcas | XP_015835531 | yes | yes (+) |
| NorpA | Tcas | XP_001812780 | yes | yes (+/-) |
| Pkc53E | Dmel | NP_476682 | yes | yes (+) |
| Rab6 | Tcas | XP_972453 | yes | yes (+) |
| RdgC | Tcas | XP_974915 | yes | yes (+/-) |
| Trp | Tcas | XP_968670 | yes | yes (+) |
| Trpl | Tcas | XP_968598 | yes | no |

'yes' indicates that the genes were detected at e-value $< 1 \times e^{-20}$. 'yes*' indicates that the

gene was detected at e-value $\geq 1 \times e^{-20}$. 'no' indicates that the genes were not detected in

BLAST-hit sequences. In BLAST-search for the transcripts, '+' and '-' indicate that
deleterious mutations (premature stop codons or frameshift mutations) were absent (+)
and present (-) in the sequences hit at. '+/-' indicates that we identified both transcripts
without and with deleterious mutations. Species name abbreviation: Pcha, *Pogonus*
*chalceus*; Tcas, *Tribolium castaneum*; Dmel, *Drosophila melanogaster*.
429
430
431     Table 3
432     The test of selection in opsin genes of *T. kuznetsovi*.
433

| | Test 1 | | Test 2 | |
| --- | --- | --- | --- | --- |
| Gene | Comparison | Ka/Ks ratio | Comparison | *k* value in Relax |
| *lw opsin* | Tkuz-Ancestor | 0.158131 | Tkuz-Pcha | 0.61 |
| | Pcha-Ancestor | 0.0275072 | | |
| *uv opsin* | Tkuz-Ancestor | 0.138044 | Tkuz-Pcha | 2.14 |
| | Pcha-Ancestor | 0.0695552 | | |

434
435
Legends
Fig. 1.
The habitat and morphology of *T. kuznetsovi*. (A) A v-shaped valley of a mountain area
at Yûbari City in Hokkaido. (B) The upper hypogean zone under the slope of a v-shaped
valley. It consists of small rocks and clay. (C) Dorsal view of *T. kuznetsovi*. Scale bar
indicates 3 mm. (D) Lateral view of *T. kuznetsovi* head. The bracket indicates the
vestigial compound eye. The black part is the pigmentation of the cuticle behind a
compound eye (ocular ridge). The arrowhead indicates a hole on the ocular ridge,
through which the nerve bundle is presumed to pass. There is no pigmentation in cells
inside the compound eye structure. Scale bar indicates 0.2 mm.
446
447     Fig. 2.
448     The structure of opsin genes in *T. kuznetsovi*. (A) The putative structure of *lw opsin*
449     gene, which consists of six coding exons. (B) The putative structure *uv opsin* gene,
450     which consists of six coding exons.

451
Fig. 3.
A maximum likelihood tree for visual opsin genes. Bootstrap probabilities are provided
on nodes. OTU names consist of the abbreviated species names, gene names and
accession numbers: Tkuz, *Trechiama kuznetsovi*; Pcha, *Pogonus chalceus*; Gmar,
*Gyrinus marinus*; Tmar, *Thermonectus marmoratus*; Tcas, *Tribolium castaneum*; Amel,
*Apis mellifera.*
458
Fig. 4.
The internal structure and nerve tissue of *T. kuznetsovi* head. (A) Transverse section of a
compound eye, dorsal is up. Scale bar indicates 20 μm. CO, cornea; R, putative
photoreceptor cells; OR, ocular ridge. (B) Dorsal view of dissected head. Nerve tissue
was stained with methylene blue solution. Scale bar indicates 0.5 mm. CE, compound
eye; NB, nerve bundle; BR, brain (supraesophageal ganglion).
465
466 Table S1.
467 The result of genome and transcript sequencing in *T. kuznetsovi*.
468
469 Table S2.
470 BLAST search for opsin genes to the assembled genome and transcripts in *T. kuznetsovi*.
471
472 Table S3.
473 Primers used in PCR and RT-PCR to amplify opsin genes in *T. kuznetsovi*. *T* (□),
474 annealing temperature.
475
476 Table S4.
477 BLAST search for opsin genes to non-redundant protein sequences in NCBI database.
478
479

480 References
481 1. McGaugh SE, Gross JB, Aken B, Blin M, Borowsky R, Chalopin D, et al. The
482 cavefish genome reveals candidate genes for eye loss. Nat Commun. 2014;5:1-10.
483 https://doi.org/10.1038/ncomms6307.
484
2. Mammola S, Cardoso P, Culver DC, Deharveng L, Ferreira RL, Fišer C, et al.
Scientists' warning on the conservation of subterranean ecosystems. BioScience.

2019;69:641-650. https://doi.org/10.1093/biosci/biz064.

3. Mammola S, Amorim IR, Bichuette ME, Borges PAV, Cheeptham N, Cooper SJB, et
al. Fundamental research questions in subterranean biology. Biol Rev.
2020;95:1855-1872. https://doi.org/10.1111/brv.12642.

4. Culver DC, Pipan T. Superficial subterranean habitats – gateway to the subterranean
realm. Cave Karst Sci. 2009;35:5-12.

5. Soares D, Niemiller ML. Sensory adaptations of fishes to subterranean environments.
BioScience. 2013;63:274-283. https://doi.org/10.1525/bio.2013.63.4.7.

6. Jeffery WR. Regressive evolution in *Astyanax* cavefish. Annu Rev Genet.
2009;43:25-47. https://doi.org/10.1146/annurev-genet-102108-134216.

7. Friedrich M. Biological clocks and visual systems in cave-adapted animals at the
dawn of speleogenomics. Integr Comp Biol. 2013;53:50-67.
https://doi.org/10.1093/icb/ict058.

8. Simon N, Fujita S, Porter M, Yoshizawa M. Expression of extraocular opsin genes
and light-dependent basal activity of blind cavefish. PeerJ. 2019;7:e8148.
https://doi.org/10.7717/peerj.8148.

9. Policarpo M, Fumey J, Lafargeas P, Naquin D, Thermes C, Naville M, et al.
Contrasting gene decay in subterranean vertebrates: insights from cavefishes and
fossorial mammals. Mol Biol Evol. 2021;38:589-605.
https://doi.org/10.1093/molbev/msaa249.

10. Yamamoto Y, Jeffery WR. Central role for the lens in cave fish eye degeneration.
Science. 2000;289:631-633. https://doi.org/10.1126/science.289.5479.631.

11. Tierney SM, Langille B, Humphreys WF, Austin AD, Cooper SJB. Massive parallel
regression: a précis of genetic mechanisms for vision loss in diving beetles. Integr
Comp Biol. 2018;58:465-479. https://doi.org/10.1093/icb/icy035.

12. Tierney SM, Cooper SJB, Saint KM, Bertozzi T, Hyde J, Humphreys WF, et al.

Opsin transcripts of predatory diving beetles: a comparison of surface and subterranean photic niches. R Soc Open Sci. 2015;2:140386. https://doi.org/10.1098/rsos.140386.

13. Langille BL, Hyde J, Saint KM, Bradford TM, Stringer DN, Tierney SM, et al. Evidence for speciation underground in diving beetles (Dytiscidae) from a subterranean archipelago. Evolution. 2021;75:166-175. https://doi.org/10.1111/evo.14135.

14. Langille BL, Tierney SM, Bertozzi T, Beasley-Hall PG, Bradford TM, Fagan-Jeffries EP, et al. Parallel decay of vision genes in subterranean water beetles. Mol Phylogenet Evol. 2022;173:107522. https://doi.org/10.1016/j.ympev.2022.107522.

15. Ortuño VM, Cuesta E, Gilgado JD, Ledesma E. A new hypogean *Trechus* Clairville (Coleoptera, Carabidae, Trechini) discovered in a non-calcareous superficial subterranean habitat of the Iberian system (Central Spain). Zootaxa. 2014;3802:359-372. https://doi.org/10.11646/zootaxa.3802.3.5.

16. Uéno SI, Lafer GS. A new anophthalmic *Trechiama* (Coleoptera, Trechinae) from Central Hokkaido, Northeast Japan. Elytra. 1992;20:137-143.

17. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018;34:i884-i890. https://doi.org/10.1093/bioinformatics/bty560.

18. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 2014;30:2114-2120. https://doi.org/10.1093/bioinformatics/btu170.

19. Chikhi R, Medvedev P. Informed and automated *k*-mer size selection for genome assembly. Bioinformatics. 2014;30:31-37. https://doi.org/10.1093/bioinformatics/btt310.

20. Kajitani R, Toshimoto K, Noguchi H, Toyoda A, Ogura Y, Okuno M, et al. Efficient de novo assembly of highly heterozygous genomes from whole-genome shotgun short reads. Genome Res. 2014;24:1384-1395. https://doi.org/10.1101/gr.170720.113.

21. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol. 2011;29:644-652. https://doi.org/10.1038/nbt.1883.

22. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. *De novo* transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. Nat Protoc. 2013;8:1494-1512. https://doi.org/10.1038/nprot.2013.084.

23. Van Belleghem SM, Vangestel C, De Wolf K, De Corte Z, Möst M, Rastas P, et al. Evolution at two time frames: polymorphisms from an ancient singular divergence event fuel contemporary parallel evolution. PLoS Genet. 2018;14:e1007796. https://doi.org/10.1371/journal.pgen.1007796.

24. Nishimura O, Hara Y, Kuraku S. gVolante for standardizing completeness assessment of genome and transcriptome assemblies. Bioinformatics. 2017;33:3635-3637. https://doi.org/10.1093/bioinformatics/btx445.

25. Shen W, Le S, Li Y, Hu F. SeqKit: a cross-platform and ultrafast toolkit for FASTA/Q file manipulation. PloS One. 2016;11:e0163962. https://doi.org/10.1371/journal.pone.0163962.

26. Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements. Nat Methods. 2015;12:357-360. https://doi.org/10.1038/nmeth.3317.

27. Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. Brief Bioinform. 2013;14:178-192. https://doi.org/10.1093/bib/bbs017.

28. Bao R, Friedrich M. Molecular evolution of the *Drosophila* retinome: exceptional gene gain in the higher Diptera. Mol Biol Evol. 2009;26:1273–1287. https://doi.org/10.1093/molbev/msp039.

29. Friedrich M, Chen R, Daines B, Bao R, Caravas J, Rai PK, et al. Phototransduction and clock gene expression in the troglobiont beetle *Ptomaphagus hirtus* of Mammoth cave. J Exp Biol. 2011;214:3532-3541. https://doi.org/10.1242/jeb.060368.

30. Tsuji T, Gotoh H, Morita S, Hirata J, Minakuchi Y, Yaginuma T, et al. Molecular characterization of eye pigmentation-related ABC transporter genes in the ladybird

beetle *Harmonia axyridis* reveals striking gene duplication of the white gene. Zool Sci.
2018;35:260-267. https://doi.org/10.2108/zs170166.

31. Futahashi R, Kawahara-Miki R, Kinoshita M, Yoshitake K, Yajima S, Arikawa K, et
al. Extraordinary diversity of visual opsin genes in dragonflies. Proc Natl Acad Sci U S
A. 2015;112:E1247-E1256. https://doi.org/10.1073/pnas.1424670112.

32. Sharkey CR, Fujimoto MS, Lord NP, Shin S, McKenna DD, Suvorov A, et al.
Overcoming the loss of blue sensitivity through opsin duplication in the largest animal
group, beetles. Sci Rep. 2017;7:8. https://doi.org/10.1038/s41598-017-00061-7.

33. Guignard Q, Allison JD, Slippers B. The evolution of insect visual opsin genes with
specific consideration of the influence of ocelli and life history traits. BMC Ecol Evol.
2022;22:2. https://doi.org/10.1186/s12862-022-01960-8.

34. Leung NY, Montell C. Unconventional roles of opsins. Annu Rev Cell Dev Biol.
2017;33:241-264. https://doi.org/10.1146/annurev-cellbio-100616-060432.

35. Leung NY, Thakur DP, Gurav AS, Kim SH, Pizio AD, Niv MY, et al. Functions of
opsins in *Drosophila* taste. Curr Biol. 2020;30:1367-1379.e6.
https://doi.org/10.1016/j.cub.2020.01.068.

36. Slater GSC, Birney E. Automated generation of heuristics for biological sequence
comparison. BMC Bioinf. 2005;6:1-11. https://doi.org/10.1186/1471-2105-6-31.

37. Smith AF, Posakony JW, Rebeiz M. Automated tools for comparative sequence
analysis of genic regions using the GenePalette application. Dev Biol.
2017;429:158-164. https://doi.org/10.1016/j.ydbio.2017.06.033.

38. Tamura K, Stecher G, Kumar S. MEGA11: molecular evolutionary genetics analysis
version 11. Mol Biol Evol. 2021;38:3022-3027.
https://doi.org/10.1093/molbev/msab120.

39. Goldman N, Yang Z. A codon-based model of nucleotide substitution for
protein-coding DNA sequences. Mol Biol Evol. 1994;11:725-736.
https://doi.org/10.1093/oxfordjournals.molbev.a040153.

40. Zhang Z. KaKs_Calculator 3.0: calculating selective pressure on coding and
non-coding sequences. Genom Proteom Bioinform. 2022;20:536-540.
https://doi.org/10.1016/j.gpb.2021.12.002.

41. Wertheim JO, Murrell B, Smith MD, Kosakovsky Pond SL, Scheffler K. RELAX:
detecting relaxed selection in a phylogenetic framework. Mol Biol Evol.
2015;32:820-832. https://doi.org/10.1093/molbev/msu400.

42. Cornette R, Matsumoto T, Miura T. Histological analysis of fat body development
and molting events during soldier differentiation in the damp-wood termite,
*Hodotermopsis sjostedti* (Isoptera, Termopsidae). Zool Sci. 2007;24:1066-1074.
https://doi.org/10.2108/zsj.24.1066.

43. Oguchi K, Miura T. Unique morphogenesis in the damp-wood termite: abscission of
the stylus during female reproductive caste differentiation. Zool Sci. 2019;36:380-386.
https://doi.org/10.2108/zs190056.

44. Home EM. The fine structure of some carabid beetle eyes, with particular reference
to ciliary structures in the retinula cells. Tissue Cell. 1976;8: 311-333.
https://doi.org/10.1016/0040-8166(76)90055-0.

45. van der Kooi CJ, Stavenga DG, Arikawa K, Belušič G, Kelber A. Evolution of
insect color vision: from spectral sensitivity to visual ecology. Annu Rev Entomol.
2021;66:435-461. https://doi.org/10.1146/annurev-ento-061720-071644.

46. Whited JL, Cassell A, Brouillette M, Garrity PA. Dynactin is required to maintain
nuclear position within postmitotic *Drosophila* photoreceptor neurons. Development.
2004;131:4677-4686. https://doi.org/10.1242/dev.01366.

47. Langille BL, Tierney SM, Austin AD, Humphreys WF, Cooper SJB. How blind are
they? Phototactic responses in stygobiont diving beetles (Coleoptera: Dytiscidae) from
calcrete aquifers of Western Australia. Austral Entomol. 2019;58:425-431.
https://doi.org/10.1111/aen.12330.

48. Dhakal A, Tsuchiya S, Ohsaka O. Long term change in landslide area in Ikawa Lake

catchment using aerial photographs and GIS. J Erosion Contr. 2007;59:25-31.
https://doi.org/10.11475/sabo1973.59.6_25.
669

670 49. Tsuchiya S, Imaizumi F. Large sediment movement caused by the catastrophic
671 Ohya-kuzure landslide. J Disaster Res. 2010;5:257-263.
672 https://doi.org/10.20965/jdr.2010.p0257.
673

674 50. Matsuoka N, Imaizumi F, Nishii R. Geomorphic dynamics and sediment budget in
675 the southern Japanese Alps: Recent studies and prospects. J Geogr. 2013;122:591-614.
676 https://doi.org/10.5026/jgeography.122.591.
677

678 51. Niemiller ML, Fitzpatrick BM, Shah P, Schmitz L, Near TJ. Evidence for repeated
679 loss of selective constraint in rhodopsin of amblyopsid cavefishes (Teleostei:
680 Amblyopsidae). Evolution. 2013;67:732-748.
681 https://doi.org/10.1111/j.1558-5646.2012.01822.x.
682

683 52. Luo XZ, Wipfler B, Ribera I, Liang HB, Tian MY, Ge SQ, et al. The cephalic
684 morphology of free-living and cave-dwelling species of trechine ground beetles from
685 China (Coleoptera, Carabidae). Org Divers Evol. 2018;18:125-142.
686 https://doi.org/10.1007/s13127-017-0351-5.
687

688 53. Protas M, Jeffery WR. Evolution and development in cave animals: from fish to
689 crustaceans. Wiley Interdiscip Rev Dev Biol. 2012;1:823-845.
690 https://doi.org/10.1002/wdev.61.
691

692 54. Bilandžija H, Laslo M, Porter ML, Fong DW. Melanization in response to wounding
693 is ancestral in arthropods and conserved in albino cave species. Sci Rep. 2017;7:1-11.
694 https://doi.org/10.1038/s41598-017-17471-2.

Fig. 1

A

5'                                                                          3'

1000 bp

B

5'                                                                          3'

1000 bp

Fig. 2

Fig. 3

Fig. 4