Mice with renal-specific alterations of stem cell-associated

signaling develop symptoms of chronic kidney disease but

surprisingly no tumors

Stem cell-associated signaling in the kidney

Adam Myszczyszyn[1,*], Oliver Popp[2], Severine Kunz[3], Anje Sporbert[4], Simone Jung[1],

Louis C. Penning[5], Annika Fendler[1], Philipp Mertins[2,6], Walter Birchmeier[1,6]

1 Cancer Research Program, Max Delbrueck Center for Molecular Medicine (MDC) in the

Helmholtz Association, Robert-Roessle-Strasse 10, 13125 Berlin, Germany

2 Proteomics, Max Delbrueck Center for Molecular Medicine (MDC) in the Helmholtz

Association, Robert-Roessle-Strasse 10, 13125 Berlin, Germany

3 Electron Microscopy, Max Delbrueck Center for Molecular Medicine (MDC) in the

Helmholtz Association, Robert-Roessle-Strasse 10, 13125 Berlin, Germany

4 Advanced Light Microscopy, Max Delbrueck Center for Molecular Medicine (MDC) in the

Helmholtz Association, Robert-Roessle-Strasse 10, 13125 Berlin, Germany

5 Faculty of Veterinary Medicine, Utrecht University, Yalelaan 1, 3584 CL Utrecht, The

Netherlands

6 Senior author

* Corresponding author: adam.myszczyszyn@gmail.com

Abstract

Previously, we found that Wnt and Notch signaling govern stem cells of clear cell kidney cancer (ccRCC) in patients. To mimic stem cell responses in the normal kidney in vitro in a marker-unbiased fashion, we have established organoids from total single adult mouse kidney epithelial cells in Matrigel and serum-free conditions. Deep proteomic and phosphoproteomic analyses revealed that the organoids resembled renewal of adult kidney tubular epithelia, since organoid cells displayed activity of Wnt and Notch signaling, long-term proliferation and expression of markers of proximal and distal nephron lineages. In our wish to model stem cell-derived human ccRCC, we have generated two types of genetic double kidney mutants in mice: Wnt-β-catenin-GOF together with Notch-GOF and Wnt-β-catenin-GOF together with a most common alteration in ccRCC, Vhl-LOF. An inducible Pax8-rtTA-LC1-Cre was used to drive recombination specifically in adult kidney epithelial cells. We confirmed mutagenesis of β-catenin, Notch and Vhl alleles on DNA, protein and mRNA target gene levels. Surprisingly, we observed symptoms of chronic kidney disease (CKD) in mutant mice, but no increased proliferation and tumorigenesis. Thus, the responses of kidney stem cells in the organoid and genetic systems produced different phenotypes, i.e. enhanced renewal versus CKD.

Introduction

Over the past few decades, Wnt-β-catenin and Notch have emerged as fundamental signal transduction systems, which regulate maintenance and differentiation of stem cells in homeostasis and regeneration of adult organs (1,2). In the mouse kidney, Wnt and Notch signaling are upregulated in Sox9-positive progenitors that contribute to regeneration of proximal tubule, Loop of Henle and distal tubule (3). Unipotent progenitors specific for proximal, distal and collecting duct segments were reported in *Rainbow* mice, which drive homeostatic maintenance and regeneration of the kidney and are Wnt-responsive (4). Moreover, deletion of the Wnt mediator β-catenin in kidney tubular cells and blocking Notch with a γ-secretase inhibitor delayed kidney recovery upon injury in mice (5,6). Together, activity of Wnt and Notch signaling seems to be a common feature of different kidney stem/progenitor cell populations.

Organoids are emerging as powerful tools to study properties of stem/progenitor cells in adult epithelial tissues. Organoids are 3D structures grown from stem/progenitor cells in culture with Matrigel and well-defined growth factors. During organoid formation, cells self-renew, differentiate and self-organize by cell sorting and spatially restricted lineage commitment in a manner similar to natural organs. Thus, organoids recapitulate stem/progenitor cell behaviors during homeostatic maintenance and regeneration of adult epithelia (7,8). Adult kidney organoids were created from normal human and mouse kidneys (9,10). However, characterization of stemness in these organoids has not been pushed forward to a satisfying level.

Clear cell renal cell carcinoma (ccRCC) is the most frequent and aggressive human kidney cancer (11). Inactivation of the Von Hippel-Lindau (*VHL*) tumor suppressor gene followed by a HIF-1α- and HIF-2α-mediated hypoxic state occurs in the majority of ccRCC cases (12). However, deletion of Vhl in mouse kidneys is insufficient for tumorigenesis (13). Our

laboratory identified CXCR4-, MET- and CD44-enriched cancer stem cells (CSCs) in human

ccRCCs (14). The CSCs displayed upregulation of Wnt and Notch signaling and

quantitatively correlated with advanced stages, aggressiveness and metastasis of tumors.

Treatment with small-molecule inhibitors of Wnt and Notch blocked self-renewal of the CSCs

in patient-derived models, i.e. in subcutaneous and orthotopic xenografts, and in organoids

and non-adherent spheres in cultures. This points to a crucial role of Wnt-high and Notch-

high stem/progenitor cells as actors in kidney tumorigenesis.

Upregulation of Wnt signaling in combination with genetic interferences at p53 or p21 in

mouse kidneys using embryonic Cres was sufficient to overcome Wnt-induced p53-p21-

mediated senescence and resulted in development of malignant lesions that displayed

characteristics of non-ccRCCs (15,16). Concomitant upregulation of Wnt and K-ras signaling

in embryonic mouse kidneys also drove formation of malignant lesions that resembled non-

ccRCCs, including pediatric tumors (17,18). Hyperactivation of Notch signaling in

combination with Vhl loss in adult mouse kidneys contributed to development of premalignant

lesions with some hallmarks of ccRCC (19,20). However, upregulation of either Wnt or Notch

in adult Cd133-positive kidney cells in mice failed to produce tumors (21). Answer to a

fundamental question remains elusive on whether the functional interplay of Wnt and Notch

drives tumorigenesis in the kidney, also in the context of Vhl inactivation.

To reveal the importance of Wnt and Notch signaling for adult renal stem/progenitor cells in

vitro without enriching populations positive for a specific marker, we have established tubular

organoids from whole mouse kidneys in Matrigel in serum-free conditions. Then, to mimic

development of human ccRCC from Wnt- and Notch-dependent stem cells, we have

generated genetic mutants in mice, i.e. Wnt-β-catenin-GOF together with Notch-GOF or with

Vhl-LOF, using an adult kidney tubular epithelium-specific Cre to circumvent embryonic or

postnatal kidney defects and non-renal confounding influences.

Materials and Methods

Mice

The Landesamt für Gesundheit und Soziales (LaGeSo) in Berlin approved the mouse study (G0342/13). Mice were housed and bred in the IVC cages type II at 21°C ambient temperature, at 50-60% humidity, in a 12-h dark/light cycle and in pathogen-free conditions. LC1-Cre mice (22) were a kind gift from Klaus Rajewsky (MDC). β-catenin-loxP (23), Notch-loxP (24) and LacZ-loxP (25) mice came from in-house colonies. Pax8-rtTA B6.Cg-Tg(Pax8-rtTA2S*M2)1Koes/J (26) and Vhl-loxP B6.129S4(C)-Vhltm1Jae/J (27) mice were purchased from the Jackson Laboratory. The studies in the organoids and genetic mouse models were performed using both male and female mice of a mixed C57BL/6J and FVB/NJ background, unless otherwise stated. To establish the genetic mouse models, adult one-month-old Pax8-rtTA-LC1-Cre-loxP mice and controls carrying loxP alleles, but no Pax8-rtTA and/or LC1-Cre (both males and females) were exposed for 5 days to doxycycline (Sigma-Aldrich) at the concentration of 0.2 mg/ml in drinking water supplemented with 5% sucrose. According to a score sheet, mice were checked at least twice per week until illness symptoms were observed. Sick mice were immediately sacrificed and biological material was collected.

Preparation of kidney cell suspensions

Kidneys were harvested and transferred to MEM with Earle's Salts (with 2.2 g/l NaHCO3, without L-Glutamine) medium (Biochrom) supplemented with 10% FCS (Life Technologies), 1x Non-Essential Amino Acids (NEAA, Thermo Fisher Scientific), 2 mM L-Glutamine (Biochrom), 1x Penicillin-Streptomycin (Life Technologies) and 125 µg/ml Amphotericin B (Biomol). Whole kidneys were minced into pieces smaller than 1 mm3, digested in 2 mg/ml Collagenase P (Roche) for 45 min and filtered through sieves with 70 and 40 µm pore size. Erythrocytes were lysed in RBC lysis buffer and lymphocytes were depleted by MACS using CD45 microbeads (Miltenyi Biotec) according to the manufacturer's instructions.

Kidney organoid cultures

Freshly isolated cell suspensions were seeded at 10^5 cells per well in 25 µl Matrigel (75%, Corning) lenses in 48-well plates. Matrigel drops were overlayed with 250 µl organoid medium (Table 1). Medium was changed every other day. The organoids were passaged for the first time after 14 days and subsequently once per week. Matrigel lenses were mechanically broken and the organoids were collected and dissociated to small cell clusters using TrypLE Select (Life Technologies) for 10 min. In the first two passages, collected organoids were additionally filtered through a sieve with 40 µm pore size. Cultures were passaged in 1:3 ratios. If single cell suspensions were needed, the organoids were dissociated in TrypLE for 1 hr. Cell clusters from the organoids were cryopreserved at an early passage in 1 ml Recovery Cell Culture Freezing Medium (Thermo Fisher Scientific) according to the manufacturer's instructions. BrdU (Thermo Fisher Scientific) treatment (at 10 µM) was performed for 4 hrs prior to organoid collection for staining. For functional assays, 100 ng/ml human Wnt3a (R&D Systems) was added. For the examination of structural changes of the organoids upon R-spondin-1 withdrawal or ICG-001 treatment, single cells were seeded at 45,000 cells in 48-well plates and cultured 7 days in organoid medium without R-spondin-1 or cultured 24h in normal medium and 6 days in medium containing ICG-001 (Biochem-partner) at IC50. Brightfield and fluorescence images were acquired with a Leica DMI 6000 B microscope using the Leica LAS X software.

Table 1. **Medium components for organoid culture.**

Component	Function	Concentration	Supplier
Advanced DMEM/F12 with GlutaMax	Basal medium	–	Life Technologies
Penicillin-Streptomycin	Antibiotics	1x	Life Technologies
Amphotericin B	Antifungal	125 µg/ml	Biomol

HEPES	Buffer	10 mM	Life Technologies
N-acetylcysteine	Antioxidant	1.25 mM	Sigma-Aldrich
B27	Serum-free growth supplement	1x	Life Technologies
N2	Serum-free growth supplement	1x	Life Technologies
Nicotinamide	Increasing organoid formation and lifespan via inhibiting sirtuin activity involved in apoptosis and differentiation (28,29)	10 mM	Sigma-Aldrich
Human EGF	Stimulating proliferation, inhibiting apoptosis (28,29)	50 ng/ml	Peprotech
A83-01	Blocking differentiation and promoting long-term culture of stem cells via inhibiting TGFβ signaling (28,29)	500 nM	Sigma-Aldrich
Human R-spondin-1	Maintaining stem cells via agonizing Wnt signaling (28,29)	500 ng/ml	Peprotech
Hydrocortisone (HC)	Acting anti-inflammatory via repressing activity of NF-κB and AP-1 (30), inducing Yap signaling (31)	0.5 µg/ml	Sigma-Aldrich
Prostaglandin E2 (PGE2)	Preventing anoikis and promoting stem cell survival via enhancing Wnt signaling (28)	1 µM	Sigma-Aldrich

Proteomics and phosphoproteomics

Three independent replicates of early (OEP) and long-term (OLP) passage organoids were

released from Matrigel by mechanical disruption, collected in a conical tube and incubated

with the non-enzymatic Cell Recovery Solution (Corning) according to the manufacturer's

instructions to remove Matrigel rests. Three independent replicates of Epcam-positive kidney

epithelial cells (control kidney, CK) were MAC-sorted from freshly isolated single cell

suspensions from whole mouse kidneys using microbeads (Miltenyi Biotec) according to the

manufacturer's instructions. Global proteomic and phosphoproteomic characterization of the

samples was achieved by the tandem mass tag (TMT)-based quantitation using the TMT 10-

plex reagents (Thermo Fisher Scien-tific), as previously described (32). Briefly, samples were

lysed in urea lysis buffer containing protease and phosphatase inhibitors, subjected to tryptic

in-solution digest and labeled with the TMT 10-plex reagents. After pooling, samples were

separated into 24 fractions for the proteomic and 12 fractions for the phosphoproteomic

analysis using basic reversed-phase HPLC. Phosphopeptides were enriched from each

fraction using robot-assisted iron-based IMAC on an AssayMap Bravo system (Agilent).

Proteomic and phosphoproteomic samples were measured on a Q Exactive HF-X orbitrap

mass spectrometer (Thermo Fisher Scientific) connected to an EASY-nLC system (Thermo

Fisher Scientific) applying a 110-min online HPLC gradient. MS acquisition was performed at

a resolution of 60,000 in a scan range from 350 to 1500 Th. Data-dependent MS2 scans

were carried out at a resolution of 45,000 with an isolation window of 0.7 Th and a maximum

injection time of 86 ms using the top 20 peaks from each precursor scan for HCD

fragmentation. For the data analysis, the MaxQuant software package version 1.6.10.43 (33)

was used. A FDR cutoff of 0.01 was applied for peptides and proteins and database search

was performed using the mouse Uniprot database (July 2018), including isoforms. Variable

modifications included phosphorylation on serine, threonine and tyrosine, methionine-

oxidation, acetylated N-termini and deamidation of asparagine and glutamine, while the

TMT10-plex reporter ion quantitation was turned on using a PIF setting of 0.5. Log2-

transformed and median-MAD centered corrected reporter ion intensities were used for

quantitation. Protein groups were filtered for proteins, which were identified by at least one

unique peptide and at least two peptides in total and had valid values across all samples.

Phosphorylation site tables were filtered for valid values across all samples. For significance

calling, a two-sample moderated t-testing (limma R package) (34) was applied. A multiple

comparison correction was done using the Benjamini-Hochberg method (adjusted P-values).

The KEGG PATHWAY (mmu04310) and a previously published list (35) served as a

combined reference protein list for the Wnt signaling heatmap. The KEGG PATHWAY

(mmu04330) served as a reference protein list for the Notch signaling heatmap. A previously

published list (36) served as a reference protein list for the Yap signaling heatmap.

Section staining

The organoids were released from Matrigel by mechanical disruption, collected in a conical

tube and fixed in 10% (v/v) Neutral-buffered Formalin overnight. The organoids were

transferred to a 2-ml tube and overlayed with 1.5% agarose. The resulting agarose beads

were dehydrated and embedded in paraffin. All stainings were performed on 5-μm sections.

For haematoxylin-eosin (H&E) staining, sections were rehydrated and stained with

haematoxylin (Fluka) for 4 min and with eosin (Merck) for 2 min. For immunofluorescence,

sections were rehydrated, and antigen retrieval in TRIS-EDTA and blocking with 10% donkey

serum (Bio-Rad) were performed. Sections were incubated overnight with rat anti-BrdU

(1:100, Abcam, ab6326), mouse anti-E-cad (1:300, BD Biosciences, 610181), rabbit anti-

Laminin (1:500, Abcam, ab11575), rat anti-Zo-1 (1:200, Santa Cruz, sc-33725), mouse anti-

MDR1 (1:50, Santa Cruz, sc-55510) and mouse anti-AQP3 (1:50, Santa Cruz, sc-518001)

primary antibodies or with Biotinylated LTL (1:300, Vector Laboratories, B-1325) and

Rhodamine-conjugated PNA (1:300, Vector Laboratories, RL-1072) lectins, followed by

incubation with donkey fluorescent dye-conjugated Alexa488 anti-mouse, Alexa488 anti-

rabbit, Alexa488 anti-rat and Cy3 anti-rat (1:250, Jackson) secondary antibodies or with

Alexa647-conjugated Streptavidin (1:200, Thermo Fisher Scientific) and by counterstaining

with DAPI (0.2 μg/ml, Thermo Fisher Scientific) for 1 hr. For immunohistochemistry, sections

were rehydrated, antigen retrieval in TRIS-EDTA was performed, endogenous peroxidase

activity was blocked in 3.5% (v/v) hydrogen peroxide and blocking with 10% donkey serum

(Bio-Rad) was performed. Sections were incubated overnight with rabbit anti-Pax8 primary

antibody (1:2000, Proteintech, 10336-1-AP), followed by incubation with peroxidase-

conjugated donkey anti-rabbit secondary antibody from a kit (Dako) for 1 hr. Sections were

developed using the substrate-chromogen system (Dako) and counterstained with

haematoxylin (Fluka) for 2 min. Kidneys and spleens were fixed in 10% (v/v) Neutral-buffered

Formalin overnight, dehydrated and embedded in paraffin. All stainings were performed on 5

µm sections. For immunohistochemistry, sections were rehydrated, antigen retrieval in TRIS-

EDTA was performed, endogenous peroxidase activity was blocked in 3.5% (v/v) hydrogen

peroxide and blocking with 10% donkey serum (Bio-Rad) was performed. Sections were

incubated overnight with rabbit anti-N1icd (1:100, Cell Signaling, 4147S), rabbit anti-Ki67

(1:200, Thermo Fisher Scientific, RM-9106-S), mouse anti-phospho-γH2AX (1:500, Merck

Millipore, 05-636), mouse anti-p53 (1:100, Santa Cruz, sc-126) and mouse anti-p21 (1:100,

Santa Cruz, sc-6246) primary antibodies, followed by incubation with peroxidase-conjugated

donkey anti-rabbit and anti-mouse secondary antibodies from a kit (Dako) for 1 hr. Sections

were developed using the substrate-chromogen system (Dako) and counterstained with

haematoxylin (Fluka) for 2 min. For PAS staining, sections were rehydrated, stained using

the Periodic Acid-Schiff (PAS) Kit (Sigma-Aldrich) according to the manufacturer's

instructions and counterstained with haematoxylin (Fluka) for 2 min. For LacZ staining,

cryosections were prepared. Kidneys were placed in 30% sucrose in PBS overnight and

embedded in OCT compound (Sakura Finetek). 10-µm sections were fixed in 0.2% (v/v)

glutaraldehyde for 10 min, washed three times in washing solution (PBS, 2 mM MgCl2,

0.01% sodium deoxycholate and 0.02% NP-40), incubated overnight in staining solution

(PBS, 2 mM MgCl2, 0.01% sodium deoxycholate, 0.02% NP-40, 5 mM K3Fe(CN)6, 5 mM

K4Fe(CN)6 and 1 mg/ml X-gal (Roth)) and counterstained with nuclear fast red (Merck) for

5 min. Images were taken with an Axio Imager.Z1 microscope (for immunofluorescence) or

an Axio Scope.A1 microscope (for H&E staining, PAS staining, LacZ staining and

immunohistochemistry) using the Zen software (all Zeiss). Selected immunofluorescence

images were processed with Fiji ImageJ.

Whole mount staining

The organoids were released from Matrigel by mechanical disruption, collected, transferred

to a 2-ml tube, fixed in 10% (v/v) Neutral-buffered Formalin for 1.5 hrs and blocked in 10%

donkey serum (Bio-Rad) for 2 hrs. The organoids were incubated overnight with mouse anti-

E-cad primary antibody (1:300, BD Biosciences, 610181), Biotinylated LTL lectin (1:300,

Vector Laboratories, B-1325) and Rhodamine-conjugated PNA lectin (1:300, Vector

Laboratories, RL-1072), and then overnight with donkey Alexa488 anti-mouse secondary

antibody (1:250, Jackson), Alexa647-conjugated Streptavidin (1:200, Thermo Fisher

Scientific) and DAPI (0.2 µg/ml, Thermo Fisher Scientific). The organoids were taken up in

1.5% low-melting agarose and transferred to chamber slides. Confocal z-stacks were

acquired with a LSM700 or LSM710 microscope (Zeiss) using a long working distance C-

Achroplan 32x/0.85 water immersion objective and Immersol W (Zeiss). DAPI, Alexa488,

Cy3 and Alexa647 fluorophores were excited with a 405 nm, 488 nm, 561 nm and 633 nm

laser, respectively, and detected by sequential scanning using BP filters 410-460 nm, 500-

540 nm, 565-620 nm and 640-720 nm, respectively. Images were acquired with a pixel size

of 0.116 or 0.142 µm (lateral) and 1.11 or 1.16 µm (axial) with 12-bit and line average 2. If

necessary, detector gain and laser power were slightly adjusted with z-depth to keep a

constant signal. Restoration of confocal z-stacks was performed with the Huygens

Deconvolution software (SVI) running on a dedicated data processing workstation (Acquifer)

using a CMLE algorithm with 60 iterations and a SNR of 12. After deconvolution, confocal z-

stacks were 3D reconstructed using the Imaris 9.1 software (Bitplane/Oxford Instruments)

and either the volume reconstruction and clipping planes or orthogonal sections were used to

visualize and explore the 3D structure of the organoids.

Transmission electron microscopy

The organoids were released from Matrigel by mechanical disruption, collected in a conical

tube and fixed in 2% (w/v) formaldehyde and 2.5% (v/v) glutaraldehyde in 0.1 M phosphate

buffer for 2 hrs. After embedding in 10% agarose in a 2-ml tube, samples were post-fixed

with 1% (v/v) osmium tetroxide (Sigma-Aldrich), dehydrated in a graded series of EtOH and

embedded in PolyBed® 812 resin (Polysciences). Ultrathin sections (60-80 nm) were stained

with uranyl acetate (Polysciences) and lead citrate (Sigma-Aldrich) and examined at 80 kV

with an EM 910 electron microscope (Zeiss). Acquisition was performed with a Quemesa

CCD camera and the iTEM software (Emsis).

TUNEL assay

Kidneys were fixed in 10% (v/v) Neutral-buffered Formalin overnight, dehydrated and

embedded in paraffin. 5-µm sections were rehydrated, stained using the ApopTag Plus

Peroxidase In Situ Apoptosis Detection Kit (Merck Millipore) according to the manufacturer's

instructions and counterstained with haematoxylin (Fluka) for 2 min. Images were taken with

an Axio Scope.A1 microscope using the Zen software (Zeiss).

Immunoblotting

Whole kidneys were homogenized and protein was extracted from 20 mg of tissue in 1 ml

RIPA buffer supplemented with protease inhibitors (Roche) for 2 hrs. 10 µg of protein was

separated by SDS-PAGE using a 10% separating and a 4.5% stacking gel and transferred to

a PVDF membrane with 0.2 µm pore size (Roth) via a semidry transfer. The membrane was

blocked with 5% skim milk in TBS-T, cut in half and probed with the mouse anti-β-catenin

(1:1000, BD Transduction Laboratories, 610153) or mouse anti-α-Tubulin (1:10000, Sigma-

Aldrich, T9026) antibody overnight and with the peroxidase-conjugated donkey anti-mouse

antibody (1:5000, Jackson) for 1 hr. The antibodies were diluted in 5% BSA in TBS-T. Bands

were visualized using the Western Lightning Plus-ECL substrate (PerkinElmer) for 5 min and

a Fusion SL imaging system (Vilber).

Real-time qPCR

Total RNA from the organoids and whole mouse kidneys was isolated using TRIzol

(Invitrogen), and 1 µg/20 µl of RNA was reversely transcribed to cDNA using MMLV reverse

transcriptase (Promega) according to the manufacturer's instructions. Real-time qPCR

reactions were performed in a CFX96-C1000 thermal cycler (Bio-Rad) with the PowerUp

SYBR Green Master Mix (Applied Biosystems) and 10 μM exon-exon junction-spanning

primers, using a standard protocol with 44 cycles according to the manufacturer's

instructions. Primer sequences are listed in the Table 2. Primer specificity was tested by

melting curve analyses and running reactions with a negative control (without cDNA).

Relative mRNA expression values were normalized to the endogenous control Gapdh using

the 2-ΔΔCt method.

Table 2. **Primer sequences for real-time qPCR.**

Gene	Forward primer 5'-3'	Reverse primer 5'-3'
Krt8	CTCAAAGGCCAGAGGGCATC	TTAATGGCCATCTCCCCACG
Krt18	ACTGGTCTCAGCAGATTGAGG	CCGAGGCTGTTCTCCAAGTT
Cldn4	CTTCATCGGCAGCAACATCG	GATGACCATAAGGGCTCGGG
Abcb1b	AGTGGCTCTTGAAGCCGTAA	AAACTCCATCACCACCTCACG
Slc3a1	AAAATGCCTTGACTGGTGGCA	CCTCAACAGCGTATCTGAAGTCT
Slc40a1	GAGCCAGTGTCCCCAACTAC	CTTGCAGCAACTGTGTCACC
Aqp3	ATCGTTGTGGGGAGATGCTT	ACCAAGATGCCAAGGGTGAC
Atp6ap2	GGCAAAACAAGAGAACACCCA	CCAAGGCCAAGCCGATCATA
Wnk1	CCTCAAGTATGGCACAGGGG	GCTGTATTCCCTGCTGCTGA
Dkk1	CGGGGGATGGATATCCCAGA	ACGGAGCCTTCTTGTCCTTTG

Axin2	GCGCTTTGATAAGGTCCTGG	TCATGTGAGCCTCCTCTCTTTT
Cyclin D1	CTGGATGCTGGAGGTCTGTGA	AGGGGGTCCTTGTTTAGCCAG
Myc	TTGGAAACCCCGCAGACAG	GCTGTACGGAGTCGTAGTCG
Hey1	GAGCGTGAGTGGGATCAGTG	GCTTAGCAGATCCCTGCTTCT
Hes1	CTGGTGCTGATAACAGCGGA	GGAATGCCGGGAGCTATCTT
Hey3 (Heyl)	GTCTTGCAGATGACCGTGGA	CGGGCATCAAAGAACCCTGT
Ca9	GTCATTGGAGCTATGGAGG	CTCATAACCCAGAAGTTCCAG
Hk2	GTGACAGACAATGGTCTCCAGAG	GCCAGGCATTCGGCAATG
Pdk1	GCAGTTCCTGGACTTCG	CAATCTAACAGGCAACTCTTG
Ldha	GATGGATCTCCAGCATGGCAG	GTGATAATGACCAGCTTGGAGTTCG
Glut1	GCTATAACACTGGTGTCATCAACG	CGTGGTGAGTGTGGTGGATG
Vegfa	CAGGCTGCTGTAACGATGAA	TTTCTTGCGCTTTCGTTTTT
Ngal	ACAACCAGTTCGCCATGGTA	AAGCGGGTGAAACGTTCCTT
Kim-1	CAGGGTCTCCTTCACAGCAG	CCACCACCCCCTTTACTTCC
Gsta1	AGCCCGTGCTTCACTACTTC	CAATCTCCACCATGGGCACT
Il1b	TTCAGGCAGGCAGTATCA	CCAGCAGGTTATCATCATCA
Cd11b	CCACACTAGCATCAAGGGCA	GCTTCACACTGCCACCGT

Ly6c	GCAGTGCTACGAGTGCTATGG	ACTGACGGGTCTTTAGTTTCCTT
Icam1 (Cd54)	GTCCGCTGTGCTTTGAGAAC	GAGGTCCTTGCCTACTTGCT
Cd68	ACTTCGGGCCATGTTTCTCT	GCTGGTAGGTTGATTGTCGT
Ccr2 (Cd192)	GCCATCATAAAGGAGCCATACC	ATGCCGTGGATGAACTGAGG
Vim	TGGATCAGCTCACCAACGAC	AAGGTCAAGACGTGCCAGAG
Fn1	ATGAGAAGCCTGGATCCCCT	GGAAGGGTAACCAGTTGGGG
a-sma	ACATCAAGGAGAAGCTGTGCT	TTTCGTGGATGCCCGCTG
Col1a1	CATGAGCCGAAGCTAACCCC	GGGTTTCCACGTCTCACCAT
Col3a1	AGTGGGCATCCAGGTCCTAT	GGGTGAAAAGCCACCAGACT

Genomic DNA PCR

Genomic DNA was isolated from whole kidneys using the GeneJET Genomic DNA Purification Kit (Thermo Fisher Scientific) according to the manufacturer's instructions. β-catenin and Vhl alleles were amplified using 100 ng of DNA, 10 μM primers and a standard protocol with 50 and 40 cycles and annealing temperature of 65°C and 56°C, respectively. A forward primer 5'-3': GGTACCTGAAGCTCAGCG-CACAGCTG and a reverse primer 5'-3': ACGTGTGGCAAGTTCCGCGTCATCC were used to amplify the β-catenin allele (23). A forward primer 1 5'-3': CTGGTACCCAC-GAAACTGTC, a forward primer 2 5'-3': CTAGGCACCGAGCTTAGAGGTTTGCG and a reverse primer 5'-3': CTGACTTCCACTGATGCTTGTCACAG were used to amplify the Vhl allele (37).

Drug response assay

Single cells were seeded at 15,000 cells per well in 9 µl Matrigel (75%, Corning) lenses in 96-well plates and overlayed with 100 µl organoid medium. After 24 hrs, medium was replaced with fresh medium containing ICG-001 (Biochempartner) or LF3 (Selleckchem). Medium was changed every other day. CellTiterGlo assay (Promega) was performed after 6 days according to the manufacturer's instructions.

Blood assays

Blood was collected from hearts of freshly sacrificed mice to tubes coated with 0.5 M EDTA pH 8 (Invitrogen) to prevent clotting. BUN concentrations were measured using the i-STAT 1 System with the CHEM8+ cartridges for patient testing (Abbott) according to the manufacturer's instructions. For plasma collection, cells and platelets were pelleted from the blood by centrifugation for 15 min at 2000 g. EPO concentrations in supernatants (plasma) were measured using the Quantikine Mouse Erythro-poietin Immunoassay (R&D Systems) according to the manufacturer's instructions.

Descriptive statistics and significance testing

The Pearson correlation matrix (coefficient) was calculated by the cor function of the base R package between all samples of mouse kidney epithelia (control kidney, CK), early passage organoids (OEP) and long-term passage organoids (OLP) based on normalized log2 intensity values for 9,000 proteins and 16,000 phosphorylation sites detected in the mass spectrometry analysis. The matrix was plotted using the pheatmap R package. For the functional annotation clustering of the most upregulated proteins using DAVID bioinformatic tool (DAVID Bioinformatics Resources 6.8, NIAID/NIH), 9,000 proteins detected in the proteomic analysis were subjected to both the 0.1% FDR cutoff (adjusted P-value < 0.001) and log2 fold change cutoff of > 0.5 for both OEP over CK and OLP over CK. 722 proteins were selected. Using the entire mouse (Mus musculus) proteome as a background for the UNIPROT_ACCESSION identifier, 671 proteins (DAVID IDs) were identified and analyzed.

Defined DAVID defaults were used and the Enrichment Thresholds (EASE Scores, modified

Fisher-exact P-values) and adjusted P-values (Benjamini-Hochberg correction) for all terms

in each cluster were subjected to the cutoff of < 0.001. 10 clusters were determined. The

overall enrichment score for each cluster was calculated based on the Enrichment

Thresholds for all terms. The most enriched term was selected to represent each cluster

based on the lowest adjusted P-value. All other statistical analyses were performed in

GraphPad Prism (GraphPad). All data are presented as mean ± SD (error bars), unless

otherwise stated. To assess the normal distribution of the data, the Shapiro-Wilk test (α =

0.05) was performed. The unpaired, two-tailed Student's t-test and the ordinary one-way

ANOVA followed by the Dunnett's multiple comparison were used to analyze data, which

passed the normality test. To compare groups, which did not pass the normality test, the

alternative non-parametric Kruskal-Wallis test followed by the Dunn's multiple comparison

were performed. The ordinary one-way ANOVA fol-lowed by the Dunnett's multiple

comparison were used to analyze the real-time qPCR data on the organoids. A P-value <

0.05 was considered statistically significant. IC50 values were calculated by a non-linear

regression analysis of the response and log10 of the inhibitor concentration fitting a curve

with a variable slope (four parameters). Statistical details of the experiments can be found in

the figure legends.

Results

Self-renewal of long-term organoid cultures from adult mouse kidneys

We isolated single epithelial cells from whole adult mouse kidneys and established 3D organoid cultures with high efficiency in Matrigel and serum-free conditions (n = 20). The protocol was adapted to specific needs of kidney stem/progenitor cells (Table 1). Within 14 days of culture, single cells grew continuously and formed organoids with sizes up to 1.5 mm in diameter (Fig 1A). By H&E staining, we classified three organoid types: cystic organoids with one or more cell layers and a single lumen (with few organoids containing one or two additional smaller lumina), solid filled organoids, and alveolar organoids with multiple lumina, accounting for 65%, 25% and 10% (Fig 1B).

Organoid formation from single cells was five times higher in the first passage in comparison to freshly seeded cultures (Fig 1C and D), indicating self-renewal capacity of cells. For maintenance, the organoids were serially passaged starting from small cell clusters (Fig 1E), which allowed to culture them for at least 3.5 months (passage 12, S1A Fig). Organoid cultures were established from mice with a mixed C57BL/6J and FVB/NJ background, but they could also be generated from C57BL/6J mice with similar efficiency (S1B Fig). Hence, we created self-renewing long-term organoids from single adult mouse kidney epithelial cells, which predominantly exhibited cystic morphologies, consistent with organoid cultures derived from other adult organs (8).

Fig 1. Self-renewal of organoid cultures from adult mouse kidneys. (A) Brightfield images acquired on consecutive days of an organoid formed from a single freshly seeded cell. (B) H&E staining showing freshly seeded organoids with cystic (left), solid (middle) and alveolar (right) morphology. (C) Brightfield images of a freshly seeded (left) and a once passaged (right) organoid culture grown from 10^5 single cells. (D) Quantification of organoid numbers in a freshly seeded and passage 1 culture. (E) Brightfield image of a serially passaged organoid culture after 5 passages. Data information: scale bars, 100 μm in A, 50 μm in B, 500 μm in C and E. 20 freshly seeded organoid cultures were established in total. In A, one independent replicate was examined. For quantification in B, 180 organoids were counted in total. One independent replicate was examined. For quantification in C and D, 10^5

cells were seeded and the organoids with diameters ≥ 100 µm were counted after 14 (freshly seeded cells) or 7

(passaged cells) days of culture (6 wells, technical replicates). One independent replicate was examined. In D,

the graph shows mean ± SD (error bars). Data passed the Shapiro-Wilk normality test (α = 0.05). The unpaired,

two-tailed Student's t-test was performed; P-value, < 0.0001, ****p < 0.0001. In E, three independent replicates

were examined.

Examination of kidney organoids by deep proteomic and phosphoproteomic analyses

revealed that the cultures closely resembled adult mouse kidney epithelia and were

phenotypically stable over time

We performed mass spectrometry-based deep bulk proteomics and phosphoproteomics.

Three independent replicates of adult kidney epithelial cells (control kidney), early passage

organoids (passage 2, 1-month culture) and long-term passage organoids (passage 12, 3.5-

month culture) were compared. To prepare control samples, Epcam-positive kidney epithelial

cells were MAC-sorted from freshly isolated single cell suspensions from whole mouse

kidneys, similar to a protocol for preparing adult human kidney organoids (10). We could

detect 9,000 proteins and 16,000 phosphorylation sites of these proteins, which is the

maximum coverage currently possible (32). We calculated Pearson correlations between the

samples and found high correlations, at least 0.8, between adult kidney epithelial cells

(control kidney, CK) and both early passage organoids (OEP) and long-term passage

organoids (OLP) for both proteome (Fig 2A) and phosphoproteome (Fig 2B). This indicated

that the organoids closely resembled kidney epithelia. Moreover, OLP recapitulated OEP for

both proteome (Fig 2A) and phosphoproteome (Fig 2B), as shown by correlations of around

1, confirming that organoid cultures were phenotypically stable over time.

Kidney organoids displayed enhanced proliferation, and upregulation of stem cell-

associated Wnt, Notch and Yap signaling

We also recorded the importance of many upregulated proteins in the organoids in

comparison to kidney epithelia. Using a bioinformatic tool for functional annotation clustering,

DAVID, we discovered 10 biological processes in the organoids, which were governed by

upregulated proteins (Fig 2C). The top three enriched processes were cell cycle, ribosomal

activity and DNA replication, which indicated the presence of proliferating cells. We

confirmed this by BrdU incorporation in nuclei (marked in magenta, Fig 2D). In line with the

self-renewal capacity, enhanced proliferation suggested enrichment in stem/progenitor cells

in the organoids and thus the recapitulation of the potential of the adult kidney for renewal.

DAVID analysis also revealed enrichment in cell polarity-related proteins of adherens

junctions and basement membranes (Fig 2C).

By manual curation, proteomic analysis in both OEP and OLP revealed upregulation of

components of stem cell-associated Wnt (Fig 2E) and Notch (Fig 2F) signaling. Upregulated

Wnt components included ligands Wnt4, Wnt7b and Wnt10a, co-receptors Lrp5 and Lrp6,

the destruction complex disassembly-mediating protein Dvl2 and targets Axin2, Birc5, Cd44

and Cyclin D1 (Ccnd1). Upregulated Notch components included the ligand Jag2, receptors

Notch1, Notch2 and Notch3, the S2 cleavage-mediating metalloproteinase Adam17, S3

cleavage γ-secretase complex-related Aph1a and Ncstn as well as co-activators Rbpj,

Maml1, Maml2 and Kat2b. Proteomic analysis also revealed in both OEP and OLP

upregulation of another stem cell-associated signaling system, Yap (S2 Fig). Upregulated

Yap components were direct Yap targets. Thus, the organoids and adult mouse kidney

epithelia largely shared proteome and phosphoproteome patterns, which remained

unchanged over passages. Moreover, strong proliferation of the organoids and upregulation

of the Wnt, Notch and Yap signaling systems were the signs of the enrichment in

stem/progenitor cells.

Fig 2. Kidney organoids closely resemble adult mouse kidney epithelia, are phenotypically stable over

time, and display enhanced proliferation and stem cell-associated Wnt and Notch signaling. (A) Heatmap

for the Pearson correlation matrix (coefficient) of the expression levels of 9,000 proteins between all samples of

mouse kidney epithelia (control kidney, CK), early passage organoids (OEP) and long-term passage organoids

(OLP). The experimental columns were correlated against each other. (B) Heatmap for the Pearson correlation

matrix (coefficient) of the phosphorylation levels of 16,000 corresponding phosphorylation sites between all

samples of mouse kidney epithelia (control kidney, CK), early passage organoids (OEP) and long-term passage

organoids (OLP). The experimental columns were correlated against each other. (C) Functional annotation

clustering of the most upregulated proteins in the organoids using DAVID bi-oinformatic tool. Shown are 10 most

enriched clusters, based on the overall enrichment scores, with their representative terms (biological processes),

based on the adjusted P-values. (D) 2D immunofluorescence for BrdU (magenta) of an organoid culture. (E)

Proteomic heatmap for components of Wnt signaling in both early (OEP) and long-term (OLP) passage organoids

in comparison to mouse kidney epithelia (control kidney, CK). (F) Proteomic heatmap for components of Notch

signaling in both early (OEP) and long-term (OLP) passage organoids in comparison to mouse kidney epithelia

(control kidney, CK). Data information: scale bars in D, 50 μm. Nuclei are counterstained with DAPI. Three

independent replicates were examined. In E and F, the heatmaps show normalized log2 intensity values for three

independent replicates of CK, OEP and OLP. A 5% FDR (adjusted P-value < 0.05) cutoff and a log2 fold change

cutoff of > 0 were applied for both OEP over CK and OLP over CK. The values were scaled (z-score by row) with

breaks from ≤ -2 to ≥ 2.

Kidney organoids contained differentiated and polarized tubular epithelial cells

Ubiquitous staining of Pax8 in nuclei (marked in brown) confirmed that organoid cells were of

kidney origin (Fig 3A). The organoids contained differentiated kidney epithelial cells, as

indicated by the presence of adherens junctions between lateral membranes of neighboring

cells, which were positive for E-cadherin (E-cad, marked in orange, Fig 3B), and by

upregulation of Keratin 8 (Krt8), Claudin 4 (Cldn4) and *Keratin 18 (Krt18)* (Fig 3C). In line

with DAVID analysis, cells displayed pronounced epithelial polarity, as shown by the location

of Laminin in the basal laminae (marked in green) and of Zo-1-positive tight junctions at the

apical sides (marked in red) (Fig 3D).

Cells were positive either for Lotus Tetragonolobus Lectin (LTL, marked in red), a marker of

proximal tubular epithelial cells, or Peanut Agglutinin (PNA, marked in green), a marker of

epithelial cells of distal nephrons, i.e. distal tubules and collecting ducts, and both markers

were detected within the same organoids (Fig 3E). Proteomic analysis revealed enrichment

in 48 membrane transporters in both OEP and OLP (Fig 3F), including known transporters of

proximal tubular cells such as amino acid transporters Slc1a4 and Slc1a5, the glucose

transporter Slc2a1, the myo-inositol transporter Slc5a11, ion transporters Slc12a6 and

Slc26a2, zinc transporters Slc30a6 and Slc39a10 and efflux transporters Abcb1b, Abcc3 and

Abcc4. In addition, known transporters of distal nephrons were upregulated, including

Slc6a9, a glycine transporter of distal and connecting tubules, and ion transporters of

collecting ducts, Slc4a1ap, Slc4a7, Atp6ap2 and Trpv4. Proton transporters Atp6v0a1 and

Atp6v0a2, which are present in both proximal tubules and collecting ducts, were also

enriched. Proteomic analysis also revealed upregulation of Wnk1, a common cytoplasmic

marker (a kinase) of distal and connecting tubules and of cortical collecting ducts, and Gata3,

a nuclear marker (a transcription factor) of collecting ducts (Fig 3G). In LTL/PNA-double-

positive organoids, we confirmed in a subset of LTL-positive cells expression of Abcb1b

(marked in cyan), another marker of proximal tubular cells, which did not overlap with PNA-

positive cells of distal nephrons (Fig 3H). In these organoids, we also detected in a subset of

PNA-positive cells expression of Aquaporin 3 (Aqp3, marked in magenta), a marker of

principal cells of collecting ducts, which was mutually exclusive with LTL-positive cells (Fig

3I). We also observed mutually exclusive expression of LTL and PNA in solid and alveolar

organoids (S3A Fig). Whole-mount confocal microscopy confirmed expression of LTL, PNA

and E-cad in 3D reconstructed cystic organoids (S3B Fig and C, and S1 Video).

Fig 3. Kidney organoids exhibit markers of differentiated and polarized tubular epithelial cells. (A)

Immunohistochemistry for Pax8 (brown) of an organoid culture. (B) 2D immunofluorescence for E-cad (orange) of

an organoid culture. (C) Proteomic heatmaps for Krt8 and Cldn4 in both early (OEP) and long-term (OLP)

passage organoids in comparison to mouse kidney epithelia (control kidney, CK; upper part), and real-time qPCR

analysis of gene expression of Krt18 in both early (OEP) and long-term (OLP) passage organoids in comparison

to whole mouse kidney cells (control kidney, CK; lower part). (D) 2D immunofluorescence for Laminin (green) and

Zo-1 (red) of an organoid culture. (E) 2D immunofluorescence for LTL (red) and PNA (green) of an organoid

culture. (F) Proteomic heatmap for 48 membrane transporters in both early (OEP) and long-term (OLP) passage

organoids in comparison to mouse kidney epithelia (control kidney, CK). (G) Proteomic heatmaps for Wnk1 and

Gata3 in both early (OEP) and long-term (OLP) passage organoids in comparison to mouse kidney epithelia

(control kidney, CK). (H) 2D immunofluorescence for Abcb1b (cyan), LTL (red) and PNA (green) of an organoid

culture. (I) 2D immunofluorescence for Aqp3 (magenta), LTL (red) and PNA (green) of an organoid culture. Data

information: scale bars in A/B/D/E/H/I, 50 μm. In A, nuclei are counterstained with haematoxylin; in B/D/E/H/I,

nuclei are counterstained with DAPI. In C, the graph shows mean ± SD (error bars). The ordinary one-way

ANOVA followed by the Dunnett's multiple comparison to CK were performed; both P-values, 0.0001, ****p⊥<

0.0001. In C/F/G, the heatmaps show normalized log2 intensity values for three independent replicates of CK,

OEP and OLP. A 5% FDR (adjusted P-value < 0.05) cutoff and a log2 fold change cutoff of > 0 were applied for

both OEP over CK and OLP over CK. Values were scaled (z-score by row) with breaks from ≤ -2 to ≥ 2. In

A/B/D/E/H/I, three independent replicates were examined. For the real-time qPCR analysis of Krt18 in C, three

independent replicates in technical triplicates were examined.

Transmission electron microscopy confirmed epithelial cell polarity and complexity of the

organoids. Cystic organoids developed microvilli (MV) at apical membranes towards the

lumen (L), a typical feature of proximal tubular cells (S4A Fig). At the basal side, microvilli-

free basal laminae (arrowhead) and filopodia (F) were seen (S4A Fig). Junctional complexes

(JC) between neighboring cells, which are involved in barrier formation, were present at

lateral membranes at the luminal side. JC included from the apical to basal side: tight

junctions (zonula occludens, black asterisk), adherens junctions (belt desmosomes, zonula

adhaerens, white asterisk), spot desmosomes (macula adhaerens, red asterisks) and closely

aligned or nearly fused lateral membranes (white and black arrowheads, respectively) (S4B

Fig). At the basal side, we observed spot desmosomes (red asterisk) and closely aligned or

nearly fused lateral membranes (white and black arrowheads, respectively) (S4C Fig).

Endocytic events, i.e. clathrin-coated pits with diameters of 100 nm, which are typical for

proximal tubular cells (38), were observed at apical and lateral membranes (marked by

arrowheads, S4D Fig). Together, apart from stem/progenitor cells, kidney organoids

contained differentiated tubular epithelial cells with apical and basal polarity and complex

intercellular junctions.

Wnt signaling controlled self-renewal, growth and differentiation of kidney organoids

We examined the dependence of the organoids on Wnt signaling by removing or adding

single Wnt activators and by blocking Wnt with small-molecule inhibitors. Withdrawal of the

Wnt activator R-spondin-1 reduced organoid numbers (self-renewal) and sizes. Addition of

the Wnt ligand Wnt3a did not improve organoid yield (Fig 4A and B), possibly due to

autocrine supply of Wnt4, Wnt7b or Wnt10a. Treatments with ICG-001, an inhibitor that

blocks β-catenin-Tcf-mediated transcription, and with LF3, another β-catenin-Tcf inhibitor

developed in our laboratory (39) (Fig 4C), produced concentration-dependent inhibition of

organoid growth (Fig 4D and F) and concentration-dependent decrease in cell viability (Fig

4E and G). By H&E staining, we observed upon R-spondin-1 removal a switch from

predominant cystic to fully solid organoids, while treatment with ICG-001 at IC50 did not

result in morphological changes (Fig 4H). Transition to solid morphologies induced by R-

spondin-1 withdrawal prompted to study the importance of Wnt signaling for the control of

organoid differentiation. By real-time qPCR, we examined in the organoids upon R-spondin-1

removal or ICG-001 treatment expression of selected genes, which are induced in

differentiated kidney tubular epithelial cells. We found increases in abundance of *Krt8*, *Krt18*

and *Cldn4* (Fig 4I). We also observed upregulation of proximal tubule genes *Abcb1b*, *Slc3a1*

and *Slc40a1* (Fig 4J) as well as of distal nephron genes *Aqp3*, *Atp6ap2* and *Wnk1* (Fig 4K).

Immunofluorescence staining confirmed upregulation of Aqp3 in these organoids (Fig 4L).

Altogether, mouse kidney organoid cultures required Wnt signaling to promote self-renewal,

growth and differentiation into proximal and distal nephron lineages.

Fig 4. Wnt signaling controls kidney organoid formation, growth and differentiation. (A) Organoid numbers

per well with indicated growth factor combinations (withdrawal or addition). (B) Diameters (μm) of the organoids

(single dots) from wells with indicated growth factor combinations (withdrawal or addition). (C) Scheme of Wnt

signaling with a common target for the small-molecule inhibitors ICG-001 and LF3. (D and F) Brightfield images of

organoid cultures at indicated ICG-001 (D) and LF3 (F) concentrations (μM) on day 6 of treatment. (E and G)

Curves showing the dependence of organoid viability (fold change ATP luminescence) on log10 ICG-001 (E) and

LF3 (G) concentration (μM) on day 6 of treatment. (H) H&E staining of organoid cultures upon R-spondin-1

removal or treatment with ICG-001 at IC50. (I) Real-time qPCR analysis of gene expression of differentiation

markers Krt8, Krt18 and Cldn4 in the organoids upon R-spondin-1 removal or treatment with ICG-001 at IC50. (J)

Real-time qPCR analysis of gene expression of markers of proximal tubular cells, Abcb1b, Slc3a1 and Slc40a1, in

the organoids upon R-spondin-1 removal or treatment with ICG-001 at IC50. (K) Real-time qPCR analysis of gene

expression of markers of distal nephrons, Aqp3, Atp6ap2 and Wnk1, in the organoids upon R-spondin-1 removal

or treatment with ICG-001 at IC50. (L) 2D immunofluorescence for Aqp3 (magenta) in the organoids upon R-

spondin-1 removal or treatment with ICG-001 at IC50. Data information: in A and B, culture medium containing

DMEM/F12 with GlutaMax, HEPES, N-acetylcysteine, B27, N2, nicotinamide, hydrocortisone (HC) and

prostaglandin E2 (PGE2) is common to all the conditions. For quantification, 10^5 cells were freshly seeded and

the organoids with diameters ≥ 200 μm were counted after 14 days of culture. The graphs show mean ± SD (error

bars). In A, data passed the Shapiro-Wilk normality test (α = 0.05). The ordinary one-way ANOVA followed by the

Dunnett's multiple comparison to complete organoid medium (Ctrl, first from the left) were performed; P-values

from left to right, 0.0001, 0.2132; ****p⊥< 0.0001, ns: non-significant. In B, data did not pass the Shapiro-Wilk

normality test (α = 0.05). The alternative non-parametric Kruskal-Wallis test followed by the Dunn's multiple

comparison to complete organoid medium (Ctrl, first from the left) were performed; P-values from left to right,

0.0090, 0.0129; **p⊥< 0.01, *p⊥< 0.05. In A and B, three independent replicates with 6 technical replicates

(wells) were examined and collectively shown. Ctrl in D-G, the organoids cultured in complete medium and

treated with DMSO at re-spective concentrations. Scale bars in D and F, 500 μm. In E and G, the curves show

mean ± SD (error bars). In E, ICG-001 produced the IC50 of 14.66 μM; in G, LF3 produced the IC50 of 15.72 μM.

In D-G, three independent replicates in technical triplicates (wells) were examined. Ctrl in H-L, the organoids

cultured in complete medium and treated or not with DMSO at a respective concentration. Scale bars in H, 100

μm. Three independent replicates were examined. In I-K, the graphs show mean ± SD (error bars). The ordinary

one-way ANOVA followed by the Dunnett's multiple comparison to Ctrl were performed; P-values from left to right

in I; 0.0004, 0.0001 for Krt8; 0.0153, 0.0002 for Krt18; 0.0002, 0.0003 for Cldn4; in J: 0.0001, 0.0001 for Abcb1b;

0.0005, 0.0164 for Slc3a1; 0.0910, 0.0001 for Slc40a1; in K: 0.0001, 0.0001 for Aqp3; 0.0014, 0.0001 for

Atp6ap2; 0.001, 0.0001 for Wnk1; ****p⊥< 0.0001, ***p⊥<⊥0.001, **p⊥<⊥0.01, *p⊥<⊥0.05, ns: non-significant.

Three independent replicates in technical triplicates were examined. Scale bars in L, 50 μm. Three independent

replicates were examined.

Generation of double kidney mutants in mice to model stem cell-driven development

of human ccRCC

To investigate a relation between Wnt-high and Notch-high kidney stem/progenitor cells

enriched in our mouse kidney organoids and in human ccRCC, we genetically modified Wnt

and Notch signaling in mice. We generated two types of double mouse mutants by the loxP-

mediated recombination of adult kidney epithelial cells: i) Wnt-β-catenin gain of function

combined with Notch1 intracellular domain (N1icd) gain of function, hereafter called β-

catenin-GOF; Notch-GOF, and ii) Von Hip-pel-Lindau loss of function combined with Wnt-β-

catenin gain of function, hereafter called Vhl-LOF; β-catenin-GOF. For the details of the

generation of the genetic system and mutants, see Fig 5A. Vhl-LOF; Notch-GOF mutant

mice had already been established using different Cre systems (19,20).

To examine cellular patterns of recombination, we established Pax8-rtTA(+); LC1-Cre(+);

LacZ(+) reporter mice, which carried a loxP-flanked stop cassette upstream of the LacZ

sequence encoding β-galactosidase. After doxycycline treatment of one-month-old mice,

immunohistochemistry revealed Cre expressed in nuclei of kidney epithelial cells in both

cortical and medullary parts, including proximal tubular cells (Fig 5B). The Cre pattern was

confirmed by cytoplasmic LacZ staining after treatment with X-gal. Glomeruli (marked by

black lines) remained negative for Cre and LacZ. We did not observe Cre- and LacZ-positive

cells in doxycycline-treated Pax8-rtTA(-); LC1-Cre(+); LacZ(+) control kidneys, indicating that

the genetic system was not leaky. Then, we treated with doxycycline critical one-month-old

Pax8-rtTA(+); LC1-Cre(+); β-catenin loxP/wt; Notch loxP/wt (β-catenin-GOF; Notch-GOF)

mice and Pax8-rtTA(+); LC1-Cre(+); Vhl loxP/loxP; β-catenin loxP/wt (Vhl-LOF; β-catenin-

GOF) mice (n = 20 per line). We also treated controls carrying β-catenin- and Notch- or Vhl-

and β-catenin-loxP alleles, but no Pax8-rtTA or LC1-Cre (n = 20).

We examined in kidney mutants the recombination of β-catenin, Notch and Vhl alleles on

genomic DNA, protein and mRNA target gene levels. We detected DNA bands of

recombined (700 bp) and wild-type (900 bp) β-catenin alleles in both mutants, as compared

to control kidneys with only wt alleles (Fig 5C). In Vhl-LOF; β-catenin-GOF mutant, we

detected DNA bands of recombined (260 bp) and non-recombined loxP (460 bp) Vhl alleles,

as opposed to control kidneys with only loxP alleles (Fig 5D). Recombined β-catenin protein

accumulated in both mutants (72 kDa band), as compared to controls with only wt protein (95

kDa band) (Fig 5E). We observed nuclear N1icd in β-catenin-GOF; Notch-GOF mutant, but

not in controls (Fig 5F). We found upregulation of β-catenin target genes *Dkk1*, *Axin2*, *Cyclin*

D1 and *Myc* in β-catenin-GOF; Notch-GOF (Fig 5G) and Vhl-LOF; β-catenin-GOF (Fig-ure

5H) mutants, as opposed to controls. Expression of Notch target genes *Hey1*, *Hes1* and

Hey3 in β-catenin-GOF; Notch-GOF mutant (Fig 5I) and of Hif-1α/2α target genes *Ca9*, *Hk2*,

Pdk1, *Myc*, *Ldha*, *Cyclin D1*, *Glut1* and *Vegfa* in Vhl-LOF; β-catenin-GOF mutant (Fig 5J)

also increased, as compared to controls. Thus, our data show that in mice targeted

mutagenesis in adult kidney epithelial cells occurred and that mutant cells were not cleared

over time.

Fig 5. Generation of double kidney mutants in mice. (A) Scheme of the genetic system to generate mutant

kidneys. (B) Time course and pattern of recombination in kidneys of Pax8-rtTA(+); LC1-Cre(+); LacZ(+) reporter

mice. (C) PCR for DNA recombination of the loxP-flanked exon 3 sequence of β-catenin gene in β-catenin-GOF;

Notch-GOF and Vhl-LOF; β-catenin-GOF mutant kidneys (700 bp) versus controls (900 bp). (D) PCR for DNA

recombination of the loxP-flanked promoter-exon 1 sequence of Vhl gene in Vhl-LOF; β-catenin-GOF mutant

kidneys (260 bp) versus controls (460 bp). (E) Immunoblotting for recombination of β-catenin in β-catenin-GOF;

Notch-GOF and Vhl-LOF; β-catenin-GOF mutant kidneys (72 kDa) versus controls (95 kDa). (F)

Immunohistochemistry for nuclear N1icd in β-catenin-GOF; Notch-GOF mutant cortical kidneys versus controls.

(G) Fold change of expression of selected β-catenin target genes in β-catenin-GOF; Notch-GOF mutant kidneys

versus controls. (H) Fold change of expression of selected β-catenin target genes in Vhl-LOF; β-catenin-GOF

mutant kidneys versus controls. (I) Fold change of expression of selected Notch target genes in β-catenin-GOF;

Notch-GOF mutant kidneys versus controls. (J) Fold change of expression of selected Hif-1α/2α target genes in

Vhl-LOF; β-catenin-GOF mutant kidneys versus controls. Data information: abbreviation in A; dox, doxycycline.

Scale bars in B, 100 μm. In the upper panel, nuclei are counterstained with haematoxylin; in the lower panel,

nuclei are counterstained with nuclear fast red. Glomeruli are marked by black lines. Two independent replicates

of Pax8-rtTA(+); LC1-Cre(+); LacZ(+) mice and one of Pax8-rtTA(-); LC1-Cre(+); LacZ(+) controls were examined.

Abbreviations in C-E; wt, wild-type; ex3 del, exon 3 deletion; prom-ex1 loxP/del, promoter-exon 1 loxP/deletion.

Scale bars in F, 100 μm. Nuclei are counterstained with haematoxylin. In C-F, three independent replicates per

line were examined. In G-J, graphs show mean + SD (error bars). Three independent replicates (mice, dots) in

technical triplicates were analyzed. No statistical test was performed because of high inter-mouse variance.

However, increase trends were observed.

No tumorigenesis was observed in double mutant kidneys

Mutant mice were examined 1-8 months after the doxycycline pulse and exhibited severe

illness symptoms. These included reduced body weight (Fig 6A) and enlarged spleens,

called splenomegaly (Fig 6B and D). Mutant kidneys appeared to be of smaller sizes (Fig

6B), but their relative weight did not change (Fig 6C). Kidneys and spleens of mutant mice

were pale, suggesting anemia (Fig 6B).

We performed Periodic acid-Schiff (PAS) staining of mutant kidneys, which detects tissues

with high proportions of sugar macromolecules such as tubular and glomerular basement

membranes and brush borders of proximal tubules. Examination by pathologists did not

reveal tumors or premalignant lesions such as dysplasia or adenomas, even after 8 months

after the doxycycline pulse. Non-neoplastic pathological changes of kidney tissues like cysts,

hypertrophy, necrosis, dilation or atrophy of tubules as well as tubule-interstitial fibrosis were

neither present (Fig 6E). Nuclear crowding was observed in some tubules in the cortex in β-

catenin-GOF; Notch-GOF, but not in Vhl-LOF; β-catenin-GOF kidneys (Fig 6F). We

conducted Ki67 staining for proliferating cells at the G1, S, G2 and M phases of the cell

cycle. None of mutant kidneys, even after 8 months after the doxycycline pulse, exhibited

higher numbers of Ki67-positive nuclei than controls (Fig 6G and H), despite upregulation of

the genes of Cyclin D1 and Myc involved in cell cycle progression.

Fig 6. No tumorigenesis was observed in double mutant kidneys. (A) Body weight of β-catenin-GOF; Notch-

GOF and Vhl-LOF; β-catenin-GOF mutant mice versus controls. (B) Overview of kidneys and spleens of β-

catenin-GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF mutant mice versus controls. (C) Kidney weight to body

weight ratio of β-catenin-GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF mutant mice versus controls. (D) Spleen

weight to body weight ratio of β-catenin-GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF mutant mice versus

controls. (E) PAS staining in β-catenin-GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF mutant kidneys versus

controls (in the cortex and medulla) after 8 months after the doxycycline pulse. (F) PAS-stained outer part of the

cortex with nuclear crowding (arrowheads) in some tubules in β-catenin-GOF; Notch-GOF, but not in Vhl-LOF; β-

catenin-GOF mutant kidneys. (G) Immunohistochemistry for nuclear Ki67 in β-catenin-GOF; Notch-GOF and Vhl-

LOF; β-catenin-GOF mutant cortical kidneys versus controls after 8 months after the doxycycline pulse. (H)

Quantification of Ki67-positive nuclei in β-catenin-GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF mutant cortical

kidneys versus controls after 8 months after the doxycycline pulse. Data information: in A, C and D, 5

independent replicates (mice, dots) per line were examined. Graphs show mean ± SD (error bars). Data passed

the Shapiro-Wilk normality test (α = 0.05). The ordinary one-way ANOVA followed by the Dunnett's multiple

comparison to controls was performed; P-values from left to right in A; 0.0001, 0.0002; ****p⊥< 0.0001,

***p⊥<⊥0.001; in C; 0.4879, 0.8354; ns: non-significant; in D; 0.0001, 0,0229; ****p⊥< 0.0001, *p⊥<⊥0.05. Scale

bar in B, 1 cm. 20 independent replicates (all mice induced) per line were examined. Scale bars in E-G, 100 μm.

Nuclei are counterstained with haematoxylin. In E and G, insets are enlarged on the right. 20 independent

replicates (all mice induced) per line were examined. In H, three independent replicates (mice, dots) with 10

technical replicates (fields per kidney) each per line were examined. Graphs show mean + SD (error bars). Data

passed the Shapiro-Wilk normality test (α = 0.05). The ordinary one-way ANOVA followed by the Dunnett's

multiple comparison to controls was performed; P-values from left to right, 0.6019, 0.9871; ns: non-significant.

We neither detected accumulation of phospho-γH2AX nor p53 stabilization and the activation

of its downstream target p21 in nuclei of mutant kidney cells (S5A Fig). We also did not

observe increases in the numbers of apoptotic cells as examined by TUNEL assays (S5B

Fig). Together, we neither found signs of malignant transformation nor of DNA damage,

senescence and cell death in mutant kidneys.

Mutant mice displayed phenotypes of CKD

To confirm anemia in mutant mice, we determined the hematocrit, which significantly

decreased in mutant mice (Fig 7A). We also measured the concentration of erythropoietin

(EPO) in the blood plasma, which strongly rose in mutant mice (Fig 7B). Pathological

examination of the enlarged spleens of mutant mice revealed extramedullary hematopoiesis

by the presence of megakaryocytes (Fig 7C).

We found upregulation of tubular injury markers *Ngal*, *Kim-1* and *Gsta1* in mutant kidneys

(Fig 7D). Moreover, the concentration of another marker of kidney injury, blood urea nitrogen

(BUN), was increased in mutant mice (Fig 7E). We also examined in mutant kidneys

expression of the inflammatory cytokine *Il1b* and of a panel of markers for myeloid lineage

immune cells, *Cd11b*, *Ly6c*, *Ccr2* and *Cd68*, as well as of *Icam1*, a ligand for *Cd11b*.

Expression of *Il1b* and *Cd11b* was upregulated in both kidney mutants, while expression of

Ly6c, *Ccr2*, *Cd68* and *Icam1* was increased in Vhl-LOF; β-catenin-GOF mutant, but not in β-

catenin-GOF; Notch-GOF mutant (Fig 7F). Expression of fibrotic markers, i.e. markers of

activated myofibroblasts, *a-sma* and *Vim*, and markers of extracellular matrix, *Fn1*, *Col1a1*

and *Col3a1*, was not enhanced in mutant kidneys (Fig 7G). We conclude that mutant mice

exhibited lethal features of CKD such as anemia as well as kidney injury and kidney

inflammation. Thus, we decided not to proceed with the generation of Vhl-LOF; β-catenin-

GOF; Notch-GOF triple mutant mice.

Fig 7. Mutant mice displayed phenotypes of chronic kidney disease (CKD). (A) Hematocrit of β-catenin-

GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF mutant mice versus controls. (B) Concentration of plasma EPO in

β-catenin-GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF mutant mice versus controls. (C) PAS staining showing

extramedullary hematopoiesis in spleens of β-catenin-GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF mutant

mice versus controls. Megakaryocytes are marked by arrowheads. (D) Fold change of gene expression of kidney

injury markers in β-catenin-GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF mutant kidneys versus controls. (E)

Concentration of blood urea nitrogen (BUN) in β-catenin-GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF mutant

mice versus controls. (F) Fold change of gene expression of inflammation markers in β-catenin-GOF; Notch-GOF

and Vhl-LOF; β-catenin-GOF mutant kidneys versus controls. (G) Fold change of gene expression of fibrotic

markers of myofibroblasts and extracellular matrix in β-catenin-GOF; Notch-GOF and Vhl-LOF; β-catenin-GOF

mutant kidneys versus controls. Data information: in A, 5 independent replicates (mice, dots) with one technical

replicate each per line were examined. Graphs show mean ± SD (error bars). Data passed the Shapiro-Wilk

normality test (α = 0.05). The ordinary one-way ANOVA followed by the Dunnett's multiple comparison to controls

was performed; P-values from left to right, 0.0001, 0.0001; ****p⊥< 0.0001. In B, 5 independent replicates (mice,

dots) in technical triplicates per line were examined. Graphs show mean ± SD (error bars). No statistical test was

performed because of high inter-mouse variance. However, increase trends were observed. Scale bars in C, 100

μm. Nuclei are counterstained with haematoxylin. An undisrupted follicular B-cell compartment in the white pulp in

controls is marked by white lines. Insets are enlarged on the right. Three independent replicates per line were

examined. In D, three independent replicates (mice, dots) in technical triplicates were examined. Graphs show

mean + SD (error bars). No statistical test was performed because of high inter-mouse variance. However,

increase trends were observed. In E, 5 independent replicates (mice, dots) with one technical replicate each per

line were examined. Graphs show mean ± SD (error bars). Data passed the Shapiro-Wilk normality test (α =

0.05). The ordinary one-way ANOVA followed by the Dunnett's multiple comparison to controls was performed; P-

values from left to right, 0.0109, 0.0888; *p⊾<⊾0.05, ns: non-significant. Although the difference in average

concentration of BUN between the Vhl-LOF; β-catenin-GOF mutant mice and controls was non-significant

(p⊾<⊾0.05), an increase trend was observed. In F and G, three independent replicates (mice, dots) in technical

triplicates were examined. Graphs show mean + SD (error bars). No statistical test was performed because of

high inter-mouse variance. However, increase trends were observed in F and no increase trends were observed

in G.

Discussion

Tubular organoid cultures from adult mouse kidneys enrich for stem/progenitor cells with active Wnt and Notch signaling

We have established robust organoid cultures from single epithelial cells from adult mouse kidneys and have examined their cellular components and stem cell-associated signaling systems at protein levels by deep bulk proteomic and phosphoproteomic analyses. These approaches represent so far the most thorough recording of important proteins and phosphorylation sites performed in kidney organoids. We found that the organoids shared proteome and phosphoprotome patterns with adult kidney tubular epithelia to a great extent and were phenotypically stable over long-term passages. Proteomic analysis revealed enhanced proliferation and upregulation of stem cell-associated Wnt, Notch and Yap signaling in the organoids, which indicated the recapitulation of kidney renewal response, similar to mice upon kidney injury (3–6,40). Functionally, Wnt was crucial for both self-renewal and differentiation of organoid cells. Despite high coverage, we were not able to detect important proteins such as collecting duct-specific Aquaporin 3, which we detected using other techniques. To characterize the organoids more thoroughly, single cell RNA sequencing should be performed in the future. Once optimized, single cell proteomics will remarkably improve a global analysis of protein levels in individual organoid cells (41). We did not examine the functional importance of phosphorylation patterns of detected proteins, because this went beyond the scope of our manuscript. However, we believe that the phosphoproteomic data set holds a lot of interesting information for other researchers regarding signaling systems of kidney stem cells. In future studies, the or-ganoid model will assist analyses of Wnt- and Notch-dependent self-renewal of marker-sorted kidney stem/progenitor cell populations, which will then possibly link these cells with their malignant counterparts, for instance, with the ones identified by our group (14).

Our findings in genetic models of adult mouse kidneys challenge the current understanding of the involvement of stem cell-associated Wnt and Notch signaling in development of human ccRCC

Surprisingly, the two types of double kidney mutants in mice, i.e. β-cat-GOF with Notch-GOF or with Vhl-LOF, did not produce tumors or premalignant lesions. This occurred despite a rational hypothesis on the involvement of Wnt-high and Notch-high stem cells in development of human ccRCC, and despite using a well-designed genetic system to target adult proximal tubular epithelial cells, the cells of tumor origin. However, the mouse mutants displayed strong fatal phenotypes that resembled CKD (42,43). The significance of the interplay of Wnt and Notch as well as of Wnt and Vhl loss in development of CKD has not previously been defined.

Wnt hyperactivation using embryonic Cres in other mouse models resulted in development of polycystic kidney disease (44,45). Wnt upregulation in tubular cells or podocytes of adult mouse kidneys contributed to either tubular and glomerular damage and tubule-interstitial inflammation or to albuminuria, indicating CKD (46,47). In another mouse model, adult kidneys with Notch hyperactivation exhibited CKD with pronounced tubular degeneration and dilation as well as tubule-interstitial inflammation and fibrosis (48). By single cell RNA sequencing, plasticity of collecting duct cells in the mouse kidney was revealed, which upon Notch upregulation was disturbed and correlated with metabolic acidosis observed in CKD patients (49). Moreover, in numerous rodent models of kidney injury, transient activation of Wnt and Notch accelerated regeneration, but sustained activation promoted maladaptive behaviors and CKD progression (50). Of note, Notch upregulation in mouse kidneys in another study led to development of both CKD and a relatively rare papillary kidney cancer, and this process was accelerated by kidney injury (51). In addition, the authors found a correlation of a kidney injury episode with subsequent occurrence of papillary cancer but not ccRCC in patients.

Lack of fibrosis in our models might suggest that mutant mice have not yet progressed to

end-stage CKD. Remarkably, mutant mice produced additional features of CKD, which so far

have not been reported in other studies that examined hyperactivation of either Wnt or Notch

in adult mouse kidneys (47,48). Mutant mice exhibited dramatic anemic phenotypes, despite

strong increases in plasma erythropoietin levels. This compensative positive feedback on

EPO production suggests a mechanism of anemia development downstream of EPO supply.

Inflammation that we detected in mutant mice could be a driver of anemia in CKD, either

directly through inhibition of EPO-mediated erythropoiesis or indirectly via disrupted iron

regulation that limits hemoglobin synthesis in erythroid cells in the bone marrow (43).

Moreover, we documented extramedullary hematopoiesis in enlarged spleens of mutant

mice, which may be induced in response to anemia and involves expansion and

differentiation of erythroid, myeloid and platelet precursors into effector cells outside the bone

marrow (52).

We suspect that focal recombination in adult kidney epithelial cells in mice through

decreased doses of doxycycline might support tumor development by overcoming severe

CKD phenotypes (53,54). Thus, our reproducible and penetrant genetic system can be used

for further studies of downstream effectors of Wnt and Notch signaling in the context of the

pathobiology of kidney stem/progenitor cells, irrespective whether this results in CKD or

tumors. We conclude that the organoid model displayed responses of kidney stem cells that

resulted in high self-renewal, in contrast to the genetic mouse mutants that produced CKD

phenotypes, but no elevated proliferation and tumors.

Author Contributions

Conceptualization, A.M., A.F., P.M. and W.B.; investigation, A.M., O.P., S.K. and S.J.; formal analysis and visualization, A.M., O.P., S.K. and A.S.; writing, A.M., L.C.P. and W.B.; supervision, P.M. and W.B.

Acknowledgments

We would like to thank Prof. Dr. med. Wolfgang Schneider and Dr. med. Ann-Christin von Brünneck from the Institute of Pathology at the Charité-Medical University Berlin for the histopathological examinations. Further, we thank Prof. Dr. med. Friedrich Luft from the Experimental and Clinical Research Center (ECRC) in Berlin for giving us important experimental suggestions. We would also like to acknowledge Dr. Bart Spee from the Faculty of Veterinary Medicine at the Utrecht University for his valuable comments on the manuscript.

Funding

A.M. and A.F. were funded in part by the Urological Research Foundation (Stiftung Urologische Forschung) in Berlin.

Data Availability Statement

The mass spectrometry proteomics and phosphoproteomics data were deposited to the ProteomeXchange Consortium via the PRIDE partner depository (55) with the dataset identifier PXD023491.

References

1. Grigoryan T, Wend P, Klaus A, Birchmeier W. Deciphering the function of canonical Wnt signals in

development and disease: conditional loss- and gain-of-function mutations of -catenin in mice. Genes Dev

[Internet]. 2008 Sep 1;22(17):2308–41. Available from:

http://genesdev.cshlp.org/cgi/doi/10.1101/gad.1686208

2. Koch U, Lehal R, Radtke F. Stem cells living with a Notch. Development [Internet]. 2013 Feb

15;140(4):689–704. Available from: http://dev.biologists.org/cgi/doi/10.1242/dev.080614

3. Kang HM, Huang S, Reidy K, Han SH, Chinga F, Susztak K. Sox9-Positive Progenitor Cells Play a Key

Role in Renal Tubule Epithelial Regeneration in Mice. Cell Rep [Internet]. 2016 Feb;14(4):861–71.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S2211124715015259

4. Rinkevich Y, Montoro DT, Contreras-Trujillo H, Harari-Steinberg O, Newman AM, Tsai JM, et al. In Vivo

Clonal Analysis Reveals Lineage-Restricted Progenitor Characteristics in Mammalian Kidney

Development, Maintenance, and Regeneration. Cell Rep [Internet]. 2014 May;7(4):1270–83. Available

from: https://linkinghub.elsevier.com/retrieve/pii/S2211124714003052

5. Chen J, Chen J-K, Conway EM, Harris RC. Survivin Mediates Renal Proximal Tubule Recovery from AKI.

J Am Soc Nephrol [Internet]. 2013 Dec;24(12):2023–33. Available from:

https://jasn.asnjournals.org/lookup/doi/10.1681/ASN.2013010076

6. Zhou D, Li Y, Lin L, Zhou L, Igarashi P, Liu Y. Tubule-specific ablation of endogenous β-catenin

aggravates acute kidney injury in mice. Kidney Int [Internet]. 2012 Sep;82(5):537–47. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0085253815555956

7. Lancaster MA, Knoblich JA. Organogenesis in a dish: Modeling development and disease using organoid

technologies. Science (80-) [Internet]. 2014 Jul 18;345(6194):1247125–1247125. Available from:

https://www.sciencemag.org/lookup/doi/10.1126/science.1247125

8. Schutgens F, Clevers H. Human Organoids: Tools for Understanding Biology and Treating Diseases.

Annu Rev Pathol Mech Dis [Internet]. 2020 Jan 24;15(1):211–34. Available from:

https://www.annualreviews.org/doi/10.1146/annurev-pathmechdis-012419-032611

9. Grassi L, Alfonsi R, Francescangeli F, Signore M, De Angelis ML, Addario A, et al. Organoids as a new

model for improving regenerative medicine and cancer personalized therapy in renal diseases. Cell Death

Dis [Internet]. 2019 Mar 27;10(3):201. Available from: http://www.nature.com/articles/s41419-019-1453-0

10. Schutgens F, Rookmaaker MB, Margaritis T, Rios A, Ammerlaan C, Jansen J, et al. Tubuloids derived

from human adult kidney and urine for personalized disease modeling. Nat Biotechnol [Internet]. 2019

Mar 4;37(3):303–13. Available from: http://www.nature.com/articles/s41587-019-0048-8

11. Hsieh JJ, Purdue MP, Signoretti S, Swanton C, Albiges L, Schmidinger M, et al. Renal cell carcinoma.

886   Nat Rev Dis Prim [Internet]. 2017 Dec 21;3(1):17009. Available from:

887   http://www.nature.com/articles/nrdp20179

888   12.   Gossage L, Eisen T, Maher ER. VHL, the story of a tumour suppressor gene. Nat Rev Cancer [Internet].

889   2015 Jan 23;15(1):55–64. Available from: http://www.nature.com/articles/nrc3844

890   13.   Hou W, Ji Z. Generation of autochthonous mouse models of clear cell renal cell carcinoma: mouse

891   models of renal cell carcinoma. Exp Mol Med [Internet]. 2018 Apr 13;50(4):30. Available from:

892   http://www.nature.com/articles/s12276-018-0059-4

893   14.   Fendler A, Bauer D, Busch J, Jung K, Wulf-Goldenberg A, Kunz S, et al. Inhibiting WNT and NOTCH in

894   renal cancer stem cells and the implications for human patients. Nat Commun. 2020 Dec 17;11(1):929.

895   15.   Cole AM, Ridgway RA, Derkits SE, Parry L, Barker N, Clevers H, et al. p21 loss blocks senescence

896   following Apc loss and provokes tumourigenesis in the renal but not the intestinal epithelium. EMBO Mol

897   Med [Internet]. 2010 Nov 26;2(11):472–86. Available from:

898   https://onlinelibrary.wiley.com/doi/10.1002/emmm.201000101

899   16.   Sansom OJ, Griffiths DFR, Reed KR, Winton DJ, Clarke AR. Apc deficiency predisposes to renal

900   carcinoma in the mouse. Oncogene [Internet]. 2005 Dec 22;24(55):8205–10. Available from:

901   http://www.nature.com/articles/1208956

902   17.   Clark PE, Polosukhina D, Love H, Correa H, Coffin C, Perlman EJ, et al. β-Catenin and K-RAS Synergize

903   to Form Primitive Renal Epithelial Tumors with Features of Epithelial Wilms' Tumors. Am J Pathol

904   [Internet]. 2011 Dec;179(6):3045–55. Available from:

905   https://linkinghub.elsevier.com/retrieve/pii/S000294401100808X

906   18.   Sansom OJ, Meniel V, Wilkins JA, Cole AM, Oien KA, Marsh V, et al. Loss of Apc allows phenotypic

907   manifestation of the transforming properties of an endogenous K-ras oncogene in vivo. Proc Natl Acad

908   Sci [Internet]. 2006 Sep 19;103(38):14122–7. Available from:

909   http://www.pnas.org/cgi/doi/10.1073/pnas.0604130103

910   19.   Bhagat TD, Zou Y, Huang S, Park J, Palmer MB, Hu C, et al. Notch Pathway Is Activated via Genetic and

911   Epigenetic Alterations and Is a Therapeutic Target in Clear Cell Renal Cancer. J Biol Chem [Internet].

912   2017 Jan;292(3):837–46. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0021925820402649

913   20.   Johansson E, Rönö B, Johansson M, Lindgren D, Möller C, Axelson H, et al. Simultaneous targeted

914   activation of Notch1 and Vhl-disruption in the kidney proximal epithelial tubular cells in mice. Sci Rep

915   [Internet]. 2016 Aug 5;6(1):30739. Available from: http://www.nature.com/articles/srep30739

916   21.   Zhu L, Finkelstein D, Gao C, Shi L, Wang Y, López-Terrada D, et al. Multi-organ Mapping of Cancer Risk.

917   Cell [Internet]. 2016 Aug;166(5):1132-1146.e7. Available from:

918   https://linkinghub.elsevier.com/retrieve/pii/S0092867416309953

22.   Schönig K, Schwenk F, Rajewsky K, Bujard H. Stringent doxycycline dependent control of CRE

recombinase in vivo. Nucleic Acids Res. 2002;

23.   Harada N. Intestinal polyposis in mice with a dominant stable mutation of the beta -catenin gene. EMBO J

[Internet]. 1999 Nov 1;18(21):5931–42. Available from:

http://emboj.embopress.org/cgi/doi/10.1093/emboj/18.21.5931

24.   Murtaugh LC, Stanger BZ, Kwan KM, Melton DA. Notch signaling controls multiple steps of pancreatic

differentiation. Proc Natl Acad Sci [Internet]. 2003 Dec 9;100(25):14920–5. Available from:

http://www.pnas.org/cgi/doi/10.1073/pnas.2436557100

25.   Soriano P. Generalized lacZ expression with the ROSA26 Cre reporter strain. Nat Genet [Internet]. 1999

Jan;21(1):70–1. Available from: http://www.nature.com/articles/ng0199_70

26.   Traykova-Brauch M, Schönig K, Greiner O, Miloud T, Jauch A, Bode M, et al. An efficient and versatile

system for acute and chronic modulation of renal tubular function in transgenic mice. Nat Med. 2008 Sep

24;14(9):979–84.

27.   Haase VH, Glickman JN, Socolovsky M, Jaenisch R. Vascular tumors in livers with targeted inactivation

of the von Hippel-Lindau tumor suppressor. Proc Natl Acad Sci [Internet]. 2001 Feb 13;98(4):1583–8.

Available from: http://www.pnas.org/cgi/doi/10.1073/pnas.98.4.1583

28.   Holmberg FE, Seidelin JB, Yin X, Mead BE, Tong Z, Li Y, et al. Culturing human intestinal stem cells for

regenerative applications in the treatment of inflammatory bowel disease. EMBO Mol Med [Internet]. 2017

May 10;9(5):558–70. Available from: https://onlinelibrary.wiley.com/doi/abs/10.15252/emmm.201607260

29.   Sato T, Stange DE, Ferrante M, Vries RGJ, van Es JH, van den Brink S, et al. Long-term Expansion of

Epithelial Organoids From Human Colon, Adenoma, Adenocarcinoma, and Barrett's Epithelium.

Gastroenterology [Internet]. 2011 Nov;141(5):1762–72. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0016508511011085

30.   Cato ACB, Nestl A, Mink S. Rapid Actions of Steroid Receptors in Cellular Signaling Pathways. Sci Signal

[Internet]. 2002 Jun 25;2002(138):re9–re9. Available from:

https://stke.sciencemag.org/lookup/doi/10.1126/stke.2002.138.re9

31.   Sorrentino G, Ruggeri N, Zannini A, Ingallina E, Bertolio R, Marotta C, et al. Glucocorticoid receptor

signalling activates YAP in breast cancer. Nat Commun [Internet]. 2017 Apr 19;8(1):14073. Available

from: http://www.nature.com/articles/ncomms14073

32.   Mertins P, Tang LC, Krug K, Clark DJ, Gritsenko MA, Chen L, et al. Reproducible workflow for

multiplexed deep-scale proteome and phosphoproteome analysis of tumor tissues by liquid

chromatography–mass spectrometry. Nat Protoc [Internet]. 2018 Jul 9;13(7):1632–61. Available from:

http://www.nature.com/articles/s41596-018-0006-9

33.   Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass

accuracies and proteome-wide protein quantification. Nat Biotechnol [Internet]. 2008 Dec

30;26(12):1367–72. Available from: http://www.nature.com/articles/nbt.1511

34.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression analyses

for RNA-sequencing and microarray studies. Nucleic Acids Res [Internet]. 2015 Apr 20;43(7):e47–e47.

Available from: http://academic.oup.com/nar/article/43/7/e47/2414268/limma-powers-differential-

expression-analyses-for

35.   Herbst A, Jurinovic V, Krebs S, Thieme SE, Blum H, Göke B, et al. Comprehensive analysis of β-catenin

target genes in colorectal carcinoma cell lines with deregulated Wnt/β-catenin signaling. BMC Genomics

[Internet]. 2014;15(1):74. Available from: http://bmcgenomics.biomedcentral.com/articles/10.1186/1471-

2164-15-74

36.   Zanconato F, Forcato M, Battilana G, Azzolin L, Quaranta E, Bodega B, et al. Genome-wide association

between YAP/TAZ/TEAD and AP-1 at enhancers drives oncogenic growth. Nat Cell Biol [Internet]. 2015

Sep 10;17(9):1218–27. Available from: http://www.nature.com/articles/ncb3216

37.   Rankin EB, Tomaszewski JE, Haase VH. Renal Cyst Development in Mice with Conditional Inactivation of

the von Hippel-Lindau Tumor Suppressor. Cancer Res [Internet]. 2006 Mar 1;66(5):2576–83. Available

from: http://cancerres.aacrjournals.org/lookup/doi/10.1158/0008-5472.CAN-05-3241

38.   Nielsen R, Christensen EI, Birn H. Megalin and cubilin in proximal tubule protein reabsorption: from

experimental models to human disease. Kidney Int [Internet]. 2016 Jan;89(1):58–67. Available from:

https://linkinghub.elsevier.com/retrieve/pii/S0085253815000551

39.   Fang L, Zhu Q, Neuenschwander M, Specker E, Wulf-Goldenberg A, Weis WI, et al. A Small-Molecule

Antagonist of the β-Catenin/TCF4 Interaction Blocks the Self-Renewal of Cancer Stem Cells and

Suppresses Tumorigenesis. Cancer Res [Internet]. 2016 Feb 15;76(4):891–901. Available from:

http://cancerres.aacrjournals.org/lookup/doi/10.1158/0008-5472.CAN-15-1519

40.   Chen J, You H, Li Y, Xu Y, He Q, Harris RC. EGF Receptor–Dependent YAP Activation Is Important for

Renal Recovery from AKI. J Am Soc Nephrol [Internet]. 2018 Sep;29(9):2372–85. Available from:

https://jasn.asnjournals.org/lookup/doi/10.1681/ASN.2017121272

41.   Marx V. A dream of single-cell proteomics. Nat Methods [Internet]. 2019 Sep 12;16(9):809–12. Available

from: http://www.nature.com/articles/s41592-019-0540-6

42.   Ferenbach DA, Bonventre J V. Mechanisms of maladaptive repair after AKI leading to accelerated kidney

ageing and CKD. Nat Rev Nephrol [Internet]. 2015 May 3;11(5):264–76. Available from:

http://www.nature.com/articles/nrneph.2015.3

43.   Koury MJ, Haase VH. Anaemia in kidney disease: harnessing hypoxia responses for therapy. Nat Rev

Nephrol [Internet]. 2015 Jul 9;11(7):394–410. Available from:

http://www.nature.com/articles/nrneph.2015.82

44.   Qian C-N, Knol J, Igarashi P, Lin F, Zylstra U, Teh BT, et al. Cystic Renal Neoplasia Following

Conditional Inactivation of Apc in Mouse Renal Tubular Epithelium. J Biol Chem [Internet]. 2005

Feb;280(5):3938–45. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0021925820762666

45.   Saadi-Kheddouci S, Berrebi D, Romagnolo B, Cluzeaud F, Peuchmaur M, Kahn A, et al. Early

development of polycystic kidney disease in transgenic mice expressing an activated mutant of the β-

catenin gene. Oncogene [Internet]. 2001 Sep 9;20(42):5972–81. Available from:

http://www.nature.com/articles/1204825

46.   Kato H, Gruenwald A, Suh JH, Miner JH, Barisoni-Thomas L, Taketo MM, et al. Wnt/β-Catenin Pathway

in Podocytes Integrates Cell Adhesion, Differentiation, and Survival. J Biol Chem [Internet]. 2011

Jul;286(29):26003–15. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0021925819485876

47.   Wong DWL, Yiu WH, Chan KW, Li Y, Li B, Lok SWY, et al. Activated renal tubular Wnt/β-catenin signaling

triggers renal inflammation during overload proteinuria. Kidney Int [Internet]. 2018 Jun;93(6):1367–83.

Available from: https://linkinghub.elsevier.com/retrieve/pii/S0085253818300838

48.   Bielesz B, Sirin Y, Si H, Niranjan T, Gruenwald A, Ahn S, et al. Epithelial Notch signaling regulates

interstitial fibrosis development in the kidneys of mice and humans. J Clin Invest [Internet]. 2010 Nov

1;120(11):4040–54. Available from: http://www.jci.org/articles/view/43025

49.   Park J, Shrestha R, Qiu C, Kondo A, Huang S, Werth M, et al. Single-cell transcriptomics of the mouse

kidney reveals potential cellular targets of kidney disease. Science (80-) [Internet]. 2018 May

18;360(6390):758–63. Available from: https://www.sciencemag.org/lookup/doi/10.1126/science.aar2131

50.   Edeling M, Ragi G, Huang S, Pavenstädt H, Susztak K. Developmental signalling pathways in renal

fibrosis: the roles of Notch, Wnt and Hedgehog. Nat Rev Nephrol [Internet]. 2016 Jul 3;12(7):426–39.

Available from: http://www.nature.com/articles/nrneph.2016.54

51.   Peired AJ, Antonelli G, Angelotti ML, Allinovi M, Guzzi F, Sisti A, et al. Acute kidney injury promotes

development of papillary renal cell adenoma and carcinoma from renal progenitor cells. Sci Transl Med.

2020;

52.   Chiu S-C, Liu H-H, Chen C-L, Chen P-R, Liu M-C, Lin S-Z, et al. Extramedullary Hematopoiesis (EMH) in

Laboratory Animals: Offering an Insight into Stem Cell Research. Cell Transplant [Internet]. 2015

Mar;24(3):349–66. Available from: http://journals.sagepub.com/doi/full/10.3727/096368915X686850

53.   Carter P, Schnell U, Chaney C, Tong B, Pan X, Ye J, et al. Deletion of Lats1/2 in adult kidney epithelia

leads to renal cell carcinoma. J Clin Invest [Internet]. 2021 Jun 1;131(11). Available from:

https://www.jci.org/articles/view/144108

54.   Harada N, Miyoshi H, Murai N, Oshima H, Tamai Y, Oshima M, et al. Lack of tumorigenesis in the mouse

liver after adenovirus-mediated expression of a dominant stable mutant of β-catenin. Cancer Res. 2002;

55.   Perez-Riverol Y, Csordas A, Bai J, Bernal-Llinares M, Hewapathirana S, Kundu DJ, et al. The PRIDE

database and related tools and resources in 2019: improving support for quantification data. Nucleic

Acids Res [Internet]. 2019 Jan 8;47(D1):D442–50. Available from:

https://academic.oup.com/nar/article/47/D1/D442/5160986

1024

Supporting Information

S1 Figure: Long-term culture of kidney organoids and establishing kidney organoids of different mouse backgrounds.

S2 Figure: Yap signaling is upregulated in kidney organoids.

S3 Figure: Markers of tubular epithelial cells are expressed in 3D reconstructed kidney organoids.

S4 Figure: Kidney organoids display tubular epithelial polarity and complexity.

S5 Figure: No DNA damage, growth arrest or senescence and increased apoptosis was observed in mutant kidneys.

S1 Video: Markers of tubular epithelial cells are expressed in 3D reconstructed kidney organoids.

A

0 1 2 3 4 7 8 9 10 14 days

B

Cystic (65%) | Solid (25%) | Alveolar (10%)

C

Freshly seeded | Passage 1

D

Organoid numbers per well

****

Freshly seeded | Passage 1

E

Serial passage

A Pax8/Haematoxylin

B E-cad/DAPI

C Krt8

Cldn4

Annotation: CK / OEP / OLP

Krt18

mRNA fold change

CK   OEP   OLP

D Zo-1/Laminin/DAPI

E LTL/PNA/DAPI

F Transporters

Protein

Annotation: CK / OEP / OLP

Atp6ap1
Slc1a5
Atp6v0a2
Slc12a2
Atp6v0a1
Abcc3
Clcc1
Slc37a1
Atp2a2
Slc25a24
Atp2c2
Slc35f5
Atp6ap2
Slc2a1
Slc25a17
Slc16a1
Slc7a4
Slc16a12
Slc39a7
Abcc4
Slc12a6
Slc39a4
Abcf3
Trpv4
Slc4a7
Slc27a3
Slc7a6os
Slc5a11
Slc1a4
Abce1
Slc26a2
Slc35b3
Abcf1
Atp8b1
Slc30a6
Slc4a1ap
Abcb1b
Clcn3
Abcc5
Atp13a3
Slc7a11
Slc30a7
Slc7a1
Slc38a2
Slc39a6
Slc39a10
Slc6a9
Abcf2

G Wnk1

Gata3

Annotation: CK / OEP / OLP

H Abcb1b/LTL/DAPI   Abcb1b/PNA/DAPI   Abcb1b/LTL/PNA/DAPI

I Aqp3/LTL/DAPI   Aqp3/PNA/DAPI   Aqp3/LTL/PNA/DAPI

A

Kidney epithelial cells

Pax8 promoter → rtTA

Exogenous locus

rtTA — Dox

P_tet promoter → Cre

LC1 locus

Recombination

Endogenous locus: loxP β-catenin exon 3 loxP

Rosa26 locus: loxP Stop loxP Notch1 icd (N1icd)

Endogenous locus: loxP Vhl promoter-exon 1 loxP

• β-catenin-GOF; Notch-GOF

• Vhl-LOF; β-catenin-GOF

→ Clear cell kidney cancer?

B

Doxycycline 0.2 mg/ml — Day 0

Cre/LacZ expression analysis — Day 5

Pax8-rtTA(+); LC1-Cre(+); LacZ(+)

Ctrl | Cortex | Medulla

Cre

LacZ

C
Ctrl | β-catenin-GOF; Notch-GOF | Ctrl | Vhl-LOF; β-catenin-GOF
1000 bp
500 bp
Wt β-catenin (900 bp)
Ex3 del β-catenin (700 bp)

D
Ctrl | Vhl-LOF; β-catenin-GOF
500 bp
200 bp
Prom-ex1 loxP Vhl (460 bp)
Prom-ex1 del Vhl (260 bp)

E
β-catenin-GOF; Notch-GOF | Vhl-LOF; β-catenin-GOF | Ctrl
Wt β-catenin (95 kDa)
Ex3 del β-catenin (72 kDa)
α-Tubulin (52 kDa)

F
Ctrl | β-catenin-GOF; Notch-GOF
N1icd

G β-catenin-GOF; Notch-GOF
mRNA fold change
Ctrl, Dkk1, Axin2, Cyclin D1, Myc
Wnt-β-catenin target genes

H Vhl-LOF; β-catenin-GOF
mRNA fold change
Ctrl, Dkk1, Axin2, Myc, Cyclin D1
Wnt-β-catenin target genes

I β-catenin-GOF; Notch-GOF
mRNA fold change
Ctrl, Hey1, Hes1, Hey3 (Heyl)
Notch target genes

J Vhl-LOF; β-catenin-GOF
mRNA fold change
Ctrl, Ca9, Hk2, Pdk1, Myc, Ldha, Cyclin D1, Glut1, Vegfa
Hif-1α/2α target genes