CellBin: a highly accurate single-cell gene expression processing pipeline for high-resolution spatial transcriptomics

Mei Li[1,2,†], Huanlin Liu[1,†], Qiang Kang[1,†], Shuangsang Fang[1,3,‡], Min Li[1], Jiajun Zhang[1,4], Fei Teng[1,4], Dan Wang[5,1], Weixuan Cen[1], Zepeng Li[1], Ning Feng[1], Jing Guo[1], Qiqi He[1], Leying Wang[1], Tiantong Zheng[1], Shengkang Li[1,6], Yinqi Bai[1], Min Xie[4,1], Yong Bai[1], Sha Liao[1,4], Ao Chen[1,4], Yong Zhang[1,6,7], Susanne Brix[2], Xun Xu[1,7,*], Yuxiang Li[1,6,7,*]

[1] BGI Research, Shenzhen 518083, China

[2] Department of Biotechnology and Biomedicine, Technical University of Denmark, 2800 Kgs. Lyngby, Denmark

[3] BGI Research, Beijing 102601, China

[4] BGI Research, Chongqing 401329, China

[5] Guangdong Provincial Key Laboratory of Interdisciplinary Research and Application for Data Science, BNU-HKBU United International College, Zhuhai 519087, China

[6] BGI Research, Wuhan 430074, China

[7] Guangdong Bigdata Engineering Technology Research Center for Life Sciences, Shenzhen 518083, China

[†] These authors contributed equally as the first authors.

[‡] Senior author.

[*] e-mail: xuxun@genomics.cn; liyuxiang@genomics.cn

Abstract

Background

Owing to recent advances in resolution and field-of-view, spatially resolved transcriptomics

sequencing, such as Stereo-seq, has emerged as a cutting-edge technology for the interpretation of

large tissues at the single-cell level. To generate accurate single-cell spatial gene expression

profiles from high-resolution spatial omics data, a powerful computational tool is required.

Findings

We present CellBin, an image-facilitated one-stop pipeline for high-resolution and large

field-of-view spatial transcriptomic data of Stereo-seq. CellBin provides a comprehensive and

systematic platform for generating high-confidence single-cell spatial gene expression profiles,

which specifically includes image stitching, image registration, tissue segmentation, nuclei

segmentation and molecule labeling. CellBin is user-friendly and does not require a specific level

of omics and image analysis expertise.

Conclusions

During image stitching and molecule labeling, CellBin delivers better-performing algorithms to

reduce stitching error and time, in addition to improving the signal-to-noise ratio of single-cell

gene expression data, in comparison with existing methods. Additionally, CellBin has been shown

to obtain highly accurate single-cell spatial data using mouse brain tissue, which facilitated

clustering and annotation.

Keywords: spatially resolved transcriptomics, image stitching, cell segmentation, molecule

labeling, gene expression profile generation

Introduction

Spatially resolved technology generates comprehensive data regarding the distribution of molecules that can be used to identify the location and function of cells within tissues, which helps to broaden our understanding of organ development [1], tumor heterogeneity [2] and cancer evolution [3,4]. This rapidly developing field is focused on obtaining detailed molecular information at single-cell resolution, as well as spatial information per molecule of tissue in a large field-of-view [5]. Single-cell resolution technologies [6-8], make it possible to explore spatial omics data at the single-cell, or even subcellular level. When combined with large field-of-view technologies [6], it allows for generation of 3D maps representing biological functions within single cells at the organ level.

Our previous work developed the high-resolution and large field-of-view spatially resolved technology, Stereo-seq, which enabled whole-organ sequencing of most tissues in model organisms (e.g., mouse brain (1cm×1cm) and mouse embryo (1cm×2cm)) [6]. The Stereo-seq technology generates two main outputs: spot-level gene expression data and high-content staining tissue images. The former is based upon detection of unique molecular identifiers captured by each DNA nanoball (DNB) on the sequencing chip while the latter provides detailed image-based (visual) information about the cellular matrix within tissue samples. Solutions that merge the two to obtain accurate spatial gene expression profiles at single-cell level in large field-of view settings will advance spatially resolved technologies and be an important steppingstone for downstream analyses.

Although spatially resolved technologies vary in resolution and field-of-view, one common feature is their ability to produce traditional high-content tissue images using dyes such as

fluorescence, hematoxylin-eosin (H&E) or 4,6-diamidino-2-phenylindole (DAPI) for cellular or

nucleus staining. The generated images provide important information regarding tissue and cell

morphology, however, two main difficulties exist: insufficient precision in image mosaics and

inaccuracy of molecule labeling.

The first difficulty is limited by the small imaging area, mechanical tolerance, and imperfect

calibration of microscopes (e.g., 10×imaging lens, ~1mm×1mm/tile), which makes them sufficient

for visual inspection of tissue images, but not accurate enough for quantitative single-cell

analytical approaches. For example, a 1cm×1cm object generates approximately 10×10 image tiles,

while a 3cm×3cm object generates thousands of image tiles. Existing image stitching methods,

such as ASHLAR [9] and MIST [10], can meet most stitching accuracy requirements. However,

stitching misalignment is a major issue, and it is difficult to collect evaluation datasets to improve

the accuracy of image mosaics in a large field-of-view, which is an obstacle in expanding their

application.

The second difficulty, which is the key to achieving high-precision single-cell resolution, is

classifying the transcripts and other molecules into the owner cell, a procedure termed "molecule

labeling". However, molecule labeling is not an easy task due to the low efficiency of molecule

capture and the problem of diffusion of molecules outside cells during wet-lab procedures, which

greatly affects the correct annotation of transcripts to the corresponding cell. Recent technology

has considered this issue, such as 10X Xenium [11], which includes more transcripts by extending

the distance outside the nucleus (15μm). Since the diameter of immune, stromal, and tumor cells

varies, it is however challenging to obtain accurate single-cell data using such a generalized model

that introduces varying levels of noise depending on the cell type. Some available methods,

including Baysor [12] and Pixel-seq [13], require high-quality data as input, such as the number of

captured molecules and their density distribution, which may not be applicable to most

sequencing-based spatial omics data. Furthermore, existing spatial data analysis frameworks, such

as Seurat [14], RCTD [15], Cell2location [16], Squidpy [17], and Spateo [18], only focus on the

downstream analysis tasks or one specific aspect of the spatial data analysis or consider only cell

nuclei segmentation.

Here we present CellBin, the first one-stop pipeline to generate single-cell spatial gene

expression profiles for whole transcriptome based on high-resolution and large field-of-view

spatially resolved sequencing. For image stitching, CellBin employs multiple Fast Fourier

Transform (FFT) [19] weighted stitching algorithm, named MFWS, to reduce stitching errors in

large-field-of-view datasets. The morphological tissue image stitching procedure of CellBin is

accurate and reliable for single-cell identification and is flexible and convenient in terms of run

time. Its high-precision stitching algorithm is useful for the correction of stitching to almost

subcellular precision. For molecule labeling, CellBin employs the cell morphology and Gaussian

Mixture Model (GMM) [20] algorithm, named MLCG, to increase the signal-to-noise ratio of the

single-cell spatial gene expression profile, which yields more reliable analysis of cell clustering

and annotation. CellBin has been successfully applied to datasets of various organs (such as brain,

heart, embryo, artery, testis, kidney, liver and lymph) from different organisms (such as *Homo*

sapiens, *Mus musculus*, *Macaca mulatta* and *Leporidae*). The molecule labeling of CellBin is also

applicable to the datasets from multiple platforms other than Stereo-seq when the nuclei mask and

spatial gene expression data are given. CellBin's rich documentation in the form of a functional

application programming interface, examples, and tutorial workflows, is easy to navigate and

accessible to both experienced developers and beginner analysts. CellBin has the potential to serve

as a bridge between the fields of image analysis and molecular omics, providing a foundation for

the development of next generation computational methods for spatially resolved technologies.

Results

Overview of CellBin

CellBin is an image-facilitated one-stop pipeline for Stereo-seq transcriptomic data. CellBin

combines large field-of-view tissue images and high-resolution spatial gene expression data to

obtain high-confidence single-cell spatial gene expression profiles. CellBin provides a

straightforward and systematic platform, which includes image stitching, image registration, tissue

segmentation, nuclei segmentation and molecule labeling (Fig. 1a). We use MFWS for image

stitching, which is based on frequency domain information and allows the accurate stitching of

high-resolution images in a wide field-of-view, in addition to improvements in cell dislocation

errors caused by inaccurate image stitching and low efficiency caused by the huge number of

image tiles (Fig. 1b). We transform the spatial gene expression data into a map. The stitched image

is registered with the spatial gene expression map based on "track lines" (marker lines designed on

Stereo-seq chip), which includes translation, scaling, flipping and rotation (Fig. 1c). We train the

deep learning models for tissue segmentation (Fig. 1d) and nuclei segmentation (Fig. 1e) from the

registered image. We enable molecule labeling though MLCG, which applies GMM to fit the

molecules within nuclei mask for accurate assignment of surrounding molecules to the most

probable cell, obtaining a gene expression profile at the single-cell level (Fig. 1f).

We provide a fixed set of hyperparameters for CellBin that are suitable for most Stereo-seq

datasets. This helps prevent users from feeling overwhelmed by the numerous hyperparameters

involved in the all-in-one process. We examine the application of CellBin on various tissue

datasets and compare it with other state-of-the-art methods. We validate the efficiency and

effectiveness of MFWS on Stereo-seq and public datasets, comparing it with ASHLAR [9] and

MIST [10]. The accuracy of the generated single-cell spatial gene expression profile is evaluated

on a mouse olfactory bulb dataset, confirming the efficacy of MLCG. We analyze the cortex

structure in the generated single-cell spatial gene expression profile on a mouse brain dataset,

comparing the results with those obtained using the Bin$X$ method (Bin$X$ means a bin with $X \times X$ of

DNBs [6]). As there is currently no image-based one-stop pipeline like CellBin available for

Stereo-seq data, we compare CellBin with SCS [21], which generates spatial gene expression

profiles based on nuclei segmentation masks but lacks image stitching and registration capabilities,

and Baysor [12], which can generate spatial gene expression profiles without using staining

images. The main experimental results are shown below, and additional experimental results can

be found in Supplementary materials and Supplementary Fig. S1.

CellBin processes high-precision morphological tissue images from image tiles

MFWS of CellBin takes a folder of all image tiles obtained by microscopy and information files as

the input, and outputs a stitched mosaic image of whole tissue. The datasets from different

field-of-views (4 Stereo-seq mouse brain datasets with the chip sizes of 1cm×1cm, 1cm×2cm,

2cm×2cm and 2cm×3cm respectively and a public dataset [22], and their grid sizes (11,9), (15,21),

(25,21), (23,29) and (10,10), respectively) are collected. For each dataset, the standards are

designed (Supplementary materials) to calculate the relative offset error between each two

adjacent image tiles and the absolute offset error of the entire stitched image in the stitching

results of different methods. The relative offset errors are statistically analyzed by Wilcoxon

signed rank test [23] to examine the significant differences. The runtime to obtain the absolute

offset error for each method is recorded.

Processing a morphological image from a tissue slide requires the stitching of an array of

multiple image tiles generated by microscopy. Microscopes can automatically capture the image

tiles of a tissue one by one and stitch the image tiles together using the built-in stitching method.

However, overlapping areas between adjacent image tiles generated during microscope movement

may be imprecise due to mechanical tolerance and imperfect calibration. Such stitching errors are

common in mosaic images and need to be removed if the goal is to achieve single-cell resolution

in spatial omics applications. As an example, a mouse brain mosaic image on a chip with a size of

~2cm×2cm and a tile size of 2424pixels×2031pixels are displayed (Fig. 2a), corresponding to a

physical size of ~1.2mm×1.0mm. To visualize seams and stitching errors, we select two adjacent

image tiles from the image (Fig. 2b). For accurate stitching results, the lower part of tile 1 (dotted

area) and the upper part of tile 2 (dotted area) should be accurately overlaid. Shadows that

represent inaccurate overlap of cells can be easily seen in the stitching results produced by the

microscope (Fig. 2c, left), but MFWS of CellBin is able to stitch the image tiles accurately,

resulting in the absence of shadows (Fig. 2c, right). The stitching results of two image tiles within

non-tissue areas (full-colored part) where individual "track lines" can be clearly seen using

auto-contrast adjustment as also shown (Fig. 2c). The applied Stereo-seq chip contains periodic

"track lines" with a size of 1500 nm (~3 pixels), which is apparent from the image (Fig. 2c, f). We

use a collection of data to evaluate the accuracy and efficiency of existing stitching algorithms as

compared with MFWS.

We apply MFWS to the public dataset, and the results show that the relative offset errors of

MFWS are comparable with those of MIST [10], both being concentrated within 5 pixels, while

ASHLAR [9] has larger offsets (>10-pixels) (Fig. 2d). Moreover, the offset error distribution is

much more concentrated by MFWS. Next, we calculate the relative and absolute offset errors on 4

Stereo-seq mouse brain datasets. MFWS is shown to perform significantly better than ASHLAR [9]

and MIST [10] with respect to the relative offset errors for all image size combinations (Fig. 2e). A

10-pixel (~5μm) dislocation roughly corresponds to half a cell (Fig. 2f). In all the datasets, the

errors generated by MFWS remain within a maximum of 10-pixels in the tissue area, while the

other methods produce a shift greater than 10-pixels, which may misplace the cell (Fig. 2g). As the

number of image tiles increases, the run time becomes a significant factor for stitching algorithms.

The run time of MFWS is significantly shorter than that of the other methods, mostly due to the

embedded spectral calculation based on FFT (Fig. 2h).

CellBin generates highly accurate single-cell gene expression profiles of spatial

omics datasets

Using CellBin, we first obtain a nuclei mask and a cell mask, and then output their single-cell

spatial gene expression profiles. To demonstrate the improvement in transcript assignment to cells

by using the cell mask generated by CellBin, we here compare the gene expression profile output

using each of these masks on Stereo-seq data generated of mouse olfactory bulb, involving

analysis of spatial gene expression data containing 37,288,344 molecules and 143 image tiles. The

generated profiles are input into Stereopy (v6.0) [24] (a downstream analysis tool) for analysis.

The silhouette coefficient [25] is used to evaluate clustering results. The moran's I (calculating by

Scanpy [26] package) is used to evaluate spatial correlation. The details of the data analysis

process and evaluation metrics are provided in Supplementary materials.

After obtaining the nuclei mask, the molecules located inside the nucleus are assigned to the

corresponding cell, and MLCG of CellBin fits the molecular density distribution to each cell

nucleus and re-labels the molecules outside the nucleus to finally generate a cell mask (Fig. 3a).

By applying the cell mask, the uniquely expressed genes and total gene counts in a single cell

increased by approximately 2.34 and 2.56 times, respectively, as compared to the nuclei mask

alone (Fig. 3b). Utilization of the cell mask provides better overall clustering (Fig. 3c), and its

silhouette coefficient is higher than that of the nuclei mask. We manually annotate the cell types

by comparing the differentially expressed genes in each cluster with the marker genes of cell types

inferred by a reference dataset [27]. Of note, there are fewer scattered points and a more

concentrated distribution in the spatial position of the astrocyte layer (Moran's I: 0.40 vs. 0.30)

and dopaminergic neuron layer (Moran's I: 0.52 vs. 0.37) after molecule labeling using the cell

mask (Fig. 3c). We further explore the marker gene distribution and expression in the annotated

cell types, and all the marker genes provided by the reference dataset [27] display higher

expression using the cell mask as compared with the nuclei mask (Fig. 3d). We compare subtypes

of granule cells discovered using the nuclei mask vs. the cell mask. The nuclei mask is able to

identify two subtypes of granule cells: granule cell 0 and granule cell 3, while the cell mask

enables identification of three granule cell subtypes: granule cell 0, granule cell 1, and granule cell

3. We explore the expression of the granule cell 1 subtype-marker genes Syt1, Scg2, and Cplx1,

among all granule cells. There is no obvious difference in expression when using nuclei mask,

while higher expression is found in granule cell 1 as compared with the other two subtypes when

using cell mask (Fig. 3e). This supports that the cell mask captures a higher number of transcript

signals from rare cells, facilitating better annotation and thereby enhances single-cell spatial

resolution.

We estimate and compare the similarity of spatial gene expression between a reference

single-cell dataset of the mouse olfactory bulb [27] and our Stereo-seq based mouse olfactory bulb

dataset using the CellBin nuclei mask vs. the cell mask to generate single-cell expression profiles.

Tissue cells are auto-annotated by Tangram [28] and correlation analysis is performed between the

expression matrixes of single cells using each of the masks and the single-cell sequencing

expression profile of the reference [27]. The Spearman correlation coefficient using annotations

based on the cell mask is higher (0.4% to 8.9% better) than that based on the nuclei mask. Overall,

the single-cell expression matrix generated by the cell mask shows a higher correlation with the

single-cell reference (Fig. 3f), indicating that the cell mask provides a spatial single cell-level gene

expression profile closer to that generated by single-cell sequencing.

CellBin enables dissection of the structural composition of mouse brain cortex

data at single-cell resolution

Non-cell-based binning methods where adjacent tissue regions are divided into regions (bins) of

specified sizes, are sometimes used in spatial transcriptomics analysis pipelines. To compare the

CellBin cell mask output to the Bin$X$ method, we here apply a Stereo-seq mouse brain dataset

containing 131,990,020 molecules and 117 image tiles, where image tiles where split using Bin100,

Bin50, Bin20. The generated profiles are input into Stereopy (v6.0) [24] for analysis, and more

details are provided in Supplementary materials. We export the spatial gene expression map,

nuclei-stained image, cell mask and Bin20 outlines to visually demonstrate the segmentation

effect of the two methods. We also compare the abilities of CellBin and the Bin approach to

reconstruct known cellular regions in the mouse brain.

The clustering of data generated by CellBin vs. Bin100, Bin50, and Bin20 is shown (Fig. 4a).

It appears that splitting using Bin100 results in difficulties in identifying several important areas in

the tissue, such as the brain hub and hippocampus. For Bin50, the tissue cortex and blood cells are

poorly identified. Both Bin20 and CellBin can identify important areas in the tissue, but CellBin

obtains the highest silhouette coefficient in evaluating the clustering results using several methods

(Fig. 4b). We calculate the uniquely expressed genes and total gene counts of CellBin vs. the

differently sized bins (Fig. 4c). The visualization shows that Bin20 more often divides a single cell

into two or more cells, while CellBin more accurately divides the cell area and is consistent with

the actual cell distribution in the tissue (Fig. 4d). Bin20 results in splitting of approximately 90%

of the cells, with only ~10% of the nuclei being completely covered in the nuclei-stained image,

while only ~2% of the cells are split into two or more cells using CellBin (Fig. 4e). The resulting

single-cell data derived from cell identification using Bin20 and CellBin were individually

annotated using Spatial-ID [29] (Fig. 4f) with adolescent mouse brain as a reference [30]. Bin20

annotates 29 different cell types, while CellBin can annotate 37 different cell types. Within the

ACTE series, both Bin20 and CellBin annotate 2 subtypes. In the MEINH series, Bin20 annotates

2 subtypes, while CellBin annotates 3 subtypes. Within the TEGLU series, Bin20 and CellBin

annotate 10 and 13 subtypes, respectively. In TEINH series, Bin20 annotates 5 subtypes and

CellBin annotates 4 subtypes, which is the only case where fewer subtypes were annotated by

CellBin. CellBin also annotates 2 subtypes within the ACNT series and 4 subtypes within the

TEINH series. When zooming in on the cortical area (Fig. 4g), it appears that the staining position

of CellBin and the nucleus are better aligned than seen for Bin20, and the cell state and tissue

structure are more in line with the actual brain tissue map. Moreover, in the gear gyrus and cortical

regions of the mouse brain, CellBin performs better than Bin20 in matching the annotation results

to the Allen mouse brain dataset (Fig. 4h). Although Bin20 is capable of annotating the gear gyrus

and cortex to align with the location in the Allen mouse brain atlas, the division between layers is

not as accurate. Expression of marker genes in the gear gyrus (DGGRC2) and cortex (TEGLU3,

TEGLU4, TEGLU7, and TEGLU8) is fully captured using the CellBin algorithm, which is not the

case when using Bin20 (Fig. 4i). Overall, CellBin provides single-cell spatial gene expression

profiles with a higher signal-to-noise ratio compared to approaches based on differently sized bins.

Moreover, the profile generated using CellBin aligns well with cellular annotation within different

brain regions and can provide a valuable reference for studies of cellular interaction networks in

health and disease.

Discussions

In the present study, we demonstrate that auxiliary staining can be extremely valuable to resolve

difficult cases. The molecule labeling or cell segmentation results will likely be improved using

better staining, such as labeling of outer membranes, and better techniques for segmenting such

images. Additionally, better photo capturing using enhanced AI recognition models will advance

our insight into biological traits and speed up the identification of features that enable us to

distinguish between healthy and diseased tissue. It is noteworthy that the entire CellBin

framework can be applied to generate single-cell gene expression for Stereo-seq, and most of its

modules are also suitable for other spatial transcriptomics technology.

CellBin has quality requirements for the input data. For Stereo-seq transcriptomic data, the

gene expression counts in each Bin200 should not be less than 5000. For the image tiles taken by

the microscope, the "track lines" clarity is required to pass our image quality control. The

registration of CellBin involves aligning the data of two different modalities, i.e. spatial gene

expression data and staining image. To ensure higher accuracy and stronger robustness in the

registration process, the assistance of "track lines" is still indispensable and irreplaceable at

present.

CellBin is equipped with an extensive testing suite, implemented in a robust continuous

integration pipeline that contains a graphic user interface for manual image stitching, image

registration, and molecule labeling. CellBin offers a rich documentation, including a functional

application programming interface, examples, and tutorial workflows, making it easy to navigate

and accessible for diverse developers and analysts. CellBin can be used on desktops, laptops,

workstations, computer clusters or cloud computing platforms. To ensure the normal operation of

CellBin, we recommend a hardware configuration of at least 8-core CPU and 16 GB of RAM. The

use of GPU can greatly accelerate the tissue and nuclei segmentation steps. The amount of disk

storage required depends on the input dataset. The information and timing records on more

different tissue datasets are shown in Supplementary Table S1, which can assist the users in

estimating the required time for their specific situations.

Conclusions

There is still a long way to go before the potential of spatially resolved technology can be fully utilized, and continued improvement of both protocols and analytical methods for processing spatial omics data is required. Here, we present CellBin, an image-facilitated cell segmentation framework for high-resolution and large-field-of-view spatial omics that provides a complete and systematic platform for obtaining high-confidence single-cell spatial data. The implemented morphological tissue image stitching procedure proved to be accurate and reliable for single-cell identification and was flexible and convenient in terms of run time. The high-precision stitching method based on FFT and was found to be useful for the correction of stitching to almost subcellular precision. We implemented molecule labeling based on GMM to generate spatial single-cell gene expression profile with a higher signal-to-noise ratio, which yielded more reliable analysis of cell clustering and end-to-end annotation. Moreover, we optimized the processing modules used for image registration to include "track lines" on the chip, nuclei segmentation, and tissue segmentation.

The goal here is to develop a general pipeline for profile generation and single-cell level analysis of spatial omics data. The data presented herein are based on spatial transcriptomics alone. In future studies, we will combine spatial information with additional biological characteristics to integrate multi-omics information at the single-cell spatial level, improve the signal-to-noise ratio, and develop interfaces and extensions that support user-defined data and methods.

Methods

Image stitching

The image tiles are stitched into a large mosaic image of the whole tissue by MFWS (Fig. 1b). The

file name of each image tile needs to reflect the row and column, such as "0000_0001.tif" that

reflects a tile at row 0 and column 1. Firstly, we calculate the actual overlap value with the

neighbors in the horizontal and vertical directions for each image tile. A pair of image tiles in the

vertical direction is taken as an example. (i) The overlapping regions of adjacent image tiles are

marked as A_f and A_m respectively, and they are divided into N sub-region images as marked $\{f_i\}$ and

$\{m_i\}$ respectively. (ii) The FFT [19] algorithm is applied to obtain $2 \times N$ frequency spectra based on

$\{f_i\}$ and $\{m_i\}$. (iii) The sub-region images in A_f and A_m are paired, such as f_i and m_i, and N

cross-power spectra are obtained using the formula for image cross-power calculation. Theoretically,

any one of these spectra can reflect the true offset information. (iv) Due to the low signal-to-noise

ratio of some sub-regions in the previous step, the calculated value differs greatly from the actual

value. The overall accuracy of the algorithm can be significantly improved by weighted

enhancement of partial blocks and reduction of residual sub-blocks. Our weighting coefficient is

based on the peak value of the mutual power spectrum because experiments show that the greater

the peak value, the higher the precision and reliability of the overlap value. A unique cross-power

spectrum is obtained after the weighted superposition of N spectra. Then, the corresponding spatial

domain correlation graph can be obtained through Inverse FFT. The coordinate of the peak value

represents the required offset value. (v) Carrying out the above steps for each pair of adjacent image

tiles to generate the offset matrices O_H and O_V. Secondly, transformation of local coordinates to

stitching coordinates is required, since image stitching demands the unique coordinates of each tile

in the reference system, i.e., the global coordinate matrix L. In the O_H and O_V matrixes, offsets

corresponding to low-confidence values are not credible that need to be eliminated. However, it

leads to the existence of single or multiple connected domains. Therefore, when obtaining the

coordinates, we first find the center of each connected domain, complete the splicing in the

connected domain according to the relative relationship among the neighbors, and then use the

experience value to fill multiple connected domains to complete the splicing of the entire image.

Thirdly, according to the image tiles and the coordinate matrix L, the size of the mosaic image is

obtained, the value is traversed, and the seam fusion is completed synchronously during the

stitching process.

Image registration

The spatial gene expression data is read and transformed into a map, in which the intensity value

of each pixel is proportional to the total count of molecules of all genes expressed at position DNB.

The stitched image is registered to the transformed map (Fig. 1c). Sequencing chips with sample

tissue attached, imaging, and sequencing are designed with periodic "track lines" (horizontal and

vertical) to assist base calling and image registration. The "track lines" are displayed as dark

straight lines in both the stitched image and transformed map. Image registration is achieved using

the transformed map as a template and performing an affine transformation on the stitched image.

The "track lines" are detected in the transformed map and stitched image by line searching

algorithms, and the intersections of the horizontal and vertical "track lines" are located. The

scaling parameter is calculated by comparing the length of the line segments between intersections

within the transformed map and stitched image. The rotation angle is defined as the difference

between the horizontal "track lines" and the horizontal direction of the image coordinate system.

For the offset between them, since the "track line" patterns are periodic, we first calculate the

center of gravity of the tissue regions within the transformed map and stitched image, roughly

match them together, and then match the "track lines" intersections to fine-tune the image

registration.

Tissue segmentation

The tissue can be segmented from the registered image. The tissue segmentation process consists

of two steps: preprocessing and model inference (Fig. 1d). In addition, a self-scoring mechanism

is added to optimize the processing time and eliminate the effort of manually checking the results.

The histogram enhancement method is used to improve the separation of tissue and background

during the preprocessing, and then, Bi-Directional ConvLSTM U-Net [31] is used to predict tissue

masks during the model inference. Compared with the original U-Net [32], the following

improvements are incorporated. (i) The convolution sequence is replaced with a densely connected

convolution in the final step of the encoding path to encourage feature reuse. (ii) The feature maps

extracted from the corresponding encoding path are combined with the previous decoding

up-convolutional layer in a non-linear manner. (iii) The batch normalization is added to accelerate

the convergence of the network. When dealing with a large amount of unstable data, the model

cannot achieve ideal results with very complex or poor-quality tissue images, and the

self-evaluation mechanism help to filter these potentially unsatisfactory results. When the

registered image possesses a high gray value, we use the threshold value obtained in the histogram

enhancement step as a reference value and assume that the pixel points higher than this value are

valid pixels. The effective pixel densities of the tissue inside the mask and in the area surrounding

the mask are calculated in sections, and the ratio of the two values is used as the basis for

evaluation.

Nuclei segmentation

The nuclei can be determined from the registered image due to nuclei staining. Nuclei

segmentation consists of three steps (Fig. 1e). The first step is image preprocessing. The median

filtering is employed to smooth the noise that may be present in the input image. To both enhance

the cell and homogenize the image background, the output of the median filtering operation is

input into the pixel-wise subtraction process. This step facilitates the relative conspicuity of nuclei

against the background. The second step is segmentation model inference. We use U-Net [32] for

segmenting the cells. Some optimizations are made to U-Net that, Residual U-Net [33] is used as a

feature extractor and the Pyramid Squeeze Attention [34] module is used instead of the 3×3

convolution in the bottleneck as the basic feature-extracting unit of Residual U-Net, which ensures

the model to pay more attention to the highlighted feature representation. The third step is mask

post-processing. To obtain a more reliable segmentation, the mask generated by the segmentation

model is input to the area filtering operation. Further, the opening operation (erode and dilatate) is

applied to revise the cells shape and boundaries.

Molecule labeling

The single-cell spatial gene expression profile can be obtained by combining the nuclei mask and

spatial gene expression data (Fig. 1f). Firstly, the morphological tissue image is used to identify

nuclear boundaries as described in nuclei segmentation. The specific transcripts that are contained

within each nucleus are then assigned. A GMM [20] algorithm is used to estimate the probability

of each transcript belonging to a given cell based on the nuclei mask. It is performed by modeling

each cell as a GMM distribution, combining its spatial position, transcript count, and density.

Simultaneously, we also estimate the probability scores of transcripts in the acellular region. In our

model, the probability of assignment of a transcript to a given cell declines progressively with

distance from the cell center but increases with transcript count and density. A mixture model is a

probabilistic model that can be used to represent a population distribution with K sub-distributions.

GMM can be regarded as a model composed of K single Gaussian models. Taking a single cell as

an example, based on the result of nuclei segmentation, we can locate the spatial position of the

nucleus. Firstly, GMM is used to fit the molecular distribution of the current cells. We expanded

the nuclei mask boundary and captured as much as possible of the true distribution of data in cells

rather than nuclei to develop the cell mask. According to the empirical value of the cell area, we

finally determine the fitting range of 100pixels×100pixels. Subsequently, GMM is used to

calculate the probability score of extracellular molecules within the fitting range. Finally,

according to the maximum and minimum probability scores within the fitting range, the quartile is

used as the adaptive threshold of the current cell. When the probability score is greater than the

threshold, the molecule is re-divided into cells. In this way, molecules are assigned to their

corresponding cell with high confidence.

424   Availability of Source Code and Requirements

Project name: CellBin

Project homepage: https://github.com/STOmics/CellBin/tree/dev

Operating system(s): Windows or Linux

Programming language: Python

Other requirements: See https://github.com/STOmics/CellBin/blob/dev/requirements.txt for

details

License: MIT License

RRID: SCR_025413

BiotoolsID: CellBin

Data availability

435 The data that support the findings of this study have been deposited into Spatial Transcript Omics

436 DataBase (STOmics DB) of China National GeneBank DataBase (CNGBdb) with accession

437 number STT0000027: https://db.cngb.org/stomics/project/STT0000027.

Addition Files

Supplementary materials. Stitching standard design, data analysis process, evaluation metrics,

additional experiments and information of different tissue datasets.

Supplementary Fig. S1. CellBin can be applied to diverse tissue datasets and outperform other

state-of-the-art methods. **a.** The clustering results of single-cell spatial gene expression profiles

obtained by CellBin on diverse tissue datasets. **b.** The result comparison of distance and area

errors obtained by MFWS, MIST and ASHLAR on a public dataset. **c.** The nuclei staining image

of Subset 1 and result comparison of number of cells, number of clusters, silhouette coefficient,

Moran's I and running time obtained by CellBin, SCS and Baysor on Subset 1. **d.** The nuclei

staining image of Subset 2 and result comparison of number of cells, number of clusters,

silhouette coefficient, Moran's I and running time obtained by CellBin, SCS and Baysor on Subset

2.

Supplementary Table S1. Timing values of CellBin on different tissue datasets.

Funding

452　This work was supported by the National Key R&D Program of China (2022YFC3400400).

Authors' contributions

Conceptualization: Mei Li, Huanlin Liu, and Shuangsang Fang.

Project administration and supervision: Susanne Brix, Xun Xu, Yong Zhang, and Yuxiang Li.

Soft development and implementation: Huanlin Liu, and Min Li.

Data collection, processing, and application: Mei Li, Huanlin Liu, Qiang Kang, Jiajun Zhang, Fei

Teng, Weixuan Cen, Zepeng Li, Ning Feng, Jing Guo, Qiqi He, and Leying Wang.

Method comparisons: Huanlin Liu, and Qiang Kang.

Biological interpretation: Qiang Kang, Shuangsang Fang, Jiajun Zhang, and Fei Teng.

Data visualization: Tiantong Zheng and Shengkang Li.

Project coordination: Mei Li, Shuangsang Fang, Sha Liao, and Ao Chen.

Manuscript writing and figure generation: Mei Li, Huanlin Liu, Qiang Kang, Shuangsang Fang,

Jiajun Zhang, Fei Teng, and Dan Wang.

Manuscript review: Susanne Brix, Yinqi Bai, Min Xie, and Yong Bai.

Competing interests

The authors declare they have no competing interests.

Abbreviations

DNB: DNA nanoball; H&E: hematoxylin-eosin; DAPI: 4,6-diamidino-2-phenylindole; FFT: Fast Fourier Transform; MFWS: multiple Fast Fourier Transform weighted stitching algorithm; GMM: Gaussian Mixture Model; MLCG: cell morphology and Gaussian Mixture Model algorithm.

Acknowledgements

This work is part of the "SpatioTemporal Omics Consortium" (STOC) paper package. A list of STOC members is available at http://sto-consortium.org. We would like to thank the MOTIC China Group, in addition to Bichao Chen, Canqiang Xu, Xiangxiang Liao, Kui Su, Zhonghan Deng, Yiwen Wu, Bohan Zhang, Chao Liu, Mengnan Cheng, Qing Zhou, Marija Ivanovic, Vladimir Kovacevic, Lei Han, Xing Guo, and Wei Peng for their help. This work was supported by Wuhan Supercomputing Center for the data training. We thank China National GeneBank for providing technical support. Dan Wang is supported by Zhuhai Basic and Applied Basic Research Foundation (2220004002717).

References

1. Fang S, Chen B, Zhang Y, et al. Computational approaches and challenges in spatial transcriptomics. Genomics Proteomics Bioinformatics 2022;21(1):24-47. https://doi.org/10.1016/j.gpb.2022.10.001.

2. Lu T, Ang CE, Zhuang X. Spatially resolved epigenomic profiling of single cells in complex

tissues. Cell 2022;185:4448-4464. https://doi.org/10.1016/j.cell.2023.04.006.

3. Erickson A, He X, Berglund E, et al. Spatially resolved clonal copy number alterations in

benign and malignant tissue. Nature 2022;608:360-367.

https://doi.org/10.1038/s41586-022-05023-2.

4. Grünwald BT, Devisme A, Andrieux G, et al. Spatially confined sub-tumor microenvironments

in pancreatic cancer. Cell 2021;184:5577-5592. https://doi.org/10.1016/j.cell.2021.09.022.

5. Zhou DC, Jayasinghe RG, Chen S, et al. Spatially restricted drivers and transitional cell

populations cooperate with the microenvironment in untreated and chemo-resistant pancreatic

cancer. Nat Genet 2022;54:1390-1405. https://doi.org/10.1038/s41588-022-01157-1.

6. Chen A, Liao S, Cheng M, et al. Spatiotemporal transcriptomic atlas of mouse organogenesis

using DNA nanoball-patterned arrays. Cell 2022;185:1777-1792.

https://doi.org/10.1016/j.cell.2022.04.003.

7. Chen KH, Boettiger AN, Moffitt JR, et al. Spatially resolved, highly multiplexed RNA profiling

in single cells. Science 2015;348:aaa6090. https://doi.org/10.1126/science.aaa6090.

8. Rodriques SG, Stickels RR, Goeva A, et al. Slide-seq: a scalable technology for measuring

genome-wide expression at high spatial resolution. Science 2019;363:1463-1467.

https://doi.org/10.1126/science.aaw1219.

9. Muhlich JL, Chen YA, Yapp C, et al. Stitching and registering highly multiplexed whole-slide

images of tissues and tumors using ASHLAR. Bioinformatics 2022;38(19):4613-4621.

https://doi.org/10.1093/bioinformatics/btac544.

10. Chalfoun J, Majurski M, Blattner T, et al. MIST: Accurate and scalable microscopy image

stitching tool with stage modeling and error minimization. Sci Rep 2017;7:4988.

https://doi.org/10.1038/s41598-017-04567-y.

11. Janesick A, Shelansky R, Gottscho AD, et al. High resolution mapping of the tumor

microenvironment using integrated single-cell, spatial and in situ analysis. Nat Commun

2023;14:7640. https://doi.org/10.1038/s41467-023-43458-x.

12. Petukhov V, Xu RJ, Soldatov RA, et al. Cell segmentation in imaging-based spatial

transcriptomics. Nat Biotechnol 2022;40:345-354. https://doi.org/10.1038/s41587-021-01044-w.

13. Fu X, Sun L, Dong R, et al. Polony gels enable amplifiable DNA stamping and spatial

transcriptomics of chronic pain. Cell 2022;185:4621-4633.e17.

https://doi.org/10.1016/j.cell.2022.10.021.

14. Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell data.

Cell 2021;184(13):3573-3587. https://doi.org/10.1016/j.cell.2021.04.048.

15. Cable DM, Murray E, Zou LS, et al. Robust decomposition of cell type mixtures in spatial

transcriptomics. Nat Biotechnol 2022;40:517-526. https://doi.org/10.1038/s41587-021-00830-w.

16. Kleshchevnikov V, Shmatko A, Dann E, et al. Cell2location maps fine-grained cell types in

spatial transcriptomics. Nat Biotechnol 2022;40:661-671.

https://doi.org/10.1038/s41587-021-01139-4.

17. Palla G, Spitzer H, Klein M, et al. Squidpy: a scalable framework for spatial omics analysis.

Nat Methods 2022;19:171-178. https://doi.org/10.1038/s41592-021-01358-2.

18. Qiu X, Zhu DY, Yao J, et al. Spateo: multidimensional spatiotemporal modeling of single-cell

spatial transcriptomics. bioRxiv 2022. https://doi.org/10.1101/2022.12.07.519417.

19. Duhamel P, Vetterli M. Fast fourier transforms: a tutorial review and a state of the art. Signal

Process 1990;19:259-299. https://doi.org/10.1016/0165-1684(90)90158-U.

20.   Reynolds   D.   Gaussian   Mixture   Models.   Encycl   Biom   2009;741:659-663.

https://doi.org/10.1007/978-0-387-73003-5_196.

21. Chen H, Li D, Bar-Joseph Z. SCS: cell segmentation for high-resolution spatial

transcriptomics. Nat Methods 2023;20:1237-1243. https://doi.org/10.1038/s41592-023-01939-3.

22. Phase_Image_Tiles. https://isg.nist.gov/BII_2015/Stitching/Phase_Image_Tiles.zip.

23. Taheri SM, Hesamian G. A generalization of the Wilcoxon signed-rank test and its applications.

Stat Papers 2013;54:457-470. https://doi.org/10.1007/s00362-012-0443-4.

24. Fang S, Xu M, Cao L, et al. Stereopy: modeling comparative and spatiotemporal cellular

heterogeneity   via   multi-sample   spatial   transcriptomics.   bioRxiv   2023.

https://doi.org/10.1101/2023.12.04.569485.

25. Rousseeuw PJ. Silhouettes: a graphical aid to the interpretation and validation of cluster

analysis. J Comput Appl Math. 1987;20:53-65. https://doi.org/10.1016/0377-0427(87)90125-7.

26. Wolf FA, Angerer P, Theis FJ. SCANPY: large-scale single-cell gene expression data analysis.

Genome Biol 2018;19:15. https://doi.org/10.1186/s13059-017-1382-0.

27. Brann DH, Tsukahara T, Weinreb C, et al. Non-neuronal expression of SARS-CoV-2 entry

genes in the olfactory system suggests mechanisms underlying COVID-19-associated anosmia.

Sci Adv 2020;6(31):eabc5801. https://doi.org/10.1126/sciadv.abc5801.

28. Biancalani T, Scalia G, Buffoni L, et al. Deep learning and alignment of spatially resolved

single-cell   transcriptomes   with   Tangram.   Nat   Methods   2021;18:1352-1362.

https://doi.org/10.1038/s41592-021-01264-7.

29. Shen R, Liu L, Wu Z, et al. Spatial-ID: a cell typing method for spatially resolved

transcriptomics via transfer learning and spatial embedding. Nat Commun 2022;13:7640.

https://doi.org/10.1038/s41467-022-35288-0.

30. Zeisel A, Hochgerner H, Lönnerberg P, et al. Molecular architecture of the mouse nervous

system. Cell 2018;174:999-1014.e22. https://doi.org/10.1016/j.cell.2018.06.021.

31. Azad R, Asadi-Aghbolaghi M, Fathy M, et al. Bi-Directional ConvLSTM U-Net with densley

connected convolutions. arxiv 2019. https://arxiv.org/abs/1909.00166.

32. Ronneberger O, Fischer P, Brox T. U-Net: convolutional networks for biomedical image

segmentation. In Proceedings of the International Conference on Medical Image Computing and

Computer-Assisted Intervention (MICCAI 2015). Springer;2015;Vol.9351.

https://doi.org/10.1007/978-3-319-24574-4_28.

33. Zhang Z, Liu Q, Wang Y. Road extraction by deep residual U-Net. IEEE Geosci Remote S

2018;15:749-753. https://doi.org/10.1109/LGRS.2018.2802944.

34. Zhang H, Zu K, Lu J, et al. EPSANet: An Efficient Pyramid Squeeze Attention Block on

Convolutional Neural Network. In Proceedings of the 16th Asian Conference on Computer Vision

(ACCV 2022). Springer;2023;Vol.13843. https://doi.org/10.1007/978-3-031-26313-2_33.

Figure legends

Figure 1: The CellBin workflow and applied algorithms. **a.** Schematic overview of the CellBin

pipeline. The spatial gene expression data and the morphological (nuclei-stained) image of tissue

are obtained using spatial transcriptomics technology and microscopy, respectively. The image

tiles obtained by microscopy are stitched together to generate a large mosaic image of the whole

tissue, the spatial gene expression data is transformed to a map, and the stitched image and spatial

gene expression map are registered. Tissue and nuclei segmentations are performed on the

registered image to obtain the tissue mask and nuclei mask. Molecule labeling is adopted to obtain

single-cell spatial gene expression profile. **b.** Cutting the overlap regions of the adjacent image

tiles, and calculating the spectral information based on the Fast Fourier Transform (FFT)

algorithm. Weighted spectral information of the overlap cuts pairs to take the maximum spectral

position as the offset of the adjacent image tiles. The image tiles are stitched using the offset value.

c. Registration with spatial gene expression map and stitched image based on "track lines". The

spatial gene expression map is the fixed image, and the stitched image is the moving one. We

detect the "track lines" on two images, the "track lines" on the spatial gene expression map

represent the template, for the stitched image, we calculate the scale and rotation parameters using

the "track lines". We then use the morphological features to get a rough registration of the offset,

flip, and 90° rotation. "Track lines" are used to fine-tune registration. **d.** Bi-Directional

ConvLSTM U-Net is used to obtain the tissue mask on registered image. **e.** U-Net is used to

obtain the nuclei mask on registered image. **f.** Molecules within the nuclear boundary are assigned

to a given cell, while molecules outside the nucleus are labeled to the cell that gains the largest

probability from Gaussian Mixture Model fitting.

Figure 2: CellBin includes a high-precision mosaic method for image stitching, enabling large

field-of-view morphological image analysis at single-cell spatial accuracy. **a.** Mosaic image of

mouse brain tissue on a 2cm×2cm chip. **b.** The red box indicates the edge of the tissue (containing

cells and non-tissue), and two neighboring image tiles (in the vertical direction) are shown in

different colors. The part that needs to be overlaid by stitching is the lower part of tile 1 (within

the dotted area) and the upper part of tile 2 (within the dotted area). **c.** The stitching results for the

two image tiles (yellow box in **b**) using the stitching method built into the microscope (left) and

MFWS (right). In each sub-figure, the left part shows the overlap of cells after stitching and the

right part shows the "track lines" in the non-tissue area. The contrast of the background was

increased to clearly show the "track lines" incorporated into the Stereo-seq chip. **d.** Comparison of

the relative errors produced by MIST, ASHLAR, and MFWS on a public dataset. **e.** Comparison

of the relative errors produced by MIST, ASHLAR, and MFWS on Stereo-seq mouse brain

datasets, analyzed using different chip sizes from 1cm×1cm to 2cm×3cm, corresponding to

number of image tiles from 11×9 to 23×29. The evaluation is based on the ground truth calculated

using the "track lines" on the Stereo-seq chip. **f.** An example of the stitching error in pixels in the

zoomed-in image. A dislocation of 10-pixels roughly corresponds to half a cell. **g.** Bar graph

illustrating the maximal accumulation of stitching errors produced by MIST, ASHLAR, and

MFWS on the datasets from **e**. **h.** Line graph displaying the run time of MIST, ASHLAR, and

MFWS on Stereo-seq mouse brain datasets.

Figure 3: CellBin provides single-cell spatial data with a higher signal-to-noise ratio that

facilitates finer cell clustering and annotation of the mouse olfactory bulb. **a.** Flowchart of MLCG

processing of Stereo-seq mouse olfactory bulb dataset. **b.** Comparison of the number of uniquely

expressed genes and the total gene count per cell using nuclei mask vs. cell mask on Stereo-seq

mouse olfactory bulb dataset. Top: density plot of the total gene counts per cell. Bottom: density

plot of the number of uniquely expressed genes per cell. **c.** Comparison of the clustering results

generated using the "Leiden" algorithm on nuclei mask vs. cell mask. Top row, left: spatial

clustering results of mouse olfactory bulb data generated using nuclei mask vs. cell mask; right:

comparison of unique cells estimated by Silhouette coefficient. Middle and bottom rows, left:

spatial distribution of cells in the inferred clusters of astrocytes and dopaminergic neurons

generated using nuclei mask vs. cell mask; right: comparison of the spatial autocorrelation

estimated by Moran's I. **d**. Comparison of marker gene expression in the main cellular clusters

inferred by the "Leiden" algorithm generated using nuclei mask vs. cell mask. **e.** Comparison of

cellular subtype identification using nuclei mask vs. cell mask. First column: subtypes of granule

cells identified using nuclei mask vs. cell mask. Other columns: gene expression heat maps of

Syt1, Scg2, and Cplx1 respectively, which have been reported to be marker genes for granule cell

1, using nuclei mask vs. cell mask. **f**. Spearman correlation between gene expression in similar

cells in tissue slides of the mouse olfactory bulb using nuclei mask vs. cell mask to define the gene

expression in single cells and a single-cell reference dataset of the mouse olfactory bulb. **c**, **d**, **f**:

AONM/T cell: anterior olfactory nucleus mitral/tufted cell, VIPP neuron: vasoactive intestinal

peptide positive neuron.

Figure 4: Clustering and annotation of mouse brain tissue by CellBin vs. differently sized bins. **a.**

Identification of gene expression clusters using differently sized bins vs. CellBin on Stereo-seq

mouse brain dataset. **b.** Silhouette coefficient evaluation of clustering performance. **c.** Violin plots

displaying the total gene counts and number of uniquely expressed genes per cell captured using

differently sized bins vs. CellBin. **d.** Comparison of the cell boundary generated using Bin20 and

CellBin. The first image shows the spatial gene expression map and nuclei-stained image, which

are merged in green and blue, respectively. The second image displays an enlarged map of the

local area from the first image. The third image is a comparison of the cell boundaries obtained

using Bin20 (red) vs. CellBin (green). **e.** Comparison of the proportion of cell nuclei covered or

intersected using Bin20 vs. CellBin.   **f.** Comparison of end-to-end annotation results of Bin20 vs.

CellBin. Colors represent the same cell type in the Bin20 and CellBin annotation charts. **g.** The

enlarged local comparison map of the cortex shows the spatial distribution and boundary

information of different cortical cells. **h.** Comparison of cell annotation based on Bin20 vs.

CellBin using the Allen mouse brain atlas as a reference. The spatial distribution of the gear gyrus

(DGGRC2, black) and the cortex (TEGLU3, TEGLU4, TEGLU7, and TEGLU8) is shown. **i.** Heat

map displaying the consistency of marker gene expression within the different substructures in **h**

(gear gyrus (DGGRC2) and cortex (TEGLU3, TEGLU4, TEGLU7, and TEGLU8) layers)). The

upper part: results of Bin20, lower part: results of CellBin.

Fig. 1

a CellBin pipeline

Tissue — Microscope

Sequenator — Tiles

Stereo-seq analysis workflow — **Image stitching**

Spatial gene expression map — Stitched image

Image registration

Registered image

Tissue segmentation

Tissue mask

Nuclei segmentation

Nuclei mask

Molecule labeling

Single-cell spatial gene expression profile

b Image stitching

Overlap Region

Tile1 — Tile2

0.2 mm — 0.2 mm

MFWS

FFT

Crop and weight

0.2 mm

Stitched image

d Tissue segmentation

Registered image

Bi-Directional ConvLSTM U-Net

Tissue mask

c Image registration

Moving image — Fixed image

Detect marker — Detect marker

Stained image — Gene expression map

rotation & resize

Confirm chip placement direction

Align by offset

best position — output

e Nuclei segmentation

Registered image

U-Net

Nuclei mask

f Molecule labeling

Gene expression data

Nuclei mask

Nuclei mask

Cell mask

MLCG

Nuclei genes density fitting

GMM

Cell mask

Fig. 2

a Stained image

b Two tiles

Tile1 — 0.05 mm
Tile2 — 0.05 mm

c Microscope — 0.05 mm / MFWS — 0.05 mm

d Public dataset

e 1cm×1cm   1cm×2cm   2cm×2cm   2cm×3cm

f Stitching example — 10pix — 0.05 mm

g Maximal accumulation error (pix)

h Runtime comparison of different methods (s)

Fig. 3

Fig. 4