1 **Title**
2
3 Meta-Research: understudied genes are lost in a leaky pipeline between genome-wide assays
4 and reporting of results
5

6 **Authors**
7
8 Reese AK Richardson[1,2], Heliodoro Tejedor Navarro[2,3], Luis A Nunes Amaral[2,3,4,5]*, Thomas
9 Stoeger[2,6,7]*
10

[1]Interdisciplinary Biological Sciences, Northwestern University

[2]Department of Chemical and Biological Engineering, Northwestern University

[3]Northwestern Institute on Complex Systems, Northwestern University

[4]Department of Physics and Astronomy, Northwestern University

[5]Department of Molecular Biosciences, Northwestern University

[6]The Potocsnak Longevity Institute, Northwestern University

[7]Simpson Querrey Lung Institute for Translational Science, Northwestern University

*Correspondence to:

L.A.N.A. (amaral@northwestern.edu), T.S. (thomas.stoeger@northwestern.edu)

23
24

25 **Abstract**
26
Present-day publications on human genes primarily feature genes that already appeared in
many publications prior to completion of the Human Genome Project in 2003. These patterns
persist despite the subsequent adoption of high-throughput technologies, which routinely
identify novel genes associated with biological processes and disease. Although several
hypotheses for bias in the selection of genes as research targets have been proposed, their
explanatory powers have not yet been compared. Our analysis suggests that understudied
genes are systematically abandoned in favor of better-studied genes between the completion
of -omics experiments and the reporting of results. Understudied genes remain abandoned by
studies that cite these -omics experiments. Conversely, we find that publications on
understudied genes may even accrue a greater number of citations. Among 45 biological and
experimental factors previously proposed to affect which genes are being studied, we find that

33 are significantly associated with the choice of hit genes presented in titles and abstracts of -
omics studies. To promote the investigation of understudied genes we condense our insights
into a tool, *find my understudied genes* (FMUG), that allows scientists to engage with potential
bias during the selection of hits. We demonstrate the utility of FMUG through the identification
of genes that remain understudied in vertebrate aging. FMUG is developed in Flutter and is
available for download at fmug.amaral.northwestern.edu as a MacOS/Windows app.
44
45 **Introduction**
46
Research into human genes concentrates on a subset of genes that were already frequently
investigated prior to the completion of the Human Genome Project in 2003[1-5]. This
concentration stems from historically acquired research patterns rather than present-day
experimental possibilities[6,7]. For most human diseases, these patterns lead to little correlation
between the volume of literature published on individual genes and the strength of supporting
evidence from genome-wide approaches[8-13]. For instance, we found that 44% of the genes
identified as promising Alzheimer's disease targets by the U.S. National Institutes of Health
(NIH) Accelerating Medicine Partnership for Alzheimer's Disease (AMP-AD) initiative have never
appeared in the title or abstract of any publication on Alzheimer's disease[13]. Furthermore,
when comparing gene-disease pairs, there is no correlation between the ranks of support by
transcriptomics and occurrence in annotation databases[9].
58
Although -omics technologies can provide insights on numerous genes across the genome at a
time and thus offer the promise to counter historically acquired research patterns[14-17], this
discrepancy has persisted[9,18-22] even as the popularity of -omics technologies has risen[5,23,24]. We
therefore sought to use bibliometric data to delineate where and why understudied human
protein-coding genes are abandoned as research targets following -omics experiments. In the
absence of any prior quantitative testing of existing hypotheses, it remains unclear whether
policies to promote the exploration of a greater set of disease-related genes should focus on
how experiments are conducted, how results are reported, or how these results are
subsequently received by other scientists.
68
69 **Data**
70
We considered 450 genome-wide association studies (GWAS, from studies indexed by the
NHGRI-EBI GWAS catalog[25]), 296 studies using affinity purification–mass spectrometry (AP-MS,
indexed by BioGRID[26]), 148 transcriptomic studies (indexed by the EBI Gene Expression Atlas,
EBI-GXA[27]), and 15 genome-wide screens using CRISPR (indexed by BioGRID Open Repository of
CRISPR Screens, BioGRID ORCS[26]) (see PRISMA diagrams in **Figures S1-S4**). We denote genes
that are found to have statistically significant changes in expression or associations with a
phenotype as 'hit' genes.
78
As a surrogate for a given gene having been investigated closer, we consider whether it was
reported in the title or abstract of a research article. We determined which genes were
mentioned in the title or abstract of articles using annotations from gene2pubmed[28] and

PubTator[29]. We used NIH iCite v32 for citations[30]. For determining which gene properties were
associated with selection as research targets, we synthesized quantitative measures from a
variety of authoritative sources (see **Methods**).
85
86   **Results**
87
88   *Understudied genes are abandoned at synthesis/writing stage*
89
We sought to identify at which point in the scientific process understudied genes are ignored as
research targets in investigations using -omics experiments (**Figure 1A**). To receive scholarly
attention, a gene must travel through a pipeline from biological reality to experimental results
to write-up of those results. These results must be extended by subsequent research by other
scholars. Understudied genes do not progress all the way through the pipeline, but it is unclear
where this leak primarily occurs. The first possibility is that some genes are less studied because
they are rarely identified as hits in experiments. Prior studies have, however, shown that
understudied genes are frequent hits in high-throughput experiments[8,9,31], suggesting that this
is not the case. The second possibility is that understudied genes are frequently found as hits in
high-throughput experiments but are not investigated further by the authors. The final
possibility is that subsequent studies do not continue work on understudied genes revealed by
the initial study.
102
103 Evaluating the first possibility, we found that understudied genes were frequently found as hits
104 in high-throughput experiments (**Figure 1B**). This demonstrates, in line with earlier studies[8-13],
105 that the lack of publications on some genes is not explained by a lack of underlying biological
106 experimental evidence.
107
Evaluating the second possibility, we found that hit genes that are highlighted in the title or
abstract are over-represented among the 20% highest-studied genes in all biomedical literature
(**Figure 1B**). These trends are independent of significance threshold (**Figure S5**) and (except for
CRISPR screens) whether we considered the current scientific literature or literature published
before 2003, before any of these articles had been published (**Figure S6**). We also find that this
effect holds when controlling for the number of genes in each title/abstract by only considering
one randomly-chosen gene per title/abstract (**Figure S7**).
115
Understudied genes are least frequently highlighted in the title/abstract in transcriptomics
experiments and most frequently highlighted in the title/abstract in CRISPR screens. GWAS
studies tend to return better-studied genes as hits; the median hit gene in GWAS studies was
more studied in the biomedical literature than 75% of genes. Hit median gene highlighted in the
title/abstract in GWAS studies was more studied in the biomedical literature than 85% of all
protein coding genes. This may explain the prior observation that the total number of articles
on individual genes partially correlates with the total number of occurrences as a hit in GWAS
studies[32].
124

125 Evaluating the final possibility, we found that the reception of -omics studies in later scientific
126 literature either reproduced authors' initial selection of highly studied genes or slightly
127 mitigated it. Jointly, the above findings reinforce that understudied genes become abandoned
128 between the completion of -omics experiments and the reporting of results, rather than being
129 abandoned by later research.
130
131 *Subsequent reception by other scientists does not penalize studies on understudied genes*
132
133 The abandonment of understudied genes could be driven by the valid concern of biomedical
134 researchers that focusing on less-investigated genes will yield articles with lower impact[17], as
135 observed around the turn of the millenium[33]. If this were the case, preemptively avoiding
136 understudied hits would be the rational decision for authors of -omics studies.
137
We thus decided to complement our preceding analysis by an analysis explicitly focused on
citation impact. Notably, we found that the concern of publications on understudied genes
receiving fewer citations does not hold for present-day research on human genes; in biomedical
literature at-large, articles focusing on less-investigated genes typically accumulate more
citations, an effect that has held consistently since 2001 (**Figure 2**). Further, since 1990, articles
about the least popular genes have at times been 3 to 4 times more likely to be among the
most cited articles than articles on the most popular genes whereas articles on the most
popular genes have been slightly less to be highly cited than lowly cited (**Figure S8**).
146
To rule out that these macroscopic observations stem from us having aggregated over different
diseases, we separately analyzed 602 disease-related MeSH terms (**Figure S9**). We found 29
MeSH terms with a statistically significant Spearman correlation using Benjamini-Hochberg FDR
< 0.01 (**Table S4**), of which 27 showed a negative association and only 2 a positive association.
This result again opposes the hypothesis that less-investigated genes will yield articles with
lower impact.
153
Returning to our observation that understudied hits from high-throughput assays are not
promoted to the title and abstract of the resulting publication, we next tested if different
experimental approaches demonstrated distinct associations between gene popularity and
citations (**Figure S10**). Among 264 technique-related MeSH terms tested, there were 20 MeSH
terms with a statistically significant Spearman correlation using Benjamini-Hochberg FDR < 0.01
(**Table S5**), of which 16 showed a negative association and only 4 a positive association.
Notably, MeSH terms representing high-throughput techniques (e.g. D055106:Genome-Wide
Association Study and D020869:Gene Expression Profiling) showed no significant association.
This finding suggests that authors of high-throughput studies have little to gain or lose citation-
wise by highlighting understudied genes.
164
To summarize, our investigations are reminiscent of the previously described separation
between "large-scale" and "small-scale" biological research[34-36]. Authors of high-throughput
studies do not highlight understudied genes in the title or abstract of their publications, the

sections of the publication most accessible to other scientists. While, overall, understudied
genes (and high-throughput assays themselves[5]) correlate with increased citation impact, for
high-throughput studies any potential gain in citations is either absent or too small to be
significant. Thus, there may not be any incentive for authors of high-throughput studies to
highlight understudied genes.

Identification of biological and experimental factors associated with selection of highlighted
genes

To illuminate why understudied genes are abandoned between experimental results and the
write-up of results, we performed a literature review to identify factors that have been
proposed to limit studies of understudied genes (**Table S1**). These factors range from
evolutionary factors (e.g., whether a gene only has homologs in primates), to chemical factors
(e.g., gene length or hydrophobicity of protein product), to historical factors (e.g., whether a
gene's sequence has previously been patented) to materialistic factors affecting experimental
design (e.g., whether designed antibodies are robust for immunohistochemistry).

As any of these factors could plausibly affect gene selection within individual domains of
biomedical research, we returned to the -omics data described above (**Figure 1B**) and measured
how much these factors align with the selective highlighting of hit genes in the title or abstract
of GWAS, AP-MS, transcriptomics, and CRISPR studies.

We identified 45 factors that relate to genes and found 33 (12 out of 23 binary factors and 21
out of 22 continuous factors) associated with selection in at least one assay type at Benjamini-
Hochberg FDR < 0.001 (**Figure 3, Table S2,** and **Table S3**). Across the four assay types, the most
informative binary factor describes whether there is a plasmid available for a gene in the
AddGene plasmid catalog. We cautiously hypothesize that this might reflect on many different
research groups producing reagents surrounding the genes that they actively study. The most
informative continuous factor is the number of research articles about a gene (**Figure 1B**).

To better understand how all 45 factors are related, we performed a cluster analysis of the
collected factors (**Figure S11** and **Figure S12**). This clustering suggests that many factors
influencing the abandonment of understudied genes are not independent. For instance, we find
that the number of articles about a gene is heavily correlated with the number of annotations
for that gene in all surveyed databases. In another case, gene length is heavily correlated with
the number of GWAS annotations for a gene, as described before in terms of transcript length
and single-nucleotide polymoprhisms[37].

Data-driven design of a tool to promote the investigation of understudied genes

To promote the investigation of understudied genes, we combined all the above insights to
create a tool we denoted *find my understudied genes* (FMUG). Our literature review revealed
several tools and resources aiming to promote research of understudied genes by publicizing
understudied genes[38-45] or by providing information about hit genes[7,46-50]. However, we noted

the absence of tools enabling scientists to actively engage with factors that align with gene
selection. Although such factors are largely correlated when considering all genes (**Figure S11**
and **Figure S12**), some factors cluster together and the influence of specific factors could vary
across laboratories. For instance, scientists could vary in their ability to perform proteomics, or
ability to explore orthologous genes in *C. elegans*, or ability to leverage human population data,
or perform standardized mouse assays.
218
Our tool makes selection bias explicit, while acknowledging that different laboratories vary in
their techniques and capabilities for follow-up research. Rather than telling scientists about the
existence of biases, FMUG aims to prompt scientists to make bias-aware informed decisions to
identify and potentially tackle important gaps in knowledge that they are well-suited to
address. For this reason, we believe that FMUG will not be of value only to scientists engaging
in high-throughput studies, but also by scientists wishing to mine existing datasets for hit genes
that they would be well-positioned to investigate further.
226
227 FMUG takes a list of genes from the user (ostensibly a hit list from a high-throughput -omics
228 experiment) and provides the kind of information that will allow a user to select genes for
229 further study.
230
Users can employ filters that reflect the factors identified in our literature review and
supported by our analysis. The default information provided to users consists of factors that are
representative of the identified clusters (**Figure S11**) and strongly associated with gene
selection in high-throughput experiments (**Figure 3, Table S2,** and **Table S3**). In extended
options, users can select any factor that demonstrated a significant association with the
selection of genes. For instance, a user may need to decide whether loss-of-function intolerant
genes should be considered for further research or not, or whether there should be robust
evidence that a gene is protein-coding. Some of these filters are context aware. For instance, a
user may select genes that have already been studied in the general biomedical literature but
not yet within the literature of their disease of interest.
241
To provide real-time feedback, users are, in parallel, presented the number of articles about
genes in their initial input list and the number of articles about genes that passed their filters.
Users can then export their filtered list of genes. In the interest of researcher privacy, FMUG
keeps all information local to the user's machine. Usage of FMUG is illustrated in **Figure 4A** and
demonstrated in **Movie S1**. FMUG is developed in Flutter and is available for download at
fmug.amaral.northwestern.edu as a MacOS/Windows app. For the development of custom
software and analytical code, we provide the data underlying FMUG at
github.com/amarallab/fmug_analysis.
250
To determine the practical usefulness of FMUG to scientists we used an early prototype of
FMUG to identify understudied genes associated with aging. One of these genes was Splicing
factor, proline- and glutamine-rich (*Sfpq*), which had not yet been investigated toward its role in
biological aging. We found *Sfpq* to be transcriptionally downregulated during murine aging.
Others had shown *Sfpq* to be required for the transcriptional elongation of long genes[51]. This

led us to hypothesize that during vertebrate aging, the transcripts of long genes become
downregulated in most tissues **(Figure 4B).** We found this hypothesis to be supported through a
multi-species analysis which we published in December 2022 in Nature Aging[52], with another
group publishing so in January 2023 in Nature Genetics[53], and a third group in iScience in March
2023[54].
261
262
263   **Study Limitations**
264
Our study has several limitations. First, all analysis is subject to annotation errors in the various
databases we employ. While these should be rare and not affect our overall findings, they may
affect users who are interested in genes with discordant annotations. Second, we focus only on
human genes. Different patterns of selection may exist for research on genes in other
organisms. Third, we take a gene being mentioned in the title or abstract of an article as a proxy
for a gene receiving attention by the article's authors. The title and abstract are space-limited
and thus cannot accommodate discussion of large numbers of genes.
272
Fourth, our literature review also identified further factors that we could not test more directly
because of absent access to fitting data. These are: experts' tendency to deepen their
expertise[3], a perceived lack of accuracy of -omics studies[17,55], -omics serving research purposes
beyond target gene identification[22], the absence of good protocols for mass spectrometry[56], the
electronic distribution and reading of research articles[57], rates of reproducibility[17], career
prospects of investigators[7,58], authors beginning manuscripts with something familiar before
introducing something new[59], and the human tendency to fall back to simplifying heuristics
when making decisions under conditions with uncertainty[60]. Fifth, we cannot resolve further
which specific step between the conduct of an experiment and the writing of a research article
leads to the abandonment of hit genes. Finally, we interpret the results of high-throughput
experiments based on their representation in the NHGRI-EBI GWAS, BioGRID, EBI-GXA and
BioGRID ORCS databases. The authors of the original studies may have processed their data
differently, obtaining different results.
286
287
288
289   **Discussion**
290
Efforts to address the gaps in detailed knowledge about most genes have crystallized as
initiatives promoting the investigation of understudied sets of genes[16,61-65], an approach to gene
scholarship recently termed 'unknomics'[66]. The insight that understudied genes are lost to titles
and abstracts of research articles in a leaky pipeline between genome-wide assays and
reporting of results, and FMUG, have already been useful in guiding our own unknomics
research **(Figure 4B).**
297
As our present analysis is correlative, it also is tempting to propose controlled trials where
published manuscripts on high-throughput studies randomly report hit genes in the abstract

even if not investigated further by the authors. This intervention would need to be carefully
designed since abstracts are limited in their size. Further, the observed discrepancy between
the popularity of hits highlighted by GWAS versus other technologies suggests that some -omics
technologies may be more powerful than others for characterizing understudied genes. This
possibility merits further research and researchers participating in unknomics should consider
the relative strengths of each technology towards providing tractable results for follow-up.
306
307 We believe that enabling scientists to consciously engage with bias in research target selection
308 will enable more biomedical researchers to participate in unknomics, to the potential benefit of
309 their own research impact and towards the advancement of our collective understanding of the
310 entire human genome.
311
312 **Materials and Methods**
313
314 *Genes information*
315
316 Homo sapiens gene information was downloaded from NCBI Gene on Aug 16, 2022
317 [ftp.ncbi.nlm.nih.gov/gene/DATA/GENE_INFO/All_Data.gene_info.gz]. Only genes with an
318 unambiguous mapping of Entrez ID to Ensembl ID were used (n = 36,035). Number of gene
319 synonyms, protein-coding status, and official gene symbol were derived from this dataset. A
320 gene symbol was considered undefined if the gene's entry for HGNC gene symbol was "-".
321
322 *Genes in title/abstract of primary research articles*
323
Homo sapiens gene information was downloaded from NCBI Gene on Aug 16, 2022
[ftp.ncbi.nlm.nih.gov/gene/DATA/GENE_INFO/Mammalia/Homo_sapiens.gene_info.gz].
gene2pubmed was download from NCBI Gene on Aug 16, 2022
[ftp.ncbi.nlm.nih.gov/gene/DATA/gene2pubmed.gz][28]. PubTator gene annotations were
downloaded from NIH-NLM on July 12, 2022
[https://ftp.ncbi.nlm.nih.gov/pub/lu/PubTatorCentral/][28,67]. PubMed was downloaded on Dec
17, 2021 [https://ftp.ncbi.nlm.nih.gov/pubmed/baseline/].
331
332 Only using PMIDs annotated as primary research articles, a human gene was considered as
333 mentioned in the title/abstract of the publication if gene was annotated as being in the
334 title/abstract by PubTator and the article appeared in gene2pubmed.
335
336 *CRISPR articles*
337
BioGRID ORCS[26] v1.1.6 was downloaded on April 25, 2022
[https://downloads.thebiogrid.org/BioGRID-ORCS/Release-Archive/BIOGRID-ORCS-1.1.6/]. Any
genome-wide CRISPR knockout screens in human with an associated PubMed ID in which hit
genes were mentioned in the title or abstract was considered (n = 15). 9,268 unique genes were
found as hits. Of these, 18 (0.19%) were highlighted in titles/abstracts in the reporting articles

and 19 (0.21%) were highlighted in titles/abstracts in citing articles. A full list of PubMed IDs is
available in **Table S6**.
345
346 *Transcriptomics articles*
347
EBI-GXA[27] release 36 was downloaded on Sep 15, 2020
[https://web.archive.org/web/20201022184159/https://www.ebi.ac.uk/gxa/download]. This is
the most recent release of EBI-GXA available as a bulk download. Any transcriptomics
comparisons with an associated PubMed ID in which hit genes were mentioned in the title or
abstract was considered (n= 148). Analysis was restricted to protein-coding genes (some
screens featured non-protein-coding genes, but this was not common to all analyses). DE was
called at Benjamini-Hochberg FDR q < 0.05. 18,295 unique genes were found as hits. Of these,
161 (0.88%) were highlighted in titles/abstracts in the reporting articles and 692 (3.78%) were
highlighted in titles/abstracts in citing articles. A full list of PubMed IDs is available in **Table S6**.
357
358 *Affinity purification–mass spectrometry articles*
359
BioGRID[26] v3.5.186 was downloaded on April 25, 2022
[https://downloads.thebiogrid.org/BioGRID/Release-Archive/BIOGRID-3.5.186/]. Any
interactions involving a human gene as the prey protein with an experimental evidence code of
'Affinity Capture-MS' labeled as 'High-Throughput' that had an associated PubMed ID in which
hit genes were mentioned in the title or abstract was considered (n= 296). Prey proteins in
these interactions were considered hits. 7,919 unique genes were found as hits. Of these, 311
(3.93%) were highlighted in titles/abstracts in reporting articles and 407 (5.14%) were
highlighted in titles/abstracts in citing articles. A full list of PubMed IDs is available in **Table S6**.
368
369 *GWAS articles*
370
The NHGRI-EBI GWAS catalog[25] (associations and studies) was download on Aug 17, 2022
[https://www.ebi.ac.uk/gwas/docs/file-downloads]. Any GWAS screens with an associated
PubMed ID in which hit genes were mentioned in the title or abstract was considered (n= 450).
Only SNPs occurring within a gene were considered hits. 1,043 unique genes were found as hits.
Of these, 413 (39.6%) were highlighted in titles/abstracts in reporting articles and 319 (30.6%)
were highlighted in titles/abstracts in citing articles. A full list of PubMed IDs is available in
Table S6
378
379 *Citing articles*
380
381 NIH iCite v32 was downloaded on Aug 25, 2022[30]
382 [https://nih.figshare.com/collections/iCite_Database_Snapshots_NIH_Open_Citation_Collectio
383 n_/4586573/32].
384
385 *Functional annotations*
386

387 Mapping of genes to Gene Ontology / Protein Interaction Database / WikiPathways / Reactome
388 / Kyoto Encyclopedia of Genes and Genomes / Human Phenotype Ontology / BioCarta
389 categories was derived from MSigDB v7.5 Entrez ID .gmt files, downloaded on Apr 12, 2022
390 [http://www.gsea-msigdb.org/gsea/downloads_archive.jsp].
391
392 *Between-species homology*
393
394 Homologene Build 68 was used to determine interspecies homology
395 [ftp.ncbi.nih.gov/pub/HomoloGene/build68/]. Human = taxid:9606, mouse = taxid:10090, rat =
396 taxid:10116, c. elegans = taxid:6239, d. melanogaster = taxid:7227, yeast = taxid:559292,
397 zebrafish = taxid:7955.
398
399 *Primate specificity*
400
401 Human genes were considered primate-specific if the only other members of their homology
402 group belonged to primate genomes. Primate taxonomy ids were downloaded from NCBI
403 Taxonomy on Sep 20, 2022
404 [https://www.ncbi.nlm.nih.gov/taxonomy/?term=txid9443[Subtree]].
405
406 *Number of publications in model organisms*
407
408 Gene information was downloaded from NCBI Gene on Aug 16, 2022
409 [ftp.ncbi.nlm.nih.gov/gene/DATA/GENE_INFO/All_Data.gene_info.gz].
410
411 Only using PMIDs annotated as primary research articles, genes was considered as mentioned
412 in the title/abstract of the publication if gene was annotated as being in the title/abstract by
413 PubTator and the article appeared in gene2pubmed.
414
415 Genes in model organisms were mapped to human genes and the number of articles on those
416 mapping to human genes were counted. If a model organism's gene had homology to human
417 but no associated publications, the number of publications was resolved to zero. Otherwise,
418 counts were listed as NA.
419
420 *Mouse phenotype hits*
421
422 International Mouse Phenotyping Consortium data release 17.0 was downloaded on Aug 18,
423 2022 [https://www.mousephenotype.org/data/release]. Mouse genes were matched to human
424 genes with Homologene.
425
426 *Gene Expression Atlas (EBI-GXA)*
427
EBI-GXA release 36 was downloaded on Sep 15, 2020
[https://web.archive.org/web/20201022184159/https://www.ebi.ac.uk/gxa/download]. This is

the most recent release of EBI-GXA available as a bulk download. For probability of DE, only
RNA-seq comparisons were considered and DE was called at Benjamini-Hochberg $q < 0.05$.

Global RNA expression

RNA consensus tissue gene data from HPA release 21.1 was downloaded on Sep 20, 2022
[https://www.proteinatlas.org/about/download]. Global RNA expression was estimated by
taking the median expression (nTPM) across tissues for each gene and the proportion of tissues
with detectable (≥1 nTPM) expression for each gene.

Expression in HeLa cells

RNA cell line gene data from HPA release 21.1 was downloaded on Sep 20, 2022
[https://www.proteinatlas.org/about/download]. Expression is in nTPM.

Previous patent activity

Genes with patent activity were defined from Table S1 of Rosenfeld and Mason, 2013[68]. Genes
were mapped with their HGNC symbol. This analysis aligned sequences in patents to the human
genome to estimate patent coverage of human coding sequences. Although this does not
necessarily reflect whether the mapped genes were claimed directly by the patent holder, as
noted by others[69], this analysis remains the most comprehensive available for determining
patent coverage of the human genome.

Druggability

Druggable genes were identified from Table S1 of Finan et al., 2017[70]. Genes were mapped with
their Ensembl identifier.

Gene length

GenBank was downloaded in spring 2017 (genome version GRCh38.p10). Gene length is defined
here as the span of the longest transcript on the chromosome. This aligns with the model of
gene length used in Stoeger et al., 2018[7].

Solubility

SwissProt protein sequences and mapping tables to Entrez GeneIDs were downloaded from
Uniprot in spring 2017. Protein GRAVY score (ignoring Pyrrolysine and Selenocysteine) was
estimated with BioPython[71].

Loss of function intolerance

Data was obtained from Karczewski et al.[72]. pLI scores > 0.9 on main transcripts, as flagged by
authors, were considered as highly loss-of-function intolerant as described by Lek et al.[73].
475
476 *Number of GWAS hits*
477
478 EBI GWAS catalog[25] (associations and studies) was download on Aug 17, 2022
479 [https://www.ebi.ac.uk/gwas/docs/file-downloads]. Loci were mapped to the nearest gene.
480
481 *Status as understudied protein*
482
483 The Illuminating the Druggable Genome understudied protein list was downloaded on Sep 20,
484 2022
485 [https://github.com/druggablegenome/IDGTargets/blob/master/IDG_TargetList_CurrentVersio
486 n.json].
487
488 *Human Protein Atlas*
489
HPA release 21.1 was downloaded on Sep 20, 2022 [https://www.proteinatlas.org/search].
Evidence for a protein's existence, as determined by NeXtProt, HPA, or UniProt was resolved as
True if the respective evidence entry was annotated as "Evidence at protein level". Status as a
membrane protein was determined by whether the 'Protein class' column contained the string
'membrane protein'. Antibodies were considered available for each protein if the protein's
entry in the 'Antibody' column was not null.
496
497 *Availability of plasmids*
498
499 The AddGene plasmid catalog was downloaded on Aug 12, 2022
500 [https://www.addgene.org/browse/gene/gene-list-data/?_=1666368044314].
501
502 *Availability of compounds*
503
504 The catalog of gene targets was downloaded from ChEMBL on Sep 20, 2022
505 [https://www.ebi.ac.uk/chembl/g/#browse/targets]. UniProt IDs were converted to Entrez IDs
506 to identify which human genes were affected by any compound.
507
508 *Mendelian inheritance*
509
510 Autosomal dominant [https://hpo.jax.org/app/browse/term/HP:0000006] and autosomal
511 recessive [https://hpo.jax.org/app/browse/term/HP:0000007] inherited disease-gene
512 associations were downloaded from the Human Phenotype Ontology on Sep 20, 2022. Genes
513 were considered to have evidence of Mendelian inheritance if they appeared in these lists of
514 associations.
515
516 *Code*

517
518 Code for analysis is available at github.com/amarallab/fmug_analysis. Code for FMUG is
519 available at github.com/amarallab/fmug.
520
521
522

523
Figure 1: A shift in focus towards well-studied genes occurs during the summarization and
write-up of results and remains in subsequent studies. **a,** Conceptual diagram depicting possible
points of abandonment for understudied genes in studies using high-throughput -omics
experiments. **b,** We identified articles reporting on genome-wide CRISPR screens (CRISPR, 15
focus articles and 18 citing articles), transcriptomics (T-omics, 148 focus articles and 1,678 citing
articles), affinity purification–mass spectrometry (AP-MS, 296 focus articles and 1,320 citing
articles), and GWAS (450 focus articles and 3,524 citing articles). Focusing only on protein-
coding genes (white box plot), we retrieved data uploaded to repositories describing which
genes came up as "hits" in each experiment (first colored box plot). We then retrieved the hits
mentioned in the titles and abstracts of those articles (second colored box plot) and hits
mentioned in the titles and abstracts of articles citing those articles (third colored box plot).
Unique hit genes are only counted once. Bibliometric data reveals that understudied genes are
frequently hits in -omics experiments but are not typically highlighted in the title/abstract of
reporting articles, nor in the title/abstract or articles citing reporting articles. ** denotes $p <$
0.01 and *** denotes $p < 0.001$ by two-sided Mann-Whitney U test, comparing genes
highlighted in title/abstract to genes present in hit lists.
540

Figure 2: Articles focusing on less popular genes tend to accrue more citations. a, Density plot

shows correlation between articles per gene before 2015 and median citations to articles

published in 2015. Contours correspond to deciles in density. Solid red line shows locally

weighted scatterplot smoothing (LOWESS) regression. ρ is Spearman rank correlation and p the

significance values of the Spearman rank correlation as described by Kendall and Stuart[74]. We

forgo depicting more recent years than 2015 to allow for citations to accumulate over multiple

years, providing a more sensitive and robust readout of long-term impact. **b,** Spearman

correlation of previous gene popularity (i.e. number of articles) to median citations per year

since 1990. Solid blue line indicates nominal Spearman correlation, shaded region indicates

bootstrapped 95% confidence interval (n=1,000). Only articles with a single gene in the

title/abstract are considered, excluding the 30.4% of gene-focused studies which feature more

than one gene in the title/abstract. For more recent years, where articles have had less time to

accumulate citations, insufficient signal may cause correlation to converge toward zero.

Figure 3: We evaluated which gene-related factors are associated with elevation to the

title/abstract of an article featuring a high-throughput experiment. a) Association between

factors with binary (True/False) identities and highlighting hits in title/abstract of reporting

articles. Values represent the odds ratio between hits in the collected articles and hits
mentioned in the title or abstract of collected articles (e.g. hits with a compound known to
affect gene activity are 4.262 times as likely to be mentioned in the title/abstract in an article
using transcriptomics, corresponding to an odds ratio of 4.331). Collected articles are described
in **Figure 1B** and **Figures S5**, **S6** and **S7.** 95% confidence interval of odds ratio is shown in
parentheses. * = Benjamini-Hochberg FDR < 0.05, ** = FDR < 0.01, and *** = FDR < 0.001 by
two-sided Fisher exact test. Results are shown numerically in **Table S2**. For consistency between
studies, hits were restricted to protein-coding genes. Thus, status as a protein-coding gene
could not be tested. †No genes without a defined HUGO symbol were found as hits in GWAS or
transcriptomics studies. **b)** Association with factors with continuous identities and highlighting
hits in title/abstract of reporting articles. Values represent F, the common-language effect size
(equivalent to AUROC, where ~0.5 indicates little effect, >0.5 indicates positive effect and <0.5
indicates negative effect) of being mentioned in the titles/abstracts of the collected articles
described in **Figure 1B** and **Figures S5**, **S6** and **S7.** * = Benjamini-Hochberg FDR < 0.05, ** = FDR
< 0.01, and *** = FDR < 0.001 by two-sided Mann-Whitney U test. Results are shown
numerically in **Table S3**.

Figure 4: We created FMUG to help researchers identify understudied genes among their genes of interest and characterize their tractability for future research. a, Diagram describing use of FMUG. **b,** An early prototype of FMUG led us to the hypothesis that transcript length negatively correlates with up-regulation during aging. First, we identified genes that strongly associate with age-dependent transcriptional change across multiple cohorts. We then performed a literature review for each of these genes to identify the most direct way the genes (or evolutionarily closely related genes or functionally closely related partner proteins) had been studied in aging. 64% had been functionally investigated in aging, 15% shown to change a measure of gene expression, 3% functionally investigated in a biological domain close to aging (such as senescence), and 5% shown to change a measure of gene expression in a biological

domain close to aging. For genes reported by others to change expression with age, we identified tissues in which transcripts of the genes change during aging. We computed 'feasibility scores' scientific strategies (GEM: G: strong genetic support, E: and experimental potential, M: homolog in invertebrate model organism) as described by Stoeger et al.[7] and total number of publications in MEDLINE. Splicing factor, proline- and glutamine-rich (*Sfpq*) had previously been demonstrated by Takeuchi et al. to be required for the transcriptional elongation of long genes[51]. When performing a data-driven analysis of factors that could possibly explain age-dependent changes of the entire transcriptome, we thus included gene and transcript lengths, and subsequently found them to be more informative than transcription factors or microRNAs[52].

Data Availability

All underlying data for figures are available at github.com/amarallab/fmug_analysis.

Acknowledgements

We thank Xiaojing Sui for testing FMUG and Northwestern Information Technology for technical assistance. RAKR was supported in part by the National Institutes of Health Training Grant (T32GM008449) through Northwestern University's Biotechnology Training Program. RAKR also acknowledges support from the Dr. John N. Nicholson fellowship from Northwestern University and Moderna Inc., "Identifying bias and improving reproducibility in RNA-seq computational pipelines". LANA was supported by NSF 1956338, NIH U19AI135964 and Simons Foundation DMS-1764421. TS was supported by NIH K99AG068544. We thank Alexander Misharin, Richard Morimoto, and Scott Budinger for feedback on an early prototype of FMUG which we used as part of our shared research into the biology of aging.

References

Hoffmann, R. & Valencia, A. Life cycles of successful genes. *Trends Genet.* **19**, 79-81 (2003). https://doi.org:10.1016/S0168-9525(02)00014-8

Su, A. I. & Hogenesch, J. B. Power-law-like distributions in biomedical publications and research funding. *Genome Biol.* **8**, 404 (2007). https://doi.org:10.1186/gb-2007-8-4-404

Edwards, A. M. *et al.* Too many roads not taken. *Nature* **470**, 163-165 (2011). https://doi.org:10.1038/470163a

Gillis, J. & Pavlidis, P. Assessing identity, redundancy and confounds in Gene Ontology annotations over time. *Bioinformatics* **29**, 476-482 (2013). https://doi.org:10.1093/bioinformatics/bts727

Stoeger, T. & Nunes Amaral, L. A. The characteristics of early-stage research into human genes are substantially different from subsequent research. *PLoS Biol* **20**, e3001520 (2022). https://doi.org:10.1371/journal.pbio.3001520

Grueneberg, D. A. *et al.* Kinase requirements in human cells: I. Comparing kinase requirements across various cell types. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 16472-16477 (2008). https://doi.org:10.1073/pnas.0808019105

7   Stoeger, T., Gerlach, M., Morimoto, R. I. & Nunes Amaral, L. A. Large-scale investigation
of the reasons why potentially important genes are ignored. *PLoS Biol* **16**, e2006643
(2018). https://doi.org:10.1371/journal.pbio.2006643
8   Riba, M. *et al.* Revealing the acute asthma ignorome: characterization and validation of
uninvestigated gene networks. *Sci Rep* **6**, 24647 (2016).
https://doi.org:10.1038/srep24647
9   Haynes, W. A., Tomczak, A. & Khatri, P. Gene annotation bias impedes biomedical
research. *Sci Rep* **8**, 1362 (2018). https://doi.org:10.1038/s41598-018-19333-x
10   Border, R. *et al.* No support for historical candidate gene or candidate gene-by-
interaction hypotheses for major depression across multiple large samples. *American
Journal of Psychiatry* **176**, 376-387 (2019).
11   Stoeger, T. & Nunes Amaral, L. A. COVID-19 research risks ignoring important host genes
due to pre-established research patterns. *Elife* **9** (2020).
https://doi.org:10.7554/eLife.61981
12   Zhang, D. *et al.* Incomplete annotation has a disproportionate impact on our
understanding of Mendelian and complex neurogenetic disorders. *Science Advances* **6**,
eaay8299 (2020).
13   Byrne, J. A. *et al.* Protection of the human gene research literature from contract
cheating organizations known as research paper mills. *Nucleic Acids Research* **50**, 12058-
12070 (2022). https://doi.org:10.1093/nar/gkac1139
14   Collins, F. S., Green, E. D., Guttmacher, A. E., Guyer, M. S. & Institute, U. S. N. H. G. R. A
vision for the future of genomics research. *Nature* **422**, 835-847 (2003).
https://doi.org:10.1038/nature01626
15   Shendure, J., Findlay, G. M. & Snyder, M. W. Genomic Medicine-Progress, Pitfalls, and
Promise. *Cell* **177**, 45-57 (2019). https://doi.org:10.1016/j.cell.2019.02.003
16   Lloyd, K. C. K. *et al.* The Deep Genome Project. *Genome Biol* **21**, 18 (2020).
https://doi.org:10.1186/s13059-020-1931-9
17   Kustatscher, G. *et al.* Understudied proteins: opportunities and challenges for functional
proteomics. *Nat Methods* **19**, 774-779 (2022). https://doi.org:10.1038/s41592-022-
01454-x
18   Rodriguez-Esteban, R. & Jiang, X. Differential gene expression in disease: a comparison
between high-throughput studies and the literature. *BMC Med Genomics* **10**, 59 (2017).
https://doi.org:10.1186/s12920-017-0293-y
19   Oprea, T. I. *et al.* Unexplored therapeutic opportunities in the human genome. *Nat Rev
Drug Discov* **17**, 317-332 (2018). https://doi.org:10.1038/nrd.2018.14
20   Sinha, S., Eisenhaber, B., Jensen, L. J., Kalbuaji, B. & Eisenhaber, F. Darkness in the
Human Gene and Protein Function Space: Widely Modest or Absent Illumination by the
Life Science Literature and the Trend for Fewer Protein Function Discoveries Since 2000.
Proteomics **18**, e1800093 (2018). https://doi.org:10.1002/pmic.201800093
21   Wood, V. *et al.* Hidden in plain sight: what remains to be discovered in the eukaryotic
proteome? *Open Biol* **9**, 180241 (2019). https://doi.org:10.1098/rsob.180241
22   Donohue, C. & Love, A. Perspectives on the Human Genome Project and genomics.
Minnesota Studies in the Philosophy of Science **23** (in press).

23 Pena-Castillo, L. & Hughes, T. R. Why are there still over 1000 uncharacterized yeast
genes? *Genetics* **176**, 7-14 (2007). https://doi.org:10.1534/genetics.107.074468
24 Ellens, K. W. *et al.* Confronting the catalytic dark matter encoded by sequenced
genomes. *Nucleic Acids Res* **45**, 11495-11514 (2017).
https://doi.org:10.1093/nar/gkx937
25 Buniello, A. *et al.* The NHGRI-EBI GWAS Catalog of published genome-wide association
studies, targeted arrays and summary statistics 2019. *Nucleic Acids Res* **47**, D1005-
D1012 (2019). https://doi.org:10.1093/nar/gky1120
26 Oughtred, R. *et al.* The BioGRID database: A comprehensive biomedical resource of
curated protein, genetic, and chemical interactions. *Protein Science* **30**, 187-200 (2021).
27 Papatheodorou, I. *et al.* Expression Atlas: gene and protein expression across multiple
studies and organisms. *Nucleic acids research* **46**, D246-D251 (2018).
28 Maglott, D., Ostell, J., Pruitt, K. D. & Tatusova, T. Entrez Gene: gene-centered
information at NCBI. *Nucleic acids research* **35**, D26-D31 (2007).
29 Wei, C.-H., Allot, A., Leaman, R. & Lu, Z. PubTator central: automated concept
annotation for biomedical full text articles. *Nucleic acids research* **47**, W587-W593
(2019).
30 Hutchins, B. I., Santangelo, George. *iCite*, <https://doi.org/10.35092/yhjc.c.4586573 >
(2019).
31 Stoeger, T. & Nunes Amaral, L. A. COVID-19 research risks ignoring important host genes
due to pre-established research patterns. *Elife* **9**, e61981 (2020).
32 Stoeger, T., Gerlach, M., Morimoto, R. I. & Nunes Amaral, L. A. Large-scale investigation
of the reasons why potentially important genes are ignored. *PLoS biology* **16**, e2006643
(2018).
33 Pfeiffer, T. & Hoffmann, R. Temporal patterns of genes in scientific publications. *Proc.*
Natl. Acad. Sci. U. S. A. **104**, 12052-12056 (2007).
https://doi.org:10.1073/pnas.0701315104
34 Knorr Cetina, K. *Epistemic Cultures*. (Harvard University Press, 1999).
35 Alberts, B. M. Limits to growth: In biology, small science is good science. *Cell*
(Cambridge) **41**, 337-338 (1985).
36 Richardson, S. S. a. S., Hallam.   (Duke University Press, 2015).
37 Lopes, I., Altab, G., Raina, P. & De Magalhães, J. P. Gene size matters: an analysis of gene
length in the human genome. *Frontiers in Genetics* **12**, 559998 (2021).
38 Duek, P., Gateau, A., Bairoch, A. & Lane, L. Exploring the Uncharacterized Human
Proteome Using neXtProt. *J Proteome Res* **17**, 4211-4226 (2018).
https://doi.org:10.1021/acs.jproteome.8b00537
39 Crow, M., Lim, N., Ballouz, S., Pavlidis, P. & Gillis, J. Predictability of human differential
gene expression. *Proc Natl Acad Sci U S A* **116**, 6491-6500 (2019).
https://doi.org:10.1073/pnas.1802973116
40 Perdigao, N. & Rosa, A. Dark Proteome Database: Studies on Dark Proteins. *High*
Throughput **8** (2019). https://doi.org:10.3390/ht8020008
41 Essegian, D., Khurana, R., Stathias, V. & Schurer, S. C. The Clinical Kinase Index: A
Method to Prioritize Understudied Kinases as Drug Targets for the Treatment of Cancer.
Cell Rep Med **1**, 100128 (2020). https://doi.org:10.1016/j.xcrm.2020.100128

42 Sheils, T. K. *et al.* TCRD and Pharos 2021: mining the human proteome for disease
biology. *Nucleic Acids Res* **49**, D1334-D1346 (2021).
https://doi.org:10.1093/nar/gkaa993
43 Rocha, J. *et al.* Functional unknomics: closing the knowledge gap to accelerate
biomedical research. *bioRxiv* (2022).
44 Higgins, D. P., Weisman, C. M., Lui, D. S., D'Agostino, F. A. & Walker, A. K. Defining
characteristics and conservation of poorly annotated genes in Caenorhabditis elegans
using WormCat 2.0. *Genetics* **221** (2022). https://doi.org:10.1093/genetics/iyac085
45 Wainberg, M. *et al.* A genome-wide atlas of co-essential modules assigns function to
uncharacterized genes. *Nat Genet* **53**, 638-649 (2021). https://doi.org:10.1038/s41588-
021-00840-z
46 Rebhan, M., Chalifa-Caspi, V., Prilusky, J. & Lancet, D. GeneCards: a novel functional
genomics compendium with automated data mining and query reformulation support.
Bioinformatics **14**, 656-664 (1998). https://doi.org:10.1093/bioinformatics/14.8.656
47 Tan, J. *et al.* ADAGE signature analysis: differential expression analysis with data-defined
gene sets. *Bmc Bioinformatics* **18**, 512 (2017). https://doi.org:10.1186/s12859-017-
1905-4
48 Kustatscher, G. *et al.* Co-regulation map of the human proteome enables identification
of protein functions. *Nat Biotechnol* **37**, 1361-1371 (2019).
https://doi.org:10.1038/s41587-019-0298-5
49 Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
Innovation (Camb) **2**, 100141 (2021). https://doi.org:10.1016/j.xinn.2021.100141
50 Jiang, J. *et al.* Systematic illumination of druggable genes in cancer genomes. *Cell Rep*
38, 110400 (2022). https://doi.org:10.1016/j.celrep.2022.110400
51 Takeuchi, A. *et al.* Loss of Sfpq Causes Long-Gene Transcriptopathy in the Brain. *Cell Rep*
23, 1326-1341 (2018). https://doi.org:10.1016/j.celrep.2018.03.141
52 Stoeger, T. *et al.* Aging is associated with a systemic length-associated transcriptome
imbalance. *Nature Aging* **2**, 1191-1206 (2022). https://doi.org:10.1038/s43587-022-
00317-6
53 Gyenis, A. *et al.* Genome-wide RNA polymerase stalling shapes the transcriptome during
aging. *Nat Genet* **55**, 268-279 (2023). https://doi.org:10.1038/s41588-022-01279-6
54 Ibañez-Solé, O., Barrio, I. & Izeta, A. Age or lifestyle-induced accumulation of
genotoxicity is associated with a length-dependent decrease in gene expression.
iScience, 106368 (2023). https://doi.org:https://doi.org/10.1016/j.isci.2023.106368
55 Brown, L. A. & Peirson, S. N. Improving Reproducibility and Candidate Selection in
Transcriptomics Using Meta-analysis. *Journal of Experimental Neuroscience* **12**,
1179069518756296 (2018). https://doi.org:10.1177/1179069518756296
56 Cesar-Razquin, A. *et al.* A Call for Systematic Research on Solute Carriers. *Cell* **162**, 478-
487 (2015). https://doi.org:10.1016/j.cell.2015.07.022
57 Evans, J. A. Electronic publication and the narrowing of science and scholarship. *Science*
321, 395-399 (2008). https://doi.org:10.1126/science.1150473
58 Alberts, B., Kirschner, M. W., Tilghman, S. & Varmus, H. Rescuing US biomedical
research from its systemic flaws. *Proc Natl Acad Sci U S A* **111**, 5773-5777 (2014).
https://doi.org:10.1073/pnas.1404402111

59 Uzzi, B., Mukherjee, S., Stringer, M. & Jones, B. Atypical combinations and scientific
impact. *Science* **342**, 468-472 (2013). https://doi.org:10.1126/science.1240474
60 Gilovich, T., Griffin, D. W. & Kahneman, D. *Heuristics and biases : the psychology of*
intuitive judgment. (Cambridge University Press, 2002).
61 Gerlt, J. A. *et al.* The Enzyme Function Initiative. *Biochemistry* **50**, 9950-9962 (2011).
https://doi.org:10.1021/bi201312u
62 Carter, A. J. *et al.* Target 2035: probing the human proteome. *Drug Discov Today* **24**,
2111-2115 (2019). https://doi.org:10.1016/j.drudis.2019.06.020
63 Rodgers, G. *et al.* Glimmers in illuminating the druggable genome. *Nat Rev Drug Discov*
17, 301-302 (2018). https://doi.org:10.1038/nrd.2017.252
64 Kustatscher, G. *et al.* An open invitation to the Understudied Proteins Initiative. *Nat*
Biotechnol **40**, 815-817 (2022). https://doi.org:10.1038/s41587-022-01316-z
65 *EUbOPEN*, <https://www.eubopen.org/> (
66 Rocha, J. J. *et al.* Functional unknomics: Systematic screening of conserved genes of
unknown function. *PLoS biology* **21**, e3002222 (2023).
67 Wei, C. H., Allot, A., Leaman, R. & Lu, Z. PubTator central: automated concept
annotation for biomedical full text articles. *Nucleic Acids Res* **47**, W587-W593 (2019).
https://doi.org:10.1093/nar/gkz389
68 Rosenfeld, J. A. & Mason, C. E. Pervasive sequence patents cover the entire human
genome. *Genome Med.* **5**, 27 (2013). https://doi.org:10.1186/gm431
69 Tu, S. *et al.* Response to 'pervasive sequence patents cover the entire human genome'.
Genome medicine **6**, 1-3 (2014).
70 Finan, C. *et al.* The druggable genome and support for target identification and
validation in drug development. *Science translational medicine* **9**, eaag1166 (2017).
71 Cock, P. J. A. *et al.* Biopython: freely available Python tools for computational molecular
biology and bioinformatics. *Bioinformatics* **25**, 1422-1423 (2009).
https://doi.org:10.1093/bioinformatics/btp163
72 Karczewski, K. J. *et al.* The mutational constraint spectrum quantified from variation in
141,456 humans. *Nature* **581**, 434-443 (2020). https://doi.org:10.1038/s41586-020-
2308-7
73 Lek, M. *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* **536**,
285-291 (2016). https://doi.org:10.1038/nature19057
74 Kendall, M. G. & Stuart, A. Inference and Relationship *The Advanced Theory of Statistics*
2 (1973).
75 Ward Jr, J. H. Hierarchical grouping to optimize an objective function. *Journal of the*
American statistical association **58**, 236-244 (1963).
797