Early striatal hyperexcitability in an *in vitro* human striatal microcircuit model

carrying the Parkinson's *GBA-N370S* mutation

Quyen B. Do[1,2,3], Bryan Ng[1,2], Ricardo Marquez Gomez[1,2,3], Dayne Beccano-Kelly[1,2], Naroa Ibarra-

Aizpura[1,2], Maria-Claudia Caiazza[1,2, 3], Charmaine Lang[1,2, 3], Jimena Baleriola[4,6], Nora Bengoa-

Vergniory[1,2,3, 4,5,6*] and Richard Wade-Martins[1,2,3*]

[1] Oxford Parkinson's Disease Centre and Department of Physiology, Anatomy and Genetics, University

of Oxford, South Park Road, Oxford OX1 3QU, United Kingdom

[2] Kavli Institute for Neuroscience Discovery, University of Oxford, Dorothy Crowfoot Hodgkin Building,

South Park Road, Oxford OX1 3QU, United Kingdom

[3] Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD,

20815, USA

[4] Achucarro Basque Center for Neuroscience, Leioa, Spain

[5] University of the Basque Country (UPV/EHU), Department of Neuroscience, Leioa, Spain

[6] Ikerbasque - Basque Foundation for Science, Bilbao, Spain

*Corresponding authors:

richard.wade-martins@dpag.ox.ac.uk

nora.bengoa-vergniory@dpag.ox.ac.uk

25
26

Abstract

Understanding medium spiny neuron (MSN) physiology is essential to understand motor impairments

in Parkinson's disease (PD) given the architecture of the basal ganglia. Here, we developed a custom

three-chamber microfluidic platform and established a cortico-striato-nigral microcircuit

recapitulating the striatal presynaptic triad *in vitro* using induced pluripotent stem cell (iPSC)-derived

neurons. We found that, although cortical glutamatergic projections facilitated MSN synaptic activity,

dopaminergic transmission was essential for excitability maturation of MSNs *in vitro*. Replacement of

wild-type iPSC-dopamine neurons (iPSC-DaNs) in the striatal microcircuit with those carrying the PD-

related *GBA-N370S* mutation induced early hyperexcitability in iPSC-MSNs through reduction of

voltage-gated sodium and potassium intrinsic currents. Such deficits were resolved in aged cultures

or with antagonism of protein kinase A activity in nigrostriatal iPSC-DaNs. Hence, our results highlight

the unique utility of modelling striatal neurons in a modular and highly physiological circuit which is

essential to reveal mechanistic insights of the loss of electrical functional integrity in the striata of

GBA1 PD patients.

Introduction

The striatum is comprised of 90% medium-sized spiny neurons (MSNs) and represents the primary input nucleus of the basal ganglia [1]. The striatum receives converging glutamatergic and dopaminergic inputs from the cortex and midbrain, respectively, forming a presynaptic triad critical for regulation of MSN functional activity. Striatal dysfunction plays a key role in the motor features of Parkinson's disease (PD), although the exact underlying mechanisms remain under intense study with divergent findings [2–4]. The consensus suggests that the characteristic preferential degeneration of substantia nigra *pars compacta* (SNc) dopaminergic neurons (DaNs) results in aberrant striatal activity patterns which subsequently alter the balanced output of the basal ganglia governing motor behaviour [5]. However, the specific effects on the two parallel principle striatal pathways, namely the direct pathway and indirect pathway, are contentious. The classical premise suggested functionally opposed roles of direct and indirect pathway neurons in regulating action and their diametrical changes in firing rate in PD [2], although a more recent observation of co-activation of both pathways at the initiation of movement has challenged this view [6].

The advent of human induced pluripotent stem cell (iPSC) technology has revolutionized the ability to investigate the pathophysiology of neurological disease in human neurons, revealing a number of potential cellular mechanisms underpinning preferential degeneration of SNc DaNs [7–12]. However, much less is known about how the presence of Parkinson's mutations changes DaN function and how this in turn impacts MSN function. *In vitro* modelling of iPSC-derived human striatal neurons has developed from original striatal monocultures [13–15] to more recent three-dimensional cortico-striatal assembloids [16], however such models have not included dopaminergic neurons.

Microfluidic neuronal culture technology has tremendously assisted efforts in modelling and understanding biological systems, primarily enabling custom and precise control of neurite isolation [17,18] as well as orchestrated neuronal circuitry [19,20]. In this work we have established a microfluidic

platform comprising an all iPSC-derived striatal presynaptic microcircuit to recapitulate the

endogenous connectivity of cortical and dopamine neurons onto MSNs. This setup is permissive to

long-term cultures of up to 100 days and allows for the investigation of temporal changes in neuronal

physiology. With this system we have dissected the importance of dual presynaptic glutamatergic and

dopaminergic inputs in facilitating both electrophysiological maturation of *in vitro* MSNs as well as the

early striatal pathology affected by the PD-related *GBA-N370S* genetic mutation.

73
74

Results

Generation of functional iPSC-derived medium spiny neurons (MSNs)

We first applied a defined cocktail of small molecules to recreate the temporal *in vivo* development

of MSNs to reflect their telencephalic and lateral ganglionic eminence (LGE) lineage (Figure 1A). To

induce forebrain development, we used modulators of SMAD, namely LDN and SB [21], and an inhibitor

of the WNT signalling pathway (XAV) at the start of the differentiation [22]. Activin A, which was

independently reported to enhance dorsal striatal patterning [14,16], was introduced from DIV12-23 to

promote LGE identity. Activin A from DIV 12 increased the expression of selected markers for LGE and

MSN progenitors two days post treatment (Supplementary Figure 1A) as well as the proportion of

post-mitotic neurons co-expressing DARRP32 and CTIP2 on DIV 35 (Supplementary Figure 1B).

Temporal examination of transcript abundance of LGE and MSN progenitor markers using RT-qPCR

suggested gradual patterning of iPSCs towards MSNs reminiscent of their *in vivo* developmental

trajectory (Figure 1B-D). LGE-specific genes such as *DLX2, ASCL1*, and *GSX2* were enriched in early

time-points and this enrichment was subsequently reduced in post-mitotic MSNs (Figure 1B).

Meanwhile, expression of selected markers of MSN progenitors including *ISL1, DLX5, FOP2,* and *MEIS2*

increased over time and remained abundant in post-mitotic MSN cultures (Figure 1C). *CTIP2, GAD61,*

and *DARPP32*, which are indicative of post-mitotic MSNs, were virtually absent in early cultures and

only subsequently expressed in great abundance from DIV 14 (Figure 1D). Tyrosine hydroxylase (*TH*)

transcript was also detected in late cultures, implying the presence of potentially non-MSNs, TH-

expressing neurons (Figure 1D). Notably, expression of *NKX2.*1, the medial ganglionic eminence

marker, was barely detectable throughout the culture development (Figure 1B), indicating a strong

LGE enrichment in the patterning.

More than 50% of MAP2+ iPSC-MSNs at DIV 40 expressed CTIP2, DARPP32, or GAD67 and up to 37%

co-expressed both canonical markers CTIP2 and DARPP32 (Figure 1E-F). We also observed expression

of selected markers specific to either the direct- or indirect-pathway MSNs (Supplementary Figure 2A-

B). Immunostaining confirmed expression of MSN-subtype specific neurotransmitter and

neurotransmitter precursors including PENK, PDYN and SUBSTANCE P at DIV 40 (Supplementary Figure

2C). Quantification of GABAergic neurons co-expressing either PKEN, PDYN or SUBSTANCE P suggested

the dominant presence of direct pathway MSNs intermingled with a modest population of indirect

pathway MSNs in our iPSC-MSN cultures (Supplementary Figure 2D).

We next performed whole cell patch-clamping to examine functional maturation of DARPP32-positive

iPSC-derived MSNs (Figure 1H). Our iPSC-derived MSNs displayed hyperpolarised resting membrane

potential (DIV 40-45: -51.2 ± 1.62 mV, n = 25 neurons; DIV 55-60: -53.3 ± 1.47, n = 43 neurons; DIV 70-

75: -55.1 ± 1.36 mV, n = 29 neurons), large cell capacitance (DIV 40-45: 24.0 ± 1.26 pA, n = 24 neurons;

DIV 55-60: 28 ± 1.11 pA, n = 42 neurons; DIV 70-75: 31.6 ± 2.31 pA, n = 26 neurons), and large input

resistance (R_{in}) (DIV 40-45: 1.39 ± 0.15 GΩ, n = 20 neurons; DIV 55-60: 1.29 ± 0.10 GΩ, n = 41 neurons;

DIV 70-75: 1.0 ± 0.08 GΩ, n = 27 neurons) (Figure 1I-L). There was no significant change of these

intrinsic properties of MSNs from DIV 40-75 (RMP: p = 0.265; R_{in}: p = 0.09; all were analysed with one-

way ANOVA with Bonferroni multiple comparison correction test) with the exception of cell

capacitance between DIV 40-45 and DIV 70-75 (p = 0.009). Moreover, we observed firing of action

potentials (APs) with the characteristic delayed first spike (Figure 1G) upon current injections. The

proportion of functionally mature neurons that displayed evoked APs compared to those that were

unresponsive increased as the culture aged (DIV 40-45: 7/10 neurons; DIV 55-60: 20/14; DIV 70-75:

14/6) (Figure 1L). These data indicated robust generation of functionally active iPSC-MSN

monocultures that are electrophysiologically stable from DIV 40 to DIV 75.

***In vitro* recapitulation of the striatal microcircuit on custom microfluidic devices**

Despite the functionally active status of iPSC-derived MSNs in monocultures, we noted a more

depolarised RMP and larger R_{in} in *in vitro* MSNs relative to *in vivo* MSNs [23]. We hypothesised that such

discrepancy was principally due to the lack of non-autologous synapses that endogenously provide

valuable developmental cues to modulates MSN functional maturation.

To address the lack of maturational signalling within MSN monocultures, we reconstructed the striatal

microcircuit *in vitro* mimicking part of the pre-synaptic complexity of the glutamatergic inputs on

MSNs using polydimethylsiloxane-based open three-chamber microfluidic chips. The device consisted

of three rectangle chambers (4*10 mm) connected via an array of approximately 120 microgrooves of

5 μm wide, 450 μm long and 10 μm high (Figure 2A). The narrow width of the micro-channels

compartmentalised the neuronal soma in its respective chamber while DaN and CN axons, but not

their dendrites, traversed the full 450 μm length of the microchannels to make connections with MSNs

in the middle chamber (Figure 2D).

iPSC-derived DaNs, CNs, MSNs and rat astrocytes were sequentially plated into their respective

chambers so as to recapitulate the cortico-striato-nigral triad in the striatum (Figure 2C-D). DaN and

CN axons were physically guided toward the middle MSN chamber via flow driven by pressure

difference (Figure 2B). We exploited the previous observation that BDNF and laminin are selectively

critical for axon polarisation and outgrowth [24,25] by setting up artificial BDNF and laminin gradients

across the channels to chemically attract axonal projections from the CN and DaN chambers into

middle MSN chambers (Figure 2B) prior to MSN deposition. Dopaminergic and cortical axons took

approximately 10 days *in vitro* to fully cross and occupy all available microgrooves. Growth of MSN

axons travelling out to the DaN and CN chambers was restricted by delaying seeding of iPSC-MSNs

until after all microgrooves had been populated with DaNs or CNs axons (Figure 2D). By DIV 0 of the

microcircuit culture, the axons of both the iPSC-DaNs and iPSC-CNs had densely projected into the

MSN chamber (Figure Di-iii) to regulate MSN progenitor development. In addition to attaining directed

connectivity, the culturing schedule was also designed to incorporate known temporal features of

these neurons as characterised above in their respective monocultures. Specifically, once the culture

platform was established, we applied the strategy of adding DIV 16 iPSC-MSN progenitors to complete

the circuit with post-mitotic and functionally active iPSC-DaNs and iPSC-CNs. Cultures were then left

until our experimental timepoints (i.e. DIV 40-70 MSNs) when all three neuronal populations were

individually electrically active [26].

Glutamatergic input facilitates synaptic maturation of *in vitro* MSNs

Cortical and dopaminergic afferents form characteristic presynaptic triads with MSN dendrites,

specifically on MSN dendritic spine heads and necks, respectively [27,28]. Corticostriatal terminals

represent the major source of glutamatergic input to regulate striatal GABAergic output while

dopamine is thought to exert a modulatory influence on the responsiveness of MSNs to excitatory

synaptic signals via the regulation of the cell intrinsic excitability state [23].

To examine the physiological role cortical collaterals exerted onto iPSC-MSNs in our *in vitro*

microcircuit, we compared various electrophysiological parameters between iPSC-MSNs grown on

coverslips in monoculture (i.e. CN⁻MSN) versus those co-cultured with iPSC-CNs (i.e. CN⁺MSN) on

microfluidic devices (Figure 3A). We found no differences in RMP across time or between culture

conditions over the period from DIV 40-70 (RMP: culturing condition p = 0.46; DIV p = 0.29). No

significant effect of culturing condition was observed for both cell capacitance and rheobase current

at all three timepoints DIV 40-45, 55-60 and 70-75 (CN$^-$MSN versus CN$^+$MSN: all p > 0.05) (Figure 3B-

D). CN$^+$MSN also displayed no differences in spontaneous inhibitory post-synaptic current (sIPSCs) (all

p > 0.05) (Figure 3E, F and I). However, coculture with CNs exerted significant effect on both frequency

and magnitude of spontaneous excitatory post-synaptic currents (sEPSCs), particularly more

pronounced in late cultures (sEPSC frequency: DIV 40-45 and 70-75 p > 0.9, DIV 55-60 p = 0.012; sEPSC

magnitude: DIV 40-45 p > 0.9, DIV 55-60 p = 0.039, DIV 70-75 p = 0.025) (Figure 3G, H and J). These

results imply that our microcircuit setup enables cortical afferents to make functional synapses with

MSNs and enhance MSN excitatory synaptic activities.

Dendritic spines are the principal sites of synaptic integration and their architecture is highly dynamic,

reflecting the complex synaptic communication between neurons. Knowing that glutamatergic input

profoundly modulates dendritic morphology of MSNs [29], we examined whether we could observe a

modified phenotype in dendritic spines in the presence of cortical glutamatergic innervation. We

found that cortical excitatory afferents promoted early dendritic spine formation in MSNs which was

detectable from DIV 40 to 75 (Figure 3K and L). On the contrary, dendritic spines were virtually absent

in early culture of MSNs devoid of cortical projections, but emerged gradually over time and attained

a similar density as in CN$^+$MSN at DIV 70-75 despite CN$^-$MSN remaining relatively synaptically more

quiescent (DIV 40-45: p = 0.001; DIV 55-60: p = 0.0254, DIV 70-75: p = 0.607 CN$^-$MSN vs. CN$^+$MSN)

(Figure 3K and 2D). These results suggested the importance of cortical afferents in facilitating synaptic

maturation of MSNs by prompting not only the formation but also the maturation of dendritic spines

of striatal neurons, indicating the utility of our microcircuit in mimicking functional significance of the

network connectivity.

Dopaminergic signalling promoted excitability maturation in a time-sensitive manner

To elucidate the direct effect of dopaminergic projections onto corticostriatal neurons, comparisons

were made between CN$^+$MSN (i.e. DaN$^-$CN$^+$MSN) and those circuited with DaNs (i.e. DaN$^+$CN$^+$MSN,

Figure 4A). We found that RMP of corticostriatal neurons became significantly more hyperpolarised

with the long-term presence of DaN projections (DIV 40-45 DaN$^+$CN$^+$MSN vs DIV 70-75 DaN$^+$CN$^+$MSN

p = 0.016), resulting in a significantly more hyperpolarised RMP of DaN$^+$CN$^+$MSN in late, but not early,

cultures (DaN$^-$CN$^+$MSN RMP vs. DaN$^+$CN$^+$MSN RMP: DIV 40-45 p = 0.4041, DIV 55-60 p = 0.1120, DIV

70-75 p = 0.044) (Figure 4B). This time-dependent hyperpolarisation of DaN$^+$CN$^+$MSN RMP was,

however, not coupled with differences in rheobase currents and input resistance of MSNs receiving

dopaminergic inputs (Rheobase current: culturing condition p = 0.4607; input resistance: culturing

condition p = 0.3802) (Figure 4C and D). Observing the trends toward reduced intrinsic excitability

produced by dopaminergic afferents, we investigated the constituent voltage-gated currents of

sodium (Na$_v$) and potassium (K$_v$) to delineate specific ionic channels responsible for the subtle

differences. The former underlies membrane depolarisation and hence, firing probability [30], while the

extensive expression of the latter is fundamental for the resting hyperpolarised state of *in vivo* MSNs

[31]. We found an increase in both the fast A-type and slow activating K$_v$ in late cultures at DIV 70-75

(DaN$^-$CN$^+$MSN vs DaN$^+$CN$^+$MSN fast A-type K$_v$: DIV 40-45 p = 0.508, DIV 55-60 p = 0.317, DIV 70-75 p

= 0.0061; DaN$^-$CN$^+$MSN vs DaN$^+$CN$^+$MSN fast slow activating K$_v$: DIV 40-45 p = 0.694, DIV 55-60 p =

0.293, DIV 70-75 p = 0.0023) (Figure 4F-H) but no significant temporal changes in Na$_v$ currents (DaN$^-$

CN$^+$MSN vs DaN$^+$CN$^+$MSN: DIV 40-45 p = 0.927, DIV 55-60 p = 0.794, DIV 70-75 p = 0.235) (Figure 4E),

in agreement with the more hyperpolarised RMP but similar rheobase current. The difference in K$_v$

can result from either increased K$_v$ of DaN$^+$CN$^+$MSN, or decreased K$_v$ of DaN$^-$CN$^+$MSN, or both. When

examining each condition independently over the period of DIV 40-70, we observed a gradual increase

in K$_v$ among DaN$^+$CN$^+$MSN while K$_v$ of DaN$^-$CN$^+$MSN increased gradually between DIV 40-65 but

subsequently dropped at DIV 70-75 (Supplementary Figure 3). These data suggest that MSN

excitability matures in DA-independent (early) and DA-dependent (late) stages of which the latter is

likely to be regulated via extensive DA-mediated K$_v$ conductances.

Early transient electrophysiological deficits of corticostriatal neurons circuited with *GBA-N370S*

DaNs

GBA1 mutations are a major genetic risk factor for PD with the N370S (the c.1226A > G) variant as one

of the most common worldwide [32]. Multiple previous studies have proposed autonomous cellular

dysfunction of DaNs harbouring *GBA-N370S* with a growing literature suggesting failure of the

mitochondria and endolysosomal pathway [7,8,12]. However, it remains elusive as to whether and how

the failing mutant DaNs contribute towards physiological perturbation of the striatal output and the

eventual motor impairment in PD.

To dissect the impact of *GBA-N370S* iPSC-DaNs on the physiology of corticostriatal output neurons,

we established the micro-striatal circuit using iPSC-derived CNs and MSNs from healthy controls

connected with iPSC-DaNs from either healthy controls or PD patients carrying the *GBA-N370S*

mutation (Figure 5A). We compared the electrophysiological properties between corticostriatal

neurons in microcircuit with either healthy DaNs (DaN$^+$CN$^+$MSN) or *GBA-N370S* DaNs (*GBA-N370S*

DaN$^+$CN$^+$MSN). We found that while both groups exhibited increasingly hyperpolarised RMP over time,

the RMP of DaN$^+$CN$^+$MSN was always more hyperpolarised than that of *GBA-N370S* DaN$^+$CN$^+$MSN,

particularly statistically pronounced in early culture (DIV 40-45: $p = 0.0022$, DIV 55-60 $p = 0.0204$, DIV

70-75 $p = 0.252$) (Figure 5B). Similarly, a significant genotype effect was observed for rheobase current

at DIV 40-45, but not in later cultures (DIV 40-45: $p = 0.0143$, DIV 55-60 $p > 0.99$, DIV 70-75 $p = 0.319$)

(Figure 5C).

Neuronal excitability is tightly controlled by coordinated activity of both Na$_v$ and K$_v$. The inward flux of

sodium ions depolarises the cell membrane towards action potential threshold and triggers

spontaneous firing while outward currents generated K$_v$ underly resting membrane potentials and

action potential afterhyperpolarisation [30]. Hence, we next explored whether the altered intrinsic

excitability was coupled with relevant ionic channels. Unsurprisingly, we observed a substantial

reduction in current density of both Na_v and K_v among the *GBA-N370S* DaN$^+$CN$^+$MSN (DIV 40-45: Na_v

and slow activating K_v, DaN genotype p <0.0001; fast A-type K_v, DaN genotype p = 0.0001) (Figure 5Di,

Ei, Fi, and G). Recordings of iPSC-MSNs from intermediate and late cultures between DIV 55-70

revealed resolution of genotypic effects previously observed at DIV 40-45 (Figure 5D-F). These data

suggest an early pathological role of *GBA-N370S* mutation in iPSC-DaNs in altering striatal GABAergic

output via early hyperexcitability of post-synaptic iPSC-MSNs independent of overt degeneration of

dopaminergic afferents.

PKA antagonism resolved early and transient ionic current deficits in corticostriatal neurons

circuited with *GBA-N370S* iPSC-DaNs

The electrophysiological activity of many cellular channels, including voltage-gated sodium and

potassium channels, is robustly regulated by PKA-mediated signalling pathway in response to

fluctuations of external stimuli [33,34]. In MSNs both $K_v4.2$ and $K_v1.2$, the principal channel contributing

towards fast A-type and slowly inactivating potassium current, respectively, as well as Na_v, are

negatively modulated by PKA activation [35–38] To examine whether PKA activity is involved in the early

deficits of corticostriatal neurons affected by *GBA-N370S* iPSC-DaNs, we applied the cell-permeable

PKA inhibitor H89 directly to iPSC-MSNs in the striatal chambers of the striatal microcircuit and

performed whole-cell patch clamping of iPSC-MSNs at DIV 40-45 (Figure 6A and B). PKA activity assay

data suggested an intrinsic doubling of PKA activity in iPSC-MSNs circuited with *GBA-N370S* iPSC-DaNs

(i.e. *GBA-N370S* DaN$^+$CN$^+$MSN), as compared to those with healthy control-derived iPSC-DaNs (i.e.

DaN$^+$CN$^+$MSN) (DaN$^+$CN$^+$MSN vs *GBA-N370S* DaN$^+$CN$^+$MSN p = 0.0426) (Figure 6C). The increase in PKA

activity in the iPSC-MSNs was then reduced to the control level upon administration of 10 μM H89 to

iPSC-MSNs for 15 mins (DaN$^+$CN$^+$MSN vs *GBA-N370S* DaN$^+$CN$^+$MSN + 10 μM H89 p > 0.99, *GBA-N370S*

DaN$^+$CN$^+$MSN vs *GBA-N370S* DaN$^+$CN$^+$MSN + 10 μM H89 p = 0.05) (Figure 6C). These data

demonstrated the phenotypic increase in PKA activity in MSNs synapsed with *GBA-N370S* DaNs and

the ability of H89 to normalise striatal PKA activity *in vitro*. DaN lysates were collected from the DaN

chambers from the same devices as the healthy control-derived striatal lysates, and a significant

reduction in PKA activity was seen in *GBA-N370S* DaNs (i.e. MSN$^+$CN$^+$*GBA-N370S* DaN) relative to

healthy control-derived DaNs (i.e. MSN$^+$CN$^+$DaN) (MSN$^+$CN$^+$DaN vs MSN$^+$CN$^+$*GBA-N370S* DaN p =

0.0179) (Figure 6D). This deficit in PKA activity in *GBA-N370S* DaNs remained in the mutant *GBA-N370S*

DaNs synapsed with iPSC-MSNs treated with H89 (p = 0.0373 relative to MSN$^+$CN$^+$DaN), confirming

the fluidic isolation of different neuronal compartments. Treatment of iPSC-MSNs with H89 resolved

the genetic effect of *GBA-N370S* DaNs on MSNs Na$_v$ and K$_v$ current density at DIV 40-45 to healthy

control levels (Figure 6E-G), indicating the involvement of PKA in mediating intrinsic excitability of

corticostriatal neurons via ionic conductances.

Discussion

The present study represents the first attempt at recapitulating an *in vitro* neuronal circuit model of

the human striatal synaptic triad using an open chamber-based microfluidic platform to reveal the

physiological importance of connectivity-driven inputs on human striatal functions. Our data suggest

an early nonautonomous striatal electrical dysfunction in MSNs in the presence of the *GBA-N370S*

mutation within presynaptic DaNs that was associated with diminished striatal PKA activity.

Interestingly, in this model striatal neurons showed dynamic cell-autonomous PKA-dependent

modulation of ionic channel physiology as illustrated by the resolution of early striatal deficits by

specific antagonism of striatal PKA activity, despite the persistent presence of mutant DaNs.

Previous strategies to investigate human striatal neuron (patho)physiology remain largely limited in

monocultures of a single neuronal type [20], with more recent success in developing a neuronal network

architecture [16]. To date, there are no cellular models that capture the complex triad of non-autologous

synapses onto striatal neurons which are critical for the emergence of functional characteristics of

300 MSNs. Indeed, although our MSNs in monoculture displayed a temporal marker expression profile

301 reminiscent of the developmental trajectory reported from the Human Brain Transcriptome

302 (https://hbatlas.org), combined cortical glutamatergic and dopaminergic inputs further facilitated

303 functional maturation of the MSNs beyond the benefits of long-term culturing. Our results

304 corroborated the relevance of cortical glutamate in promoting both inhibitory and excitatory synaptic

305 maturity of MSNs [39] as well as the time-sensitive influence of dopaminergic transmission on MSNs

306 excitability maturation previously reported in rodent models [23]. It should also be noted that the

307 glutamatergic input onto MSNs originates from not only the cortex but also the thalamus which makes

308 up ~40% of glutamatergic synapses on MSN dendrites and converges with corticostriatal afferents

309 indiscriminately onto both MSN subtypes [40].

311 A significant highlight of this open microfluidic-based circuit approach is its customisability. The

312 modular nature of this setup allows great flexibility for reconstructing the compartmentalisation of

313 different neuronal populations and the directionality of their specific connectivity. Hence, virtually any

314 microcircuit can be established applying a similar design concept. Targeted manipulation to specific

315 compartments is also permitted with great ease as exemplified by our striatal PKA treatment

316 paradigm. Immunostaining and electrophysiological results indicate that our approach has captured

317 the oriented connectivity resembling that *in vivo* and that functional glutamate and dopamine

318 transmission exists.

320 Because the ATP competitive analogue H89 produced similar amplification of both Na_v and K_v in our

321 study, the observed elevation of striatal PKA activity likely occurred upstream of the cAMP in the PKA

322 signalling cascade, potentially through excessive Dopamine 1 receptor (D1R) activation followed by

323 stimulation of adenylyl cyclase-cAMP cascade [41]. The D1R antagonist SKF83566 has been shown

324 previously to decrease PKA activity in direct pathway MSNs [42]. Voltage clamp work has confirmed that

325 D1R stimulation reduced Na_v currents by limiting channel availability [43] while lessening K_v peak current

amplitude through altering channel inactivation kinetics [38]. Furthermore, the D1R agonist SFK81297

reversibly reproduced the reduction in Na_v and K_v current amplitude, similarly through PKA-mediated

phosphorylation of channel subunits, and improved striatal firing probability [44,45].

The majority of our culture was composed of direct pathway MSNs (dMSNs) expressing mainly

substance P or pro-dynorphin (Supplementary Figure 2) implicating a potential role for D1R.

Considering that the striatal excitability in the absence of dopaminergic projections at DIV 40-45

(Figure 4Ei, Fi, and Gi) did not phenocopy the early striatal deficit synapsed with *GBA-N370S* DaNs at

the same time point (Figure 5Di, Ei, and Fi), we speculate that increased, rather than reduced, tonic

DA content at the nigrostriatal synapses is a more logical explanation of D1R over-stimulation. This

hypothesis aligns with previous work on other models of PD-related familial mutations reporting early

cell-autonomous hyperexcitability of midbrain DaNs prior to overt neurodegeneration [46–48] which

could result in increased PKA activity in dMSNs [42], but contrasts with a single report on *GBA-N370S-*

associated reduced spontaneous firing and basal DA release by Woodard and colleagues [49] in

monocultured DaNs.

Epidemiological data suggests that only about 9.1% of *GBA1* mutation carriers will develop PD, often

with a relatively earlier age of onset and modest motor impairment as compared to non-carriers [32].

This is consistent with the early changes in striatal function observed in our study which was

subsequently resolved at later timepoints. As the compensation was likely to be intrinsic to non-

mutant MSNs, we propose that *GBA1* mutation-driven deficits in DaNs emerge early but in themselves

are insufficient to induce long-term dysfunction in the motor circuit regulated by the basal ganglia.

This evidence of potential compensatory capacity is in keeping with the late age of disease

presentation in PD. It is interesting to speculate that MSNs might be able to compensate for PD

disease-driving deficits elicited by DaNs, thereby masking initial disease stages and the prodrome. A

functional motor study of reversible PKA modulation *in vivo*, would be an interesting experiment to

further test these compensation effects in a PD mouse model that was not yet symptomatic for motor

impairments. While major therapeutic efforts have been channelled toward reversing DaN pathology,

our results suggest that rescue of the striatal deficits is an attractive alternative to preserve the

integrity of the basal ganglia output. If compensation is indeed functionally implicated in genetic

disease as our data suggests, then therapies targeting MSNs might be able to provide patients with

symptomatic relief. Future studies could mimic the full genetic landscape of *GBA-N370S* carriers by

deriving all three neuronal populations of the microcircuit from the same *GBA-N370S* cell lines.

Pairwise comparisons of striatal function from microcircuits in which the *GBA1* mutation burden is

carried by different, all, or none of the component cell types mixed with healthy controls could

delineate contributing pathophysiological roles of different neuronal populations of the striatal

presynaptic circuitry to the net debilitating burden of the *GBA1* mutation.

We have demonstrated a PKA activity-related mechanism by which mutant *GBA-N370S* DaNs induced

early hyperexcitability of healthy control-derived iPSC-MSNs preceding neurodegeneration. We also

observed a reduction in PKA activity in *GBA-N370S* iPSC-DaNs relative to healthy control-derived DaNs

(Figure 6D) which is consistent with the previously reported downregulation of PKRACB transcript

expression from DaN monocultures harbouring the same genetic burden [8]. Mutation of the PKA

regulatory subunit *PRKAR1B* has also been reported in a novel familial neurodegenerative condition

with parkinsonism and dementia [50], and *Prkar1b*-knockout rats exhibit body tremor [51]. Moreover,

reduced PKA activity in DaNs has been implicated in PD pathophysiology in other PD-related rodent

mutation models such as *LRRK2* R1444C/G/H [52,53] and *PINK1* [54]. Likewise, increased PKA activity was

thought to be protective against PD [42,55]. However, the exact mechanistic link between *GBA1*

mutations and PKA activity in nigral neurons remains to be further investigated. Nevertheless, it is well

documented that the large network of lysosome genes including *GBA1* is moderated by a master

regulator transcription factor EB (TFEB) [56] which in turn is the downstream substrate of PKA-mediated

phosphorylation via mTORC1 [57]. Considering our previous observation indicating dysregulation of

lysosomal/autophagic signalling in monocultured DaNs carrying *GBA-N370S* [7,8], we suggest that the

multiple cellular perturbations in nigral DaNs burdened by the *GBA-N370S* mutation might be initiated

by reduced activity of the master kinase PKA which disrupts various associated neuronal functions

potentially via lysosomes and/or lysosomal-related organelles. In addition, PKA activation is closely

related to proteasomal activation [58,59], which is a necessary function often compromised in PD [60,61]. It

is also interesting to speculate that reduced PKA activity could alter protein homeostasis through

reduced proteasomal function, and so, in turn, cause cellular dysfunction. From a preferential

vulnerability point of view, protected cells like MSNs might be able to find compensation mechanisms

and recover from these effects, while vulnerable cells such as DaNs may not be able to [62].

PKA was thought to positively modulate neurotransmitter release in most synapses via a combination

of factors including phosphorylation of various proteins involved in synaptic vesicle exocytosis [63-65]. *In*

vivo studies have suggested PKA phosphorylation-mediated increase in DA release via activation of TH

enzymatic activity [66,67]. The involvement of PKA in neurotransmitter release suggests that mutant

GBA-N370S may affect DA striatal content, and that the pathophysiology of *GBA-N370S* variant-

mediated PD may result from a disruption of the tight modulation of PKA activity.

In conclusion, we have described an open microfluidic-based neuronal microcircuit model which has

used iPSC-derived neurons to recapitulate *in vitro* the presynaptic network relevant to *in vivo* striatal

neurons. This approach allowed us to explore emerging electrophysiological features of human MSNs

over time in both the healthy and diseased state. We envision greater use of this modular circuit-

based model in addressing sophisticated functional questions of human physiology and pathology.

400
401

Method details

1.   Acquisition and maintenance of iPS cell lines

iPS cell lines used in this study were derived from human skin biopsy fibroblast acquired with informed

consent and ethical approval (Ethics committee: National Health Service, Health Research Authority,

NRES Committee South Central, Berkshire, UK, REC 10/H0505/71). The iPS cell lines have been

previously characterized: 3 control iPSC lines SFC067-03-01 (RRID:CVCL_RD75), SFC156-03-01 (EBiSC

Cat# STBCi101-A, RRID:CVCL_RD71) [8], and SFC856-03-04 (RRID:CVCL_RC81), 3 *GBA-N370S* PD lines

MK082-26 (RRID:CVCL_IJ04) [8]; MK088-01 (EBiSC Cat# UOXFi003-A), and MK071-03 (EBiSC Cat#

UOXFi001-B) [7]. All iPSCs were maintained on Matrigel™ (Corning)-coated plates under feeder-free

culture condition with mTSER1™ (StemCell Technology) supplemented with 1%

penicillin/streptomycin (P/S) (Life Technologies). Medium was changed daily, and cultures were

passaged 1:2 every other day using TrypLE Express™ (Life Technologies) when they reached 90-100%

confluency in the presence of Rock inhibitor (Bio-Techne).

2.   Generation of iPSC-derived MSNs

3 iPSC control lines (SFC067-03-01, SFC156-03-01, and SFC856-03-04) were differentiated into MSNs

using condition modified from previously established protocols [14,15]. In brief, iPSCs were plated at

75,000 cells/cm^2 and maintained until 85% confluence. Culture was switched to neural induction

medium containing DMEM/F12 (Life Technologies), 1% MEM Non-Essential Amino Acids (NEAA), 1%

Glutamax, 2% B27 without vitamin A (all from ThermoFisher), 1% P/S, 1 µM LDN193189 (Sigma),

10µM SB431542 (Abcam) and 4 µM XAV (Tocris). On DIV 04, culture was pre-incubated with ROCK

Inhibitor (Y-27632) for 1 hour before being passaged in 1:2 ratio. Culture was passaged in 1:2 on DIV

8 and switched to differentiation medium containing DMEM/F12, 1% P/S, 1% NEAA, 2 mM L-glutamine

(Life Technologies), 2% B27, 200 nM LDN193189, and 4 µM XAV. Activin A was added to from DIV 12

till DIV 23.

On day 16, the culture was either cryopreserved for future experiments or replated at 65,000 cells/cm^2

onto freshly coated plates or coverslips. Coating included 10 and 100 μg/ml Poly-D-Lysine (Sigma-

Aldrich) overnight for plastic plates and glass coverslips, respectively, followed by 1:100 Matrigel™ for

1 hour. Neuronal progenitor cells were cultured in the maturation medium SynaptoJuice® as

previously reported [15]. Cultures were treated with 1 μM AraC on DIV 18 to arrest proliferation of non-

neuronal cells capable of DNA synthesis. AraC concentration in the culture was gradually diluted with

subsequent media changes.

3.   Generation of iPSC-derived CNs

Generation of CNs from 3 iPSC control lines ((SFC067-03-01, SFC156-03-0, and SFC856-03-04) was

adapted from a previously detailed protocol [26]. In brief, neural development was induced with dual

SMAD inhibitors 10 μM SB431542 (Tocris) and 118 nM LDN (Sigma). Following formation of a dense

neuroepithelial sheet, the whole sheet was enzymatically lifted using Dispase (Life Technologies) and

broken down into smaller clumps to encourage development of neural rosettes. Rosettes were

maintained and expanded on laminin-coated wells till DIV 30 at which point cultures were either

cryopreserved or dissociated into single cells for further differentiation. At DIV 30 cortical

neurogenesis was induced by doxycyline-dependent overexpression of *NGN2* via co-transduction of

LV-TetO-mNgn2-T2A-Puro and LV-Ubiq-rTA lentiviruses into the cultures. Transduced cells were

incubated for 2 days before being selected with fresh media supplemented with 1 μg/ml puromycin

for another 48 hours prior to final replating.

4.   Generation of iPSC-derived DaNs.

All 3 control and 3 *GBA N370S* PD iPS cell lines were differentiated into DaNs as previously described

[8] with added modifications [68]. In brief, iPSCs were dissociated into single cells, plated on Geltrex-coated

plastic plates at 125,000 cells/cm^2 and cultured till confluence. iPSCs were then patterned toward

floor-ventral midbrain precursors for 10 days. Day 10 precursors were expanded with weekly

passaging for the subsequent 20 days. These cells can be used immediately or cryopreserved. Post-

expansion DIV 29 precursors were further differentiated and matured into post-mitotic midbrain

neurons until DIV 38 for final replating onto microfluidic devices.

5. Immunocytochemistry

Cultures were fixed with 2% PFA for 20 mins at room temperature (RT), rinsed with PBS, and antigen

retrieved in citrate buffer pH 6.0 (ThermoFisher) in water bath at 80ºC for 5 mins. Samples were left

to rest for 10 mins at room temperature before being permeabilised and blocked in PBS, 10% donkey

serum and 0.01% of Triton X-100 for 10 mins. Incubation of primary antibody in PBS and 10% donkey

serum was performed overnight at 4ºC. Samples were washed with PBS and subsequently incubated

in species-appropriate Alexa Fluor© secondary antibody in PBS with 10% donkey serum for 1 hour at

RT. The list of primary and secondary antibodies used for immunostaining was tabulated in

supplementary table 2 and 3, respectively. Images were acquired on the Opera Phenix High-content

Screening system (PerkinElmer) or the Invitrogen EVOS ™ FL Auto (ThermoFisher) cell imaging system

and subsequently processed on Harmony (Perkin Elmer; RRID:SCR_018809) or ImageJ

(RRID:SCR_003070), respectively.

6. Reverse transcription-qualitative polymerase chain reaction (RT-qPCR)

Total RNA of MSN cultures over time was extracted and purified from cell lysates using RNeasy Mini

kit (Qiagen) according to manufacturer's instructions. Extracted RNA was quantified and assessed on

a Nanodrop (DS-11, Denovix). RNA was reverse transcribed into cDNA using superscript III reverse

transcriptase kit (Life Technologies). qPCR was performed with standard SYBR green (ThermoFisher)

in StepOnePlus thermal cycler (Life Technologies). Full list of primers used are tabled in Supplementary

Table 1.

7.   Microfluidic fabrication.

A custom master mould containing both duo- and trio-chambers was manufactured by Microliquid

(Spain). The microfluidic device was made from the mould by pouring a silicon elastomer SylGARD

184, Dow Corning © pre-mixed with its curing agent in the 1:10 ratio. Polymerisation occurred when

incubating the PDMS-filled mould for 3 hours at 60ºC. Final devices were manually cut out from the

negative cast and washed with 100% ethanol. Cut-out devices were air-dried in tissue culture hood

and subsequently placed on ethanol –sterilised 19 mm-diameter coverslips to form an instantaneously

tight seal. Both the chambers and microchannel area were coated with poly-D-lysin (0.1 mg/ml)

overnight followed by Geltrex™ (ThermoFisher) prior to cell culturing.

8.   Electrophysiology

Whole-cell recordings of MSNs on coverslips were done at 27ºC with periodic change of external

solution following every new recording cell. The external solution contained: 2.4 mM KCl, 167 mM

NaCl, 10 mM Glucose, 10 mM HEPES, 1 mM $MgCl_2$, and 2 mM $CaCl_2$ adjusted to 300 mOsmol/l and pH

7.36 with NaOH. Neurons were patched with a borosilicate glass pipette (8-12 MΩ) pulled using a

Sutter P-97 Flaming Brown puller (Sutter Instrument Company) and filled with an internal solution

consisting of: 140 mM $C_6H_{11}KO_7$, 6 mM NaCl, 1 mM EGTA, 4 mM MgATP, 0.4 mM Na_3GTP, 10 mM

HEPES, and 0.01% Neurobiotin (SP-1120) adjusted to 290 mOsmol/l and pH 7.3 using KOH. Data were

acquired with MultiClamp 700B amplifier (Molecular Devices) and Digidata 1550B digitiser (Molecular

Devices) and analysed in Clampfit 10.7 (Molecular Devices). Cells were always clamped at -70 mV

unless otherwise stated. Access resistance was constantly monitored; cells with access resistance

greater than 50 MΩ were excluded. All recordings are made within two hours of coverslip preparation.

Cell capacitance and input resistance were automatically reported from Clampex membrane test

sampled at 33 kHz. Resting membrane potentials (RMP) were acquired immediately upon successful

break-in in current clamp with zero current injection. Liquid junction potential was estimated at 12mV

using JPCal [69] and adjusted for in RMP recordings.

Na_V and K_V currents were recorded in voltage clamp using a series of 400 ms square voltage steps of

10mV increments from -70 mV to +70 mV. Signals were sampled at 10 kHz and filtered at 2 kHz. Leak

subtraction was applied with 4 sub-sweeps and a settling time of 250 ms. Evoked action potentials

were recorded in current clamp by injecting 500 ms current steps in 10 pA increments from -10 pA to

+130 pA.

Spontaneous EPSC and IPSC were recorded in voltage-clamp mode with 20x gain at one another's

reversal potential i.e. -70 mV and 0 mV, respectively. The internal solution for spontaneous synaptic

activity recordings was Cs-based containing 140 mM $C_6H_{11}CsO_7$, 6 mM NaCl, 1 mM EGTA, 4 mM MgATP,

0.4 mM Na_3GTP, 10 mM HEPES, and 0.01% Neurobiotin adjusted to 290 mOsmol/l and pH 7.3 using

CsOH. All sEPSC and IPSC recordings were 1-4 mins in duration and low-pass Bessel 8-pole filtered

post-hoc. Spontaneous post-synaptic events and event magnitude were automatically detected based

on absolute magnitude difference from the baseline (>10 pA for sEPSCs and >15 pA for sIPSCs) using

threshold search function on Clampfit (RRID:SCR_011323). Putative EPSC or IPSC events were

excluded based on template-dependent criteria including rise time and half-width (<4 ms and <1.2 ms,

respectively) and manually validated to reject false-positive events.

H89 (Tocris) was dissolved in dH2O and were media applied to the middle chamber to the final

concentrations of 10 μM for 15 mins prior to wash off and recording.

All patched neurons were post-hoc labelled with Streptavidin-Alexa Fluor 488 conjugate

(ThermoFisher) and DARPP32 (Sigma) to ascertain the MSN identity. Only neurobiotinylated neurons

that co-expressed DARPP32 were selected for analysis.

9.   Dendritic spine visualisation and quantification

Patched coverslip of MSNs was immunolabelled for DARPP32 (as described above) and mounted onto

SlowFade ™ Diamond Antifade mountant (ThermoFisher). Detection and imaging of DARPP32-positive

Neurobiotin™-filled neurons were captured under Olympus FluoView FV1000 confocal microscope

with argon and solid-state laser with 488 nm and 559 nm excitation, respectively. Z-stacks of images

were sampled sequentially at resolution of 1024 * 1024 pixels, with 60x oil-immersion objective (NA

= 1.40), at 1.05 μm steps as optimised by Nyquist sampling theorem. Dendritic branches of

biotinylated neurons were captured at 3x software zoom. Images were segmented, reconstructed and

automatically rendered using filament and spine detection module in Imaris 9.6.0 (Bitplane, South

Windsor, CT, USA, RRID:SCR_007370) which identified dendrites and detect protrusions along the

dendritic filament length. Putative spines at branch points or disconnected dots were manually

excluded.

10. PKA activity assay

PKA activity were measured using the PKA Colorimetric Acitivity Kit (ThermoFisher) according to

manufacturer's guidelines. Protein concentrations of the same samples were quantified using

Pierce™BCA Protein Assay Kit (ThermoFisher) following manufacturer's instructions. Both plates were

read for absorbance in a PHERAstar® microplate reader (BMG Labtech). Protein concentration and

PKA activity values were inferred from their respective standard curves. Final PKA activity read-outs

were normalised against the sample protein concentrations.

11. Statistical analysis

All data was presented as mean ± Standard error of the mean (sem) unless otherwise stated. Raw data

was tested for normality and statistical comparison of the means was performed using unpaired

Student's t-test, two-way ANOVA with Bonferroni post-hoc test, or Kruskal-Wallis test or Friedman

test corrected with Dunn's multiple comparison test when appropriate. Significance difference was

considered at $p < 0.05$. All statistical analyses were performed on GraphPad Prism 6.0 (GraphPad

Software, RRID:SCR_002798).

Supplementary Table 1: Primer sequences

Gene	Forward primers	Reverse primers
PAX6	aggccagcaacacacctagt	agccagatgtgaaggaggaa
DLX2	gcctcaacaacgtcccttact	tcactatccgaatttcaggctca
GSX2	atgtcgcgctccttctatgtc	atgccaagcgggatgaagaaa
DLX5	ttcagaagactcagtacctcgc	gagttacacgccattgggtc
ASCL1	gtcctgtcgcccaccatctc	ccctcccaacgccactgac
MEIS2	gatgaaagagacggcagctcc	gggttgaggttgcatcatcg
NKX2.1	aggacaccatgaggaacagc	cccatgaagcgggagatg
FOXP1	ctaccgcttccatgggaaatc	ctgttgtcactaaggacaggg
GAPDH	ctgacttcaacagcgaccc	atgaggtccaccaccctgt
ISL1	caggttgtacgggatcaaatgc	cacacagcggaaacactcgat
CTIP2	atcctcagcccctttgttt	gccgttgttcctgaattgtt
GAD1	ccaaggtgctggactttcat	aaatcgaggatgacctgtgc
DARPP32	cctgaaggtcatcaggcagt	ggtcttccacttggtcctca
TH	tgaggagaaggaggggaagg	ccgggtctctagatggtgga
FOXP	ctaccgcttccatgggaaatc	ctgttgtcactaaggacaggg
EBF1	aatgtaagcaaggtggacgc	tcaaggtctaagccggacac
D1R	agggacatgtctttggcttcag	gggaacagtgttagcacctgtt
PDYN	caccacagcggactgcctgt	agcagggcagcctggcattg
D2R	ctggagaggcagaactggag	gaagacgatgacagcgatga
ADORA2A	aggagggctgtcaggtgaa	gttctggcagcagcatcat
PENK	atcctcgccaagcggtatg	ggttgtcccctgtttccaga

558
559

560   Supplementary table 2: Primary antibodies used

Antibody	Target	Host organism	Dilution	Source	Cat #	RRID	URL
Anti-CTIP2, monoclonal	CTIP2	rat	1:250	Abcam	Ab18465	RRID:AB_2064130	https://www.abcam.com/ctip2-antibody-25b6-ab18465.html
Anti-DARPP32, monoclonal	DARPP32	rabbit	1:250	Abcam	ab40801	RRID:AB_731843	https://www.abcam.com/darpp32-antibody-ep720y-ab40801.html
Anti-DARPP32, polyclonal	DARPP32	rabbit	1:250	Sigma-Aldrich	HPA048630	RRID:AB_2680468	https://www.sigmaaldrich.com/GB/en/product/sigma/hpa048630
Anti-GAD67, monoclonal	Glutamate decarboxylase1	mouse	1:250	Millipore	MAB5406	RRID:AB_2278725	https://www.merckmillipore.com/GB/en/product/Anti-GAD67-Antibody-clone-1G10.2,MM_NF-MAB5406
anti-Calbindin, Monoclonal	Calbindin	rabbit	1:250	Sigma-Aldrich	ab1778	RRID:AB_2068336	https://www.sigmaaldrich.com/GB/en/product/mm/ab1778
Anti-Substance P, monoclonal	Substance P	rat	1:250	Santa Cruz	sc21715	RRID:AB_628299	https://www.scbt.com/p/substance-p-antibody-nc1-34hl
Anti-PDYN, polyclonal	Prodynorphin	rabbit	1:250	Sigma-Aldrich	HPA053342	RRID:AB_2682121	https://www.sigmaaldrich.com/GB/en/product/sigma/hpa053342
anti-MAP2, Polyclonal	Microtubule-associated protein 2	chicken	1:250	Abcam	ab92434	RRID:AB_2138147	https://www.abcam.com/map2-antibody-ab92434.html
anti-TH, monoclonal	Tyrosine hydroxylase	mouse	1:250	Santa Cruz	sc-25269	RRID:AB_628422	https://www.sigmaaldrich.com/GB/en/product/mm/ab152
Anti-PKEN, polyclonal	Proenkephalin	rabbit	1:250	Sigma-Aldrich	HPA013138	RRID:AB_1855191	https://www.sigmaaldrich.com/GB/en/product/sigma/hpa013138

562 Supplementary table 3: Secondary antibodies used.
563

Antibody	Target	Host organism	Dilution	Source	Cat #	RRID	URL
Alexa-Fluor 555	IgG Mouse	Donkey	1:1000	Invitrogen	A-11057	RRID:AB_2534104	https://www.thermofisher.com/antibody/product/Donkey-anti-Goat-IgG-H-L-Cross-Adsorbed-Secondary-Antibody-Polyclonal/A-11057
Alexa-Fluor 555	IgG Rabbit	Donkey	1:1000	Invitrogen	A-31570	RRID:AB_2536180	https://www.thermofisher.com/antibody/product/Donkey-anti-Mouse-IgG-H-L-Highly-Cross-Adsorbed-Secondary-Antibody-Polyclonal/A-31570
Alexa-Fluor 648	IgG Rabbit	Donkey	1:1000	Invitrogen	A-31572	RRID:AB_162543	https://www.thermofisher.com/antibody/product/Donkey-anti-Rabbit-IgG-H-L-Highly-Cross-Adsorbed-Secondary-Antibody-Polyclonal/A-31572
Alexa Fluor 488	IgG Rat	Donkey	1:1000	Invitrogen	A-11006	RRID:AB_2534074	https://www.thermofisher.com/antibody/product/Goat-anti-Rat-IgG-H-L-Cross-Adsorbed-Secondary-Antibody-Polyclonal/A-11006
Alexa Fluor 488	IgG Chicken	Donkey	1:1000	Stratech Scientific Ltd	703-545-155-JIR-0.5mg	RRID:AB_2340375	https://www.jacksonimmuno.com/catalog/products/703-545-155
DAPI (4',6-Diamidino-2-Phenylindole, Dilactate)	DAPI	NA	1:1000	Thermo Fisher Scientific	D1306	RRID:AB_2307445 or RRID:AB_2629482	https://www.thermofisher.com/order/catalog/product/D1306
Streptavidin, Alexa Fluor™ 488 Conjugate	Neurobiotin	Donkey	1:1000	Thermo Fisher Scientific	S32354		https://www.thermofisher.com/order/catalog/product/S32354

Data Availability

The original data used in this study is available at Zenodo (DOI 10.5281/zenodo.7661355), and detailed

protocols of each method is available at protocols.io (DOI 10.17504/protocols.io.x54v9dw61g3e/v1).

Author Contributions

Q.D. designed, performed, and analysed all experimental data. N.I.A. and C.L. undertook iPSC-

dopaminergic neuron differentiations. B.N. optimised iPSC-cortical neuron differentiation. J.B.

provided expertise and assistance with fabrication of microfluidic chambers. N.B.V. and Q.D.

fabricated microfluidic chambers. N.B.V. and R.W.M conceived the project. N.B.V., R.W.M and

R.M.G. supervised the study. Q.D., N.B.V. and R.W.M. wrote the paper, with contributions from all

authors. All authors reviewed the manuscript and approved its submission.

Acknowledgements:

This research was funded in part by Aligning Science Across Parkinson's [ASAP-020370] through the

Michael J. Fox Foundation for Parkinson's Research (MJFF), and in part by the Monument Trust

Discovery Award from Parkinson's UK (J-1403). The work was supported by a National Institute for

Health Research-Medical Research Council Dementias Platform UK Equipment Award

(MR/M024962/1) to R.W.M. Q.D. was supported by a National Science Scholarship from Agency for

Science, Technology and Research in Singapore. For the purpose of open access, the author has applied

a CC BY public copyright license to all Author Accepted Manuscripts arising from this submission.

We thank Natalie Connor-Robson for her advice on MSN differentiation.

Competing interests

The authors declare no competing interests.

References

Tepper JM, Bolam JP. Functional diversity and specificity of neostriatal interneurons. *Curr Opin Neurobiol* 2004; **14**: 685–692.

Parker JG, Marshall JD, Ahanonu B, Wu Y-W, Kim TH, Grewe BF *et al.* Diametric neural ensemble dynamics in parkinsonian and dyskinetic states. *Nature* 2018; **557**: 177–182.

Cruz BF, Guiomar G, Soares S, Motiwala A, Machens CK, Paton JJ. Action suppression reveals opponent parallel control via striatal circuits. *Nature* 2022; **607**: 521–526.

Valsky D, Heiman Grosberg S, Israel Z, Boraud T, Bergman H, Deffains M. What is the true discharge rate and pattern of the striatal projection neurons in Parkinson's disease and Dystonia? *Elife* 2020; **9**. doi:10.7554/eLife.57445.

McGregor MM, Nelson AB. Circuit Mechanisms of Parkinson's Disease. *Neuron* 2019; **101**: 1042–1056.

Cui G, Jun SB, Jin X, Pham MD, Vogel SS, Lovinger DM *et al.* Concurrent activation of striatal direct and indirect pathways during action initiation. *Nature* 2013; **494**: 238–242.

Fernandes HJR, Hartfield EM, Christian HC, Emmanoulidou E, Zheng Y, Booth H *et al.* ER Stress and Autophagic Perturbations Lead to Elevated Extracellular α-Synuclein in GBA-N370S Parkinson's iPSC-Derived Dopamine Neurons. *Stem cell reports* 2016; **6**: 342–356.

Lang C, Campbell KR, Ryan BJ, Carling P, Attar M, Vowles J *et al.* Single-Cell Sequencing of iPSC-Dopamine Neurons Reconstructs Disease Progression and Identifies HDAC4 as a Regulator of Parkinson Cell Phenotypes. *Cell Stem Cell* 2019; **24**: 93-106.e6.

Nguyen HN, Byers B, Cord B, Shcheglovitov A, Byrne J, Gujar P *et al.* LRRK2 mutant iPSC-derived DA neurons demonstrate increased susceptibility to oxidative stress. *Cell Stem Cell* 2011; **8**: 267–280.

Zambon F, Cherubini M, Fernandes HJR, Lang C, Ryan BJ, Volpato V *et al.* Cellular α-synuclein pathology is associated with bioenergetic dysfunction in Parkinson's iPSC-derived dopamine neurons. *Hum Mol Genet* 2019; **28**: 2001–2013.

Ludtmann MHR, Angelova PR, Horrocks MH, Choi ML, Rodrigues M, Baev AY *et al.* α-synuclein oligomers interact with ATP synthase and open the permeability transition pore in Parkinson's disease. *Nat Commun* 2018; **9**: 2293.

Kim S, Wong YC, Gao F, Krainc D. Dysregulation of mitochondria-lysosome contacts by GBA1 dysfunction in dopaminergic neuronal models of Parkinson's disease. *Nat Commun* 2021; **12**: 1807.

Delli Carri A, Onorati M, Lelos MJ, Castiglioni V, Faedo A, Menon R *et al.* Developmentally coordinated extrinsic signals drive human pluripotent stem cell differentiation toward authentic DARPP-32+ medium-sized spiny neurons. *Development* 2013; **140**: 301–312.

Arber C, Precious S V, Cambray S, Risner-Janiczek JR, Kelly C, Noakes Z *et al.* Activin A directs striatal projection neuron differentiation of human pluripotent stem cells. *Development* 2015; **142**: 1375–1386.

Telezhkin V, Schnell C, Yarova P, Yung S, Cope E, Hughes A *et al.* Forced cell cycle exit and modulation of GABAA, CREB, and GSK3beta signaling promote functional maturation of induced pluripotent stem cell-derived neurons. *Am J Physiol Cell Physiol* 2016; **310**: C520-41.

Miura Y, Li M-Y, Birey F, Ikeda K, Revah O, Thete MV *et al.* Generation of human striatal organoids and cortico-striatal assembloids from human pluripotent stem cells. *Nat Biotechnol* 2020; **38**: 1421–1430.

Bengoa-Vergniory N, Faggiani E, Ramos-Gonzalez P, Kirkiz E, Connor-Robson N, Brown L V *et al.* CLR01 protects dopaminergic neurons in vitro and in mouse models of Parkinson's disease. *Nat Commun* 2020; **11**: 4885.

Baleriola J, Walker CA, Jean YY, Crary JF, Troy CM, Nagy PL *et al.* Axonally Synthesized ATF4 Transmits a Neurodegenerative Signal across Brain Regions. *Cell* 2014; **158**: 1159–1172.

Iannielli A, Ugolini GS, Cordiglieri C, Bido S, Rubio A, Colasante G *et al.* Reconstitution of the

Human Nigro-striatal Pathway on-a-Chip Reveals OPA1-Dependent Mitochondrial Defects
and Loss of Dopaminergic Synapses. *Cell Rep* 2019; **29**: 4646-4656.e4.

20 Virlogeux A, Moutaux E, Christaller W, Genoux A, Bruyere J, Fino E *et al.* Reconstituting
Corticostriatal Network on-a-Chip Reveals the Contribution of the Presynaptic Compartment
to Huntington's Disease. *Cell Rep* 2018; **22**: 110–122.

21 Chambers SM, Fasano CA, Papapetrou EP, Tomishima M, Sadelain M, Studer L. Highly
efficient neural conversion of human ES and iPS cells by dual inhibition of SMAD signaling.
Nat Biotechnol 2009; **27**: 275–280.

22 Gulacsi AA, Anderson SA. Beta-catenin-mediated Wnt signaling regulates neurogenesis in the
ventral telencephalon. *Nat Neurosci* 2008; **11**: 1383–1391.

23 Lieberman OJ, McGuirt AF, Mosharov E V, Pigulevskiy I, Hobson BD, Choi S *et al.* Dopamine
Triggers the Maturation of Striatal Spiny Projection Neuron Excitability during a Critical
Period. *Neuron* 2018; **99**: 540-554.e4.

24 Cheng P-L, Song A-H, Wong Y-H, Wang S, Zhang X, Poo M-M. Self-amplifing autocrine actions
of BDNF in axon development. *Proc Natl Acad Sci U S A* 2011; **108**: 18430–18435.

25 Turney SG, Bridgman PC. Laminin stimulates and guides axonal outgrowth via growth cone
myosin II activity. *Nat Neurosci* 2005; **8**: 717–719.

26 Hedegaard A, Monzón-Sandoval J, Newey SE, Whiteley ES, Webber C, Akerman CJ. Pro-
maturational Effects of Human iPSC-Derived Cortical Astrocytes upon iPSC-Derived Cortical
Neurons. *Stem cell reports* 2020; **15**: 38–51.

27 Freund TF, Powell JF, Smith AD. Tyrosine hydroxylase-immunoreactive boutons in synaptic
contact with identified striatonigral neurons, with particular reference to dendritic spines.
Neuroscience 1984; **13**: 1189–1215.

28 Smith Y, Bennett BD, Bolam JP, Parent A, Sadikot AF. Synaptic relationships between
dopaminergic afferents and cortical or thalamic input in the sensorimotor territory of the
striatum in monkey. *J Comp Neurol* 1994; **344**: 1–19.

29 Segal M, Greenberger V, Korkotian E. Formation of dendritic spines in cultured striatal
neurons depends on excitatory afferent activity. *Eur J Neurosci* 2003; **17**: 2573–2585.

30 Hogkin AL, Huxley AF. A quantitative description of membrane current and its application to
conduction and excitation in nerve. *J Physiol* 1952; **117**: 500–544.

31 Wilson CJ, Kawaguchi Y. The origins of two-state spontaneous membrane potential
fluctuations of neostriatal spiny neurons. *J Neurosci Off J Soc Neurosci* 1996; **16**: 2397–2410.

32 Riboldi GM, Di Fonzo AB. GBA, Gaucher Disease, and Parkinson's Disease: From Genetic to
Clinic to New Therapeutic Approaches. *Cells* 2019; **8**. doi:10.3390/cells8040364.

33 Jonas EA, Kaczmarek LK. Regulation of potassium channels by protein kinases. *Curr Opin*
Neurobiol 1996; **6**: 318–323.

34 Catterall WA. From ionic currents to molecular mechanisms: the structure and function of
voltage-gated sodium channels. *Neuron* 2000; **26**: 13–25.

35 Shen W, Hernandez-Lopez S, Tkatch T, Held JE, Surmeier DJ. Kv1.2-containing K+ channels
regulate subthreshold excitability of striatal medium spiny neurons. *J Neurophysiol* 2004; **91**:
1337–1349.

36 Tkatch T, Baranauskas G, Surmeier DJ. Kv4.2 mRNA Abundance and A-Type K+ Current
Amplitude Are Linearly Related in Basal Ganglia and Basal Forebrain Neurons. *J Neurosci*
2000; **20**: 579 LP – 588.

37 Scheuer T. Regulation of sodium channel activity by phosphorylation. *Semin Cell Dev Biol*
2011; **22**: 160–165.

38 Schrader LA, Anderson AE, Mayne A, Pfaffinger PJ, Sweatt JD. PKA modulation of Kv4.2-
encoded A-type potassium channels requires formation of a supramolecular complex. *J*
Neurosci 2002; **22**: 10123–10133.

39 Tu G, Sepers MD, Buren C, Parsons MP, Raymond LA. Influence of cortical synaptic input on
striatal neuronal dendritic arborization and sensitivity to excitotoxicity in corticostriatal

coculture. *J Neurophysiol* 2016; **116**: 380–390.

40   Doig NM, Moss J, Bolam JP. Cortical and Thalamic Innervation of Direct and Indirect Pathway
Medium-Sized Spiny Neurons in Mouse Striatum. *J Neurosci* 2010; **30**: 14610 LP – 14618.

41   Gerfen CR, Surmeier DJ. Modulation of striatal projection systems by dopamine. *Annu Rev*
Neurosci 2011; **34**: 441–466.

42   Ma L, Day-Cooney J, Benavides OJ, Muniak MA, Qin M, Ding JB *et al.* Locomotion activates
PKA through dopamine and adenosine in striatal neurons. *Nature* 2022; **611**: 762–768.

43   Surmeier DJ, Kitai ST. Chapter 20 D1 and D2 dopamine receptor modulation of sodium and
potassium currents in rat neostriatal neurons. In: Arbuthnott GW, Emson PCBT-P in BR (eds).
Chemical Signalling in the Basal Ganglia. Elsevier, 1993, pp 309–324.

44   Schiffmann SN, Lledo PM, Vincent JD. Dopamine D1 receptor modulates the voltage-gated
sodium current in rat striatal neurones through a protein kinase A. *J Physiol* 1995; **483 (Pt 1**:
95–107.

45   Kitai ST, Surmeier DJ. Cholinergic and dopaminergic modulation of potassium conductances in
neostriatal neurons. *Adv Neurol* 1993; **60**: 40–52.

46   Bishop MW, Chakraborty S, Matthews GAC, Dougalis A, Wood NW, Festenstein R *et al.*
Hyperexcitable substantia nigra dopamine neurons in PINK1- and HtrA2/Omi-deficient mice.
J Neurophysiol 2010; **104**: 3009–3020.

47   Carola G, Malagarriga D, Calatayud C, Pons-Espinal M, Blasco-Agell L, Richaud-Patin Y *et al.*
Parkinson's disease patient-specific neuronal networks carrying the LRRK2 G2019S mutation
unveil early functional alterations that predate neurodegeneration. *npj Park Dis* 2021; **7**: 55.

48   Subramaniam M, Althof D, Gispert S, Schwenk J, Auburger G, Kulik A *et al.* Mutant α-
Synuclein Enhances Firing Frequencies in Dopamine Substantia Nigra Neurons by Oxidative
Impairment of A-Type Potassium Channels. *J Neurosci* 2014; **34**: 13586 LP – 13599.

49   Woodard CM, Campos BA, Kuo S-H, Nirenberg MJ, Nestor MW, Zimmer M *et al.* iPSC-derived
dopamine neurons reveal differences between monozygotic twins discordant for Parkinson's
disease. *Cell Rep* 2014; **9**: 1173–1182.

50   Wong TH, Chiu WZ, Breedveld GJ, Li KW, Verkerk AJMH, Hondius D *et al.* PRKAR1B mutation
associated with a new neurodegenerative disorder with unique pathology. *Brain* 2014; **137**:
1361–1373.

51   Hoang Trung H, Yoshihara T, Nakao A, Hayashida K, Hirata Y, Shirasuna K *et al.* Deficiency of
the RIβ subunit of protein kinase A causes body tremor and impaired fear conditioning
memory in rats. *Sci Rep* 2021; **11**: 2039.

52   Muda K, Bertinetti D, Gesellchen F, Hermann JS, von Zweydorf F, Geerlof A *et al.* Parkinson-
related LRRK2 mutation R1441C/G/H impairs PKA phosphorylation of LRRK2 and disrupts its
interaction with 14-3-3. *Proc Natl Acad Sci U S A* 2014; **111**: E34-43.

53   Parisiadou L, Yu J, Sgobio C, Xie C, Liu G, Sun L *et al.* LRRK2 regulates synaptogenesis and
dopamine receptor activation through modulation of PKA activity. *Nat Neurosci* 2014; **17**:
367–376.

54   Dagda RK, Pien I, Wang R, Zhu J, Wang KZQ, Callio J *et al.* Beyond the mitochondrion:
cytosolic PINK1 remodels dendrites through protein kinase A. *J Neurochem* 2014; **128**: 864–
877.

55   Hwang R-D, Wiemerslage L, LaBreck CJ, Khan M, Kannan K, Wang X *et al.* The neuroprotective
effect of human uncoupling protein 2 (hUCP2) requires cAMP-dependent protein kinase in a
toxin model of Parkinson's disease. *Neurobiol Dis* 2014; **69**: 180–191.

56   Sardiello M, Palmieri M, di Ronza A, Medina DL, Valenza M, Gennarino VA *et al.* A gene
network regulating lysosomal biogenesis and function. *Science* 2009; **325**: 473–477.

57   Martina JA, Chen Y, Gucek M, Puertollano R. MTORC1 functions as a transcriptional regulator
of autophagy by preventing nuclear transport of TFEB. *Autophagy* 2012; **8**: 903–914.

58   Myeku N, Clelland CL, Emrani S, Kukushkin N V, Yu WH, Goldberg AL *et al.* Tau-driven 26S
proteasome impairment and cognitive dysfunction can be prevented early in disease by

activating cAMP-PKA signaling. *Nat Med* 2016; **22**: 46–53.

59   Lokireddy S, Kukushkin NV, Goldberg AL. cAMP-induced phosphorylation of 26S proteasomes
on Rpn6/PSMD11 enhances their activity and the degradation of misfolded proteins. *Proc*
Natl Acad Sci U S A 2015; **112**: E7176-85.

60   Bi M, Du X, Jiao Q, Chen X, Jiang H. Expanding the role of proteasome homeostasis in
Parkinson's disease: beyond protein breakdown. *Cell Death Dis* 2021; **12**: 154.

61   Thibaudeau TA, Anderson RT, Smith DM. A common mechanism of proteasome impairment
by neurodegenerative disease-associated oligomers. *Nat Commun* 2018; **9**: 1097.

62   Alegre-Abarrategui J, Brimblecombe KR, Roberts RF, Velentza-Almpani E, Tilley BS, Bengoa-
Vergniory N *et al.* Selective vulnerability in α-synucleinopathies. *Acta Neuropathol* 2019; **138**:
681–704.

63   Cho RW, Buhl LK, Volfson D, Tran A, Li F, Akbergenova Y *et al.* Phosphorylation of Complexin
by PKA Regulates Activity-Dependent Spontaneous Neurotransmitter Release and Structural
Synaptic Plasticity. *Neuron* 2015; **88**: 749–761.

64   Lonart G, Schoch S, Kaeser PS, Larkin CJ, Südhof TC, Linden DJ. Phosphorylation of RIM1alpha
by PKA triggers presynaptic long-term potentiation at cerebellar parallel fiber synapses. *Cell*
2003; **115**: 49–60.

65   Nagy G, Reim K, Matti U, Brose N, Binz T, Rettig J *et al.* Regulation of releasable vesicle pool
sizes by protein kinase A-dependent phosphorylation of SNAP-25. *Neuron* 2004; **41**: 417–
429.

66   Wu J, Filer D, Friedhoff AJ, Goldstein M. Site-directed mutagenesis of tyrosine hydroxylase.
Role of serine 40 in catalysis. *J Biol Chem* 1992; **267**: 25754–25758.

67   Campbell DG, Hardie DG, Vulliet PR. Identification of four phosphorylation sites in the N-
terminal region of tyrosine hydroxylase. *J Biol Chem* 1986; **261**: 10489–10492.

68   Fedele S, Collo G, Behr K, Bischofberger J, Müller S, Kunath T *et al.* Expansion of human
midbrain floor plate progenitors from induced pluripotent stem cells increases dopaminergic
neuron differentiation potential. *Sci Rep* 2017; **7**: 6036.

69   Barry PH. JPCalc, a software package for calculating liquid junction potential corrections in
patch-clamp, intracellular, epithelial and bilayer measurements and for correcting junction
potential measurements. *J Neurosci Methods* 1994; **51**: 107–116.

Figure 1: Generation and characterisation of iPSC-derived MSNs. (A) Differentiation schematic of medium spiny neurons (MSNs) from iPSC. (B-D) Temporal expression of selected markers of LGE progenitors (B), MSN progenitors (C) and post-mitotic MSNs (D) from DIV 0 till DIV 35. (E) Immunostaining for MAP2 (yellow), CTIP2 (green), DARPP32 (red), GAD67 (yellow) and DAPI (blue) and (F) quantification of their expression in percentage term at DIV 40-45. (G) Representative traces of evoked APs (iAP) showing slow-ramp depolarisation of MSNs. Blue arrow indicates delayed first spikes. (H) Whole-cell patch-clamp recording and subsequent post-hoc identification of neurobiotin-filled DARPP32+ neurons for electrophysiological analysis. (I-K) (I) Resting membrane potential (RMP), (J) cell capacitance, and (K) input resistance of iPSC-derived MSNs over time from DIV 40-75, n = 20-43 recording MSNs per condition, each dot represents a single neuron, One-way ANOVA corrected with Bonferroni post-hoc test, **$p<0.01$, ns $p>0.05$. (L) Number of MSNs that displayed evoked action potential upon current injection (iAP) and MSNs that did not (no iAP) over time. All data is presented as mean ± sem, N = 3-4 differentiation experiments of 3 iPS cell lines unless otherwise stated.

792
Figure 2: *In vitro* recapitulation of directed striatal presynaptic triad using microfluidic-based devices
(A) Schematic of custom microfluidic device which mimics orientation of the cortico-striato-nigral
circuit. (B) Side view schematics of fluid and growth factor-driven culturing paradigm of the cortico-
striato-nigral microcircuit. (C) Experimental design schematic showing the culturing workflow from
iPSC stage to reconstruct the directional connectivity of striatal presynaptic triad. (D) Representative
image showing compartmentalisation of (Di) cortical neurons (CNs), (Dii) MSNs and (Diii) dopaminergic
neurons (DaNs) in their respective chamber and uni-directional traversing of axons through
microchannels.

Figure 3: Coculturing MSNs with iPSC-derived cortical neurons improved synaptic activity and dendritic spine morphology of iPSC-derived MSNs. (A) Schematic illustrating the generation and whole cell patch-clamp recording of iPSC-derived MSNs in monoculture (CN⁻MSN) or in co-culture with cortical neurons in microfluidic devices (CN⁺MSN). (B-D) (B) Resting membrane potential (RMP), (C) rheobase current, and D) input resistance of CN⁻MSN and CN⁺MSN, n = 12-42, 7-27, and 12-42 recording MSNs per condition, respectively. (E-H) Quantification of spontaneous inhibitory post-synaptic current (sIPSC) (E) frequency (F) and magnitude over time from DIV 40-75, n = 10-37 recording MSNs per condition. (I) Representative traces of IPSC of CN⁻MSN and CN⁺MSN. (G-H) Quantification of spontaneous excitatory post-synaptic current (sEPSC) (G) frequency and (H) magnitude of CN⁻MSN and CN⁺MSN over time from DIV 40-75. N = 9-21 recording MSNs per condition. (I) Representative trace of EPSC of CN⁻MSN and CN⁺MSN. (K) Pair-wise comparative quantification between CN⁻MSN and CN⁺MSN from DIV 40-75, each dot represents a recording neuron, n = 11-29 recording MSNs per condition, unpaired two-tailed Student's t-test. (L) Representative images of dendritic spines in neurobiotin-filled DARPP32+ MSNs. All data is presented as mean ± sem, N = 3 differentiation experiments of 3 iPS cell lines, 2-way ANOVA with Bonferroni post-hoc correction test unless otherwise stated. *p<0.05, **p<0.01ßß

Figure 4: Dopaminergic neurons enhanced the intrinsic maturation of iPSC-derived MSNs during maturation via maintenance of high potassium conductances. (A) Schematic illustrating the generation and whole cell patch-clamp recording of corticostriatal neurons in the presence or absence of DaNs, i.e. DaN⁻CN⁺MSN and DaN⁺CN⁺MSN, respectively, on microfluidic devices. (B-D) (B) Resting membrane potential (RMP), (C) rheobase current, and (D) input resistance over time, n = 9-35 recording MSNs per condition. (E-G) Current density curves of (E) voltage-gated sodium channels, (F) fast A-type voltage-gated potassium, and (G) slow activating voltage-gated potassium channels over time, n = 19-30 recording MSNs per condition. (H) Representative fast A-type and and slow activating K_v currents of DaN⁻CN⁺MSN and DaN⁺CN⁺MSN at DIV 70-75. All data is presented as mean ± sem, N = 2 differentiation experiments of 3 iPS cell lines, 2-way ANOVA with Bonferroni post-hoc test, *p<0.05, **p<0.001.

847
Figure 5: Early but transient current deficits and altered excitability of MSNs circuited with *GBA-N370S*
DaNs. (A) Schematic illustrating the generation and whole cell patch-clamp recording of DaN+CN+MSN
and *GBA-N370S* DaN+CN+MSN. (B-C) (B) Resting membrane potential (RMP), and (C) rheobase current
of DaN+CN+MSN and *GBA-N370S* DaN+CN+MSN over time, n = 5-38 recording MSNs per condition. (D-
F) Current density curves of (D) voltage-gated sodium channels, (E) fast A-type voltage-gated
potassium, and (F) slow activating voltage-gated potassium channels over time. (G) Representative
differential fast A-type and and slow activating K_v currents of DaN+CN+MSN and *GBA-N370S*
DaN+CN+MSN at DIV 40-45. All data is presented as mean ± sem, N= 2-3 differentiation experiments
of 3 iPS cell lines, n = 27-32 recording MSNs at DIV 40-45, n = 14-16 recording MSNs per condition at
DIV 55-60 and DIV 70-75. 2-way ANOVA with Bonferroni post-hoc test, ns p>0.05, *p<0.05, **p<0.01,
***p<0.001, ****p<0.0001.

870
Figure 6: PKA antagonist rescued early current deficits of MSNs circuited with *GBA-N370S* DaNs. (A)
Schematic demonstrating antagonistic mechanism of H89 in PKA-mediated phosphorylation signalling
cascade. (B) Schematic illustrating treatment of PKA antagonist H89 on MSNs in the striatal
presynaptic microcircuit; C = catalytic subunit, R = regulatory subunit, cAMP = Cyclic adenosine
monophosphate, ATP = Adenosine triphosphate, ADP = Adenosine diphosphate. (C-D) Quantitative
measure of PKA activity of (C) DaN⁺CN⁺MSN, *GBA-N370S* DaN⁺CN⁺MSN and *GBA-N370S* DaN⁺CN⁺MSN
treated with PKA antagonist H89, Kruskal-Wallis test with Dunn's correction test, and (D) healthy
DaNs, mutant *GBA-N370S* DaNs synapsed with healthy untreated MSNs or mutant *GBA-N370S* DaNs
synapsed with H89-treated MSNs in their corresponding devices; Friedman test with Dunn's correction
test. N = 2 differentiations of 3 iPS cell lines each, n = 3 technical replicates; values were measured by
PKA colorimetric activity kit and normalised against total protein level extrapolated from BSA assays.
Each unit per ml (U/ml) was defined as the amount of PKA needed to mediate the transfer of 1.0pmol
phosphate from ATP to substrate at 30°C as per manufacturer's instructions. (E-G) Current density
curves of (E) Na_v, (F) fast A-type K_v, and (G) slowly activating K_v of DIV 40-45 MSNs. All data were
presented as mean ± sem, N = 2 differentiation experiments of 3 iPS cell lines, n = 17-22 recording
MSNs per condition. 2-way ANOVA with Bonferroni post-hoc test unless otherwise stated, ns $p > 0.05$,
*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$

Supplementary figure 1: Activin A promoted striatal development. (A) Relative transcript abundance
of selected LGE and MSN progenitor markers at DIV14 when cells were treated with activin A at DIV
12 normalised against those that were not; dotted line represents fold difference of 1. (B) Percentages
of post-mitotic neurons co-expressing DARPP32 and CTIP2 at DIV 40 in the presence or absence of
Activin A from DIV 12-23. N = 2-3 differentiation, n = 3-4 iPS cell lines, and 3 technical replicates.
893

Supplementary figure 2: Cultures contained a mixture of both direct and indirect MSNs. (A-B)
Transcript abundance of selected marker of (A) direct MSNs and (B) indirect MSNs over time.
Expression was averaged from 2 differentiations of 3 independent iPS cell lines. (C) Immunostaining
for GAD67 (green), SUBSTANCE P (red), PKEN (yellow), and PDYN (magenta) at DIV 50 (D) Proportion
of GAD67+ cell co-expresses different MSN subtype-specific neurotransmitter at DIV 50. All data is
presented as mean ± sem, N = 2 differentiation experiments of 3 iPS cell lines.
901

916 Supplementary Figure 3: Dopaminergic transmission mediated temporal changes of K_v current density.
917 (A-B) Temporal profile of sodium and potassium ionic currents of cortico-striatal neurons in the (A)
918 present and (B) absence of dopaminergic input.

919