Investigation of Volatile Metabolites in Sebum as Prodromal Indicators of Parkinson's Disease

Caitlin Walton-Doyle[1], Beatrice Heim[2], Eleanor Sinclair[1], Katherine A Hollywood[1], Joy Milne[1], Evi Holzknecht[2], Ambra Stefani[2], Birgit Högl[2], Klaus Seppi[2], Monty Silverdale[3], Werner Poewe[2], Perdita Barran[1], Drupad K Trivedi[1]*

[1]Manchester Institute of Biotechnology, Department of Chemistry, University of Manchester, Manchester, M1 7DN, UK.

[2]Department of Neurology, Innsbruck Medical University, Innsbruck, Austria.

[3]Department of Neurology, Salford Royal Foundation Trust, Manchester Academic Health Science Centre, University of Manchester, Manchester, UK, M6 8HD.

*Corresponding author

Email: drupad.trivedi@manchester.ac.uk

Abstract

Background: Parkinson's Disease (PD) has been associated with a distinct odour, strongest in sebum-rich areas. Thermal Desorption – Gas Chromatography – Mass Spectrometry (TD-GC-MS) has revealed volatile signatures that distinguish individuals with Parkinson's Disease (PD) from healthy controls. Here, we applied the same method, including subjects with isolated REM sleep behaviour disorder (iRBD) to examine the volatiles in sebum and compare this with that found in PD subjects and control participants. Participants with iRBD have a high likelihood for conversion to overt clinical synucleinopathies like PD, Dementia with Lewy Bodies (DLB) or (less commonly) Multiple System Atrophy (MSA).

Methods: Subjects with clinically established PD (n=16) or iRBD (n=9) as well as healthy controls (n=9) were included. Following methods established in our laboratory, sebum was sampled from each participant using cotton gauze and the headspace from these swabs, analysed directly with TD-GC-MS[1,2]. Univariate and multivariate analysis was employed to probe the differences between volatile metabolites found for each phenotype. Putative identifications were assigned using spectral matching against the Golm metabolome and NIST spectral databases.

Findings: We can completely classify each phenotype using the sampled volatilome from which we built models with logistic regression analysis. The classification between PD and control improved on previously published work, from 85% to 100%. Putatively annotated molecules include alkanes, aldehydes, fatty acid methyl esters (FAMEs), and three metabolites namely purine, tropinone and oleamide. Investigation of highly ranked features revealed 18 features that showed intermediate expression in samples from iRBD participants.

Interpretation: TD-GC-MS can differentiate volatile metabolite signatures from sebum between PD, RDB and control samples. More than 70% of the identifiable metabolites that

permit this discrimination were putatively annotated as hydrocarbons and fatty acid methyl esters (FAMEs). Our prior work indicates that these components arise from larger lipid molecules that decompose during the experiment[2]. Features putatively annotated as tropinone, oleamide and purine, have previously been linked with neuroprotection, sleep induction and antioxidation, respectively, are significantly different between the three groups of participants, along with FAMEs and hydrocarbons.

Funding: We thank Michael J Fox Foundation (grant ref:12921) and Parkinson's UK (grant ref: K-1504) for funding this study and Community of Analytical and Measurement Sciences (CAMS) for supporting DT's research position. This work was supported by the BBSRC (award BB/L015048/1) for instrumentation used in this work and by a DTA to CW-D (project ref. 2113640). We also thank our recruitment centres for their enthusiasm and rigor during the recruitment process. We are very grateful to all the participants who took part in this study as well as PIs and nurses at the recruiting centres.

Introduction

Parkinson's Disease (PD) is the second most common neurodegenerative disorder, with more than six million people affected worldwide and an expected two-fold increase in prevalence over the next generation[3]. The pathological hallmarks are neuronal α-synuclein positive inclusions and cell loss in the *substantia nigra,* other areas of the brain, and the peripheral autonomic nervous system causing the cardinal motor features of bradykinesia, rigidity and rest tremor as well as a plethora of non-motor symptoms[4]. Current diagnostic criteria for PD are anchored on clinical features, which overlap with other conditions and may result in initial misclassification[5]. Highly sensitive and specific tests, therefore, are a major need in PD - both to enhance diagnostic accuracy and to enable detection of the earliest disease stages where future disease-modifying therapies would be the most effective[5].

Clinically manifested PD evolves from a prodromal stage characterised by the occurrence of a variety of non-motor symptoms, including anosmia, constipation, anxiety, depression, change in cognitive behaviour as well as REM-Sleep Behaviour Disorder (RBD)[6,7]. Together with various risk markers, these have been used to define research criteria for prodromal PD. Within these criteria a polysomnographic diagnosis of isolated (iRBD), characterised by loss of REM-atonia and jerks or dream-enacting behaviours during REM sleep confers the highest risk for future PD with a likelihood ratio of 130[8]. Large prospective studies of cohorts with iRBD have shown conversion rates to clinically manifest synucleinopathies like PD, Dementia with Lewy Bodies (DLB) or Multiple System Atrophy (MSA) of more than 80% after follow-up for ten years or more[9]. Recent insights into prodromal features and risk markers for PD have led to the launch of population-based screening for PD risk, using multistep protocols where an initial online remote screen is followed by in-person tests, including dopamine transporter imaging[7].

Sebum is a lipid-rich substance secreted by the sebaceous glands on the skin, which are most concentrated on the face and upper back[10]. Increased sebum output (seborrhoea) and seborrheic dermatitis are well-known features of PD and changes in volatile profiles of sebum have recently been detected using Thermal Desorption – Gas Chromatography – Mass Spectrometry (TD-GC-MS)[1,2]. This method has so far not been employed on cohorts showing prodromal PD symptoms, which would be important to understand if volatile profiles of sebum have potential as biomarkers for early diagnosis. In this study, we analysed sebum samples from subjects with PSG-confirmed iRBD and compared findings to samples obtained from subjects with Parkinson's Disease (PD) and healthy controls in order to investigate differences in sebum composition and corresponding volatile metabolites profile. We describe how odour and mass spectrometry markers permit classification between the different phenotypes.

Odourous volatilome and metabolic markers can thus, be indicative of progression from iRBD to PD (Figure 1).

Figure 1: Schematic to show the processes used in this study and how they relate to the manifestation of disease phenotype. Healthy sebum biomarkers (Blue) are distinguished from iRBD (Green) and PD (Red) using both mass spectrometry and olfactory analysis. Mass spectrometer data allows us to classify phenotype from variables and determine the extent of true positives (ROC curve). By examining expression change of phenotypic features, we can determine markers that show the change from healthy to diseased. In parallel we investigated the odour from the sebum of each participant group and determined that each disease has a distinctive smell above that of healthy sebum, and remarkably that the change in odour could also be a progression indicator.

Sample collection and clinical cohort

Subjects were recruited at the Movement Disorder and Sleep Disorder out-patient clinics of the Dept. of Neurology of the Medical University Innsbruck, Austria. Site staff, patients' caregivers, and patients without neurodegenerative or dermatologic disorders were invited to participate as healthy controls. The study was reviewed and approved by the local ethics committee (EK number AN1979d 336/4.19) and all participants provided written informed consent. A total of 34 subjects were included (16 PD, 9 iRBD and 9 controls) and participant demographics are detailed in Table 1.

All samples were collected by the same individual to limit alterations in technique. Samples were collected on cotton-based gauze swabs (7.5 x 7.5 cm) that was wiped firmly on the upper back of participants, following the sampling protocol as described in earlier work[1,2]. Samples were transferred to airtight bags where they were sealed and stored at -80°C until analysis. After collection, samples were blinded and randomised for the analytical team at Manchester, UK until the data analysis and interpretation stage. Dynamic headspace was analysed from these samples in a randomised order. Raw data were deconvolved, annotated and converted to a data matrix consisting of 613 features and statistical interrogation of data was undertaken. The materials used, analytical method setup and data processing is described in supplementary information (S1, S2 and S3, respectively).

Table 1: Participants' characteristics at the time of recruitment. Nine PD participants and all iRBD participants had PSG recordings, with eight PD participants having PSG-confirmed iRBD. Two of the PSG-confirmed iRBD participants developed signs of Parkinsonism at clinical follow ups but did not show signs of Parkinsonism at the time of sampling. None of the controls showed any signs of Parkinsonism or clinical history of RBD at the time of sampling.

Phenotype	Age (mean ± SD)	Sex (M: F)	BMI (mean ± SD)	PSG confirmed RBD (*n*=)	Medication for PD (*n*=)	RBD-Medication (*n*=)
PD (*n*=16)	64.62 ± 6.38	3:1	25.78 ± 3.78	8	16 LDOPA: 14 DA: 11 MAO-BI: 9 amantadine: 5	4 clonazepam: 3 melatonin: 1
iRBD (*n*=9)	69.89 ± 11.83	8:1	27.93 ± 4.69	9	- DA agonists for RLS:1	3 clonazepam: 2 melatonin: 1
C (*n*=9)	60.44 ± 4.80	8:1	27.06 ± 3.61	0	-	-

C, controls; DA, dopamine agonist; LDOPA, Levodopa; MAO-BI, MAO-B inhibitor; PD, Parkinson's disease; RBD, REM sleep behavior disorder; RLS, restless legs syndrome.

Results and discussion

Classification of phenotypes

In order to classify between PD, iRBD and control, we applied a logistic regression model to the deconvoluted features from the mass spectrometry experiment which was able to distinguish each set of participants with 100% accuracy. The complete classification of PD and control samples validate and improve on our previously published findings[1,2]. Using this model and feature ranking method described in S3, we investigated top 25 features as our features of interest (*FoI*). Figure 2A shows clear separation and complete classification of phenotypes using only *FoI*. Multivariate receiver operator characteristic (ROC) curves show that binary classifier built using *FoI*, can completely distinguishing PD participants from iRBD as well as controls. Even when using just five *FoI* we can achieve AUC of 90% and 100% when modelling for PD against controls and PD against iRBD respectively (Figure 2B).

Figure 2: (A) *MDS plot showing clustering by phenotypes using FoI. Lines joining dots indicate which of the samples within groups were more similar than the rest, indicating common features in the disease phenotype and a far higher degree of dissimilarity in the control samples. (B) Multivariate ROC curve shows using the top five FoI, AUC=0.9 is obtained for PD vs Control classification. For both cases, using 15 or more FoIs (PD vs Control) and using five or more FoIs (PD vs iRBD) an AUC of 1 is obtained.*

Abundance of volatiles by phenotypes

Comparing means of these features across three phenotypes nine showed an increase from control to iRBD to PD and nine showed a decrease from control to iRBD to PD (Figure 2). These 18 features indicate iRBD as an intermediary phenotype between a control and a Parkinson's phenotype. There was one feature found to be highest in iRBD, six were lowest in iRBD (Supplementary S4). The *FoI* were putatively annotated as alkane (10), FAMEs (4), aldehyde (2), alkene (1), dialkyl ethers (1), purine (1) and tropinone (1). Four of the *FoI* could not be matched to library and thus, classed as unknown. Lack of identification of these species could be due to chromatographic co-elution, lack of precursor ions or lack of suitable match in the underivatized spectra in libraries – all preventative in assigning a biologically relevant annotation. Nonetheless, the complete classification of iRBD samples from control and PD, suggests that this disease has a distinct volatilome in terms of the abundance of identified features.

A sebum volatilome pattern of iRBD, that for most of its components, appears intermediate between that seen in controls and PD subjects is consistent with the notion of iRBD reflecting a prodromal state of PD. This consecutively lends further support to the sebum volatilome as a diagnostic biomarker for PD. This is best represented by viewing box plots of the identified *FoI* (Figure 3). We have plotted the *FoI* in terms of their different abundances in each participant group. In Figure 3A, we have shown compounds that increase in expression from control to PD, and in Figure 3B those that decrease from control to PD as well as some where the levels between PD and iRDB are not significantly different. It is unknown, if any of these are more sensitive than others to indicate phenoconversion from iRBD to PD or which of these changes earlier or after the phenoconversion. This will require further longitudinal studies. Also, since iRBD can evolve into Lewy Body disorders (PD and DLB) or MSA, the question remains which of the iRBD-associated features observed in this study are most

specific for conversion to either of these conditions. With our current sample size and lack of prospective data, it is not possible to comment if there are underlying volatilome signatures of iRBD (*i.e.*, within group variations or lack thereof seen in cluster analysis) that can be associated with conversion trajectory towards PD/DLB or MSA.

Figure 3: *Out of 25 FoI, 18 features had intermediate expression in the iRBD participants. These consisted of putatively annotated alkanes, aldehyde, FAMEs, tropinone and a purine species. Boxplots compare mean normalised intensities (black lines) between these phenotypes. Plots of features are ordered in their ranked importance. (A) compounds that increase in expression from control to PD (B) compounds that decrease from control to PD*

Interpretation of volatilome changes

Changes in fatty acid concentrations in brain tissue[11], blood[12] and the gut[13] have previously been associated with PD, and we have also reported alterations in lipids with fatty acid chains in sebum[14]. Analysing lipid standards with GC-MS gives rise to several resolvable features, identified as alkanes, as opposed to a single identifiable species as expected, which indicates that the ionisation process causes substantial fragmentation[2]. We hypothesise that the alkane *FoI* are derived from larger fatty acids and FAMEs in sebum. Fatty acids are a well-documented product of the breakdown of larger lipid molecules, and acylglycerols[15-17]. Similarly, aldehydes in headspace may arise due to oxidation of lipids, either endogenously or during the analysis[18].

Tropinone which was observed to decrease in iRBD and further decrease in PD, is an alkaloid derived from putrescine[19]. Studies in animal models have shown an important role of putrescine in skin physiology and neuroprotection[20]. Oleamide, an endogenous fatty acid was observed in the top 30 highly ranked features, and it had lower expression in the iRBD cohort compared to controls in this study. It is a sleep-inducing lipid, which has been found to decrease motility in animal models[21,22]. It reduces sleep apnoeic events and affects amount of REM and non-REM sleep in rat models[23,24]. Oleamide along with a few more alkanes and FAMEs were putatively annotated features that were not one of the *FoI*s due to lower ranking and contributed to only a marginal improvement to the classification model. Purine was observed to be lowest in control and highest in PD, with intermediate expression in iRBD. One of the end products of purine metabolism is urate - an antioxidant that is a risk marker for PD, and lower levels are also associated with higher severity of disease[25-27]. Alterations in the purine pathway have been researched in an investigation for PD biomarkers, as well as in comparison with other neurological dieases[28,29]. In iRBD, the intermediate expression of these molecules and

breakdown products discussed earlier, compared to PD and controls may indicate purine and tropinone along with certain classes of lipids may be markers of prodromal PD.

Interpretation of volatilome changes with regards to body odour

When presented with PD, iRBD and control gauze, Joy Milne, with her hypersensitivity to smell due to hereditary hyperosmia[30], was able to describe the presence of three distinct odours from gauze. The smell of PD as described earlier was thick, musky, that of iRBD was biscuity sweet and scent of control swabs was described as a person odour lacking any of the other smells. In this blind exploration, Joy was able to hint at mosaic presence or mix of PD and iRBD scent in some of the iRBD participants. Out of the three where hints of PD smell were described, two later showed signs of PD during clinical follow ups (Table 1). Based on these data, we further hypothesis that onset of Parkinsonism may alter body odour prior to detection by volatilome profiles measured by GC-MS or even current clinical symptoms (Figure 1) which implies a possibility of early detection of iRBD conversion to PD.

Limitations of the study

The key limitations of this study were smaller sample numbers which were unavoidable due to selective phenotypes and majority of participants were on medication as a treatment. The cohort was however, controlled for age and BMI to avoid confounding effects on the measurements. No strong associations were found between the *FoI* and known medications namely madopar, l-dopa, carbidopa, entacapone, ropinirole, rasagiline, apomorphine, amantadine, pramipexole, propranolol, clonazepame (rivotril), quetiapine, melotonin and rotigotine.

Figure 4: Pearson's correlation analysis between drugs and top 50 ranked features suggests very weak correlation (less than 0.6 and greater than -0.4) between drug intake and measured intensities. Each row refers to medication and each column is one of the top 50 ranked features. The colour of circle represents value of coefficient correlation R. The diameter of hollow circle represents narrow difference between lower and higher bounds of 95% confidence intervals (CI). The circles with no hollow inner circle represent big difference in higher and lower end of CI.

Although some associations were significant, they were not strong enough. This indicates the detected volatilome may not be contain directly measurable drug metabolites and other metabolites detected are not affected due to participant's medication history. Due to lack of non-derivatised volatilome library that covers endogenous metabolites, it was not possible to assign definite annotation which prevented meaningful metabolic pathway enrichment analysis. However, the results indicate there is a clear, discriminating volatilome signature that is present on skin swabs collected from all three cohorts.

Conclusion

This experiment has shown distinct volatilome signatures detectable from sebum in Parkinson's disease, iRBD and control phenotypes. We have shown a model that can classify with high certainty between these phenotypes using this volatilome signature. Upon further investigation of the significant features in the models, we have shown many metabolite signals to be the result of lipid degradation such as hydrocarbons and fatty acids. Thus, the detection of these maybe due to the alteration in lipid composition in sebum with the disease. The expression of some of these features in iRBD cohort is intermediate between that of PD and control. It can be speculated that change in expression of certain lipids present in human sebum and metabolites such as alkaloids and purine could be early indicators of pre-manifest PD in healthy controls as well as those suffering from iRBD.

References

1. Trivedi DK, Sinclair E, Xu Y, et al. Discovery of Volatile Biomarkers of Parkinson's Disease from Sebum. *ACS Central Science* 2019; **5**(4).
2. Sinclair E, Walton-Doyle C, Sarkar D, et al. Validating Differential Volatilome Profiles in Parkinson's Disease. *ACS central science* 2021; **7**(2): 300-6.
3. Collaborators GBDPsD. Global, regional, and national burden of Parkinson's disease, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. *The Lancet Neurology* 2018; **17**(11): 939-53.
4. Poewe W, Seppi K, Tanner CM, et al. Parkinson disease. *Nature Reviews Disease Primers* 2017; **3**(1).
5. Tolosa E, Garrido A, Scholz SW, Poewe W. Challenges in the diagnosis of Parkinson's disease. *The Lancet Neurology* 2021; **20**(5): 385-97.
6. Postuma RB, Berg D. Prodromal Parkinson's Disease: The Decade Past, the Decade to Come. *Movement Disorders* 2019; **34**(5): 665-75.
7. Mahlknecht P, Marini K, Werkmann M, Poewe W, Seppi K. Prodromal Parkinson's disease: hype or hope for disease-modification trials? *Translational neurodegeneration* 2022; **11**(1): 11.
8. Högl B, Stefani A, Videnovic A. Idiopathic REM sleep behaviour disorder and neurodegeneration - an update. *Nat Rev Neurol* 2018; **14**(1): 40-55.

9. Galbiati A, Verga L, Giora E, Zucconi M, Ferini-Strambi L. The risk of neurodegeneration in REM sleep behavior disorder: A systematic review and meta-analysis of longitudinal studies. *Sleep Med Rev* 2019; **43**: 37-46.

10. Thody AJ, Shuster S. Control and function of sebaceous glands. *Physiological Reviews* 1989; **69**(2): 383-416.

11. Fabelo N, Martín V, Santpere G, et al. Severe alterations in lipid composition of frontal cortex lipid rafts from Parkinson's disease and incidental Parkinson's disease. *Molecular medicine (Cambridge, Mass)* 2011; **17**(9-10): 1107-18.

12. Shin C, Lim Y, Lim H, Ahn TB. Plasma Short-Chain Fatty Acids in Patients With Parkinson's Disease. *Movement Disorders* 2020; **35**(6): 1021-7.

13. Unger MM, Spiegel J, Dillmann K-U, et al. Short chain fatty acids and gut microbiota differ between patients with Parkinson's disease and age-matched controls. *Parkinsonism Related Disorders* 2016; **32**: 66-72.

14. Sinclair E, Trivedi DK, Sarkar D, et al. Metabolomics of sebum reveals lipid dysregulation in Parkinson's disease. *Nature communications* 2021; **12**(1): 1592.

15. Smith RN, Braue A, Varigos GA, Mann NJ. The effect of a low glycemic load diet on acne vulgaris and the fatty acid composition of skin surface triglycerides. *Journal of Dermatological Science* 2008; **50**(1): 41-52.

16. Marzouki ZMH, Taha AM, Gomaa KS. Fatty acid profiles of sebaceous triglycerides by capillary gas chromatography with mass-selective detection. *Journal of Chromatography B: Biomedical Sciences and Applications* 1988; **425**: 11-24.

17. Shetewi T, Finnegan M, Fitzgerald S, Xu S, Duffy E, Morrin A. Investigation of the relationship between skin-emitted volatile fatty acids and skin surface acidity in healthy participants—a pilot study. *Journal of Breath Research* 2021; **15**(3): 037101.

18. Shahidi F. Headspace volatile aldehydes as indicators of lipid oxidation in foods. *Adv Exp Med Biol* 2001; **488**: 113-23.

19. Bedewitz MA, Jones AD, D'Auria JC, Barry CS. Tropinone synthesis via an atypical polyketide synthase and P450-mediated cyclization. *Nature communications* 2018; **9**(1): 5281.

20. Jänne J, Alhonen L, Keinänen TA, et al. Animal disease models generated by genetic engineering of polyamine metabolism. *Journal of cellular and molecular medicine* 2005; **9**(4): 865-82.

21. Boger DL, Sato H, Lerner AE, et al. Exceptionally potent inhibitors of fatty acid amide hydrolase: the enzyme responsible for degradation of endogenous oleamide and anandamide. *Proceedings of the National Academy of Sciences of the United States of America* 2000; **97**(10): 5044-9.

22. Roy A, Kundu M, Chakrabarti S, Patel DR, Pahan K. Oleamide, a Sleep-Inducing Supplement, Upregulates Doublecortin in Hippocampal Progenitor Cells via PPARα. *J Alzheimers Dis* 2021; **84**(4): 1747-62.

23. Carley DW, Pavlovic S, Janelidze M, Radulovacki M. Functional Role for Cannabinoids in Respiratory Stability During Sleep. *Sleep* 2002; **25**(4): 388-95.

24. Babson KA, Sottile J, Morabito D. Cannabis, Cannabinoids, and Sleep: a Review of the Literature. *Current Psychiatry Reports* 2017; **19**(4).

25. Chen X, Wu G, Schwarzschild MA. Urate in Parkinson's Disease: More Than a Biomarker? *Current Neurology and Neuroscience Reports* 2012; **12**(4): 367-75.

26. Weisskopf MG, O'Reilly E, Chen H, Schwarzschild MA, Ascherio A. Plasma urate and risk of Parkinson's disease. *American journal of epidemiology* 2007; **166**(5): 561-7.

27. Cipriani S, Chen X, Schwarzschild MA. Urate: a novel biomarker of Parkinson's disease risk, diagnosis and prognosis. *Biomarkers in medicine* 2010; **4**(5): 701-12.

28. Constantinescu R, Andreasson U, Holmberg B, Zetterberg H. Serum and cerebrospinal fluid urate levels in synucleinopathies versus tauopathies. *Acta Neurologica Scandinavica* 2012; **127**(2): e8-e12.

29. Koros C, Simitsi A-M, Papagiannakis N, et al. Serum uric acid level as a putative biomarker in Parkinson's disease patients carrying GBA1 mutations: 2-Year data from the PPMI study. *Parkinsonism Related Disorders* 2021; **84**: 1-4.

30. Morgan J. Joy of super smeller: sebum clues for PD diagnostics. *The Lancet Neurology* 2016; **15**(2): 138-9.