Title

Comparative metagenomics reveals host-specific functional adaptation of intestinal

microbiota across hominids

Authors

Rühlemann MC[1,2], Bang C[1], Gogarten JF[3,4,5,6], Hermes BM[7,8], Groussin M[1], Waschina S[9],

Poyet M[8], Ulrich M[4,5], Akoua-Koffi C[10], Deschner T[11], Muyembe-Tamfum JJ[12], Robbins MM[13],

Surbeck M[14,15], Wittig RM[16,17], Zuberbühler K[18], Baines JF[7,8], Leendertz FH[4,5], Franke A[1]

Affiliations

1 Institute of Clinical Molecular Biology, Kiel University, Kiel, Germany

2 Institute for Medical Microbiology and Hospital Epidemiology, Hannover Medical School

Hannover, Germany

3 Applied Zoology and Nature Conservation, University of Greifswald, Greifswald, Germany

4 Helmholtz Institute for One Health, Greifswald, Germany

5 Epidemiology of Highly Pathogenic Microorganisms, Robert Koch Institute, Berlin,

Germany

6 Viral Evolution, Robert Koch Institute, Berlin, Germany

7 Evolutionary Genomics, Max Planck Institute for Evolutionary Biology, Plön, Germany

8 Institute of Experimental Medicine, Kiel University, Kiel, Germany

9 Nutriinformatics Research Group, Institute for Human Nutrition and Food Science, Kiel

University, Kiel, Germany

10 Training and Research Unit for in Medical Sciences, Alassane Ouattara University /

University Teaching Hospital of Bouaké, Bouaké, Côte d'Ivoire

11 Comparative BioCognition, Institute of Cognitive Science, University of Osnabrück,

Osnabrück, Germany

12 National Institute for Biomedical Research, National Laboratory of Public Health,

Kinshasa, Democratic Republic of the Congo

13 Department of Primate Behavior and Evolution, Max Planck Institute for Evolutionary

Anthropology, Leipzig, Germany

14 Department of Human Evolutionary Biology, Harvard University, Cambridge,

Massachusetts, United States of America

15 Max Planck Institute for Evolutionary Anthropology, Leipzig, Germany

16 Institute of Cognitive Sciences, CNRS UMR5229 University Lyon 1, Bron Cedex, France

17 Taï Chimpanzee Project, CSRS, Abidjan, Côte d'Ivoire

18 Institute of Biology, University of Neuchatel, Neuchatel, Switzerland

Summary

Characterizing trajectories of the composition and function of hominid gut microbiota across diverse environments and host species can help reveal specific properties of the human microbiota, with possible implications for host evolution and health. Using shotgun metagenomic sequencing, we investigated taxonomic and functional diversity in the gut microbiota of wild-living great apes, including two gorilla subspecies (*Gorilla gorilla gorilla*, *Gorilla beringei beringei*), three chimpanzee subspecies (*Pan troglodytes verus*, *P.t. troglodytes*, *P.t. schweinfurthii*), and bonobos (*Pan paniscus*), together with human samples from Africa and Europe. We identified microbial taxonomic and functional adaptations convergent with host phylogeny at both the community and microbial genomic levels. We could show that repeated horizontal gene transfer and gene loss are processes involved in these adaptations. We hypothesize, that these adaptation processes and changes in the microbiome predispose the host to chronic inflammatory disorders, such as type 2 diabetes via altered histidine metabolism and inflammatory bowel disease indicated by adaptation of microbes to aerobic conditions. Additionally, we find multiple lines of evidence suggesting a widespread loss of microbial diversity and evolutionary conserved clades in the human microbiota, especially in the European population. Lastly, we observed patterns consistent with codivergence of hosts and microbes, particularly for the bacterial family *Dialisteraceae*, though we find that overall, co-phylogeny patterns are frequently disrupted in humans.

Introduction

Human gut microbiome research has demonstrated that numerous factors, including diet, environment, and lifestyle influence the structure of the human gut microbiota, which, in turn, has profound impacts on human health and disease (Johnson et al., 2019; Lloyd-Price et al., 2019; Vangay et al., 2018). To date, the majority of these studies have been conducted on sample collections from high-income countries and only recently, has interest shifted to include humans from diverse global populations, thereby providing an additional angle to investigate and evaluate shared and specific microbiome properties across human populations (McCall et al., 2020; Schaan et al., 2021; E. D. Sonnenburg and Sonnenburg, 2019; Vangay et al., 2018). These recent efforts provided an opportunity to discover microbiome signatures of host geography and lifestyle that go beyond differences in

microbiota diversity structures. For instance, they revealed elevated rates of horizontal gene

transfer (HGT) that correlate with the Human Development Index (HDI) of a population,

suggesting that gut microbiota constantly acquire new functions in conjunction with host

lifestyle changes (Groussin et al., 2021). However, the mechanisms that link changes in gut

microbial structure with host behavior and ecology remain largely unexplored.

Additional critical insight into understanding patterns of diversity and composition among

human gut communities can be obtained from comparative surveys of the hominid gut

microbiota. Humans, chimpanzees, bonobos, and gorillas show increasingly divergent gut

microbiota, with more distantly related species exhibiting more divergent community

composition (phylosymbiosis; (Brooks et al., 2016)). At the same time, the phylogeny of

some of their individual bacterial lineages parallels their own phylogeny (codivergence;

(Groussin et al., 2020; Suzuki et al., 2022)). Both patterns of phylosymbiosis and

codivergence are suggestive of long-term effects of hominid evolution on their communities

of symbionts (Groussin et al., 2017; Moeller et al., 2016; Ochman et al., 2010). Notably,

results from comparative marker-gene analyses suggest co-diversifying members of the

hominid gut microbial communities (both prokaryotic and phage) are lost and replaced with

human lineages, when animals leave their natural environments and are moved into

captivity (Gogarten et al., 2021; Nishida and Ochman, 2021). However, poor taxonomic

resolution and a lack of functional characterization precludes a deeper understanding of

processes driving these changes.

Functional analyses from shotgun metagenomic data revealed a conserved phylogenetic

signal across wild nonhuman primates (NHP), despite dietary changes over an individual's

lifespan and between species, suggesting that the evolution of gut microbiota within wild

NHP is partially constrained by host genetics and physiology (Amato et al., 2019). A

comparative meta-analysis of microbial functions in NHP and diverse human populations

observed a comparable loss of biodiversity in captive NHP and human populations from

regions with higher HDI (Manara et al., 2019), supporting previous findings (Moeller et al.,

2014). However, overall, the functional characterization of NHP microbiota in previous

studies has been limited to only selected microbial taxa without addressing broader scale

functional changes and specific alterations especially in African great apes and humans. As

such, robust comparative functional analyses are still needed for a comprehensive
understanding of how gut microbiota have evolved with hominids and shaped the current
structure and functional capabilities (and deficits) of the human gut microbiome.
105
Here, we present a large-scale comparative study of wild NHP and humans from
geographically distinct populations spanning two continents to better elucidate host-
microbiome interactions in the hominid gut, within an evolutionary context. To functionally
characterize hominid gut microbial communities, we performed shotgun metagenomic
sequencing from the feces of wild-living great apes, including two gorilla subspecies (*Gorilla*
gorilla gorilla, Gabon (GAB), n=8; *Gorilla beringei beringei,* Uganda (UGA), n=11), three
chimpanzee subspecies (*Pan troglodytes verus,* Côte d'Ivoire (CIV), n=55; *P.t. troglodytes*,
GAB, n=11; *P.t. schweinfurthii*, UGA, n=12), and bonobos (*Pan paniscus,* Dem. Rep. of the
Congo (DRC), n=12). Further, we included published metagenomic data from samples of *G.*
g. gorilla (n=28) and *P.t. troglodytes* (n=18) from the Republic of Congo (CG) (Campbell et
al., 2020). We additionally sequenced human fecal samples from a rural, remote village from
Côte d'Ivoire (n=12) and a rural community from the Democratic Republic of Congo (n=12),
jointly representing a spectrum of African populations with lower HDI; samples from a
German population were included in our analyses as a representative of populations with
high HDI.
121
Using this extensive data resource, we created a comprehensive catalog of high-quality
prokaryotic genomes assembled from metagenomic data, which we annotated on a
taxonomic and functional level. Using this catalog we explored patterns of diversity, host-
specificities in abundances of taxonomic groups and functions, and phylogenetic parallels
between host and microbiota on a community level, but also for individual microbial clades.
These analyses reveal intriguing patterns associated with human gut microbial communities,
convergent functional adaptations across lineages and potential mechanisms driving these.
129

An expanded catalog of microbial genomes from the hominid gut
From 203 fecal samples of humans and non-human primates with shotgun metagenomic
data, we reconstructed a total of 7,506 metagenome-assembled genomes (MAGs) ensuring
maximum completeness and low contamination using multiple binning algorithms and

dedicated curation and scoring tools ((Rühlemann et al., 2022); see Methods). The most

MAGs (quality score > 50%) were reconstructed for the most sampled subgroup of great

ape, *P.t. verus* (n=2,431), while the per sample on average the highest number of

reconstructed MAGs was the highest for *P.t. schweinfurthii* (mean=74.8). Library size /

number of sequencing reads was highly correlated with total assembly size ($\rho_{Spearman}$ =

0.644), which in turn was directly correlated with the number of bins recovered for a sample

($\rho_{Spearman}$ = 0.966).

Together with two large collection of reference species reconstructed from human fecal

metagenomes (UHGGv2; (Almeida et al., 2021)), and non-human primate fecal

metagenomes(Manara et al., 2019), MAGs were clustered into 5,648 species-level genome

bins (SGBs; 95% ANI) using stringent criteria (Supplementary Table 1). These included 1,047

novel SGBs not covered by any of the two large references, which mostly originated from

NHP samples (n=954, 91.1%; Suppl. Figure 1b). The highest-quality MAG in each SGB was

chosen as representative. Overall quality of SGB representative genomes was high (median

quality score = 94.5%; Suppl. Figure 1c). SGB representatives were used as comprehensive

reference for the estimation of per-sample abundances (Methods, Suppl. Figure 1d, Suppl.

Table 2). In total 2,934 SGBs, encompassing 21 bacterial and two archaeal phyla (Suppl.

Figure 1a), were found present in the dataset. This newly curated catalog of SGBs from

gorillas, bonobos, and chimpanzees increases the number of microbial species genomes

previously reconstructed from feces by more than ten-fold, and increases mapping success

of fecal metagenomes from NHPs to reference genomes by two- to three-fold (Suppl. Figure

1d; (Manara et al., 2019). Novelty in human fecal samples was comparably low, with 5.6%,

2.3 %, and 1.1% of novel SGBs found in samples from CIV, DRC, and GER, respectively (Suppl.

Figure 1d). For both NHPs and humans, the highest percentages of novel diversity were

observed within the phyla Bacteroidota and Spirochaetota, and, to a lesser degree, within

Firmicutes and Firmicutes A for NHPs only (Suppl. Figure 1e). Generally, recovered clusters

were highly host specific. While the 7,506 MAGs spanned 1,759 of the final SGBs, only for 48

of these SGBs MAGs were reconstructed from samples of more than a single host genus.

Within sample diversity varied considerably between host (sub-)species and with

sequencing depth (Suppl. Figure 1f). The used SGB collection covers large proportions of the

diversity found in the human gut (Suppl. Fig. 1d) and Faith's PD incorporates SGB

relatedness in the diversity calculation, which enables a better estimate of total diversity of

a community then simpler richness estimates from taxonomic group abundances.

Phylogenetic diversity (PD) at an even mapping depth of 1 Mio. reads per sample showed

significantly lower diversity in Humans compared to all African great ape hosts ($P_{Wilcoxon}$ = 6.9

x 10^{-14} ; Figure 1a). Comparing individual groups, human from GER and DRC showed lower

means than all NHP group (all $P_{Wilcoxon}$ < 0.05), while Humans from CIV exhibited high

variance and were found to have significant lower diversity than all NHP hosts ($P_{Wilcoxon}$ <

0.05), except for *G. g. gorilla* and *P.t. troglodytes* ($P_{Wilcoxon}$ > 0.05). These two great ape taxa

were found to have the lowest library sizes, low numbers of recovered genomes and lowest

mapping efficiency. Taken together, this suggests that phylogenetic diversity in these hosts

may be biased by lower representation in the reference database and that the reduced

diversity could be an artefact of this. Consequently, samples with less than 1 Million

mapped reads were removed from further analyses. This resulted in the removal of samples

from the analysis for the groups *G.b. beringei* (n=1), *G.g. gorilla* (n_{GAB}=5, n_{CG}=1), *P.t.*

troglodytes (n_{GAB}=4), *P.t. schweinfurthii* (n=1) and humans from CIV (n=1). Of note, lowest

mean values for within-sample phylogenetic diversity were found for the human subgroup

from Germany, which however were not significantly different from that of humans from

African communities (all $P_{Wilcoxon}$ > 0.05), contradicting previous reports (McCall et al., 2020;

Vangay et al., 2018). Additionally, the considerable differences in PD observed between CIV

and DRC highlight the diversity found between human populations and the need to better

characterize human diversity. Therefore, while total diversity of some host groups might not

be exhaustively sampled, especially considering lower abundant microbial clades, the

presented reference collection of high-quality metagenome reconstructed genomes likely

represents the current best resource for an in-depth taxonomic and functional assessment

of hominid fecal microbiomes and highlights that some human populations, including those

sampled in this analysis, have lost considerable microbial diversity in their guts.

Figure 1: Functional and taxonomic specificities of human and NHP fecal microbiome. (a) Phylogenetic Diversity across host groups at a sampling depth of 1 Mio. mapped reads per sample. (b) Ordination of unweighted UniFrac distances of all samples, colored by host subgroups. (c) Tanglegram of host (left) and microbiome (right) trees, the latter based on unweighted UniFrac distances. (d) SGB sharing coefficients between host group. Rows represent reference host groups, columns the groups they are sharing overlap with. Numbers in the tiles are P-values from the analysis for enrichment (↑) and depletion (↓) in the reference group. (e) Effect sizes of the abundance differences of genera in fecal samples of NHPs and humans (x-axis) and humans with African and European populations separated as well (y-axis). Points are colored as belonging to the association groups with European (dark blue) and African human populations (light blue), or as being generally enriched in NHPs (orange) or humans (mid-blue). (f) Per-sample and host group abundances of selected genera. (g) Abundance trajectories of taxa associated with human communities and NHPs (h) KEGG ortholog (KO) effect sizes in the test for differential abundance between African and European human population associated taxa. Shown are KOs in functional KEGG categories enriched among with differential abundance (Q<0.1). (i) Effect sizes (t-values) of the community-level abundance differences of KOs in fecal samples of NHPs and humans (x-axis) and humans from African and European (y-axis) (j) Effect sizes of community level KO abundance differences within KEGG modules significantly enriched in the NHP vs. human African and European population comparison. Grey points indicate non-significant values (Q>0.05).

Phylosymbiosis, a pattern in which microbial community divergence parallels that of the

hosts, can be a sign of community-level co-evolution of host and microbiota indicating a

close relationship over evolutionary timescales (Brooks et al., 2016). To investigate such

patterns, we used six different measures of beta diversity. Four were based on phylogenetic

or taxonomic distance metrics (weighted and unweighted UniFrac, as well as genus level

Aitchison and Jaccard distance), two on functional capacities of the community (KEGG

ortholog (KO) abundance and presence/absence patterns; Supplementary Figure 2). Strong signals for phylosymbiosis were found for Aitchison, Jaccard and unweighted UniFrac distances (all $P < 0.001$, Figure 1b and c, Suppl. Figure 2, Suppl. Table 3), but not for the function-based distances and the weighted UniFrac distance (all $P > 0.25$). These results suggest phylosymbiotic divergence in the general structure of hominid microbiota (Jaccard and unweighted UniFrac are based on presence/absence of microbial clades), while abundance patterns of distinct microbial clades and functions do not seem to follow this trend, which is in line with previous observations in other host systems (Brooks et al., 2016). This pattern might be explained as the result of fluctuating clade abundances and functions in response to changing environmental factors, including diet, indicating a high functional plasticity of the hominid gut microbiome in response to immediate influences, while structural changes through acquisition and loss of microbial clades are rather the result of longer-term adaptations.

Assessing shared SGBs between host groups, we found that especially human-associated SGBs were strongly depleted across multiple NHP groups (Figure 1d), while no such signal was found in the opposite direction, suggesting the widespread acquisition of novel microbial clades in the human intestinal microbiome. A similar pattern was found for *G. b. beringei* and multiple *Pan* subspecies, however not for *G.g. gorilla*. *G.g. gorilla* and *P.t. troglodytes* are sympatric species and were sampled in the same environment in the Republic of Congo and Gabon. The absence of excess strong host-specific signals between these groups might suggest effects of shared environmental factors influencing microbiome structure. SGBs from human samples from lower HDI populations were enriched in all human groups, indicating a shared conserved core microbiota, however no overrepresentation of SGBs from higher HDI populations were found in the two other groups, suggesting that specific SGBs were introduced to the human gut in association with lifestyle changes.

Fecal microbiome of high HDI populations marked by loss of evolutionarily conserved core microbiota

Abundance difference in microbial clades between humans and NHPs and between human communities with differing environments, such as living in rural or urban regions, in regions

of the world with lower or higher HDI, can give insights into microbiome-mediated
adaptations to environmental changes in the far and more recent past. We analyzed and
compared the abundance profiles of gut microbes between NHPs and humans, including
individuals living in rural, lower HDI areas of CIV and DRC as well as individuals residing in
urban, higher HDI regions within GER. For all following taxonomic and functional
comparisons, we restricted the analysis to human and *Pan* (chimpanzees and bonobos)
samples to obtain focused insights into the microbiota divergence since their hosts diverged
about 7-8 million years ago (Langergraber et al., 2012).

A total of 287 microbial genera were included in the analysis, of which 148 were found to be
differentially abundant ($Q_{Bonferroni}$<0.05, Figure 1d, Suppl. Table 4) in at least one of these
comparisons between human subgroups, or between humans and NHPs, and subsequently
sorted into one of four groups. We identified 30 taxa with increased abundances among
humans from Germany, such as *Akkermansia*, *Bacteroides*, and *Alistipes* (Figure 1d and e).
We additionally found 30 taxa that are enriched in the two African populations such as
Cryptobacteroides, Prevotella, and *Succinivibrio* (Figure 1e and f). Overall, the marker taxa of
European microbiomes that we detected are in agreement with previous findings (Jha et al.,
2018; Pasolli et al., 2019). Our approach allowed us to identify bacterial taxa that exhibit
differential abundance profiles between humans and NHPs that are independent from the
human population considered (Figure 1f). We found 63 taxa, such as *SIG603* that have
increased abundance in NHPs and 25 taxa, such as *Coprococcus*, with increased abundance
in humans. Interestingly, taxa depleted in the microbiome of European individuals are more
likely to be also abundant (>0.1%) in the microbiota of NHPs (P < 0.001; Figure 1g),
suggesting a loss of evolutionary conserved clades in these populations.

Widespread changes in fecal microbiome function between hosts and across human
communities
Taxon-specific changes reflect broad-scale differences between host-groups. A focused
analysis of microbial functions can give insights into the specific driving forces of such
community-level changes. We performed, analysis of abundance differences of 6,179 KEGG
orthologs (KOs, (Kanehisa and Goto, 2000)) in NHP (genus *Pan*) vs. human fecal microbiota
and humans in European and African societies and found significant abundance differences

in 2,030 (32.9 %) and 607 (9.82 %) KOs, respectively ($Q_{Bonferroni}$<0.05; Figure 1i, Suppl. Table

5). For the NHP vs. human comparisons, the KOs with the lowest Q-values included K20426

(acbU, 1-epi-valienol-7-phosphate kinase; Q = 6.03 x 10^{-48}) and K17723 (preA;

dihydropyrimidine dehydrogenase (NAD+) subunit PraA; Q = 2.25 x 10^{-41}) with higher

abundances in NHPs and K07267 (oprB; porin; Q = 2.11 x 10^{-50}) and K08646 (MEP; peptidyl-

Lys metalloendopeptidase; Q = 3.08 x 10^{-45}) with higher abundances in humans. K22336

(bfrB, bacterioferritin B; Q = 3.43 x 10^{-32}) and K17285 (SELENBP1, methanethiol oxidase; Q =

8.21 x 10^{-21}) were strongly increased in Germans. The strongest signals for humans with

African lifestyles were found for K12252 (aruH; arginine:pyruvate transaminase; Q = 1.01 x

10^{-12}) and K12078 (ptlA; type IV secretion system protein PtlA; Q = 1.24 x 10^{-12}).

Analysis of higher-level KEGG annotations including e.g. complete pathways

overrepresented among differentially abundant KOs revealed three annotations with

enrichment in NHPs compared to humans, and seven annotations differentiating humans in

Europe and Africa (Q_{FDR} < 0.05; Figure 1j, Suppl. Table 6). The enriched KEGG categories in

NHPs were "V/A-type ATPase, prokaryotes" and "ABC tansporters". ABC transporters are a

large family of ATP-dependent translocases (n_{KO}=377) with broad substrate spectra,

including e.g. carbohydrates, metal ions, amino acids, that are involved in nutrient uptake

(Davies et al., 2021) and export of toxic compounds (Davidson and Chen, 2004; Lee et al.,

2014). The signal is driven by a broad range of ABC transporters found with increase

abundance in NHPs (Suppl. Table 6), which might indicate adaptation to a diverse diet (Ejby

et al., 2016; Tsujikawa et al., 2021) or metabolism of plant-derived xenobiotics (Rodríguez-

Daza et al., 2021). Additionally, we found an enrichment of terms in "Microbial metabolism

in diverse environments" (map01120), an extensive pathway that serves as an umbrella

category for KEGG metabolic modules; thus, this signal can be interpreted as pointing to

substantial changes in the abundance of non-specific metabolic functions between NHPs

and humans.

Signals driven by differences between human subgroups included the KEGG module

"Helicobacter pylori pathogenicity signature, cagA pathogenicity island" (md:M00564),

driven by increased functional abundances in the fecal communities of humans living in

Africa. Individuals infected with CagA-positive H. pylori are at increased risk to develop

gastric cancer (Hatakeyama, 2014). Curiously, *H.pylori* was not found in the dataset,

however the KOs annotated for this pathway are found in the genus *Succinivibrio,* belonging

to the same phylum as *Helicobacter* (Proteobacteria), which is depleted in Europeans.

Whether *Succinivibrio* in the gut can under certain circumstances exhibit a higher risk for

other cancer types needs to be investigated.

In the European population, Vitamin B_{12} / Cobalamin biosynthesis (md:M00122, M00924,

M00925, and path:map00860) and "antimicrobial resistance genes" (br:ko01504) were

found to be enriched. Increased antimicrobial functions may be explained by a higher use of

antibiotics in human healthcare and animal husbandry, as well as by environmental

pollution (Groussin et al., 2021). The increased abundance of Vitamin B_{12}-producing

microorganisms in the feces of Europeans may be driven by higher dietary intake of meat

and dairy products from ruminants (Watanabe and Bito, 2018). These associations warrant

further experimental studies to identify their driving factors and to understand their

implications on the gut ecosystem.

Success of taxa associated to European populations relates to oxidative carbohydrate

metabolism

To elucidate the functional basis of the success of taxa in the human gut of individuals

residing within in Europe, we analyzed differences in the functional capacities of these in

comparison to other human associated taxa either enriched in African societies or without

association to this dichotomy. Compared to community-level functional analyses used to

identify higher-level functional patterns, pangenome analysis can specifically identify

individual genes and functions enriched or depleted in association with a trait of interest

within a specific microbial clade. Analyses were restricted to five bacterial families,

Acidaminococcaceae, *Bacteroidaceae,* *Lachnospiraceae,* *Oscillospiraceae,* and

Ruminococcaceae, for which sufficient numbers of SGBs (n>=20) for pangenome analysis

were recovered. SGBs in these family represent large proportions of the overall human

microbiome (mean = 61.7%), with no significant differences between the subgroups ($P_{Kruskal-Wallis}$ = 0.8). The analysis was conducted at the family-level, as higher taxonomic ranks would

increase clade-specific functional biases. To account for between-family functional

differences, functional differences of SGBs associated with Europeans were first analyzed

within families using Fisher's exact test and subsequently subjected to unweighted meta-
analysis using Z-scores to leverage shared signals.

We identified 276 enriched and 50 depleted KOs in pangenomes associated with the
European population ($Q_{Bonferroni}$<0.05; Suppl. Tables 7 and 8). Then, by using higher-level
KEGG annotations (modules, pathways, BRITE hierarchies), we found that 18 of these
annotations were overrepresented in the dataset ($Q_{Bonferroni}$ < 0.05, Figure 1h, Table 1, Suppl.
Table 9).

Among these were multiple groups involved in carbohydrate metabolism enriched in taxa
enriched in Europeans, specifically pointing at aerobic breakdown of sugar molecules for
ATP generation (Citrate cycle: md:M00009, path:map00020; Pentose phosphate pathway:
md:M00004; V/A-type ATPase, prokaryote: md:M00159). These signals strongly suggest that
the selection for taxa in the gut of humans living in Europe is connected to the adaptation to
transient microaerobic conditions in the gut environment, and to the use of oxidative
phosphorylation as a mean to release energy from nutrients, which is more efficient than
strictly anaerobic fermentation (Jurtshuk, 1996). However, reduced fermentation can
impact short-chain fatty acid production, which can, in turn, potentially negatively affect the
host's intestinal epithelial cells and metabolism (Deleu et al., 2021). Additionally, we found
multiple pathways connected to the metabolism of amino acids (Histidine metabolism:
path:map00340, md:M00026, md:M00045; Glycine, serine and threonine metabolism:
path:map00260). Especially gut microbial histidine metabolism has been discussed with
relevance to human health, as it was shown that an intermediary product of histidine
degradation, imidazole propionate, was increased in type 2 diabetic individuals in a large
study of almost 2,000 individuals, and that this increase was directly connected to the
microbiota and overall unhealthy dietary habits, however independent of dietary histidine
intake (Molinaro et al., 2020). This suggests a microbiome-driven pre-disposition to
metabolic disorders in European human communities.

We did not find any higher-level KEGG annotations significantly enriched among the taxa
not enriched in Europeans. the relative lower fitness of these taxa may not result from a
single mechanism associated to European lifestyles, but rather from multiple selective
forces specific to the individual clades.

Convergent host-specific adaptations are found across microbial families

Figure 2: (a) KO terms consistently enriched in human- (blue diamonds) and NHP-associated (orange diamonds). Individual
genus-level results are shown in points, colored by phylum. (b) Prevalence patterns of KO terms in human- (blue) and NPH-
associated (orange) Prevotella SGBs. Filled shapes represent KOs with significant (Q<0.05) differences in the statistical test.
(c) Tanglegram of phylogenetic tree of cydA-positive Prevotella and cydA-gene tree. Colored lines highlight common
subtrees. Both trees were rooted using Paraprevotella as outgroup.

Shared gene gains or losses across multiple microbial clades can indicate convergent

adaptations to specific host intestinal environments. We performed a pangenome analysis

of genera shared between humans and NHPs (n=37) to identify such patterns. To control for

higher-level clade effects, functional repertoires (KEGG Ontology terms) were compared at

the genus-level between SGBs from NHPs (Suppl. Table 10) and humans and then combined

in a meta-analysis across genera. In total, 123 KO terms were identified as carrying

signatures of convergent adaptation to the respective host group, with 44 of these

signatures associating with humans and 79 associating with NHPs ($Q_{Bonferroni}$<0.05; Figure 2a,

Suppl. Table S10). Among the human-associated KOs, we found multiple functional groups

hinting again at an adaptation to increased oxygen by utilization of oxygen as an electron

acceptor within the respiratory chain, such as cytochrome *bd* ubiquinol oxidase subunits

(cydA, cydB), as well as adaptation to increased oxidative stress through ferritin (ftnA) and

thioredoxin-dependent peroxiredoxin (BCP). Among the KOs enriched in NHP pangenomes

we found 1-epi-valienol-7-phosphate kinase (acbU), phosphoserine phosphatase (rsbU/P),

two KOs annotated as polyketide synthases (rhiA, pksN) and a phage terminase (xtmB).

Products produced by polyketide synthases have diverse functions, including antibiotic

activity, virulence and support of symbiotic relationships (Ridley et al., 2008).

Prevotella represented the largest genus-level clade in the dataset (n=113, 5.2% of all SGBs).

While we found this genus across all host species, it is largely decreased in Europeans. Thus,

we selected this genus for further analysis to elucidate potential functional mechanisms

driving the observed patterns. Enrichment analysis revealed 55 KOs with distinct prevalence

patterns (Q<0.05; Figure 2b). The most striking difference was found for the cytochrome *bd*

ubiquinol oxidases subunits 1 and 2 (cydA and cydB), which were ubiquitous in all 31

human-associated *Prevotella* SGBs, but present in only two out of 62 SGBs from NHPs. It is

important to note that these prevalence differences are not driven by single lineages within

the *Prevotella* genus that would have distribution ranges restricted to single hosts. Instead,

they are observed across multiple sibling clades spanning the entire phylogenetic tree of the

taxon. Cytochrome *bd* oxidases are involved in stress responses, most prominently in

transiently microaerobic environments (Giuffrè et al., 2014). Comparison of the *Prevotella*

species tree (reconstructed using all SGB representatives recovered from the dataset (n =

191; including UHGGv2)) and the cydA gene phylogeny exhibit widespread incongruencies

between the tree topologies (Figure 2c). We performed a tree reconciliation using a

duplication-transfer-loss (DTL; (Kundu and Bansal, 2018)) model between the *Prevotella* and

cydA phylogenies, which revealed a high rate of gene transfer ($\bar{T}$ = 32.227) and loss ($\bar{L}$ =

35.769) driving the evolution of the cydA gene family. The gene transfer events and

mappings were robust across 1,000 reconciliations with different starting seeds, with 94% of

all events and 88% of all mappings found with 100% consistency. These results suggest that

the enrichment of cytochrome *bd* ubiquinol oxidases observed in humans compared to

NHPs are the result of multiple individual events of gene loss and horizontal gene transfer

within the hominid gut.

Co-phylogeny is rare in spore-forming microbes and disrupted in humans

Patterns of co-phylogeny between host and microbes can result from close interaction, or even interdependence in extreme cases, and congruent metabolic pathways from co-evolutionary trajectories. Using stringent selection criteria, we subjected 222 subtrees of the SGB phylogeny for co-phylogeny analysis (see Methods). The subtrees spanned 45 families and 961 (32.8%) SGBs present in the dataset in addition to 90 SGBs from the UHGGv2 catalog. We used a Mantel-test based framework and permutation to detect co-phylogeny signals ((Hommola et al., 2009), see Methods for details). When defining co-phylogeny candidates based on a mean P-value < 0.05 across all permutations, 53 of 222 subtrees (23.9%) qualified as exhibiting co-phylogenetic patterns (Suppl. Table S11). These

subtrees cover 303 of the 1051 SGBs (28.8%) included in the analysis and 5.7% of the total

5,316 hominid SGBs (excluding the SGBs from Manara et al.). All results and subtrees can be

inspected online (https://mruehlemann.shinyapps.io/great_apes_shiny_app). By visually

inspecting subtrees with co-phylogeny signals, we find many candidates microbial

phylogenies that do not follow host phylogeny, *e.g.,* in a subgroup of the genus

Cryptobacteroides (Figure 3a). Such signals suggest that co-phylogeny within the *Gorilla* and

Pan clades can result in statistically significant Mantel tests, despite topological

incongruences of human-derived sequences, for which no host sister (sub-)species from the

same genus is available.

Human-derived genomes were found in 160 (72.1%) of the tested subtrees, of which 35

(21.9%) were co-phylogeny candidates. In comparison, 18 of 62 (29%) tested subtrees

without human-derived representatives exhibited co-phylogeny signals, *e.g.,* W0P29-013

spp. (Figure 3a). Overall, 16.3% (n = 66 out of 404) of human-derived SGBs in the analysis

were found in co-phylogenetic subtrees, which is significantly less compared to 34.6% (n =

227 out of 656) of NHP-derived SGBs ($P_{Fisher} = 6 \times 10^{-11}$). Co-phylogeny signals, defined based

on the ratio of SGBs in trees exhibiting co-phylogeny patterns and the total SGBs in the

family, differed strongly. Six families showed excessive signals of co-phylogeny and four

families a significant depletion ($Q_{Fisher, FDR} < 0.05$; Figure 3b, Suppl. Table S12). The strongest

co-phylogeny ratio was found for the family *Dialisteraceae* (Cophyl-Ratio = 61.22%, Q = 4.44

$x 10^{-7}$), which in the analysis were entirely represented by members of the genus *Dialister*.

Dialister are a common, but rather neglected members of the gut microbiota which have

been found increased (Vals-Delgado et al., 2021; Zheng et al., 2018) and decreased

(Joossens et al., 2011) with various human diseases, hence their relation to human health

remains unclear. However, on species, *Dialister invisus,* was found to be moderately

transmissible between human mother-infant pairs and within households in a large meta-

analysis (Valles-Colomer et al., 2023). The strongest depletions were found for *e.g.* the

families *Lachnospiraceae* and *Treponemataceae*, the former confirming previous results for

this clade (Moeller et al., 2016). The latter, *Treponemataceae*, especially the genus

Treponema D, were found depleted in humans living in Germany and occur in anaerobic

sediments (Thingholm et al., 2021), serving as (intermediary) reservoirs for transmission to

humans and NHPs, which can disrupt co-phylogenetic signals by constant re-introduction to
the community.

When comparing host groups, the proportion of SGBs with co-phylogeny signal is
significantly reduced in all human subgroups compared to the NHP hosts (all $Q_{Wilcoxon} < 0.05$;
Figure 3c). Further, humans from Germany exhibited even lower proportions of co-
phylogeny SGBs compared to the two African human populations ($Q_{Wilcoxon} < 0.05$). The
human subgroups from Africa did not differ in their co-phylogeny proportions ($P_{Wilcoxon} =$
0.583). These results clearly suggest a loss of wild great ape associated clades with
"humanization" of the intestinal microbiota and the introduction of novel microbial partners
with changing environment and lifestyle, confirming again findings from previous analyses.

Previous analyses suggested that spore-forming clades are less likely to be exhibiting co-
phylogenic patterns, due to their ability to survive outside of the gut, facilitating dispersal
between hosts (Groussin et al., 2020; Moeller et al., 2016). We find a negative correlation
between the number of sporulation-associated genes and family-level cophylogeny-ratio (P
$< 1.4 \times 10^{-4}$, Figure 3d, Suppl. Table S13), confirming this hypothesis, also supporting
previous findings of increased direct host-to-host transmission of non-spore forming
microbial clades in baboons (Tung et al., 2015).

Although not enriched, we found multiple (n=10 out of 28) subgroups in *Bacteroidaceae*
with co-phylogeny signals, consistent with previous findings based on *gyrB* amplicon data
(Moeller et al., 2016). However, no evidence for strict co-phylogeny was found in
Bifidobacteriaceae, which is inconsistent with findings from the same previous report
(Moeller et al., 2016). Comparing the phylogeny of metagenome-derived *gyrB* sequences
and the GTDB marker-gene phylogeny for *Bifidobacterium spp.* revealed clear incongruences
between both approaches (Figure 3e), which can likely explain the differences in the
presented analysis and previous findings.

Discussion
Here, we present the largest curated dataset of fecal metagenomes derived from wild
African great apes and human populations. For this, we surveyed and reconstructed high-

quality microbial genomes from the feces wild nonhuman primates, including gorillas

(*Gorilla gorilla gorilla*; *Gorilla beringei beringei*), chimpanzees (*Pan troglodytes verus*; *P.t.*

troglodytes; *P.t. schweinfurthii*), and bonobos (*Pan paniscus*) as well as human populations

from Africa and Germany. We identified signals of phylosymbiosis across the included

hominids, indicating a conserved evolutionary relationship of microbial communities with

their host species. Moreover, by employing a comparative approach, we found extensive

changes of microbial taxonomic and functional abundances across the intestinal microbiota

of NHPs and humans. Previous studies have pointed to Western lifestyles as an important

factor influencing the intestinal microbiota in humans. Within our human sample

population, we were able to confirm differential signals of prokaryotes associated with the

German human population. Importantly, using a comparative dataset of great ape taxa

showed that microbial clades lost in the German population in comparison to African human

populations are also found in wild great ape populations. Thus, we suggest that the loss of

these taxa might be regarded as a mass extinction event of evolutionary conserved

members of hominid-associated gut microbiota. While it is tempting to link these changes to

industrialization (as previous studies have done), there are many differences between these

human populations (e.g., genomic diversity, diet, exercise, sunlight exposure, exposure to

antibiotics, population bottlenecks) and it was certainly not possible with the sampling

regime here, to determine the particular factors responsible for the variation observed

between the human populations sampled here. Despite this caveat, that fact that

considerable variation exists between human populations is notable and highlights the need

for much higher resolution sampling of human associated microbial diversity.

In a pangenome analysis to identify individual genes or functions enriched or depleted in

genomes of taxa associated with different human populations, we identified numerous

functional traits involved in aerobic respiration associated with the German population in

the analysis. We hypothesize that taxa found to be enriched in the fecal samples of humans

from Germany might have a selective advantage via their clade-independent ability to

survive or even utilize aerobic conditions in the intestinal tract. More specifically, we

propose that these taxa have undergone convergent adaptation to tolerate high oxygen

concentrations. Such aerotolerance could increase microbial fitness, whereby bacteria can

withstand high oxygen concentrations to metabolize mucus layers for energy (Zheng et al.,

2015). However, the depletion of this mucus by bacteria diminishes an important physical

and immunological barrier that protects the human host against microbial assaults and

allows for direct interaction between host epithelial cells and microbiota, potentially

triggering (auto-)immune processes (Costa et al., 2016; Matute et al., 2023). Notably, we

showed that the introduction of novel microbes associated with industrialization related to

vast differences in the community assembly of fecal microbiota in the German population.

Thus, it is possible that susceptibility to intestinal inflammation might be potentiated by

specific taxa found in this population. Accordingly, we found increased abundances of well-

characterized mucin-degrading taxa, including *Akkermansia* and *Bacteroides*, in the German

cohort. These findings are congruent with previous reports suggesting that there is

increased mucus degradation by intestinal microbiota in human populations with direct

access to industrial food systems, which may relate to higher incidences of inflammatory

bowels diseases observed in developed economies (E. D. Sonnenburg and Sonnenburg,

2019).

Comparatively, only a few pathways showed conserved enrichment in the opposite

direction, suggesting clade specific mechanisms for their loss in some human societies. In

particular, we found the taxon *Prevotella* is depleted in Germans but conserved across

hominids, despite representing a diversity of host clades and diets. *Prevotella* is a major

determining taxon of one of the human enterotypes, a concept used to define fecal

microbial communities (Arumugam et al., 2011). It remains controversial as to whether the

human gut microbiome is best classified using such discrete categories, or rather along a

dynamic, continuous gradient (Cheng 2019, Knight 2014). Nevertheless, previous reports

have shown that individuals who access industrialized food systems (i.e., consume so-called

"Westernized" diets) generally display a *Bacteroides*-dominant enterotype. *Bacteroides*-

enterotypes have been previously associated with a multitude of intestinal (Vieira-Silva et

al., 2019) and extra-intestinal inflammatory diseases (Valles-Colomer et al., 2019).

Conversely, individuals who rely on rural and traditional subsistence strategies (i.e.,

consume plant-rich diets) tend to exhibit a *Prevotella*-dominant enterotype (Vangay et al.,

2018). This enterotype is also displayed in about 20% of individuals living within Western

societies (Costea et al., 2018). Interestingly, there are conflicting reports concerning

Prevotella and host health. While it has been shown that *Prevotella* may improve glucose

metabolism (Kovatcheva-Datchary et al., 2015), other reports have linked high abundances

of *Prevotella* spp. with autoimmune diseases and intestinal inflammation (Iljazovic et al.,

2021). While results from model systems have suggested *Prevotella* likely plays a role in

autoimmunity, these studies largely relied on mono-colonization of germ-free animals and

thus may be biased due to a lack of microbial interaction partners and an aberrant host

physiology (Iljazovic et al., 2021). Within human studies, no convincing link between

increased *Prevotella* spp. and inflammatory bowel disease has yet been shown (Iljazovic et

al., 2021).

Here, we used an evolutionarily-informed framework to extend the enterotype concept to

elucidate the functional dynamics involved in the assembly of the human gut microbiome

over evolutionary timescales. Such insights may better inform how changes in the gut

microbiome might affect human health. We find the taxon *Prevotella* to be conserved across

the sampled hominids. Moreover, the sheer diversity of *Prevotella* displayed across all

hominid clades clearly suggests an evolutionary conservation and long-standing interaction

of this microbial clade with the host, as further revealed by host-specific microbial functions

identified in the metagenomic pangenome analysis. In other words, we find the *Prevotella*

clade to be an integral member of the intestinal microbial community of all hominids.

Therefore, we propose that the *Prevotella*-enterotype represents an evolutionary ancestral

community state for the human gut microbiome. Rather than a discrete enterotype, a

reduced abundance and diversity of *Prevotella* may better regarded a key biomarker for

disease risk (Gorvitovskaia et al., 2016) or for microbiota insufficiency syndrome (J. L.

Sonnenburg and Sonnenburg, 2019), which seems to be partly driven by changes associated

with Western lifestyles. Additional research and large-scale strain collections for *Prevotella*

are needed for an in-depth analysis and evaluation of this diverse taxonomic group with

regard to host health and its role in inflammation. Such research must consider *Prevotella*

spp. as members of a complex consortium of interacting microorganisms and as, we argue, a

potential target for pre- and probiotic intervention in chronic inflammatory disorders.

Lastly, we leveraged our catalog of high-quality metagenome-assembled genomes from

hominid fecal samples together with existing data to investigate co-phylogenetic patterns

across the sampled hosts. Overall, co-phylogeny showed highly clade-specific enrichments

and depletions. In addition, human-derived MAGs were found significantly less often among
co-phylogenetic groups than MAGs from NHPs. Since we included human-derived data from
global reference datasets (Almeida et al., 2021), this effect is unlikely to be an artifact of
non-exhaustive coverage of human microbiome members. Additionally, we provide
evidence for the previously proposed effect of sporulation capabilities as a mechanism for
prokaryotes to become independent of direct host-to-host transmission, disrupting co-
phylogeny patterns, potentially through repeated environmental uptake of spore-forming
bacteria (Groussin et al., 2020).

Our study has limitations. The co-phylogeny analysis relies heavily on genome-sequences
recovered from shotgun metagenomic sequencing (MAGs), which are potentially
contaminated and incomplete, which could bias tree structure and thus, co-phylogeny
estimates, and also the can potentially under- (or over-)estimate the functional capacities of
recovered microbial genomes. To address the potential shortcomings of MAGs, we
implemented stringent data processing pipelines and quality control to achieve the high-
quality MAGs and a host-specific pan-genome based functional annotation framework
incorporating information from multiple MAGs per species representative to reduce
potential genome gaps (see Methods). Additionally, the commonly used estimates of
divergence times of the hominid hosts included in the analysis set the timeframe of the split
from a shared ancestor to 8-19 million years ago (Langergraber et al., 2012; Scally et al.,
2012). Bacterial speciation events happen in the timeframe of 10-100 million years, or
longer (Marin et al., 2017; McDonald and Currie, 2017; Ochman et al., 1999), and thus, co-
phylogeny in hominids is expected to be observed within microbial species or possibly
genera. In the presented dataset, species-level sharing of MAGs between host genera was
low (2.73%; n = 48 out of 1,759 newly reconstructed SGBs, not including Manara *et al.* and
UHGGv2). SGBs were defined on 95% average nucleotide identity, a measure generally
regarded as appropriate (Jain et al., 2018), but it is nevertheless prone to clade specific
biases, potentially further influenced by altered speciation dynamics in association with
(evolutionary) changes in host lifestyle (Lawrence and Retchless, 2009), *i.e.* previously
demonstrated increases in horizontal gene transfer (HGT) within individual microbiomes
(Groussin et al., 2021). Accounting for these potential biases, we relaxed the threshold to a
shared genus-level annotations for subgroups to be included in the co-phylogeny testing,

while keeping the number of tested sub-phylogenies to a minimum through the definition of
stringent inclusion criteria (see Methods). Despite these considerations, the observation of
signals of co-phylogeny across hominids is supported by a robust statistical framework.

Our work here lays the foundation for the analysis of disease-associated changes in the
human intestinal microbiome in an evolutionarily informed framework, thereby allowing
researchers to evaluate microbiome-associated inflammatory disorders from a point of view
that considers both proximal and evolutionary influences. Future investigations should
consider in-depth analysis of horizontal gene transfer events within or even between
primate hosts to shed further light on also cross-species dynamics and transition of
microbes. Such analyses however require either microbial isolate genomes or at least long-
read sequencing data to increase confidence in detection events. Additionally, time series
data for host groups sharing the same habitat, *e.g. G. g. gorilla* and *P. t. troglodytes*, could
give additional insights into cross-species sharing dynamics which cannot be appropriately
elucidated based on single-timepoint data.

In summary, we present an in-depth taxonomic and functional description and analysis of
hominid-associated fecal communities spanning about ten millions of years of evolution and
host-microbiome interactions in the gut of humans and African great apes. Western lifestyle
and maybe more precisely industrialization associated changes in human gut microbiota
have been previously suggested as a driver of microbiome insufficiency syndrome, whereby
an incompatibility between quickly adapting microbiota and slowly evolving host genes
leads to chronic inflammatory diseases such as metabolic syndrome, type 2 diabetes, and
inflammatory bowel disease (E. D. Sonnenburg and Sonnenburg, 2019; Wallenborn and
Vonaesch, 2022). Thus, a comparative analysis of human and NHP intestinal microbiota that
considers evolutionary forces as presented herein provides a powerful platform to advance
our understanding of human-associated microbiota and guide the development of
personalized, targeted interventions to prevent and treat chronic inflammatory disorders.

Tables
670
671 Table1: KEGG functional groups with significant enrichment (Q < 0.05) in the gut
672 microbiome of humans living in Germany.
673

KEGG ID	Name	# of KOs	Mean Z-Score	P-value	Q-Value
path:map01230	Biosynthesis of amino acids	238	2.43	2.80E-17	2.29E-14
path:map00340	Histidine metabolism	47	4.69	3.92E-16	3.20E-13
md:M00159	V/A-type ATPase, prokaryotes	9	13.61	5.08E-15	4.15E-12
md:M00026	Histidine biosynthesis, PRPP => histidine	20	4.56	3.05E-13	2.49E-10
path:map00630	Glyoxylate and dicarboxylate metabolism	101	3.27	3.26E-11	2.66E-08
br:br01601	Enzymes of 2-oxocarboxylic acid metabolism	60	5.39	1.33E-08	1.09E-05
md:M00010	Citrate cycle, first carbon oxidation, oxaloacetate => 2-oxoglutarate	6	5.98	3.68E-07	3.01E-04
md:M00004	Pentose phosphate pathway (Pentose phosphate cycle)	15	7.29	8.40E-07	6.87E-04
md:M00045	Histidine degradation, histidine => N-formiminoglutamate => glutamate	8	4.7	1.68E-06	1.38E-03
md:M00631	D-Galacturonate degradation (bacteria), D-galacturonate => pyruvate + D-glyceraldehyde 3P	8	4.92	1.68E-06	1.38E-03
path:map00520	Amino sugar and nucleotide sugar metabolism	156	2.38	4.74E-06	3.87E-03
path:map00620	Pyruvate metabolism	132	3.41	6.86E-06	5.60E-03
path:map00260	Glycine, serine and threonine metabolism	109	2.54	9.95E-06	8.13E-03
md:M00009	Citrate cycle (TCA cycle, Krebs cycle)	42	3.81	1.37E-05	1.12E-02
path:map00020	Citrate cycle (TCA cycle)	66	4.35	1.94E-05	1.59E-02
md:M00008	Entner-Doudoroff pathway, glucose-6P => glyceraldehyde-3P + pyruvate	5	7.59	1.97E-05	1.61E-02
path:map00190	Oxidative phosphorylation	223	8.71	2.74E-05	2.24E-02
md:M00006	Pentose phosphate pathway, oxidative phase, glucose 6P => ribulose 5P	6	8.44	3.90E-05	3.19E-02

674
675
Acknowledgements

We thank the IKMB Microbiome and NGS laboratories for excellent technical support. We

thank Sébastien Calvignac-Spencer and Grit Schubert for their valuable contributions to the

analysis and the manuscript. This study is an outcome of the Collaborative Research Center

1182, Origin and Function of Metaorganisms (https://www.metaorganism-research.com,

no. SFB1182, Projects A2 and Z2) and received infrastructure support from the DFG

Excellence Cluster 2167 "Precision Medicine in Chronic Inflammation" (PMI). JFG was

supported by the DFG grant "The ecology and evolution of primate phageomes" (GO

3443/1-1: Project number: 453352748) and the Deutsche Forschungsgemeinschaft (DFG)

Research Group "Sociality and Health in Primates" (FOR2136; CA 1108/3-1). Research at

Kokolopori was supported by the Max Planck Society and Harvard University. Research at
Ozouga and Taï National Park was supported by the Max Planck Society. Research at Loango
was supported by US Fish and Wildlife Service, Tusk Trust, Berggorilla Regenwald Direkthilfe,
and the Max Planck Society. Research at Bwindi was supported by the Max Planck Society.
The survey of the Bwindi mountain gorillas was funded by the International Gorilla
Conservation Programme coalition members with supplemental funding from the Wildlife
Conservation Society. We are extremely grateful to the organizations that facilitated work at
these field sites and granted permission for sample collection: Bwindi—The mountain gorilla
survey was conducted by the Uganda Wildlife Authority, l'Institut Congolais pour la
Conservation de la Nature, the Rwanda Development Board, the International Gorilla
Conservation Programme, the Max Planck Institute for Evolutionary Anthropology,
Conservation Through Public Health, the Mountain Gorilla Veterinary Project, the Institute
for Tropical Forest Conservation, and The Dian Fossey Gorilla Fund and was conducted in
compliance with the regulations of and permission of the Uganda National Council for
Science and Technology and the Uganda Wildlife Authority; Kokolopori—permission was
granted through the Ministere de Recherche Scientifique et Technologie, Democratic
Republic of the Congo, and work was supported by Vie Sauvage, the Bonobo Conservation
Initiative; Loango—permission was granted by the Agence Nationale des Parcs Nationaux,
the Centre National de la Recherche Scientifique et Technique of Gabon; Taï National Park—
permission was granted by the Ministère de l'Enseignement Supérieur et de la Recherche
Scientifique, the Ministère des Eaux et Fôrets in Côte d'Ivoire, and the Office Ivoirien des
Parcs et Réserves, and work was supported by the Centre Suisse de Recherches Scientifiques
en Côte d'Ivoire and the staff members of the Taï Chimpanzee Project. Ethical approval for
work on human samples was obtained from the Local Ethics Committee Germany, Kiel
(reference number A156/03), and we thank the national and local health authorities in Côte
d'Ivoire and Democratic Republic of Congo as well as the according ethics commission for
granting permission for this work. We thank Taylor Hermes for support in the writing of the
manuscript.
714
Data and code availability

All metagenomic sequencing data is available via the NCBI BioProject accession IDs

PRJNA692042 and PRJNA539933. All code to process sequencing files to generate the

presented results is available via https://github.com/mruehlemann/greatapes_mgx_scripts.

Conflicts of interest

All authors declare no conflicts of interest.

Material and Methods

Fecal Sampling, DNA extraction and data generation

Sampling procedures for collecting feces from humans (n=48) and wild non-human primates

(n=109) have been previously described (Gogarten et al., 2021). Briefly, fecal samples were

collected immediately after defecation, and, depending on the local infrastructure, either

stored in RNAlater and frozen at -20°C or stored in a cryotube, cooled in a thermos until

return to the field laboratory, and subsequently snap frozen in liquid nitrogen. Appropriate

government permits and permission to conduct research on wild primates were granted by

the relevant authorities (see Acknowledgments for site-specific details). Human fecal

samples from the Democratic Republic of Congo (n = 12) were stored in RNAlater and frozen

at -20°C. Human fecal samples from Côte d'Ivoire (n=12) were stored in a cryotube, cooled

in a thermos until return to the field laboratory and subsequently snap frozen in liquid

nitrogen. Human fecal samples from Germany were collected at home by the participant in

standard fecal collection tubes, mailed to the study center, and stored at -80°C. Ethical

approval was obtained from the Local Ethics Committee (Kiel, Germany; reference number

A156/03). DNA extraction from fecal samples was performed from 200mg of stool

transferred to 0.70mm Garnet Bead tubes (Qiagen) with 1.1 mL ASL buffer, followed by

bead beating in a SpeedMill PLUS (Analytik Jena AG) for 45 s at 50 Hz. Samples were heated

to 95°C for 5 min and centrifuged, retaining 200 µl of the supernatant for DNA extraction

with the QIAamp DNA Stool Mini Kit (Qiagen) automated on a QIAcube system (Qiagen)

according to the manufacturer's protocol. DNA quality was assessed by Qubit and Genomic

DNA ScreenTape (Agilent). Illumina Nextera DNA Library Preparation Kit was used to

construct shotgun metagenomic libraries, and subsequently sequenced with either 2 × 125

bp reads on a HiSeq 2500 platform or with 2 × 150 bp reads on a HiSeq 4000 machine

(Illumina).

Data processing, assembly and metagenomic binning

Raw sequencing FastQ files were quality controlled and preprocessed using the BBMap software suite ("BBMap," n.d.). Host reads were removed using a human reference database and a lenient threshold of 95% identity to account for a broader host range using bbmap.sh. Metagenomic contigs were assembled with metaSPAdes and contigs >= 2000 bp were retained (Bankevich et al., 2012). Reads were mapped to the contigs of the respective samples using Minimap2 (Li, 2018), converted to BAM files with Samtools (Li et al., 2009) and used to estimate per-contig mapping depth with the jgi_summarize_bam_contig__depths binary from the MetaBAT2 binning tool (Kang et al., 2019). Contig binning for individual samples was performed with MetaBAT2 (Kang et al., 2019), MaxBin2 (Wu et al., 2016) and CONCOCT (Alneberg et al., 2014). In addition, the VAMB binning tool (Nissen et al., 2021) was used on a cross-mapping catalog of the merged contigs from all samples within each host group. Individual binning results were refined using MAGScoT (Rühlemann et al., 2022) to acquire high quality metagenome-assembled genomes (MAGs) for each sample. Clustering of MAGs to species-level genome bins (SGBs) was performed with dRep (Olm et al., 2017) in a multi-step approach to control for inflated SGBs due to low MAG quality. First, MAGs were dereplicated to 97% similarity within each host group, choosing the MAG with the highest score (calculated by MAGScoT based on completeness and contamination) as cluster representative. High and good quality representatives (score > =0.7) from all host groups together with representative sequences from the UHGG v2 were then clustered into 95% SGBs using dRep, again selecting the highest quality MAG as representatives. Medium quality (scores between 0.5 and 0.7) 97% representatives from previous clustering step were then compared to SGB representatives using fastANI (Jain et al., 2018), assigning MAGs with high similarity (>=95%) to the respective SGB. Medium quality 97% representatives without hits to the high quality SGB library were then clustered into 95% SGBs and added to the catalog in the case of at least two genomes in the cluster, discarding singleton clusters. The final catalog of SGB representatives was used to quantify contig abundances in all samples using Salmon in metagenome mode (Patro et al., 2017). Taxonomic annotations were performed using the GTDBtk (v2.1) and GTDB release 207v2 (Chaumeil et al., 2022; Parks et al., 2022). For SGBs

780 without genus- and/or species-level assignments, the SGB ID was used as taxonomic label.
781 GTDBtk marker gene alignments were used to generate a phylogenetic tree of all SGB
782 representatives using the respective "infer" function of the GTDBtk. All data processing
783 scripts are available online: https://github.com/mruehlemann/greatapes_mgx_scripts

Pangenome catalog creation, annotation and analysis

All MAGs underwent calling of coding sequences using prodigal (v2.6.3) (Hyatt et al., 2010).
Protein sequences were clustered based on 95% similarity using MMseqs (Hauser et al.,
2016; Steinegger and Söding, 2017) and annotated using the emapper.py script of the
eggNOG-mapper v2 (Cantalapiedra et al., 2021) annotation tool with the eggNOG 5.0
reference database (Huerta-Cepas et al., 2019). MAG level functional profiles based on
KEGG Ortholog annotations were collapsed into SGB-level pangenomes for each host genus
(*Homo, Gorilla*, and *Pan*). In the case that no MAGs of an SGB were recovered from a given
host genus, functional profiles were inferred from MAGs across the other host groups,
accounting for host-specificities in the inferred accessory genomes/functions by considering
a function to be present if it was present in all host-specific pangenomes of the respective
SGB with MAGs recovered from the metagenomic data.

Calculation of microbial clade and functional abundances

All downstream data processing and statistical calculations were performed in R v4.2 (R
Core Team, 2022) and using the tidyverse library (Wickham et al., 2019). Per-sample contig
abundances for the SGB representatives from Salmon were used to estimate SGB
abundances. Salmon output includes total mapped reads per contig and mapping reads
adjusted for library size and total sequencing depth as transcripts per million (TPM), a
measure from the transcriptomics field which can be directly transferred to metagenomic
libraries. Individual contig coverages were calculated from the number of mapped reads and
the effective lengths of the Salmon mapping output, considering contigs with > 10%
coverage as present. An SGB was considered present when at least 20% of its total length
was in contigs marked as "present" and if at least 1,000 total reads and 250 TPM mapped to
it. Final SGB abundances were calculated as TPM, calculated from the reads mapping to the
SGBs present in the respective sample, thus representing a normalized abundance across all
samples. Combining SGB abundances with taxonomic assignments, domain- to species-level

abundances were calculated as cumulative TPM abundances within the respective

taxonomic bins. Rarefactions were calculated based on 5-fold repeated subsampling of

contig level mapped reads at 100k, 250k, 500k, 1M, 2.5M, 5M, and 10M reads, followed by

TPM calculations as described above. By rarefying reads and not TPM we realistically

simulate sampling effects introduced by low coverage and low abundances of SGBs affecting

especially samples with small library sizes. Community level functional profiles were

calculated be multiplying TPM abundances of SGBs with the respective host-genus specific

functional profiles of the SGBs and summarizing the per-SGB values into a sample-level

abundance of functional annotations.

Alpha and beta diversity

Faith's phylogenetic diversity (PD) (Faith, 1992) was used as measure of alpha diversity,

calculated from the phylogenetic tree based on GTDBtk marker genes using the pd()

function of the picante package for R (Kembel et al., 2010). Genus-level increase of PD from

novel SGBs was calculated from the differences of PDs with and without novel SGB

annotated as the respective genus. Sample level PDs were calculated from the SGB

presence/absence patterns. Beta diversity was assessed as unweighted and weighted

UniFrac distances (Lozupone et al., 2011) using the UniFrac() function of the phyloseq

package for R (McMurdie and Holmes, 2013) and SGB abundances and the phylogenetic tree

based on GTDBtk marker genes as input. Aitchison distance (Aitchison, 1982) was calculated

from the CLR-transformed genus-level TPM abundances obtained from the clr() function

from the compositions package for R (van den Boogaart and Tolosana-Delgado, 2008) and

adding a pseudocount of 1 to all abundances, setting all CLR-transformed abundances below

zero to zero. Jaccard distances (Jaccard, 1912) were also calculated on genus-level

presence/absence patterns using the vegdist() function from the vegan package for R

(Oksanen et al., 2022). Genus-level abundances were chosen for Aitchison and Jaccard

distance, as SGBs are highly host-specific, thus would lead to high beta-diversities simply

due to host exclusive SGBs, grouping at genus-level prevents from this and UniFrac distances

use phylogenetic relations between SGBs.

Cross-host sharing analysis

Analysis of excessive and reduced sharing of SGBs between host groups were based on the

mean SGB abundances of the five rarefaction of 1M mapped reads to account for

differences in library depth impacting SGB richness. For each host group, 100-fold sampling

of five samples from this group were drawn and the SGBs found in the host were analyzed

for their presence in five random samples of each of the other host groups, calculating the

relative amount of shared SGBs. The mean of all 100 samplings was used as relative sharing

coefficient for all host pairs in both directions. Excess and reduced sharing was analyzed by

1000-fold drawing of five random samples accounting for differences of host groups and the

repetition of above calculations for relative sharing with all host groups. P-values were

calculated from the proportions of random samplings exceeding/falling below the true

sharing coefficients.

Phylosymbiosis analysis

Phylosymbiosis was assessed using five measures for community level diversity, unweighted

and weighted UniFrac, genus-level Aitchison, and Jaccard distances, as well as KEGG

ortholog (KO) abundance based Aitchison distance, and following the approach described in

Brooks et al. (2016). Briefly, host group differences were used to infer microbiome

dendrograms by UPGMA clustering. Branch support was calculated from 1000-fold jackknife

sampling. Robinson-Foulds distances between microbiome trees and host phylogeny were

calculated using the RF.dist() function from the phangorn package for R (Schliep, 2011).

Significance of phylosymbiosis was assessed using the host phylogeny and 100,000 random

trees as comparison for the microbiome trees. Tanglegrams were created with the ggtree

and cowplot packages for R (Yu et al., 2017).

Assessment of between-group abundance differences

Taxonomic and functional abundance differences between Humans and NHPs, as well as

between humans living outside and within industrialized systems were based on CLR-

transformed abundances to account for the compositionality of microbiome data (Gloor et

al., 2017). Included in the calculations were all genera with a prevalence > 20% and relative

abundance (before CLR-transformation) of > 0.1% in at least of the host groups and all KO

abundances with a prevalence > 20% and CLR-transformed abundance of > 1 in at least of

the host groups. Abundance differences were assessed using linear regression. Log-fold

differences were calculated using group mean abundances and a pseudocount of 0.01. P-
values were adjusted for multiple testing using Bonferroni correction. Genera with
significant differences with lifestyle were grouped into BloSSUM (Bloom or Selected in
Societies of Urbanization/Modernization) and VANISH (Volatile and/or Associated
Negatively with Industrialized Societies of Humans) taxa. Remaining genera with significant
differences between humans and NHPs were grouped according to these patterns. Genera
without abundance differences in any of these comparisons were grouped as "other".
Accordingly, significantly different KO abundances were grouped into association with
human subgroups or humans vs. NHPs depending on the statistical significance and
direction of the association.
885
Functional pangenome differences between groups
Pangenome catalogs of human-associated SGBs were compared within microbial families
between SGBs in BloSSUM and non-BloSSUM taxa, the latter comprising all VANISH taxa and
taxa with higher abundance in humans compared to NHPs, however not associated with the
industrialization gradient. KEGG Ontology (KO) term annotations were used as functional
groups and their prevalence differences between groups were assessed using Fisher's exact
test. Effect sizes were calculated using the log2 of the ratio of prevalences in the two groups
and a pseudo count of 0.01. Per-family Z-Scores of KOs were calculated from P-values and
the direction of the effects. The sum of the Z-Scores were added and divided by the square-
root of the total number of families the respective KOs were found in to obtain a Z_{Meta} for
each KO term, used to calculated P_{Meta}. P_{Meta}-values were adjusted for multiple testing using
Holm-correction. KO terms with $Q < 0.05$ and present in at least two of the microbial
families in the analysis were considered as functions with differential prevalence. A similar
approach was employed to assess functional differences between NHP- and Human-
associated SGBs, however in this case, SGB pangenome differences were compared on
genus level and the meta-analysis was performed combining signals from all genera, and
specifically across the genera within particular phyla.
903
Tree reconciliation analysis
Proteins from the representative SGBs of the genus *Prevotella* annotated with the
annotation "cydA" (cytochrome *bd* oxidase subunit 1) as "Preferred name" in the

emapper/eggNOG annotation were extracted from the unclustered protein sequence
catalog. The same procedure was followed for *Paraprevotella,* which was included as an
outgroup. Incomplete cydA sequences were removed using a length threshold of 200.
Protein sequences were aligned using Clustal Omega (Sievers et al., 2011). The alignment
was used to reconstruct the phylogenetic tree using IQTREE2 (Minh et al., 2020) and a WAG
model (Whelan and Goldman, 2001), calculating branch support using UFBoot (Minh et al.,
2013) and performing SH-aLRT test (Guindon et al., 2010). Alignments of GTDBtk marker
protein sequences for *Prevotella* and *Paraprevotella* SGBs were used to reconstruct a
genome-level species phylogeny in the same respective way as described above for the cydA
sequences. Very low confident branches (< 60% bootstrap support) in the cydA phylogeny
were resolved together with the species tree using the OptResolutions supplementary
program of the RANGER-DTL 2.0 software (Bansal et al., 2018) resulting in 495 equally
probable trees with optimized duplication-transfer-loss costs using default values
(duplication: 2, loss: 1, transfer: 3). A randomly chosen output tree was using in the
reconciliation analysis with the species tree in RANGER-DTL 2.0 using default values and
1,000 random starting seeds in parallel (Tange, 2011) to assess robustness. Resulting
sampling outcomes were summarized using the AggregateRanger tool of the RANGER-DTL
2.0 software package.

Co-phylogeny analysis
Host phylogenetic trees were obtained from the 10kTrees website ((Arnold et al., 2010);
https://10ktrees.nunn-lab.org/). To assure high quality microbial phylogenies for the co-
phylogeny analysis, family-level maximum-likelihood trees were reconstructed from the
GTDBtk marker gene alignments with the IQTREE2 software (Minh et al., 2020) and a WAG
model including a random SGB outside the respective families as outgroups. Family level
trees were rooted and for each SGB traced from tip to root to identify for each SGB the
smallest subtree which covered 4, 5, 6, and 7 host groups. Combining information from all
SGBs, the overall set of smallest trees to be included in the co-phylogeny analysis were
identified, discarding subtrees for which the inclusion criterion was fulfilled already for a
smaller tree starting from a different tip. In addition, subtrees spanning more than a single
genus were excluded from the analysis, as divergence times of microbial genera predate
divergence of the included hosts (Ochman et al., 1999). For all subgroups included in the

analysis, maximum-likelihood distances and trees using a WAG model in IQTREE2 were

inferred from the marker gene alignment of all MAGs assigned to the SGBs in the respective

subgroups. Co-phylogeny of the subgroup was assessed by randomly selecting one MAG per

host, calculating congruence with the host tree by Robinson-Foulds metric and by Mantel-

test (Hommola et al., 2009). Tiplabels were permutated 999-fold and P-values calculated.

This process starting from the random selection of one MAG per host was repeated 999

times to obtain final P-values. Family-level co-phylogeny ratios were calculated based on the

ratio of SGBs within subtrees with co-phylogeny signal and total SGBs in the respective

family that were included in the analysis. Enrichment of co-phylogeny for each microbial

family was calculated by using Fisher's exact test on the SGBs in the analysis dividing them

into four groups based on family membership and being in a subtree with co-phylogeny

signal. All P-values were adjusted using FDR correction.

Correlation between co-phylogeny ratio and sporulation genes

Genes were selected using the KEGG Ontology annotations, selecting KOs with the words

"spore" or "sporulation" in their description. This resulted in a total of 149 unique KOs. The

presence of these KOs in an SGB was assessed based on the pangenome annotations (see

above). The number of KOs was summed up for each genome and genus-level averages

were regressed against the family-level co-phylogeny ratios. The P-value of the correlation

was calculated from the two-sided t-statistic calculated from regression coefficient and

standard error.

References

Aitchison, J., 1982. The Statistical Analysis of Compositional Data. Journal of the Royal
Statistical Society: Series B (Methodological) 44, 139–160.
https://doi.org/10.1111/j.2517-6161.1982.tb01195.x
Almeida, A., Nayfach, S., Boland, M., Strozzi, F., Beracochea, M., Shi, Z.J., Pollard, K.S.,
Sakharova, E., Parks, D.H., Hugenholtz, P., Segata, N., Kyrpides, N.C., Finn, R.D., 2021.
A unified catalog of 204,938 reference genomes from the human gut microbiome.
Nat Biotechnol 39, 105–114. https://doi.org/10.1038/s41587-020-0603-3
Alneberg, J., Bjarnason, B.S., de Bruijn, I., Schirmer, M., Quick, J., Ijaz, U.Z., Lahti, L., Loman,
N.J., Andersson, A.F., Quince, C., 2014. Binning metagenomic contigs by coverage
and composition. Nat Methods 11, 1144–1146. https://doi.org/10.1038/nmeth.3103
Amato, K.R., Mallott, E.K., McDonald, D., Dominy, N.J., Goldberg, T., Lambert, J.E., Swedell,
L., Metcalf, J.L., Gomez, A., Britton, G.A.O., Stumpf, R.M., Leigh, S.R., Knight, R., 2019.
Convergence of human and Old World monkey gut microbiomes demonstrates the

importance of human ecology over phylogeny. Genome Biology 20, 201.
https://doi.org/10.1186/s13059-019-1807-z

Arnold, C., Matthews, L.J., Nunn, C.L., 2010. The 10kTrees website: A new online resource
for primate phylogeny. Evolutionary Anthropology: Issues, News, and Reviews 19,
114–118. https://doi.org/10.1002/evan.20251

Arumugam, M., Raes, J., Pelletier, E., Le Paslier, D., Yamada, T., Mende, D.R., Fernandes,
G.R., Tap, J., Bruls, T., Batto, J.-M., Bertalan, M., Borruel, N., Casellas, F., Fernandez,
L., Gautier, L., Hansen, T., Hattori, M., Hayashi, T., Kleerebezem, M., Kurokawa, K.,
Leclerc, M., Levenez, F., Manichanh, C., Nielsen, H.B., Nielsen, T., Pons, N., Poulain,
J., Qin, J., Sicheritz-Ponten, T., Tims, S., Torrents, D., Ugarte, E., Zoetendal, E.G.,
Wang, J., Guarner, F., Pedersen, O., de Vos, W.M., Brunak, S., Doré, J., MetaHIT
Consortium, Antolín, M., Artiguenave, F., Blottiere, H.M., Almeida, M., Brechot, C.,
Cara, C., Chervaux, C., Cultrone, A., Delorme, C., Denariaz, G., Dervyn, R., Foerstner,
K.U., Friss, C., van de Guchte, M., Guedon, E., Haimet, F., Huber, W., van Hylckama-
Vlieg, J., Jamet, A., Juste, C., Kaci, G., Knol, J., Lakhdari, O., Layec, S., Le Roux, K.,
Maguin, E., Mérieux, A., Melo Minardi, R., M'rini, C., Muller, J., Oozeer, R., Parkhill, J.,
Renault, P., Rescigno, M., Sanchez, N., Sunagawa, S., Torrejon, A., Turner, K.,
Vandemeulebrouck, G., Varela, E., Winogradsky, Y., Zeller, G., Weissenbach, J.,
Ehrlich, S.D., Bork, P., 2011. Enterotypes of the human gut microbiome. Nature 473,
174–180. https://doi.org/10.1038/nature09944

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M.,
Nikolenko, S.I., Pham, S., Prjibelski, A.D., Pyshkin, A.V., Sirotkin, A.V., Vyahhi, N.,
Tesler, G., Alekseyev, M.A., Pevzner, P.A., 2012. SPAdes: A New Genome Assembly
Algorithm and Its Applications to Single-Cell Sequencing. J Comput Biol 19, 455–477.
https://doi.org/10.1089/cmb.2012.0021

Bansal, M.S., Kellis, M., Kordi, M., Kundu, S., 2018. RANGER-DTL 2.0: rigorous reconstruction
of gene-family evolution by duplication, transfer and loss. Bioinformatics 34, 3214–
3216. https://doi.org/10.1093/bioinformatics/bty314

BBMap [WWW Document], n.d. . SourceForge. URL
https://sourceforge.net/projects/bbmap/ (accessed 12.1.22).

Brooks, A.W., Kohl, K.D., Brucker, R.M., Opstal, E.J. van, Bordenstein, S.R., 2016.
Phylosymbiosis: Relationships and Functional Effects of Microbial Communities
across Host Evolutionary History. PLOS Biology 14, e2000225.
https://doi.org/10.1371/journal.pbio.2000225

Campbell, T.P., Sun, X., Patel, V.H., Sanz, C., Morgan, D., Dantas, G., 2020. The microbiome
and resistome of chimpanzees, gorillas, and humans across host lifestyle and
geography. ISME J 14, 1584–1599. https://doi.org/10.1038/s41396-020-0634-2

Cantalapiedra, C.P., Hernández-Plaza, A., Letunic, I., Bork, P., Huerta-Cepas, J., 2021.
eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain
Prediction at the Metagenomic Scale. Molecular Biology and Evolution 38, 5825–
5829. https://doi.org/10.1093/molbev/msab293

Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., Parks, D.H., 2022. GTDB-Tk v2: memory
friendly classification with the genome taxonomy database. Bioinformatics 38, 5315–
5316. https://doi.org/10.1093/bioinformatics/btac672

Costa, F.R.C., Françozo, M.C.S., de Oliveira, G.G., Ignacio, A., Castoldi, A., Zamboni, D.S.,
Ramos, S.G., Câmara, N.O., de Zoete, M.R., Palm, N.W., Flavell, R.A., Silva, J.S., Carlos,
D., 2016. Gut microbiota translocation to the pancreatic lymph nodes triggers NOD2

activation and contributes to T1D onset. J Exp Med 213, 1223–1239.
https://doi.org/10.1084/jem.20150744
Costea, P.I., Hildebrand, F., Arumugam, M., Bäckhed, F., Blaser, M.J., Bushman, F.D., de Vos,
W.M., Ehrlich, S.D., Fraser, C.M., Hattori, M., Huttenhower, C., Jeffery, I.B., Knights,
D., Lewis, J.D., Ley, R.E., Ochman, H., O'Toole, P.W., Quince, C., Relman, D.A.,
Shanahan, F., Sunagawa, S., Wang, J., Weinstock, G.M., Wu, G.D., Zeller, G., Zhao, L.,
Raes, J., Knight, R., Bork, P., 2018. Enterotypes in the landscape of gut microbial
community composition. Nat Microbiol 3, 8–16. https://doi.org/10.1038/s41564-
017-0072-8
Davidson, A.L., Chen, J., 2004. ATP-binding cassette transporters in bacteria. Annu Rev
Biochem 73, 241–268. https://doi.org/10.1146/annurev.biochem.73.011303.073626
Davies, J.S., Currie, M.J., Wright, J.D., Newton-Vesty, M.C., North, R.A., Mace, P.D., Allison,
J.R., Dobson, R.C.J., 2021. Selective Nutrient Transport in Bacteria: Multicomponent
Transporter Systems Reign Supreme. Frontiers in Molecular Biosciences 8.
Deleu, S., Machiels, K., Raes, J., Verbeke, K., Vermeire, S., 2021. Short chain fatty acids and
its producing organisms: An overlooked therapy for IBD? EBioMedicine 66, 103293.
https://doi.org/10.1016/j.ebiom.2021.103293
Ejby, M., Fredslund, F., Andersen, J.M., Vujičić Žagar, A., Henriksen, J.R., Andersen, T.L.,
Svensson, B., Slotboom, D.J., Abou Hachem, M., 2016. An ATP Binding Cassette
Transporter Mediates the Uptake of α-(1,6)-Linked Dietary Oligosaccharides in
Bifidobacterium and Correlates with Competitive Growth on These Substrates. J Biol
Chem 291, 20220–20231. https://doi.org/10.1074/jbc.M116.746529
Faith, D.P., 1992. Conservation evaluation and phylogenetic diversity. Biological
Conservation 61, 1–10. https://doi.org/10.1016/0006-3207(92)91201-3
Giuffrè, A., Borisov, V.B., Arese, M., Sarti, P., Forte, E., 2014. Cytochrome bd oxidase and
bacterial tolerance to oxidative and nitrosative stress. Biochimica et Biophysica Acta
(BBA) - Bioenergetics, 18th European Bioenergetics Conference 2014 Lisbon,
Portugal 1837, 1178–1187. https://doi.org/10.1016/j.bbabio.2014.01.016
Gloor, G.B., Macklaim, J.M., Pawlowsky-Glahn, V., Egozcue, J.J., 2017. Microbiome Datasets
Are Compositional: And This Is Not Optional. Frontiers in Microbiology 8.
Gogarten, J.F., Rühlemann, M., Archie, E., Tung, J., Akoua-Koffi, C., Bang, C., Deschner, T.,
Muyembe-Tamfun, J.-J., Robbins, M.M., Schubert, G., Surbeck, M., Wittig, R.M.,
Zuberbühler, K., Baines, J.F., Franke, A., Leendertz, F.H., Calvignac-Spencer, S., 2021.
Primate phageomes are structured by superhost phylogeny and environment.
Proceedings of the National Academy of Sciences 118, e2013535118.
https://doi.org/10.1073/pnas.2013535118
Gorvitovskaia, A., Holmes, S.P., Huse, S.M., 2016. Interpreting Prevotella and Bacteroides as
biomarkers of diet and lifestyle. Microbiome 4, 15. https://doi.org/10.1186/s40168-
016-0160-7
Groussin, M., Mazel, F., Alm, E.J., 2020. Co-evolution and Co-speciation of Host-Gut Bacteria
Systems. Cell Host & Microbe 28, 12–22.
https://doi.org/10.1016/j.chom.2020.06.013
Groussin, M., Mazel, F., Sanders, J.G., Smillie, C.S., Lavergne, S., Thuiller, W., Alm, E.J., 2017.
Unraveling the processes shaping mammalian gut microbiomes over evolutionary
time. Nat Commun 8, 14319. https://doi.org/10.1038/ncomms14319
Groussin, M., Poyet, M., Sistiaga, A., Kearney, S.M., Moniz, K., Noel, M., Hooker, J., Gibbons,
S.M., Segurel, L., Froment, A., Mohamed, R.S., Fezeu, A., Juimo, V.A., Lafosse, S.,

Tabe, F.E., Girard, C., Iqaluk, D., Nguyen, L.T.T., Shapiro, B.J., Lehtimäki, J.,
Ruokolainen, L., Kettunen, P.P., Vatanen, T., Sigwazi, S., Mabulla, A., Domínguez-
Rodrigo, M., Nartey, Y.A., Agyei-Nkansah, A., Duah, A., Awuku, Y.A., Valles, K.A.,
Asibey, S.O., Afihene, M.Y., Roberts, L.R., Plymoth, A., Onyekwere, C.A., Summons,
R.E., Xavier, R.J., Alm, E.J., 2021. Elevated rates of horizontal gene transfer in the
industrialized human microbiome. Cell 184, 2053-2067.e18.
https://doi.org/10.1016/j.cell.2021.02.052

Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010. New
Algorithms and Methods to Estimate Maximum-Likelihood Phylogenies: Assessing
the Performance of PhyML 3.0. Systematic Biology 59, 307–321.
https://doi.org/10.1093/sysbio/syq010

Hatakeyama, M., 2014. Helicobacter pylori CagA and gastric cancer: a paradigm for hit-and-
run carcinogenesis. Cell Host Microbe 15, 306–316.
https://doi.org/10.1016/j.chom.2014.02.008

Hauser, M., Steinegger, M., Söding, J., 2016. MMseqs software suite for fast and deep
clustering and searching of large protein sequence sets. Bioinformatics 32, 1323–
1330. https://doi.org/10.1093/bioinformatics/btw006

Hommola, K., Smith, J.E., Qiu, Y., Gilks, W.R., 2009. A permutation test of host-parasite
cospeciation. Mol Biol Evol 26, 1457–1468. https://doi.org/10.1093/molbev/msp062

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S.K., Cook, H.,
Mende, D.R., Letunic, I., Rattei, T., Jensen, L.J., von Mering, C., Bork, P., 2019.
eggNOG 5.0: a hierarchical, functionally and phylogenetically annotated orthology
resource based on 5090 organisms and 2502 viruses. Nucleic Acids Res 47, D309–
D314. https://doi.org/10.1093/nar/gky1085

Hyatt, D., Chen, G.-L., LoCascio, P.F., Land, M.L., Larimer, F.W., Hauser, L.J., 2010. Prodigal:
prokaryotic gene recognition and translation initiation site identification. BMC
Bioinformatics 11, 119. https://doi.org/10.1186/1471-2105-11-119

Iljazovic, A., Amend, L., Galvez, E.J.C., de Oliveira, R., Strowig, T., 2021. Modulation of
inflammatory responses by gastrointestinal Prevotella spp. – From associations to
functional studies. International Journal of Medical Microbiology 311, 151472.
https://doi.org/10.1016/j.ijmm.2021.151472

Jaccard, P., 1912. The Distribution of the Flora in the Alpine Zone.1. New Phytologist 11, 37–
50. https://doi.org/10.1111/j.1469-8137.1912.tb05611.x

Jain, C., Rodriguez-R, L.M., Phillippy, A.M., Konstantinidis, K.T., Aluru, S., 2018. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species
boundaries. Nat Commun 9, 5114. https://doi.org/10.1038/s41467-018-07641-9

Jha, A.R., Davenport, E.R., Gautam, Y., Bhandari, D., Tandukar, S., Ng, K.M., Fragiadakis, G.K.,
Holmes, S., Gautam, G.P., Leach, J., Sherchand, J.B., Bustamante, C.D., Sonnenburg,
J.L., 2018. Gut microbiome transition across a lifestyle gradient in Himalaya. PLoS
Biol 16, e2005396. https://doi.org/10.1371/journal.pbio.2005396

Johnson, A.J., Vangay, P., Al-Ghalith, G.A., Hillmann, B.M., Ward, T.L., Shields-Cutler, R.R.,
Kim, A.D., Shmagel, A.K., Syed, A.N., Walter, J., Menon, R., Koecher, K., Knights, D.,
2019. Daily Sampling Reveals Personalized Diet-Microbiome Associations in Humans.
Cell Host & Microbe 25, 789-802.e5. https://doi.org/10.1016/j.chom.2019.05.005

Joossens, M., Huys, G., Cnockaert, M., Preter, V.D., Verbeke, K., Rutgeerts, P., Vandamme,
P., Vermeire, S., 2011. Dysbiosis of the faecal microbiota in patients with Crohn's

disease and their unaffected relatives. Gut 60, 631–637.
https://doi.org/10.1136/gut.2010.223263
Jurtshuk, P., 1996. Bacterial Metabolism, in: Baron, S. (Ed.), Medical Microbiology.
University of Texas Medical Branch at Galveston, Galveston (TX).
Kanehisa, M., Goto, S., 2000. KEGG: kyoto encyclopedia of genes and genomes. Nucleic
Acids Res 28, 27–30. https://doi.org/10.1093/nar/28.1.27
Kang, D.D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., Wang, Z., 2019. MetaBAT 2: an
adaptive binning algorithm for robust and efficient genome reconstruction from
metagenome assemblies. PeerJ 7, e7359. https://doi.org/10.7717/peerj.7359
Kembel, S.W., Cowan, P.D., Helmus, M.R., Cornwell, W.K., Morlon, H., Ackerly, D.D.,
Blomberg, S.P., Webb, C.O., 2010. Picante: R tools for integrating phylogenies and
ecology. Bioinformatics 26, 1463–1464.
https://doi.org/10.1093/bioinformatics/btq166
Kovatcheva-Datchary, P., Nilsson, A., Akrami, R., Lee, Y.S., De Vadder, F., Arora, T., Hallen, A.,
Martens, E., Björck, I., Bäckhed, F., 2015. Dietary Fiber-Induced Improvement in
Glucose Metabolism Is Associated with Increased Abundance of Prevotella. Cell
Metab 22, 971–982. https://doi.org/10.1016/j.cmet.2015.10.001
Kundu, S., Bansal, M.S., 2018. On the impact of uncertain gene tree rooting on duplication-
transfer-loss reconciliation. BMC Bioinformatics 19, 290.
https://doi.org/10.1186/s12859-018-2269-0
Langergraber, K.E., Prüfer, K., Rowney, C., Boesch, C., Crockford, C., Fawcett, K., Inoue, E.,
Inoue-Muruyama, M., Mitani, J.C., Muller, M.N., Robbins, M.M., Schubert, G.,
Stoinski, T.S., Viola, B., Watts, D., Wittig, R.M., Wrangham, R.W., Zuberbühler, K.,
Pääbo, S., Vigilant, L., 2012. Generation times in wild chimpanzees and gorillas
suggest earlier divergence times in great ape and human evolution. Proc Natl Acad
Sci U S A 109, 15716–15721. https://doi.org/10.1073/pnas.1211740109
Lawrence, J.G., Retchless, A.C., 2009. The Interplay of Homologous Recombination and
Horizontal Gene Transfer in Bacterial Speciation, in: Gogarten, M.B., Gogarten, J.P.,
Olendzenski, L.C. (Eds.), Horizontal Gene Transfer: Genomes in Flux, Methods in
Molecular Biology. Humana Press, Totowa, NJ, pp. 29–53.
https://doi.org/10.1007/978-1-60327-853-9_3
Lee, J., Yang, J., Zhitnitsky, D., Lewinson, O., Rees, D., 2014. Structural and functional
characterization of a heavy metal detoxifying ABC transporter (997.2). The FASEB
Journal 28, 997.2. https://doi.org/10.1096/fasebj.28.1_supplement.997.2
Li, H., 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34,
3094–3100. https://doi.org/10.1093/bioinformatics/bty191
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352
Lloyd-Price, J., Arze, C., Ananthakrishnan, A.N., Schirmer, M., Avila-Pacheco, J., Poon, T.W.,
Andrews, E., Ajami, N.J., Bonham, K.S., Brislawn, C.J., Casero, D., Courtney, H.,
Gonzalez, A., Graeber, T.G., Hall, A.B., Lake, K., Landers, C.J., Mallick, H., Plichta, D.R.,
Prasad, M., Rahnavard, G., Sauk, J., Shungin, D., Vázquez-Baeza, Y., White, R.A.,
Braun, J., Denson, L.A., Jansson, J.K., Knight, R., Kugathasan, S., McGovern, D.P.B.,
Petrosino, J.F., Stappenbeck, T.S., Winter, H.S., Clish, C.B., Franzosa, E.A., Vlamakis,
H., Xavier, R.J., Huttenhower, C., 2019. Multi-omics of the gut microbial ecosystem in

inflammatory bowel diseases. Nature 569, 655–662.
https://doi.org/10.1038/s41586-019-1237-9
Lozupone, C., Lladser, M.E., Knights, D., Stombaugh, J., Knight, R., 2011. UniFrac: an effective
distance metric for microbial community comparison. ISME J 5, 169–172.
https://doi.org/10.1038/ismej.2010.133
Manara, S., Asnicar, F., Beghini, F., Bazzani, D., Cumbo, F., Zolfo, M., Nigro, E., Karcher, N.,
Manghi, P., Metzger, M.I., Pasolli, E., Segata, N., 2019. Microbial genomes from non-
human primate gut metagenomes expand the primate-associated bacterial tree of
life with over 1000 novel species. Genome Biology 20, 299.
https://doi.org/10.1186/s13059-019-1923-9
Marin, J., Battistuzzi, F.U., Brown, A.C., Hedges, S.B., 2017. The Timetree of Prokaryotes:
New Insights into Their Evolution and Speciation. Molecular Biology and Evolution
34, 437–446. https://doi.org/10.1093/molbev/msw245
Matute, J.D., Duan, J., Flak, M.B., Griebel, P., Tascon-Arcila, J.A., Doms, S., Hanley, T.,
Antanaviciute, A., Gundrum, J., Mark Welch, J.L., Sit, B., Abtahi, S., Fuhler, G.M.,
Grootjans, J., Tran, F., Stengel, S.T., White, J.R., Krupka, N., Haller, D., Clare, S.,
Lawley, T.D., Kaser, A., Simmons, A., Glickman, J.N., Bry, L., Rosenstiel, P., Borisy, G.,
Waldor, M.K., Baines, J.F., Turner, J.R., Blumberg, R.S., 2023. Intelectin-1 binds and
alters the localization of the mucus barrier-modifying bacterium Akkermansia
muciniphila. J Exp Med 220, e20211938. https://doi.org/10.1084/jem.20211938
McCall, L.-I., Callewaert, C., Zhu, Q., Song, S.J., Bouslimani, A., Minich, J.J., Ernst, M., Ruiz-
Calderon, J.F., Cavallin, H., Pereira, H.S., Novoselac, A., Hernandez, J., Rios, R.,
Branch, O.H., Blaser, M.J., Paulino, L.C., Dorrestein, P.C., Knight, R., Dominguez-Bello,
M.G., 2020. Home chemical and microbial transitions across urbanization. Nature
microbiology 5, 108. https://doi.org/10.1038/s41564-019-0593-4
McDonald, B.R., Currie, C.R., 2017. Lateral Gene Transfer Dynamics in the Ancient Bacterial
Genus Streptomyces. mBio 8, e00644-17. https://doi.org/10.1128/mBio.00644-17
McMurdie, P.J., Holmes, S., 2013. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLOS ONE 8, e61217.
https://doi.org/10.1371/journal.pone.0061217
Minh, B.Q., Nguyen, M.A.T., von Haeseler, A., 2013. Ultrafast Approximation for
Phylogenetic Bootstrap. Molecular Biology and Evolution 30, 1188–1195.
https://doi.org/10.1093/molbev/mst024
Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A.,
Lanfear, R., 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. Molecular Biology and Evolution 37, 1530–1534.
https://doi.org/10.1093/molbev/msaa015
Moeller, A.H., Caro-Quintero, A., Mjungu, D., Georgiev, A.V., Lonsdorf, E.V., Muller, M.N.,
Pusey, A.E., Peeters, M., Hahn, B.H., Ochman, H., 2016. Cospeciation of gut
microbiota with hominids. Science 353, 380–382.
https://doi.org/10.1126/science.aaf3951
Moeller, A.H., Li, Y., Mpoudi Ngole, E., Ahuka-Mundeke, S., Lonsdorf, E.V., Pusey, A.E.,
Peeters, M., Hahn, B.H., Ochman, H., 2014. Rapid changes in the gut microbiome
during human evolution. Proceedings of the National Academy of Sciences 111,
16431–16435. https://doi.org/10.1073/pnas.1419136111
Molinaro, A., Bel Lassen, P., Henricsson, M., Wu, H., Adriouch, S., Belda, E., Chakaroun, R.,
Nielsen, T., Bergh, P.-O., Rouault, C., André, S., Marquet, F., Andreelli, F., Salem, J.-E.,

Assmann, K., Bastard, J.-P., Forslund, S., Le Chatelier, E., Falony, G., Pons, N., Prifti, E.,
Quinquis, B., Roume, H., Vieira-Silva, S., Hansen, T.H., Pedersen, H.K., Lewinter, C.,
Sønderskov, N.B., Køber, L., Vestergaard, H., Hansen, T., Zucker, J.-D., Galan, P.,
Dumas, M.-E., Raes, J., Oppert, J.-M., Letunic, I., Nielsen, J., Bork, P., Ehrlich, S.D.,
Stumvoll, M., Pedersen, O., Aron-Wisnewsky, J., Clément, K., Bäckhed, F., 2020.
Imidazole propionate is increased in diabetes and associated with dietary patterns
and altered microbial ecology. Nat Commun 11, 5881.
https://doi.org/10.1038/s41467-020-19589-w
Nishida, A.H., Ochman, H., 2021. Captivity and the co-diversification of great ape
microbiomes. Nat Commun 12, 5632. https://doi.org/10.1038/s41467-021-25732-y
Nissen, J.N., Johansen, J., Allesøe, R.L., Sønderby, C.K., Armenteros, J.J.A., Grønbech, C.H.,
Jensen, L.J., Nielsen, H.B., Petersen, T.N., Winther, O., Rasmussen, S., 2021.
Improved metagenome binning and assembly using deep variational autoencoders.
Nat Biotechnol 39, 555–560. https://doi.org/10.1038/s41587-020-00777-4
Ochman, H., Elwyn, S., Moran, N.A., 1999. Calibrating bacterial evolution. Proceedings of the
National Academy of Sciences 96, 12638–12643.
https://doi.org/10.1073/pnas.96.22.12638
Ochman, H., Worobey, M., Kuo, C.-H., Ndjango, J.-B.N., Peeters, M., Hahn, B.H., Hugenholtz,
P., 2010. Evolutionary Relationships of Wild Hominids Recapitulated by Gut Microbial
Communities. PLOS Biology 8, e1000546.
https://doi.org/10.1371/journal.pbio.1000546
Oksanen, J., Simpson, G.L., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., O'Hara, R.B.,
Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M.,
Bolker, B., Borcard, D., Carvalho, G., Chirico, M., Caceres, M.D., Durand, S.,
Evangelista, H.B.A., FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M.O.,
Lahti, L., McGlinn, D., Ouellette, M.-H., Cunha, E.R., Smith, T., Stier, A., Braak, C.J.F.T.,
Weedon, J., 2022. vegan: Community Ecology Package.
Olm, M.R., Brown, C.T., Brooks, B., Banfield, J.F., 2017. dRep: a tool for fast and accurate
genomic comparisons that enables improved genome recovery from metagenomes
through de-replication. ISME J 11, 2864–2868.
https://doi.org/10.1038/ismej.2017.126
Parks, D.H., Chuvochina, M., Rinke, C., Mussig, A.J., Chaumeil, P.-A., Hugenholtz, P., 2022.
GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based
taxonomy. Nucleic Acids Res 50, D785–D794. https://doi.org/10.1093/nar/gkab776
Pasolli, E., Asnicar, F., Manara, S., Zolfo, M., Karcher, N., Armanini, F., Beghini, F., Manghi, P.,
Tett, A., Ghensi, P., Collado, M.C., Rice, B.L., DuLong, C., Morgan, X.C., Golden, C.D.,
Quince, C., Huttenhower, C., Segata, N., 2019. Extensive Unexplored Human
Microbiome Diversity Revealed by Over 150,000 Genomes from Metagenomes
Spanning Age, Geography, and Lifestyle. Cell 176, 649-662.e20.
https://doi.org/10.1016/j.cell.2019.01.001
Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., Kingsford, C., 2017. Salmon: fast and bias-
aware quantification of transcript expression using dual-phase inference. Nat
Methods 14, 417–419. https://doi.org/10.1038/nmeth.4197
R Core Team, 2022. R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria.

Ridley, C.P., Lee, H.Y., Khosla, C., 2008. Evolution of polyketide synthases in bacteria.
Proceedings of the National Academy of Sciences 105, 4595–4600.
https://doi.org/10.1073/pnas.0710107105
Rodríguez-Daza, M.C., Pulido-Mateos, E.C., Lupien-Meilleur, J., Guyonnet, D., Desjardins, Y.,
Roy, D., 2021. Polyphenol-Mediated Gut Microbiota Modulation: Toward Prebiotics
and Further. Frontiers in Nutrition 8.
Rühlemann, M.C., Wacker, E.M., Ellinghaus, D., Franke, A., 2022. MAGScoT - a fast,
lightweight, and accurate bin-refinement tool. Bioinformatics btac694.
https://doi.org/10.1093/bioinformatics/btac694
Scally, A., Dutheil, J.Y., Hillier, L.W., Jordan, G.E., Goodhead, I., Herrero, J., Hobolth, A.,
Lappalainen, T., Mailund, T., Marques-Bonet, T., McCarthy, S., Montgomery, S.H.,
Schwalie, P.C., Tang, Y.A., Ward, M.C., Xue, Y., Yngvadottir, B., Alkan, C., Andersen,
L.N., Ayub, Q., Ball, E.V., Beal, K., Bradley, B.J., Chen, Y., Clee, C.M., Fitzgerald, S.,
Graves, T.A., Gu, Y., Heath, P., Heger, A., Karakoc, E., Kolb-Kokocinski, A., Laird, G.K.,
Lunter, G., Meader, S., Mort, M., Mullikin, J.C., Munch, K., O'Connor, T.D., Phillips,
A.D., Prado-Martinez, J., Rogers, A.S., Sajjadian, S., Schmidt, D., Shaw, K., Simpson,
J.T., Stenson, P.D., Turner, D.J., Vigilant, L., Vilella, A.J., Whitener, W., Zhu, B.,
Cooper, D.N., de Jong, P., Dermitzakis, E.T., Eichler, E.E., Flicek, P., Goldman, N.,
Mundy, N.I., Ning, Z., Odom, D.T., Ponting, C.P., Quail, M.A., Ryder, O.A., Searle,
S.M., Warren, W.C., Wilson, R.K., Schierup, M.H., Rogers, J., Tyler-Smith, C., Durbin,
R., 2012. Insights into hominid evolution from the gorilla genome sequence. Nature
483, 169–175. https://doi.org/10.1038/nature10842
Schaan, A.P., Sarquis, D., Cavalcante, G.C., Magalhães, L., Sacuena, E.R.P., Costa, J., Fonseca,
D., Mello, V.J., Guerreiro, J.F., Ribeiro-dos-Santos, Â., 2021. The structure of Brazilian
Amazonian gut microbiomes in the process of urbanisation. npj Biofilms
Microbiomes 7, 1–12. https://doi.org/10.1038/s41522-021-00237-0
Schliep, K.P., 2011. phangorn: phylogenetic analysis in R. Bioinformatics 27, 592–593.
https://doi.org/10.1093/bioinformatics/btq706
Sievers, F., Wilm, A., Dineen, D., Gibson, T.J., Karplus, K., Li, W., Lopez, R., McWilliam, H.,
Remmert, M., Söding, J., Thompson, J.D., Higgins, D.G., 2011. Fast, scalable
generation of high-quality protein multiple sequence alignments using Clustal
Omega. Mol Syst Biol 7, 539. https://doi.org/10.1038/msb.2011.75
Sonnenburg, E.D., Sonnenburg, J.L., 2019. The ancestral and industrialized gut microbiota
and implications for human health. Nat Rev Microbiol 17, 383–390.
https://doi.org/10.1038/s41579-019-0191-8
Sonnenburg, J.L., Sonnenburg, E.D., 2019. Vulnerability of the industrialized microbiota.
Science 366, eaaw9255. https://doi.org/10.1126/science.aaw9255
Steinegger, M., Söding, J., 2017. MMseqs2 enables sensitive protein sequence searching for
the analysis of massive data sets. Nat Biotechnol 35, 1026–1028.
https://doi.org/10.1038/nbt.3988
Suzuki, T.A., Fitzstevens, J.L., Schmidt, V.T., Enav, H., Huus, K.E., Mbong Ngwese, M.,
Grießhammer, A., Pfleiderer, A., Adegbite, B.R., Zinsou, J.F., Esen, M., Velavan, T.P.,
Adegnika, A.A., Song, L.H., Spector, T.D., Muehlbauer, A.L., Marchi, N., Kang, H.,
Maier, L., Blekhman, R., Ségurel, L., Ko, G., Youngblut, N.D., Kremsner, P., Ley, R.E.,
2022. Codiversification of gut microbiota with humans. Science 377, 1328–1332.
https://doi.org/10.1126/science.abm7759

Tange, O., 2011. GNU Parallel - The Command-Line Power Tool. ;login: The USENIX Magazine
36, 42–47.
Thingholm, L.B., Bang, C., Rühlemann, M.C., Starke, A., Sicks, F., Kaspari, V., Jandowsky, A.,
Frölich, K., Ismer, G., Bernhard, A., Bombis, C., Struve, B., Rausch, P., Franke, A.,
2021. Ecology impacts the decrease of Spirochaetes and Prevotella in the fecal gut
microbiota of urban humans. BMC microbiology 21. https://doi.org/10.1186/s12866-
021-02337-5
Tsujikawa, Y., Ishikawa, S., Sakane, I., Yoshida, K., Osawa, R., 2021. Identification of genes
encoding a novel ABC transporter in Lactobacillus delbrueckii for inulin polymers
uptake. Sci Rep 11, 16007. https://doi.org/10.1038/s41598-021-95356-1
Tung, J., Barreiro, L.B., Burns, M.B., Grenier, J.-C., Lynch, J., Grieneisen, L.E., Altmann, J.,
Alberts, S.C., Blekhman, R., Archie, E.A., 2015. Social networks predict gut
microbiome composition in wild baboons. eLife 4, e05224.
https://doi.org/10.7554/eLife.05224
Valles-Colomer, M., Blanco-Míguez, A., Manghi, P., Asnicar, F., Dubois, L., Golzato, D.,
Armanini, F., Cumbo, F., Huang, K.D., Manara, S., Masetti, G., Pinto, F., Piperni, E.,
Punčochář, M., Ricci, L., Zolfo, M., Farrant, O., Goncalves, A., Selma-Royo, M., Binetti,
A.G., Becerra, J.E., Han, B., Lusingu, J., Amuasi, J., Amoroso, L., Visconti, A., Steves,
C.M., Falchi, M., Filosi, M., Tett, A., Last, A., Xu, Q., Qin, N., Qin, H., May, J., Eibach,
D., Corrias, M.V., Ponzoni, M., Pasolli, E., Spector, T.D., Domenici, E., Collado, M.C.,
Segata, N., 2023. The person-to-person transmission landscape of the gut and oral
microbiomes. Nature 614, 125–135. https://doi.org/10.1038/s41586-022-05620-1
Valles-Colomer, M., Falony, G., Darzi, Y., Tigchelaar, E.F., Wang, J., Tito, R.Y., Schiweck, C.,
Kurilshikov, A., Joossens, M., Wijmenga, C., Claes, S., Van Oudenhove, L.,
Zhernakova, A., Vieira-Silva, S., Raes, J., 2019. The neuroactive potential of the
human gut microbiota in quality of life and depression. Nat Microbiol 4, 623–632.
https://doi.org/10.1038/s41564-018-0337-x
Vals-Delgado, C., Alcala-Diaz, J.F., Molina-Abril, H., Roncero-Ramos, I., Caspers, M.P.M.,
Schuren, F.H.J., Van den Broek, T.J., Luque, R., Perez-Martinez, P., Katsiki, N.,
Delgado-Lista, J., Ordovas, J.M., van Ommen, B., Camargo, A., Lopez-Miranda, J.,
2021. An altered microbiota pattern precedes Type 2 diabetes mellitus development:
From the CORDIOPREV study. J Adv Res 35, 99–108.
https://doi.org/10.1016/j.jare.2021.05.001
van den Boogaart, K.G., Tolosana-Delgado, R., 2008. "compositions": A unified R package to
analyze compositional data. Computers & Geosciences 34, 320–338.
https://doi.org/10.1016/j.cageo.2006.11.017
Vangay, P., Johnson, A.J., Ward, T.L., Al-Ghalith, G.A., Shields-Cutler, R.R., Hillmann, B.M.,
Lucas, S.K., Beura, L.K., Thompson, E.A., Till, L.M., Batres, R., Paw, B., Pergament, S.L.,
Saenyakul, P., Xiong, M., Kim, A.D., Kim, G., Masopust, D., Martens, E.C.,
Angkurawaranon, C., McGready, R., Kashyap, P.C., Culhane-Pera, K.A., Knights, D.,
2018. US Immigration Westernizes the Human Gut Microbiome. Cell 175, 962-
972.e10. https://doi.org/10.1016/j.cell.2018.10.029
Vieira-Silva, S., Sabino, J., Valles-Colomer, M., Falony, G., Kathagen, G., Caenepeel, C.,
Cleynen, I., van der Merwe, S., Vermeire, S., Raes, J., 2019. Quantitative microbiome
profiling disentangles inflammation- and bile duct obstruction-associated microbiota
alterations across PSC/IBD diagnoses. Nat Microbiol 4, 1826–1831.
https://doi.org/10.1038/s41564-019-0483-9

Wallenborn, J.T., Vonaesch, P., 2022. Intestinal microbiota research from a global
perspective. Gastroenterology Report 10, goac010.
https://doi.org/10.1093/gastro/goac010
Watanabe, F., Bito, T., 2018. Vitamin B12 sources and microbial interaction. Exp Biol Med
(Maywood) 243, 148–158. https://doi.org/10.1177/1535370217746612
Whelan, S., Goldman, N., 2001. A General Empirical Model of Protein Evolution Derived
from Multiple Protein Families Using a Maximum-Likelihood Approach. Molecular
Biology and Evolution 18, 691–699.
https://doi.org/10.1093/oxfordjournals.molbev.a003851
Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemund, G.,
Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Bache, S.M.,
Müller, K., Ooms, J., Robinson, D., Seidel, D.P., Spinu, V., Takahashi, K., Vaughan, D.,
Wilke, C., Woo, K., Yutani, H., 2019. Welcome to the Tidyverse. Journal of Open
Source Software 4, 1686. https://doi.org/10.21105/joss.01686
Wu, Y.-W., Simmons, B.A., Singer, S.W., 2016. MaxBin 2.0: an automated binning algorithm
to recover genomes from multiple metagenomic datasets. Bioinformatics 32, 605–
607. https://doi.org/10.1093/bioinformatics/btv638
Yu, G., Smith, D.K., Zhu, H., Guan, Y., Lam, T.T.-Y., 2017. ggtree: an r package for visualization
and annotation of phylogenetic trees with their covariates and other associated
data. Methods in Ecology and Evolution 8, 28–36. https://doi.org/10.1111/2041-
210X.12628
Zheng, L., Kelly, C.J., Colgan, S.P., 2015. Physiologic hypoxia and oxygen homeostasis in the
healthy intestine. A Review in the Theme: Cellular Responses to Hypoxia. Am J
Physiol Cell Physiol 309, C350–C360. https://doi.org/10.1152/ajpcell.00191.2015
Zheng, P., Li, Z., Zhou, Z., 2018. Gut microbiome in type 1 diabetes: A comprehensive review.
Diabetes/Metabolism Research and Reviews 34, e3043.
https://doi.org/10.1002/dmrr.3043
1375
1376