Logic-based mechanistic machine learning on high-content images reveals how drugs differentially regulate cardiac fibroblasts

Anders R. Nelson[1], Steven L. Christiansen[1,2], Kristen M. Naegle[1], Jeffrey J. Saucerman[1*]

1) University of Virginia School of Medicine, Charlottesville, VA 22903

2) Brigham Young University Department of Biochemistry, Provo, UT 84602

*University of Virginia School of Medicine, Charlottesville, VA 2290

(e-mail: jsaucerman@virginia.edu)

PO Box 800759, Charlottesville, VA 22908, 434-924-5095

Classification: Biological Sciences; Systems Biology

Keywords: Fibrosis, Fibroblast, Machine learning, Cell Biology, Differentiation, Drugs

Abstract

Fibroblasts are essential regulators of extracellular matrix deposition following cardiac injury. These cells exhibit highly plastic responses in phenotype during fibrosis in response to environmental stimuli. Here, we test whether and how candidate anti-fibrotic drugs differentially regulate measures of cardiac fibroblast phenotype, which may help identify treatments for cardiac fibrosis. We conducted a high content microscopy screen of human cardiac fibroblasts treated with 13 clinically relevant drugs in the context of TGFβ and/or IL-1β, measuring phenotype across 137 single-cell features. We used the phenotypic data from our high content imaging to train a logic-based mechanistic machine learning model (LogiMML) for fibroblast signaling. The model predicted how pirfenidone and Src inhibitor WH-4-023 reduce F-actin assembly and F-actin stress fiber formation, respectively. Validating the LogiMML model prediction that PI3K partially mediates the effects of Src inhibition, we found that PI3K inhibition reduces F-actin fiber formation and procollagen I production in human cardiac fibroblasts. In this study, we establish a modeling approach combining the strengths of logic-based network models and regularized regression models, apply this approach to predict mechanisms that mediate the differential effects of drugs on fibroblasts, revealing Src inhibition acting via PI3K as a potential therapy for cardiac fibrosis.

Significance

Cardiac fibrosis is a dysregulation of the normal wound healing response, resulting in excessive scarring and cardiac dysfunction. As cardiac fibroblasts primarily regulate this process, we explored how candidate anti-fibrotic drugs alter the fibroblast phenotype. We identify a set of 137 phenotypic features that change in response to drug treatments. Using a new computational modeling approach termed logic-based mechanistic machine learning, we predict how pirfenidone and Src inhibition affect the regulation of the phenotypic features F-actin assembly and F-actin stress fiber formation. We also show that inhibition of PI3K reduces F-actin fiber formation and procollagen I production in human cardiac fibroblasts, supporting a role for PI3K as a mechanism by which Src inhibition may suppress fibrosis.

Introduction

Cardiac fibroblasts are the primary regulators of remodeling following cardiac injury[1]. Extracellular matrix (ECM) deposition by activated myofibroblasts is essential to this response, but excessive deposition can lead to ventricular stiffness, diastolic dysfunction, and heart failure[1]. While fibroblasts are critical to the wound healing response, current standard-of-care

therapeutics for cardiac injury, such as myocardial infarction (MI), affect downstream
symptoms but do not specifically target fibroblast signaling[2]. Recent drug discovery and
development has focused on identifying drugs such as Entresto (sacubitril/valsartan) that
reduce fibrosis in part by modulating fibroblast signaling[3,4].

Collagen secretion, αSMA expression, and F-actin assembly are traditional markers for a
profibrotic fibroblast phenotype[5,6]. While high expression of these markers provides an initial
indication of myofibroblast activation, traditional marker expression is inconsistent and does
not fully capture the fibrotic response[7]. Recent studies of fibroblast phenotype have shown that
fibroblasts exhibit high phenotypic heterogeneity across many facets in response to injury, and
that phenotypic changes are also sensitive to drug perturbations[8–11]. Identifying drugs that
regulate fibroblast signaling may provide targeted control of fibrosis.

Previously, we developed a logic-based mechanistic network model of fibroblast signaling and
applied it to perform virtual screens for anti-fibrotic drugs[12,13]. That study predicted and
experimentally validated an antifibrotic role for the TGFβ receptor inhibitor galunisertib[13].
While the fibroblast network model predicts a number of drugs that modulate fibroblast
activation, substantial experimental characterization is needed to capture phenotypic
responses to drugs that were not captured by prior modeling.

In this study, we combined high content microscopy, network modeling, and machine learning
to identify drugs that differentially regulate fibroblast phenotypic metrics and predict their
underlying network mechanisms. We used image-based feature extraction to more deeply
characterize drug response and fibroblast phenotype, capturing drug-induced changes across a
set of single-cell metrics relevant to fibrosis. Using a novel logic-based mechanistic machine
learning approach, LogiMML, we predicted signaling pathways that determine how drugs
regulate fibroblast phenotype. Finally, we experimentally validated the main pathway
mechanism predicted by the LogiMML model that mediates how Src inhibition suppresses
fibrotic responses.

Results

An *in vitro* screen for candidate fibrosis drugs

Previously, we applied our published cardiac fibroblast network model[12] to identify candidate
therapies predicted to reduce cardiac fibrosis[13]. This logic-based differential equation network
model was developed from a wide range of fibroblast signaling relationships from *in vitro*
studies in the literature. The model predicts changes in fibrotic outputs including collagen I and
III, αSMA, EDA fibronectin, matrix metalloproteases, and F-actin in response to changes in
extracellular signaling contexts and drug treatment[12]. This model was previously integrated
with the drug-target database DrugBank to make predict the response of fibroblasts to 121
FDA-approved or investigational drugs that have targets in this network[13].

To expand upon the *in silico* modeling work done in that previous study[13], we aimed to develop
a list of drug candidates to test experimentally for their ability to reduce fibrosis in cardiac
fibroblasts *in vitro*. As the model predicted many drugs to reduce fibrosis effectively[13], we
included drug selection criteria outside of our modeling results alone to further narrow-down a
list of candidate drugs. First, we prioritized pathway diversity of the drug targets to ensure that
we would perturb fibrotic signaling comprehensively and avoid testing redundant drugs in our
experiments. As drug repurposing has become an increasingly effective and efficient strategy
for treating cardiovascular disease, we next looked to prioritize drugs that had previous clinical
indications for other disease areas[14,15]. Using these criteria, we selected thirteen drugs to
evaluate experimentally: anakinra, valsartan, defactinib, HW-4-023, glutathione, CW-HM12,
salbutamol, marimistat, fasudil, SB203580, pirfenidone, brain natriuretic peptide (BNP), and a
combination of valsartan and BNP (Table S1). Among the list of candidate drug targets are
regulators for inflammatory signaling, mechanical stretch response, non-canonical TGFβ
signaling, and modification of secreted proteins.

We next aimed to test these candidate drugs for their ability to quantitatively reduce fibrosis as
characterized by image-based single-cell profiling of procollagen I, α-smooth muscle actin
(αSMA), and F-actin. In injury signaling conditions, such as following myocardial infarction (MI),
myocardial cells are exposed to elevated proinflammatory and profibrotic stimuli[16–18]. To
represent these signaling contexts in an *in vitro* system, we included IL1β and TGFβ, shown to
be elevated following cardiac injury, in our treatment conditions to represent proinflammatory
and profibrotic contexts respectively[19–21]. We tested our candidate drugs under four total
cytokine contexts (baseline context with no added cytokine, fibrotic context represent by TGFβ,
inflammatory context represented by IL1β, and combined context represent by both TGFβ and
IL1β)[19–21]. In total, we used 108 treatment conditions consisting of one of the thirteen drugs at
a low, medium, or high dose combined with one of the four cytokine contexts. We also included
treatments of each cytokine context with no drug to establish a control baseline for cell
responses to cytokines. We imaged and quantified single-cell protein expression of three
fibrotic markers, procollagen I, α-smooth muscle actin (αSMA), and F-actin using high-content
microscopy and a custom CellProfiler software pipeline[22].

Interestingly, the antifibrotic drugs in our screen induced differential effects on fibrosis. Of the
13 candidate drugs, WH-4-023, fasudil, and defactinib caused the strongest reduction of
procollagen I, F-actin, and αSMA expression in a TGFβ signaling context, even at the lowest
dose (Figure 1A). Conversely, a second set of drugs including anakinra and glutathione
increased fibrotic marker expression in both TGFβ and combined TGFβ/IL1β contexts when
applied directly to fibroblasts. In a previous clinical study, anakinra, an IL1 receptor inhibitor,
was shown to improve cardiac function and reduced heart failure incidence following acute MI
in human patients[23]. While anakinra has been shown to reduce infarct scar area in a mouse MI
model, it also exhibits other beneficial cardiac effects post-MI including inhibition of post-MI
myocyte apoptosis and reduction in systemic inflammation[24,25]. Based on these previous
studies, it is likely that anakinra has a net antifibrotic effect on fibroblasts in the presence of

other myocardial cell types even though anakinra treatment increased fibrotic marker

expression in this experiment. A third set of drugs showed more selective antifibrotic effects.

For example, while fasudil significantly reduced expression of all three fibrosis markers in a

TGFβ signaling context, pirfenidone only significantly reduced F-actin (Figure 1 B-E). This third

set of drugs is of particular interest as it contains drugs that differentially regulate markers for

fibrosis. Given the recent clinical effectiveness of pirfenidone for lung fibrosis, and success in

diseases models for cardiac fibrosis[26,27], we further investigated the mechanisms by which it

regulates F-actin in cardiac fibroblasts.

Figure 1: **High-content microscopy screen for drugs that module fibroblast activation.** A) Expression of fibroblast activation markers procollagen I, F-actin, and αSMA in human cardiac fibroblasts upon treatment of 13 drugs at 3 doses, under environmental contexts of TGFβ, IL1β, or both. Fold change values show 'drug vs. no drug' Integrated Intensities for each protein. Panels B and C show quantification and representative images of the effects of pirfenidone, a non-specific inhibitor of TGFβ expression, which consistently regulates fibrotic protein expression. Panels D and E show quantification and representative images of the effects of fasudil, a Rho-kinase inhibitor, which differentially regulates fibrotic protein expression. *p≤0.05 ANOVA with Tukey's post-hoc.

LogiMML: logic-based mechanistic machine learning model predicts how drugs regulate fibroblast phenotype

Assembled F-actin filaments play a key role in contractility as fibroblasts transition to become
myofibroblasts[28]. Therefore, we asked whether the previous mechanistic computational model
of the fibroblast signaling network[12] could predict our experimentally measured inhibition of F-
actin by pirfenidone from Figure 1D. While the model had correctly predicted responses to a
number of drugs including galunisertib[13], here, the original mechanistic model did not capture
the ability of pirfenidone to suppress F-actin in a TGFβ signaling context (Figure 2 A).

Given the limitations of a model based only on prior knowledge, we asked whether drug
predictions could be improved by combining the mechanistic model with a machine learning
model that leverages data from the drug screen. Motivated by 'white-box' machine learning
strategies that combine mechanistic models with machine learning[29,30], we designed a logic-
based mechanistic machine learning (LogiMML) model to predict key regulators that conduct
signaling from network model inputs and simulated drugs to experimentally measured
phenotypic outputs (Figure 2 B). As the 108 treatments were insufficient to infer new links to
phenotypic outputs from all 91 model nodes, we reduced the model's dimensionality by
clustering nodes into modules. Eleven signaling modules were computed based on a combined
influence and sensitivity analysis, grouping nodes with similar predicted behavior across
signaling contexts. The machine learning component was then trained by mapping the model-
predicted activity of each network module for each of the 108 drug+cytokine treatments to
respective experimentally measured outputs. Regularized ridge regression was selected for the
machine learning layer of the LogiMML model to reduce the likelihood of overfitting[31]. As
measured experimentally, the LogiMML model correctly predicted the respective induction and
suppression of F-actin by TGFβ and pirfenidone (Figure 2 C).

We next asked whether the LogiMML model could provide new mechanistic insights into how
F-actin is regulated by pirfenidone. First, we used the LogiMML model's ridge regression
coefficients to predict the modules that most influence F-actin. 'PI3K' and 'Smad3' modules
were predicted to be the top positive regulators of F-actin, while the 'P38_Calcium' module was
predicted as the top negative regulator (Figure 2 E). These predictions for fibroblasts are
consistent with previous studies with other cell types showing that members of the 'Smad3'

 and 'P38_Calcium' signaling modules regulate F-actin filament assembly in endothelial cells and

Figure 2: **LogiMML logic-based mechanistic machine learning approach guides model revision and predicts network mechanisms underlying pirfenidone suppression of F-actin.** A) Original fibroblast network model predicts no change in F-actin upon TGFβ or pirfenidone treatment. Experimental data shows pirfenidone significantly reverses the increase of F-actin by TGFβ (data previously shown in figure 1 D). B) Schematic of the LogiMML approach for integrating logic-based network modeling with machine learning to predict network mechanisms for cell phenotypes. The average activity within each network module is mapped to predict fibroblast phenotypic features via a Ridge regression layer. C) The Coupled LogiMML model predicts TGFβ and pirfenidone effects on F-actin that qualitatively match experimental data shown in panel A. D) LogiMML ridge regression coefficients show predicted relative influence of network modules on F-actin. E) LogiMML node knockdown sensitivity analysis in the context of TGFβ+pirfenidone. Nodes from most influential modules are sequentially knocked down, predicting change in F-actin upon knockdown. F) Schematic of the network mechanisms predicted for the actions of pirfenidone on F-actin, derived from sensitivity analysis in panel E.

that members of the 'PI3K' signaling module promote actin filament remodeling during
migration in embryonic fibroblasts [32–34]. To identify which individual signaling nodes within
these three modules most regulate F-actin, we performed a virtual knockdown screen of the
LogiMML model for regulators of F-actin in a 'TGFβ+pirfenidone' signaling context (Figure 2 E).
Summarizing these analyses, the LogiMML model predicts that pirfenidone regulation of F-actin
is positively regulated by p38, Akt, and CBP, while negatively regulated by ROS and NOX (Figure
2 F).

Drugs and pathways controlling fibroblast morphology and texture
Given the differential regulation of fibrosis marker protein expression, we asked whether other
aspects of fibroblast phenotype may also be differentially regulated by drugs and cytokines.
Qualitatively, we observed morphological changes in cell shape, stress fiber formation,
intracellular protein distribution, and cell area (e.g. for pirfenidone treatment see Figure 1 E).
To measure these characteristics of fibroblast phenotype, we developed a custom CellProfiler
image analysis pipeline quantifying 137 total single-cell cell features[22,35]. Integrated intensities
for the three fibrotic marker proteins, procollagen I, F-actin, and αSMA clustered relatively
close to each other across the feature space (Figure 3 A). As expected, expression of these
marker proteins and similar features were high under TGFβ and TGFβ-like treatments, and low
under negative control and IL1β conditions. While the central rows of the heatmap contain
many features with similar treatment responses, the features at the top and bottom regions of
the heatmap show high heterogeneity in response to drugs.

To gain a comprehensive understanding of fibroblast phenotypic responses to drugs and
cytokines, we applied principal component analysis (PCA) dimensionality reduction to the data
(Figure 3 B-C). To mitigate feature redundancy and improve PCA performance, we first reduced
the feature set from 137 total features to 18 representative features. These features were

selected by clustering a correlation matrix of the feature set into 15 clusters, selecting one
feature from each cluster (Figure S1, Figure S2) as key representatives of that cluster's
information. The three integrated intensity features for procollagen I, F-actin, and αSMA were
also retained.

Negative control treatments had a negative score on the first principal component (PC1), while
cells treated with TGFβ showed a high positive score on PC1, indicating that the first principal
component correlates with an axis of classical fibroblast activation (Figure 3 B, Figure S3 A). This
was further supported by the PCA loading values for integrated procollagen I, F-actin, and
αSMA (Figure 3 C, Figure S3 B). These three features are expected to be relatively high in
activated myofibroblasts and indeed have strong positive loadings on PC1. On the PCA scores,

any of the 'TGFβ + Drug' groups deviated from the control-TGFβ axis defined on PC1, implying

Figure 3: **Survey of single-cell fibroblast phenotypic features in response to 13 drugs at 3 doses and 4 environmental contexts.** A) 137 single-cell fibroblast features that quantify protein intensity, protein localization, cell morphology, and fiber texture. This heatmap was organized on treatment and feature axes by agglomerative hierarchical clustering. B) Principal component scores of experimental data reduced to a set of 18 representative fibroblast features. C) Principal component loadings the reduced of PCA scores and loadings define a primary axis of fibroblast activation with correlated protein expression of procollagen, αSMA, and F-actin that is modulated by many drugs. Off-axis, the Src inhibitor WH-4-023 modulated the cell texture feature Actin Long Angular Second Moment, which motivated further study.

that drugs induce phenotypic changes distinct from a simple reversal of TGFβ's effects. To
further investigate drug-induced changes in phenotype, we analyzed the PCA scores and
loadings to infer links between drugs and the features they regulate. Notably, the Src inhibitor
WH-4-023 (WH) showed directionality on the scores plot similar to that of Actin Long Angular
Second Moment (Actin Long ASM, a measure for F-actin uniformity) on the loadings plot. Actin
stress fibers, composed of multiple F-actin filaments along with other proteins, contribute to
pathological fibrosis and myofibroblast differentiation[36–38]. This feature and treatment pair
showed a negative value on PC1 and a positive value on PC2 relative to the TGFβ and control
groups, respectively. The similar directionality of WH and Actin Long ASM suggests that Src
inhibition may modulate F-actin uniformity.

Based on the initial inference from the PCA, we revisited the images from the high-content
microscopy experiment. Fibroblasts treated with TGFβ exhibited discrete F-actin stress fibers,
and stress fibers were qualitatively reduced when WH-4-023 (WH) was added (Figure 4 A).
Quantitative analysis of F-actin uniformity (inversely correlated with stress fibers) using Actin
Long Angular Second Moment (ASM) further supported that TGFβ increased and Src inhibitor
WH reduced F-actin uniformity (Figure 4 B).

To predict the signaling pathways that specifically regulate F-actin stress fibers, we again
applied the LogiMML coupled modeling approach, but this time training the ridge regression
layer of the model on experimental measurements of Actin Long ASM. The LogiMML model
regression coefficients predicted that the 'Mechanical' module was the top positive regulator of
Actin Long ASM and that the 'PI3K' module was the top negative regulator of Actin Long ASM
(Figure 4 C). To identify which individual signaling nodes within these two modules most
regulate Actin Long ASM, we performed a virtual knockdown screen of the LogiMML network
model for regulators of Actin Long ASM in the context of 'TGFβ+WH-4-023' and predicted that
Rho, MKK4, and Akt are proximal regulators of Actin Long ASM and actin stress fiber formation
(Figure 4 D-E).

A

Control TGFβ TGFβ+WH

75 um

B

Actin Long Angular Second Moment

×10⁻³

TGFβ — + +
WH — — +

C

Relative influence of modules on
Actin Long Angular Second Moment

Smad3
Mech
Ras_Raf
Autocrine
PI3K
STAT
P38_Calcium
PKA
PDGF
MKK3
NP

-0.5 -0.4 -0.3 -0.2 -0.1 0.0 0.1 0.2 0.3

D

Knockdown sensitivity for WH+TGFB context

0.10

0.05

0.00

-0.05

-0.10

MKK4 Rac1 ROCK FA SRF FAK TRAF Rho Akt

Module
PI3K
Mechanical

E

WH-4-023

Signaling Modules

Mech
Rho
MKK4

TNFa
PI3K
Akt

Actin Long Angular
Second Moment

F

200 um

Control LY

G

Actin Long Angular Second Moment

×10⁻⁴

LY — +

F-actin Intensity (AU)

LY — +

Procollagen I Intensity (AU)

LY — +

Figure 4: **Logic-based mechanistic machine learning predicts the PI3K module to mediate how Src inhibitor suppresses stress fibers, validated by subsequent experiments.** A) Images of human cardiac fibroblasts treated with baseline control stimulus, TGFβ, or TGFβ + WH-4-023. B) Quantification of Actin Long Angular Second Moment (ASM), a measure of F-actin uniformity and reduced stress fibers based on images in panel A. C) Regression coefficients from the LogiMML mechanistic machine learning model that predicts network modules that regulate actin long ASM. D) Knockdown sensitivity analysis predicting individual proteins that regulate actin long ASM in the TGFβ+WH-4-023 signaling context. E) Signaling schematic for WH-4-023 effect on actin long ASM, derived from sensitivity analysis in panel D. F) Human cardiac fibroblasts treated with PI3K inhibitor LY294002 or baseline control stimulus, measuring F-actin and procollagen expression. G) Quantification of long actin Angular Second Moment (measure of F-actin uniformity), F-actin integrated intensity, and Procollagen I integrated intensity. *$p \leq 0.05$ ANOVA with Tukey's post-hoc in panel B, and *$p \leq 0.05$ Student's T-test in panel G.

PI3K signaling stimulates F-actin stress fiber formation and collagen expression

After deriving a putative signaling schematic for Actin Long ASM using the LogiMML model, we
aimed to experimentally validate the prediction that inhibition of PI3K/Akt would suppress
stress fiber formation and thereby increase Actin Long ASM (Figure 4 E). In previous studies
using PI3K inhibitors, PI3K was shown to regulate fibroblast contractility, fibroblast-to-
myofibroblast transition, and TGFβ-induced αSMA and collagen production[39,40]. Given these
previously implicated roles for PI3K in myofibroblast activation and fibrosis, we investigated if
PI3K regulates F-actin stress fiber formation in cardiac fibroblasts. We treated human cardiac
fibroblasts with either a negative control condition or a 20 μM dose of the PI3K inhibitor
LY294002 (LY). Treatment with LY significantly increased Actin Long ASM, but notably, it had no
significant effect on the total assembly of F-actin in each cell, measured by integrated F-actin
intensity (Figure 4 F-G). This selective effect of PI3K inhibition on stress fiber formation, while
having no significant effect on total F-actin, suggests that F-actin assembly and stress fiber
formation are differentially regulated processes. PI3K inhibition also significantly reduced
integrated procollagen I intensity, demonstrating a role for PI3K signaling in cardiac fibroblast
collagen production (Figure 4G).

Discussion

Cardiac fibroblasts are central regulators and promising therapeutic targets following cardiac
injury. To identify how clinically relevant drugs regulate diverse aspects of fibroblast
phenotype, we performed high-content screening of 13 drugs in 4 environmental contexts. We
expanded our high-content microscopy feature set to 137 single-cell features, measuring
fibrotic marker protein intensity, intracellular protein distribution, fiber texture, and cell
morphology. After reducing the feature space and dimensionality of our experimental data, we
found that many aspects of fibroblast phenotype are uniquely induced by drug and cytokine
treatments. Notably, when administered with TGFβ, the drugs WH-4-023, defactinib, fasudil,
and pirfenidone induced phenotypes that deviated from the PCA axis corresponding to classical

TGFβ response. The differences between these phenotypes can be partially explained by
differential drug regulation of features capturing procollagen I and αSMA expression, and F-
actin assembly and stress fiber formation. To predict how drugs regulate cell signaling and
influence phenotype, we developed the logic-based mechanistic machine learning (LogiMML)
approach which coupled the logic-based fibroblast network model with a ridge regression
model trained on the high-content drug screen. Using this expanded LogiMML model, we
predicted regulatory mechanisms for pirfenidone and Src inhibitor WH-4-023 on F-actin. We
predicted that pirfenidone regulates F-actin assembly via the 'P38_Calcium', 'Smad3', and 'PI3K'
signaling modules, with Akt, p38, and CBP predicted to be positive drivers of F-actin assembly
within these modules. We also predicted that WH-4-023 regulates F-actin stress fiber formation
via the 'PI3k' and 'Mechanical' signaling modules. As predicted by the LogiMML model, we
experimentally validated that PI3K inhibition reduces F-actin stress fiber formation in human
cardiac fibroblasts. These studies validate the ability of the LogiMML approach to predict
signaling mechanisms from a phenotypic screen.

Differential regulation of fibroblast phenotype by drugs and the development of targeted
antifibrotic therapies
Drugs that specifically target fibroblast signaling may provide directed control over the fibrotic
response. A major challenge in therapeutic development for fibrosis is that many drugs capable
of reducing fibrosis target non-specific regulatory pathways outside of the fibrotic response. For
example, the ALK5 inhibitor galunisertib targets the TGFβ receptor and shows promising
therapeutic reduction of fibrosis across organs[41–43]. While TGFβ receptor inhibition can reduce
fibrosis, recent efforts in target discovery have successfully identified new approaches to
mitigate fibrosis that are more fibroblast specific. For example, it was shown that activating
fibroblast-specific TLR4 in mice can drive the development of skin and lung fibrosis and that
TLR4 inhibition reduces αSMA expression and collagen production in fibroblasts [44]. Another
study showed that fibroblast-specific knockout of STAT3 ameliorates skin fibrosis, and that
pharmacological inhibition of STAT3 successfully reduces myofibroblast activation, collagen
accumulation, and dermal thickening in experimental fibrosis in mice[45]. Future work can
advance our understanding of how candidate drugs regulate specific components of the fibrotic
response in fibroblasts and provide targeted control of fibrosis.

Features of cardiac fibroblast phenotype
Following the reduction of the original set of 137 single-cell features from our high content
image analysis, we identified a set of 18 phenotypic features of fibroblasts that exhibit high
heterogeneity in response to drug treatments (Figure S 2, Table S 2). Notably, many of the
features measuring fiber texture for αSMA and F-actin show different response patterns
compared to features measuring overall expression level for those respective proteins (i.e.
aSMA integrated intensity versus aSMA long correlation). This distribution of features indicates
that the expression and organization of aSMA and F-actin are independently regulated by
candidate drugs. The processes of αSMA protein expression and fiber assembly have different

degrees of contribution to pathological fibrosis. For example, a recent study showed that
fibroblasts can compensate for the loss of *Acta2* transcription and form stress fibers using
similar proteins, implying that stress fiber formation is more important than αSMA production
for the fibrotic response[46]. Incorporating an expanded set of measurements in future fibrosis
studies may provide greater resolution of the fibrotic phenotype in response to therapies and
help evaluate changes in pathologically relevant features beyond protein expression.

Contributions of the LogiMML mechanistic machine learning approach

Mechanistic logic-based differential equation models have enabled systematic prediction of
drug action, yet these models are limited by the availability of priori knowledge [13,47–49]. An
alternative is machine learning, although 'black-box' ML approaches like artificial neural
networks predict input-output relationships without mechanistic insight. In contrast, two recent
studies combined mechanistic modeling with machine learning models like regression and
visible neural networks to predict antibiotic stress on metabolism and drug synergies for
cancer[29,50]. These 'white-box' approaches provide greater transparency of the intermediate
layers between input and output[51].

Building on such advances for logic-based biological networks, our LogiMML mechanistic
machine learning approach combines the flexible trainability of a machine learning model with
the robust experimentally-determined internal network structure of a mechanistic model. In
this study, we used the LogiMML model to predict signaling mechanisms that mediate how
drugs regulate F-actin assembly and stress fiber formation in cardiac fibroblasts. However, this
is just one of many possible applications for this modeling framework. The LogiMML approach
is designed to work across multiple mechanistic modeling formalisms and types of experimental
data, coupling the mechanistic model and data to predict mechanisms for the phenotype of
interest. The flexible nature of LogiMML presents promising future applications to elucidate cell
signaling that regulates diverse cellular phenotypes.

Src kinase as a therapeutic target for fibrosis

Of the 13 drugs used in this study, the Src inhibitor WH-4-023 (WH) was one of three drugs that
showed a strong reversal in TGFβ-induced F-actin, αSMA, and procollagen I expression. WH was
also effective at reversing the formation of F-actin stress fibers in response to TGFβ. Src
inhibitors dasatinib, ponatinib, and saracatinib have all been used in clinical trials across
different types of cancer[52–56]. In cancer, Src has been shown to promote proliferation and
metastasis through many signaling targets including FAK, Akt, Ras, and PI3K[57–61].

Given that Src signaling affects many central regulatory pathways, recent studies have tested
the potential for Src inhibition as a therapy for fibrotic disease. In a renal fibrosis study, blocking
Src kinase using PP1 was shown to inhibit TGFβ-induced expression of collagen I, αSMA, and
fibronectin[62]. In that study, Src inhibition was also shown the reduce the development of renal
fibrosis in obstructed kidneys *in vivo* in mice, indicating Src inhibition as a potential renal
fibrosis and chronic kidney disease therapy. Another study focusing on lung fibrosis showed

that TGFβ induces Src kinase activity in lung fibroblasts and that Src is required for
myofibroblast contraction[63]. Further, inhibition of Src kinase *in vivo* with AZD0530 reduced scar
area and αSMA expression in mice with bleomycin-induced lung fibrosis[63].

In this study, we applied the LogiMML network to investigate how Src contributes to F-actin
stress fiber formation induced by TGFβ. We predicted that PI3K signaling contributes to
profibrotic Src signaling in cardiac fibrosis. This proposed mechanism is supported by previous
studies, showing that PI3K regulates fibroblast contractility and myofibroblast activation in skin
fibroblasts, and TGFβ-induced αSMA and collagen production in lung fibroblasts[39,40]. To validate
this proposed profibrotic role for PI3K, we show that PI3K inhibition reduced procollagen I
production and F-actin stress fiber organization in HCFs. While previous work has shown that
mechanical stretch, Rho-kinase, and myosin light chain kinase (MLCK) positively regulate the
organization of F-actin filaments into stress fibers, the role of PI3K's regulation of F-actin stress
fiber formation has not been thoroughly explored[64,65]. Here, we show that treatment with PI3K
inhibitor LY294002 (LY) significantly reduces stress fiber formation without affecting the total
amount of assembled F-actin, implying PI3K specifically enhances stress fiber formation in
cardiac fibroblasts. Future studies should explore if Src kinase inhibitors mitigate cardiac fibrosis
in vivo, and to what degree PI3K kinase contributes to the regulation of cardiac fibrosis by Src.

Limitations and future directions
The main limitation of this study is that our modeling and experimental approaches address cell
signaling in cardiac fibroblasts *in vitro*, but do not address how fibroblasts respond to drugs in
an *in vivo* signaling environment. Our experimental data also captures some key fibrotic
proteins, but does not measure other fibrotic outputs of interest, like EDA fibronectin, and does
not capture a comprehensive signaling profile of the fibroblast. Despite these limitations, the
LogiMML framework was sufficient to predict a validated role for PI3K in promoting stress fiber
formation. Experimentally, future work could include proteomics or RNA-seq analysis of
fibroblasts to measure how drugs differentially regulate intracellular molecular profiles. Future
modeling work could include simulated conditions for *in vivo* or *in vitro* co-culture conditions to
incorporate the signaling influence of other cell types. Given the flexibility of the LogiMML
modeling approach, these simulated data could be feasibly paired with respective experimental
data to make predictions for fibroblast signaling under new conditions.

Conclusions
In this study, we showed that drugs differentially regulate cardiac fibroblast phenotype and
work via distinct mechanisms that can be predicted by logic-based mechanistic machine
learning. By expanding the microscopy feature set in the high content imaging pipeline, we
captured greater resolution of the fibroblast phenotype and measured how phenotypic
features changed in response to drugs. Using our LogiMML modeling approach, we predicted
signaling mechanisms for how pirfenidone and Src inhibitor WH-4-023 affect F-actin protein
assembly and stress fiber formation, respectively. We predicted that PI3K regulates F-actin
stress fiber formation, which we experimentally validated in human cardiac fibroblasts. This

study presents new features of fibroblast phenotype to be further explored in fibrosis,
identifies specific roles for PI3K in cardiac fibroblast signaling, and demonstrates an adaptable
mechanistic machine learning approach to predict signaling outcomes for fibrosis that can be
expanded to other diseases.

Methods

In vitro experiments in human cardiac fibroblasts

Primary human ventricular cardiac fibroblasts were purchased from PromoCell (PromoCell C-
12375; PromoCell GmbH, Germany). Cells were cultured in DMEM containing 10% FBS and 1%
Pen/Strep, and were kept in an incubator maintained at 5% CO_2. Cells were plated in a 96-well
plate at 5,000 cells/well and then grown in 10% FBS for 24 hours, serum starved for 24 hours,
and then treated with the following cytokine conditions for 96 hours: 0% FBS control media, 0%
FBS media with 20ng/mL TGFβ1 (Cell Signaling Technology, 8915LC), and 0% FBS media with 10
ng/mL human IL1β (Cell Signaling Technology, 8900SC), or TGFβ1 and IL1β combined. Cells were
treated with these conditions either alone or with 1 of 13 compounds at 1 of 3 concentrations.
We determined drug concentrations via a literature search, prioritizing concentrations that
yielded significant effects in vitro in fibroblasts or similar cell types. The drugs with their
respective concentrations are as follows: [0.25,1,2] µg/ml of anakinra (Kineret, SOBI Inc.),
[1,5,10] µM valsartan (Sigma-Aldrich, SML0142-10MG), [0.2,1,2] µM BNP (Sigma-Aldrich,
B5900-.5MG), [1,5,10]µM valsartan combos respectively with [0.2,1,2] µM BNP, [10,30,60]mM
glutathione (Sigma-Aldrich, G4251-1G), [1,3,5] µM CW-HM12 (Cayman Chemical Company,
19480), [10,20,50] µM salbutamol (Sigma-Aldrich, S8260-25MG), [5,10,25] µM marimistat
(Sigma-Aldrich, M2699-5Mg), [1,5,10] µM galunisertib (Selleck Chemicals, S2230), [12.5,25,50]
µM fasudil (Sigma-Aldrich, CDS021620-10MG), [10,25,50]µM SB203580 (Sigma-Aldrich, S8307-
1MG), [1,5,10] mg/mL pirfenidone (Sigma-Aldrich, P2116-10MG), [5,10,20] µM defactinib
(MedChem Express, HY-12289A), [5,10,20] µM WH-4-023 (Sigma-Aldrich, SML1334-5MG), and
20 µM LY294002 (Selleck Chemicals, S1105). Cells were grown in these conditions for 72 hours.

Cells were then fixed in 4% PFA in PBS for 30 minutes, permeabilized and blocked for 1 hour in
a solution containing 3% BSA and 0.2% Triton, and then stained overnight at 4°C with a 1:500
Anti-Collagen I antibody (Abcam, ab34710). After overnight incubation, cells were washed 3x in
PBS and stained with 1:5000 Dapi, 1:1000 Phalloidin CruzFluor 647 Conjugate (Santa Cruz
Biotechnology, sc-363797), 1:250 α-Smooth Muscle Actin antibody (Sigma-Aldrich, C6198), and
1:1000 Goat-anti-Rabbit (ThermoFisher Scientific, A-11034).

Microscopy and single-cell quantification

96-well plates we imaged using the Operetta CLS High-Content Analysis System (Perkin Elmer).
All three replicate wells for each condition were imaged and quantified. To quantify αSMA
expression, an automated image analysis pipeline was employed in CellProfiler (Broad
Institute)[22]. Fibroblast nuclei were identified by the DAPI signal. Next, the collagen-positive

region corresponding to each nucleus was segmented using the "propagate" algorithm, using
the segmented nucleus as the seed. Next, Fibroblast boundaries were segmented using the
"propagate" algorithm, musing the segmented collagen region as the seed. αSMA signal was
integrated within each cell's boundary. Short, medium, and long texture feature information
was derived using the MeasureTexture module in CellProflier using texture scales of 2, 6, and
10 pixels respectively. Texture feature values were calculated by subtracting the smallest angle
value of a given feature from the largest angle value of that same feature for each cell. F-actin
and procollagen expressions were quantified similarly.

Statistics

Feature values for each well were determined by taking the median value of the feature across
all cells in the center tile of each well. Well median values were used as replicates (n=3).
Significance was determined using an ANOVA with Tukey's posthoc in comparisons between
more than two groups, and Student's T-test in comparisons between two groups. Automated
data analysis and statistical calculations were performed using Python 3.8.5 and the
'statsmodels' Python module version 0.13.2.

Model Simulations

Drug simulations in the fibroblast network model were performed as previously described using
MATLAB version 2022a[12,13,66]. Predicted node activity is calculated using logic-based Hill
differential equations. Agonist and antagonist drug relationships were represented by altering
the activation function of the target node, representing either competitive or non-competitive
drug interactions with the respective target. To better represent the cell-to-cell variability
observed in *in vitro* cell responses to treatments, we employed a previously developed
ensemble modeling approach combining multiple simulations with random normally distributed
parameters[66]. Ensemble simulations were performed using the MATLAB 'normrnd' function
from the 'Statistics and Machine Learning' toolbox to randomly sample parameters within a
normal distribution and simulation n of 100. The randomly sampled parameters and means of
the sampling ranges are as follows: baseline ligand inputs (0.25), mechanical input (0.85), drug
dose (0.85), and raised ligand inputs (0.6). The sampling range for each parameter was
calculated by $paramMean \pm COV * paramMean$ where COV=0.0396. This COV value, used to
scale stochasticity in the model was determined by taking the average coefficient of variation in
F-actin, procollagen I, and αSMA expression in human cardiac fibroblasts treated with TGFβ
from our *in vitro* experiments.

LogiMML Network-Regression Coupling

The LogiMML mechanistic machine learning model is comprised of a network model layer and a
Ridge regression layer. The independent 'X' variables used to train the regression model are
node activity values from the network model predicted under each simulated drug and
environmental condition. To reduce model complexity, network nodes were clustered into 11
signaling modules derived from k-means clustering on a combined sensitivity and influence
analysis on the network model[12]. The node activity values were averaged within each module,

and these modules' mean activity values were fed into the regression layer. The dependent 'Y'
variables for this model were experimentally measured values from our high-content imaging
experiments in human cardiac fibroblasts. Sensitivity knockout analysis was performed by
simulating a given drug and cytokine context int network model i.e. 'TGFβ+pirfenidone' and
sequentially setting each node ymax value to 0, measuring reduction or increase in the
dependent variable e.g. 'F-actin Intensity' upon knockdown. Leave-one-out cross validation
(LOOCV) was performed on the LogiMML model to evaluate performance across variations in
the experimental data set. The means and standard deviations of the LOOCV MSE values were
0.022 and 0.080 for the F-actin Intensity model and 0.083 and 0.142 for the Long Actin ASM
model.

Acknowledgements

478 We acknowledge Dr. Sarah Ewald at the University of Virginia Carter Immunology Center for
479 providing anakinra used in this study.

Funding

482 This study was funded by the NIH (HL137755, HL007284, HL160665, HL162925, 1S10OD021723-
483 01A1).

References

1. Burgos Villar, K. N., Liu, X. & Small, E. M. Transcriptional regulation of cardiac fibroblast
phenotypic plasticity. *Curr. Opin. Physiol.* **28**, 100556 (2022).

2. Jacobs, A. K. *et al.* Systems of Care for ST-Segment-Elevation Myocardial Infarction: A
Policy Statement From the American Heart Association. *Circulation* **144**, e310–e327
(2021).

3. Burke, R. M., Lighthouse, J. K., Mickelsen, D. M. & Small, E. M. Sacubitril/Valsartan
Decreases Cardiac Fibrosis in Left Ventricle Pressure Overload by Restoring PKG Signaling
in Cardiac Fibroblasts. *Circ. Hear. Fail.* **12**, (2019).

4. Fan, Z. & Guan, J. Antifibrotic therapies to control cardiac fibrosis. *Biomater. Res. 2016*
201 **20**, 1–13 (2016).

5. Welch, M. P., Odland, G. F. & Clark, R. A. F. Temporal relationships of F-actin bundle
formation, collagen and fibronectin matrix assembly, and fibronectin receptor expression
to wound contraction. *J. Cell Biol.* **110**, 133–145 (1990).

6. Travers, J. G., Kamal, F. A., Robbins, J., Yutzey, K. E. & Blaxall, B. C. Cardiac Fibrosis: The
Fibroblast Awakens. *Circ. Res.* **118**, 1021–40 (2016).

7.   Sun, K. H., Chang, Y., Reed, N. I. & Sheppard, D. α-smooth muscle actin is an inconsistent
marker of fibroblasts responsible for force-dependent TGFβ activation or collagen
production across multiple models of organ fibrosis. *Am. J. Physiol. - Lung Cell. Mol.*
Physiol. **310**, L824–L836 (2016).

8.   Avery, D. *et al.* Extracellular Matrix Directs Phenotypic Heterogeneity of Activated
Fibroblasts. *Matrix Biol.* **67**, 90 (2018).

9.   Deng, C. C. *et al.* Single-cell RNA-seq reveals fibroblast heterogeneity and increased
mesenchymal fibroblasts in human fibrotic skin diseases. *Nat. Commun. 2021 121* **12**, 1–
16 (2021).

10. Gerckens, M. *et al.* Phenotypic drug screening in a human fibrosis model identified a
novel class of antifibrotic therapeutics. *Sci. Adv.* **7**, 3673 (2021).

11. Fu, X. *et al.* Specialized fibroblast differentiated states underlie scar formation in the
infarcted mouse heart. *J. Clin. Invest.* **128**, 2127–2143 (2018).

12. Zeigler, A. C., Richardson, W. J., Holmes, J. W. & Saucerman, J. J. A computational model
of cardiac fibroblast signaling predicts context-dependent drivers of myofibroblast
differentiation. *J. Mol. Cell. Cardiol.* **94**, 72–81 (2016).

13. Zeigler, A. C. *et al.* Network model-based screen for FDA-approved drugs affecting
cardiac fibrosis. *CPT Pharmacometrics Syst. Pharmacol.* **10**, 377–388 (2021).

14. Kort, E. & Jovinge, S. Drug Repurposing: Claiming the Full Benefit from Drug
Development. *Curr. Cardiol. Rep.* **23**, 1–7 (2021).

15. Abdelsayed, M., Kort, E. J., Jovinge, S. & Mercola, M. Repurposing drugs to treat
cardiovascular disease in the era of precision medicine. doi:10.1038/s41569-022-00717-6

16. Khalil, H. *et al.* Fibroblast-specific TGF-β-Smad2/3 signaling underlies cardiac fibrosis. *J.*
Clin. Invest. **127**, 3770–3783 (2017).

17. Turner, N. A. *et al.* Interleukin-1α stimulates proinflammatory cytokine expression in
human cardiac myofibroblasts. *Am. J. Physiol. Circ. Physiol.* **297**, H1117–H1127 (2009).

18. Francis Stuart, S. D., De Jesus, N. M., Lindsey, M. L. & Ripplinger, C. M. The crossroads of
inflammation, fibrosis, and arrhythmia following myocardial infarction. *J. Mol. Cell.*
Cardiol. **91**, 114–22 (2016).

19. Deten, A., Volz, H. C., Briest, W. & Zimmer, H.-G. Cardiac cytokine expression is
upregulated in the acute phase after myocardial infarction. Experimental studies in rats.
Cardiovasc. Res. **55**, 329–340 (2002).

20. Bujak, M., Frangogiannis, N. G. & Hirszfeld, © L. The role of IL-1 in the pathogenesis of
heart disease THE IL–1 FAMILY OF CYTOKINES. doi:10.1007/s00005-009-0024-y

21. BUJAK, M. & FRANGOGIANNIS, N. The role of TGF-β signaling in myocardial infarction and
cardiac remodeling. *Cardiovasc. Res.* **74**, 184–195 (2007).

22.   Stirling, D. R. *et al.* CellProfiler 4: improvements in speed, utility and usability. *BMC*
Bioinformatics **22**, 1–11 (2021).

23.   Abbate, A. *et al.* Effects of interleukin-1 blockade with anakinra on adverse cardiac
remodeling and heart failure after acute myocardial infarction [from the virginia
commonwealth university-anakinra remodeling trial (2) (vcu-art2) pilot study]. *Am. J.*
Cardiol. **111**, (2013).

24.   Abbate, A. *et al.* Anakinra, a recombinant human interleukin-1 receptor antagonist,
inhibits apoptosis in experimental acute myocardial infarction. *Circulation* **117**, (2008).

25.   Abbate, A. *et al.* Interleukin-1 blockade inhibits the acute inflammatory response in
patients with st-segment–elevation myocardial infarction. *J. Am. Heart Assoc.* **9**, (2020).

26.   Bracco Gartner, T. C. L. *et al.* Pirfenidone Has Anti-fibrotic Effects in a Tissue-Engineered
Model of Human Cardiac Fibrosis. *Front. Cardiovasc. Med.* **9**, 459 (2022).

27.   Behr, J. *et al.* Pirfenidone in patients with progressive fibrotic interstitial lung diseases
other than idiopathic pulmonary fibrosis (RELIEF): a double-blind, randomised, placebo-
controlled, phase 2b trial. *Lancet Respir. Med.* **9**, 476–486 (2021).

28.   Sandbo, N. & Dulin, N. The actin cytoskeleton in myofibroblast differentiation:
Ultrastructure defining form and driving function. *Transl. Res.* **158**, 181 (2011).

29.   Yang, J. H. *et al.* A White-Box Machine Learning Approach for Revealing Antibiotic
Mechanisms of Action. *Cell* **177**, 1649-1661.e9 (2019).

30.   Rieg, T., Frick, J., Baumgartl, H. & Buettner, R. Demonstration of the potential of white-
box machine learning approaches to gain insights from cardiovascular disease
electrocardiograms. *PLoS One* **15**, e0243615 (2020).

31.   McNeish, D. M. Using Lasso for Predictor Selection and to Assuage Overfitting: A Method
Long Overlooked in Behavioral Sciences. *Multivariate Behav. Res.* **50**, 471–484 (2015).

32.   Hu, T. *et al.* Reactive oxygen species production via NADPH oxidase mediates TGF-β-
induced cytoskeletal alterations in endothelial cells. *Am. J. Physiol. - Ren. Physiol.* **289**,
816–825 (2005).

33.   Huot, J. *et al.* SAPK2/p38-dependent F-Actin Reorganization Regulates Early Membrane
Blebbing during Stress-induced Apoptosis. *J. Cell Biol.* **143**, 1361–1373 (1998).

34.   Qian, Y. *et al.* PI3K induced actin filament remodeling through Akt and p70S6K1:
implication of essential role in cell migration. *Am. J. Physiol. Cell Physiol.* **286**, (2004).

35.   Haralick, R. M., Dinstein, I. & Shanmugam, K. Textural Features for Image Classification.
IEEE Trans. Syst. Man Cybern. **SMC**-**3**, 610–621 (1973).

36.   Nagaraju, C. K. *et al.* Myofibroblast Phenotype and Reversibility of Fibrosis in Patients
With End-Stage Heart Failure. *J. Am. Coll. Cardiol.* **73**, 2267–2282 (2019).

37.   Sandbo, N. *et al.* Delayed stress fiber formation mediates pulmonary myofibroblast

differentiation in response to TGF-β. *Am. J. Physiol. - Lung Cell. Mol. Physiol.* **301**, 656–666 (2011).

38. Naumanen, P., Lappalainen, P. & Hotulainen, P. Mechanisms of actin stress fibre assembly. *J. Microsc.* **231**, 446–454 (2008).

39. Conte, E. *et al.* Inhibition of PI3K prevents the proliferation and differentiation of human lung fibroblasts into myofibroblasts: The role of class I P110 isoforms. *PLoS One* **6**, (2011).

40. Li, G., Li, Y. Y., Sun, J. E., Lin, W. H. & Zhou, R. X. ILK-PI3K/AKT pathway participates in cutaneous wound contraction by regulating fibroblast migration and differentiation to myofibroblast. *Lab. Investig.* **96**, 741–751 (2016).

41. Luangmonkong, T. *et al.* Evaluating the antifibrotic potency of galunisertib in a human ex vivo model of liver fibrosis. *Br. J. Pharmacol.* **174**, (2017).

42. Masuda, A. *et al.* Promotion of liver regeneration and anti-fibrotic effects of the TGF-β receptor kinase inhibitor galunisertib in CCl4-treated mice. *Int. J. Mol. Med.* **46**, (2020).

43. Dadrich, M. *et al.* Combined inhibition of TGFβ and PDGF signaling attenuates radiation-induced pulmonary fibrosis. *Oncoimmunology* **5**, (2016).

44. Bhattacharyya, S. *et al.* TLR4-dependent fibroblast activation drives persistent organ fibrosis in skin and lung. *JCI insight* **3**, (2018).

45. Chakraborty, D. *et al.* Activation of STAT3 integrates common profibrotic pathways to promote fibroblast activation and tissue fibrosis. *Nat. Commun. 2017 81* **8**, 1–16 (2017).

46. Li, Y. *et al.* Loss of Acta2 in cardiac fibroblasts does not prevent the myofibroblast differentiation or affect the cardiac repair after myocardial infarction. *J. Mol. Cell. Cardiol.* **171**, 117–132 (2022).

47. Tan, P. M., Buchholz, K. S., Omens, J. H., McCulloch, A. D. & Saucerman, J. J. Predictive model identifies key network regulators of cardiomyocyte mechano-signaling. *PLoS Comput. Biol.* **13**, (2017).

48. Ryall, K. A. *et al.* Network reconstruction and systems analysis of cardiac myocyte hypertrophy signaling. *J. Biol. Chem.* **287**, 42259–42268 (2012).

49. Kraeutler, M. J., Soltis, A. R. & Saucerman, J. J. Modeling cardiac β-adrenergic signaling with normalized-Hill differential equations: comparison with a biochemical model. *BMC Syst. Biol.* **4**, 157 (2010).

50. Kuenzi, B. M. *et al.* Predicting Drug Response and Synergy Using a Deep Learning Model of Human Cancer Cells. *Cancer Cell* **38**, 672-684.e6 (2020).

51. Petch, J., Di, S. & Nelson, W. Opening the Black Box: The Promise and Limitations of Explainable Machine Learning in Cardiology. *Can. J. Cardiol.* **38**, 204–213 (2022).

52. Belli, S. *et al.* C-Src and EGFR inhibition in molecular cancer therapy: What else can we improve? *Cancers* **12**, (2020).

53. Gucalp, A. *et al.* Phase II trial of saracatinib (AZD0530), an oral SRC-inhibitor for the
treatment of patients with hormone receptor-negative metastatic breast cancer. *Clin.*
Breast Cancer **11**, (2011).

54. Cortes, J. *et al.* Ponatinib dose-ranging study in chronic-phase chronic myeloid leukemia:
a randomized, open-label phase 2 clinical trial. *Blood* **138**, (2021).

55. Haura, E. B. *et al.* Phase I/II study of the Src inhibitor dasatinib in combination with
erlotinib in advanced non-small-cell lung cancer. *J. Clin. Oncol.* **28**, (2010).

56. Twardowski, P. W. *et al.* A phase II trial of dasatinib in patients with metastatic
castration-resistant prostate cancer treated previously with chemotherapy. *Anticancer.*
Drugs **24**, (2013).

57. Beadnell, T. C. *et al.* Src-mediated regulation of the PI3K pathway in advanced papillary
and anaplastic thyroid cancer. *Oncogenesis* **7**, (2018).

58. Bunda, S. *et al.* Src promotes GTPase activity of Ras via tyrosine 32 phosphorylation.
Proc. Natl. Acad. Sci. U. S. A. **111**, (2014).

59. Xiao, X. *et al.* Diallyl disulfide suppresses SRC/Ras/ERK signaling-mediated proliferation
and metastasis in human breast cancer by up-regulating miR-34a. *PLoS One* **9**, (2014).

60. Westhoff, M. A., Serrels, B., Fincham, V. J., Frame, M. C. & Carragher, N. O. Src-Mediated
Phosphorylation of Focal Adhesion Kinase Couples Actin and Adhesion Dynamics to
Survival Signaling. *Mol. Cell. Biol.* **24**, (2004).

61. Mahajan, K. & Mahajan, N. P. PI3K-independent AKT activation in cancers: A treasure
trove for novel therapeutics. *Journal of Cellular Physiology* **227**, (2012).

62. Yan, Y. *et al.* Src inhibition blocks renal interstitial fibroblast activation and ameliorates
renal fibrosis. *Kidney Int.* **89**, (2016).

63. Hu, M. *et al.* Therapeutic Targeting of Src Kinase in Myofibroblast Differentiation and
Pulmonary Fibrosis. *J. Pharmacol. Exp. Ther.* **351**, 87 (2014).

64. Katoh, K., Kano, Y., Amano, M., Kaibuchi, K. & Fujiwara, K. Stress fiber organization
regulated by MLCK and Rho-kinase in cultured human fibroblasts. *Am. J. Physiol. - Cell*
Physiol. **280**, (2001).

65. Kaunas, R., Nguyen, P., Usami, S. & Chien, S. Cooperative effects of Rho and mechanical
stretch on stress fiber organization. *Proc. Natl. Acad. Sci. U. S. A.* **102**, (2005).

66. Zeigler, A. C. *et al.* Computational Model Predicts Paracrine and Intracellular Drivers of
Fibroblast Phenotype After Myocardial Infarction. *Matrix Biol.* (2020).
doi:10.1016/j.matbio.2020.03.007