Logic-based mechanistic machine learning on high-content images reveals how drugs differentially regulate cardiac fibroblasts

Anders R. Nelson[1], Steven L. Christiansen[1,2], Kristen M. Naegle[1], Jeffrey J. Saucerman[1*]

1) University of Virginia School of Medicine, Charlottesville, VA 22903

2) Brigham Young University Department of Biochemistry, Provo, UT 84602

*University of Virginia School of Medicine, Charlottesville, VA 2290

(e-mail: jsaucerman@virginia.edu)

PO Box 800759, Charlottesville, VA 22908, 434-924-5095

Classification: Biological Sciences; Systems Biology

Keywords: Fibrosis, Fibroblast, Machine learning, Cell Biology, Differentiation, Drugs

Abstract

Fibroblasts are essential regulators of extracellular matrix deposition following cardiac injury. These cells exhibit highly plastic responses in phenotype during fibrosis in response to environmental stimuli. Here, we test whether and how candidate anti-fibrotic drugs differentially regulate measures of cardiac fibroblast phenotype, which may help identify treatments for cardiac fibrosis. We conducted a high content microscopy screen of human cardiac fibroblasts treated with 13 clinically relevant drugs in the context of TGFβ and/or IL-1β, measuring phenotype across 137 single-cell features. We used the phenotypic data from our high content imaging to train a logic-based mechanistic machine learning model (LogiMML) for fibroblast signaling. The model predicted how pirfenidone and Src inhibitor WH-4-023 reduce actin filament assembly and actin-myosin stress fiber formation, respectively. Validating the LogiMML model prediction that PI3K partially mediates the effects of Src inhibition, we found that PI3K inhibition reduces actin-myosin stress fiber formation and procollagen I production in human cardiac fibroblasts. In this study, we establish a modeling approach combining the strengths of logic-based network models and regularized regression models, apply this approach to predict mechanisms that mediate the differential effects of drugs on fibroblasts, revealing Src inhibition acting via PI3K as a potential therapy for cardiac fibrosis.

Significance

Cardiac fibrosis is a dysregulation of the normal wound healing response, resulting in excessive scarring and cardiac dysfunction. As cardiac fibroblasts primarily regulate this process, we explored how candidate anti-fibrotic drugs alter the fibroblast phenotype. We identify a set of 137 phenotypic features that change in response to drug treatments. Using a new computational modeling approach termed logic-based mechanistic machine learning, we predict how pirfenidone and Src inhibition affect the regulation of the phenotypic features actin filament assembly and actin-myosin stress fiber formation. We also show that inhibition of PI3K reduces actin-myosin stress fiber formation and procollagen I production in human cardiac fibroblasts, supporting a role for PI3K as a mechanism by which Src inhibition may suppress fibrosis.

Introduction

Cardiac fibroblasts are the primary regulators of remodeling following cardiac injury[1]. Extracellular matrix (ECM) deposition by activated myofibroblasts is essential to this response, but excessive deposition can lead to ventricular stiffness, diastolic dysfunction, and heart failure[1]. While fibroblasts are critical to the wound healing response, current standard-of-care therapeutics for cardiac injury, such as myocardial infarction (MI), affect downstream symptoms but do not specifically target fibroblast signaling[2]. Recent drug discovery and development has focused on identifying drugs such as Entresto (sacubitril/valsartan) that reduce fibrosis in part by modulating fibroblast signaling[3,4].

Collagen secretion, αSMA expression, and actin filaments (F-actin) are traditional markers for a
profibrotic fibroblast phenotype[5,6]. While high expression of these markers provides an initial
indication of myofibroblast activation, traditional marker expression is inconsistent and does
not fully capture the fibrotic response[7]. Recent studies of fibroblast phenotype have shown that
fibroblasts exhibit high phenotypic heterogeneity across many facets in response to injury, and
that phenotypic changes are also sensitive to drug perturbations[8–11]. Identifying drugs that
regulate fibroblast signaling may provide targeted control of fibrosis.

Previously, we developed a logic-based mechanistic network model of fibroblast signaling and
applied it to perform virtual screens for anti-fibrotic drugs[12,13]. That study predicted and
experimentally validated an antifibrotic role for the TGFβ receptor inhibitor galunisertib[13].
While the fibroblast network model predicts a number of drugs that modulate fibroblast
activation, substantial experimental characterization is needed to capture phenotypic
responses to drugs that were not captured by prior modeling.

In this study, we combined high content microscopy, network modeling, and machine learning
to identify drugs that differentially regulate fibroblast phenotypic metrics and predict their
underlying network mechanisms. We used image-based feature extraction to more deeply
characterize drug response and fibroblast phenotype, capturing drug-induced changes across a
set of single-cell metrics relevant to fibrosis. Using a novel logic-based mechanistic machine
learning approach, LogiMML, we predicted signaling pathways that determine how drugs
regulate fibroblast phenotype. Finally, we experimentally validated the main pathway
mechanism predicted by the LogiMML model that mediates how Src inhibition suppresses
fibrotic responses.

Results

An *in vitro* screen for candidate fibrosis drugs

Previously, we applied our published cardiac fibroblast network model[12] to identify candidate
therapies predicted to reduce cardiac fibrosis[13]. This logic-based differential equation network
model was developed from a wide range of fibroblast signaling relationships from *in vitro*
studies in the literature. The model predicts changes in fibrotic outputs including collagen I and
III, αSMA, EDA fibronectin, matrix metalloproteases, and F-actin in response to changes in
extracellular signaling contexts and drug treatment[12]. This model was previously integrated with
the drug-target database DrugBank to make predict the response of fibroblasts to 121 FDA-
approved or investigational drugs that have targets in this network[13].

To expand upon the *in silico* modeling work done in this previous study[13], we aimed to develop
a list of drug candidates to test experimentally for their ability to reduce fibrosis in cardiac
fibroblasts *in vitro*. As the model predicted many drugs to reduce fibrosis to similar quantitative
degrees[13], we included drug selection criteria outside of our modeling results alone to further

narrow-down a list of candidate drugs. First, we prioritized pathway diversity of the drug
targets to ensure that we would perturb fibrotic signaling comprehensively and avoid testing
redundant drugs in our experiments. As drug repurposing has become an increasingly effective
and efficient strategy for treating cardiovascular disease, we next looked to prioritize drugs that
had previous clinical indications for other disease areas[14,15]. Using these selection criteria, we
developed the following list of thirteen drugs to evaluate experimentally: anakinra, valsartan,
defactinib, HW-4-023, glutathione, CW-HM12, salbutamol, marimistat, fasudil, SB203580,
pirfenidone, brain natriuretic peptide (BNP), and a combination of valsartan and BNP (Table S1).
Among the list of candidate drug targets are regulators for inflammatory signaling, mechanical
stretch response, non-canonical TGFβ signaling, and modification of secreted proteins.

We next aimed to test these candidate drugs for their ability to quantitatively reduce fibrosis as
characterized by image-based single-cell profiling of procollagen I, α-smooth muscle actin
(αSMA), and F-actin. In injury signaling conditions, such as following myocardial infarction (MI),
myocardial cells are exposed to elevated proinflammatory[16–18]. To represent these signaling
contexts in an *in vitro* system, we included IL1β and TGFβ, shown to be elevated following
cardiac injury, in our treatment conditions to represent proinflammatory and profibrotic
contexts respectively[19–21]. We tested our candidate drugs under four total cytokine contexts
(baseline context with no added cytokine, fibrotic context represent by TGFβ, inflammatory
context represented by IL1β, and combined context represent by both TGFβ and IL1β)[19–21]. In
total, we used 108 treatment conditions consisting of one of the thirteen drugs at a low,
medium, or high dose combined with one of the four cytokine contexts. We also included
treatments of each cytokine context with no drug to establish a control baseline for cell
responses to cytokines. We imaged and quantified single-cell protein expression of three
fibrotic markers, procollagen I, α-smooth muscle actin (αSMA), and F-actin using high-content
microscopy and a custom CellProfiler software pipeline[22].

Interestingly, the antifibrotic drugs in our screen induced differential effects on fibrosis. Of the
13 candidate drugs, WH-4-023, fasudil, and defactinib caused the strongest reduction of
procollagen I, F-actin, and αSMA expression in a TGFβ signaling context, even at the lowest
dose (Figure 1A). Conversely, a second set of drugs including anakinra and glutathione
increased fibrotic marker expression in both TGFβ and combined TGFβ/IL1β contexts when
applied directly to fibroblasts. In a previous clinical study, anakinra, an IL1 receptor inhibitor,
was shown to improve cardiac function and prevented heart failure following acute MI[23]. While
anakinra has been shown to reduce infarct scar area in a mouse MI model, it also exhibits other
beneficial cardiac effects post-MI including inhibition of post-MI myocyte apoptosis and
reduction in systemic inflammat[24,25]. Based on these previous studies, it is likely that anakinra
has a net antifibrotic effect on fibroblasts in the presence of other myocardial cell types even
though anakinra treatment increased fibrotic marker expression in this experiment. A third set
of drugs showed more selective antifibrotic effects. For example, while fasudil significantly
reduced expression of all three fibrosis markers in a TGFβ signaling context, pirfenidone only
significantly reduced F-actin (Figure 1 B-E). This third set of drugs is of particular interest as it

contains drugs that differentially regulate markers for fibrosis. Given the recent clinical
effectiveness of pirfenidone for lung fibrosis, and success in diseases models for cardiac
fibrosis[26,27], we further investigated the mechanisms by which it regulates F-actin in cardiac
fibroblasts.

LogiMML: logic-based mechanistic machine learning model predicts how drugs
regulate fibroblast phenotype

Assembled actin filaments play a key role in contractility as fibroblasts transition to become
myofibroblasts[28]. Therefore, we asked whether the previous mechanistic computational model
of the fibroblast signaling network[12] could predict our experimentally measured inhibition of
filament assembly by pirfenidone from Figure 1D. While the model had correctly predicted
responses to a number of drugs including galunisertib[13], here, the original mechanistic model
did not capture the ability of pirfenidone to suppress actin filament assembly in a TGFβ
signaling context (Figure 2 A).

Given the limitations of a model based only on prior knowledge, we asked whether drug
predictions could be improved by combining the mechanistic model with a machine learning
model that leverages data from the drug screen. Motivated by 'white-box' machine learning
strategies that combine mechanistic models with machine learning[29,30], we designed a logic-
based mechanistic machine learning (LogiMML) model to predict key regulators that conduct
signaling from network model inputs and simulated drugs to experimentally measured
phenotypic outputs (Figure 2 B, Figure S 1). As the 108 treatments were insufficient to infer
new links to phenotypic outputs from all 91 model nodes, we reduced the model's
dimensionality by clustering nodes into modules. Eleven signaling modules were computed
based on a combined influence and sensitivity analysis, grouping nodes with similar predicted
behavior across signaling contexts. The machine learning component was then trained by
mapping the model-predicted activity of each network module for each of the 108
drug+cytokine treatments to respective experimentally measured outputs. Regularized ridge
regression was selected for the machine learning layer of the LogiMML model to reduce the
likelihood of overfitting[31]. As measured experimentally, the LogiMML model correctly predicted
the respective induction and suppression of F-actin by TGFβ and pirfenidone (Figure 2 C). Leave-
one-out cross validation (LOOCV) was performed on the LogiMML model to evaluate
performance across variations in the experimental data set. The means and standard deviation
of the LOOCV MSE values were 0.022 and 0.080 for the F-actin Integrated Intensity model.

We next asked whether the LogiMML model could provide new mechanistic insights into how
F-actin is regulated by pirfenidone. First, we used the LogiMML model's ridge regression
coefficients to predict the modules that most influence F-actin. We used the β coefficients from
the LogiMML model to predict the influence of a given signaling module on the cell feature of
interest. 'PI3K' and 'Smad3' modules were predicted to be the top positive regulators of F-
actin, while the 'P38_Calcium' module was predicted as the top negative regulator (Figure 2 D).
These predictions for fibroblasts are consistent with previous studies with other cell types

showing that members of the 'Smad3' and 'P38_Calcium' signaling modules regulate F-actin
filament assembly in endothelial cells and that members of the 'PI3K' signaling module promote
actin filament remodeling during migration in embryonic fibroblasts [32–34]. To identify which
individual signaling nodes within these three modules most regulate F-actin, we performed a
virtual knockdown screen of the mechanistic network model for regulators of F-actin in a
'TGFβ+pirfenidone' signaling context (Figure 2 E). In these analyses, the quantitative outputs of
the model are normalized outputs that can be compared to determine predicted increases or
decreases in a cell feature in response to a perturbation. Summarizing these analyses, the
LogiMML model predicts that pirfenidone regulation of F-actin is positively regulated by p38,
Akt, and CBP, while negatively regulated by ROS and NOX (Figure 2 F).

Drugs and pathways controlling fibroblast morphology and texture

Given the differential regulation of fibrosis marker protein expression, we asked whether other
aspects of fibroblast phenotype may also be differentially regulated by drugs and cytokines.
Qualitatively, we observed morphological changes in cell shape, actin-myosin stress fiber
formation, intracellular protein distribution, and cell area (e.g. for pirfenidone treatment see
Figure 1 E). To measure these characteristics of fibroblast phenotype, we developed a custom
CellProfiler image analysis pipeline quantifying 137 total single-cell cell features[22,35]. Integrated
intensities for the three fibrotic marker proteins, procollagen I, F-actin, and αSMA clustered
relatively close to each other across the feature space (Figure 3 A). As expected, expression of
these marker proteins and similar features were high under TGFβ and TGFβ-like treatments,
and low under negative control and IL1β conditions. While the central rows of the heatmap
contain many features with similar treatment responses, the features at the top and bottom
regions of the heatmap show high heterogeneity in response to drugs. The significance of the
overall correlation between actin, aSMA, and collagen expression is two-fold: that some drugs
such as fasudil suppress a well-studied canonical myofibroblast activation program, and that the
responses to other drugs revealed that the overall phenotype space of fibroblasts is much more
diverse and can be specifically targeted with drugs like WH-4-023. Even within the
actin/aSMA/collagen cluster, the hierarchical clustering shows some examples where only in the
context of TGFβ treatment, some drugs up-regulated aSMA/collagen features while down-
regulating actin features.

The extraction of fibrotic marker proteins and the large degree of information about those
fibrotic features is a rich dataset with which we next wished to understand more directly how
they relate to each other and to treatment. Given the risk that some of our features carry
redundant information, we calculated the correlation between all features and clustered the
correlation matrix. This identified 15 strong sets of feature clusters. We selected one
representative feature of each cluster (Figure S2, Figure S3) based on choosing the cluster
member that demonstrated high variance across samples and low correlation with features from
other clusters. In addition to the 15 representative features, we retained the three integrated
intensity features for procollagen I, F-actin, and αSMA, yielding a final set of 18 distinct features

of the original 137 features, which have the potential to represent the complexity of the larger
dataset.

In order to interpret the overall underlying relationships in the 18 selected features and how
they relate to treatments, we performed Principal Component Analysis (PCA) (Figure 3B-C).
Negative control treatments had a negative score on the first principal component (PC1), while
cells treated with TGFβ showed a high positive score on PC1, indicating that the first principal
component correlates with an axis of classical fibroblast activation (Figure 4 B, Figure S4 A). This
was further supported by the PCA loading values for integrated procollagen I, F-actin, and
αSMA (Figure 3 C, Figure S4 B). These three features are expected to be relatively high in
activated myofibroblasts and indeed have strong positive loadings on PC1. On the PCA scores,
many of the 'TGFβ + Drug' groups deviated from the control-TGFβ axis defined on PC1, implying
that drugs induce phenotypic changes distinct from a simple reversal of TGFβ's effects. To
further investigate drug-induced changes in phenotype, we analyzed the PCA scores and
loadings to infer links between drugs and the features they regulate. Notably, the Src inhibitor
WH-4-023 (WH) showed directionality on the scores plot similar to that of Actin Long Angular
Second Moment (Actin Long ASM, a measure for actin uniformity) on the loadings plot. Actin-
myosin stress fibers, composed of multiple actin filaments along with other proteins, contribute
to pathological fibrosis and myofibroblast differentiation[36–38]. This feature and treatment pair
showed a negative value on PC1 and a positive value on PC2 relative to the TGFβ and control
groups, respectively. The similar directionality of WH and Actin Long ASM suggests that Src
inhibition may modulate actin uniformity.

Based on the initial inference from the PCA, we revisited the images from the high-content
microscopy experiment. Fibroblasts treated with TGFβ exhibited discrete actin-myosin stress
fibers, and stress fibers were qualitatively reduced when WH-4-023 (WH) was added (Figure 4
A). Quantitative analysis of actin uniformity (inversely correlated with stress fibers) using Actin
Long Angular Second Moment (ASM) further supported that TGFβ increased and Src inhibitor
WH reduced actin uniformity (Figure 4 B). The full dose response for Long Actin ASM to WH-4-
023 is shown in Figure S5.

To predict the signaling pathways that specifically regulate actin-myosin stress fibers, we again
applied the LogiMML coupled modeling approach, but this time training the ridge regression
layer of the model on experimental measurements of Actin Long ASM. The means and standard
deviation of the LOOCV MSE values were 0.083 and 0.142 for the Long Actin ASM model. The
LogiMML model regression coefficients predicted that the 'Mechanical' module was the top
positive regulator of Actin Long ASM and that the 'PI3K' module was the top negative regulator
of Actin Long ASM (Figure 4 C). To identify which individual signaling nodes within these two
modules most regulate Actin Long ASM, we performed a virtual knockdown screen of the
mechanistic network model for regulators of Actin Long ASM in the context of 'TGFβ+WH-4-
023' and predicted that Rho, MKK4, and Akt are proximal regulators of Actin Long ASM and
actin-myosin stress fiber formation (Figure 4 D-E).

PI3K signaling stimulates actin-myosin stress fiber formation and collagen expression

After deriving a putative signaling schematic for Actin Long ASM using the LogiMML model, we aimed to experimentally validate the prediction that inhibition of PI3K/Akt would suppress stress fiber formation and thereby increase Actin Long ASM (Figure 4 E). In previous studies using PI3K inhibitors, PI3K was shown to regulate fibroblast contractility, fibroblast-to-myofibroblast transition, and TGFβ-induced αSMA and collagen production[39,40]. Given these previously implicated roles for PI3K in myofibroblast activation and fibrosis, we wanted to investigate if PI3K has a regulatory role for actin-myosin stress fiber formation in cardiac fibroblasts. We treated human cardiac fibroblasts with either a negative control condition or a 20 μM dose of the PI3K inhibitor LY294002 (LY). Treatment with LY significantly increased Actin Long ASM, but notably, it had no significant effect on the total assembly of actin filaments in each cell, measured by integrated F-actin intensity (Figure 4 F-G). This selective effect of PI3K inhibition on stress fiber formation, while having no significant effect on total F-actin, suggests that actin filament assembly and stress fiber formation are differentially regulated processes. PI3K inhibition also significantly reduced integrated procollagen I intensity, demonstrating a role for PI3K signaling in cardiac fibroblast collagen production (Figure 4G).

Discussion

Cardiac fibroblasts are central regulators and promising therapeutic targets following cardiac injury. To identify how clinically relevant drugs regulate diverse aspects of fibroblast phenotype, we performed high-content screening of 13 drugs in 4 environmental contexts. We expanded our high-content microscopy feature set to 137 single-cell features, measuring fibrotic marker protein intensity, intracellular protein distribution, fiber texture, and cell morphology. After reducing the feature space and dimensionality of our experimental data, we found that many aspects of fibroblast phenotype are uniquely induced by drug and cytokine treatments. Notably, when administered with TGFβ, the drugs WH-4-023, defactinib, fasudil, and pirfenidone induced phenotypes that deviated from the PCA axis corresponding to classical TGFβ response. The differences between these phenotypes can be partially explained by differential drug regulation of features capturing Procollagen I and αSMA expression, and actin filament assembly and actin-myosin stress fiber formation. To predict how drugs regulate cell signaling and influence phenotype, we developed the logic-based mechanistic machine learning (LogiMML) approach which coupled the logic-based fibroblast network model with a ridge regression model trained on the high-content drug screen. Using this expanded LogiMML model, we predicted regulatory mechanisms for pirfenidone and Src inhibitor WH-4-023 on actin filaments. We predicted that pirfenidone regulates actin filament assembly via the 'P38_Calcium', 'Smad3', and 'PI3K' signaling modules, with Akt, p38, and CBP predicted to be positive drivers of actin filament assembly within these modules. We also predicted that WH-4-023 regulates actin-myosin stress fiber formation via the 'PI3k' and 'Mechanical' signaling

300   modules. As predicted by the LogiMML model, we experimentally validated that PI3K inhibition
301   reduces actin-myosin stress fiber formation in human cardiac fibroblasts. These studies validate
302   the ability of the LogiMML approach to predict signaling mechanisms from a phenotypic screen.

Differential regulation of fibroblast phenotype by drugs and the development of targeted antifibrotic therapies

Drugs that specifically target fibroblast signaling may provide directed control over the fibrotic
response. A major challenge in therapeutic development for fibrosis is that many drugs capable
of reducing fibrosis target non-specific regulatory pathways outside of the fibrotic response. For
example, the ALK5 inhibitor galunisertib targets the TGFβ receptor and shows promising
therapeutic reduction of fibrosis across organs[41–43]. While TGFβ receptor inhibition can reduce
fibrosis, recent efforts in target discovery have successfully identified new approaches to
mitigate fibrosis that are more fibroblast specific. For example, it was shown that activating
fibroblast-specific TLR4 in mice can drive the development of skin and lung fibrosis and that
TLR4 inhibition reduces αSMA expression and collagen production in fibroblasts [44]. Another
study showed that fibroblast-specific knockout of STAT3 ameliorates skin fibrosis, and that
pharmacological inhibition of STAT3 successfully reduces myofibroblast activation, collagen
accumulation, and dermal thickening in experimental fibrosis in mice[45]. Future work can
advance our understanding of how candidate drugs regulate specific components of the fibrotic
response in fibroblasts and provide targeted control of fibrosis.

Features of cardiac fibroblast phenotype

Traditional fibrotic markers are not always expressed in fibroblasts and exhibit significant
heterogeneity and context dependence, in this study we aimed to explore multiple features of
the fibrotic response[46]. Following the reduction of the original set of 137 single-cell features
from our high content image analysis, we identified a set of 18 phenotypic features of
fibroblasts that exhibit high heterogeneity in response to drug treatments (Figure S 3, Table S
2). Notably, many of the features measuring fiber texture for αSMA and F-actin show different
response patterns compared to features measuring overall expression level for those respective
proteins (i.e. aSMA integrated intensity versus aSMA long correlation). This distribution of
features indicates that the expression and organization of aSMA and actin filaments are
independently regulated by candidate drugs. The processes of αSMA protein expression and
fiber assembly have different degrees of contribution to pathological fibrosis. For example, a
recent study showed that fibroblasts can compensate for the loss of *Acta2* transcription and
form stress fibers using similar proteins, implying that stress fiber formation is more important
than αSMA production for the fibrotic response[47]. Incorporating an expanded set of
measurements in future fibrosis studies may provide greater resolution of the fibrotic
phenotype in response to therapies and help evaluate changes in pathologically relevant
features beyond protein expression.

Contributions of the LogiMML mechanistic machine learning approach

Mechanistic logic-based differential equation models have enabled systematic prediction of drug action, yet these models are limited by the availability of priori knowledge [13,48–50]. An alternative is machine learning, although 'black-box' ML approaches like artificial neural networks predict input-output relationships without mechanistic insight. In contrast, two recent studies combined mechanistic modeling with machine learning models like regression and visible neural networks to predict antibiotic stress on metabolism and drug synergies for cancer[29,51]. These 'white-box' approaches provide greater transparency of the intermediate layers between input and output[52].

The prior approach most similar to the LogiMML framework is 'white-box' machine learning proposed by Yang et. al[53]. In that study, a flux balance model of E. Coli metabolism with simple linear regression to predict metabolic reactions important to growth on particular carbon sources. In this study, we propose a method that combines regularized regression with a logic-based model to predict signaling pathways in response to signaling perturbations representative of drug effects.

Building on such advances for logic-based biological networks, our LogiMML mechanistic machine learning approach combines the flexible trainability of a machine learning model with the robust experimentally-determined internal network structure of a mechanistic model. In this study, we used the LogiMML model to predict signaling mechanisms that mediate how drugs regulate F-actin assembly and stress fiber formation in cardiac fibroblasts. However, this is just one of many possible applications for this modeling framework. The LogiMML approach is designed to work across multiple mechanistic modeling formalisms and types of experimental data, coupling the mechanistic model and data to predict mechanisms for the phenotype of interest. The flexible nature of LogiMML presents promising future applications to elucidate cell signaling across many disease areas.

Src kinase as a therapeutic target for fibrosis

Of the 13 drugs used in this study, the Src inhibitor WH-4-023 (WH) was one of three drugs that showed a strong reversal in TGFβ-induced actin filament assembly, αSMA, and procollagen I expression. WH was also effective at reversing the formation of actin-myosin stress fibers in response to TGFβ. Src inhibitors dasatinib, ponatinib, and saracatinib have all been used in clinical trials across different types of cancer[54–58]. In cancer, Src has been shown to promote proliferation and metastasis through many signaling targets including FAK, Akt, Ras, and PI3K[59–63].

Given that Src signaling affects many central regulatory pathways, recent studies have tested the potential for Src inhibition as a therapy for fibrotic disease. In a renal fibrosis study, blocking Src kinase using PP1 was shown to inhibit TGFβ-induced expression of collagen I, αSMA, and fibronectin[64]. In that study, Src inhibition was also shown the reduce the development of renal fibrosis in obstructed kidneys *in vivo* in mice, indicating Src inhibition as a potential renal

fibrosis and chronic kidney disease therapy. Another study focusing on lung fibrosis showed
that TGFβ induces Src kinase activity in lung fibroblasts and that Src is required for
myofibroblast contraction[65]. Further, inhibition of Src kinase *in vivo* with AZD0530 reduced scar
area and αSMA expression in mice with bleomycin-induced lung fibrosis[65].

While PI3K signaling has established relevance in fibrotic pathologies, this signaling pathway

has been heavily implicated in other disease models including cancer. Aberrant activation of
PI3K signaling has been shown to contribute to tumor progression in multiple cancers including
breast, lung, and ovarian cancers[66]. Changes in extracellular matrix remodeling have been
shown to influence many classically defined hallmarks of cancer, with ECM adhesion-induced
PI3K signaling shown to regulate self-sufficient cell growth via FAK signaling[67,68] . Given the large
overlap between regulators of cytoskeletal restructuring and tumor progression, therapeutic
targets like PI3K could be efficient future drug targets that can modulate pathways governing
multiple diseases. The interplay between cytoskeleton regulation and cancer progression
should be further explored to identify other central regulators that may modulate both cancer
and fibrotic disease progression.

In this study, we applied the LogiMML network to investigate how Src contributes to actin-
myosin stress fiber formation induced by TGFβ. We predicted that PI3K signaling contributes to
profibrotic Src signaling in cardiac fibrosis. This proposed mechanism is supported by previous
studies, showing that PI3K regulates fibroblast contractility and myofibroblast activation in skin
fibroblasts, and TGFβ-induced αSMA and collagen production in lung fibroblasts[39,40]. To validate
this proposed profibrotic role for PI3K, we show that PI3K inhibition reduced procollagen I
production and actin-myosin stress fiber organization in HCFs. While previous work has shown
that mechanical stretch, Rho-kinase, and myosin light chain kinase (MLCK) positively regulate
the organization of actin filaments into stress fibers, the role of PI3K's regulation of actin-
myosin stress fiber formation has not been thoroughly explored[69,70]. Here, we show that
treatment with PI3K inhibitor LY294002 (LY) significantly reduces stress fiber formation without
affecting the total amount of assembled actin filaments, implying PI3K positively and
specifically regulates actin-myosin stress fiber formation in cardiac fibroblasts. Future studies
should explore if Src kinase inhibitors mitigate cardiac fibrosis *in vivo*, and to what degree PI3K
kinase contributes to the regulation of cardiac fibrosis by Src.

Limitations and future directions

The main limitation of this study is that our modeling and experimental approaches address cell
signaling in cardiac fibroblasts *in vitro*, but do not address how fibroblasts respond to drugs in
an *in vivo* signaling environment. Our experimental data also captures some key fibrotic
proteins, but does not measure other fibrotic outputs of interest, like EDA fibronectin, and does
not capture a comprehensive signaling profile of the fibroblast. Despite these limitations, the
LogiMML framework was sufficient to predict a validated role for PI3K in promoting stress fiber
formation. Experimentally, future work could include proteomics or RNA-seq analysis of
fibroblasts to measure how drugs differentially regulate intracellular molecular profiles. To

maximize reproducibility across the drug screen, we used 2D culture on multi-well plates
treated with CellBind. More focused follow-on studies could perform secondary validations
with various extracellular matrix, stiffness or mechanical stretch. Future modeling work could
include simulated conditions for *in vivo* or *in vitro* co-culture conditions to incorporate the
signaling influence of other cell types. Given the flexibility of the LogiMML modeling approach,
these simulated data could be feasibly paired with respective experimental data to make
predictions for fibroblast signaling under new conditions.

Conclusions

421
In this study, we showed that drugs exhibit differential effects on cardiac fibroblast phenotype
and work via distinct mechanisms that can be predicted by logic-based mechanistic machine
learning. By expanding the microscopy feature set in the high content imaging pipeline, we
captured greater resolution of the fibroblast phenotype and measured how phenotypic
features changed in response to drugs. Using our LogiMML modeling approach, we predicted
signaling mechanisms for how pirfenidone and Src inhibitor WH-4-023 affect actin filament
assembly and actin-myosin stress fiber formation, respectively. We predicted that PI3K
regulates F-actin stress fiber formation, which we experimentally validated in human cardiac
fibroblasts. This study presents new features of fibroblast phenotype to be further explored in
fibrosis, identifies specific roles for PI3K in cardiac fibroblast signaling, and demonstrates an
adaptable mechanistic machine learning approach to predict signaling outcomes for fibrosis
that can be expanded to other diseases.

Methods

435
In vitro experiments in human cardiac fibroblasts

436
Primary human ventricular cardiac fibroblasts were purchased from PromoCell (PromoCell C-12375;
PromoCell GmbH, Germany). Cells were cultured in DMEM containing 10% FBS and 1% Pen/Strep, and
were kept in an incubator maintained at 5% CO_2. Cells were plated in a 96-well plate at 5,000 cells/well
and then grown in 10% FBS for 24 hours, serum starved for 24 hours, and then treated with the
following cytokine conditions for 96 hours: 0% FBS control media, 0% FBS media with 20ng/mL TGFβ1
(Cell Signaling Technology, 8915LC), and 0% FBS media with 10 ng/mL human IL1β (Cell Signaling
Technology, 8900SC), or TGFβ1 and IL1β combined. Cells were treated with these conditions either
alone or with 1 of 13 compounds at 1 of 3 concentrations. We determined drug concentrations via a
literature search, prioritizing concentrations that yielded significant effects *in vitro* in fibroblasts or
similar cell types. The drugs with their respective concentrations are as follows: [0.25,1,2] μg/ml of
anakinra (Kineret, SOBI Inc.), [1,5,10] μM valsartan (Sigma-Aldrich, SML0142-10MG), [0.2,1,2] μM BNP
(Sigma-Aldrich, B5900-.5MG), [1,5,10]μM valsartan combos respectively with [0.2,1,2] μM BNP,
[10,30,60]mM glutathione (Sigma-Aldrich, G4251-1G), [1,3,5] μM CW-HM12 (Cayman Chemical
Company, 19480), [10,20,50] μM salbutamol (Sigma-Aldrich, S8260-25MG), [5,10,25] μM marimistat
(Sigma-Aldrich, M2699-5Mg), [1,5,10] μM galunisertib (Selleck Chemicals, S2230), [12.5,25,50] μM
fasudil (Sigma-Aldrich, CDS021620-10MG), [10,25,50]μM SB203580 (Sigma-Aldrich, S8307-1MG),
[1,5,10] mg/mL pirfenidone (Sigma-Aldrich, P2116-10MG), [5,10,20] μM defactinib (MedChem Express,

HY-12289A), [5,10,20] μM WH-4-023 (Sigma-Aldrich, SML1334-5MG), and 20 μM LY294002 (Selleck
Chemicals, S1105). Cells were grown in these conditions for 72 hours.

Cells were then fixed in 4% PFA in PBS for 30 minutes, permeabilized and blocked for 1 hour in a solution
containing 3% BSA and 0.2% Triton, and then stained overnight at 4°C with a 1:500 Anti-Collagen I
antibody (Abcam, ab34710). After overnight incubation, cells were washed 3x in PBS and stained with
1:5000 Dapi, 1:1000 Phalloidin CruzFluor 647 Conjugate (Santa Cruz Biotechnology, sc-363797), 1:250 α-
Smooth Muscle Actin antibody (Sigma-Aldrich, C6198), and 1:1000 Goat-anti-Rabbit (ThermoFisher
Scientific, A-11034).

Microscopy and single-cell quantification

96-well plates we imaged using the Operetta CLS High-Content Analysis System (Perkin Elmer). All three
replicate wells for each condition were imaged and quantified. To quantify αSMA expression, an
automated image analysis pipeline was employed in CellProfiler (Broad Institute)[22]. Fibroblast nuclei
were identified by the DAPI signal. Next, the collagen-positive region corresponding to each nucleus was
segmented using the "propagate" algorithm, using the segmented nucleus as the seed. Next, Fibroblast
boundaries were segmented using the "propagate" algorithm, musing the segmented collagen region as
the seed. αSMA signal was integrated within each cell's boundary. Short, medium, and long texture
feature information was derived using the MeasureTexture module in CellProflier using texture scales of
2, 6, and 10 pixels respectively. Texture feature values were calculated by subtracting the smallest angle
value of a given feature from the largest angle value of that same feature for each cell. F-actin and
procollagen expressions were quantified similarly.

Statistics

475   Feature values for each well were determined by taking the median value of the feature across all cells
476   in the center tile of each well. Well median values were used as replicates (n=3). Significance was
477   determined using an ANOVA with Tukey's posthoc in comparisons between more than two groups, and
478   Student's T-test in comparisons between two groups. Automated data analysis and statistical
479   calculations were performed using Python 3.8.5 and the 'statsmodels' Python module version 0.13.2.

Model Simulations

Drug simulations in the fibroblast network model were performed as previously described using
MATLAB version 2022a[12,13,71]. Predicted node activity is calculated using logic-based Hill
differential equations. Agonist and antagonist drug relationships were represented by altering
the activation function of the target node, representing either competitive or non-competitive
drug interactions with the respective target. To better represent the cell-to-cell variability
observed in *in vitro* cell responses to treatments, we employed a previously developed
ensemble modeling approach combining multiple simulations with random normally distributed
parameters[71]. Ensemble simulations were performed using the MATLAB 'normrnd' function
from the 'Statistics and Machine Learning' toolbox to randomly sample parameters within a
normal distribution and simulation n of 100. The randomly sampled parameters and means of
the sampling ranges are as follows: baseline ligand inputs (0.25), mechanical input (0.85), drug
dose (0.85), and raised ligand inputs (0.6). The sampling range for each parameter was
calculated by $paramMean \pm COV * paramMean$ where COV=0.0396. This COV value, used to

scale stochasticity in the model was determined by taking the average coefficient of variation in
F-actin, procollagen I, and αSMA expression in human cardiac fibroblasts treated with TGFβ
from our *in vitro* experiments. Code for all modeling, regression and data analysis is available at
at https://github.com/andersnelson/Logic-based_MML.

LogiMML Network-Regression Coupling
The LogiMML mechanistic machine learning model is comprised of a network model layer and a
Ridge regression layer. The independent 'X' variables used to train the regression model are
node activity values from the network model predicted under each simulated drug and
environmental condition. To reduce model complexity, network nodes were clustered into 11
signaling modules derived from k-means clustering on a combined sensitivity and influence
analysis on the network model[12]. Sensitivity analysis was performed by systematically
perturbing individual node values and measuring the change in all other nodes in response to
the perturbed node. The influence matrix, the transposition of the sensitivity matrix, was
combined with the sensitivity matrix and this combined matrix was used for the k-means
clustering. The node activity values were averaged within each module, and these modules'
mean activity values were fed into the regression layer. The dependent 'Y' variables for this
model were experimentally measured values from our high-content imaging experiments in
human cardiac fibroblasts. Sensitivity knockout analysis was performed by simulating a given
drug and cytokine context int network model i.e. 'TGFβ+pirfenidone' and sequentially setting
each node ymax value to 0, measuring reduction or increase in the dependent variable e.g. 'F-
actin Intensity' upon knockdown.

Acknowledgements
We acknowledge Dr. Sarah Ewald at the University of Virginia Carter Immunology Center for
providing anakinra used in this study.

Funding
This study was funded by the NIH (HL137755, HL007284, HL160665, HL162925, 1S10OD021723-
01A1).

References
1.   Burgos Villar, K. N., Liu, X. & Small, E. M. Transcriptional regulation of cardiac fibroblast
phenotypic plasticity. *Curr Opin Physiol* **28**, 100556 (2022).

2.   Jacobs, A. K. *et al.* Systems of Care for ST-Segment-Elevation Myocardial Infarction: A Policy
Statement From the American Heart Association. *Circulation* **144**, e310–e327 (2021).

3.   Burke, R. M., Lighthouse, J. K., Mickelsen, D. M. & Small, E. M. Sacubitril/Valsartan Decreases
Cardiac Fibrosis in Left Ventricle Pressure Overload by Restoring PKG Signaling in Cardiac
Fibroblasts. *Circ Heart Fail* **12**, (2019).

4.   Fan, Z. & Guan, J. Antifibrotic therapies to control cardiac fibrosis. *Biomaterials Research 2016*
20:1 **20**, 1–13 (2016).

5.   Welch, M. P., Odland, G. F. & Clark, R. A. F. Temporal relationships of F-actin bundle formation,
collagen and fibronectin matrix assembly, and fibronectin receptor expression to wound
contraction. *J Cell Biol* **110**, 133–145 (1990).

6.   Travers, J. G., Kamal, F. A., Robbins, J., Yutzey, K. E. & Blaxall, B. C. Cardiac Fibrosis: The Fibroblast
Awakens. *Circ Res* **118**, 1021–40 (2016).

7.   Sun, K. H., Chang, Y., Reed, N. I. & Sheppard, D. α-smooth muscle actin is an inconsistent marker
of fibroblasts responsible for force-dependent TGFβ activation or collagen production across
multiple models of organ fibrosis. *Am J Physiol Lung Cell Mol Physiol* **310**, L824–L836 (2016).

8.   Avery, D. *et al.* Extracellular Matrix Directs Phenotypic Heterogeneity of Activated Fibroblasts.
Matrix Biol **67**, 90 (2018).

9.   Deng, C. C. *et al.* Single-cell RNA-seq reveals fibroblast heterogeneity and increased mesenchymal
fibroblasts in human fibrotic skin diseases. *Nature Communications 2021 12:1* **12**, 1–16 (2021).

10.   Gerckens, M. *et al.* Phenotypic drug screening in a human fibrosis model identified a novel class
of antifibrotic therapeutics. *Sci Adv* **7**, 3673 (2021).

11.   Fu, X. *et al.* Specialized fibroblast differentiated states underlie scar formation in the infarcted
mouse heart. *J Clin Invest* **128**, 2127–2143 (2018).

12.   Zeigler, A. C., Richardson, W. J., Holmes, J. W. & Saucerman, J. J. A computational model of
cardiac fibroblast signaling predicts context-dependent drivers of myofibroblast differentiation. *J*
Mol Cell Cardiol **94**, 72–81 (2016).

13.   Zeigler, A. C. *et al.* Network model-based screen for FDA-approved drugs affecting cardiac
fibrosis. *CPT Pharmacometrics Syst Pharmacol* **10**, 377–388 (2021).

14.   Kort, E. & Jovinge, S. Drug Repurposing: Claiming the Full Benefit from Drug Development. *Curr*
Cardiol Rep **23**, 1–7 (2021).

15.   Abdelsayed, M., Kort, E. J., Jovinge, S. & Mercola, M. Repurposing drugs to treat cardiovascular
disease in the era of precision medicine. doi:10.1038/s41569-022-00717-6.

16.   Khalil, H. *et al.* Fibroblast-specific TGF-β-Smad2/3 signaling underlies cardiac fibrosis. *J Clin Invest*
127, 3770–3783 (2017).

17.   Turner, N. A. *et al.* Interleukin-1α stimulates proinflammatory cytokine expression in human
cardiac myofibroblasts. *American Journal of Physiology-Heart and Circulatory Physiology* **297**,
H1117–H1127 (2009).

18.　Francis Stuart, S. D., De Jesus, N. M., Lindsey, M. L. & Ripplinger, C. M. The crossroads of
inflammation, fibrosis, and arrhythmia following myocardial infarction. *J Mol Cell Cardiol* **91**, 114–
22 (2016).

19.　Deten, A., Volz, H. C., Briest, W. & Zimmer, H.-G. Cardiac cytokine expression is upregulated in
the acute phase after myocardial infarction. Experimental studies in rats. *Cardiovasc Res* **55**, 329–
340 (2002).

20.　Bujak, M., Frangogiannis, N. G. & Hirszfeld, © L. The role of IL-1 in the pathogenesis of heart
disease THE IL–1 FAMILY OF CYTOKINES. doi:10.1007/s00005-009-0024-y.

21.　BUJAK, M. & FRANGOGIANNIS, N. The role of TGF-β signaling in myocardial infarction and cardiac
remodeling. *Cardiovasc Res* **74**, 184–195 (2007).

22.　Stirling, D. R. *et al.* CellProfiler 4: improvements in speed, utility and usability. *BMC
Bioinformatics* **22**, 1–11 (2021).

23.　Abbate, A. *et al.* Effects of interleukin-1 blockade with anakinra on adverse cardiac remodeling
and heart failure after acute myocardial infarction [from the virginia commonwealth university-
anakinra remodeling trial (2) (vcu-art2) pilot study]. *American Journal of Cardiology* **111**, (2013).

24.　Abbate, A. *et al.* Anakinra, a recombinant human interleukin-1 receptor antagonist, inhibits
apoptosis in experimental acute myocardial infarction. *Circulation* **117**, (2008).

25.　Abbate, A. *et al.* Interleukin-1 blockade inhibits the acute inflammatory response in patients with
st-segment–elevation myocardial infarction. *J Am Heart Assoc* **9**, (2020).

26.　Bracco Gartner, T. C. L. *et al.* Pirfenidone Has Anti-fibrotic Effects in a Tissue-Engineered Model of
Human Cardiac Fibrosis. *Front Cardiovasc Med* **9**, 459 (2022).

27.　Behr, J. *et al.* Pirfenidone in patients with progressive fibrotic interstitial lung diseases other than
idiopathic pulmonary fibrosis (RELIEF): a double-blind, randomised, placebo-controlled, phase 2b
trial. *Lancet Respir Med* **9**, 476–486 (2021).

28.　Sandbo, N. & Dulin, N. The actin cytoskeleton in myofibroblast differentiation: Ultrastructure
defining form and driving function. *Translational Research* **158**, 181 (2011).

29.　Yang, J. H. *et al.* A White-Box Machine Learning Approach for Revealing Antibiotic Mechanisms of
Action. *Cell* **177**, 1649-1661.e9 (2019).

30.　Rieg, T., Frick, J., Baumgartl, H. & Buettner, R. Demonstration of the potential of white-box
machine learning approaches to gain insights from cardiovascular disease electrocardiograms.
PLoS One **15**, e0243615 (2020).

31.　McNeish, D. M. Using Lasso for Predictor Selection and to Assuage Overfitting: A Method Long
Overlooked in Behavioral Sciences. *Multivariate Behav Res* **50**, 471–484 (2015).

32.　Hu, T. *et al.* Reactive oxygen species production via NADPH oxidase mediates TGF-β-induced
cytoskeletal alterations in endothelial cells. *Am J Physiol Renal Physiol* **289**, 816–825 (2005).

33. Huot, J. *et al.* SAPK2/p38-dependent F-Actin Reorganization Regulates Early Membrane Blebbing
during Stress-induced Apoptosis. *Journal of Cell Biology* **143**, 1361–1373 (1998).

34. Qian, Y. *et al.* PI3K induced actin filament remodeling through Akt and p70S6K1: implication of
essential role in cell migration. *Am J Physiol Cell Physiol* **286**, (2004).

35. Haralick, R. M., Dinstein, I. & Shanmugam, K. Textural Features for Image Classification. *IEEE*
Trans Syst Man Cybern **SMC-3**, 610–621 (1973).

36. Nagaraju, C. K. *et al.* Myofibroblast Phenotype and Reversibility of Fibrosis in Patients With End-
Stage Heart Failure. *J Am Coll Cardiol* **73**, 2267–2282 (2019).

37. Sandbo, N. *et al.* Delayed stress fiber formation mediates pulmonary myofibroblast
differentiation in response to TGF-β. *Am J Physiol Lung Cell Mol Physiol* **301**, 656–666 (2011).

38. Naumanen, P., Lappalainen, P. & Hotulainen, P. Mechanisms of actin stress fibre assembly. *J*
Microsc **231**, 446–454 (2008).

39. Conte, E. *et al.* Inhibition of PI3K prevents the proliferation and differentiation of human lung
fibroblasts into myofibroblasts: The role of class I P110 isoforms. *PLoS One* **6**, (2011).

40. Li, G., Li, Y. Y., Sun, J. E., Lin, W. H. & Zhou, R. X. ILK-PI3K/AKT pathway participates in cutaneous
wound contraction by regulating fibroblast migration and differentiation to myofibroblast.
Laboratory Investigation **96**, 741–751 (2016).

41. Luangmonkong, T. *et al.* Evaluating the antifibrotic potency of galunisertib in a human ex vivo
model of liver fibrosis. *Br J Pharmacol* **174**, (2017).

42. Masuda, A. *et al.* Promotion of liver regeneration and anti-fibrotic effects of the TGF-β receptor
kinase inhibitor galunisertib in CCl4-treated mice. *Int J Mol Med* **46**, (2020).

43. Dadrich, M. *et al.* Combined inhibition of TGFβ and PDGF signaling attenuates radiation-induced
pulmonary fibrosis. *Oncoimmunology* **5**, (2016).

44. Bhattacharyya, S. *et al.* TLR4-dependent fibroblast activation drives persistent organ fibrosis in
skin and lung. *JCI Insight* **3**, (2018).

45. Chakraborty, D. *et al.* Activation of STAT3 integrates common profibrotic pathways to promote
fibroblast activation and tissue fibrosis. *Nature Communications 2017 8:1* **8**, 1–16 (2017).

46. Ivey, M. J. & Tallquist, M. D. Defining the cardiac fibroblast. *Circulation Journal* vol. 80 2269–2276
Preprint at https://doi.org/10.1253/circj.CJ-16-1003 (2016).

47. Li, Y. *et al.* Loss of Acta2 in cardiac fibroblasts does not prevent the myofibroblast differentiation
or affect the cardiac repair after myocardial infarction. *J Mol Cell Cardiol* **171**, 117–132 (2022).

48. Tan, P. M., Buchholz, K. S., Omens, J. H., McCulloch, A. D. & Saucerman, J. J. Predictive model
identifies key network regulators of cardiomyocyte mechano-signaling. *PLoS Comput Biol* **13**,
(2017).

49. Ryall, K. A. *et al.* Network reconstruction and systems analysis of cardiac myocyte hypertrophy
signaling. *Journal of Biological Chemistry* **287**, 42259–42268 (2012).

50. Kraeutler, M. J., Soltis, A. R. & Saucerman, J. J. Modeling cardiac β-adrenergic signaling with
normalized-Hill differential equations: comparison with a biochemical model. *BMC Syst Biol* **4**,
157 (2010).

51. Kuenzi, B. M. *et al.* Predicting Drug Response and Synergy Using a Deep Learning Model of
Human Cancer Cells. *Cancer Cell* **38**, 672-684.e6 (2020).

52. Petch, J., Di, S. & Nelson, W. Opening the Black Box: The Promise and Limitations of Explainable
Machine Learning in Cardiology. *Canadian Journal of Cardiology* **38**, 204–213 (2022).

53. Yang, J. H. *et al.* A White-Box Machine Learning Approach for Revealing Antibiotic Mechanisms of
Action. *Cell* **177**, 1649-1661.e9 (2019).

54. Belli, S. *et al.* C-Src and EGFR inhibition in molecular cancer therapy: What else can we improve?
Cancers vol. 12 Preprint at https://doi.org/10.3390/cancers12061489 (2020).

55. Gucalp, A. *et al.* Phase II trial of saracatinib (AZD0530), an oral SRC-inhibitor for the treatment of
patients with hormone receptor-negative metastatic breast cancer. *Clin Breast Cancer* **11**, (2011).

56. Cortes, J. *et al.* Ponatinib dose-ranging study in chronic-phase chronic myeloid leukemia: a
randomized, open-label phase 2 clinical trial. *Blood* **138**, (2021).

57. Haura, E. B. *et al.* Phase I/II study of the Src inhibitor dasatinib in combination with erlotinib in
advanced non-small-cell lung cancer. *Journal of Clinical Oncology* **28**, (2010).

58. Twardowski, P. W. *et al.* A phase II trial of dasatinib in patients with metastatic castration-
resistant prostate cancer treated previously with chemotherapy. *Anticancer Drugs* **24**, (2013).

59. Beadnell, T. C. *et al.* Src-mediated regulation of the PI3K pathway in advanced papillary and
anaplastic thyroid cancer. *Oncogenesis* **7**, (2018).

60. Bunda, S. *et al.* Src promotes GTPase activity of Ras via tyrosine 32 phosphorylation. *Proc Natl
Acad Sci U S A* **111**, (2014).

61. Xiao, X. *et al.* Diallyl disulfide suppresses SRC/Ras/ERK signaling-mediated proliferation and
metastasis in human breast cancer by up-regulating miR-34a. *PLoS One* **9**, (2014).

62. Westhoff, M. A., Serrels, B., Fincham, V. J., Frame, M. C. & Carragher, N. O. Src-Mediated
Phosphorylation of Focal Adhesion Kinase Couples Actin and Adhesion Dynamics to Survival
Signaling. *Mol Cell Biol* **24**, (2004).

63. Mahajan, K. & Mahajan, N. P. PI3K-independent AKT activation in cancers: A treasure trove for
novel therapeutics. *Journal of Cellular Physiology* vol. 227 Preprint at
https://doi.org/10.1002/jcp.24065 (2012).

64. Yan, Y. *et al.* Src inhibition blocks renal interstitial fibroblast activation and ameliorates renal
fibrosis. *Kidney Int* **89**, (2016).

65.   Hu, M. *et al.* Therapeutic Targeting of Src Kinase in Myofibroblast Differentiation and Pulmonary
Fibrosis. *J Pharmacol Exp Ther* **351**, 87 (2014).

66.   Liu, P., Cheng, H., Roberts, T. M. & Zhao, J. J. Targeting the phosphoinositide 3-kinase pathway in
cancer. *Nature Reviews Drug Discovery* vol. 8 Preprint at https://doi.org/10.1038/nrd2926
(2009).

67.   Pylayeva, Y. *et al.* Ras- and PI3K-dependent breast tumorigenesis in mice and humans requires
focal adhesion kinase signaling. *Journal of Clinical Investigation* **119**, (2009).

68.   Pickup, M. W., Mouw, J. K. & Weaver, V. M. The extracellular matrix modulates the hallmarks of
cancer. *EMBO Rep* **15**, (2014).

69.   Katoh, K., Kano, Y., Amano, M., Kaibuchi, K. & Fujiwara, K. Stress fiber organization regulated by
MLCK and Rho-kinase in cultured human fibroblasts. *Am J Physiol Cell Physiol* **280**, (2001).

70.   Kaunas, R., Nguyen, P., Usami, S. & Chien, S. Cooperative effects of Rho and mechanical stretch
on stress fiber organization. *Proc Natl Acad Sci U S A* **102**, (2005).

71.   Zeigler, A. C. *et al.* Computational Model Predicts Paracrine and Intracellular Drivers of Fibroblast
Phenotype After Myocardial Infarction. *Matrix Biology* (2020) doi:10.1016/j.matbio.2020.03.007.

Figure 1: High-content microscopy screen for drugs that module fibroblast activation. A) Expression of
fibroblast activation markers procollagen I, F-actin, and αSMA in human cardiac fibroblasts upon
treatment of 13 drugs at 3 doses, under environmental contexts of TGFβ, IL1β, or both. Fold change
values show 'drug vs. no drug' Integrated Intensities for each protein. Panels B and C show
quantification and representative images of the effects of 50 μM fasudil, a Rho-kinase inhibitor, which
differentially regulates fibrotic protein expression. Panels D and E show quantification and
representative images of the effects of 10mg/mL pirfenidone, a non-specific inhibitor of TGFβ
expression, which consistently regulates fibrotic protein expression. *p≤0.05 ANOVA with Tukey's post-
hoc.

Figure 2: LogiMML logic-based mechanistic machine learning approach guides model revision and
predicts network mechanisms underlying pirfenidone suppression of F-actin. A) Original fibroblast
network model predicts no change in F-actin upon TGFβ or pirfenidone treatment. Experimental data
shows pirfenidone significantly reverses the increase of F-actin by TGFβ (data previously shown in figure
1 D). B) Schematic of the LogiMML approach for integrating logic-based network modeling with machine
learning to predict network mechanisms for cell phenotypes. The average activity within each network
module is mapped to predict fibroblast phenotypic features via a Ridge regression layer. C) The Coupled
LogiMML model predicts TGFβ and pirfenidone effects on F-actin that qualitatively match experimental
data shown in panel A. D) LogiMML ridge regression coefficients show predicted relative influence of
network modules on F-actin. E) LogiMML node knockdown sensitivity analysis in the context of
TGFβ+pirfenidone. Nodes from most influential modules are sequentially knocked down, predicting
change in F-actin upon knockdown. F) Schematic of the network mechanisms predicted for the actions
of pirfenidone on F-actin, derived from sensitivity analysis in panel E.

Figure 3: Survey of single-cell fibroblast phenotypic features in response to 13 drugs at 3 doses and 4
environmental contexts. A) 137 single-cell fibroblast features that quantify protein intensity, protein
localization, cell morphology, and fiber texture. This heatmap was organized on treatment and feature
axes by agglomerative hierarchical clustering. B) Principal component scores of experimental data
reduced to a set of 18 representative fibroblast features. C) Principal component loadings the reduced
of PCA scores and loadings define a primary axis of fibroblast activation with correlated protein
expression of procollagen, αSMA, and F-actin that is modulated by many drugs. Off-axis, the Src inhibitor
WH-4-023 modulated the cell texture feature Actin Long Angular Second Moment, which motivated
further study.

Figure 4: Logic-based mechanistic machine learning predicts the PI3K module to mediat how Src
inhibitor suppresses stress fibers, validated by subsequent experiments. A) Images of human cardiac
fibroblasts treated with baseline control stimulus, TGFβ, or TGFβ + 20 μM WH-4-023. B) Quantification
of Actin Long Angular Second Moment (ASM), a measure of actin uniformity and reduced stress fibers
based on images in panel A. C) Regression coefficients from the LogiMML mechanistic machine learning
model that predicts network modules that regulate actin long ASM. D) Knockdown sensitivity analysis
predicting individual proteins that regulate actin long ASM in the TGFβ+WH-4-023 signaling context. E)
Signaling schematic for WH-4-023 effect on actin long ASM, derived from sensitivity analysis in panel D.
F) Human cardiac fibroblasts treated with 20 μM PI3K inhibitor LY294002 or baseline control stimulus,
measuring F-actin and procollagen expression. G) Quantification of long actin Angular Second Moment
(measure of actin uniformity), F-actin integrated intensity, and Procollagen I integrated intensity.
*p≤0.05 ANOVA with Tukey's post-hoc in panel B, and *p≤0.05 Student's T-test in panel G.

A Network Model

B Experiment

D Coupled Model

C Network Model

Ridge Regression Model

$$Y = \beta_1 X_1 + \beta_2 X_2 + \cdots \beta_{11} X_{11}$$

Phenotypic Features

F-actin Assembly

E Relative influence of modules on F-actin

F Knockdown sensitivity for Pirfenidone+TGFB context

G Signaling Modules

Pirfenidone

ETAR → ROS → p38

TNFa → PI3K → Akt

TGFB → CBP, NOX

F-actin

A

Control　TGFβ　TGFβ+WH

75 um

B

Actin Long Angular Second Moment

×10⁻³

TGFβ	−	+	+
WH	−	−	+

C

Relative influence of modules on
Actin Long Angular Second Moment

Smad3
Mech
Ras_Raf
Autocrine
PI3K
STAT
P38_Calcium
PKA
PDGF
MKK3
NP

−0.5 −0.4 −0.3 −0.2 −0.1 0.0 0.1 0.2 0.3

D

Knockdown sensitivity for WH+TGFB context

0.10

0.05

0.00

−0.05

−0.10

MKK4 Rac1 ROCK FA SRF FAK TRAF Rho Akt

Module
PI3K
Mechanical

E

WH-4-023

Signaling Modules

Mech → Rho → MKK4

TNFa → PI3K → Akt

Actin Long Angular
Second Moment

F

Control　LY

200 um

G

Actin Long Angular Second Moment

×10⁻³

LY	−	+

F-actin Intensity (AU)

LY	−	+

Procollagen I Intensity (AU)

LY	−	+