Commensal Bacteria Maintain a Qa-1[b]-restricted Unconventional CD8[+] T Population in Gut Epithelium

Jian Guan[1,*], Michael Manoharan Valerio[2], J. David Peske[1], Chansu Park[1], Ellen A. Robey[2,3], Scheherazade Sadegh-Nasseri[1,3], Nilabh Shastri[1,3]

[1]Department of Pathology and the Institute of Cell Engineering, Johns Hopkins University School of Medicine, Baltimore, MD 21287, USA
[2]Division of Immunology and Molecular Medicine, Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA 94720, USA
[3]Co-senior author
*Correspondence: jguan5@jh.edu

Abstract

Intestinal intraepithelial lymphocytes (IELs) are characterized by an unusual phenotype and developmental pathway, yet their specific ligands and functions remain largely unknown. Here by analysis of QFL T cells, a population of CD8[+] T cells critical for monitoring the MHC I antigen processing pathway, we established that unconventional Qa-1[b]-restricted CD8[+] T cells are abundant in intestinal epithelium. We found that QFL T cells showed a Qa-1[b]-dependent antigen-experienced phenotype in the spleen and small intestine of naïve wild-type mice. The splenic QFL T cells showed innate-like functionality exemplified by rapid response to cytokines or antigen, while the gut population was refractory to stimuli. Microbiota was required for the maintenance, but not the initial gut homing of QFL T cells. Interestingly, monocolonization with *Pediococcus pentosaceus,* which expresses a peptide that cross-activated QFL T cells, was sufficient to maintain QFL T cells in the intestine. Thus, microbiota are critical for shaping the Qa-1[b]-restricted IEL landscape.

Introduction

The display of peptides by MHC class I (MHC I) molecules on the cell surface is critical for CD8[+] T cell immune surveillance[1]. Generation of a peptide repertoire which accurately reflects intracellular events, such as viral infection or mutations, relies on a functional antigen processing and presentation pathway. After cytosolic cleavage of protein precursors and transport of peptides into the endoplasmic reticulum (ER), the peptide intermediates are further customized by ER aminopeptidase associated with antigen processing (ERAAP) until 'ideal' peptides that fit the groove of MHC I molecules are eventually shuttled and displayed on the cell surface[2, 3, 4]. Disruption of each step of the pathway can lead to immunological dysfunction[5, 6, 7]. The critical role of ERAAP in the antigen processing pathway has been established through extensive studies of ERAAP deficient (ERAAP-KO) cells and mice[8, 9, 10]. Loss of ERAAP severely disrupts the peptide repertoire presented by both the classical MHC Ia and the nonclassical MHC Ib molecules [11, 12]. Qa-1[b], a nonclassical MHC Ib molecule, has been shown to present a significantly increased number of peptides on the cell surface of ERAAP-KO cells as compared with wild-type (WT) cells[13]. One peptide presented by Qa-1[b], FYAEATPML (FL9) (with the Qa-1[b]-FL9 complex termed QFL), was identified as an immunodominant ligand uniquely presented on ERAAP deficient cells[14]. The CD8[+] T cells that specifically recognize this ligand are thus named QFL-specific T (QFL T) cells.

Early analysis of QFL T cells revealed the unusual nature of these cytotoxic CD8[+] T cells. Unlike conventional antigen-specific CD8[+] T cells which are typically detected at a frequency of 1 in $10^5 \sim 10^6$, QFL T cells are present at a frequency 10-fold higher in the spleen of naïve mice. Interestingly, the bulk splenic QFL T population displayed a CD44[hi]CD122[+] antigen-experienced phenotype[14]. T cell receptor (TCR) analysis of QFL T cells revealed that a large proportion of the QFL T population expresses an invariant TCRα chain Vα3.2Jα21[15]. These traits of QFL T cells indicate their potential similarity to other unconventional T cells, such as invariant NKT (iNKT) and mucosal associated invariant T(MAIT) cells, which are typically characterized by recognition of ligands present by non-classical MHC Ib, expression of invariant TCRs, residence in nonlymphoid tissues and innate-like functions[16, 17]. The tissue distribution and functions of QFL T cells remains to be fully elucidated.

The gut mucosa is an immunologically complex niche with abundant lymphoid populations [18]. The small intestinal intraepithelial lymphocyte (siIEL) compartment is populated by antigen-experienced T cells of both TCRγδ+ and TCRαβ+ lineage with characteristic CD8αα expression. The development of the CD8αα+CD4-CD8αβ-TCRαβ+ population (CD8αα+ IEL), categorized as natural IELs (natIELs) because they acquire their activated phenotype in the thymus, has been extensively studied [19], yet little is known about their antigen specificity and TCR repertoire. Emerging evidence shows that nonclassical MHC Ib molecules are important for the development and effector function of CD8αα+ IELs[20]. For instance, while loss of classical MHC Ia molecules showed little impact on the CD8αα+ IEL population, these cells were decreased in mice deficient for Qa-2 [21, 22, 23]. Furthermore, a population of CD8αα+ IEL precursors which preferentially expresses Vα3.2 was shown to be decreased in number in the absence of CD1d[24]. However, whether Qa-1b is involved in shaping the CD8αα+ IEL population remains to be studied. In addition to MHC molecules, gut microbiota plays a critical role in the establishment and shaping of the gut immune system. Studies of germ-free (GF) mice have shown extensive defects in gut-associated lymphoid tissues together with morphological changes in the intestine associated with the absence of gut microbiota[25]. Notably, despite the critical role of gut microbiota in the establishment of the gut immune system, natIELs that do not rely on cognate antigens in the periphery are believed to home to the gut independent of microbes[26] .

Here we found abundant antigen-experienced QFL T cells in both the spleen and small intestinal IEL (siIEL) compartment of naïve WT mice. The splenic Vα3.2+QFL T cells expressed high level of CD44 and showed typical innate-like functions including hyperresponsiveness to cytokines and rapid IFN-γ production in response to antigen. In contrast, the population of the same antigen specificity in the gut phenotypically resembled natIEL and were likewise functionally quiescent. Analysis of Qa-1b deficient mice revealed that while Qa-1b was not absolutely required for the development/survival of splenic or siIEL QFL T cells, it was essential for imprinting their unconventional phenotype. Furthermore, analysis of GF mice showed that gut microbiota was needed for the long-term maintenance of QFL T cells. Interestingly, we found that maintenance of the gut Vα3.2+QFL T was associated with colonization by the commensal bacterium *Pediococcus pentosaceus* (*P. pentosaceus*) which expresses an FL9 homologue that could cross-activate QFL cells. Overall, these results establish Vα3.2+QFL T

cells as a unique population of unconventional CD8$^+$ T cells that rely on nonclassical MHC Ib to acquire their phenotype, and gut microbiota to be properly maintained in the intestine.

Results

Vα3.2$^+$QFL T cells are abundant in the spleen and gut of naïve wild-type mice

To identify QFL T cells in tissues of naïve WT mice, we generated Qa-1^b-FL9 dextramers (QFL-Dex) based on the 'dextran-doping' technique[27]. In comparison with Qa-1^b-FL9 tetramers (QFL-Tet), QFL-Dex showed improved sensitivity and specificity for QFL T cell detection (**Supplementary Fig. 1**), and further allowed enrichment of QFL T cells using magnetic beads. The average number of QFL T cells – defined as CD45$^+$CD19$^-$TCRβ$^+$CD4$^-$QFL-Dex-PE$^+$APC$^+$ (**Fig.1a**) – detected in the siIEL compartment of naïve WT mice was comparable to the population in the spleen. Strikingly, however, QFL T cells were present at a frequency of ~1 in 1,000 of the siIEL CD8$^+$ T cells, which was 10 times more frequent than in the spleen (**Fig.1 b, c**). Both the splenic and siIEL QFL T cells were essentially all CD8$^+$ T cells, as these cells were barely detectable within the CD4$^+$ T population (**Supplementary Fig. 2**). Consistent with prior studies, ~80% of the splenic QFL T cells expressed the 'invariant' TCR Vα segment Vα3.2 which was significantly higher than the proportion of Vα3.2-expressing cells within the total CD8$^+$ T population in the spleen (<5%) (**Fig.1 d,e**) [15]. Interestingly, the QFL T cells in the siIEL compartment also contained a relatively larger proportion of Vα3.2$^+$ cells (~60%) compared to the total CD8$^+$ TCRαβ siIELs (<30%) (**Fig.1 f,g**). The average percentage of Vα3.2$^+$ cells within the QFL T population (20.1%) was lower in the siIEL compartment than in the spleen (63.3%). Vα3.2$^+$ QFL T cells were nevertheless abundantly present in both the spleen and gut of naïve WT mice.

The unconventional phenotype of Vα3.2$^+$QFL T cells

We have previously observed a high percentage of CD44^{hi} cells within the bulk splenic QFL T population of naïve WT mice[14, 15]. Here we further investigated the association between preferred Vα3.2 usage by QFL T cells and their phenotype. Strikingly, we observed that

Vα3.2[+]QFL T cells contained a significantly larger proportion of CD44[hi] cells (~90%) than the Vα3.2[-]QFL T cells, suggesting that the antigen experienced phenotype of the QFL T population was mainly contributed by the Vα3.2-expressing subpopulation (**Fig.2 a, b**). In contrast, only around 20% of the total Vα3.2[+]CD8[+] T population expressed high CD44. Interestingly, the percentage of CD44[hi] cells within the Vα3.2[+]CD8[+] T population was slightly higher compared to the Vα3.2[-] population (**Supplementary Fig. 3 a**), which was in line with a previous report [28]. We thus conclude that although both Vα3.2[+] and Vα3.2[-] cells were detected within the splenic QFL T population, a memory phenotype was specifically enriched on Vα3.2[+]QFL T cells.

Given the strikingly high frequency of QFL T cells in the gut, we further investigated the phenotype of these cells in this location. We first assessed CD8αα expression, a hallmark of the unconventional phenotype of natIELs, on Vα3.2[+] and Vα3.2[-] QFL T cells in the siIEL compartment of naïve WT mice. A relatively larger fraction of Vα3.2[+]QFL T cells lacked CD4 and CD8αβ expression but expressed CD8αα compared to the Vα3.2[-] subpopulation suggesting that the majority of Vα3.2[+] QFL T cells were CD8αα[+] natIELs (**Fig.3 a, b**). No such difference was observed between the total Vα3.2[+] and Vα3.2[-] CD8[+] IEL populations (**Supplementary Fig. 3 b**). These observations indicate that the Vα3.2[+] QFL T cells in both the spleen and siIEL compartment are unconventional CD8[+] T cells which likely are derived from the same precursors in thymus.

The Qa-1[b]/ERAAP/TAP-dependent unconventional phenotype of Vα3.2[+]QFL T cells

We further investigate what drives the unconventional phenotype of Vα3.2[+] QFL T cells by analyzing mice deficient of molecules crucial for QFL antigen presentation including Qa-1[b], ERAAP or TAP (Qa-1[b]-KO, ERAAP-KO, TAP-KO). While we consistently observed loss of both the splenic and siIEL Vα3.2[+]QFL T cells in TAP-KO mice, the populations were detectable in both tissues of Qa-1[b]-KO or ERAAP-KO mice with minorly reduced population size in the spleen of Qa-1[b]-KO mice (**Fig.3 a**). Strikingly, Vα3.2[+] QFL T populations showed reduced percentages of CD44[hi] cells in the spleen of Qa-1[b] and ERAAP deficient mice with the reduction being more significant in Qa-1[b]-KO mice. In contrast, the phenotype of Vα3.2[-] QFL T cells was

not significantly affected (**Fig.3 b, c**). Interestingly, similar pattern of phenotype change was observed for the siIEL Vα3.2$^+$ QFL T cells which showed almost complete loss of the unconventional natIEL phenotype including CD8αα expression in Qa-1^b-KO mice (**Fig.3 d,e, Supplementary Fig. 4**). These results demonstrate TAP, Qa-1^b and ERAAP all play a role in the establishment the unconventional Vα3.2$^+$ QFL T population with TAP being required for the presence of the population, Qa-1^b being absolutely required and ERAAP being partially required for the unconventional phenotype imprinting of these cells. It is worth noting that background splenic CD8$^+$ T cells showed a similar loss of memory phenotype among the Vα3.2$^+$CD8αβ$^+$, but not Vα3.2$^-$CD8αβ$^+$, population in Qa-1^b-KO and ERAAP-KO mice (**Supplementary Fig. 3 c**). Thus, the unconventional Vα3.2$^+$QFL T cells specifically, and some of the background Vα3.2$^+$ splenocytes more broadly, require Qa-1^b and ERAAP to acquire their memory phenotype.

Vα3.2$^+$QFL T cells are functionally innate-like in the spleen but quiescent in gut

To further investigate the functional features of Vα3.2$^+$QFL T cells, we utilized transgenic mice (QFLTg) expressing the predominant invariant TCR found on QFL T cells (*Vα3.2Jα21, Vβ1Dβ1Jβ2-7*), referred to hereafter as QFLTg cells. Abundant QFLTg cells or control OT-1 cells were detected in both the spleen and siIEL compartment of the respective TCR transgenic mice (**Supplementary Fig. 5 a**). Phenotypically, the QFLTg cells recapitulated the memory phenotype of polyclonal Vα3.2$^+$QFL T cells with high CD44 expression in the spleen and CD8αα expression in the siIEL compartment. In contrast, OT-1 cells were phenotypically similar to conventional naïve CD8$^+$ T cells, as they expressed low to intermediate levels of CD44 in the spleen and lacked CD8αα expression in gut (**Supplementary Fig. 5 b,c**).

Cytokines including IL15 and the combination of IL7/IL18 have been implicated in the maintenance and survival of both 'virtual' memory T cells and IELs [28, 29, 30, 31]. We thus compared the proliferation of QFLTg and OT-1 cells in response to these cytokines. In keeping with their virtual memory surface marker phenotype, splenic QFLTg cells were significantly more responsive to IL15 or IL7/IL18 stimulation than OT-1 cells (**Fig.4 a**). At 72hs, ~80% of QFLTg cells had proliferated in response to the cytokines, compared with only ~50% of OT-1 cells (**Fig.4 b**). Interestingly, T cells isolated from the siIEL compartment of QFLTg or OT-1

mice showed a distinct pattern of cytokine responsiveness from the splenic T cells. First, both QFLTg and OT-1 cells from siIEL were less responsive to IL15 than the splenic population. Second, siIEL QFLTg cells showed significantly reduced proliferation in response to the cytokines, especially to the combination of IL7/IL18, as compared with OT-1 cells (**Fig.4 c, d**).

To ensure that altered effector function was not skewed by the high specific TCR frequency in TCR transgenics, we generated partial hematopoietic chimeric mice with physiological precursor frequencies (QFLTg_WT) by transferring bone marrow cells from GFP[+]QFLTg mice into nonirradiated naïve WT neonates[32]. OT-1_WT chimera mice were generated in parallel as representative of conventional CD8[+] T cells (**Fig.4 e**). This experimental design allowed us to track the GFP[+] donor-derived T cells (GFP[+]CD45[+]CD19[-]TCRβ[+]CD4[-]Vα3.2[+]) separately from the recipient's endogenous T cells (**Fig.4 f**). Based on the hyperresponsiveness of splenic QFLTg cells to cytokines, we hypothesized that these cells would exert effector functions more rapidly than conventional CD8 T cells, similar to the known innate-like T cells such as iNKT. We measured IFN-γ production by QFLTg or OT-1 cells from splenocytes or siIEL cells of naïve QFLTg_WT or OT-1_WT chimera mice stimulated the with 2μM FL9 or SIINFEKL(SL8) peptide respectively for 4.5 hours directly *ex vivo*. A strikingly large fraction (~60%) of splenic QFLTg cells expressed high levels of IFN-γ under these conditions, while only 6% of OT-1 cells did so (**Fig.4 g,h**). Notably, despite the low responsiveness to epitopes, siIEL QFLTg cells showed relatively higher basal levels of IFN-γ expression than OT-1 cells (**Fig.4 i,j**). In contrast to the hyperresponsiveness of splenic QFLTg cells to their ligand, neither QFLTg nor OT-1 cells from the siIEL compartment showed significant IFN-γ production at 4.5 hours. The low IFN-γ production of IEL QFLTg cells was in keeping with their natIEL phenotype -antigen-experienced yet functionally quiescent[19, 33]. We conclude that Vα3.2[+]QFL T cells in the spleen are functionally innate-like, whereas the population with the same TCR specificity in the small intestine shows functional features of natIEL .

Gut microbiota is associated with retention but not homing of QFL T cells

The establishment and function of the intestinal immune system is intimately associated with homeostasis of the gut microbial community, as evidenced by an altered gut immune cell

composition in germ-free mice [25, 34, 35]. To investigate if an association exists between gut microbiota and gut QFL T cells, we compared the numbers and frequencies of QFL T cells in the spleen and siIEL compartment of naïve specific-pathogen-free (SPF) and germ-free (GF) WT mice of various ages. First, we found the absolute number of QFL T cells in the spleen was unaffected by the absence of gut microbiota in either young or old mice. The frequency of QFL T cells was relatively higher in GF than in SPF mice, as a result of a decreased number of non-QFL CD8$^+$ T cells in the spleen of GF mice (**Fig.5 a, b**). Interestingly, the percentage of CD44^{hi} cells within the total CD8$^+$ T population was proportionally higher in the spleen of GF WT than in SPF WT mice regardless of their Vα3.2 expression (**Supplementary Fig. 6 a**). This observation suggested that gut microbiota might be more closely associated with the presence of conventional naïve CD8$^+$ T cells rather than the memory phenotype CD8$^+$ T population in the spleen. Second, while the number and frequency of QFL T cells in siIEL compartment showed no significant difference between GF and SPF mice of relatively young age (8~17-week), the population was gradually lost in old GF mice (18~22-week) (**Fig.5 c, d**), with the decrease in GF mice starting from around 16 weeks of age (data not shown). In addition, the QFL T cells detected in the spleen or siIEL compartment of young SPF or GF mice showed similar expressions of Vα3.2, CD44 and CD8αα (**Supplementary Fig. 7**). We conclude that although QFL T cells could home to gut in the absence of microbiota, microbe-derived antigens or signals are required for their maintenance there. Interestingly, we observed that the average percentage of CD8αα$^+$ cells among the total TCRβ$^+$CD4$^-$ T population was proportionally larger in GF mice than SPF mice regardless of Vα3.2 expression, although the Vα3.2$^-$ population showed a more significant increase (**Supplementary Fig. 6 b**). This was mainly a result of a decreased number of CD8αβ$^+$ T cells in the siIEL of GF mice (data not shown). Nevertheless, here we showed that Qa-1^b restricted T cells could home to the small intestine epithelium and display a natIEL phenotype independent of gut microbiota.

QFL T cells cross-react with a microbial antigen and are maintained by commensal bacteria

We next investigated how the QFL T population was retained in the gut of SPF WT mice, hypothesizing that the presence of specific antigen might play a role. QFL T cells specifically

recognize the Qa-1^b-FL9 complex which is uniquely presented on ERAAP deficient cells. Because it is unlikely this ligand is constantly presented in gut epithelium under homeostatic conditions, we reasoned that QFL T cells might be retained in the gut through exposure to FL9 homologue peptide(s) expressed by commensal bacteria that colonize small intestine.

To identify potentially cross-reactive peptides, we first determined the key residues of the FYAEATPML(FL9) peptide that affected TCR recognition by QFL T cells. Due to the lack of structural information on QFL-TCR complex, we hypothesized that the glutamic acid at P4 and threonine at P6 were likely to be critical based on the nature and position of amino acids. The QFL T hybridoma cell line BEko8Z which expresses TCR-induced β-galactosidase (LacZ) was used to test whether FL9 peptide variants with the hypothesized key residues substituted by alanine altered T cell responses. By measuring the LacZ production of BEko8Z cells in response to Qa-1^b-expressing Lmtk$^-$ (L-Qa-1^b) cells pulsed with FYAAATPML(FL9-P4A) or FYAEAAPML(FL9-P6A), we found that replacement of the P6 threonine reduced TCR recognition, while replacement of the P4 glutamic acid led to the complete loss of the BEko8Z response (**Supplementary Fig. 8 a**). We thus conclude that both the P4 and P6 positions are important with the P4 glutamic acid being particularly critical for QFL T recognition.

To identify potential bacterial FL9 homologous peptides, we aligned the FL9 peptide sequence with proteomes of the five dominant gut commensal bacterial phyla, including Actinobacteria, Bacteroidetes, Firmicutes, Proteobacteria, Verrucomicrobia from the NCBI Swissprot Non-redundant UniProtKB database[36]. The alignment result was further curated to 30 candidate peptides based on their homology with FL9 peptide at position P4 and P6 (**Supplementary Fig. 8 b, Supplementary Table.1**). By measuring the LacZ production of BEko8Z cells in response to L-Qa-1^b cells pulsed with 50nM of each of the candidate peptides, we identified a single stimulatory peptide FYAEDGTPIL(FL10) that is expressed in *Pediococcus pentosaceus* (*P. pentosaceus*), a gram-positive lactic acid bacteria(LAB) that colonizes the small intestine (**Fig.6 a**). Sensitivity to the FL9 peptide was ~1,000 fold greater than to FL10, but BEko8Z cells could recognize FL10 at concentrations as low as 1.5nM. In contrast, FYAEDDTPIV(FV10), a FL10 homologous peptide expressed in *Lactobacillus johnsonii* which is a common LAB that colonize GI tract, failed to activate BEko8Z cells (**Fig.6 b, Supplementary Fig. 8 b**).

To further investigate the association between *P. pentosaceus* and Vα3.2$^+$QFL T cells in gut, we first generated QFLTg_WT partial hematopoietic chimeras using germ-free WT mice (GF QFLTg_WT) and monocolonized the GF chimera mice with *P. pentosaceus* at 8 weeks of age. The mice were analyzed 8 weeks later (**Fig.6 c**). The donor-derived QFLTg cells and the recipient-derived Vα3.2$^+$QFL T cells were distinguished by GFP expression and further defined as the CD45$^+$CD19$^-$TCRβ$^+$CD4$^-$Vα3.2$^+$QFL-Dex-PE$^+$ population on flow cytometry as was described earlier. While the donor-derived QFLTg populations were present and phenotypically unaltered in the spleen and siIEL compartment of GF QFLTg_WT chimera mice at 8 weeks of age, they were barely detectable in the GF chimera mice at 16 weeks of age (**Supplementary Fig. 9, Fig.6 d**). Strikingly, monocolonization of the GF QFLTg_WT chimera mice with *P. pentosaceus* completely restored the donor-derived QFLTg population in both tissues in 16-week GF chimera mice (**Fig.6 d**). We then further tested whether restoration of the gut QFL T population was specificaly associated with *P. pentosaceus* by monocolonizing GF WT mice with *P. pentosaceus* or *L. johnsonii* (**Fig.6 c**). While the endogenous Vα3.2$^+$QFL T population in the siIEL compartment was retained by *P. pentosaceus* colonization of GF mice at 16 weeks of age, *L. johnsonii* colonization failed to restore the population to level comparable with SPF or *P. pentosaceus* colonized mice. On the other hand, maintenance of the endogenous splenic QFL T population was independent of microbes, in line with our earlier observations (**Fig.6 e**). Donor-derived QFLTg and endogenous Vα3.2$^+$QFL T cells in the spleen likely showed discrepant results due to different sources of precursors for the two populations. Unlike the endogenous population which was constantly replenished, the donor-derived QFLTg cells rose from a fixed number of bone marrow cells, and gradually decreased in number as the animal age (**Supplementary Fig. 10**). Thus, presence of the ligand which was supplied by peptide derived from *P. pentosaceus* might be required for the donor-derived QFLTg cells to be retained in the spleen, whereas the endogenous splenic population was unaffected by microbes. Phenotype analysis revealed that both the donor-derived QFLTg cells and the endogenous Vα3.2$^+$QFL T cells from the colonized GF chimera mice showed a memory phenotype identical to that seen in the SPF chimera mice (**Supplementary Fig. 11**). We thus conclude that the late-life maintenance of Vα3.2$^+$QFL T cells in the small intestine was dependent on the commensal bacterium *P. pentosaceus* which expresses a FL9 homologue peptide. While the presence of the bacterial

FL10 peptide likely plays a major role in retention of gut QFL T cells, we do not exclude the possible minor contributions of other commensal bacterial factors. In addition, the memory phenotype of QFL T cells was likely acquired through microbe-independent mechanisms.

Discussion

The small intestine epithelium harbors a highly heterogenous intraepithelial lymphocyte population which is comprised of a large proportion of $CD4^-CD8\alpha\beta^-CD8\alpha\alpha^+$ T cells. Due to the heterogeneity, it has been challenging to identify or study a particular antigen-specific T cell clone from the population. Here we found that QFL T cells, a $V\alpha3.2$-expressing $CD8^+$ T cell population which specifically recognizes the $Qa-1^b$-FL9 ligand presented on ERAAP-KO cells, naturally resided in both the spleen and small intestine epithelium of naïve WT mice. Further characterization of QFL T cells revealed their unconventional phenotype, functionality and intimate association with gut microbiota.

Unlike conventional memory T cells which are generated upon encounter with their cognate peptide antigens on classical MHC molecules, 'innate' memory T cells rise under homeostatic conditions in response to neonatal lymphopenia, high levels of IL-4 or exposure to self-antigens in naïve mice [37, 38]. Here, we found that splenic $V\alpha3.2^+$QFL T cells display a memory phenotype and corresponding rapid effector function, but acquire this phenotype in a novel $Qa-1^b$ dependent manner. It is worth mentioning that ERAAP deficiency which leads to universal expression of the QFL epitope only showed minor effect on the $V\alpha3.2^+$QFL T cells indicating potential agonist selection of the population. The unique developmental mechanism of QFL T cells is elaborated by Manoharan Manoharan Valerio et al. Given the unique dependence on $Qa-1^b$ expression for the phenotype imprinting of QFL T cells, it is possible that QFL T cells are exposed to transiently induced QFL epitope on cells due to various intracellular stressors that might affect ERAAP function. Alternatively, QFL T cells might cross-react with $Qa-1^b$ presented FL9 homologue peptide(s). Indeed, we found that $V\alpha3.2^+$QFL T cells cross-reacted with one such variant of FL9 peptide expressed in commensal bacteria and presented by $Qa-1^b$. However, the $Qa-1^b$-FL10 ligand itself is unlikely to be directly associated with the memory imprinting for QFL T cells as these cells showed unaltered phenotype in germ-free mice. There might be other

unidentified self-peptide(s) presented by Qa-1[b] that are involved in the process. It is also possible that there is some level of degeneracy in the TCR-Qa-1[b] interaction, so the precise peptide being presented is less critical than the presence of any peptide-Qa-1[b] complex. Deciphering among these intriguing possibilities remains to be investigated. Notably, in line with a previous study, we observed that the Vα3.2-expressing CD8αβ[+] T population was comprised of a relatively larger proportion of memory phenotype cells in the spleen than the non-Vα3.2 expressing population[28]. In addition, these memory phenotype non-QFL Vα3.2[+]CD8αβ[+] T cells were lost in Qa-1[b]-KO mice. These results indicate that Vα3.2[+]QFL T cells are likely only one prototypical example of Qa-1[b] restricted Vα3.2[+]CD8[+] T cells clones with memory phenotype and innate-like function.

Strikingly, we detected abundant Vα3.2[+]QFL T cells in small intestine epithelium which showed the signature phenotype of natIELs including expression of CD8αα and lack of CD4, CD8αβ, CD5 and CD90. Similar to the splenic population, although the population size was unaffected by Qa-1[b] deficiency, the CD8αα[+] phenotype of Vα3.2[+]QFL T cells in the gut was strongly Qa-1[b]-dependent. It is so far reported that CD8αα[+] IELs can be restricted to K[b]D[b], CD1d or even unknown MHC Ib molecules[24, 39]. Here by characterization of the gut Vα3.2[+]QFL T cells, we showed evidence of abundant Qa-1[b]-restricted TCRαβ[+] natIELs being present in gut.

Functionally, the gut Vα3.2[+]QFL T cells showed features consistent with their CD8αα[+] natIEL phenotype, which is generally associated with a suppressed state. However, it is worth emphasizing that although the siIEL Vα3.2[+]QFL T cells showed delayed and reduced IFN-γ production in response to the cognate peptide, they showed higher basal levels of intracellular IFN-γ than CD8αβ[+] IELs. It has been proposed that spontaneous IFN-γ section by IELs may be an important component of immunosurveillance at the mucosal surface, capable of identifying and eliminating transformed cells[40, 41]. siIEL QFL T cells may likewise be poised for immediate function, eliminating cells with dysfunctional ERAAP. CD8αα[+] IELs are intrinsically highly cytotoxic, as was shown by PMA/ionomycin induced production of high level of IFN-γ[42]. However, our observation of the relative hyporesponsiveness of siIEL Vα3.2[+]QFL T to stimuli indicated that under physiological conditions, the activation of IELs was more complex and highly regulated as TCRs and inhibitory coreceptors such as CD8αα was engaged[43, 44]. It is not

surprising that the gut QFL T cells were functionally quiescent as these cells are constantly challenged by the microbial or dietary antigens from gut lumen[33]. Cross-reaction between QFL T cells and peptide expressed in the commensal bacterium *P. pentosaceus* further supports such phenomenon. Strikingly, monocolonization of the GF mice with *P. pentosaceus* was sufficient to restore gut Vα3.2$^+$QFL T cells which were lost in old GF mice. Cross-reaction of Vα3.2$^+$QFL T cells with the FL10 peptide expressed by *P. pentosaceus* might play a crucial role in the process as colonization of GF mice with a common lactate acid bacterium *L. johnsonni* which lack FL10 expression did not show a similar effect. Yet other signals generated from colonization of the commensal bacterium, such as secretion of bacteriocins, should not be excluded.

In conclusion, Vα3.2-expressing QFL T cells represent an unconventional population of Qa-1^b-restricted CD8$^+$ T cells which naturally reside at high frequency in small intestine epithelium. These cells showed memory phenotype in the spleen and characteristics of natIELs in gut. In addition, the fact that Vα3.2$^+$QFL T cross-reacts with FL9 homologue peptide expressed in commensal bacterium suggests that these cells might be functionally versatile besides elimination of ERAAP deficient cells. Further study of the biological significance of these cells could potentially reveal the delicate equilibrium between gut microenvironment and immune system.

Acknowledgements
We thank P. Rhee at Shastri Lab, H. Ding at the JHU germ-free core facility for technical support. We thank Dr. L. Coscoy at University of California, Berkeley for providing cellular biology expertise, Dr. X. Li at Ragon Institute of MGH, MIT and Harvard for providing insights on structural biology. This work was supported by grants from the National Institutes of Health (R01AI130210, R37AI060040).

Methods
Mice. SPF wild-type C57BL/6J, H-2T23-deficient Qa-1^b-KO, C57BL/6-Tg (TcraTcrb)1100Mjb/J OT-1, or C57BL/6-Tg (UBC-GFP)30Scha/J B6.GFP mice were obtained from the Jackson Laboratory. QFL T TCR transgenic mice QFLTg were generated in the laboratory of E. Robey (University of California Berkeley, Berkeley, CA) and housed in our

facility. GFP[+]QFLTg mice were generated in our facility by crossing B6.GFP mice with QFLTg mice. SPF QFLTg_WT, SPF OT-1_WT and GF QFLTg_WT partial hematopoietic chimera mice were generated as previously described[32]. GF wild-type C57BL/6J and GF QFLTg_WT chimera mice were generated and maintained in the Johns Hopkins Germ-free Mouse Core Facility. Mice were housed and all procedures were done in accordance with protocols approved by Animal Care and Use Committee of the Johns Hopkins University School of Medicine.

Generation of the QFLTg mouse. The QFLTg TCR α-and β-chain sequences were cloned and amplified from the genomic DNA of BEko8Z hybridoma[14, 15]. The TCR α-chain was amplified with the forward primer 5'-AAAACCCGGGCCAAGGCTCAGCCATGCTCCTGG-3' and the reverse primer 5'-AAAAGCGGCCGCATACAACATTGGACAAGGATCCAAGCTAAAGAGAACTC-3'. The TCR β-chain was cloned with the forward primer 5'-AAAACTCGAGCCCGTCTGGAGCCTGATTCCA-3' and the reverse primer 5'-AAAACCGCGGGGGACCCAGGAATTTGGGTGGA-3'. The TCR α-chain DNA fragment was cloned into pTα cassette vector by inserting it between the Xmal and NotI sites, while the TCR β-chain DNA fragment was cloned into pTβ cassette vector in between the XhoI and SacII sites[45]. The ampicillin resistance gene was removed from pTα and pTβ cassette by EarI enzyme digest. The QFLTg mice were generated on C57BL/6J background in the Cancer Research Laboratory Gene Targeting Facility at UC Berkeley under standard procedures. Founder mice were identified by flow cytometry and PCR genotyping of tail genomic DNA using primers mentioned above.

Antibodies, cell lines and peptides. Antibody for flow cytometry were from BioLegend (anti-CD45(30-F11), anti-CD19(1D3/CD19), anti-TCRβ(H57-597), anti-CD8β(53-5.8), anti-Vα3.2(RR3-16), anti-Vα2(B20.1), anti-CD90.2(53-2.1), anti-CD5(53-7.3), anti-IFN-ɣ(XMG1.2), anti-CD62L(MEL-14) and BD Biosciences (anti-CD4(RM4-5), anti-CD8α (53-6.7), anti-CD44(IM7)). BEko8Z, L-Qa-1[b] or Lmtk[-] cells were maintained as previously described[14]. Peptides were obtained from GenScript. The purity of SL8, FL9 and FL10 peptides were ⩾98%.

Generation of the QFL-dextramer. Qa-1[b]-FL9 (QFL) monomers were synthesized by the Tetramer Core Facility of the US National Institutes of Health. Phycoerythrin (PE)- or allophycocyanin (APC)-conjugated streptavidin were obtained from Agilent. The QFL dextramers were generated following the 'dextran doping' technique[27]. The QFL monomers were incubated with PE or APC conjugated streptavidin at a molar ratio of 3:1 at 4°C for 10min followed by addition of biotinylated dextran molecular weight 500kDa at a molar ratio of 1:20 with respect to streptavidin. The mixture was further incubated at 4°C for $\geqq$1h before being used in experiments.

Isolation of IELs. Small intestinal IELs were isolated following an established protocol with minor modifications [46]. In brief, the small intestine with Payer's patches removed were cut into appropriate length. The fecal content and mucus were removed by expelling with the flat side of forceps followed by flushing with PBS. The intestine was cut open longitudinally to reveal the epithelium and further cut laterally into ~2cm pieces. The intestine pieces were then placed in 25ml warmed dithioerythritol (DTE) solution (Ca^{2+}- and Mg^{2+}-free Hanks balanced salt solution, HEPES-bicarbonate buffer, 10% FCS) in 50ml conical tube and were shaken at 75rpm 37°C for 20min. The tube was vortexed for 10s before the supernatant was transferred into a new 50ml conical tube through a 70µM cell strainer. The cells were palleted and resuspend in 44% Percoll solution (44% Percoll in RPMI 1640). Density gradient was generated by underlaying the cell suspension with 67% Percoll solution (67% Percoll in RPMI 1640). The gradient cell suspension was further centrifuged at 1600 × g for 20 min at RT without using the brake. The layer of cells at the 44% and 67% interphase were thus collected as IELs.

Enrichment for dextramer-positive cells. Splenic dextramer-positive cells were enriched as previously described[14]. IELs were resuspend in 100µl sorter buffer (0.1% sodium azide and 5% FCS in PBS). PE or APC labeled QFL dextramers were added at a final dilution of 1:100. Cells were incubated at room temperature for 45 min, then were washed twice with 3ml of sorter buffer. 20µl of anti-PE and 20µl of anti-APC microbeads (Miltenyi Biotec) were added into cells resuspended in 450µl of sorters buffer, followed by incubation of 20 min at 4°C. Cells were washed twice with 3ml sorter buffer. The PE- and APC-labeled cells were positively selected by

passing through LS magnetic columns (Miltenyi Biotec). The entire isolated population was stained with anti-CD45, anti-CD19, anti-TCRβ, anti-CD4, anti-CD8α and anti-CD8β. QFL T cells were gated as CD45$^+$CD19$^-$TCRβ$^+$CD4$^-$QFL-Dex-PE$^+$QFL-Dex-APC$^+$ population. The absolute numbers and frequencies of the cells were calculated based on a fixed number of CountBright Beads (Thermo Fisher) added into each sample.

Cytokine stimulation and proliferation assay. Splenocytes or small intestinal IELs isolated from QFLTg or OT-1 mice were depleted of non-T cells using Pan T isolation Kit II (Miltenyi Biotec). The negatively selected T cells were then labeled with 5µM Cell Tracing Violet (CTV) (Thermo Fisher) and adjusted to the concentration of 10^6/ml. 10^6 of the CTV labeled T cells were stimulated *in vitro* for 3 or 5 days in cultures containing 100ng/ml IL15 (BioLegend) or a combination of 100ng/ml IL7 (BioLegend) and 100ng/ml IL18 (BioLegend) in 24-well plate at 37°C. Proliferation of cells were measured by dilution of CTV on flow cytometer.

CTL assay. Lymphocytes isolated from the spleen or siIEL compartment of naïve QFLTg_WT or OT-1_WT chimera mice were cultured *in vitro* with 2µM of FL9 or SL8 peptide respectively together with GolgiPlug (BD Biosciences) for 4.5h. WT splenocytes were supplemented into siIEL cultures as APCs. Cells were then washed and stained for surface markers. For measurement of IFN-ɣ production, cells were fixed and permeabilized using Fixation/Permeabilization Kit (BD Biosciences) and stained for intracellular IFN-ɣ.

Hybridoma assay. 10^5 of BEko8Z hybridoma cells were cocultured with 10^5 of L-Qa-1^b or Lmtk- cells in medium containing peptides of indicated concentrations in 96-well plates at 37°C for 12 hours. T cell response was determined by cleavage of the chromogenic LacZ substrate chlorophenol red-β-D-galactopyranoside (CPRG, Millipore Sigma) by TCR-induced LacZ and measure by spectrophotometry as presented by absorbance at 595 nm (A595) [47]. Background signals were subtracted by measurement of absorbance at 655nm.

Commensal culture and colonization. *Pediococcus pentosaceus Mees*(33314) and *Lactobacillus johnsonii*(33200) were obtained from ATCC. The bacteria were grown overnight at 37°C in 416 Lactobacilli MRS Broth (BD Biosciences). GF QFLTg_WT chimera mice were

gavage once with 200µl of $2x10^8$/ml CFU of *P. pentosaceus Mees* at 8 weeks of age. GF WT mice were gavage twice with 200µl of $2x10^8$/ml CFU of *P. pentosaceus Mees* or *L. johnsonii* at 8 and 10 weeks of age. Fecal pallets collected from the colonized mice were homogenized in 416 Lactobacilli MRS Broth, and the associated mice were determined by plating the appropriate dilution of homogenate on 416 Lactobacilli MRS agar plates.

1.   Shastri, N., Schwab, S. & Serwold, T. Producing nature's gene-chips: the generation of peptides for display by MHC class I molecules. *Annu Rev Immunol* **20**, 463-493 (2002).

2.   Blum, J.S., Wearsch, P.A. & Cresswell, P. Pathways of antigen processing. *Annu Rev Immunol* **31**, 443-473 (2013).

3.   Shastri, N., Cardinaud, S., Schwab, S.R., Serwold, T. & Kunisawa, J. All the peptides that fit: the beginning, the middle, and the end of the MHC class I antigen-processing pathway. *Immunol Rev* **207**, 31-41 (2005).

4.   Serwold, T., Gonzalez, F., Kim, J., Jacob, R. & Shastri, N. ERAAP customizes peptides for MHC class I molecules in the endoplasmic reticulum. *Nature* **419**, 480-483 (2002).

5.   Van Kaer, L. *et al.* Altered peptidase and viral-specific T cell response in LMP2 mutant mice. *Immunity* **1**, 533-541 (1994).

6.   Grandea, A.G., 3rd *et al.* Impaired assembly yet normal trafficking of MHC class I molecules in Tapasin mutant mice. *Immunity* **13**, 213-222 (2000).

7.   Van Kaer, L., Ashton-Rickardt, P.G., Ploegh, H.L. & Tonegawa, S. TAP1 mutant mice are deficient in antigen presentation, surface class I molecules, and CD4-8+ T cells. *Cell* **71**, 1205-1214 (1992).

8.   Blanchard, N. & Shastri, N. Coping with loss of perfection in the MHC class I peptide repertoire. *Curr Opin Immunol* **20**, 82-88 (2008).

9.   Hammer, G.E., Kanaseki, T. & Shastri, N. The final touches make perfect the peptide-MHC class I repertoire. *Immunity* **26**, 397-406 (2007).

10.   Guan, J., Peske, J.D., Taylor, J.A. & Shastri, N. The nonclassical immune surveillance for ERAAP function. *Curr Opin Immunol* **70**, 105-111 (2021).

11.   Hammer, G.E., Gonzalez, F., Champsaur, M., Cado, D. & Shastri, N. The aminopeptidase ERAAP shapes the peptide repertoire displayed by major histocompatibility complex class I molecules. *Nat Immunol* **7**, 103-112 (2006).

12.   Hammer, G.E., Gonzalez, F., James, E., Nolla, H. & Shastri, N. In the absence of aminopeptidase ERAAP, MHC class I molecules present many unstable and highly immunogenic peptides. *Nat Immunol* **8**, 101-108 (2007).

13.   Nagarajan, N.A. *et al.* ERAAP Shapes the Peptidome Associated with Classical and Nonclassical MHC Class I Molecules. *J Immunol* **197**, 1035-1043 (2016).

14.   Nagarajan, N.A., Gonzalez, F. & Shastri, N. Nonclassical MHC class Ib-restricted cytotoxic T cells monitor antigen processing in the endoplasmic reticulum. *Nat Immunol* **13**, 579-586 (2012).

15.   Guan, J., Yang, S.J., Gonzalez, F., Yin, Y. & Shastri, N. Antigen Processing in the Endoplasmic Reticulum Is Monitored by Semi-Invariant αβ TCRs Specific for a Conserved Peptide-Qa-1b MHC Class Ib Ligand. *J Immunol* **198**, 2017-2027 (2017).

16.   Godfrey, D.I., Uldrich, A.P., McCluskey, J., Rossjohn, J. & Moody, D.B. The burgeoning family of unconventional T cells. *Nat Immunol* **16**, 1114-1123 (2015).

17.   Salio, M., Silk, J.D., Jones, E.Y. & Cerundolo, V. Biology of CD1- and MR1-restricted T cells. *Annu Rev Immunol* **32**, 323-366 (2014).

18.   Faria, A.M.C., Reis, B.S. & Mucida, D. Tissue adaptation: Implications for gut immunity and tolerance. *J Exp Med* **214**, 1211-1226 (2017).

19.   Cheroutre, H., Lambolez, F. & Mucida, D. The light and dark sides of intestinal intraepithelial lymphocytes. *Nat Rev Immunol* **11**, 445-456 (2011).

20.   Das, G. & Janeway, C.A., Jr. MHC specificity of iIELs. *Trends Immunol* **24**, 88-93 (2003).

21.   Das, G. *et al.* Qa-2-dependent selection of CD8αα T cell receptor αβ+ cells in murine intestinal intraepithelial lymphocytes. *J Exp Med* **192**, 1521-1528 (2000).

22.   Das, G. & Janeway, C.A., Jr. Development of CD8αα and CD8αβ T cells in major histocompatibility complex class I-deficient mice. *J Exp Med* **190**, 881-884 (1999).

23.   Park, S.H. *et al.* Selection and expansion of CD8αα+ T cell receptor αβ+ intestinal intraepithelial lymphocytes in the absence of both classical major histocompatibility complex class I and nonclassical CD1 molecules. *J Exp Med* **190**, 885-890 (1999).

24.   Ruscher, R., Kummer, R.L., Lee, Y.J., Jameson, S.C. & Hogquist, K.A. CD8αα intraepithelial lymphocytes arise from two main thymic precursors. *Nat Immunol* **18**, 771-779 (2017).

25.   Round, J.L. & Mazmanian, S.K. The gut microbiota shapes intestinal immune responses during health and disease. *Nat Rev Immunol* **9**, 313-323 (2009).

26.   Mota-Santos, T. *et al.* Divergency in the specificity of the induction and maintenance of neonatal suppression. *Eur J Immunol* **20**, 1717-1721 (1990).

27.   Bethune, M.T., Comin-Anduix, B., Fu, Y.-H.H., Ribas, A. & Baltimore, D. Preparation of peptide–MHC and T-cell receptor dextramers by biotinylated dextran doping. *BioTechniques* **62**, 123-130 (2017).

28.   Prasad, M. *et al.* Expansion of an Unusual Virtual Memory CD8+ Subpopulation Bearing Valpha3.2 TCR in Themis-Deficient Mice. *Front Immunol* **12**, 644483 (2021).

29.   Schluns, K.S. & Lefrancois, L. Cytokine control of memory T-cell development and survival. *Nat Rev Immunol* **3**, 269-279 (2003).

30.   Okazawa, A. *et al.* Human intestinal epithelial cell-derived interleukin (IL)-18, along with IL-2, IL-7 and IL-15, is a potent synergistic factor for the proliferation of intraepithelial lymphocytes. *Clin Exp Immunol* **136**, 269-276 (2004).

31.   Klose, C.S. *et al.* The transcription factor T-bet is induced by IL-15 and thymic agonist selection and controls CD8αα+ intraepithelial lymphocyte development. *Immunity* **41**, 230-243 (2014).

32.   Ladi, E., Herzmark, P. & Robey, E. In situ imaging of the mouse thymus using 2-photon microscopy. *J Vis Exp* (2008).

33.   Denning, T.L. *et al.* Mouse TCRαβ+CD8αα intraepithelial lymphocytes express genes that down-regulate their antigen reactivity and suppress immune responses. *J Immunol* **178**, 4230-4239 (2007).

34.   Macpherson, A.J. & Harris, N.L. Interactions between commensal intestinal bacteria and the immune system. *Nature Reviews Immunology* **4**, 478-485 (2004).

35.   Belkaid, Y. & Hand, T.W. Role of the microbiota in immunity and inflammation. *Cell* **157**, 121-141 (2014).

36.   Donaldson, G.P., Lee, S.M. & Mazmanian, S.K. Gut biogeography of the bacterial microbiota. *Nat Rev Microbiol* **14**, 20-32 (2016).

37.   Jameson, S.C., Lee, Y.J. & Hogquist, K.A. Innate memory T cells. *Adv Immunol* **126**, 173-213 (2015).

38.   Sprent, J. & Surh, C.D. Normal T cell homeostasis: the conversion of naive cells into memory-phenotype cells. *Nat Immunol* **12**, 478-484 (2011).

39.   Mayans, S. *et al.* αβT Cell Receptors Expressed by CD4–CD8αβ– Intraepithelial T Cells Drive Their Fate into a Unique Lineage with Unusual MHC Reactivities. *Immunity* **41**, 207-218 (2014).

40.   Roberts, A.I., O'Connell, S.M., Biancone, L., Brolin, R.E. & Ebert, E.C. Spontaneous cytotoxicity of intestinal intraepithelial lymphocytes: clues to the mechanism. *Clin Exp Immunol* **94**, 527-532 (1993).

41.   Carol, M. *et al.* Spontaneous secretion of interferon gamma and interleukin 4 by human intraepithelial and lamina propria gut lymphocytes. *Gut* **42**, 643-649 (1998).

42.   Van Kaer, L. *et al.* CD8αα+ innate-type lymphocytes in the intestinal epithelium mediate mucosal immunity. *Immunity* **41**, 451-464 (2014).

43.   Madakamutil, L.T. *et al.* CD8αα-mediated survival and differentiation of CD8 memory T cell precursors. *Science* **304**, 590-593 (2004).

44.   Cheroutre, H. & Lambolez, F. Doubting the TCR coreceptor function of CD8αα. *Immunity* **28**, 149-159 (2008).

45.   Kouskoff, V., Signorelli, K., Benoist, C. & Mathis, D. Cassette vectors directing expression of T cell receptor genes in transgenic mice. *J Immunol Methods* **180**, 273-280 (1995).

46.   Qiu, Z. & Sheridan, B.S. Isolating Lymphocytes from the Mouse Small Intestinal Immune System. *J Vis Exp* (2018).

47.   Sanderson, S. & Shastri, N. LacZ inducible, antigen/MHC-specific T cell hybrids. *Int Immunol* **6**, 369-376 (1994).

Figure 1 Abundant Vα3.2⁺ QFL T cells in both the spleen and siIEL compartment of naïve wild-type mice. (**a**) Definition of QFL T cells by flow cytometry. Cells from the spleen or small intestinal IEL (siIEL) compartment were stained with Qa-1^b-FL9 (QFL) dextramers labeled with phycoerythrin (QFL-Dex-PE) or allophycocyanin (QFL-Dex-APC) and analyzed for before (Unenriched) or after (Enriched) magnetic enrichment of dextramer-positive cells. QFL T cells are defined as the CD45⁺CD19⁻TCRβ⁺CD4⁻QFL-Dex-PE⁺APC⁺ population. Plots representing siIELs from naïve wild-type(WT) mice after enrichment for QFL-Dex⁺ cells. (**b**) Flow cytometry of cells from the spleen (Sp) and siIEL compartment of naïve WT mice 'Unenriched' or 'Enriched' for dextramer-positive cells. Numbers in plots indicate absolute numbers of QFL-Dex⁺ cells detected after enrichment. (**c**) Absolute numbers (left) and frequencies (right) of QFL-Dex⁺ cells detected (as in **b**) among TCRβ⁺CD4⁻ population in the spleen or siIEL compartment. **P=0.0069 (**d**) Analysis of Vα3.2 expression on QFL-Dex⁺(QFL) or total CD8⁺ T (Total) cells in the spleen. Numbers in plots indicate average percentages of Vα3.2⁺ cells within the indicated populations. (**e**) Frequencies of Vα3.2⁺ cells detected as in **d.** ****P<0.0001 (**f**) Analysis of Vα3.2 expression on the QFL-Dex⁺(QFL) or total CD8⁺ T (Total) cells in the siIEL compartment. Numbers in plots indicate average percentages of Vα3.2⁺ cells within the indicated populations. (**g**) Frequencies of Vα3.2⁺ cells detected as in **f.** **P=0.0058 Representative data are shown in flow plots(**a,b,d,f**) and the number of replicates is specified in the bar graphs(**c,e,g**). Each symbol represents data collected from the indicated tissue isolated from an individual

mouse. P values were calculated with Student's *t* test. 'ns' indicates the comparison was not significant.

Figure 2 Unconventional phenotype of Vα3.2$^+$ QFL T cells. (**a**) Analysis of CD44 expression on Vα3.2$^+$ or Vα3.2$^-$ QFL-Dex$^+$ cells enriched from the spleen of naïve WT mice. Numbers in plots indicate average percentages of CD44^{hi} cells detected among the Vα3.2$^+$ or Vα3.2$^-$ QFL-Dex$^+$ populations. (**b**) Frequencies of CD44^{hi} cells detected as in **a**.****P<0.0001 (**c**) Analysis of CD8α and CD8β expression on Vα3.2$^+$ or Vα3.2$^-$ QFL-Dex$^+$ cells enriched from the siIEL compartment of naïve WT mice. Numbers in plots indicate average percentages of CD8αα$^+$ cells detected among the Vα3.2$^+$ or Vα3.2$^-$ QFL-Dex$^+$ populations. (**d**) Frequencies of CD8αα$^+$ cells detected as in **a**. **P=0.0065 Samples of <20 cells in the Vα3.2$^+$QFL-Dex$^+$ gate were excluded in the phenotype analysis. Representative data are shown in flow plots(**a,d**) and the number of replicates is specified in the bar graphs(**b,c,e**). Each symbol represents data collected from the indicated tissue isolated from an individual mouse. P values were calculated with Student's *t* test or Mann-Whitney test depending on the normality of data sets. 'ns' indicates the comparison was not significant.

Figure 3.

Figure 3 Phenotype of Vα3.2⁺ QFL T cells in mice of various genotypes. (**a**) Absolute numbers of Vα3.2⁺QFL-Dex⁺ cells detected in the spleen (left) or siIEL compartment (right) of Qa-1^b, ERAAP or TAP deficient (Qa-1^b-KO, ERAAP-KO or TAP-KO) mice in comparison with WT mice.** P<0.009 ***P=0.0006 Symbols on x-axis indicate that the cells were undetectable in TAP-KO mice. (**b**) Analysis of CD44 expression on splenic Vα3.2⁺QFL-Dex⁺ cells enriched from naïve WT, Qa-1^b-KO or ERAAP-KO mice. Numbers in plots indicate average percentages of CD44^{hi} cells. (**c**) Frequencies of CD44^{hi} cells detected among Vα3.2⁺(as in **d**) or Vα3.2⁻ QFL T cells.***P=0.0004*P=0.0388 (**d**) Flow cytometry analysis of CD8α and CD8β expression on the Vα3.2⁺QFL-Dex⁺ cells enriched from the siIEL compartment of naïve WT, Qa-1^b-KO or

ERAAP-KO mice. Numbers in plots indicate average percentages of CD8αα$^+$ cells. (**e**) Percentages of CD8αα$^+$ cells detected among Vα3.2$^+$(as in **d**) or Vα3.2$^-$ QFL T cells. **P=0.0082 *P=0.0335 Representative data are shown in flow plots (**b,d**) and the number of replicates is specified in the bar graphs (**a,c,e**). Each symbol represents data collected from the indicated tissue isolated from an individual mouse. P values were calculated with Student's *t* test or Mann-Whitney test depending on the normality of data sets. 'ns' indicates the comparison was not significant.

Figure 4 Distinct functional features of QFLTg cells in the spleen and gut. (**a**) Proliferation of splenic QFLTg or OT-1 cells from QFLTg or OT-1 mice in response to IL15 (left) or a combination of IL7 and IL18 (right) stimulation. Cells were tracked using Cell Tracing Violate (CTV). Numbers in plots indicate average percentages of proliferated splenic QFLTg cells. (**b**) Percentages of proliferated splenic QFLTg or OT-1 cells as detected in **a**. ****P<0.0001 **P=0.0052 (**c**) Proliferation of siIEL QFLTg or OT-1 cells from QFLTg or OT-1 mice in response to IL15 (left) or a combination of IL7 and IL18 (right) stimulation. Cells were tracked using Cell Tracing Violate (CTV). Numbers in plots indicate average percentages of proliferated

QFLTg cells from siIEL compartment. (**d**) Percentages of proliferated siIEL QFLTg or OT-1 as detected in **c**. ***P=0.0009 *P=0.0101 (**e**) Generation of QFLTg_WT or OT-1_WT partial hematopoietic chimera mice. 2x10⁶ of the bone marrow (BM) cells from GFP⁺QFLTg or GFP⁺OT-1 mice were transferred into WT neonates at 3~5 days of age. The chimera mice were analyzed at 8~10 weeks of age. (**f**) Gating strategy for the QFLTg cells originated from donor bone marrow cells in QFLTg_WT chimera mice. Plots representing donor-derived QFLTg population in the spleen being gated as GFP⁺CD45⁺CD19⁻TCRβ⁺CD4⁻Vα3.2⁺ cells. (**g**) Flow cytometry measurement of IFN-ɣ production by QFLTg or OT-1 cells isolated from the spleen of naïve chimera mice stimulated with or without 2μM FL9 or SL8 peptide respectively for 4.5h. Numbers in plots indicate average percentages of IFN-ɣ⁺ cells. (**h**) Percentages of IFN-ɣ⁺ cells detected as in **g**. ****P<0.0001 ***P=0.0001 (**i**) Flow cytometry measurement of IFN-ɣ production by QFLTg or OT-1 cells isolated from the siIEL compartment of naïve chimera mice stimulated with or without 2μM FL9 or SL8 peptide respectively for 4.5h. Numbers in plots indicate average percentages of IFN-ɣ⁺ cells. (**j**) Percentages of IFN-ɣ⁺ cells detected as in **i**. ***P=0.0004 *P=0.0247 Representative data are shown in flow plots (**a,c,f,g,i**) and the number of replicates is specified in the bar graphs (**b,d,h,j**). Each symbol represents data collected from the indicated tissue isolated from an individual mouse. P values were calculated with Student's *t* test. 'ns' indicates the comparison was not significant.

Figure 5 Gut microbiota is associated with retention but not homing of QFL T cells in the siIEL compartment. (**a**) Flow cytometry of QFL-Dex⁺ cells enriched from the spleen of specific-pathogen-free (SPF) or germ-free (GF) WT mice at 10 or 18 weeks of age. Numbers in plots indicate absolute numbers of QFL-Dex⁺ cells. (**b**) Absolute numbers(left) and frequencies of QFL-Dex⁺ cells detected within the total splenic CD8⁺ population in SPF or GF WT mice of 8~17 weeks or 18~22 weeks of age.***P=0.009 ****P<0.0001 (**c**) Flow cytometry of QFL-Dex⁺

cells enriched from the siIEL compartment of SPF or GF WT mice at 10 or 18 weeks of age. Numbers in plots indicate absolute numbers of QFL-Dex+ cells. (**d**) Absolute numbers and frequencies of QFL-Dex+ cells detected among the total siIEL TCRβ+CD4- population in SPF or GF WT mice of 8~17 weeks or 18~22 weeks of age. **P=0.005 *P=0.037 Representative data are shown in flow plots (**a,c**) and the number of replicates is specified in the bar graphs (**b,d**). Each symbol represents data collected from the indicated tissue isolated from an individual mouse. P values were calculated with Student's *t* test. 'ns' indicates the comparison was not significant.

Figure 6

Figure 6 Association between Vα3.2+QFL T cells and the commensal bacterium *P. pentosaceus*. (**a**) Identification of the QFL T cross-reactive peptide FYAEDGTPIL(FL10). LacZ response of BEko8Z hybridoma cells to the Qa-1b-expressing Lmtk- (L-Qa-1b) cells or the non-Qa-1b-expressing Lmtk- cells pulsed with 50nM of the candidate FL9 or its homolog peptides measured by spectrophotometry. (**b**) Response of BEko8Z hybridoma to L-Qa-1b cells pulsed

with FL9, FL10 or FYAEDDTPIV (FV10) peptide of various concentrations. (**c**) Colonization of GF mice with individual commensal bacterium. GF QFLTg_WT chimera mice received oral gavage of *P. pentosaceus* at 8 weeks of age and were analyzed for both the donor-derived QFLTg cells and the endogenous Vα3.2$^+$QFL T cells at 16 weeks of age. GF WT mice received oral gavage of *P. pentosaceus* or *L. johnsonii* at 8 and 10 weeks of age and were analyzed for the endogenous Vα3.2$^+$QFL T cells at 16 weeks of age. (**d**) Absolute numbers of donor-derived QFLTg cells in the spleen (left) or siIEL compartment (right) of 16-week GF QFLTg_WT chimera mice colonized with *P. pentosaceus* in comparison with SPF or uncolonized GF chimera mice. Symbols on x-axis indicate that QFLTg cells were undetectable in the indicated group. *P<0.03 (**e**) Absolute numbers of endogenous Vα3.2$^+$QFL T cells in the spleen (left) or siIEL compartment (right) of 16-week colonized GF mice compared with SPF or uncolonized GF chimera mice. Endogenous Vα3.2$^+$QFL T data collected from the GF QFLTg_WT chimeric mice and the non-chimeric WT mice were pooled for this analysis. **P<0.008 (**a,b**) Data are from one experiment representative of three experiments. (**d,e**) The number of replicates is specified in the bar graphs with each symbol representing data collected from the indicated tissue isolated from an individual mouse. P values were calculated with Mann-Whitney test. 'ns' indicates the comparison was not significant.