1 **Commensal Bacteria Maintain a Qa-1^b-restricted Unconventional CD8$^+$ T Population in**

2 **Gut Epithelium**

3

4 Jian Guan[1,2*], J. David Peske[1,2], Michael Manoharan Valerio[3], Chansu Park[1,2], Ellen A. Robey[3,4],

5 Scheherazade Sadegh-Nasseri[1,4,*]

6

7 [1] Department of Pathology, Johns Hopkins University School of Medicine, Baltimore, MD

8 21287, USA

9

10 [2] Institute of Cell Engineering, Johns Hopkins University School of Medicine, Baltimore, MD

11 21287, USA

12

13 [3] Division of Immunology and Molecular Medicine, Department of Molecular and Cell Biology,

14 University of California, Berkeley, Berkeley, CA 94720, USA

15

16 [4] Co-senior authors

17

18 *Lead contacts: jian.guan.cc@outlook.com (J.G) and ssadegh@jhmi.edu (SS-N)

19 This paper is dedicated to the memory of Dr. Nilabh Shastri who laid foundation for the studies

20 on ERAP1 immune surveillance and QFL T cells. He had offered his insights and reviewed the

21 bulk of the data presented in this study, but unfortunately passed away before finalization of the

22 work.

23
24

Abstract

Intestinal intraepithelial lymphocytes (IELs) are characterized by an unusual phenotype and

developmental pathway, yet their specific ligands and functions remain largely unknown. Here

by analysis of QFL T cells, a population of CD8$^+$ T cells critical for monitoring the MHC I

antigen processing pathway, we established that unconventional Qa-1^b-restricted CD8$^+$ T cells

are abundant in intestinal epithelium. We found that QFL T cells showed a Qa-1^b-dependent

unconventional phenotype in the spleen and small intestine of naïve wild-type mice. The splenic

QFL T cells showed innate-like functionality exemplified by rapid response to cytokines or

antigen, while the gut population was refractory to stimuli. Microbiota was required for the

maintenance, but not the initial gut homing of QFL T cells. Moreover, monocolonization with

Pediococcus pentosaceus, which expresses a peptide that cross-activated QFL T cells, was

sufficient to maintain QFL T cells in the intestine. Thus, microbiota is critical for shaping the

Qa-1^b-restricted IEL landscape.

39
40

Introduction

The display of peptides by MHC class I (MHC I) molecules on the cell surface is critical for

CD8$^+$ T cell immune surveillance (***Shastri et al., 2002***). Generation of a peptide repertoire which

accurately reflects intracellular events, such as viral infection or mutations, relies on a functional

antigen processing and presentation pathway. After cytosolic cleavage of protein precursors and

transport of peptides through the peptide transporter associated with antigen processing (TAP)

into the endoplasmic reticulum (ER), the peptide intermediates are further customized by ER

aminopeptidase associated with antigen processing (ERAP1) until 'ideal' peptides that fit the

groove of MHC I molecules are eventually shuttled and displayed on the cell surface (***Serwold et***

al., 2002; Shastri et al., 2005). Disruption of each step of the pathway can lead to

immunological dysfunction (***Grandea et al., 2000; Van Kaer et al., 1994; Van Kaer et al.,***

1992). The critical role of ERAP1 in the antigen processing pathway has been established

through extensive studies of ERAP1 deficient (ERAP1-KO) cells and mice (***Blanchard et al.,***

2008; Guan et al., 2021; Hammer et al., 2007b). Loss of ERAP1 severely disrupts the peptide

repertoire presented by both the classical MHC Ia and the nonclassical MHC Ib molecules

(*Hammer et al., 2006; Hammer et al., 2007a*). Qa-1^b, a nonclassical MHC Ib molecule, has been

shown to present a significantly increased number of peptides on the cell surface of ERAP1-KO

cells as compared with wild-type (WT) cells (*Nagarajan et al., 2016*). One peptide presented by

Qa-1^b, FYAEATPML (FL9) (with the Qa-1^b-FL9 complex termed QFL), was identified as an

immunodominant ligand uniquely presented on ERAP1-deficient cells (*Nagarajan et al., 2012*).

The CD8$^+$ T cells that specifically recognize this ligand are thus named QFL-specific T (QFL T)

cells.

Early analysis of QFL T cells revealed the unusual nature of these CD8$^+$ T cells. Unlike

conventional antigen-specific CD8$^+$ T cells which are typically detected at a frequency of 1 in

$10^5 \sim 10^6$, QFL T cells are present at a frequency 10-fold higher in the spleen of naïve mice with

the bulk splenic QFL T population displaying a CD44^{hi}CD122$^+$ antigen-experienced phenotype

(*Nagarajan et al., 2012*). T cell receptor (TCR) analysis of QFL T cells revealed that a large

proportion of the QFL T population expresses an invariant TCRα-chain Vα3.2Jα21 (*Guan et al.,*

2017). These traits of QFL T cells indicate their potential similarity to other unconventional T

cells, such as invariant NKT (iNKT) and mucosal associated invariant T(MAIT) cells, which are

typically characterized by recognition of ligands presented by non-classical MHC Ib, expression

of invariant TCRs, residence in nonlymphoid tissues and innate-like functions (*Godfrey et al.,*

2015; Salio et al., 2014). The tissue distribution and functions of QFL T cells remains to be fully

elucidated.

The gut mucosa is an immunologically complex niche with abundant lymphoid populations

(*Faria et al., 2017*). The small intestinal intraepithelial lymphocyte (siIEL) compartment is

populated by unconventional T cells of both TCRγδ$^+$ and TCRαβ$^+$ lineage which express CD8αα

but lack CD4, CD8αβ, CD5 and CD90 expression. The development of the CD8αα$^+$CD4$^-$CD8αβ$^-$

TCRαβ$^+$ population (CD8αα$^+$ IEL), categorized as natural IELs (natIELs) because they acquire

their activated phenotype in the thymus, has been extensively studied (*Cheroutre et al., 2011*).

Yet little is known about their antigen specificity and TCR repertoire. Emerging evidence shows

that nonclassical MHC Ib molecules are important for the development and effector function of

CD8αα$^+$ IELs (*Das et al., 2003*). For instance, while loss of classical MHC Ia molecules showed

little impact on the CD8αα$^+$ IEL population, these cells were decreased in mice deficient for Qa-

2 (***Das et al., 2000; Das et al., 1999; Park et al., 1999***). Furthermore, a population of CD8αα[+]

IEL precursors which preferentially expresses Vα3.2 was shown to be decreased in number in

the absence of CD1d (***Ruscher et al., 2017***). However, whether Qa-1[b] is involved in shaping the

CD8αα[+] IEL population remains to be studied. In addition to MHC molecules, gut microbiota

plays a critical role in the establishment and shaping of the gut immune system. Studies of germ-

free (GF) mice have shown extensive defects in gut-associated lymphoid tissues together with

morphological changes in the intestine associated with the absence of gut microbiota (***Round et***

al., 2009). Notably, despite the critical role of gut microbiota in the establishment of the gut

immune system, natIELs that do not rely on cognate antigens in the periphery are believed to

home to the gut independent of microbes (***Mota-Santos et al., 1990***) .

Here we found abundant unconventional QFL T cells in both the spleen and siIEL compartment

of naïve WT mice. The splenic Vα3.2[+]QFL T cells expressed high level of CD44 and showed

typical innate-like functions including hyperresponsiveness to cytokines and rapid IFN-ɣ

production in response to antigen. In contrast, the populations of the same antigen specificity in

the gut phenotypically resembled natIEL and were likewise functionally quiescent. Analysis of

mice deficient of molecules associated with Qa-1[b]-FL9 antigen presentation revealed that TAP

was required for the presence of the splenic or siIEL QFL T cells, whereas Qa-1[b] was essential

for imprinting their unconventional phenotype. Analysis of GF mice showed that gut microbiota

was needed for the long-term maintenance of QFL T cells. Furthermore, we found that

maintenance of the gut Vα3.2[+]QFL T was associated with colonization by the commensal

bacterium *Pediococcus pentosaceus* (*P. pentosaceus*) which expresses an FL9 homologue that

could cross-activate QFL T cells. Overall, these results establish Vα3.2[+]QFL T cells as a unique

population of unconventional CD8[+] T cells that rely on nonclassical MHC Ib to acquire their

phenotype, and gut microbiota to be properly maintained in the intestine.

Results

Vα3.2[+]QFL T cells are abundant in the spleen and gut of naïve wild-type mice

To identify QFL T cells in tissues of naïve WT mice, we generated Qa-1[b]-FL9 dextramers (QFL-

Dex) based on the 'dextran-doping' technique (***Bethune et al., 2017***). Prior work of our lab

showed that unlike the Qa-1^b-Qdm tetramer which stains NK cells, the Qa-1^b-FL9 tetramer

(QFL-Tet) does not bind to the NKG2A receptor but specifically stains CD8$^+$ T cells (***Nagarajan***

et al., 2012). Because the same QFL monomers were used for generating the QFL-multimers in

this study, we reasoned that it is unlikely that the QFL-Dex will exhibit shifted specificity.

Moreover, in comparison with QFL-Tet, QFL-Dex showed improved sensitivity and specificity

for QFL T cell detection (**Supplementary Fig. 1**), and further allowed enrichment of QFL T

cells using magnetic beads. The average number of QFL T cells – defined as CD45$^+$CD19$^-$

TCRβ$^+$CD4$^-$QFL-Dex-PE$^+$APC$^+$ (**Fig.1 a**) – detected in the siIEL compartment of naïve WT

mice was comparable to the population in the spleen. Additionally, QFL T cells were present at a

frequency of ~1 in 1,000 of the siIEL CD8$^+$ T cells, which was 10 times more frequent than in

the spleen (**Fig.1 b, c**). Both the splenic and siIEL QFL T cells were essentially all CD8$^+$ T cells,

as these cells were barely detectable within the CD4$^+$ T population (**Supplementary Fig. 2**).

Consistent with prior studies, ~80% of the splenic QFL T cells expressed the 'invariant' TCR Vα

segment Vα3.2 which was significantly higher than the proportion of Vα3.2-expressing cells

within the total CD8$^+$ T population in the spleen (<5%) (**Fig.1 d, e**) (***Guan et al., 2017***). Notably,

QFL T cells in the siIEL compartment also contained a relatively larger proportion of Vα3.2$^+$

cells (20.1%) compared to the total CD8$^+$ TCRαβ siIELs (<10%) (**Fig.1 f, g**). The average

percentage of Vα3.2$^+$ cells within the QFL T population (20.1%) was lower in the siIEL

compartment than in the spleen (63.3%). Vα3.2$^+$ QFL T cells were nevertheless abundantly

present in both the spleen and gut of naïve WT mice.

The unconventional phenotype of Vα3.2$^+$QFL T cells

We have previously observed a high percentage of CD44^{hi} cells within the bulk splenic QFL T

population of naïve WT mice(***Guan et al., 2017; Nagarajan et al., 2012***). Here we further

investigated the association between preferred Vα3.2 usage by QFL T cells and their phenotype.

We observed that Vα3.2$^+$QFL T cells contained a significantly larger proportion of CD44^{hi} cells

(~90%) than the Vα3.2$^-$QFL T cells, suggesting that the antigen experienced phenotype of the

QFL T population was mainly contributed by the Vα3.2-expressing subpopulation (**Fig.2 a, b**).

In contrast, only around 20% of the total Vα3.2$^+$CD8$^+$ T population expressed high CD44. It is

worth noting that the percentage of CD44^{hi} cells within the total Vα3.2$^+$CD8$^+$ T population was

slightly higher compared to the Vα3.2$^-$ population (**Supplementary Fig. 3 a**), which was in line

with a previous report (***Prasad et al., 2021***). We thus conclude that although both Vα3.2$^+$ and

Vα3.2$^-$ cells were detected within the splenic QFL T population, a memory phenotype was

specifically enriched in Vα3.2$^+$QFL T cells.

Given the high frequency of QFL T cells in the gut, we further investigated their phenotype. We

assessed CD8αα expression, a hallmark of the unconventional phenotype of natIELs, on Vα3.2$^+$

and Vα3.2$^-$ QFL T cells in the siIEL compartment of naïve WT mice. A relatively larger fraction

of Vα3.2$^+$QFL T cells lacked CD4 and CD8αβ expression but expressed CD8αα compared to the

Vα3.2$^-$ subpopulation suggesting that the majority of Vα3.2$^+$ QFL T cells were CD8αα$^+$ natIELs

(**Fig.2 c, d**). No such difference was observed between the total Vα3.2$^+$ and Vα3.2$^-$ CD8$^+$ IEL

populations (**Supplementary Fig. 3 b**). These observations indicate that the Vα3.2$^+$ QFL T cells

in both the spleen and siIEL compartment are unconventional CD8$^+$ T cells which likely are

derived from the same precursors in thymus.

The impact of Qa-1^b, ERAP1, and TAP on Vα3.2$^+$QFL T cells

We further investigated what drives the unconventional phenotype of Vα3.2$^+$ QFL T cells by

analyzing mice deficient of molecules that impact Qa-1^b-FL9 antigen presentation including Qa-

1^b, ERAP1 or TAP (Qa-1^b-KO, ERAP1-KO, TAP-KO). While we consistently observed loss of

both the splenic and siIEL Vα3.2$^+$QFL T cells in TAP-KO mice, the populations were detectable

in both tissues of Qa-1^b-KO or ERAP1-KO mice with a slightly reduced population size in the

spleen of Qa-1^b-KO mice (**Fig.3 a**). Vα3.2$^+$ QFL T populations showed reduced percentages of

CD44^{hi} cells in the spleen of Qa-1^b and ERAP1 deficient mice with the reduction being more

significant in Qa-1^b-KO mice. In contrast, the phenotype of splenic Vα3.2$^-$ QFL T cells was not

significantly affected (**Fig.3 b, c**). Similar pattern of phenotype change was observed for the

siIEL Vα3.2$^+$ QFL T cells which showed almost complete loss of the unconventional natIEL

phenotype including CD8αα expression in Qa-1^b-KO mice (**Fig.3 d, e, Supplementary Fig. 4**).

These results demonstrate that TAP and Qa-1^b both play a role in the establishment of the

unconventional Vα3.2$^+$ QFL T population, with TAP being required for the presence of the

population, and Qa-1^b being required for the unconventional phenotype imprinting of these cells.

Paradoxically, loss of ERAP1, which would be expected to increase the presentation of FL9, also

led to a reduction of memory and unconventional phenotypes in QFL T cells, perhaps due to the

loss of certain QFL T clones as a result of tolerance mechanisms, such as negative selection. It is

also worth noting that total splenic Vα3.2$^+$CD8αβ$^+$ T cells showed a similar loss of memory

phenotype in Qa-1^b-KO and ERAP1-KO mice, whereas Vα3.2$^-$CD8αβ$^+$T cells did not

(**Supplementary Fig. 3 c**). Thus, the unconventional Vα3.2$^+$QFL T cells specifically, and a

substantial proportion of total Vα3.2$^+$ splenocytes, are impacted by Qa-1^b and ERAP1.

Vα3.2$^+$QFL T cells are functionally innate-like in the spleen but are quiescent in the gut

To further investigate the functional features of Vα3.2$^+$QFL T cells, we utilized transgenic mice

(QFLTg) expressing the predominant invariant TCR found on QFL T cells (*Vα3.2Jα21,*

Vβ1Dβ1Jβ2-7), referred to hereafter as QFLTg cells. Abundant QFLTg cells or control OT-1

cells were detected in both the spleen and siIEL compartment of the respective TCR transgenic

mice (**Supplementary Fig. 5 a**). Phenotypically, the QFLTg cells recapitulated the

unconventional phenotype of polyclonal Vα3.2$^+$QFL T cells with high CD44 expression in the

spleen and CD8αα expression in the siIEL compartment. In contrast, OT-1 cells were

phenotypically similar to conventional naïve CD8$^+$ T cells, as they expressed low to intermediate

levels of CD44 in the spleen and lacked CD8αα expression in gut (**Supplementary Fig. 5 b, c**).

Cytokines including IL-15 and the combination of IL-7/IL-18 have been implicated in the

maintenance and survival of both 'virtual' memory T cells and IELs (***Klose et al., 2014;***

Okazawa et al., 2004; Prasad et al., 2021; Schluns et al., 2003). We thus compared the

proliferation of QFLTg and OT-1 cells in response to these cytokines. In keeping with their

virtual memory surface marker phenotype, splenic QFLTg cells were significantly more

responsive to IL-15 or IL-7/IL-18 stimulation than OT-1 cells (**Fig.4 a**). At 72hs, ~80% of

QFLTg cells had proliferated in response to the cytokines, compared with only ~50% of OT-1

cells (**Fig.4 b**). In contrast, T cells isolated from the siIEL compartment of QFLTg or OT-1 mice

showed a distinct pattern of cytokine responsiveness from the splenic T cells. First, both QFLTg

and OT-1 cells from siIEL were less responsive to IL-15 than the splenic population. Second,

siIEL QFLTg cells showed significantly reduced proliferation in response to the cytokines,

especially to the combination of IL-7/IL-18, as compared with OT-1 cells (**Fig.4 c, d**).

To ensure that altered effector function was not skewed by the high specific TCR frequency in
TCR transgenics, we generated partial hematopoietic chimeric mice with physiological precursor
frequencies (QFLTg_WT) by transferring bone marrow cells from GFP$^+$QFLTg mice into
nonirradiated naïve WT neonates (***Ladi et al., 2008***). OT-1_WT chimera mice were generated in
parallel as representative of conventional CD8$^+$ T cells (**Fig.4 e**). This experimental design
allowed us to track the GFP$^+$ donor-derived T cells (GFP$^+$CD45$^+$CD19$^-$TCRβ$^+$CD4$^-$Vα3.2$^+$)
separately from the recipient's endogenous T cells (**Fig.4 f**). Based on the hyperresponsiveness
of splenic QFLTg cells to cytokines, we hypothesized that these cells would exert effector
functions more rapidly than conventional CD8 T cells, similar to the known innate-like T cells
such as iNKT. We measured IFN-ɣ production by QFLTg or OT-1 cells from splenocytes or
siIEL cells of naïve QFLTg_WT or OT-1_WT chimera mice stimulated the with 2µM FL9 or
SIINFEKL(SL8) peptide respectively for 4.5 hours directly *ex vivo*. A large fraction (~60%) of
splenic QFLTg cells expressed high levels of IFN-ɣ under these conditions, while only 6% of
OT-1 cells did so (**Fig.4 g, h**). Notably, despite the low responsiveness to epitopes, siIEL QFLTg
cells showed relatively higher basal levels of IFN-ɣ expression than OT-1 cells (**Fig.4 i, j**). In
contrast to the hyperresponsiveness of splenic QFLTg cells to their ligand, neither QFLTg nor
OT-1 cells from the siIEL compartment showed significant IFN-ɣ production at 4.5 hours. The
low IFN-ɣ production of IEL QFLTg cells was in keeping with their natIEL phenotype - antigen-
experienced yet functionally quiescent (***Cheroutre et al., 2011; Denning et al., 2007***). We
conclude that Vα3.2$^+$QFL T cells in the spleen are functionally innate-like, whereas the
population with the same TCR specificity in the small intestine shows functional features of
natIEL.
232
Gut microbiota is associated with retention but not homing of QFL T cells
The establishment and function of the intestinal immune system is intimately associated with
homeostasis of the gut microbial community, as evidenced by an altered gut immune cell
composition in germ-free (GF) mice (***Belkaid et al., 2014; Macpherson et al., 2004; Round and***
Mazmanian, 2009). To investigate if an association exists between gut microbiota and gut QFL
T cells, we compared the numbers and frequencies of QFL T cells in the spleen and siIEL
compartment of naïve specific-pathogen-free (SPF) and GF WT mice of various ages. First, we
found the absolute number of QFL T cells in the spleen was unaffected by the absence of gut

microbiota in either young or old mice. The frequency of QFL T cells was relatively higher in

GF than in SPF mice, as a result of a decreased number of non-QFL CD8$^+$ T cells in the spleen

of GF mice (**Fig.5 a, b**). However, the percentage of CD44^{hi} cells within the total CD8$^+$ T

population was proportionally higher in the spleen of GF WT than in SPF WT mice regardless of

their Vα3.2 expression (**Supplementary Fig. 6 a**). This observation suggested that gut

microbiota might be more closely associated with the presence of conventional naïve CD8$^+$ T

cells rather than the memory phenotype CD8$^+$ T population in the spleen. Second, while the

number and frequency of QFL T cells in siIEL compartment showed no significant difference

between GF and SPF mice of relatively young age (8~17-week), the population was gradually

lost in old GF mice (18~22-week) (**Fig.5 c, d**), with the decrease in GF mice starting from

around 16 weeks of age (data not shown). In addition, the QFL T cells detected in the spleen or

siIEL compartment of young SPF or GF mice showed similar expressions of Vα3.2, CD44 and

CD8αα (**Supplementary Fig. 7**). We conclude that although QFL T cells could home to gut in

the absence of microbiota, microbe-derived antigens or signals are required for their maintenance

there. Additionally, we observed that the average percentage of CD8αα$^+$ cells among the total

TCRβ$^+$CD4$^-$ T population was proportionally larger in GF mice than SPF mice regardless of

Vα3.2 expression, although the Vα3.2$^-$ population showed a more significant increase

(**Supplementary Fig. 6 b**). This was mainly a result of a decreased number of CD8αβ$^+$ T cells in

the siIEL of GF mice (data not shown). Nevertheless, here we showed that Qa-1^b restricted T

cells could home to the small intestine epithelium and display a natIEL phenotype independent

of gut microbiota.

QFL T cells cross-react with a microbial antigen

We next investigated how the QFL T population was retained in the gut of SPF WT mice,

hypothesizing that the presence of specific antigen might play a role. QFL T cells specifically

recognize the Qa-1^b-FL9 complex which is uniquely presented on ERAP1 deficient cells.

Because it is unlikely this ligand is constantly presented in gut epithelium under homeostatic

conditions (**Supplementary Fig. 8 a**), we reasoned that QFL T cells might be retained in the gut

through exposure to FL9 homologue peptide(s) expressed by commensal bacteria that colonize

small intestine.

To identify potentially cross-reactive peptides, we first determined the key residues of the

FYAEATPML(FL9) peptide that could potentially affect activation of QFL T cells. Due to the

lack of structural information on QFL-TCR:FL9:Qa-1^b ternary complex , we started by

predicting the structure using AlphaFold2 (***Mirdita et al., 2022***). The sequences of the α1 and α2

domain of Qa-1^b α-chain, FL9 and Vα3.2Jα21, Vβ1Dβ1Jβ2-7 were submitted as input. Most of

the residues in the predicted structure have high to intermediate confidence scores as was

measured by the predicted local distance difference test (pLDDT). The output structure model of

the top ranked pLDDT score ('rank_1' = 90.8) was selected for further visualization

(**Supplementary Fig. 8 b**). The simulated QFL-TCR:FL9:Qa-1^b complex model suggests a

similar TCR docking as observed in TCRαβ complexed with classical MHC I (**Fig.6 a**). While

we fully appreciated the limitations in accurately predicting TCR docking, prediction of the

pMHC complex is nevertheless relatively reliable based on the well-defined principles (***Nielsen***

et al., 2020). Superimposition of simulated FL9:Qa-1^b model onto the crystallographic structure

of Qdm:Qa-1^b (PDB 3VJ6) suggested similar main chain conformations between FL9 and Qdm

(**Fig.6 b**). We thus refer to the reported structure of Qdm:Qa-1^b whose anchoring residues were

P2, P3, P6, P7 and P9 of AMAPRTLLL(Qdm) for determination of putative key residues on FL9

(***Zeng et al., 2012***). Among the rest of the non-anchoring residues, we hypothesized that the

glutamic acid at P4 of FL9 may be critical for interacting with the QFL TCR. Notably, the

importance of P4 for TCR interaction was also highlighted in recently published studies where

crystal structures of several disease-related TCR:pMHC I complexes were reported including an

αβTCR complexed with an HIV peptide loaded in HLA-B5301 (***Li et al., 2023***) and a pre-TCR

complexed with vesicular stomatitis virus octapeptide(VSV8) loaded in H-2K^b (***Li et al., 2021***).

To test the hypothesis, we used the QFL T hybridoma cell line BEko8Z which expresses TCR-

induced β-galactosidase (LacZ) to test whether the mutant FL9 peptide with the P4 residue

substituted by alanine altered T cell responses. The P6 threonine whose side chain basically

bends towards to the binding groove with relatively limited access to TCR engagement was

tested in parallel as a putative anchoring residue. By measuring the LacZ production of BEko8Z

cells in response to Qa-1^b-expressing Lmtk$^-$ (L-Qa-1^b) cells pulsed with FYAAATPML (FL9-

P4A) or FYAEAAPML (FL9-P6A), we found that in agreement with the prediction, replacement

of the P6 threonine reduced TCR recognition, while replacement of the P4 glutamic acid led to

the complete loss of the BEko8Z response (**Fig. 6c**). Additionally, we performed Qa-1^b binding

predictions for various FL9 mutant peptides using NetMHCpan4.1 (***Reynisson et al., 2020***). The

predicted binding scores suggested that substitution of the P9 Leu with Ala should decrease

binding of FL9 peptide to Qa-1[b], whereas mutation of the rest of the residues were less likely to

have such effect. Hence, Leu at P9 position may serve as a key anchoring residue

(**Supplementary Fig. 8 c**). Thus, P4, P6 and C-terminus positions were considered the key

residues with the P4 Glu being particularly critical for QFL T recognition. Based on these data,

the presence of glutamic acid at P4 was considered to be important, while the C-terminus leucine

and P6 threonine were considered secondary. Although it is possible that the residues that were

not tested in the T cell assays are involved in peptide binding or TCR interaction, data on the P4,

P6 and P9 residues of FL9 peptide were sufficient for determining peptide homology and

establishing a testable library of bacterial FL9 homologs.

To identify potential bacterial FL9 homologous peptides, we aligned the FL9 peptide sequence

with proteomes of the five dominant gut commensal bacterial phyla, including Actinobacteria,

Bacteroidetes, Firmicutes, Proteobacteria, Verrucomicrobia from the NCBI Swissprot Non-

redundant UniProtKB database (***Donaldson et al., 2016***). The alignment result was further

curated to 30 candidate peptides based on their homology with FL9 peptide as described above

(**Fig. 8 d, Supplementary Table.1**). By measuring LacZ production of BEko8Z cells in response

to L-Qa-1[b] cells pulsed with 50nM of each candidate peptide, we identified a single stimulatory

peptide FYAEDGTPIL (FL10) that is expressed in *P. pentosaceus*, a gram-positive lactic acid

bacteria (LAB) that colonizes in the small intestine (**Fig.6 e**). While sensitivity to FL10 was

~1,000 folds less than FL9 peptide, BEko8Z cells could recognize FL10 at concentrations as low

as 1.5nM. In contrast, FYAEDDTPIV(FV10), a FL10 homologous peptide expressed in

Lactobacillus johnsonii which is a common LAB that colonize GI tract, failed to activate

BEko8Z cells (**Fig.6 f, Supplementary Fig. 8 d**).

QFL T cells are retained in gut by a commensal bacterium

To further investigate the association between *P. pentosaceus* and Vα3.2[+]QFL T cells in gut, we

first generated QFLTg_WT partial hematopoietic chimeras using germ-free WT mice (GF

QFLTg_WT) and monocolonized the GF chimera mice with *P. pentosaceus* at 8 weeks of age.

The mice were analyzed 8 weeks later (**Fig.7 a**). The donor-derived QFLTg cells and the

recipient-derived Vα3.2$^+$QFL T cells were distinguished by GFP expression and further defined

as the CD45$^+$CD19$^-$TCRβ$^+$CD4$^-$Vα3.2$^+$QFL-Dex-PE$^+$ population on flow cytometry as was

described earlier. While the donor-derived QFLTg populations were present and phenotypically

unaltered in the spleen and siIEL compartment of GF QFLTg_WT chimera mice at 8 weeks of

age, they were barely detectable in the GF chimera mice at 16 weeks of age (**Supplementary**

Fig. 9, Fig.7 b). Notably, monocolonization of the GF QFLTg_WT chimera mice with *P.*

pentosaceus completely restored the donor-derived QFLTg population in both tissues in 16-week

GF chimera mice (**Fig.7 b**). We then further tested whether restoration of the gut QFL T

population was specifically associated with *P. pentosaceus* by monocolonizing GF WT mice

with *P. pentosaceus* or *L. johnsonii* (**Fig.7 a**). Analysis of the microbiome composition of fecal

pallets showed that both species efficiently colonized the GF mice (**Fig.7 c**). While the

endogenous Vα3.2$^+$QFL T population in the siIEL compartment was retained by *P. pentosaceus*

colonization of GF mice at 16 weeks of age, *L. johnsonii* colonization failed to restore the

population to level comparable with SPF or *P. pentosaceus* colonized mice. On the other hand,

maintenance of the endogenous splenic QFL T population was independent of microbes, in line

with our earlier observations (**Fig.7 d**). Donor-derived QFLTg and endogenous Vα3.2$^+$QFL T

cells in the spleen likely showed discrepant results due to different sources of precursors for the

two populations. Unlike the endogenous populations which were constantly replenished, the

donor-derived QFLTg cells rose from a fixed number of bone marrow cells, and gradually

decreased in number as the animal age (**Supplementary Fig. 10**). Thus, presence of the ligand

which was supplied by peptide derived from *P. pentosaceus* might be required for the donor-

derived QFLTg cells to be retained in the spleen, whereas the endogenous splenic population

was unaffected by microbes. Phenotype analysis revealed that both the donor-derived QFLTg

cells and the endogenous Vα3.2$^+$QFL T cells from the colonized GF chimera mice showed a

memory phenotype identical to that seen in the SPF chimera mice (**Supplementary Fig. 11**). We

thus conclude that the late-life maintenance of Vα3.2$^+$QFL T cells in the small intestine was

dependent on the commensal bacterium *P. pentosaceus* which expresses a FL9 homologue

peptide. While the presence of the bacterial FL10 peptide likely plays an important role in

retention of gut QFL T cells, we do not exclude the possible contributions of other commensal

bacterial factors. In addition, the unconventional phenotype of QFL T cells was likely acquired

through microbe-independent mechanisms.

Discussion

The small intestine epithelium harbors a highly heterogenous intraepithelial lymphocyte

population which is comprised of a large proportion of CD4$^-$CD8αβ$^-$CD8αα$^+$ T cells. Due to the

heterogeneity, it has been challenging to identify or study a particular antigen-specific T cell

clone from the population. Here we found that QFL T cells, a Vα3.2-expressing CD8$^+$ T cell

population which specifically recognizes the Qa-1^b-FL9 ligand presented on ERAP1-KO cells,

naturally resided in both the spleen and small intestine epithelium of naïve WT mice. Further

characterization of QFL T cells revealed their unconventional phenotype, functionality and

intimate association with gut microbiota (**Table 1**).

Unlike conventional memory T cells which are generated upon encounter with their cognate

peptide antigens on classical MHC molecules, 'innate' memory T cells arise under homeostatic

conditions in response to neonatal lymphopenia, high levels of IL-4 or exposure to self-antigens

in naïve mice (***Jameson et al., 2015; Sprent et al., 2011***). Here, we found that splenic

Vα3.2$^+$QFL T cells display a memory phenotype and the corresponding rapid effector function,

but acquire this phenotype in a Qa-1^b dependent manner. Given the unique dependence on Qa-1^b

expression for the phenotype imprinting of QFL T cells, it is possible that QFL T cells are

exposed to transiently induced QFL epitope on cells due to various intracellular stressors that

might affect ERAP1 function. Alternatively, QFL T cells might cross-react with Qa-1^b presented

FL9 homologue peptide(s). Indeed, we found that Vα3.2$^+$QFL T cells cross-reacted with one

such variant of FL9 peptide (FL10) expressed in a commensal bacterium and presented by Qa-1^b.

However, the Qa-1^b-FL10 ligand itself is unlikely to be directly associated with the memory

imprinting for QFL T cells as these cells showed unaltered phenotype in germ-free mice. There

might be other unidentified self-peptide(s) presented by Qa-1^b that are involved in the process. It

is also possible that there is some level of degeneracy in the TCR-Qa-1^b interaction, so the

precise peptide being presented is less critical than the presence of any peptide-Qa-1^b complex. It

is worth noting that the predicted structure of QFL-TCR:FL9:Qa-1^b complex indicated that the

FL9 peptide binds to Qa-1^b in a manner potentially similar to Qdm. Because the binding groove

of Qa-1^b and its human homolog HLA-E display remarkable structural homology (***Zeng et al.,***

2012), it is possible that HLA-E can present FL9 or its homolog peptides and further support the

establishment of a QFL T-like CD8 T population. Studies on cytomegalovirus-vectored vaccines

have shown that HLA-E-restricted CD8 T cells could recognize a broad repertoire of peptides

presented upon vaccination and provide protection against HIV-1 infection (**Hansen et al., 2016;**

Yang et al., 2021). Thus, analysis of the HLA-E peptide repertoire and investigation on the

presence of HLA-E-restricted human version of QFL T cells will be promising and meaningful

future directions.

In line with a previous study, we observed that the Vα3.2-expressing CD8αβ$^+$ T population was

comprised of a relatively larger proportion of memory phenotype cells in the spleen than the

non-Vα3.2 expressing population (**Prasad et al., 2021**). In addition, these memory phenotype

non-QFL Vα3.2$^+$CD8αβ$^+$ T cells were lost in Qa-1^b-KO mice. These results indicate that

Vα3.2$^+$QFL T cells are likely only one prototypical example of Qa-1^b restricted Vα3.2$^+$CD8$^+$ T

cells clones with memory phenotype and innate-like function. We detected abundant Vα3.2$^+$QFL

T cells in small intestine epithelium which showed the signature phenotype of natIELs including

expression of CD8αα and lack of CD4, CD8αβ, CD5 and CD90. Similar to the splenic

population, while TAP is required for the presence of the population, the unconventional

phenotype of Vα3.2$^+$QFL T cells in the gut is strongly Qa-1^b-dependent. It is worth noting that

compared with the other Qa-1^b-restrict T cells reported, such as a population of semi-invariant T

cells which recognize TAP-independent peptides (**Doorduijn et al., 2018**) or an insulin peptide-

specific CD8 T population (**Sullivan et al., 2002**), QFL T is likely a distinct population which

undergoes alternative developmental stages which requires TAP-dependent presentation of FL9

(**Nagarajan et al., 2012**) (or other alternative essential peptides) for positive selection and Qa-1^b

for acquiring their unconventional features. Additionally, it is so far reported that CD8αα$^+$ IELs

can be restricted to K^{b}D^b, CD1d or even unknown MHC Ib molecules (**Mayans et al., 2014;**

Ruscher et al., 2017). Here by characterization of the gut Vα3.2$^+$QFL T cells, we showed

evidence of abundant Qa-1^b-restricted TCRαβ$^+$ natIELs being present in the gut.

Functionally, the gut Vα3.2$^+$QFL T cells showed features consistent with their CD8αα$^+$ natIEL

phenotype. Notably, although the siIEL Vα3.2$^+$QFL T cells showed delayed and reduced IFN-ɣ

production in response to the cognate peptide, they displayed higher basal levels of intracellular

IFN-ɣ than CD8αβ$^+$ IELs. It has been proposed that spontaneous IFN-ɣ secretion by IELs may

be an important component of immunosurveillance at the mucosal surface, capable of identifying

and eliminating transformed cells (***Carol et al., 1998; Roberts et al., 1993***). Thus, siIEL QFL T

cells may likewise be poised for immediate elimination of abnormal cells.

CD8αα⁺ IELs are intrinsically programmed for innate functionality, as was shown by

PMA/ionomycin induced production of high level of IFN-ɣ, CXCL2 etc. (***Van Kaer et al.,***

2014). However, our observation of the relative hyporesponsiveness of siIEL Vα3.2⁺QFL T to

stimuli indicated that under physiological conditions, the activation of IELs was more complex

and highly regulated as TCRs and inhibitory coreceptors such as CD8αα were engaged

(***Cheroutre et al., 2008; Madakamutil et al., 2004***). It is not surprising that the gut QFL T cells

were functionally quiescent as these cells are constantly challenged by the microbial or dietary

antigens from gut lumen (***Denning et al., 2007***). Cross-reaction between QFL T cells and peptide

expressed in the commensal bacterium *P. pentosaceus* further supports this notion. Surprisingly,

monocolonization of the GF mice with *P. pentosaceus* was sufficient to restore gut Vα3.2⁺QFL

T cells that were lost in old GF mice. Of note, the control bacterium *L. johnsonni,* which lack

FL10 expression, did not restore gut Vα3.2⁺QFL T cells. Despite the evidence we remain

cautious of other possible signals generated from colonization of the commensal bacterium, such

as secretion of bacteriocins.

In conclusion, Vα3.2-expressing QFL T cells represent an unconventional population of

Qa-1ᵇ-restricted CD8⁺ T cells which naturally reside at high frequency in small intestine

epithelium. These cells have memory phenotype in the spleen and reveal the characteristics of

natIELs in the gut. In addition, the fact that Vα3.2⁺QFL T cross-reacts with FL9 homologue

peptide expressed in a commensal bacterium suggests that these cells might be functionally

versatile. Although we have not yet detected steady-state FL9 presentation, it remains to be

assessed whether FL9 peptide can be transiently presented under certain conditions at particular

locations or within certain time windows. Further study of the biological significance of these

cells could potentially reveal the delicate equilibrium between gut microenvironment and

immune system.

Acknowledgements

We thank Dr. X. Li at the University of Science and Technology of China and the Ragon

Institute of MGH, MIT and Harvard for providing insights on prediction of peptide binding to

unconventional MHC I molecules, Dr. L. Coscoy at University of California, Berkeley for

providing cellular biology expertise, and Mr. P. Rhee at Shastri Lab, and Dr. H. Ding at the JHU

germ-free core facility for technical support. This work was supported by grants R01AI130210,

R37AI060040 and R01AI149341 from the National Institutes of Health to NS.

Methods

Mice. SPF wild-type C57BL/6J, H-2T23-deficient Qa-1^b-KO, C57BL/6-Tg

(TcraTcrb)1100Mjb/J OT-1, or C57BL/6-Tg (UBC-GFP)30Scha/J B6.GFP mice were obtained

from the Jackson Laboratory. QFL T TCR transgenic mice QFLTg were generated in the

laboratory of E. Robey (University of California Berkeley, Berkeley, CA) and housed in our

facility. GFP$^+$QFLTg mice were generated in our facility by crossing B6.GFP mice with QFLTg

mice. SPF QFLTg_WT, SPF OT-1_WT and GF QFLTg_WT partial hematopoietic chimera mice

were generated as previously described(***Ladi et al., 2008***). GF wild-type C57BL/6J and GF

QFLTg_WT chimera mice were generated and maintained in the Johns Hopkins Germ-free

Mouse Core Facility. Mice were housed and all procedures were done in accordance with

protocols approved by Animal Care and Use Committee of the Johns Hopkins University School

of Medicine.

Generation of the QFLTg mouse. The QFLTg TCR α-and β-chain sequences were cloned and

amplified from the genomic DNA of BEko8Z hybridoma(***Guan et al., 2017; Nagarajan et al.,***

2012). The TCR α-chain was amplified with the forward primer 5'-

AAAACCCGGGCCAAGGCTCAGCCATGCTCCTGG-3' and the reverse primer 5'-

AAAAGCGGCCGCATACAACATTGGACAAGGATCCAAGCTAAAGAGAACTC-3'. The

TCR β-chain was cloned with the forward primer 5'-

AAAACTCGAGCCCGTCTGGAGCCTGATTCCA-3' and the reverse primer 5'-

AAAACCGCGGGGGACCCAGGAATTTGGGTGGA-3'. The TCR α-chain DNA fragment

was cloned into pTα cassette vector by inserting it between the Xmal and NotI sites, while the

TCR β-chain DNA fragment was cloned into pTβ cassette vector in between the XhoI and SacII

sites(***Kouskoff et al., 1995***). The ampicillin resistance gene was removed from pTα and pTβ

489 cassette by EarI enzyme digest. The QFLTg mice were generated on C57BL/6J background in

490 the Cancer Research Laboratory Gene Targeting Facility at UC Berkeley under standard

491 procedures. Founder mice were identified by flow cytometry and PCR genotyping of tail

492 genomic DNA using primers mentioned above.

494 **Antibodies, cell lines and peptides.** Antibody for flow cytometry were from BioLegend (anti-

495 CD45(30-F11), anti-CD19(1D3/CD19), anti-TCRβ(H57-597), anti-CD8β(53-5.8), anti-

496 Vα3.2(RR3-16), anti-Vα2(B20.1), anti-CD90.2(53-2.1), anti-CD5(53-7.3), anti-IFN-ɣ(XMG1.2),

497 anti-CD62L(MEL-14) and BD Biosciences (anti-CD4(RM4-5), anti-CD8α (53-6.7), anti-

498 CD44(IM7)). BEko8Z, L-Qa-1^b or Lmtk$^-$ cells were maintained as previously

499 described(*Nagarajan et al., 2012*). Peptides were obtained from GenScript. The purity of SL8,

500 FL9 and FL10 peptides were ⩾98%.

502 **Generation of the QFL-dextramer.** Qa-1^b-FL9 (QFL) monomers were synthesized by the

503 Tetramer Core Facility of the US National Institutes of Health. Phycoerythrin (PE)- or

504 allophycocyanin (APC)-conjugated streptavidin were obtained from Agilent. The QFL

505 dextramers were generated following the 'dextran doping' technique(*Bethune et al., 2017*). The

506 QFL monomers were incubated with PE or APC conjugated streptavidin at a molar ratio of 3:1 at

507 4℃ for 10min followed by addition of biotinylated dextran molecular weight 500kDa at a molar

508 ratio of 1:20 with respect to streptavidin. The mixture was further incubated at 4℃ for ≧1h

509 before being used in experiments.

511 **Isolation of IELs.** Small intestinal IELs were isolated following an established protocol with

512 minor modifications (*Qiu et al., 2018*). In brief, the small intestine with Payer's patches removed

513 were cut into appropriate length. The fecal content and mucus were removed by expelling with

514 the flat side of forceps followed by flushing with PBS. The intestine was cut open longitudinally

515 to reveal the epithelium and further cut laterally into ~2cm pieces. The intestine pieces were then

516 placed in 25ml warmed dithioerythritol (DTE) solution (Ca^{2+}- and Mg^{2+}-free Hanks balanced salt

517 solution, HEPES-bicarbonate buffer, 10% FCS) in 50ml conical tube and were shaken at 75rpm

518 37℃ for 20min. The tube was vortexed for 10s before the supernatant was transferred into a new

519 50ml conical tube through a 70µM cell strainer. The cells were palleted and resuspend in 44%

520   Percoll solution (44% Percoll in RPMI 1640). Density gradient was generated by underlaying the

521   cell suspension with 67% Percoll solution (67% Percoll in RPMI 1640). The gradient cell

522   suspension was further centrifuged at 1600 × g for 20 min at RT without using the brake. The

523   layer of cells at the 44% and 67% interphase were thus collected as IELs.

524
525

Enrichment for dextramer-positive cells. Splenic dextramer-positive cells were enriched as

previously described(*Nagarajan et al., 2012*). IELs were resuspend in 100µl sorter buffer (0.1%

sodium azide and 5% FCS in PBS). PE or APC labeled QFL dextramers were added at a final

dilution of 1:100. Cells were incubated at room temperature for 45 min, then were washed twice

with 3ml of sorter buffer. 20µl of anti-PE and 20µl of anti-APC microbeads (Miltenyi Biotec)

were added into cells resuspended in 450µl of sorters buffer, followed by incubation of 20 min at

4°C. Cells were washed twice with 3ml sorter buffer. The PE- and APC-labeled cells were

positively selected by passing through LS magnetic columns (Miltenyi Biotec). The entire

isolated population was stained with anti-CD45, anti-CD19, anti-TCRβ, anti-CD4, anti-CD8α

and anti-CD8β. QFL T cells were gated as CD45$^+$CD19$^-$TCRβ$^+$CD4$^-$QFL-Dex-PE$^+$QFL-Dex-

APC$^+$ population. The absolute numbers and frequencies of the cells were calculated based on a

fixed number of CountBright Beads (Thermo Fisher) added into each sample.

Cytokine stimulation and proliferation assay. Splenocytes or small intestinal IELs isolated

from QFLTg or OT-1 mice were depleted of non-T cells using Pan T isolation Kit II (Miltenyi

Biotec). The negatively selected T cells were then labeled with 5µM Cell Tracing Violet (CTV)

(Thermo Fisher) and adjusted to the concentration of 10^6/ml. 10^6 of the CTV labeled T cells were

stimulated *in vitro* for 3 or 5 days in cultures containing 100ng/ml IL-15 (BioLegend) or a

combination of 100ng/ml IL-7 (BioLegend) and 100ng/ml IL-18 (BioLegend) in 24-well plate at

37°C. Proliferation of cells were measured by dilution of CTV on flow cytometer.

CTL assay. Lymphocytes isolated from the spleen or siIEL compartment of naïve QFLTg_WT

or OT-1_WT chimera mice were cultured *in vitro* with 2µM of FL9 or SL8 peptide respectively

together with GolgiPlug (BD Biosciences) for 4.5h. WT splenocytes were supplemented into

siIEL cultures as APCs. Cells were then washed and stained for surface markers. For

measurement of IFN-ɣ production, cells were fixed and permeabilized using

Fixation/Permeabilization Kit (BD Biosciences) and stained for intracellular IFN-ɣ.

Hybridoma assay. 10^5 of BEko8Z hybridoma cells were cocultured with 10^5 of L-Qa-1^b or

Lmtk- cells in medium containing peptides of indicated concentrations in 96-well plates at 37℃

for 12 hours. T cell response was determined by cleavage of the chromogenic LacZ substrate

chlorophenol red-β-D-galactopyranoside (CPRG, Millipore Sigma) by TCR-induced LacZ and

measure by spectrophotometry as presented by absorbance at 595 nm (A595) (**Sanderson et al.,**

1994). Background signals were subtracted by measurement of absorbance at 655nm.

Commensal culture and colonization. *Pediococcus pentosaceus Mees* (33314) and

Lactobacillus johnsonii (33200) were obtained from ATCC. The bacteria were grown overnight

at 37℃ in 416 Lactobacilli MRS Broth (BD Biosciences). GF QFLTg_WT chimera mice were

gavaged with 200µl of $2x10^8$/ml CFU of *P. pentosaceus Mees* at 8 weeks of age. GF WT mice

were gavaged with 200µl of $2x10^8$/ml CFU of *P. pentosaceus Mees* or *L. johnsonii* at 8 and 10

weeks of age. Fecal pallets collected from the colonized mice were homogenized in 416

Lactobacilli MRS Broth, and the associated mice were determined by plating the appropriate

dilution of homogenate on 416 Lactobacilli MRS agar plates. Bacterial colonization efficiency

was determined by whole shotgun metagenomic sequencing of fecal pallets collected from the

colonized mice at 16 weeks of age (TransnetYX Microbiome).

AlphaFold2 structure and Qa-1^b binding predictions. The predicted structure of QFL

TCR:FL9:Qa-1^b complex was obtained by submitting the sequences of the α1 and α2 domain of

Qa-1^b α-chain, FL9 and Vα3.2Jα21, Vβ1Dβ1Jβ2-7 as 'hetero-oligomer' to ColabFold v1.5.2-

patch: AlphaFold2 using MMseqs2 at

https://colab.research.google.com/github/sokrypton/ColabFold/blob/main/AlphaFold2.ipynb.

The prediction was conducted using the alphafold2_multimer_v3 model on default settings.

Visualization and alignment of the protein structures was conducted on the PyMOL software.

The Qa-1^b binding prediction was conducted by submitting the sequences of Qdm, FL9 or

mutant FL9 peptides and Qa-1^b (UniProt: P06339) to NetMHCpan4.1 at

https://services.healthtech.dtu.dk/services/NetMHCpan-4.1/.

References

Belkaid, Y., and Hand, T.W. (2014). Role of the microbiota in immunity and inflammation. *Cell*
157, 121-141.10.1016/j.cell.2014.03.011, PMID:24679531

Bethune, M.T., Comin-Anduix, B., Fu, Y.-H.H., Ribas, A., and Baltimore, D. (2017).
Preparation of peptide–MHC and T-cell receptor dextramers by biotinylated dextran doping.
Biotechniques **62**, 123-130.10.2144/000114525, PMID:28298179

Blanchard, N., and Shastri, N. (2008). Coping with loss of perfection in the MHC class I peptide
repertoire. *Curr Opin Immunol* **20**, 82-88.10.1016/j.coi.2007.12.004, PMID:18243675

Carol, M., Lambrechts, A., Van Gossum, A., Libin, M., Goldman, M., and Mascart-Lemone, F.
(1998). Spontaneous secretion of interferon gamma and interleukin 4 by human intraepithelial
and lamina propria gut lymphocytes. *Gut* **42**, 643-649.10.1136/gut.42.5.643, PMID:9659157

Cheroutre, H., and Lambolez, F. (2008). Doubting the TCR coreceptor function of CD8αα.
Immunity **28**, 149-159.10.1016/j.immuni.2008.01.005, PMID:18275828

Cheroutre, H., Lambolez, F., and Mucida, D. (2011). The light and dark sides of intestinal
intraepithelial lymphocytes. *Nat Rev Immunol* **11**, 445-456.10.1038/nri3007, PMID:21681197

Das, G., Gould, D.S., Augustine, M.M., Fragoso, G., Sciutto, E., Stroynowski, I., Van Kaer, L.,
Schust, D.J., Ploegh, H., and Janeway, C.A., Jr. (2000). Qa-2-dependent selection of CD8αα T
cell receptor αβ+ cells in murine intestinal intraepithelial lymphocytes. *J Exp Med* **192**, 1521-
1528.10.1084/jem.192.10.1521, PMID:11085754

Das, G., and Janeway, C.A., Jr. (1999). Development of CD8αα and CD8αβ T cells in major
histocompatibility complex class I-deficient mice. *J Exp Med* **190**, 881-
884.10.1084/jem.190.6.881, PMID:10499926

Das, G., and Janeway, C.A., Jr. (2003). MHC specificity of iIELs. *Trends Immunol* **24**, 88-
93.10.1016/s1471-4906(02)00036-4, PMID:12547506

Denning, T.L., Granger, S.W., Mucida, D., Graddy, R., Leclercq, G., Zhang, W., Honey, K.,
Rasmussen, J.P., Cheroutre, H., Rudensky, A.Y.*, et al.* (2007). Mouse TCRαβ+CD8αα
intraepithelial lymphocytes express genes that down-regulate their antigen reactivity and
suppress immune responses. *J Immunol* **178**, 4230-4239.10.4049/jimmunol.178.7.4230,
PMID:17371979

Donaldson, G.P., Lee, S.M., and Mazmanian, S.K. (2016). Gut biogeography of the bacterial
microbiota. *Nat Rev Microbiol* **14**, 20-32.10.1038/nrmicro3552, PMID:26499895

Doorduijn, E.M., Sluijter, M., Querido, B.J., Seidel, U.J.E., Oliveira, C.C., van der Burg, S.H.,
and van Hall, T. (2018). T Cells Engaging the Conserved MHC Class Ib Molecule Qa-1(b) with
TAP-Independent Peptides Are Semi-Invariant Lymphocytes. *Front Immunol* **9**,
60.10.3389/fimmu.2018.00060, PMID:29422902

Faria, A.M.C., Reis, B.S., and Mucida, D. (2017). Tissue adaptation: Implications for gut
immunity and tolerance. *J Exp Med* **214**, 1211-1226.10.1084/jem.20162014, PMID:28432200

Godfrey, D.I., Uldrich, A.P., McCluskey, J., Rossjohn, J., and Moody, D.B. (2015). The
burgeoning family of unconventional T cells. *Nat Immunol* **16**, 1114-1123.10.1038/ni.3298,
PMID:26482978

Grandea, A.G., 3rd, Golovina, T.N., Hamilton, S.E., Sriram, V., Spies, T., Brutkiewicz, R.R.,
Harty, J.T., Eisenlohr, L.C., and Van Kaer, L. (2000). Impaired assembly yet normal trafficking
of MHC class I molecules in Tapasin mutant mice. *Immunity* **13**, 213-222.10.1016/s1074-
7613(00)00021-2, PMID:10981964

Guan, J., Peske, J.D., Taylor, J.A., and Shastri, N. (2021). The nonclassical immune surveillance
for ERAAP function. *Curr Opin Immunol* **70**, 105-111.10.1016/j.coi.2021.05.008,
PMID:34098489

Guan, J., Yang, S.J., Gonzalez, F., Yin, Y., and Shastri, N. (2017). Antigen Processing in the
Endoplasmic Reticulum Is Monitored by Semi-Invariant αβ TCRs Specific for a Conserved
Peptide-Qa-1b MHC Class Ib Ligand. *J Immunol* **198**, 2017-2027.10.4049/jimmunol.1600764,
PMID:28108559

Hammer, G.E., Gonzalez, F., Champsaur, M., Cado, D., and Shastri, N. (2006). The
aminopeptidase ERAAP shapes the peptide repertoire displayed by major histocompatibility
complex class I molecules. *Nat Immunol* **7**, 103-112.10.1038/ni1286, PMID:16299505

Hammer, G.E., Gonzalez, F., James, E., Nolla, H., and Shastri, N. (2007a). In the absence of
aminopeptidase ERAAP, MHC class I molecules present many unstable and highly
immunogenic peptides. *Nat Immunol* **8**, 101-108.10.1038/ni1409, PMID:17128277

Hammer, G.E., Kanaseki, T., and Shastri, N. (2007b). The final touches make perfect the
peptide-MHC class I repertoire. *Immunity* **26**, 397-406.10.1016/j.immuni.2007.04.003,
PMID:17459809

Hansen, S.G., Wu, H.L., Burwitz, B.J., Hughes, C.M., Hammond, K.B., Ventura, A.B., Reed,
J.S., Gilbride, R.M., Ainslie, E., Morrow, D.W.*, et al.* (2016). Broadly targeted CD8(+) T cell
responses restricted by major histocompatibility complex E. *Science* **351**, 714-
720.10.1126/science.aac9475, PMID:26797147

Jameson, S.C., Lee, Y.J., and Hogquist, K.A. (2015). Innate memory T cells. *Adv Immunol* **126**,
173-213.10.1016/bs.ai.2014.12.001, PMID:25727290

Klose, C.S., Blatz, K., d'Hargues, Y., Hernandez, P.P., Kofoed-Nielsen, M., Ripka, J.F., Ebert,
K., Arnold, S.J., Diefenbach, A., Palmer, E.*, et al.* (2014). The transcription factor T-bet is
induced by IL-15 and thymic agonist selection and controls CD8αα+ intraepithelial lymphocyte
development. *Immunity* **41**, 230-243.10.1016/j.immuni.2014.06.018, PMID:25148024

Kouskoff, V., Signorelli, K., Benoist, C., and Mathis, D. (1995). Cassette vectors directing
expression of T cell receptor genes in transgenic mice. *J Immunol Methods* **180**, 273-
280.10.1016/0022-1759(95)00002-r, PMID:7714342

Ladi, E., Herzmark, P., and Robey, E. (2008). In situ imaging of the mouse thymus using 2-
photon microscopy. *J Vis Exp*.10.3791/652, PMID:19066556

Li, X., Mizsei, R., Tan, K., Mallis, R.J., Duke-Cohan, J.S., Akitsu, A., Tetteh, P.W., Dubey, A.,
Hwang, W., Wagner, G.*, et al.* (2021). Pre-T cell receptors topologically sample self-ligands
during thymocyte beta-selection. *Science* **371**, 181-185.10.1126/science.abe0918,
PMID:33335016

Li, X., Singh, N.K., Collins, D.R., Ng, R., Zhang, A., Lamothe-Molina, P.A., Shahinian, P., Xu,
S., Tan, K., Piechocka-Trocha, A.*, et al.* (2023). Molecular basis of differential HLA class I-

restricted T cell recognition of a highly networked HIV peptide. *Nat Commun* **14**,
2929.10.1038/s41467-023-38573-8, PMID:37217466

Macpherson, A.J., and Harris, N.L. (2004). Interactions between commensal intestinal bacteria
and the immune system. *Nature Reviews Immunology* **4**, 478-485.10.1038/nri1373,

Madakamutil, L.T., Christen, U., Lena, C.J., Wang-Zhu, Y., Attinger, A., Sundarrajan, M.,
Ellmeier, W., von Herrath, M.G., Jensen, P., Littman, D.R*., et al.* (2004). CD8αα-mediated
survival and differentiation of CD8 memory T cell precursors. *Science* **304**, 590-
593.10.1126/science.1092316, PMID:15105501

Mayans, S., Stepniak, D., Palida, Sakina F., Larange, A., Dreux, J., Arlian, Britni M.,
Shinnakasu, R., Kronenberg, M., Cheroutre, H., and Lambolez, F. (2014). αβT Cell Receptors
Expressed by CD4–CD8αβ– Intraepithelial T Cells Drive Their Fate into a Unique Lineage with
Unusual MHC Reactivities. *Immunity* **41**, 207-218.10.1016/j.immuni.2014.07.010,

Mirdita, M., Schütze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M. (2022).
ColabFold: making protein folding accessible to all. *Nature Methods* **19**, 679-
682.10.1038/s41592-022-01488-1,

Mota-Santos, T., Masmoudi, H., Voegtle, D., Freitas, A., Coutinho, A., and Cazenave, P.A.
(1990). Divergency in the specificity of the induction and maintenance of neonatal suppression.
Eur J Immunol **20**, 1717-1721.10.1002/eji.1830200814, PMID:2209685

Nagarajan, N.A., de Verteuil, D.A., Sriranganadane, D., Yahyaoui, W., Thibault, P., Perreault,
C., and Shastri, N. (2016). ERAAP Shapes the Peptidome Associated with Classical and
Nonclassical MHC Class I Molecules. *J Immunol* **197**, 1035-1043.10.4049/jimmunol.1500654,
PMID:27371725

Nagarajan, N.A., Gonzalez, F., and Shastri, N. (2012). Nonclassical MHC class Ib-restricted
cytotoxic T cells monitor antigen processing in the endoplasmic reticulum. *Nat Immunol* **13**,
579-586.10.1038/ni.2282, PMID:22522492

Nielsen, M., Andreatta, M., Peters, B., and Buus, S. (2020). Immunoinformatics: Predicting
Peptide-MHC Binding. *Annu Rev Biomed Data Sci* **3**, 191-215.10.1146/annurev-biodatasci-
021920-100259, PMID:37427310

Okazawa, A., Kanai, T., Nakamaru, K., Sato, T., Inoue, N., Ogata, H., Iwao, Y., Ikeda, M.,
Kawamura, T., Makita, S*., et al.* (2004). Human intestinal epithelial cell-derived interleukin (IL)-
18, along with IL-2, IL-7 and IL-15, is a potent synergistic factor for the proliferation of
intraepithelial lymphocytes. *Clin Exp Immunol* **136**, 269-276.10.1111/j.1365-2249.2004.02431.x,
PMID:15086390

Park, S.H., Guy-Grand, D., Lemonnier, F.A., Wang, C.R., Bendelac, A., and Jabri, B. (1999).
Selection and expansion of CD8αα+ T cell receptor αβ+ intestinal intraepithelial lymphocytes in
the absence of both classical major histocompatibility complex class I and nonclassical CD1
molecules. *J Exp Med* **190**, 885-890.10.1084/jem.190.6.885, PMID:10499927

Prasad, M., Wojciech, L., Brzostek, J., Hu, J., Chua, Y.L., Tung, D.W.H., Yap, J., Rybakin, V.,
and Gascoigne, N.R.J. (2021). Expansion of an Unusual Virtual Memory CD8+ Subpopulation
Bearing Valpha3.2 TCR in Themis-Deficient Mice. *Front Immunol* **12**,
644483.10.3389/fimmu.2021.644483, PMID:33897691

Qiu, Z., and Sheridan, B.S. (2018). Isolating Lymphocytes from the Mouse Small Intestinal
Immune System. *J Vis Exp*.10.3791/57281, PMID:29553537

Reynisson, B., Alvarez, B., Paul, S., Peters, B., and Nielsen, M. (2020). NetMHCpan-4.1 and
NetMHCIIpan-4.0: improved predictions of MHC antigen presentation by concurrent motif
deconvolution and integration of MS MHC eluted ligand data. *Nucleic Acids Research* **48**,
W449-W454.10.1093/nar/gkaa379,

Roberts, A.I., O'Connell, S.M., Biancone, L., Brolin, R.E., and Ebert, E.C. (1993). Spontaneous
cytotoxicity of intestinal intraepithelial lymphocytes: clues to the mechanism. *Clin Exp Immunol*
94, 527-532.10.1111/j.1365-2249.1993.tb08229.x, PMID:8252812

Round, J.L., and Mazmanian, S.K. (2009). The gut microbiota shapes intestinal immune
responses during health and disease. *Nat Rev Immunol* **9**, 313-323.10.1038/nri2515,
PMID:19343057

Ruscher, R., Kummer, R.L., Lee, Y.J., Jameson, S.C., and Hogquist, K.A. (2017). CD8αα
intraepithelial lymphocytes arise from two main thymic precursors. *Nat Immunol* **18**, 771-
779.10.1038/ni.3751, PMID:28530714

Salio, M., Silk, J.D., Jones, E.Y., and Cerundolo, V. (2014). Biology of CD1- and MR1-
restricted T cells. *Annu Rev Immunol* **32**, 323-366.10.1146/annurev-immunol-032713-120243,
PMID:24499274

Sanderson, S., and Shastri, N. (1994). LacZ inducible, antigen/MHC-specific T cell hybrids. *Int*
Immunol **6**, 369-376.10.1093/intimm/6.3.369, PMID:8186188

Schluns, K.S., and Lefrancois, L. (2003). Cytokine control of memory T-cell development and
survival. *Nat Rev Immunol* **3**, 269-279.10.1038/nri1052, PMID:12669018

Serwold, T., Gonzalez, F., Kim, J., Jacob, R., and Shastri, N. (2002). ERAAP customizes
peptides for MHC class I molecules in the endoplasmic reticulum. *Nature* **419**, 480-
483.10.1038/nature01074, PMID:12368856

Shastri, N., Cardinaud, S., Schwab, S.R., Serwold, T., and Kunisawa, J. (2005). All the peptides
that fit: the beginning, the middle, and the end of the MHC class I antigen-processing pathway.
Immunol Rev **207**, 31-41.10.1111/j.0105-2896.2005.00321.x, PMID:16181325

Shastri, N., Schwab, S., and Serwold, T. (2002). Producing nature's gene-chips: the generation of
peptides for display by MHC class I molecules. *Annu Rev Immunol* **20**, 463-
493.10.1146/annurev.immunol.20.100301.064819, PMID:11861610

Sprent, J., and Surh, C.D. (2011). Normal T cell homeostasis: the conversion of naive cells into
memory-phenotype cells. *Nat Immunol* **12**, 478-484.10.1038/ni.2018, PMID:21739670

Sullivan, B.A., Kraj, P., Weber, D.A., Ignatowicz, L., and Jensen, P.E. (2002). Positive selection
of a Qa-1-restricted T cell receptor with specificity for insulin. *Immunity* **17**, 95-
105.10.1016/s1074-7613(02)00343-6, PMID:12150895

Van Kaer, L., Algood, H.M.S., Singh, K., Parekh, V.V., Greer, M.J., Piazuelo, M.B., Weitkamp,
J.H., Matta, P., Chaturvedi, R., Wilson, K.T*., et al.* (2014). CD8αα[+] innate-type lymphocytes in
the intestinal epithelium mediate mucosal immunity. *Immunity* **41**, 451-
464.10.1016/j.immuni.2014.08.010, PMID:25220211

Van Kaer, L., Ashton-Rickardt, P.G., Eichelberger, M., Gaczynska, M., Nagashima, K., Rock,
K.L., Goldberg, A.L., Doherty, P.C., and Tonegawa, S. (1994). Altered peptidase and viral-
specific T cell response in LMP2 mutant mice. *Immunity* **1**, 533-541.10.1016/1074-
7613(94)90043-4, PMID:7600282

Van Kaer, L., Ashton-Rickardt, P.G., Ploegh, H.L., and Tonegawa, S. (1992). TAP1 mutant mice
are deficient in antigen presentation, surface class I molecules, and CD4-8+ T cells. *Cell* **71**,
1205-1214.10.1016/s0092-8674(05)80068-6, PMID:1473153

Yang, H., Rei, M., Brackenridge, S., Brenna, E., Sun, H., Abdulhaqq, S., Liu, M.K.P., Ma, W.,
Kurupati, P., Xu, X.*, et al.* (2021). HLA-E-restricted, Gag-specific CD8(+) T cells can suppress
HIV-1 infection, offering vaccine opportunities. *Sci Immunol* **6**.10.1126/sciimmunol.abg1703,
PMID:33766848

Zeng, L., Sullivan, L.C., Vivian, J.P., Walpole, N.G., Harpur, C.M., Rossjohn, J., Clements,
C.S., and Brooks, A.G. (2012). A structural basis for antigen presentation by the MHC class Ib
molecule, Qa-1b. *J Immunol* **188**, 302-310.10.4049/jimmunol.1102379, PMID:22131332

760
Figure 1 Abundant Vα3.2⁺ QFL T cells in both the spleen and siIEL compartment of naïve
WT mice. (**a**) Definition of QFL T cells by flow cytometry. Cells from the spleen or siIEL
compartment were stained with Qa-1^b-FL9 (QFL) dextramers labeled with phycoerythrin (QFL-
Dex-PE) or allophycocyanin (QFL-Dex-APC) and analyzed for before (Unenriched) or after
(Enriched) magnetic enrichment of dextramer-positive cells. QFL T cells are defined as the
CD45⁺CD19⁻TCRβ⁺CD4⁻QFL-Dex-PE⁺APC⁺ population. Plots representing siIELs from naïve
WT mice after enrichment for QFL-Dex⁺ cells. (**b**) Flow cytometry of cells from the spleen (Sp)
and siIEL compartment of naïve WT mice 'Unenriched' or 'Enriched' for dextramer-positive
cells. Numbers in plots indicate absolute numbers of QFL-Dex⁺ cells detected after enrichment.
(**c**) Absolute numbers (left) and frequencies (right) of QFL-Dex⁺ cells detected (as in **b**) among
TCRβ⁺CD4⁻ population in the spleen or siIEL compartment. **P=0.0069 (**d, f**) Analysis of
Vα3.2 expression on QFL-Dex⁺(QFL) or total CD8⁺ T (Total) cells from the spleen (**d**) or the
siIEL compartment (**f**). Numbers in plots indicate average percentages of Vα3.2⁺ cells within the
indicated populations. (**e**) Frequencies of Vα3.2⁺ cells detected as in **d**. ****P<0.0001 (**g**)
Frequencies of Vα3.2⁺ cells detected as in **f**. **P=0.0058 Representative data are shown in flow
plots (**a, b, d, f**) and the number of replicates is specified in the bar graphs (**c, e, g**). Each symbol
represents data collected from the indicated tissue isolated from an individual mouse. Data from
≧5 independent experiments were pooled for statistical analysis. P values were calculated
with Student's *t* test. 'ns' indicates the comparison was not significant.
780

782
Figure 2 The unconventional phenotype of Vα3.2+ QFL T cells. (**a**) Analysis of CD44
expression on Vα3.2+ or Vα3.2- QFL-Dex+ cells enriched from the spleen of naïve WT mice.
Numbers in plots indicate average percentages of CD44hi cells detected among the Vα3.2+ or
Vα3.2- QFL-Dex+ populations. (**b**) Frequencies of CD44hi cells detected as in **a**.****P<0.0001
(**c**) Analysis of CD8α and CD8β expression on Vα3.2+ or Vα3.2- QFL-Dex+ cells enriched from
the siIEL compartment of naïve WT mice. Numbers in plots indicate average percentages of
CD8αα+ cells detected among the Vα3.2+ or Vα3.2- QFL-Dex+ populations. (**d**) Frequencies of
CD8αα+ cells detected as in **c**. **P=0.0065 Samples of <20 cells in the Vα3.2+QFL-Dex+ gate
were excluded in the phenotype analysis. Representative data are shown in flow plots (**a, c**) and
the number of replicates is specified in the bar graphs (**b, d**). Each symbol represents data
collected from the indicated tissue isolated from an individual mouse. Data from ≧11
independent experiments were pooled for statistical analysis. P values were calculated with
Student's *t* test. 'ns' indicates the comparison was not significant.

796
Figure 3 Phenotype of Vα3.2⁺ QFL T cells in mice of various genotypes. (**a**) Absolute
numbers of Vα3.2⁺QFL-Dex⁺ cells detected in the spleen (left) or siIEL compartment (right) of
Qa-1ᵇ, ERAP1 or TAP deficient (Qa-1ᵇ-KO, ERAP1-KO or TAP-KO) mice in comparison with
WT mice.** P<0.009 ***P=0.0006 Symbols on x-axis indicate that the cells were undetectable
in TAP-KO mice. (**b**) Analysis of CD44 expression on splenic Vα3.2⁺QFL-Dex⁺ cells enriched
from naïve WT, Qa-1ᵇ-KO or ERAP1-KO mice. Numbers in plots indicate average percentages
of CD44ʰⁱ cells. (**c**) Frequencies of CD44ʰⁱ cells detected among Vα3.2⁺(as in **b**) or Vα3.2⁻ QFL
T cells.***P=0.0004 *P=0.0388 (**d**) Flow cytometry analysis of CD8α and CD8β expression on
the Vα3.2⁺QFL-Dex⁺ cells enriched from the siIEL compartment of naïve WT, Qa-1ᵇ-KO or
ERAP1-KO mice. Numbers in plots indicate average percentages of CD8αα⁺ cells. (**e**)
Percentages of CD8αα⁺ cells detected among Vα3.2⁺(as in **d**) or Vα3.2⁻ QFL T cells. **P=0.0082

*P=0.0335 Representative data are shown in flow plots (**b, d**) and the number of replicates is
specified in the bar graphs (**a, c, e**). Each symbol represents data collected from the indicated
tissue isolated from an individual mouse. Data from $\geqq 4$ independent experiments were
pooled for statistical analysis. P values were calculated with Student's *t* test. 'ns' indicates the
comparison was not significant.
813

814
Figure 4 Distinct functional features of QFLTg cells in the spleen and gut. (**a, c**)
Proliferation of splenic (**a**) or siIEL (**c**) QFLTg or OT-1 cells from QFLTg or OT-1 mice in
response to IL-15 (left) or a combination of IL-7 and IL-18 (right) stimulation. Cells were
tracked using Cell Tracing Violate (CTV). Numbers in plots indicate average percentages of
proliferated QFLTg cells. (**b**) Percentages of proliferated splenic QFLTg or OT-1 cells as
detected in **a**. ****P<0.0001 **P=0.0052 (**d**) Percentages of proliferated siIEL QFLTg or OT-1
as detected in **c**. ***P=0.0009 *P=0.0101 (**e**) Generation of QFLTg_WT or OT-1_WT partial
hematopoietic chimera mice. 2x10^6 of the bone marrow (BM) cells from GFP$^+$QFLTg or
GFP$^+$OT-1 mice were transferred into WT neonates at 3~5 days of age. The chimera mice were
analyzed at 8~10 weeks of age. (**f**) Gating strategy for the QFLTg cells originated from donor
bone marrow cells in QFLTg_WT chimera mice. Plots representing donor-derived QFLTg
population in the spleen being gated as GFP$^+$CD45$^+$CD19$^-$TCRβ$^+$CD4$^-$Vα3.2$^+$ cells. (**g, i**) Flow
cytometry measurement of IFN-γ production by QFLTg or OT-1 cells isolated from the spleen
(**g**) or siIEL compartment (**i**) of naïve chimera mice stimulated with or without 2μM FL9 or SL8
peptide respectively for 4.5h. Numbers in plots indicate average percentages of IFN-γ$^+$ cells. (**h**)
Percentages of IFN-γ$^+$ cells detected as in **g**. ****P<0.0001 ***P=0.0001 (**j**) Percentages of

IFN-γ⁺ cells detected as in **i**. ***P=0.0004 *P=0.0247 Representative data are shown in flow
plots (**a, c, f, g, i**) and the number of replicates is specified in the bar graphs (**b, d, h, j**). Data
from 2(**b, d**) or 3 (**h, j**) independent experiments were pooled for statistical analysis. Each
symbol represents data collected from the indicated tissue isolated from an individual mouse. P
values were calculated with Student's *t* test. 'ns' indicates the comparison was not significant.
836

837
Figure 5 Gut microbiota is associated with retention but not homing of QFL T cells in the
siIEL compartment. (**a, c**) Flow cytometry of QFL-Dex⁺ cells enriched from the spleen (**a**) or
siIEL compartment (**c**) of specific-pathogen-free (SPF) or germ-free (GF) WT mice at 10 or 18
weeks of age. Numbers in plots indicate absolute numbers of QFL-Dex⁺ cells. (**b**) Absolute
numbers (left) and frequencies (right) of QFL-Dex⁺ cells detected within the total splenic CD8⁺
population in SPF or GF WT mice of 8~17 weeks or 18~22 weeks of age.***P=0.009
****P<0.0001 (**d**) Absolute numbers and frequencies of QFL-Dex⁺ cells detected among the
total siIEL TCRβ⁺CD4⁻ population in SPF or GF WT mice of 8~17 weeks or 18~22 weeks of
age. **P=0.005 *P=0.037 Representative data are shown in flow plots (**a, c**) and the number of
replicates is specified in the bar graphs (**b, d**). Each symbol represents data collected from the
indicated tissue isolated from an individual mouse. Data from ≧5 independent experiments
were pooled for statistical analysis. P values were calculated with Student's *t* test. 'ns'
indicates the comparison was not significant.
851
852

853
Figure 6 Identification of a QFL T cross-reactive FL9 homolog peptide expressed in
commensal bacterium *P. pentosaceus*. (**a**) Structure model of the QFL TCR:FL9:Qa-1^b
complex predicted by AlphaFold2. Detailed view of the binding groove is shown in the right
panel. (**b**) Conformation alignment of FL9 (red) and Qdm (periwinkle). (**c**) Alanine screen of the
predicted key residues (P4 and P6) in FL9. Activation of QFL T cells was assessed by measuring
the LacZ response of BEko8Z hybridoma cells to the Qa-1^b-expressing Lmtk$^-$ (L-Qa-1^b) cells
pulsed with mutant FL9 peptides including FYAEAAPML (FL9-P6A) and FYAAATPML(FL9-
P4A) or the wild-type FYAEATPML(FL9) using spectrophotometry. (**d**) Workflow for
identification of commensal bacterial FL9 homolog peptide which cross-activate QFL T cells. (**e**)
Identification of the QFL T cross-reactive peptide FYAEDGTPIL(FL10). BEko8Z response to
the L-Qa-1^b cells or the non-Qa-1^b-expressing Lmtk$^-$ cells pulsed with 50nM of the FL9 or the
candidate homolog peptides. (**f**) Response of BEko8Z hybridoma to L-Qa-1^b cells pulsed with
FL9, FL10 or FYAEDDTPIV (FV10) peptide of various concentrations. (**c, e, f**) Data are from
one experiment representative of three independent experiments.
868

Figure 7 Association between Vα3.2⁺QFL T cells and the commensal bacterium *P. pentosaceus*. (**a**) Colonization of GF mice with commensal bacteria. GF QFLTg_WT chimera mice received oral gavage of *P. pentosaceus* at 8 weeks of age and were analyzed for both the donor-derived QFLTg cells and the endogenous Vα3.2⁺QFL T cells at 16 weeks of age. GF WT mice received oral gavage of *P. pentosaceus* or *L. johnsonii* at 8 and 10 weeks of age and were analyzed for the endogenous Vα3.2⁺QFL T cells at 16 weeks of age. (**b**) Absolute numbers of donor-derived QFLTg cells in the spleen (left) or siIEL compartment (right) of 16-week GF QFLTg_WT chimera mice colonized with *P. pentosaceus* in comparison with SPF or uncolonized GF chimera mice. Symbols on x-axis indicate that QFLTg cells were undetectable in the indicated group. *P<0.03 (**c**) Relative abundance of *P. Pentosaceus* or *L. Johnsonii* determined by whole shotgun metagenomic sequencing of microbiota composition in pooled fecal pallets collected from each cage. (**d**) Absolute numbers of endogenous Vα3.2⁺QFL T cells in the spleen (left) or siIEL compartment (right) of 16-week colonized GF mice compared with SPF or uncolonized GF chimera mice. Endogenous Vα3.2⁺QFL T data collected from the GF QFLTg_WT chimeric mice and the non-chimeric WT mice were pooled for this analysis.

886   **P<0.008 The number of replicates is specified in the bar graphs with each symbol representing
887   data collected from the indicated sample/tissue isolated from an individual cage (**c**) or mouse (**b**,
888   **d**). Data from ≧3 independent experiments were pooled for statistical analysis. P values
889   were calculated with Student's *t* test. 'ns' indicates the comparison was not significant.
890
891

Table 1. Characteristics of Vα3.2$^+$ QFL T cells in the spleen and IEL

Tissue	Cell type	Phenotype	Microbiota	Critical molecules
Spleen	Innate-like T	CD44^{hi}	Independent	Population establishment: Qa-1^b (partially required), TAP
IEL	Natural IELs	CD8αα$^+$ CD5$^-$ CD90$^-$	Homing: Independent Retention: Dependent	Phenotype imprinting: Qa-1^b, ERAP1 (partially required)

893   **Table 1.** Summary of the characteristic features of the splenic and IEL Vα3.2$^+$ QFL T cells

894   including their cell type, expression of cell surface markers, dependence on microbiota for tissue

895   homing/retention and the critical molecules required for population establishment or phenotype

896   imprinting.

897