1    Original Article

**2    The role of intraspecific crop-weed hybridization in the evolution of weediness and**

**3    invasiveness: cultivated and weedy radish (*Raphanus sativus* L.) as a case study**

Román B. Vercellino[1], Fernando Hernández[1,2], Alejandro Presotto[1]

[1]Departamento de Agronomía, CERZOS, Universidad Nacional del Sur (UNS)-CONICET,

San Andrés 800, 8000, Bahía Blanca, Argentina

[2]Department of Zoology and Biodiversity Research Centre, University of British Columbia,

Vancouver, British Columbia, Canada

**Correspondence:** R B Vercellino, Departamento de Agronomía, Universidad Nacional del

Sur, San Andrés 800, 8000, Bahía Blanca, Argentina. Tel: 0054 0291 4595102 (4345); E-

mail: rbvercellino@cerzos-conicet.gob.ar

**Short title:** Intraspecific crop-weed hybridization in *Raphanus sativus*

**ORCIDs**

Román B Vercellino          0000-0002-9313-3501

Fernando Hernández          0000-0001-5158-2029

Alejandro Presotto          0000-0002-9085-9673

**PREMISE:** Crop-wild/weed hybrids usually exhibit intermediate and maladapted

phenotypes compared to their parents; however, hybridization has sometimes been associated

with increased fitness, potentially leading to enhanced weediness and invasiveness. Since the

ecological context and maternal genetic effects may affect hybrid fitness, they could

influence the evolutionary outcomes of hybridization. Here, we evaluated the performance of

first-generation crop-weed hybrids of *Raphanus sativus* L. and their parents under two

contrasting ecological conditions.

**METHODS:** Using experimental hybridization and common garden experiments under field

conditions, we assessed the differences in time to flowering, survival to maturity, plant

biomass and reproductive components between bidirectional crop-weed hybrids and their

parents, under two contrasting ecological conditions, agrestal (wheat cultivation, fertilization,

weeding) and ruderal (human-disturbed uncultivated area) over two years.

**RESULTS:** Crop, weeds and bidirectional hybrids overlapped -at least partially- during the

flowering period, indicating a high probability of gene flow. Hybrids survived to maturity at

rates at least as successful as their parents and showed higher plant biomass and fecundity,

which resulted in higher fitness compared to their parents in both contrasting environments,

without any differences associated with the direction of the hybridization.

**CONCLUSIONS:** Intraspecific crop-weed hybridization, regardless of the cross direction,

has the potential to promote weediness in weedy *R. sativus* both in agrestal and ruderal

environments, increasing the chances of the introgression of crop alleles into weed

populations. This is the first report of intraspecific crop-weed hybridization in *R. sativus*.

**KEY WORDS:** Brassicaceae; crop-wild hybridization; gene flow; hybrid fitness; invasives;

weeds; admixture.

## INTRODUCTION

Hybridization is increasingly recognized as an important evolutionary phenomenon that can influence the evolution of both weeds and invasives (Ellstrand et al., 2010, 2013; Vercellino et al., 2023). When hybridization occurs, the hybrid lineages harbor alleles from both progenitors, and the process of introgression will depend on the fitness of the hybrids relative to their parents (Mercer et al., 2007; Presotto et al., 2019). Hybrids commonly exhibit intermediate and maladapted phenotypes and reduced fitness compared to their parents (Mercer et al., 2006; Sahoo et al., 2010; Hooftman et al., 2015; Martin et al., 2019; Yue et al., 2021), with nonviable or sterile offspring in extreme cases (Rieseberg et al., 2007). If novel alleles reduce hybrid fitness, these alleles may not prevail or introgress. However, hybridization can also lead to adaptive evolution through evolutionary novelty, increased genetic variation, hybrid vigor (i.e., heterosis) and dumping genetic load, increasing weediness and invasiveness (Schierenbeck and Ellstrand, 2009; Hovick and Whitney, 2014). Spontaneous hybridization between crop and their wild/weedy relatives has increased the weediness and invasiveness of existing weeds and invasives, e.g., weedy *Brassica rapa* (Ureta et al., 2017; Pandolfo et al., 2018b) and California wild radish (Snow et al., 2010), and has resulted in the evolution of new weeds and invasives, e.g., European weed beet (Bartsch, 2010), weedy rice (Vigueira et al., 2019), European teosinte (Le Corre et al., 2020) and weedy sunflower (Presotto et al., 2017; Hernández et al., 2022). In addition, in some cases, hybrid populations have replaced their progenitors locally (Hegde et al., 2006). Conversely, in many instances, hybridization has not affected weediness or invasiveness (Magnussen and Hauser, 2007; Whitney et al., 2009; Martin et al., 2019; Page et al., 2019).

Hybrid performance relative to their parents may be influenced by the environment (Campbell and Snow, 2007; Arnold and Martin, 2010; Hovick and Whitney, 2014; Presotto et al., 2019). Due to this environmental dependence, evolutionary outcomes of hybridization in

agrestal (i.e., agricultural) environments may differ from those in ruderal (human-modified

uncultivated area) habitats (Mercer et al., 2006; Presotto et al., 2019; Mitchell et al., 2021).

The fitness of wild/weeds could be enhanced by the acquisition of crop traits under

agricultural environments. This is especially true for traits conferring resistance to herbicides,

insects or diseases, or tolerance to abiotic stresses (Xia et al., 2016; Ureta et al., 2017;

Pandolfo et al., 2018b; Nam et al., 2020; Kreiner et al., 2022), but also for other typical crop

traits, such as early flowering, high growth rate and seed production (Campbell and Snow,

2007; Presotto et al., 2017; Le Corre et al., 2020). However, the acquisition of crop traits

which have been shaped to suit the agricultural environment by artificial selection, frequently

create maladapted crop-wild/weed hybrids with lower fitness under less selective

environments (e.g., ruderal environments), limiting the adaptive evolution of weeds and

invasives (Mercer et al., 2007; Hovick et al., 2012; Presotto et al., 2019; Vercellino et al.,

2021a).

Hybrid performance may also be influenced by the phenotypes of its maternal parent (known

as maternal genetic effects) (Kimball et al., 2008; Hernández et al., 2021). Maternal genetic

effects (hereafter maternal effects) are defined as the contribution of the phenotype or

genotype of the maternal parent to the phenotype of its offspring beyond the equal

chromosomal contribution expected from each parent (Roach and Wulff, 1987). Numerous

studies have investigated the ecological and evolutionary consequences of crop-to-wild/weed

hybridization (Campbell et al., 2006; Mercer et al., 2006, 2007; Snow et al., 2010; Hovick et

al., 2012; Presotto et al., 2019). However, much less attention has been paid to hybridization

in the opposite direction (i.e., wild/weed-to-crop hybridization) and how maternal genetic

effects may influence the fitness of their offspring immediately after hybridization (Kimball

et al., 2008; Piskurewicz et al., 2016; Singh et al., 2017; Hernández et al., 2021). If such

maternal effects exist, crop-weed hybrids produced on crop and weed plants will show

differences in performance. Common garden studies comparing the fitness of bidirectional

crop–wild/weed hybrids relative to both parents under contrasting ecological conditions are

essential to deepen our understanding on the role of crop-wild/weed hybridization on

weediness and invasiveness (Mercer et al., 2006; Campbell and Snow, 2007).

In this study, we performed common garden experiments using experimental populations of

the crop-weed complex of radish (*Raphanus sativus* L.) as a model system to assess the

hypothesis that bidirectional intraspecific crop-weed hybridization increases weediness and

invasiveness (Bourgeois et al., 2019). These studies allow the acceleration of natural

processes in order to explore local ecological scenarios under which the fitness of crop-weed

hybrids may equal or exceed the fitness of their parents, favoring the persistence of crop

alleles in wild/weedy populations and consequently enhancing weediness and invasiveness

(Mercer et al., 2006; Campbell et al., 2009). The *Raphanus* complex, formed by the

cultivated species *R. sativus*, their weedy conspecific forms and their weed related species *R.*

*raphanistrum*, is a well-established model system that has been widely used to evaluate the

ecological and evolutionary consequences of interspecific crop-weed hybridization

(Campbell et al., 2006, 2009; Hegde et al., 2006; Campbell and Snow, 2007; Ridley et al.,

2008; Snow et al., 2010; Hovick et al., 2012; Shukla et al., 2020, 2021). We compared the

time to flowering, survival to maturity, plant biomass and reproductive components of

bidirectional first-generation crop-weed hybrids in relation to both parents under two

contrasting ecological environments, agrestal (wheat cultivation, fertilization, weeding) and

ruderal (human-disturbed uncultivated area) over two years. Despite that the fitness of first-

generation hybrids does not determine the ultimate fate of all crop alleles, it affects the initial

course of introgression through its effect on the frequencies of crop alleles that are

subsequently subject to selection and drift (Snow et al., 1998; Mercer et al., 2006; Presotto et

al., 2019). Our work represents the first experimental study to evaluate the effect of

bidirectional crop-weed hybridization in *R. sativus* on the evolution of weediness and

invasiveness.

**2. MATERIALS AND METHODS**

*Study system-*

Conspecific cultivated and weedy radish (*Raphanus sativus* L., Brassicaceae) are annual, self-

incompatible, insect-pollinated plants, that can hybridize over long distances (Ellstrand et al.,

1989; Snow and Campbell, 2005). Cultivated radish is an ancient crop domesticated

independently in both Europe and South Asia (Zhang et al., 2021) and selected for multiple

purposes, including human consumption as roots (botanically hypocotyl plus root), forage

and cover crops (Snow and Campbell, 2005). *R. sativus* is not known in the wild (Warwick,

2011); however, spontaneous populations of radish have been found as a weed infesting crops

in several parts of the world, including North America, South America and South Africa

(Hegde et al., 2006; Barnaud et al., 2013; Vercellino et al., 2018; Costa et al., 2021; Heap,

2023). In North America, its invasive capacity is attributed to the introgression with a closely

related species *R. raphanistrum* (known as California wild radish) (Hegde et al., 2006;

Ellstrand et al., 2010), and in South America, it can be found in Argentina, Brazil, Chile,

Uruguay and Paraguay (Pandolfo et al., 2018a). In Argentina, weedy radish is present in 20

out of 23 provinces and has been considered a widespread invasive weed since at least the

1930s (Ibarra, 1937; Pandolfo et al., 2018a). It grows in disturbed environments, such as road

margins, rural roads and fence lines, and within agricultural fields. It is one of the most

noxious weeds in cereals, oilseeds and some horticultural and forage crops (Scursoni et al.,

2014; Vercellino et al., 2018, 2021b) and has evolved resistance to AHAS (acetohydroxyacid

synthase, also known as acetolactate synthase (ALS))-inhibiting herbicides (Pandolfo et al.,

2016; Heap, 2023). Argentine weedy *R. sativus* populations show many crop-like traits, such

as white or purple flowers, pods attached to the mother plant at maturity, and no seed

dormancy (Vercellino et al., 2018, 2019). On the other hand, weedy populations do not

harbor typical traits of *R. raphanistrum* like constricted fruits and yellow flowers (Pandolfo et

al., 2018a). However, they have an extended period of seedling emergence due to the

requirement of darkness for germination and the mechanical restriction of the pericarp

(Vercellino et al., 2019).

In southern Buenos Aires, an area severally infested with weedy radish (Scursoni et al., 2014;

Vercellino et al., 2018), the use of cultivated radish as horticultural and cover crop has grown

rapidly in recent years, leading to the formation of natural crop-weed hybrids in the field.

*Germplasm and crosses-*

In this study, bidirectional crosses between weed and cultivated *R. sativus* were used due to

the potential bidirectional hybridization in the *Raphanus* complex in Argentina. Weed

populations were collected from two sites in southern Buenos Aires: Balcarce (BAL;

37°35'25''S, 58°31'59''W) in 2008  and Pieres (PIE; 38°24'31''S, 58°35'60''W) in 2011

(Vercellino et al., 2018), before the introduction of *R. sativus* cover crop cultivars in those

areas. Balcarce and Pieres are temperate sub-humid environments, with an annual average

temperature of 13.9 and 14.3 °C (SMN 2023, https://www.smn.gob.ar/), respectively.

Cultivated materials were represented by two commercial cultivars: the European radish

'Rovi – Round Red Radish with White Tips' (Garde, Giusti y Chuchuy S.A., Ciudad

Autónoma de Buenos Aires, Argentina) cultivated for root consumption and the daikon

radish 'CCS 770 – Daikon Radish' (El Cencerro, Coronel Suarez, Buenos Aires, Argentina)

cultivated as cover crop. Radish is cultivated in urban and peri-urban farms and gardens

throughout the country, and as cover crop radish in the Pampas region and northern

Argentina, Paraguay and Brazil (Snow and Campbell, 2005). Bidirectional crosses were made

between weed populations and cultivars (referred to as CROP). The crosses (hereafter

referred to as Biotypes) were characterized according to their female and male parents.

To minimize the maternal environmental effects, all the seeds used in each experiment were

produced in a greenhouse (20 ± 5°C, natural light) in the Agronomy Department at

Universidad Nacional del Sur, Bahía Blanca, Argentina. For seed production, 60 plants of

each BAL, PIE, and CROP were grown in the greenhouse and seeds were produced under

controlled hand-pollination, creating seven different biotypes: three pure parents (BAL, PIE

and CROP) and bidirectional F1 crop-weed hybrids (BAL x CROP, PIE x CROP, CROP x

BAL and CROP x PIE). CROP was represented by the open-pollinated cultivar 'Rovi –

Round Red Radish with White Tips' in 2019, and by the cultivar 'CCS 779 Daikon Radish'

in 2020. In the greenhouse, the weedy and cultivated plants flowered simultaneously. Seeds

were produced as follows: at the flowering stage, unopened flower buds from weed plants

were opened and emasculated without damaging the pistil by using sharp-edged forceps

before pollen dehiscence. Once emasculated, the flowers were hand-pollinated with pollen

from sibling plants (weed biotypes: BAL and PIE) or pollen from cultivated plants (weed-

crop hybrids: BAL x CROP and PIE x CROP). Likewise, unopened flowers buds from

cultivated plants were opened, emasculated and hand-pollinated with pollen from sibling

plants (CROP biotype) or pollen from weed plants (crop-weed hybrids: CROP x BAL and

CROP x PIE). Prior to emasculation, all other flower buds on the racemes were removed.

Immediately after pollination, the racemes were covered with $12 \times 6$ cm pollination bags to

avoid any pollination from other plants. Pod formation from successful hand-pollination

could be observed ten days after pollination and, subsequently, the bags were removed from

the racemes. The effectiveness of pollination was >99 %. At the end of the growing season,

the mature pods were stored at room temperature and less than 10% moisture until use. For

experiments in the 2020 and 2021, seeds were produced during the 2019 and 2020 growing

seasons, respectively.

*Site and experimental design-*

We compared weed, crop, and their hybrids in two environments (ruderal and agrestal).

Common garden experiments were conducted in the Agronomy Department experimental

field at the Universidad Nacional del Sur, Bahía Blanca, Argentina (38°41´38" S, 62°14´53"

W), during two winter growing seasons (Jun-Jan 2020/21 and 2021/22, the typical life cycle

for both the crop and weedy radish). The site is characterized as semi-arid with a temperate

climate, and a mean annual temperature of 14.9 °C; the mean temperatures of the coldest

month (July) and the hottest month (January) are 7.6 and 23.0 °C, respectively. The average

annual precipitation is 641 mm (SMN 2023, https://www.smn.gob.ar/). The soil has a well-

drained loamy sand texture with 1.1% organic matter, pH 7.7, and 8.4 mg kg$^{-1}$ (0–12 cm)

available phosphorus (P) (a typical soil in the southwest of Buenos Aires province). The

experimental field was maintained as a natural field for at least two years prior to the start of

the experiment, leaving natural vegetation to grow. The site was governed by herbaceous

plants, mainly *Lolium spp.* and *Diplotaxis tenuifolia* in winter, and C*entaurea solstitialis,*

*Chenopodium album, Portulaca oleracea, Helianthus annuuss* subsp. *annuus, Cynodon*

*dactilon* and *Cenchrus incertus* in spring-early summer. One month prior to wheat sowing

(see below), a tillage operation with a disc harrow was used to prepare the soil. Then the

environments were defined. Field experiments were arranged in a split-plot design.

Environments (agrestal and ruderal) were assigned to main plots (one main plot per

environment, that is in total two main plots). Each main plot was divided to eight blocks, and

each block was divided in eight subplots. The seven biotypes (CROP, BAL, PIE, CROP x

BAL, CROP x PIE, BAL x CROP and PIE x CROP) plus empty control were assigned to

subplots within each block, whereby the assignment to subplot was randomized for each

block, environment and year. Each subplot contained ten plants of the same biotype (Appendix S1; see the Supplementary Data with this article). The whole experiment was repeated in two subsequent years.

**Appendix S1.** Experimental crosses and common garden design. Environments (main plots) are indicated in different colors, agrestal (yellow) and ruderal (green). Each environment was composed of eight blocks, each of eight subplots. Subplots were seven radish biotypes (1-BAL, 2-BAL x CROP, 3-CROP x BAL, 4-CROP, 5-CROP x PIE, 6-PIE x CROP, 7-PIE) and a control plot without radish (8). Numbers (1-8) on the field map represent each biotype or control position in year 1. In year 2, both environments were randomized again.

From the radish seeds produced in the crossing design described above, 2-3 seeds per pod

from more than 400 pods were pooled to represent each biotype. Since seeds from both weed

and cultivated radish are non-dormant (Vercellino et al., 2019), radish seeds from all seven

biotypes were hand-planted in early Jul 2020, 2-3 seeds per planting positions, 2-3 cm deep,

at 0.3-m intervals in rows spaced at 0.6-m. The subplots were two rows wide by five planting

positions long. Sixty-four subplots ((7 biotypes + 1 control without radish) x 8 blocks) made

up each main plot (i.e., each ruderal and agrestal environment). The main plots were

surrounded by a border of two rows of weedy radish to buffer against possible border effects.

Since most of the seeds germinated and emerged, at the fully expanded one-leaf stage, radish

seedlings were hand-thinned to one per position, reaching the target of 10 plants per subplot.

In 2021, since the experiment was established in a place where there was weedy radish three

years before and to avoid contamination from the seed bank, seedlings of the seven radish

biotypes were established in plastic trays containing potting mix (Grow Mix Terrafertil®),

grown in a glasshouse ($20 \pm 5°C$, natural light and automatic watering), and transplanted into

the experimental field at the 2-3-leaf stage, in early Aug, as above. Experiments were drip

irrigated, watered complementarily at the beginning to facilitate their establishment, and later

when necessary.

*Description of environments-*

*Agrestal environment-* Before the establishment of the radish biotypes in the field in late

autumn (mid-Jun), in the agrestal environment, a wheat (*Triticum aestivum* L.) crop was

established as follow: seeds were planted in 0.2-m spaced rows at 3-cm deep, to reach a target

of 160 plants $m^{-2}$ (the common stand density in southern Buenos Aires). Wheat rows were

arranged to surround rather than overlap the radish plants. The cultivars of wheat used were

ACA 360 and KLEIN MINERVA in 2020/21 and 2021/22, respectively. Fertilization

consisted of 120 kg $ha^{-1}$ diammonium phosphate at planting, 125 kg $ha^{-1}$ urea at early tillering

and 125 kg ha$^{-1}$ urea at mid-tillering. Weeds other than radish were periodically removed by

hand. At the end of each experiment, the yield of the wheat crop was estimated by hand-

harvesting and hand-threshing four replicates of 1 m$^2$, collected in subplots without radish.

The yield of the wheat was 5136 ± 201 and 5190 ± 188 kg ha$^{-1}$ in 2020/21 and 2021/22,

respectively.

*Ruderal environment-* In the ruderal environment, we did not establish any crops or fertilize

the soil. Biotypes were exposed to naturally occurring weather conditions, competing weeds

(i.e., same as described above), herbivores, and pathogens, simulating a human-disturbed

uncultivated area (e.g., field margin, fence row).

***Measurements-***

In all ten plants per subplot, we recorded daily time to flowering (days from emergence to

first flowering) and survival to reproduction (defined as plants that produced at least one

pod). At maturity, the above-ground biomass of the six central radish plants from each

subplot was hand-harvested, dried at 60°C to constant weight and weighed to determine the

total aboveground dry biomass per plant (hereafter referred to as 'plant biomass'). After that,

pod biomass per plant, pod number per plant, seed biomass per plant, seed number per plant

and seed weight were evaluated in the harvested plants following the procedure described in

Vercellino et al. (2018, 2021b). Seed number per plant was used as the proxy of fecundity.

Since all ten plants per subplot were used to evaluate survival to maturity, but only the six

central plants to estimate fecundity, the mean fitness of each subplot was estimated by

multiplying survival to maturity by the average fecundity of each subplot in order to calculate

the fitness of bidirectional crop-weed hybrids relative to weed plants (hereafter referred to as

'relative fitness'). Relative fitness (RF) was calculated as: RF = bidirectional hybrid (crop-

weed or weed-crop)/weed for each block and weed origin (BAL and PIE).

*Statistical Analysis-*

All data were analyzed using generalized linear mixed models (GLMMs) with restricted

maximum likelihood in PROC GLIMMIX (SAS University Edition). We analyze the effects

of experimental factors in the time to flowering, survival to reproduction, total aboveground

biomass, and reproductive components (i.e., pod biomass per plant, pod number per plant,

seed biomass per plant, seed number per plant and seed weight). For each year (2020/21 and

2021/22), environment (ruderal and agrestal), biotype (CROP, BAL, PIE, CROP x BAL,

CROP x PIE, BAL x CROP and PIE x CROP), and its interaction (environment x biotype)

were considered as fixed effects, while the block nested within environment effect was

considered as random. After that, we grouped the seven biotypes into four groups: WEED

(BAL and PIE), WEED x CROP (crop-weed hybrids produced on weed plants: BAL x CROP

and PIE x CROP), CROP x WEED (crop-weed hybrids produced on crop plants: CROP x

BAL and CROP x PIE), and CROP. To compare groups (WEED, WEED x CROP, CROP x

WEED, and CROP), we constructed six a priori orthogonal contrasts: (1) WEED vs. CROP x

WEED (2) WEED vs. WEED x CROP, (3) WEED vs. CROP, (4) CROP x WEED vs. WEED

x CROP, (5) CROP x WEED vs. CROP, and (6) WEED x CROP vs. CROP, and compare

them using the CONTRAST statement in PROC GLIMMIX, SAS.

Since categorical analyses do not allow for the complex model specifications needed in an

analysis of a split-plot design, survival to reproduction was analyzed as a continuous response

variable, based on averages of radish plants in each subplot. Data were arcsine-square-root

transformed prior to analyses to meet assumptions of normality (Mercer et al., 2006).

Flowering time analysis only included radish plants that reached flowering, and plant

biomass and reproductive component (pod biomass per plant, pod number per plant, seed

biomass per plant, seed number per plant and seed weight) analyses only included radish

plants that survived to reproduction. To avoid redundancy, we performed a multiple

Pearson´s correlation analysis between plant biomass and the reproductive components with

PROC CORR in SAS, and for all pairwise comparisons with r > |0.9|, and we only retained

one variable of each pair. After this filtering, we retained seed number per plant and seed

weight (Appendix S2). All traits (i.e., except survival to reproduction which was previously

indicated) were square-root transformed to satisfy assumptions regarding the normality of the

residuals. Finally, least square means of relative fitness were estimated using PROC

GLIMMIX.

**RESULTS**

*Time to flowering-*

In year 1, there were significant biotype, environment, and biotype x environment interaction

effects; therefore, we performed contrasts for each environment (agrestal and ruderal) (Table

1). In both environments, the crop flowered later than the weeds, although differences were

larger in the ruderal environment (Fig. 1a, Table 1). Overall, hybrids showed a flowering time

intermediate to that of both parents. Whereas the crop-weed hybrids produced on weed plants

flowered simultaneously with the weeds, the crop-weed hybrids produced on crop plants

flowered a little bit later (<5 days) than the weeds, resulting in significant differences

between bidirectional hybrids only in the agrestal environment (Fig. 1a, Table 1). In year 2,

there were significant biotype and environment effects, but no biotype x environment

interaction (Table 2). Crop, weeds and hybrids flowered simultaneously (Fig. 1b), although

some contrasts showed statistically significant (<3 days) differences between groups (Fig. 1b,

Table 2).

*Survival to maturity-*

In year 1, there were significant biotype, environment, and biotype x environment interaction

effects; therefore, we performed contrasts for each environment (agrestal and ruderal) (Table

1). In the agrestal environment, all the weed and hybrid plants survived to maturity, whereas

the crop survived significantly less than the weeds and hybrids (Figure 1c, Table 1). In

contrast, in the ruderal environment, there was more variation between groups; hybrids (both

cross directions) survived significantly more than weeds, and weeds survived significantly

more than the crop (Figure 1c, Table 1). There were no significant differences between

bidirectional hybrids (Figure 1a, Table 1). In year 2, most plants (> 95 %) survived to

maturity and there were no significant effects of environments, biotypes, nor environment x

biotype interaction (Figure 1b, Table 2).

*Fecundity: reproductive components-*

For the set of six variables (plant biomass, pod biomass per plant, pod number per plant, seed

biomass per plant, seed number per plant and seed weight), we found positive correlations

between seed number per plant and plant biomass (r = 0.95), pod number per plant (r = 0.93),

pod biomass per plant (r = 0.96) and seed biomass per plant (r = 0.98), while the seed weight

showed no correlation (r < 0.10) with any of these traits (Appendix S2). Therefore, seed

number per plant and seed weight were retained for further analysis.

**Appendix S2.** Correlations among plant biomass and reproductive components of weedy radish.

| Traits | Biomass plant$^{-1}$ | Pod biomass plant$^{-1}$ | Pod number plant$^{-1}$ | Seed biomass plant$^{-1}$ | Seed weight | Seed number plant$^{-1}$ |
|---|---|---|---|---|---|---|
| Biomass plant$^{-1}$ | 1 | **0.986** | **0.942** | **0.958** | 0.074 | **0.952** |
| Pod biomass plant$^{-1}$ | | 1 | **0.943** | **0.972** | 0.092 | **0.957** |
| Pod number plant$^{-1}$ | | | 1 | **0.921** | 0.015 | **0.934** |
| Seed biomass plant$^{-1}$ | | | | 1 | 0.088 | **0.979** |
| Seed weight | | | | | 1 | -0.009 |
| Seed number plant$^{-1}$ | | | | | | 1 |

Correlation coefficients are reported. The pairwise correlations with r > |0.9| are in bold. N> 1000 for
each combination.

Analysis of the seed number per plant (i.e., fecundity) revealed significant biotype and

environment effects, but no significant biotype x environment interaction in either year

(Tables 1 and 2). Overall, all plants in the agrestal environment showed 84.3 and 60.3 %

higher fecundity than the radish plants in the ruderal environment in year 1 and 2,

respectively (Figs. 1g-h, Tables 1 and 2). Fecundity of weeds was 167.5 % higher than crop

in year 1, but no differences were observed in year 2 (Figs. 1g-h, Tables 1 and 2). Fecundity

of the crop-weed hybrids were 47.0 and 169.0 % higher than fecundity of the weeds in year 1

and 2, respectively, with no differences associated with the cross direction, indicating no

maternal genetic effect (Figs. g-h, Table 1).

With respect to the seed weight, there were significant biotype and environment (only in year

1) effects, and no significant biotype x environment interaction (Tables 1 and 2). In year 1,

the seeds of the weed plants were 21.3 % heavier than the seeds of the crop plants. However,

in year 2, the seeds of the crop plants were 33.4 % heavier than the seeds of the weed plants

(Figs. 1e-f, Tables 1 and 2). In both years, the seeds of crop-weed hybrids (regardless of the

cross direction) were as heavy as the seeds of the parent with heavier seeds (Figs. 1e-f, Tables

1 and 2).

*Relative fitness-*

In both years, analysis of the relative fitness revealed a significant biotype effect but no

significant environment or biotype x environment interaction effects (Tables 1 and 2). The

relative fitness of hybrids compared to weeds was $1.86 \pm 0.21$ and $3.11 \pm 0.20$ in year 1 and

2, respectively, regardless of the environment and the direction of hybridization (Tables 1 and

2).

**Figure 1.** Boxplots of time to flowering (2020: A; 2021: B), survival to reproduction (2020: C; 2021: D), seed weight (2020/21: E; 2021/22: F) and seed number per plant (2020: G; 2021: H) for bidirectional crop-weed hybrids and their parents (WEED: BAL and PIE, CROP) of *Raphanus sativus* growing in two contrasting environments (ruderal and agrestal), and in two years. Box edges represent the 0.25 and 0.75 quartiles, solid line represents the median value, and whiskers extend to the minimum and maximum value or 1.5x the interquartile range with points for more extreme values. Black points represent outliers, and small and transparent points represent each data point. Values sharing the same letter within an environment and year are not significantly different based on contrast tests, and different letters within a panel indicate significant differences as generated by the contrasts. Note the variation in the scale of the y-axis between (H) and (G) to show large differences in seed number per plant between year 1 and 2.

**DISCUSSION**

This study shows that crop-weed hybridization has the potential to stimulate weediness in weedy *R. sativus* under both agrestal and ruderal environments. First-generation crop-weed hybrids showed hybrid vigor in all measured traits; they survived at rates that are at least as successful as the rates of both weed and crop plants and produced higher plant biomass that resulted in 47 and 169 % more fecundity than the weeds in year 1 and 2, respectively, in both ruderal and agrestal environments. Relative fitness of hybrids compared to weeds was 1.86 and 3.11 in year 1 and 2, respectively, with no differences between environments. Seeds of hybrids were as heavy as the seeds of the parental biotype with heavier seeds. Finally, we found no differences associated with the direction of hybridization, indicating the general absence of maternal genetic effects, which also supports hybrid vigor as the main genetic mechanisms of increased fitness. Our study highlights the risk of increasing weediness after hybridization in *R. sativus*, and suggests hybrid vigor as the genetic mechanisms increasing fitness of hybrid progeny.

*Evolutionary consequences of crop-weed hybridization-*

In our experiments, cultivated radish started flowering later than the weedy radish in year 1, but both cultivated and weedy radish started flowering simultaneously in year 2. This is probably associated with genetic differences between the radish cultivars. In year 1, we expected that the crop-weed hybrids would have delayed flowering compared to weedy radish, as was observed in California wild radish (hybrids between *R. sativus* and *R. raphanistrum*) (Campbell et al., 2006, 2009). While we found a trend in this direction (especially in the crop-weed hybrids produced on crop plants) (Fig. 1a), the difference was less than five days. Due to the long flowering period of the species, the flowering period of cultivated and weedy radish largely overlapped in both years in the field and in the

greenhouse during seed production (data not shown), suggesting that crop-weed hybridization is highly probable if crop and weed grow in sympatry.

Common garden studies using experimental early generation crop-weed hybrids can provide tentative predictions of the effects of hybridization on the dynamics of weed populations and the likelihood of crop allele introgression into weeds (Campbell et al., 2006; Mercer et al., 2006; Presotto et al., 2019). These studies allowed us examining effects of crop-weed hybridization on early and later life history traits over environmental conditions that realistically represent those at interfaces between crop and weed populations (Mercer et al., 2006; Presotto et al., 2019). In contrast to the common assumption (based on previous results) that early generations of crop-wild/weed hybrids suffer dramatic reductions in fitness relative to their wild/weedy counterpart (Snow et al., 1998; Cummings et al., 2002; Mercer et al., 2006; Campbell and Snow, 2007; Sahoo et al., 2010; Hooftman et al., 2015), our results showed at least equal survival and vastly higher plant biomass and fecundity in the hybrids relative to their parents in both contrasting environments and years. The increased fitness of first-generation hybrids is expected to facilitate the introgression of crop alleles into weedy populations and stimulate the evolution of weediness (Schierenbeck and Ellstrand, 2009; Vercellino et al., 2023). Similarly, potential for introgression from cultivated to their wild or weedy counterpart has been found in other similar crop-wild/weedy complexes, such as wild beet (Bartsch, 2010), wild lettuce (Uwimana et al., 2012), wild chicory (Kiær et al., 2007), wild carrot (Magnussen and Hauser, 2007), weedy *Brassica rapa* (Hauser et al., 2003) and wild radish (Hegde et al., 2006).

The assessment of the effect of crop-weed hybridization under contrasting ecological conditions was used to evaluate the environmental dependence of hybridization effects. In previous studies, the weed *R. raphanistrum* showed higher fecundity than the crop-weed hybrids under non-competitive conditions, but the crop-weed hybrids were less negatively

affected under intense competition, suggesting that hybrids may be favored under the highly

competitive condition imposed by crops in agricultural environments (Campbell and Snow,

2007; Campbell et al., 2009). Similarly, wild sunflower (*Helianthus annuus*) showed higher

fecundity than crop-wild hybrids in four contrasting environments, but the relative fitness of

hybrids was higher under competitive conditions, possibly because wild sunflower is not

adapted to agricultural environments (Mercer et al., 2006, 2007). Contrary to our expectation,

our results showed higher plant biomass and fecundity and consequently higher relative

fitness in the hybrids compared to their parents in both contrasting environments, indicating

that the result of the crop-weed hybridization was not environmental dependent, and that

bidirectional crop-weed hybridization has the potential to increase weediness in a wide range

of environmental conditions (e.g., both inside and outside the agricultural environment).

However, our inference should be studied further by evaluating the effect of crop-weed

hybridization across a greater diversity of locations in order to assess the generality of these

results.

Comparison of the direction of crop-weed hybridization was associated with the fact that

bidirectional crop-weed hybridization may contribute to the evolution of weediness in weedy

radish (Ridley et al., 2008), and previous studies have shown differential maternal genetic

effects when evaluating crop-wild/weed hybridization (Pace et al., 2015; Singh et al., 2017;

Hernández et al., 2021). Contrary to our expectation, the evaluated traits were not generally

influenced by the genotype of the maternal parent, suggesting similar evolutionary

consequences regardless of the direction of the hybridization. Nonetheless, considering that

radish was selected for larger root and leaf size -unlike crops selected for grain production

such as corn, sunflower and rice- and the lack of substantial differences between crop and

weedy radish, the absence of maternal genetic effects is not outlandish. To our knowledge,

this is the first study that has evaluated plants of bidirectional crop-weed hybrids in a common garden under field conditions.

A plausible explanation for the increased plant biomass and fecundity of hybrids is heterosis (Ellstrand and Schierenbeck, 2000). Heterosis, defined as the superiority of hybrids relative to their parents, is expected to occur after admixture between divergent populations that experienced genetic bottlenecks (Hovick et al., 2012; van Kleunen et al., 2015). Weedy radish populations from Argentina probably experienced strong genetic bottlenecks, first as a result of the species domestication, followed by species introduction into Argentina, and then as result of the feralization process (Vercellino et al., 2023). However, if current cultivars and weedy populations are divergent enough to explain heterosis effects observed in this study is unknown. Genetic studies are needed to better understand the evolutionary history of weedy radish in Argentina. Additional studies using weed-weed and crop-crop hybridization would provide control crosses where there is mixing between populations, but no specific crop-weed hybridization, and help us to disentangle the potential mechanisms of how crop-weed hybridization could lead to hybrid vigor and adaptive evolution. Theory predicts that heterosis of early generation hybrids is only transitory and it would decrease in the following generations (Rius and Darling, 2014); however, previous studies have shown that hybrid vigor could be maintained or fixed in subsequent generations (Ellstrand and Schierenbeck, 2000). For example, the superiority of interspecific crop-weed (*R. sativus* x *R. raphanistrum*) hybrids was maintained at least after four generations and under several environmental conditions (Campbell et al., 2006, 2009; Campbell and Snow, 2007; Snow et al., 2010; Hovick et al., 2012). Similar results were found in natural interspecific crop-weed radish hybrids (Hegde et al., 2006; Ridley and Ellstrand, 2009). However, additional studies should be carried out to understand the long-term consequences of intraspecific hybridization events (Campbell et al., 2009; Snow et al., 2010; Mercer et al., 2014; Mitchell et al., 2019).

A possible limitation of our study of bidirectional crop-weed hybridization is that only two

crop cultivars and two weed populations were used to create the biotypes. Despite this, we

did not find significant differences in any trait as a result of the sources of the weed

population, which could somehow support the generality of our results. On the other hand,

hybrid fitness may be influenced by factors that were not considered in our common garden

experiments, e.g., seed dormancy and seedling emergence (e.g., Hooftman et al. 2015;

Mitchell et al. 2019). However, fecundity is the most important trait contributing to hybrid

population growth in the *Raphanus* complex of North America (Campbell et al., 2014, 2016;

Teitel et al., 2016). Future studies incorporating more life-history traits would be necessary in

order to better understand the result of crop-weed hybridization. Finally, our discussion and

conclusions are based on field experiments representing the two main ecological conditions

that progenitors originally colonize and where bidirectional hybridization can occur;

however, while we adequately tested the effect of crop-weed hybridization within each

environment, comparisons between environments within year should be carried out carefully

as they were not replicated.

*Implications for crop and weed management-*

The assessment of the effect of intraspecific crop-weed hybridization under contrasting

ecological conditions was initially motivated by concerns about the risk of the evolution of

weediness after crop-weed hybridization. Here, we discuss the implications of our results for

weed evolution and management.

In recent years, *R. sativus* has been included as a cover crop in agricultural systems from

several parts of the world (Lawley et al., 2011). In the Argentine Pampas region, radish cover

crop is currently cultivated for both seed production and in farm fields in the same areas

where weedy radish has been one of the most problematic weeds for more than 80 years

(Ibarra, 1937; Scursoni et al., 2014). Considering that *R. sativus* is a self-incompatible, insect-

pollinated species (Snow and Campbell, 2005) with high gene flow over long distances

(Ellstrand et al., 1989), crop-weed hybridization is expected in areas where cultivated and

weedy radish are sympatric. Therefore, the concern regarding crop-weed hybridization and

the subsequent evolution of weediness is not whether it can happen, but the speed with which

humans accelerate this evolutionary process (reviewed in Vercellino et al. 2023).

A management practice to prevent crop-weed gene flow could be the suppression of the

cultivated radish prior to the flowering stage. Since this practice could only be carried out in

farmer fields, but not in seed production areas where the risk of dispersion is considerably

higher (e.g., weedy beet in Europe; Bartsch 2010), we propose that the introduction of

cultivated radish in areas with weedy radish should be carried out with extreme care, i.e., by

planting cultivated radish in weedy radish-free plots and by controlling weedy radish plants

in its surroundings. Considering that weed evolution after the introduction of a new crop in a

new area has been demonstrated in several crop-weed complexes, with multiple negative

impacts (reviewed in Ellstrand et al. 2010; Vercellino et al. 2023), we recommend a thorough

agroecological and environmental risk assessment of the introduction of a new crop (e.g.,

radish cover crop) before being approved for production and commercialization. In addition,

integrated weed management strategies aimed at preventing gene flow with the crop should

be a priority in order to prevent the emergence and spread of novel more problematic weedy

radish biotypes (reviewed in Vercellino et al., 2023).

Finally, crop-weed hybridization not only has negative consequences for maintaining crop

genetic uniformity and the increase in weediness, but weedy radish and hybrid plants could

also be used as genetic resources to improve crops, including crops more tolerant to biotic

and abiotic stress or more locally adapted (Wu et al., 2021; Mabry et al., 2023). In our case,

higher biomass and fitness of the hybrids could be interesting for the production of improved

cultivated radish both for cover crops or fodder, with higher production of dry matter.

However, the risk of fertilization of new cultivars should be taken into account.

**CONCLUSION**

In summary, our results using experimental populations under field conditions in a common

garden show that crop-weed hybridization has the potential to stimulate weediness and

invasiveness in the invasive weedy *R. sativus* in both ruderal and agrestal environments,

regardless of the direction of hybridization. This could promote the introgression and spread

of crop alleles -including alleles with resistance to herbicides, insects, pathogens, viruses and

tolerance to abiotic stresses (genetically modified or not)- into weed populations, promoting

the evolution of novel weed biotypes. We suggest that subsequent long-term full life-cycle

experiments in multiple locations are necessary to assess the generality of our results. Further

research should also study the phenotypic variation of functional traits on which selection can

act, as well as the strength and direction of direct and indirect selection operating on them

using phenotypic selection analysis to better understand the process of adaptive evolution that

may accompany the evolution of weeds and invasives.

**ACKNOWLEDGMENTS**

This work was supported by the Agencia Nacional de Promoción Científica y Tecnológica

(ANPCyT) PICT 2017-0473. We thank the National Research Council of Argentina

(CONICET) for a fellowship for FH and RBV, and two anonymous reviewers for their

helpful feedback that greatly improved the manuscript.

**LITERATURE CITED**

Arnold, M. L., and N. H. Martin. 2010. Hybrid fitness across time and habitats. *Trends in*

*Ecology and Evolution* 25: 530–536.

Barnaud, A., J. M. Kalwij, M. A. McGeoch, and B. J. van Vuuren. 2013. Patterns of weed

invasion: Evidence from the spatial genetic structure of *Raphanus raphanistrum*.

*Biological Invasions* 15: 2455–2465.

Bartsch, D. 2010. Gene flow in sugar beet. *Sugar Tech* 12: 201–206.

Bourgeois, B., F. Munoz, G. Fried, L. Mahaut, L. Armengot, P. Denelle, J. Storkey, et al.

2019. What makes a weed a weed? A large-scale evaluation of arable weeds through a

functional lens. *American Journal of Botany* 106: 90–100.

Campbell, L. G., and A. a. Snow. 2007. Competition alters life history and increases the

relative fecundity of crop-wild radish hybrids (*Raphanus* spp.). *New Phytologist* 173:

648–660.

Campbell, L. G., A. A. Snow, and C. E. Ridley. 2006. Weed evolution after crop gene

introgression: Greater survival and fecundity of hybrids in a new environment. *Ecology

Letters* 9: 1198–1209.

Campbell, L. G., A. A. Snow, and P. M. Sweeney. 2009. When divergent life histories

hybridize: Insights into adaptive life-history traits in an annual weed. *New Phytologist*

184: 806–818.

Campbell, L. G., Z. Teitel, and M. N. Miriti. 2016. Contemporary evolution and the dynamics

of invasion in crop-wild hybrids with heritable variation for two weedy life-histories.

*Evolutionary Applications* 9: 697–708.

Campbell, L. G., Z. Teitel, M. N. Miriti, and A. A. Snow. 2014. Context-specific enhanced

invasiveness of *Raphanus* crop–wild hybrids: A test for associations between greater

fecundity and population growth. *Canadian Journal of Plant Science* 94: 1315–1324.

Le Corre, V., M. Siol, Y. Vigouroux, M. I. Tenaillon, and C. Délye. 2020. Adaptive

introgression from maize has facilitated the establishment of teosinte as a noxious weed

in Europe. *Proceedings of the National Academy of Sciences* 117: 25618–25627.

Costa, L. O., J. Cechin, M. A. Rizzardi, S. L. Martin, C. A. Sauder, F. S. Adegas, and M. F.

Grando. 2021. Target-site resistance and cross-resistance to ALS-inhibiting herbicides in

radish and wild radish biotypes from Brazil. *Agronomy Journal* 113: 236–249.

Cummings, C. L., H. M. Alexander, A. A. Snow, L. H. Rieseberg, M. J. Kim, and T. M.

Culley. 2002. Fecundity selection in a sunflower crop-wild study: Can ecological data

predict crop allele changes? *Ecological Applications* 12: 1661–1671.

Ellstrand, N. C., B. Devlin, and D. L. Marshall. 1989. Gene flow by pollen into small

populations: Data from experimental and natural stands of wild radish. *Proceedings of

the National Academy of Sciences* 86: 9044–9047.

Ellstrand, N. C., S. M. Heredia, J. A. Leak-Garcia, J. M. Heraty, J. C. Burger, L. Yao, S.

Nohzadeh-Malakshah, and C. E. Ridley. 2010. Crops gone wild: Evolution of weeds and

invasives from domesticated ancestors. *Evolutionary Applications* 3: 494–504.

Ellstrand, N. C., P. Meirmans, J. Rong, D. Bartsch, A. Ghosh, T. J. de Jong, P. Haccou, et al.

2013. Introgression of crop alleles into wild or weedy populations. *Annual Review of

Ecology, Evolution, and Systematics* 44: 325–345.

Ellstrand, N. C., and K. A. Schierenbeck. 2000. Hybridization as a stimulus for the evolution

of invasiveness in plants? *Proceedings of the National Academy of Sciences* 97: 7043–

7050.

Exposito-Alonso, M., A. C. Brennan, C. Alonso-Blanco, and F. X. Picó. 2018. Spatio-

temporal variation in fitness responses to contrasting environments in *Arabidopsis

thaliana*. *Evolution* 72: 1570–1586.

Hauser, T. P., C. Damgaard, and R. B. Jørgensen. 2003. Frequency-dependent fitness of

hybrids between oilseed rape (*Brassica napus*) and weedy *B. rapa* (Brassicaceae).

*American Journal of Botany* 90: 571–578.

Heap, I. 2022. The International Survey of Herbicide Resistant Weeds. Website

www.weedscience.org.

Hegde, S. G., J. D. Nason, J. M. Clegg, and N. C. Ellstrand. 2006. The evolution of

California's wild radish has resulted in the extinction of its progenitors. *Evolution* 60:

1187–1197.

Hernández, F., R. B. Vercellino, I. Fanna, and A. Presotto. 2021. Maternal control of early

life history traits affects overwinter survival and seedling phenotypes in sunflower

(*Helianthus annuus* L.). *Plant Biology* 23: 307–316.

Hernández, F., R. B. Vercellino, C. Pandolfo, J. R. Mandel, and A. Presotto. 2022. Rapid

evolution of seed dormancy during sunflower de-domestication. *Journal of Heredity*

113: 288–297.

Hooftman, D. A. P., J. M. Bullock, K. Morley, C. Lamb, D. J. Hodgson, P. Bell, J. Thomas,

and R. S. Hails. 2015. Seed bank dynamics govern persistence of *Brassica* hybrids in

crop and natural habitats. *Annals of Botany* 115: 147–157.

Hovick, S. M., L. G. Campbell, A. A. Snow, and K. D. Whitney. 2012. Hybridization alters

early life-history traits and increases plant colonization success in a novel region. *The*

*American Naturalist* 179: 192–203.

Hovick, S. M., and K. D. Whitney. 2014. Hybridisation is associated with increased fecundity

and size in invasive taxa: Meta-analytic support for the hybridisation-invasion

hypothesis. *Ecology Letters* 17: 1464–1477.

Ibarra, F. 1937. Malezas más comunes del trigo y del lino. *Almanaque del Ministerio de Agricultura*.: 405–410.

Kiær, L. P., M. Philipp, R. B. Jorgensen, and T. P. Hauser. 2007. Genealogy , morphology and fitness of spontaneous hybrids between wild and cultivated chicory (*Cichorium intybus*). *Heredity* 99: 112–120.

Kimball, S., D. R. Campbell, and C. Lessin. 2008. Differential performance of reciprocal hybrids in multiple environments. *Journal of Ecology* 96: 1306–1318.

van Kleunen, M., M. Röckle, and M. Stift. 2015. Admixture between native and invasive populations may increase invasiveness of *Mimulus guttatus*. *Proceedings of the Royal Society B: Biological Sciences* 282.

Kreiner, J. M., S. E. Latorre, H. A. Burbano, J. R. Stinchcombe, S. P. Otto, D. H. Weigel, and S. I. Wright. 2022. Rapid weed adaptation and range expansion in response to agriculture over the past two centuries. *Science* 378: 1079–108.

Lawley, Y. E., R. R. Weil, and J. R. Teasdale. 2011. Forage radish cover crop suppresses winter annual weeds in fall and before corn planting. *Agronomy Journal* 103: 137–144.

Mabry, M. E., M. V Bagavathiannan, J. M. Bullock, A. L. Caicedo, C. J. Dabney, H. Wang, E. B. M. Drummond, et al. 2023. Building a feral future : Open questions in crop ferality. *Plants People Planet*. DOI: https://doi.org/10.1002/ppp3.10367

Magnussen, L. S., and T. P. Hauser. 2007. Hybrids between cultivated and wild carrots in natural populations in Denmark. *Heredity* 99: 185–192.

Martin, S. L., B. E. Lujan-Toro, C. A. Sauder, T. James, S. Ohadi, and L. M. Hall. 2019. Hybridization rate and hybrid fitness for *Camelina microcarpa* Andrz. ex DC (♀) and *Camelina sativa* (L.) Crantz (Brassicaceae) (♂). *Evolutionary Applications* 12: 443–455.

Mercer, K. L., D. A. Andow, D. L. Wyse, and R. G. Shaw. 2007. Stress and domestication

traits increase the relative fitness of crop-wild hybrids in sunflower. *Ecology Letters* 10:

383–393.

Mercer, K. L., D. J. Emry, A. A. Snow, M. A. Kost, B. A. Pace, and H. M. Alexander. 2014.

Fitness of crop-wild hybrid sunflower under competitive conditions: Implications for

crop-to-wild introgression. *PLoS ONE* 9: e109001.

Mercer, K. L., D. L. Wyse, and R. G. Shaw. 2006. Effects of competition on the fitness of

wild and crop-wild hybrid sunflower from a diversity of wild populations and crop lines.

*Evolution* 60: 2044–2055.

Mitchell, N., S. A. Chamberlain, and K. D. Whitney. 2021. Proximity to crop relatives

determines some patterns of natural selection in a wild sunflower. *Evolutionary*

*Applications* 14: 1328–1342.

Mitchell, N., G. L. Owens, S. M. Hovick, L. H. Rieseberg, and K. D. Whitney. 2019.

Hybridization speeds adaptive evolution in an eight-year field experiment. *Scientific*

*Reports* 9: 6746.

Nam, K. H., D. Y. Kim, Y. S. Moon, I. S. Pack, S. C. Jeong, H. B. Kim, and C. G. Kim.

2020. Performance of hybrids between abiotic stress-tolerant transgenic rice and its

weedy relatives under water-stressed conditions. *Scientific Reports* 10: 9319.

Pace, B. A., H. M. Alexander, J. D. Emry, and K. L. Mercer. 2015. Seed fates in crop-wild

hybrid sunflower: Crop allele and maternal effects. *Evolutionary Applications* 8: 121–

132.

Page, A., J. Gibson, R. S. Meyer, and M. A. Chapman. 2019. Eggplant domestication:

Pervasive gene flow, feralization, and transcriptomic divergence. *Molecular Biology and*

*Evolution* 36: 1359–1372.

Palacio-Lopez, K., and J. Molofsky. 2021. Phenotypic shifts following admixture in

recombinant offspring of *Arabidopsis thaliana*. *Evolutionary Ecology* 35: 575–593.

Pandolfo, C. E., A. Presotto, and M. Cantamutto. 2018a. *Raphanus sativus* L. - *Raphanus*

*raphanistrum* L. In Fernández, O. A., E. S. Leguizamón, and H. A. Acciaresi. Malezas e

Invasoras de la Argentina, EdiUNS, Bahía Blanca, Argentina, pp 625–636.

Pandolfo, C. E., A. Presotto, F. T. Carbonell, S. Ureta, M. Poverene, and M. Cantamutto.

2018b. Transgene escape and persistence in an agroecosystem: the case of glyphosate-

resistant *Brassica rapa* L. in central Argentina. *Environmental Science and Pollution*

*Research* 25: 6251–6264.

Pandolfo, C. E., A. Presotto, F. Moreno, I. Dossou, J. P. Migasso, E. Sakima, and M.

Cantamutto. 2016. Broad resistance to acetohydroxyacid-synthase-inhibiting herbicides

in feral radish (*Raphanus sativus* L.) populations from Argentina. *Pest Management*

*Science* 72: 354–361.

Piskurewicz, U., M. Iwasaki, D. Susaki, C. Megies, T. Kinoshita, and L. Lopez-Molina.

2016. Dormancy-specific imprinting underlies maternal inheritance of seed dormancy in

*Arabidopsis thaliana*. *eLife* 5: e19573.

Presotto, A., F. Hernández, M. Díaz, I. Fernández-Moroni, C. Pandolfo, J. Basualdo, S.

Cuppari, et al. 2017. Crop-wild sunflower hybridization can mediate weediness

throughout growth-stress tolerance trade-offs. *Agriculture, Ecosystems and Environment*

249: 12–21.

Presotto, A., F. Hernández, and K. L. Mercer. 2019. Phenotypic selection under two

contrasting environments in wild sunflower and its crop–wild hybrid. *Evolutionary*

*Applications* 12: 1703–1717.

Ridley, C. E., and N. C. Ellstrand. 2009. Evolution of enhanced reproduction in the hybrid-

derived invasive, California wild radish (*Raphanus sativus*). *Biological Invasions* 11:

2251–2264.

Ridley, C. E., S. C. Kim, and N. C. Ellstrand. 2008. Bidirectional history of hybridization in

California wild radish, *Raphanus sativus* (Brassicaceae), as revealed by chloroplast

DNA. *American Journal of Botany* 95: 1437–1442.

Rieseberg, L. H., S. C. Kim, R. A. Randell, K. D. Whitney, B. L. Gross, C. Lexer, and K.

Clay. 2007. Hybridization and the colonization of novel habitats by annual sunflowers.

*Genetica* 129: 149–165.

Rius, M., and J. A. Darling. 2014. How important is intraspecific genetic admixture to the

success of colonising populations? *Trends in Ecology and Evolution* 29: 233–242.

Roach, D. A., and R. D. Wulff. 1987. Maternal effects in plants. *Annual Review of Ecology*

*and Systematics* 18: 209–235.

Sahoo, L., J. J. Schmidt, J. F. Pedersen, D. J. Lee, and J. L. Lindquist. 2010. Growth and

fitness components of wild × cultivated *Sorghum bicolor* (Poaceae) hybrids in Nebraska.

*American Journal of Botany* 97: 1610–1617.

Schierenbeck, K. A., and N. C. Ellstrand. 2009. Hybridization and the evolution of

invasiveness in plants and other organisms. *Biological Invasions* 11: 1093–1105.

Scursoni, J. A., R. Gigón, A. N. Martín, M. Vigna, E. S. Leguizamón, C. Istilart, and R.

López. 2014. Changes in weed communities of spring wheat crops of Buenos Aires

province of Argentina. *Weed Science* 62: 51–62.

Shukla, K., A. E. Laursen, J. Benavides, N. Ejbari, and L. G. Campbell. 2021. Growth and

fecundity of colonizing hybrid *Raphanus* populations are environmentally dependent. *American Journal of Botany* 108: 580–597.

Shukla, K., S. Sbrizzi, A. E. Laursen, J. Benavides, and L. G. Campbell. 2020. Hybridization slows rate of evolution in crop-wild compared to wild populations of weedy *Raphanus* across a moisture gradient. *Frontiers in Agronomy* 2: 600346.

Singh, J., J. A. Clavijo Michelangeli, S. A. Gezan, H. Lee, and C. E. Vallejos. 2017. Maternal effects on seed and seedling phenotypes in reciprocal F1 hybrids of the common bean (*Phaseolus vulgaris* L.). *Frontiers in Plant Science* 8: 1–13.

Snow, A. A., and L. G. Campbell. 2005. Can feral radishes become weeds?, in Crop Ferality and Volunteerism: A Threat to Food Security in the Transgenic Era?, ed. by Gressel J. Taylor & Francis Group, Boca Raton, FL, pp. 193–207

Snow, A. A., T. M. Culley, L. G. Campbell, P. M. Sweeney, S. G. Hegde, and N. C. Ellstrand. 2010. Long-term persistence of crop alleles in weedy populations of wild radish (*Raphanus raphanistrum*). *New Phytologist* 186: 537–548.

Snow, A. A., P. Moran-Palma, L. H. Rieseberg, A. Wszelaki, and G. J. Seiler. 1998. Fecundity, phenology, and seed dormancy of F1 wild-crop hybrids in sunflower (*Helianthus annuus*, Asteraceae). *American Journal of Botany* 85: 794–801.

Teitel, Z., A. Klimowski, and L. G. Campbell. 2016. Assessing the effects of hybridization and precipitation on invasive weed demography using strength of selection on vital rates. *BMC Evolutionary Biology* 16: 1–13.

Ureta, M. S., F. Torres Carbonell, C. Pandolfo, A. D. Presotto, M. A. Cantamutto, and M. Poverene. 2017. IMI resistance associated to crop-weed hybridization in a natural *Brassica rapa* population: characterization and fate. *Environmental Monitoring and*

*Assessment* 189: 101.

Uwimana, B., M. J. M. Smulders, D. A. P. Hooftman, Y. Hartman, P. H. van Tienderen, J. Jansen, L. K. McHale, et al. 2012. Crop to wild introgression in lettuce: Following the fate of crop genome segments in backcross populations. *BMC Plant Biology* 12: 43.

Vercellino, R. B., F. Hernández, C. E. Pandolfo, M. Cantamutto, and A. Presotto. 2021a. Ecological fitness cost associated with the AHAS Trp574Leu mutation in feral *Raphanus sativus*. *Weed Research* 61: 210–220.

Vercellino, R. B., C. E. Pandolfo, G. Breccia, M. Cantamutto, and A. Presotto. 2018. AHAS Trp574Leu substitution in *Raphanus sativus* L.: screening, enzyme activity and fitness cost. *Pest Management Science* 74: 1600–1607.

Vercellino, R. B., C. E. Pandolfo, M. Cantamutto, and A. Presotto. 2021b. Interference of feral radish (*Raphanus sativus*) resistant to AHAS-Inhibiting herbicides in oilseed rape, wheat and sunflower crops. *International Journal of Pest Management*. DOI: https://doi.org/10.1080/09670874.2021.1959081

Vercellino, R. B., C. E. Pandolfo, A. Cerrota, M. Cantamutto, and A. Presotto. 2019. The roles of light and pericarp on seed dormancy and germination in feral *Raphanus sativus* (Brassicaceae). *Weed Research* 59: 396–406.

Vercellino, R., F. Hernández, C. Pandolfo, M. S. Ureta, and A. Presotto. 2023. Agricultural weeds: The contribution of domesticated species to the origin and evolution of feral weeds. *Pest Management Science* 79: 922–934.

Vigueira, C. C., X. Qi, B. K. Song, L. F. Li, A. L. Caicedo, Y. Jia, and K. M. Olsen. 2019. Call of the wild rice: *Oryza rufipogon* shapes weedy rice evolution in Southeast Asia. *Evolutionary Applications* 12: 93–104.

Warwick, S. I. S. 2011. Brassicaceae in Agriculture. In Schmidt R., and I. Bancroft [eds.],

Genetics and Genomics of the Brassicaceae, 33–65. Springer New York, New York,

NY.

Whitney, K. D., J. R. Ahern, and L. G. Campbell. 2009. Hybridization-prone plant families

do not generate more invasive species. *Biological Invasions* 11: 1205–1215.

Wu, D., S. Lao, and L. Fan. 2021. De-domestication: An extension of crop evolution. *Trends*

*in Plant Science* 26: 560–574.

Xia, H., H. Zhang, W. Wang, X. Yang, F. Wang, J. Su, H. Xia, et al. 2016. Ambient insect

pressure and recipient genotypes determine fecundity of transgenic crop-weed rice

hybrid progeny: Implications for environmental biosafety assessment. *Evolutionary*

*Applications* 9: 847–856.

Yue, J., L. Ze, W. Sheng, Y. Qi, H. Qi, and L. Sheng. 2021. Fitness of F1 hybrids between 10

maternal wild soybean populations and transgenic soybean. *Transgenic Research* 30:

105–119.

Zhang, X., T. Liu, J. Wang, P. Wang, Y. Qiu, W. Zhao, S. Pang, et al. 2021. Pan-genome of

*Raphanus* highlights genetic variation and introgression among domesticated, wild, and

weedy radishes. *Molecular Plant* 14: 2032–2055.

**Table 1.** Restricted maximum likelihood for survival to reproduction, days to flowering, seed weight and seed number per plant in crop-weed bidirectional hybrids and their progenitors (weed: BAL and PIE; CROP) in two contrasted environments (ruderal and agrestal) performed using maximum likelihood in SAS PROC GLIMMIX for year 1. Data were collected in the Agronomy Department at Universidad Nacional del Sur, Bahía Blanca, Argentina.

| Fixed effects | Time to flowering $df_{NUM}/df_{DEN}$ | F | P | Survival to maturity $df_{NUM}/df_{DEN}$ | F | P | Seed number plant-1 $df_{NUM}/df_{DEN}$ | F | P | Seed weight $df_{NUM}/df_{DEN}$ | F | P | Relative fitness $df_{NUM}/df_{DEN}$ | F | P |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Environment (E) | 1/14 | 11.39 | 0.0045 | 1/14 | 84.94 | <.0001 | 1/14 | 25.82 | 0.0002 | 1/14 | 12.53 | 0.0033 | 1/14 | 3.46 | 0.0838 |
| Biotype (B) | 6/1033 | 48.55 | <.0001 | 6/84 | 18.47 | <.0001 | 6/589 | 19.9 | <.0001 | 6/540 | 8.25 | <.0001 | 5/66 | 2.83 | 0.0224 |
| E x B | 6/1033 | 4.09 | 0.0005 | 6/84 | 6.12 | <.0001 | 6/589 | 0.4 | 0.8786 | 6/540 | 1.14 | 0.3351 | 5/66 | 0.48 | 0.7885 |
| Random effects | | Z | P | | Z | P | | Z | P | | Z | P | | Z | P |
| Block (E) | | 2.05 | 0.0203 | | 0 | . | | 2.17 | 0.015 | | 0.87 | 0.1924 | | . | . |
| Residuals | | 22.73 | <.0001 | | 7 | <.0001 | | 17.16 | <.0001 | | 16.44 | <.0001 | | 6.32 | <.0001 |

| Contrasts[1] | Ruderal $df_{NUM}/df_{DEN}$ | F | P | Agrestal $df_{NUM}/df_{DEN}$ | F | P | Ruderal $df_{NUM}/df_{DEN}$ | F | P | Agrestal $df_{NUM}/df_{DEN}$ | F | P | $df_{NUM}/df_{DEN}$ | F | P | $df_{NUM}/df_{DEN}$ | F | P | $df_{NUM}/df_{DEN}$ | F | P |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| W vs. W x C | 1/497 | 0.45 | 0.5042 | 1/536 | 0.93 | 0.3345 | 1/42 | 10.55 | 0.0023 | 1/42 | 0 | 1 | 1/589 | 23.72 | <.0001 | 1/540 | 2.19 | 0.1398 | 1/66 | 7.15 | 0.0094 |
| W vs. C x W | 1/497 | 6.55 | 0.0108 | 1/536 | 24.06 | <.0001 | 1/42 | 4.35 | 0.0431 | 1/42 | 0 | 1 | 1/589 | 20.23 | <.0001 | 1/540 | 3.63 | 0.0573 | 1/66 | 9.3 | 0.0033 |
| W vs. C | 1/497 | 140.75 | <.0001 | 1/536 | 104.09 | <.0001 | 1/42 | 32.67 | <.0001 | 1/42 | 30.57 | <.0001 | 1/589 | 37 | <.0001 | 1/540 | 28.1 | <.0001 | - | - | - |
| C x W vs. W x C | 1/497 | 3.59 | 0.0586 | 1/536 | 15.53 | <.0001 | 1/42 | 1.35 | 0.2518 | 1/42 | 0 | 1 | 1/589 | 0.16 | 0.6853 | 1/540 | 0.16 | 0.6927 | 1/66 | 0.1 | 0.7559 |
| C x W vs. C | 1/497 | 98.96 | <.0001 | 1/536 | 39.46 | <.0001 | 1/42 | 55.04 | <.0001 | 1/42 | 30.57 | <.0001 | 1/589 | 88.04 | <.0001 | 1/540 | 42.77 | <.0001 | - | - | - |
| W x C vs. C | 1/497 | 129.4 | <.0001 | 1/536 | 88.97 | <.0001 | 1/42 | 70.02 | <.0001 | 1/42 | 30.57 | <.0001 | 1/589 | 93.31 | <.0001 | 1/540 | 39.06 | <.0001 | - | - | - |

[1]Cross types for orthogonal contrasts: Weeds (W), crop (C), crop-weed hybrids produced on weed plants (WxC), crop–weed hybrids produced on crop plants (CxW)

**Table 2**. Restricted maximum likelihood for survival to reproduction, days to flowering, seed weight and seed number per plant in crop-weed bidirectional hybrids and their progenitors (weed: BAL and PIE; CROP) in two contrasted environments (ruderal and agrestal) performed using maximum likelihood in SAS PROC GLIMMIX for year 2. Data were collected in the Agronomy Department at Universidad Nacional del Sur, Bahía Blanca, Argentina.

| | Time to flowering | | | Survival to maturity | | | Seed number plant$^{-1}$ | | | Seed weight | | | Relative fitness | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Fixed effects | $df_{NUM}/df_{DEN}$ | $F$ | $P$ | $df_{NUM}/df_{DEN}$ | $F$ | $P$ | $df_{NUM}/df_{DEN}$ | $F$ | $P$ | $df_{NUM}/df_{DEN}$ | $F$ | $P$ | $df_{NUM}/df_{DEN}$ | $F$ | $P$ |
| Environment (E) | 1/14 | **5.06** | **0.0412** | 1/14 | 2.77 | 0.12 | 1/14 | **12.35** | **0.0034** | 1/14 | 1.71 | 0.21 | 1/14 | 1.72 | 0.2104 |
| Biotype (B) | 6/1028 | **11.52** | **<.0001** | 6/84 | 1.27 | 0.28 | 6/594 | **42.83** | **<.0001** | 6/581 | **25.61** | **<.0001** | 5/70 | **28.74** | **<.0001** |
| E x B | 6/1028 | 1.05 | 0.39 | 6/84 | 0.99 | 0.44 | 6/594 | 1.73 | 0.11 | 6/581 | 0.67 | 0.68 | 5/70 | 1.37 | 0.2476 |
| Random effects | | $Z$ | $P$ | | $Z$ | $P$ | | $Z$ | $P$ | | $Z$ | $P$ | | $Z$ | $P$ |
| Block (E) | | 2.34 | 0.0096 | | 0 | . | | 2.1 | 0.0181 | | 0.47 | 0.32 | | 0.93 | 0.1775 |
| Residuals | | 22.67 | <.0001 | | 7 | <.0001 | | 17.23 | <.0001 | | 17.04 | <.0001 | | 5.92 | <.0001 |
| Contrasts[1] | $df_{NUM}/df_{DEN}$ | $F$ | $P$ | | $F$ | $P$ | $df_{NUM}/df_{DEN}$ | $F$ | $P$ | $df_{NUM}/df_{DEN}$ | $F$ | $P$ | $df_{NUM}/df_{DEN}$ | $F$ | $P$ |
| W vs. W x C | 1/1028 | **12.45** | **0.0004** | | - | - | 1/594 | **144.1** | **<.0001** | 1/581 | **99.99** | **<.0001** | 1/70 | **86.54** | **<.0001** |
| W vs. C x W | 1/1028 | **3.87** | **0.0495** | | - | - | 1/594 | **166.94** | **<.0001** | 1/581 | **102.11** | **<.0001** | 1/70 | **105.2** | **<.0001** |
| W vs. C | 1/1028 | 0.97 | 0.33 | | - | - | 1/594 | 0.92 | 0.34 | 1/581 | **70.51** | **<.0001** | - | - | - |
| C x W vs. W x C | 1/1028 | 2.51 | 0.11 | | - | - | 1/594 | 0.48 | 0.49 | 1/581 | 0 | 0.99 | 1/70 | 0.91 | 0.3439 |
| C x W vs. C | 1/1028 | 0.4 | 0.53 | | - | - | 1/594 | **94.68** | **<.0001** | 1/581 | 0 | 0.95 | - | - | - |
| W x C vs. C | 1/1028 | 3.7 | 0.06 | | - | - | 1/594 | **81.77** | **<.0001** | 1/581 | 0.01 | 0.94 | - | - | - |

[1]Cross types for orthogonal contrasts: Weeds (W), crop (C), crop-weed hybrids produced on weed plants (WxC), crop–weed hybrids produced on crop plants (CxW).