Distinct cellular expression and subcellular localization of Kv2 voltage-gated K+ channel

subtypes in dorsal root ganglion neurons conserved between mice and humans

Robert G. Stewart[1], Miriam Camacena[1], Bryan A. Copits[2], Jon T. Sack[1,3]

Affiliations:

[1] Department of Physiology and Membrane Biology, University of California Davis,

Davis, CA 95616, USA.

[2] Washington University Pain Center, Washington University School of Medicine, St. Louis, MO

63110, USA; Department of Anesthesiology, Washington University School of Medicine, St.

Louis, MO 63110, USA.

[3] Department of Anesthesiology and Pain Medicine, University of California Davis,

Davis, CA 95616, USA.

Corresponding author email: jsack@ucdavis.edu

Pages in manuscript: 82

Number of figures: 11

Number of supplemental figures: 12

Number of tables: 2

Number of words in abstract: 248

Number of words in introduction: 601

Number of words in discussion: 1119

Conflict of interest statement: The authors declare no competing conflicts of interest.

Acknowledgments:

We thank James S. Trimmer (University of California, Davis) for generous provision of

antibodies, numerous discussions, and feedback on the manuscript. We thank Michael Ferns

(University of California, Davis) for feedback on the manuscript. We thank the human tissue

donor and their family for their generous donations to science. We thank Jiwon Yi, Zack Bertels,

Maria Payne, and Jon Lemen for performing hDRG extractions. GxTX was synthesized at the

Molecular Foundry of the U.S. Department of Energy under contract DE-AC02-05CH11231. This

research was supported by U.S. National Institutes of Health grant R01NS096317 to J.T. Sack

and the University of California, Davis.

Author contributions: R. G. Stewart: Conceptualization, formal analysis, investigation,

methodology, visualization, writing-original draft, writing-reviewing and editing. M. Camacena:

investigation. B. A. Copits investigation, writing-reviewing and editing. J.T. Sack:

Conceptualization, formal analysis, funding acquisition, investigation, methodology, project

administration, supervision, visualization, writing-original draft, writing-reviewing and editing.

ABSTRACT

The distinct organization of Kv2 voltage-gated potassium channels on and near the cell body of brain

neurons enables their regulation of action potentials and specialized membrane contact sites.

Somatosensory neurons have a pseudo unipolar morphology that allows transmitted action potentials

to bypass the cell body. Kv2 channels regulate action potentials in somatosensory neurons, yet little is

known about where Kv2 channels are located. Here we define the cellular and subcellular localization of

the Kv2 paralogs, Kv2.1 and Kv2.2, in dorsal root ganglia (DRG) somatosensory neurons with a panel of

antibodies, cell markers, and genetically modified mice. We find that relative to spinal cord neurons,

DRG neurons have similar levels of detectable Kv2.1, and higher levels of Kv2.2. In older mice,

detectable Kv2.2 remains similar while detectable Kv2.1 decreases. Both Kv2 subtypes adopt clustered

subcellular patterns that are distinct from central neurons. Kv2.2 is distinct in being found at

juxtaparanodes and paranodes of myelinated axons. Most DRG neurons co-express Kv2.1 and Kv2.2,

although neuron subpopulations show preferential expression of Kv2.1 or Kv2.2. We find that Kv2

protein expression and subcellular localization is similar between mouse and human DRG neurons. We

conclude that the organization of both Kv2 channels is consistent with physiological roles in the somata

and stem axons of DRG neurons. The general prevalence of Kv2.2 in DRG as compared to central

neurons and the enrichment of Kv2.2 relative to detectable Kv2.1, in older mice, proprioceptors, and

myelinated axons suggest more widespread roles for Kv2.2 in DRG neurons.

Significance statement

The subcellular distribution of Kv2 voltage-gated potassium channels enable compartment-

specific modulation of membrane excitability and organization of membrane contact sites. Here

we identify subcellular distributions of the Kv2 paralogs, Kv2.1 and Kv2.2, in somatosensory

neurons that bear similarities to and distinctions from central neurons. The distribution of Kv2

channels is similar in mouse and human somatosensory neurons. These results identify unique

locations of Kv2 channels in somatosensory neurons that could enable roles in sensory

information processing.

INTRODUCTION

The subcellular localization of voltage-gated ion channels in neurons determines how electrical signals are propagated. The two members of the Kv2 family of voltage gated potassium channels, Kv2.1 and Kv2.2, are important for modulating electrical signals in mammalian somatosensory neurons (Bocksteins et al., 2009; Lee et al., 2020; Sun et al., 2022; Tsantoulas et al., 2014; Zheng et al., 2019), yet little is known about where Kv2 channels are localized in these neurons. In central neurons, Kv2 channels are sequestered to specific subcellular regions and identification of where these channels are has helped elucidate their functional roles (Bishop et al., 2015; Du, Tao-Cheng, Zerfas, & McBain, 1998; Irie, 2021; Jensen et al., 2017; Johnson et al., 2018; Kihira, Hermanstyne, & Misonou, 2010; Kirmiz, Vierra, Palacio, & Trimmer, 2018; Misonou, Mohapatra, Menegola, & Trimmer, 2005; Muennich & Fyffe, 2004; Romer et al., 2014; Scannevin, Murakoshi, Rhodes, & Trimmer, 1996; Trimmer, 1991; Vierra, O'Dwyer, Matsumoto, Santana, & Trimmer, 2021). Establishing where Kv2 channels are in somatosensory neurons likewise identifies sites where they could function.

Among vertebrates, the Kv2 family contains two conserved paralogs, Kv2.1 and Kv2.2, which can assemble into homo- or hetero-tetramers to form voltage gated K^+ channels (Blaine & Ribera, 1998; Kihira et al., 2010). In central neurons, Kv2.1 and Kv2.2 channels have unique cellular expression, subcellular localization and show distinct physiological roles (Bishop et al., 2015; Newkirk et al., 2022). Among cortical neuron types, Kv2.1 is more broadly distributed than Kv2.2 (Bishop et al., 2015). Little is known of how Kv2.1 or Kv2.2 channels are distributed among somatosensory neuron subtypes. In central neurons, Kv2.1 and Kv2.2 channels localize in clusters on the cell soma, proximal dendrites, and axon initial segment (Bishop et al., 2015; King, Manning, & Trimmer, 2014; Sarmiere, Weigle, & Tamkun, 2008; Scannevin et al., 1996; Trimmer, 1991). Clustered Kv2 channels form endoplasmic reticulum-

plasma membrane junctions (Johnson et al., 2018; Kirmiz et al., 2018), where they perform

nonconducting functions which include mediating coupling of L-type calcium channels to ryanodine

receptors (Vierra et al., 2021), and Ca^{2+} uptake (Panzera et al., 2022). In somatosensory neurons it

remains unknown whether Kv2 channels form organized structures or are localized to specific

subcellular compartments. Identifying if channel expression is specific to neuron subtypes and within

specific subcellular domains could indicate whether Kv2.1 and Kv2.2 channels also have distinct

physiological roles in somatosensory neurons.

Kv2 channels are expressed and play important functional roles in DRG. Kv2 transcripts have been

detected in all classes of DRG neurons with a notable absence of Kv2.1 mRNA in proprioceptors (Usoskin

et al., 2015; Wangzhou et al., 2020; Zheng et al., 2019). Presence of Kv2.1 protein in DRG of mice and

rats has been identified by western blot and immunohistochemistry respectively (Sun et al., 2022;

Tsantoulas et al., 2012). Pharmacological inhibition of Kv2 channels enhances rat DRG neuron

responsiveness to sustained inputs (Tsantoulas et al., 2014). Kv2 conductances have been identified in

several mouse DRG neuron subtypes (Bocksteins et al., 2009; Regnier, Bocksteins, Van de Vijver,

Snyders, & van Bogaert, 2016; Zheng et al., 2019). Kv2 channels have also been implicated in

nociception. Kv2.1 and Kv2.2 mRNA transcript levels are decreased after peripheral axotomy of rat DRG

neurons (Tsantoulas et al., 2014). Suppression of Kv2.1 transcription by the epigenetic factor Cdyl is

implicated in regulating pain sensation (Sun et al., 2022). A missing piece of the puzzle of how Kv2

channels are involved in somatosensory neuron function is the location of the channels themselves.

Here we use antibodies against Kv2.1 and Kv2.2 to define their cellular expression and subcellular

localization in DRG neurons of both mice and humans.

MATERIALS AND METHODS

Mice

This study was approved by the UC Davis Institutional Animal Care and Use Committee and conforms to

guidelines established by the NIH. Mice were maintained on a 12 h light/dark cycle, and food and water

was provided *ad libitum.* Kv2.1 KO and Kv2.2 KO mice were from breeding Kv2.1+/- or Kv2.2+/-

heterozygous mice such that WT mice were from breeding pairs that generated KO mice. Kv2.1 and Kv2.2

DKO mice were generated from breeding Kv2.1+/- heterozygous and Kv2.2-/- homozygous mice. WT

controls for Kv2.1/Kv2.2 DKO experiments were C57BL/6J mice purchased from Jackson Laboratory (stock

#000664). The PV^{Ai14} mouse line was a generous gift from Dr. Theanne Griffith (University of California

Davis, Davis CA) and were a cross of $Rosa26^{Ai14}$ (stock #007914, MGI: J:155793) and *PVcre* (stock #008069,

MGI:J: 100886) mice. The $MrgprD^{GFP}$ mouse line was a generous gift from Dr. David Ginty (Harvard

University, Boston MA) (MGI: 3521853). Detailed information about the genotype, age, sex and level of

spinal column of mice used in each figure can be found in Table 1.

Human Tissue Collection

Human DRG (L3 spinal level) were obtained in collaboration with Mid-America Transplant (St. Louis, MO)

from a 59-year-old black male (cause of death: hemorrhagic stroke). DRG were extracted less than 2

hours after aortic cross clamp and transported to the lab where they were dissected to remove the

dura, embedded in Optimal Cutting Temperature (OCT) compound, snap frozen, and stored at -80 °C

until use (Valtcheva et al., 2016). Human DRG were obtained from organ donors with full legal consent

for use of tissue for research in compliance with procedures approved by Mid-America Transplant.

Tissue Preparation

Mice were briefly anesthetized with 3-5% isoflurane and then decapitated. The spinal column was

dissected, and excess muscle tissue removed. The spinal column was then bisected in the middle of the

L1 vertebrae identified by the 13[th] rib and drop fixed for 1 hour in ice cold PFA: 4% formaldehyde

prepared fresh from paraformaldehyde in 0.1 M phosphate buffer (PB) pH adjusted to 7.4 with NaOH.

0.1 M PB buffer was diluted from a 0.4 M PB stock solution that was made by diluting 91.37 g of

$Na_2H_2PO_4$ and 20.98 g of $Na_2H_2PO_4 \bullet H_2O$ in 2 liters of milliQ water. The relatively short fixation period

was chosen because a longer fixation period increased the off target secondary antibody

immunofluorescence from mouse isotype secondary antibodies (Figure 1-1). The spine was washed 3×

for 10 min each wash in PB and cryoprotected at 4 °C in 30% sucrose in PB for 24 hours. The spine was

cut into sections containing two vertebra per sample which were frozen in OCT (Fisher cat#4585) and

stored at -80 °C until sectioning. Vertebrae position relative to the 13[th] rib was recorded for each frozen

sample to determine the specific vertebrae position in the spinal cord. Samples were cut into 30 μm

sections on a freezing stage sliding microtome and were collected on Colorfrost Plus microscope slides

(Fisher cat#12-550-19). Slides were stored at -20 °C or immediately used for multiplex

immunofluorescence labeling.

Human DRG were cut into 30 μm sections on a freezing stage sliding microtome and were collected on

Colorfrost Plus microscope slides. Sections were briefly thawed to adhere to the slide but were

immediately returned to the cryostat kept at -20 °C. Slides were removed from the cryostat and

immediately transferred to freshly made 4% PFA pH 7.4 for 10 minutes. Slides were then placed in PB

solution for 10 minutes and immediately used for multiplex immunofluorescence labeling. Attempts at

immunolabeling Kv2 channels in human DRG tissue that was fixed in 4% PFA for 24 hours were not

successful.

Neuron Cell Culture

Cervical, thoracic and lumbar DRGs were harvested from mice and transferred to Hank's buffered saline

solution (HBSS) (Invitrogen). Ganglia were treated with collagenase (2 mg/ml; Type P, Sigma-Aldrich) in

HBSS for 15 min at 37°C followed by 0.05% Trypsin-EDTA (Gibco) for 2.5 min with gentle rotation. Trypsin

was neutralized with culture media (MEM, with L-glutamine, Phenol Red, without sodium pyruvate)

supplemented with 10% horse serum (heat-inactivated; Gibco), 10 U/ml penicillin, 10 µg/ml streptomycin,

MEM vitamin solution (Gibco), and B-27 supplement (Gibco). Serum-containing media was decanted and

neurons were triturated using a fire-polished Pasteur pipette in MEM culture media containing the

supplements listed above. Neurons were plated on laminin-treated (0.05 mg/ml, Sigma-Aldrich) 35 mm

glass bottom dishes (MatTek cat#P35G-1.5-7-C). Neurons were then incubated at 37°C in 5% CO_2. Neurons

were used for imaging experiments within 24 hours after plating.

Labeling Endogenous Kv2 in live cultured DRG neurons

Plates were removed from 37°C incubator 24 hours after plating. Neurons were washed once with 1 mL

neuronal external (NE) solution (3.5 KCl, 155 NaCl, 10 HEPES, 1.5 CaCl2, and 1 MgCl2, 10 mM glucose

adjusted to pH 7.4 with NaOH). Neurons were then incubated with NE solution supplemented with 1%

BSA for 5 minutes. Neurons were then washed once with NE solution and incubated in 5 µg/mL wheat

germ agglutinin conjugated to Alexa Fluor 405 (Biotium cat#29028-1) diluted in NE solution for 2.5

minutes. Neurons were washed twice with NE solution and imaged. NE solution was removed and then

neurons were incubated for 10 minutes in 100 nM GxTX-594 prepared as described (Thapa et al., 2021).

Neurons were then washed three times in NE solution and imaged again. The temperature of the

imaging chamber was 25 °C throughout the experiment. Experimenter was blinded to which culture

dishes contained DRG neurons from either WT or Kv2.1/Kv2.2 double knockout mice throughout the

experiment and image analysis.

Immunohistochemistry

A hydrophobic barrier was drawn around tissue sections mounted on slides as described above using a

hydrophobic barrier pen (Scientific Device cat#9804-02). Sections were incubated in 4% milk, 0.1 M

phosphate buffer (PB) and 0.2% Triton X-100 (vehicle) for 1 hour. Sections were then incubated in

vehicle containing 0.1 mg/mL IgG F(ab) polyclonal IgG antibody (Abcam cat#ab6668) for 1 hour. We

determined that a concentration of 0.1 mg/mL maximally blocked off-target secondary antibody labeling

when using isotype-specific anti-mouse secondary antibodies (Figure 1-2). Sections were washed 3× for

5 min each in vehicle and then incubated in vehicle containing primary antibodies for 1 hour. Primary

antibodies and concentrations used are listed in Table 2. Sections were then washed 3× for 5 min each

in vehicle and then incubated in vehicle containing IgG-subclass-specific secondary Abs for one hour

(Thermo Fisher). Sections were then washed 3× for 5 min each in PB and mounted with Prolong Gold

(Thermo Fisher cat#P36930) and Deckglaser cover glass (Fisher Scientific cat#NC1776158). All

incubations and washes were done at room temperature with gentle rocking. Human tissue was

immunolabeled identically to mice with the exception that the step incubating sections in vehicle

containing 0.1 mg/mL IgG F(ab) polyclonal IgG antibody was omitted and sections were incubated in

primary antibodies for 2 hours instead of 1. Primary antibodies used in Figure 2 and Figure 3 were used

at saturating concentrations (Figure 2-1).

Imaging

Images were acquired with an inverted scanning confocal and airy disk imaging system (Zeiss LSM 880

Airyscan, 410900-247-075) run by ZEN black v2.1. Laser lines were 405 nm, 488 nm, 543 nm and 633 nm.

Low-magnification images to image whole mouse DRG were acquired in confocal or airy disk imaging

mode with a 0.8 NA 20x objective (Zeiss 420650-9901) details in figure legends. Images containing both

the DRG and spinal cord were tile scan images acquired in confocal mode with the same 20x objective.

When imaging whole DRG for Figure 2 and Figure 3 the imaging plane was selected using fluorescence

from channels that did not contain anti-Kv2.1 or anti-Kv2.2 immunofluorescence. High-magnification

images were acquired in airy disk imaging mode with a 1.4 NA 63x oil objective (Zeiss 420782-9900-799).

Linear adjustments to contrast and brightness and average fluorescence intensity z-projections were

performed using ImageJ software.

Image Analysis

Images were analyzed using ImageJ software (Schindelin et al., 2012). A summary of automated analysis

for selecting neurons in DRG sections can be found in Figure 2-2. The ImageJ plugin MorphoLibJ was

used for performing watershed segmentation of images (Legland, Arganda-Carreras, & Andrey, 2016).

Unless stated otherwise the region of DRG neurons used to analyze anti-Kv2 immunofluorescence is the

outer edge (Figure 2-2 I). Immunofluorescence is defined as the raw pixel values from confocal images

and was not background subtracted. We noted that in smaller neurons the nucleus takes up a greater

percentage of the total volume and could skew fluorescence measurements if fluorescence in the entire

soma volume is measured. As Kv2 protein is not present in the nucleus and is enriched at the outer edge

(Figure 1) we found that measuring anti-Kv2 immunofluorescence at the outer edge of the neuron

reduces potential error that could be associated with soma diameter. For proteins that were not Kv2

channels we analyzed the immunofluorescence of the entire soma as these proteins did not exhibit the

same enrichment at the outer edge as Kv2 channels (Figure 10 B). Where specified manual ROIs were

used in analysis otherwise automatically generated ROIs were used. When analyzing

immunofluorescence from human neurons, ROIs were drawn around areas within the neuronal soma

that do not contain apparent lipofuscin (Figure 11-1 C). Fitting of imaging data was performed using Igor

Pro software version 8 (Wavemetrics, Lake Oswego, OR) that employs nonlinear least square curve

fitting via the Levenberg-Marquardt algorithm. Distributions of fluorescence intensity from DRG neurons

were fit with a log normal distribution:

$$f(x) = A^{-\left[\frac{\ln\left(\frac{x}{x_0}\right)}{width}\right]^2}$$ (Equation 1)

Where A = amplitude, x_0 = mean, and width = $\sqrt{2}$ times standard deviation. Concentration-effect

experiments in Figure 2-1 were fit with the Hill equation:

$$f(x) = base + \frac{max-base}{1+\left(\frac{x_{half}}{x}\right)^n}$$ (Equation 2)

Code Accessibility

In house Fiji macros and R script used to process imaging data are available for download at

https://github.com/SackLab/DRG-Image-Processing

Statistics

All statistical tests were performed in Igor Pro software version 8 (Wavemetrics, Lake Oswego, OR).

Independent replicates (n) are individual neurons while biological replicates (N) are individual mice. The

n and N values for each figure are listed in Table 1. Details of statistical tests are in the figure legends.

RESULTS

Kv2.1 and Kv2.2 immunolabeling is enriched at the apparent plasma membrane of DRG neuron

somata

To assess the cellular expression and subcellular localization of Kv2.1 and Kv2.2 voltage gated potassium

channels in DRG neurons we used a set of anti-Kv2.1 and anti-Kv2.2 antibodies that have previously

been validated in brain tissue from Kv2 knockout mice (Bishop et al., 2015) to perform multiplex

immunofluorescence labeling of DRG neurons. Anti-Kv2.1 (magenta) and anti-Kv2.2 (green)

immunofluorescence is prominent in DRG neuron somata (Figure 1 A). In most neurons we observed

anti-Kv2.1 and anti-Kv2.2 immunofluorescence enriched at the outer edge of DRG neuron somata

consistent with their plasma membrane localization (Figure 1 B arrows). This putative cell surface

localization is distinct from prior reports of anti-Kv2 immunofluorescence in DRG (Tsantoulas et al.,

2012) as well as that for other ion channel proteins such as Nav1.8 (Shields et al., 2012) and TRPV1 (Cho

& Valtschanoff, 2008), all of which are primarily in the cytoplasm. Anti-Kv2.1 or anti-Kv2.2

immunofluorescence at the edge of neuron somata was more apparent when imaging thinner optical

sections. To determine if enrichment of Kv2 immunolabeling at the outer edge of DRG neurons was an

artifact of our immunohistochemistry or imaging protocols, we additionally labeled DRG sections with a

knockout-validated antibody against Nav1.8. In the same DRG section labeled for Kv2 channels,

cytoplasmic anti-Nav1.8 immunofluorescence was prominent in small to medium diameter lumbar DRG

neurons, consistent with previous reports (He et al., 2010; Shields et al., 2012) (Figure 1 B). To quantify

enrichment of anti-Kv2 versus anti-Nav1.8 immunofluorescence at the outer edge of DRG neurons, we

manually drew regions of interest (ROIs) for each DRG soma. We generated a 2 μm wide annulus that

encompassed the outer edge and a second 2 μm wide annulus just inside the first one (Figure 1 C arrows

1 and 2 respectively). By comparing the fluorescence intensity of immunofluorescence signals within

these annuli we found that the edge of DRG neuron somata is enriched for anti-Kv2.1 and anti-Kv2.2

immunofluorescence relative to that of Nav1.8 (t-test: Kv2.1:Nav1.8 = p < 0.001; Kv2.2:Nav1.8 = p <

0.001) (Figure 1 D). Similarly, anti-Kv2.1 and anti-Kv2.2 immunofluorescence is enriched at the edge of

DRG neuron somata relative to anti-TRPV1 immunofluorescence (t-test: Kv2.1:Nav1.8 = p < 0.001;

Kv2.2:Nav1.8 = p < 0.001) (Figure 1-3).

Most dorsal root ganglion neurons express Kv2 protein

To determine whether the anti-Kv2.1 immunofluorescence signal in DRG sections is specific and

dependent on the presence of Kv2.1 protein, we compared fluorescence intensities in DRG samples

prepared in parallel from wild-type (WT) mice and Kv2.1 knock-out (KO) mice. We observed anti-Kv2.1

immunolabeling in WT DRG neurons that was absent from Kv2.1 KO, while fluorescence corresponding

to an antibody that targets βIII tubulin was similar in both WT and Kv2.1 KO mice (Figure 2 A). To

quantify fluorescence intensities of individual neurons, we used an automated method to generate

regions of interest (ROIs) at the outer edge of DRG neuron somata. This analysis confirmed a reduction

in anti-Kv2.1 but not anti-βIII tubulin immunofluorescence in Kv2.1 KO samples (Figure 2 B and C). In 5

mice, anti-Kv2.1 immunofluorescence was significantly reduced in Kv2.1 KO samples relative to WT (t-

test p < 0.0001) while no significant difference was observed in these samples for anti-Kv2.2 or anti-βIII

tubulin immunofluorescence (Figure 2 D).

We developed a method to estimate the fraction of DRG neurons that express Kv2.1 protein on their

outer edge. Fluorescence from Kv2.1 KO neurons labeled with an anti-Kv2.1 antibody had neuron-to-

neuron variability which could reasonably be fit with a log normal distribution (equation 1) (Figure 2 E).

We assumed a similar distribution of background immunofluorescence is also present in WT DRG

neurons and scaled the Kv2.1 KO log normal fit to estimate the fraction of neurons without detectable

Kv2.1 (Figure 2 F red gaussian). Using this method we compared Kv2.1 immunofluorescence in DRG

sections of age and sex matched WT and Kv2.1 KO mice from the 12[th] thoracic vertebra and estimated

that 84 ± 9% (SD) of DRG neurons have detectable Kv2.1 protein (N = 5 pairs of mice) (Figure 2 G).

Using the same type of analysis, we found that anti-Kv2.2 immunofluorescence was significantly

reduced in Kv2.2 KO mice (Two tailed t-test p < 0.001) while no significant difference in anti-NF200 or

anti-Kv2.1 immunofluorescence was observed (Figure 3 A-D). Comparing anti-Kv2.2

immunofluorescence from WT and Kv2.2 KO mice indicates that 91 ± 5% (SD) of the WT mouse DRG

neurons have detectable Kv2.2 protein (N = 8 pairs of mice) (Figure 3 E-G). We did not observe a

difference between males and females in detectable Kv2.1 or Kv2.2 protein, but animal numbers were

not sufficient for rigorous statistical comparison.

We validated this methodology with a transgenic MrgprD-GFP mouse line that expresses GFP in non-

peptidergic nociceptors that comprise approximately 19-24% of all lumbar DRG neurons (Dirajlal,

Pauers, & Stucky, 2003; Dong, Han, Zylka, Simon, & Anderson, 2001; Wang & Zylka, 2009). In comparing

the MrgprD-GFP mice to WT C57BL/6J mice we found that 23 ± 15% (SD) of lumbar neurons (N = 4 pairs

of mice) had detectable GFP (Figure 2-3).

Kv2.1 immunolabelling decreases in older mice while Kv2.2 immunolabelling remains similar

Kv2.2 transcript in mouse DRG decreases during postnatal development (Regnier et al., 2016) and Kv2.1

protein expression in mouse brain decreases with age (Cotella et al., 2012; Regnier et al., 2016). To

identify if Kv2 protein levels in DRG neurons vary between young adult and old mice we compared anti-

Kv2.1 and anti-Kv2.2 immunofluorescence from samples subjected to identical immunolabeling and

imaging protocols. We observed lower anti-Kv2.1 (Figure 4 A, B and C) in DRG sections from 50 week old

mice relative to 7 week old mice. Comparisons with Kv2.1 KO mice indicate that $50 \pm 23\%$ (SD) of DRG

neurons from 50 week old mice express detectable Kv2.1 protein, significantly less (p = 0.018) than the

$84 \pm 9\%$ (SD) of neurons with detectable Kv2.1 protein from 7-16 week old mice (Figure 4 D). In contrast,

similar anti-Kv2.2 immunofluorescence was seen in samples from young adult and old mice (Figure 4 E, F

and G). Comparisons with Kv2.2 KO mice indicate that $96 \pm 3\%$ of DRG neurons from 50 week old mice

express detectable Kv2.2 protein, a weakly significant increase (p = 0.03) from the $88 \pm 3\%$ (SD)

detectable Kv2.2 protein in DRG neurons from 7-24 week old mice (Figure 4 H).

Kv2 channels are expressed on the cell surface of acutely dissociated DRG neurons

To determine if Kv2 channels are present on neuron surfaces we labeled live acutely dissociated DRG

neurons with a cell-impermeant fluorescent probe which binds Kv2 channels. The probe is a variant of

the tarantula peptide guangxitoxin-1E conjugated to Alex Fluor 594 (GxTX-594) which binds to an

extracellular site of Kv2.1 and Kv2.2 channels (Thapa et al., 2021). We applied GxTX-594 to live acutely

dissociated DRG neurons from WT mice and double knockout (DKO) mice lacking expression of both

Kv2.1 and Kv2.2 protein (Figure 5). In WT DRG neurons, we observed fluorescence at the membrane

after application of 100 nM GxTX-594 that was not present before the addition of GxTX-594 (Figure 5 A).

Wheat germ agglutinin conjugated to Alexa 405 (WGA-405) was used to identify the surface membrane

of cultured DRG neurons and confirmed that the GxTX-594 fluorescence observed in WT neurons was

cell surface localized (Figure 5 A middle). The cell surface localized fluorescence of GxTX-594 observed in

WT neurons was not present in Kv2.1/Kv2.2 DKO neurons (Figure 5 B). To quantify the cell surface-

associated fluorescence signals of individual WT and Kv2.1/Kv2.2 DKO neurons, we used an automated

method to generate ROIs corresponding to the WGA-405 fluorescence and measured GxTX-594 labeling

within this ROI (Figure 5 C). This analysis confirmed a reduction in GxTX-594 fluorescence at the

membrane in cultured DRG neurons from Kv2.1/Kv2.2 DKO mice compared to WT mice (Figure 5 D).

Similar to the Kv2 immunofluorescence signals from KO mice (Figures 2 and 3), GxTX-594 fluorescence

from Kv2.1 /Kv2.2 DKO mice could be reasonably fit with a log normal distribution (equation 1) (Figure 5

E). Applying the same method used to estimate the fraction of neurons expressing detectable Kv2.1 and

Kv2.2 immunofluorescence we compared cultured DRG neurons of age and sex matched WT and

Kv2.1/Kv2.2 DKO mice and estimated that 76 ± 2.4% (SD) of DRG neurons have detectable cell surface

GxTX-594 fluorescence (Figure 5 F and G). This indicates that most, if not all, neurons express Kv2

channels in their surface membrane, consistent a report of Kv2 conductance in every DRG neuron type

assessed (Zheng et al., 2019) .

Anti-Kv2.2 immunofluorescence is enriched in DRG neurons relative to spinal cord neurons

In mammals, anti-Kv2.1 and anti-Kv2.2 immunofluorescence is apparent in central neurons of the brain

(Kihira et al., 2010; Bishop et al., 2015) and anti-Kv2.1 in the spinal cord (Bishop et al., 2015; Muennich &

Fyffe, 2004). We measured Kv2 immunofluorescence intensities in sections of mouse spinal column

which contain both the DRG and the spinal cord. Neurons in the ventral and dorsal horn of the spinal

cord exhibit anti-Kv2.1 immunofluorescence (Figure 6 A). We observed anti-Kv2.1 puncta on neurons in

the ventral horn (Figure 6 A inset) similar to that described in alpha motor neurons (Fletcher et al., 2017;

Romer et al., 2014). Fewer spinal cord neurons exhibit anti-Kv2.2 immunofluorescence than anti-Kv2.1

(Figure 6 A and Figure 6-1 A arrowheads). Kv2.2 channel protein in the spinal cord of mice has to our

knowledge not been assessed, but anti-Kv2.2 immunofluorescence has been observed early in

development in the ventrolateral spinal cord in the frog *Xenopus laevis* (Gravagna, Knoeckel, Taylor,

Hultgren, & Ribera, 2008). We analyzed anti-Kv2 immunofluorescence from individual neurons in the

DRG and ventral horn of five mice and found that while anti-Kv2.1 immunofluorescence is similar

between the two regions (Figure 6 B), anti-Kv2.2 immunofluorescence is significantly higher in DRG

neurons relative to ventral horn neurons (Figure 6 C). Approximately 30% of neurons in the DRG have

anti-Kv2.2 immunofluorescence at a level higher than the brightest anti-Kv2.2 immunofluorescence

observed in the spinal cord. We repeated this analysis with a different anti-Kv2.2 antibody and found

similar results (Figure 6-1).

Kv2 channels on DRG neuron somata and stem axons form clustered subcellular patterns with

similarities to and distinctions from central neurons.

In neurons throughout the brain (Trimmer, 1991, Scannevin et al., 1996, Bishop et al., 2015) and motor

neurons in the spinal cord (Muennich and Fyffe, 2004), Kv2 channels form punctate structures referred

to as clusters. In central neurons, Kv2 clusters localize to the soma, proximal dendrites and axon initial

segment (Lim, Antonucci, Scannevin, & Trimmer, 2000; Scannevin et al., 1996; Trimmer, 1991). At

clusters, Kv2 channels organize protein signaling complexes and endoplasmic reticulum-plasma

membrane junctions (Johnson et al., 2018; Kirmiz et al., 2018; Panzera et al., 2022; Vierra et al., 2021).

To determine whether Kv2 channels in DRG neurons form clusters similar to central neurons, we

compared *en face* z-projections of Kv2 clusters in DRG neurons and ventral horn neurons in the same

spinal cord section (Figure 7). Anti-Kv2.1 immunofluorescence in the ventral horn neurons of these mice

resembles previous reports of mice (Fletcher et al., 2017) and rats (Fletcher et al., 2017; Muennich &

Fyffe, 2004). In both DRG and spinal cord neurons Kv2.1 and Kv2.2 channels form dense clusters (Figure

7 A and B). Kv2 channel clusters in DRG typically appeared smaller than Kv2 clusters in spinal cord.

However, we are not certain these observations in fixed tissue represent the in vivo arrangement, as

Kv2.1 channels in brain neurons can decluster in the period between sacrifice and fixation (Misonou et

al., 2005). In some DRG neurons, we observed donut shaped Kv2.1 and Kv2.2 clusters (Figure 7 B and C

insets) which resemble donut shaped Kv2 clusters in interneurons which harbor specialized protein

machinery in their center (Vullhorst et al., 2015).

DRG neurons are distinct from central neurons as they have a single stem axon that exits the cell soma

and bifurcates into the central and peripheral axon branches (Ha, 1970). We observed punctate

immunofluorescence for both Kv2.1 and Kv2.2 on the apparent stem axons of DRG neurons (Figure 7 B,

D and E arrowheads, Figure 7-1). We observed Kv2 clusters on axons clearly protruding from C-fiber

neurons expressing soluble GFP (Figure 7-1 A), confirming stem axon localization.

While Kv2 clusters in central neurons and many DRG neurons are distributed evenly throughout the

soma, we found that DRG neurons asymmetrically distribute Kv2 clusters on the soma (Figure 7 B-E

arrows). To test if enrichment of Kv2 channel clusters is oriented to a specific region on the soma we

manually traced the surface of neurons whose apparent stem axons were visible (Figure 7 F). The

brightest anti-Kv2.1 and anti-Kv2.2 immunofluorescence is near the stem axon in 76 and 80% of DRG

neurons respectively (Figure 7 G and H). Asymmetric distribution of Kv2 channels is not frequently

observed in central neurons of mice (Bishop et al., 2018; Bishop et al., 2015; Romer et al., 2014)

suggesting a potentially unique role of Kv2 clusters in DRG neurons.

Kv2.2 channels are expressed in peripheral axons of DRG neurons at the paranodes and

juxtaparanodes

To determine if anti-Kv2 immunofluorescence is detectible in DRG axons in regions beyond the stem

axon we immunolabeled samples containing both peripheral and central axons, using NF200 or βIII

tubulin as a markers for DRG axons (Figure 8 and Figure 8-1). Neurons were not labeled with βIII tubulin

in Figure 8 because the βIII tubulin and Kv2.2 targeting antibodies were of the same isotype. We did not

observe anti-Kv2.1 immunofluorescence in peripheral axons (Figure 8-1) from neurons that did have

detectable anti-Kv2.1 immunofluorescence in the neuron soma (Figure 2). However, anti-Kv2.2

immunofluorescence is detectable in peripheral and central axons of WT but not Kv2.2 KO mice (Figure

8 B and C). Anti-Kv2.2 immunofluorescence on myelinated peripheral axons appears as discrete puncta

or as larger bands which span the width of the axon (Figure 8 B arrows). We also observed anti-Kv2.2

immunofluorescence not colocalized with NF200, suggestive of unmyelinated fibers (Figure 8 B

arrowhead). In MrgprD-GFP mice, anti-Kv2.2 immunofluorescence colocalizes with axons expressing

cytosolic GFP, identifying that unmyelinated C-fiber neurons also express Kv2.2 protein along the

peripheral axon (Figure 8 D)

We noted that anti-Kv2.2 immunofluorescence on myelinated fibers was not uniform and was in a

pattern suggestive of localization at nodes of Ranvier. To investigate if Kv2.2 channels have a nodal

subcellular localization we co-labeled peripheral axons with antibodies specific for CASPR (Einheber et

al., 1997) and Kv1.2 (Rasband et al., 1998) to identify the paranodal and juxtaparanodal regions of nodes

of Ranvier, respectively. We found punctate anti-Kv2.2 immunofluorescence associated with both the

paranode and juxtaparanode of WT DRG neuron axons that is not present in Kv2.2 KO mice (Figure 9 A).

We analyzed anti-Kv2.2 immunofluorescence at the paranode and juxtaparanode of WT and Kv2.2 KO

mice by manually drawing ROIs around CASPR and Kv1.2 immunofluorescence while blinded to anti-

Kv2.2 immunofluorescence and found that 26/34 paranodes and 20/31 juxtaparanodes had anti-Kv2.2

immunofluorescence brighter than the brightest anti-Kv2.2 immunofluorescence measured in the

paranodes and juxtaparanodes of Kv2.2 KO mice (Figure 9 B).

Anti-Kv2.1 and anti-Kv2.2 immunofluorescence is non-uniform across DRG neuron subtypes

DRG neurons are a highly heterogeneous population and different voltage gated potassium channel

families make distinct contributions to outward potassium currents in different sensory neuron subtypes

(Zheng et al., 2019). We observed subpopulations of DRG neurons that have bright anti-Kv2.1

immunofluorescence but dim anti-Kv2.2 immunofluorescence and vice-versa (Figure 10 A and C, Figure

1 A, and Figure 6 A). We hypothesized that the differences in the cellular expression patterns of Kv2.1

and Kv2.2 could potentially denote specific subtypes of sensory neurons. We assessed anti-Kv2.1 and

anti-Kv2.2 immunofluorescence in four DRG neuron subtypes, employing genetically encoded markers

for non-peptidergic nociceptors (MrgprD-GFP) or proprioceptors (PV-tdTomato), or immunolabeling to

mark peptidergic nociceptors (CGRP) or myelinated DRG neurons (NF200) (Figure 10 B and D). The

peptidergic and non-peptidergic nociceptors have similar ranges of both anti-Kv2.1 and anti-Kv2.2

immunofluorescence. Many neurons with the brightest anti-Kv2.1 or anti-Kv2.2 immunofluorescence

were NF200-positive myelinated neurons. Proprioceptors, a subset of the myelinated neurons, have a

range of anti-Kv2.2 but dim anti-Kv2.1 immunofluorescence. The neurons with bright anti-Kv2.1 and dim

anti-Kv2.2 immunofluorescence are mostly NF200-positive myelinated neurons of unknown subtype

(Figure 10 B and C). Overall, these results are consistent with mRNA transcript levels reported for Kv2.1

and Kv2.2 in DRG neuron subtypes; notably that Kv2.1 and Kv2.2 transcript levels are similar in identified

classes of DRG neurons, except for proprioceptors where Kv2.1 mRNA transcripts are low (Usoskin et al.,

2015; Zheng et al., 2019).

Kv2 channel cellular expression and subcellular localization in human DRG neurons is similar to mice

Previous work has shown that Kv2 mRNA transcripts are present in human DRG neurons (Tavares-

Ferreira et al., 2022; Wangzhou et al., 2020). As human Kv2.1 and Kv2.2 knockout controls are not

available, we compared fluorescence in DRG sections with anti-Kv2 primary antibodies to those lacking

primary antibodies to determine Kv2 channel protein localization in human DRG neurons. Control

samples lacking primary antibodies revealed strong autofluorescence at all excitation wavelengths

(Figure 11 A, Figure 11-1 A and B) consistent with other studies of human DRG neurons which have

attributed this autofluorescence to lipofuscin (Shiers et al., 2021). We found that most neurons have

anti-Kv2.1 and anti-Kv2.2 immunofluorescence brighter than controls lacking primary antibodies (Figure

11-1 D and E) suggesting that Kv2 protein is present in human DRG neurons. As a control, we used the

same methods with anti-Nav1.7 and anti-Nav1.8 antibodies and found anti-Nav1.7 immunofluorescence

in nearly all human DRG neurons while anti-Nav1.8 was in a smaller fraction of DRG neurons, consistent

with previous reports of Nav1.7 protein and Nav1.8 transcript in human DRG (Shiers, Klein, & Price,

2020) (Figure 11-1 A arrowheads, B, F and G). Similar to DRG neurons in mice, we observed

subpopulations of neurons that had bright anti-Kv2.1 immunofluorescence but low anti-Kv2.2

immunofluorescence and conversely neurons that had bright anti-Kv2.2 immunofluorescence but low

anti-Kv2.1 immunofluorescence (Figure 11 A and Figure 11-1 A and H). Attempts to label neurons with

antibodies that target parvalbumin were unsuccessful, and we were unable to identify if the population

of human DRG neurons with low anti-Kv2.1 and high anti-Kv2.2 are proprioceptors as we observed in

mice.

To identify if Kv2 protein subcellular localization in human is similar to mice we examined the pattern of

anti-Kv2.1 and anti-Kv2.2 immunofluorescence in sections of human DRG. We identified neurons in

human DRG where anti-Kv2.1 and anti-Kv2.2 immunofluorescence is enriched at the outer edge of DRG

neuron somata (Figure 11 A arrows and Figure 11-2). In some neurons, Kv2 clusters exhibited

asymmetric distribution on the neuronal soma (Figure 11 B arrows, Figure 11-2 arrows). *En face* z-

projections of human DRG neurons revealed Kv2 clusters as well as donut shapes similar to observations

in mice (Figure 11 C). We were not able to identify Kv2.1 on the stem axon of human DRG neurons but

did observe Kv2.2 on apparent stem axons (Figure 11 C arrow) along with anti-Nav1.7

immunofluorescence (Figure 11-3 arrow). To determine if Kv2.2 channels are present in human nodes of

Ranvier, we labeled human DRG sections with the same anti-Kv2.2, anti-CASPR and anti-Kv1.2 antibodies

that were used in mice. We found anti-Kv2.2 immunofluorescence at the CASPR-labeled paranode and

Kv1.2 labeled juxtaparanode of human DRG neuron fibers (Figure 11 D arrowheads and arrows

respectively) suggesting that these channels might contribute to electrical propagation or other

physiological functions in human DRG axons.

DISSCUSSION

Our results reveal overlapping yet distinct localization patterns of Kv2.1 and Kv2.2 in DRG neurons. Here,

we discuss the unique features of Kv2 localization in the DRG and how these could influence neuronal

physiology and somatosensation.

We identify differences between anti-Kv2.1 and anti-Kv2.2 immunofluorescence in DRG neurons. Kv2.2

mRNA is abundant in all DRG neuron subtypes identified, while Kv2.1 mRNA is low in proprioceptors and

abundant in other subtypes (Usoskin et al., 2015; Zheng et al., 2019). Consistent with these findings we

identify Kv2.1 and Kv2.2 immunofluorescence in most DRG neurons and find that Kv2.1

immunofluorescence is anomalously low in proprioceptors. Detection of Kv2.2 but not Kv2.1 in DRG

neuron peripheral and central axons suggests that Kv2.2 channels may be important for electrical

transmission in these axons. While detectable Kv2.2 remained constant in DRG neurons of older mice,

detectable Kv2.1 decreased, possibly contributing to the altered repolarization noted in DRG neurons

from aged mice (Scott, Leu, & Cinader, 1988). The differences between Kv2.1 and Kv2.2 in

somatosensory neurons are consistent with distinct roles for these Kv2 paralogs in somatosensory

neuron physiology, as seen in brain and pancreatic islets (Bishop et al., 2015; Johnston et al., 2008; Li et

al., 2013).

The expression and subcellular localization of Kv2 channels in DRG neurons are distinct from central

neurons. Kv2.1 is present in most brain neurons (Bishop et al., 2018; Bishop et al., 2015; Hwang, Fotuhi,

Bredt, Cunningham, & Snyder, 1993) while Kv2.2 is less broadly expressed (Bishop et al., 2015;

Hermanstyne et al., 2010; Hwang, Glatt, Bredt, Yellen, & Snyder, 1992; Johnston et al., 2008; Kihira et

al., 2010). In contrast to this, we find Kv2.2 is expressed in most if not all DRG neurons. We also identify

that Kv2.2 is enriched in the DRG relative to neurons in the spinal cord, suggesting that abundant Kv2.2

protein expression is a distinct feature of DRG neuron physiology. In central neurons, Kv2 channel

clusters are typically evenly distributed across neuron somata (Bishop et al., 2015; Kihira et al., 2010;

Muennich & Fyffe, 2004) and asymmetric subcellular localization of Kv2 clusters on neuron somata has

to our knowledge not been described. We find that Kv2 channels are enriched at the plasma membrane

in a subcellular region near the stem axon of some DRG neurons. Differences in Kv2 localization

between central neurons and the somatosensory neurons could arise from differences in the

extracellular environment surrounding DRG neuron somata. In central neurons, Kv2 channel clusters are

reported at cholinergic C-terminal synaptic sites, S-type synapses, and are apposed to astrocytic end

feet (Du et al., 1998; Muennich & Fyffe, 2004). However, synapses have not been reported in the DRG

and DRG neurons are surrounded by satellite glial cells instead of astrocytes (Matsuda et al., 2005).

Further studies could investigate if Kv2 clusters are localized to extracellular structures in the DRG. Kv2.2

has been identified in juxtaparanodes but not paranodes of axons in the osseous spiral lamina of the

cochlea (Kim & Rutherford, 2016) and previous work has also identified that Kv2.1 channels are

necessary for enabling ER Ca2+ uptake during electrical activity in both the soma and axon (Panzera et

al., 2022). To our knowledge, Kv2.2 channels have not been previously reported at the nodes of Ranvier

in the brain, spinal cord or DRG. Our results indicate that Kv2.2 channels are present at the

juxtaparanode and paranode of myelinated DRG neurons and also in the axons of unmyelinated DRG

neurons. The mechanism of Kv2.2 protein trafficking to the axon is unknown. Kv2.1 channels are

trafficked to the axon initial segment through a distinct secretory pathway (Jensen et al., 2017) allowing

neurons to regulate precise compartment specific localization of Kv2.1 protein. A similar mechanism for

Kv2.2 localization in DRG axons could allow neurons to independently control Kv2.2 protein expression

in the neuron soma and axon. The subcellular distribution of Kv2.1 and Kv2.2 subunits in DRG neurons

suggests they could each distinctly modulate electrical signals. We note that our observations could not

distinguish whether immunofluorescence in DRG axons could represent intracellular trafficking vesicles

of Kv2.2 proteins, or whether axonal Kv2.2 is on the membrane surface where it could modulate

electrical signals.

Kv2 channels in DRG neurons may play important roles in somatosensation and nociception. Kv2

conductances can modulate action potential repolarization, afterhyperpolarization and repetitive firing

(Liu & Bean, 2014; Zheng et al., 2019). Kv2 conductances modulate electrical transmission in rat DRG

neurons (Tsantoulas et al., 2014). Knockdown of Kv2.1 increases hypersensitivity to painful stimuli and

regulation of Kv2.1 protein expression through the epigenetic factor Cdyl modulates neuronal

excitability and nociception (Sun et al., 2022). Consistent with these studies, our results show that Kv2

channels are localized to DRG neuron axons, where they could potentially influence electrical

transmission.

Regulatory pore-forming subunits of the Kv5, Kv6, Kv8, and Kv9 "silent subunit" families obligately

assemble into heteromeric channels with Kv2 subunits (Bocksteins, 2016) and have been implicated in

nociception. Women with decreased sensitivity to labor pain were identified to have a mutation in the

Kv6.4 silent subunit that disrupts Kv2.1/Kv6.4 heteromer expression and is proposed to modulate

nociceptor signaling (Lee et al., 2020). Downregulation of the Kv9.1 silent subunit after axotomy evokes

hyperexcitability of DRG neurons (Tsantoulas et al., 2012). While both Kv2 channels are broadly

expressed in DRG neurons, transcript levels of silent subunits in mice and humans show much more

neuron subtype specificity (Tavares-Ferreira et al., 2022; Zheng et al., 2019), raising the possibility of

molecularly distinct Kv2-containing channels in somatosensory neuron subtypes.

Previous studies have identified that voltage gated ion channel expression can be substantially different

between mice and human DRG, notably, in the expression of Nav1.7 (Shiers et al., 2020). However, we

find expression of Kv2 channels to be similar in mouse and human DRG. An overall similarity is

supported by the presence of detectable Kv2 channels in most neurons, consistent with transcriptomics

data from both mice and humans (Tavares-Ferreira et al., 2022; Zheng et al., 2019). The subcellular

localization of Kv2 channels also appears similar between mice and humans. These similarities include

enrichment of Kv2 channels at the outer edge of neurons, Kv2 clusters on the neuron soma, and Kv2.2

on the stem axon, paranode and juxtaparanode. These similarities suggest that the functional roles of

Kv2 channels could be similar in both mouse and human DRG neurons.

Overall, we find attributes of Kv2 channel localization in somatosensory neurons that could point to

their functional roles in sensory information processing. The broad expression of Kv2.2 among DRG

neuron types and expansive axonal localization of Kv2.2 suggest a key role for this channel in the

somatosensory nervous system. The specialized divergence of Kv2.1 and Kv2.2 suggest the functional

roles of Kv2 subtypes in DRG neurons are overlapping yet distinct.

REFERENCES

Bishop, H. I., Cobb, M. M., Kirmiz, M., Parajuli, L. K., Mandikian, D., Philp, A. M., . . . Trimmer, J. S. (2018). Kv2 ion channels determine the expression and localization of the associated AMIGO-1 cell adhesion molecule in adult brain neurons. *Front Mol Neurosci, 11*, 1. doi:10.3389/fnmol.2018.00001

Bishop, H. I., Guan, D., Bocksteins, E., Parajuli, L. K., Murray, K. D., Cobb, M. M., . . . Trimmer, J. S. (2015). Distinct cell- and layer-specific expression patterns and independent regulation of Kv2 channel subtypes in cortical pyramidal neurons. *J Neurosci, 35*(44), 14922-14942. doi:10.1523/jneurosci.1897-15.2015

Blaine, J. T., & Ribera, A. B. (1998). Heteromultimeric potassium channels formed by members of the Kv2 subfamily. *J Neurosci, 18*(23), 9585-9593. doi:10.1523/jneurosci.18-23-09585.1998

Bocksteins, E. (2016). Kv5, Kv6, Kv8, and Kv9 subunits: No simple silent bystanders. *J Gen Physiol, 147*(2), 105-125. doi:10.1085/jgp.201511507

Bocksteins, E., Raes, A. L., Van de Vijver, G., Bruyns, T., Van Bogaert, P. P., & Snyders, D. J. (2009). Kv2.1 and silent Kv subunits underlie the delayed rectifier K+ current in cultured small mouse DRG neurons. *Am J Physiol Cell Physiol, 296*(6), C1271-1278. doi:10.1152/ajpcell.00088.2009

Cho, W. G., & Valtschanoff, J. G. (2008). Vanilloid receptor TRPV1-positive sensory afferents in the mouse ankle and knee joints. *Brain research, 1219*, 59-65. doi:https://doi.org/10.1016/j.brainres.2008.04.043

Cotella, D., Hernandez-Enriquez, B., Wu, X., Li, R., Pan, Z., Leveille, J., . . . Sesti, F. (2012). Toxic

role of K+ channel oxidation in mammalian brain. *J Neurosci, 32*(12), 4133-4144.

doi:10.1523/jneurosci.6153-11.2012

Dirajlal, S., Pauers, L. E., & Stucky, C. L. (2003). Differential response properties of IB(4)-positive

and -negative unmyelinated sensory neurons to protons and capsaicin. *J Neurophysiol,*

89(1), 513-524. doi:10.1152/jn.00371.2002

Dong, X., Han, S., Zylka, M. J., Simon, M. I., & Anderson, D. J. (2001). A diverse family of GPCRs

expressed in specific subsets of nociceptive sensory neurons. *Cell, 106*(5), 619-632.

doi:10.1016/s0092-8674(01)00483-4

Du, J., Tao-Cheng, J. H., Zerfas, P., & McBain, C. J. (1998). The K+ channel, Kv2.1, is apposed to

astrocytic processes and is associated with inhibitory postsynaptic membranes in

hippocampal and cortical principal neurons and inhibitory interneurons. *Neuroscience,*

84(1), 37-48. doi:https://doi.org/10.1016/S0306-4522(97)00519-8

Einheber, S., Zanazzi, G., Ching, W., Scherer, S., Milner, T. A., Peles, E., & Salzer, J. L. (1997). The

axonal membrane protein Caspr, a homologue of neurexin IV, is a component of the

septate-like paranodal junctions that assemble during myelination. *J Cell Biol, 139*(6),

1495-1506. doi:10.1083/jcb.139.6.1495

Fletcher, E. V., Simon, C. M., Pagiazitis, J. G., Chalif, J. I., Vukojicic, A., Drobac, E., . . . Mentis, G.

Z. (2017). Reduced sensory synaptic excitation impairs motor neuron function via Kv2.1

in spinal muscular atrophy. *Nat Neurosci, 20*(7), 905-916. doi:10.1038/nn.4561

Gravagna, N. G., Knoeckel, C. S., Taylor, A. D., Hultgren, B. A., & Ribera, A. B. (2008). Localization

of Kv2.2 protein in Xenopus laevis embryos and tadpoles. *J Comp Neurol, 510*(5), 508-

524. doi:10.1002/cne.21804

Ha, H. (1970). Axonal bifurcation in the dorsal root ganglion of the cat: a light and electron

microscopic study. *J Comp Neurol, 140*(2), 227-240. doi:10.1002/cne.901400206

He, X. H., Zang, Y., Chen, X., Pang, R. P., Xu, J. T., Zhou, X., . . . Liu, X. G. (2010). TNF-α

contributes to up-regulation of Nav1.3 and Nav1.8 in DRG neurons following motor fiber

injury. *PAIN, 151*(2), 266-279. doi:10.1016/j.pain.2010.06.005

Hermanstyne, T. O., Kihira, Y., Misono, K., Deitchler, A., Yanagawa, Y., & Misonou, H. (2010).

Immunolocalization of the voltage-gated potassium channel Kv2.2 in GABAergic neurons

in the basal forebrain of rats and mice. *J Comp Neurol, 518*(21), 4298-4310.

doi:10.1002/cne.22457

Hwang, P. M., Fotuhi, M., Bredt, D. S., Cunningham, A. M., & Snyder, S. H. (1993). Contrasting

immunohistochemical localizations in rat brain of two novel K+ channels of the Shab

subfamily. *J Neurosci, 13*(4), 1569-1576. doi:10.1523/jneurosci.13-04-01569.1993

Hwang, P. M., Glatt, C. E., Bredt, D. S., Yellen, G., & Snyder, S. H. (1992). A novel K+ channel with

unique localizations in mammalian brain: molecular cloning and characterization.

Neuron, 8(3), 473-481. doi:10.1016/0896-6273(92)90275-i

Irie, T. (2021). Essential role of somatic Kv2 channels in high-frequency firing in cartwheel cells

of the dorsal cochlear nucleus. *eNeuro, 8*(3). doi:10.1523/eneuro.0515-20.2021

Jensen, C. S., Watanabe, S., Stas, J. I., Klaphaak, J., Yamane, A., Schmitt, N., . . . Misonou, H.

(2017). Trafficking of Kv2.1 channels to the axon initial segment by a novel

nonconventional secretory pathway. *J Neurosci, 37*(48), 11523-11536.

doi:10.1523/jneurosci.3510-16.2017

Johnson, B., Leek, A. N., Sole, L., Maverick, E. E., Levine, T. P., & Tamkun, M. M. (2018). Kv2

potassium channels form endoplasmic reticulum/plasma membrane junctions via

interaction with VAPA and VAPB. *Proc Natl Acad Sci U S A, 115*(31), E7331-e7340.

doi:10.1073/pnas.1805757115

Johnston, J., Griffin, S. J., Baker, C., Skrzypiec, A., Chernova, T., & Forsythe, I. D. (2008). Initial

segment Kv2.2 channels mediate a slow delayed rectifier and maintain high frequency

action potential firing in medial nucleus of the trapezoid body neurons. *J Physiol,*

586(14), 3493-3509. doi:10.1113/jphysiol.2008.153734

Kihira, Y., Hermanstyne, T. O., & Misonou, H. (2010). Formation of heteromeric Kv2 channels in

mammalian brain neurons. *J Biol Chem, 285*(20), 15048-15055.

doi:10.1074/jbc.M109.074260

Kim, K. X., & Rutherford, M. A. (2016). Maturation of NaV and Kv channel topographies in the

auditory nerve spike initiator before and after developmental onset of hearing function.

J Neurosci, 36(7), 2111-2118. doi:10.1523/jneurosci.3437-15.2016

King, A. N., Manning, C. F., & Trimmer, J. S. (2014). A unique ion channel clustering domain on

the axon initial segment of mammalian neurons. *J Comp Neurol, 522*(11), 2594-2608.

doi:10.1002/cne.23551

Kirmiz, M., Vierra, N. C., Palacio, S., & Trimmer, J. S. (2018). Identification of VAPA and VAPB as

Kv2 channel-interacting proteins defining endoplasmic reticulum-plasma membrane

junctions in mammalian brain neurons. *J Neurosci*. doi:10.1523/jneurosci.0893-18.2018

Lee, M. C., Nahorski, M. S., Hockley, J. R. F., Lu, V. B., Ison, G., Pattison, L. A., . . . Woods, C. G.

(2020). Human labor pain is influenced by the voltage-gated potassium channel Kv 6.4

subunit. *Cell Rep, 32*(3), 107941. doi:10.1016/j.celrep.2020.107941

Legland, D., Arganda-Carreras, I., & Andrey, P. (2016). MorphoLibJ: integrated library and

plugins for mathematical morphology with ImageJ. *Bioinformatics, 32*(22), 3532-3534.

doi:10.1093/bioinformatics/btw413

Li, X. N., Herrington, J., Petrov, A., Ge, L., Eiermann, G., Xiong, Y., . . . Zhou, Y. P. (2013). The role

of voltage-gated potassium channels Kv2.1 and Kv2.2 in the regulation of insulin and

somatostatin release from pancreatic islets. *J Pharmacol Exp Ther, 344*(2), 407-416.

doi:10.1124/jpet.112.199083

Lim, S. T., Antonucci, D. E., Scannevin, R. H., & Trimmer, J. S. (2000). A novel targeting signal for

proximal clustering of the Kv2.1 K+ channel in hippocampal neurons. *Neuron, 25*(2), 385-

397. doi:10.1016/s0896-6273(00)80902-2

Liu, P. W., & Bean, B. P. (2014). Kv2 channel regulation of action potential repolarization and

firing patterns in superior cervical ganglion neurons and hippocampal CA1 pyramidal

neurons. *J Neurosci, 34*(14), 4991-5002. doi:10.1523/jneurosci.1925-13.2014

Matsuda, S., Kobayashi, N., Terashita, T., Shimokawa, T., Shigemoto, K., Mominoki, K., . . .

Fujiwara, T. (2005). Phylogenetic investigation of Dogiel's pericellular nests and Cajal's

initial glomeruli in the dorsal root ganglion. *J Comp Neurol, 491*(3), 234-245.

doi:10.1002/cne.20713

Misonou, H., Mohapatra, D. P., Menegola, M., & Trimmer, J. S. (2005). Calcium- and metabolic

state-dependent modulation of the voltage-dependent Kv2.1 channel regulates

695  neuronal excitability in response to ischemia. *J Neurosci, 25*(48), 11184-11193.

696  doi:10.1523/jneurosci.3370-05.2005

697 Muennich, E. A., & Fyffe, R. E. (2004). Focal aggregation of voltage-gated, Kv2.1 subunit-

698  containing, potassium channels at synaptic sites in rat spinal motoneurones. *J Physiol,*

699  *554*(Pt 3), 673-685. doi:10.1113/jphysiol.2003.056192

700 Newkirk, G. S., Guan, D., Dembrow, N., Armstrong, W. E., Foehring, R. C., & Spain, W. J. (2022).

701  Kv2.1 potassium channels regulate repetitive burst firing in extratelencephalic

702  neocortical pyramidal neurons. *Cereb Cortex, 32*(5), 1055-1076.

703  doi:10.1093/cercor/bhab266

704 Panzera, L. C., Johnson, B., Quinn, J. A., Cho, I. H., Tamkun, M. M., & Hoppa, M. B. (2022).

705  Activity-dependent endoplasmic reticulum Ca(2+) uptake depends on Kv2.1-mediated

706  endoplasmic reticulum/plasma membrane junctions to promote synaptic transmission.

707  *Proc Natl Acad Sci U S A, 119*(30), e2117135119. doi:10.1073/pnas.2117135119

708 Park, K. S., Mohapatra, D. P., Misonou, H., & Trimmer, J. S. (2006). Graded regulation of the

709  Kv2.1 potassium channel by variable phosphorylation. *Science, 313*(5789), 976-979.

710  doi:10.1126/science.1124254

711 Rasband, M. N., Trimmer, J. S., Schwarz, T. L., Levinson, S. R., Ellisman, M. H., Schachner, M., &

712  Shrager, P. (1998). Potassium channel distribution, clustering, and function in

713  remyelinating rat axons. *J Neurosci, 18*(1), 36-47. doi:10.1523/jneurosci.18-01-

714  00036.1998

Regnier, G., Bocksteins, E., Van de Vijver, G., Snyders, D. J., & van Bogaert, P. P. (2016). The

contribution of Kv2.2-mediated currents decreases during the postnatal development of

mouse dorsal root ganglion neurons. *Physiol Rep, 4*(6). doi:10.14814/phy2.12731

Romer, S. H., Dominguez, K. M., Gelpi, M. W., Deardorff, A. S., Tracy, R. C., & Fyffe, R. E. W.

(2014). Redistribution of Kv2.1 ion channels on spinal motoneurons following peripheral

nerve injury. *Brain research, 1547*, 1-15. doi:10.1016/j.brainres.2013.12.012

Sarmiere, P. D., Weigle, C. M., & Tamkun, M. M. (2008). The Kv2.1 K+ channel targets to the

axon initial segment of hippocampal and cortical neurons in culture and in situ. *BMC*

Neurosci, 9, 112. doi:10.1186/1471-2202-9-112

Scannevin, R. H., Murakoshi, H., Rhodes, K. J., & Trimmer, J. S. (1996). Identification of a

cytoplasmic domain important in the polarized expression and clustering of the Kv2.1 K+

channel. *J Cell Biol, 135*(6 Pt 1), 1619-1632. doi:10.1083/jcb.135.6.1619

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., . . . Cardona, A.

(2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods, 9*(7),

676-682. doi:10.1038/nmeth.2019

Scott, B., Leu, J., & Cinader, B. (1988). Effects of aging on neuronal electrical membrane

properties. *Mech Ageing Dev, 44*(3), 203-214. doi:10.1016/0047-6374(88)90022-x

Shields, S. D., Ahn, H. S., Yang, Y., Han, C., Seal, R. P., Wood, J. N., . . . Dib-Hajj, S. D. (2012).

Nav1.8 expression is not restricted to nociceptors in mouse peripheral nervous system.

PAIN, 153(10), 2017-2030. doi:10.1016/j.pain.2012.04.022

Shiers, S., Klein, R. M., & Price, T. J. (2020). Quantitative differences in neuronal subpopulations

between mouse and human dorsal root ganglia demonstrated with RNAscope in situ

hybridization. *PAIN, 161*(10), 2410-2424. doi:10.1097/j.pain.0000000000001973

Shiers, S., Sankaranarayanan, I., Jeevakumar, V., Cervantes, A., Reese, J. C., & Price, T. J. (2021).

Convergence of peptidergic and non-peptidergic protein markers in the human dorsal

root ganglion and spinal dorsal horn. *J Comp Neurol, 529*(10), 2771-2788.

doi:10.1002/cne.25122

Sun, Z. W., Waybright, J. M., Beldar, S., Chen, L., Foley, C. A., Norris-Drouin, J. L., . . . Wang, Y.

(2022). Cdyl deficiency brakes neuronal excitability and nociception through promoting

Kcnb1 transcription in peripheral sensory neurons. *Adv Sci (Weinh), 9*(10), e2104317.

doi:10.1002/advs.202104317

Tavares-Ferreira, D., Shiers, S., Ray, P. R., Wangzhou, A., Jeevakumar, V., Sankaranarayanan, I., .

. . Price, T. J. (2022). Spatial transcriptomics of dorsal root ganglia identifies molecular

signatures of human nociceptors. *Sci Transl Med, 14*(632), eabj8186.

doi:10.1126/scitranslmed.abj8186

Thapa, P., Stewart, R., Sepela, R. J., Vivas, O., Parajuli, L. K., Lillya, M., . . . Sack, J. T. (2021).

EVAP: A two-photon imaging tool to study conformational changes in endogenous Kv2

channels in live tissues. *J Gen Physiol, 153*(11). doi:10.1085/jgp.202012858

Trimmer, J. S. (1991). Immunological identification and characterization of a delayed rectifier K+

channel polypeptide in rat brain. *Proc Natl Acad Sci U S A, 88*(23), 10764-10768.

doi:10.1073/pnas.88.23.10764

Tsantoulas, C., Zhu, L., Shaifta, Y., Grist, J., Ward, J. P., Raouf, R., . . . McMahon, S. B. (2012).

Sensory neuron downregulation of the Kv9.1 potassium channel subunit mediates

neuropathic pain following nerve injury. *J Neurosci, 32*(48), 17502-17513.

doi:10.1523/jneurosci.3561-12.2012

Tsantoulas, C., Zhu, L., Yip, P., Grist, J., Michael, G. J., & McMahon, S. B. (2014). Kv2 dysfunction

after peripheral axotomy enhances sensory neuron responsiveness to sustained input.

Exp Neurol, 251, 115-126. doi:10.1016/j.expneurol.2013.11.011

Usoskin, D., Furlan, A., Islam, S., Abdo, H., Lönnerberg, P., Lou, D., . . . Ernfors, P. (2015).

Unbiased classification of sensory neuron types by large-scale single-cell RNA

sequencing. *Nat Neurosci, 18*(1), 145-153. doi:10.1038/nn.3881

Valtcheva, M. V., Copits, B. A., Davidson, S., Sheahan, T. D., Pullen, M. Y., McCall, J. G., . . .

Gereau, R. W. t. (2016). Surgical extraction of human dorsal root ganglia from organ

donors and preparation of primary sensory neuron cultures. *Nat Protoc, 11*(10), 1877-

1888. doi:10.1038/nprot.2016.111

Vierra, N. C., O'Dwyer, S. C., Matsumoto, C., Santana, L. F., & Trimmer, J. S. (2021). Regulation

of neuronal excitation-transcription coupling by Kv2.1-induced clustering of somatic L-

type Ca(2+) channels at ER-PM junctions. *Proc Natl Acad Sci U S A, 118*(46).

doi:10.1073/pnas.2110094118

Vullhorst, D., Mitchell, R. M., Keating, C., Roychowdhury, S., Karavanova, I., Tao-Cheng, J. H., &

Buonanno, A. (2015). A negative feedback loop controls NMDA receptor function in

cortical interneurons via neuregulin 2/ErbB4 signalling. *Nat Commun, 6*, 7222.

doi:10.1038/ncomms8222

Wang, H., & Zylka, M. J. (2009). Mrgprd-expressing polymodal nociceptive neurons innervate

most known classes of substantia gelatinosa neurons. *J Neurosci, 29*(42), 13202-13209.

doi:10.1523/jneurosci.3248-09.2009

Wangzhou, A., McIlvried, L. A., Paige, C., Barragan-Iglesias, P., Shiers, S., Ahmad, A., . . . Price, T.

J. (2020). Pharmacological target-focused transcriptomic analysis of native vs cultured

human and mouse dorsal root ganglia. *PAIN, 161*(7), 1497-1517.

doi:10.1097/j.pain.0000000000001866

Zheng, Y., Liu, P., Bai, L., Trimmer, J. S., Bean, B. P., & Ginty, D. D. (2019). Deep sequencing of

somatosensory neurons reveals molecular determinants of intrinsic physiological

properties. *Neuron, 103*(4), 598-616.e597. doi:10.1016/j.neuron.2019.05.039

Figure 1

Figure 1 legend

Kv2.1 and Kv2.2 protein are enriched at the outer edge of DRG neuron somas relative to

Nav1.8. Lumbar DRG section from an 11 week old male mouse labeled with antibodies which

target Kv2.1, Kv2.2 or Nav1.8. *A*, Anti-Kv2.1 (magenta) and anti-Kv2.2 (green)

immunofluorescence in a lumbar DRG section. Scale bar is 100 μm. *B*, Anti-Kv2.1, anti-Kv2.2 and

anti-Nav1.8 immunofluorescence from box shown in A. Arrows indicate prominent localization

of anti-Kv2 immunofluorescence at the edge of DRG neuron somas. In merge image anti-Kv2.1,

anti-Kv2.2 and anti-Nav1.8 immunofluorescence is magenta, green and blue respectively. Scale

bar is 20 μm. *C*, Representative ROIs that encompass the outer edge of DRG neurons (arrow 1)

and the region just inside the outer edge (arrow 2). *D*, Ratio of anti-Kv2.1, anti-Kv2.2 or anti-

Nav1.8 immunofluorescence from outer and inner ROIs for individual neurons from image in A.

Bars represent mean.

Figure 2

Figure 2 legend

Kv2.1 protein is detectable in most mouse DRG neurons. *A*, WT (top) and Kv2.1 KO (bottom)

DRG sections immunolabeled for Kv2.1 (magenta) and BIII tubulin (white). Images were taken

with identical imaging settings and are set to the same brightness and contrast. Scale bars are

100 μm. *B*, Distribution of fluorescence intensity from WT (black) and Kv2.1 KO (red) neurons.

Data represent fluorescence intensity of 476 WT neurons from 10 DRG sections from 1 mouse

or 235 Kv2.1 KO neurons from 5 DRG sections from 1 mouse. DRG sections were taken from 7

week old female mice and are from the 13th thoracic DRG. Images shown in A represent one

section from WT and Kv2.1 KO mice used in this data set. *C*, Distribution of BIII tubulin -

fluorescence intensity from the same WT (black) and Kv2.1 KO (red) neurons shown in B. *D*,

Mean fluorescence intensity of Kv2.1 KO neurons normalized to WT neurons labeled with anti-

Kv2.1, anti-Kv2.2 and anti-BIII tubulin antibodies. Each point represents one Kv2.1 KO mouse

normalized to one age and sex matched WT mouse which was stained simultaneously and

imaged with identical microscopy settings. The color of each point represents the same mouse

and purple points represent data from the mouse whose DRG immunofluorescence data are

shown in A, B and C. p-values are calculated by a Students T-test and compares if points have a

significant difference from 1. *E*, Kv2.1 KO data shown in B fit with a log normal distribution (red

fit). *F*, WT data shown in B fit with the Kv2.1 KO distribution (red fit) where width and mean

were constrained to the Kv2.1 KO distribution and amplitude was unconstrained (equation 1).

Red dotted line represents the mean of the Kv2.1 KO distribution. Only WT data to the left of

red dotted line was used for the fit. *G*, Percent of neurons with detectable Kv2.1 protein of 5

mice (1 female and 4 males). Point colors correspond to the WT mice analyzed in D. All mice

were compared to age and sex matched Kv2.1 KO mice.

 Figure 3

Figure 3 legend

Detectable Kv2.2 protein is present in most if not all mouse DRG neurons. **A**, WT (top) and

Kv2.2 KO (bottom) DRG sections from the 13th thoracic DRG in 7 week old male mice

immunolabeled for Kv2.2 (green) and NF200 (white). Identical imaging and display settings.

Scale bars are 100 μm. **B**, Distribution of fluorescence intensities from WT (black) and Kv2.2 KO

(red) neurons. 672 WT neurons from 1 mouse. 398 Kv2.2 KO neurons from 1 mouse. Data from

same mice shown in A. **C**, Distribution of anti-NF200 immunofluorescence intensity from the

same WT (black) and Kv2.2 KO (red) neurons shown in B. **D**, Mean fluorescence intensity of

Kv2.2 KO neurons normalized to WT neurons labeled with anti-Kv2.1, anti-Kv2.2 and anti-NF200

antibodies. Each point represents one Kv2.2 KO mouse normalized to one age and sex matched

WT mouse which was stained simultaneously and imaged with identical microscopy settings.

The color of each point represents the same mouse and purple points represent data from the

male mouse whose DRG immunofluorescence data are shown in A, B and C. Missing points in

anti-Kv2.1 column are because some sections were not labeled with anti-Kv2.1 antibodies. p-

values are calculated by a Students T-test and compares if points have a significant difference

from 1. **E**, Kv2.2 KO data shown in B fit with a log normal distribution (red fit). **F**, WT data shown

in B fit with the Kv2.2 KO distribution (red fit) where width and mean were constrained to the

Kv2.2 KO distribution and amplitude was unconstrained (equation 1). Red dotted line

represents the mean of the Kv2.2 KO distribution. Only WT data to the left of red dotted line

was used for the fit. **G**, Percent of neurons with detectable Kv2.2 protein of 8 mice (7 males and

851   1 female). Point colors correspond to the WT mice analyzed in D. All mice were compared to

852   age and sex matched Kv2.2 KO mice.

 Figure 4

Figure 4 legend

Detectable Kv2.1 protein decreases in older mice while detectable Kv2.2 does not. **A**, DRG

sections from the 13th thoracic DRG in 7 week (left) and 50 week (right) old mice

immunolabeled for Kv2.1. Vertical bar on right is pseudo coloring key for pixel intensity.

Identical imaging and display settings. Scale bars are 100 μm. **B**, Distribution of fluorescence

intensities from 7 week old WT (black) and 7 week old Kv2.1 KO (red) neurons. 609 WT neurons

from 1 mouse. 367 Kv2.1 KO neurons from 1 mouse. **C**, Distribution of fluorescence intensities

from 50 week old WT (black) and 50 week old Kv2.1 KO (red) neurons. 793 WT neurons from 1

mouse. 378 Kv2.1 KO neurons from 1 mouse. **D**, Percent of neurons with detectable Kv2.1

protein in 7-16 week old and 50 week old mice. Data from 7-16 week old mice is the same data

in Figure 2. **E**, DRG sections from the 13th thoracic DRG in 7 week (left) and 50 week (right) old

mice immunolabeled for Kv2.2. Vertical bar on right is pseudo coloring key for pixel intensity.

Identical imaging and display settings. Scale bars are 100 μm. **F**, Distribution of fluorescence

intensities from 7 week old WT (black) and 7 week old Kv2.2 KO (red) neurons. 746 WT neurons

from 1 mouse. 717 Kv2.2 KO neurons from 1 mouse. Data from same mice shown in E. **G**,

Distribution of fluorescence intensities from 50 week old WT (black) and 50 week old Kv2.2 KO

(red) neurons. 671 WT neurons from 1 mouse. 398 Kv2.2 KO neurons from 1 mouse. Data from

same mice shown in E. **H**, Percent of neurons with detectable Kv2.2 protein in 7-24 week old

and 50 week old mice. Data is the same data from Figure 3 where mice were separated into a

young group (7-24 weeks) and an old group (50 weeks).

 Figure 5

 Figure 5 legend

Kv2 channels are expressed at the surface membrane of DRG neurons. **A**, Fluorescence of live

dissociated DRG neurons excited at 594 nm before (top left) and after (bottom left) application

of 100 nM GxTX-594. Fluorescence from membrane marker WGA-405 before (top middle) and

after application of 100 nM GxTX-594 (bottom middle). Merge image shows 594 excitation

fluorescence (magenta) and 405 excitation fluorescence (green). Scale bar 20 μm. **B**,

Dissociated DRG neurons from WT (top) and Kv2.1/Kv2.2 DKO (bottom) mice before and after

the application of GxTX-594, left panel and middle panel respectively. Arrows in middle panel

indicate location of soma membrane based on WGA-405 fluorescence. Right images are merge

of 594 excitation fluorescence (magenta) and 405 excitation fluorescence (green) after

application of 100 nM GxTX-594. Scale bars 100 μm. **C**, Example of WGA-405 fluorescence (left)

used in watershed segmentation (middle) to generate annulus ROI (right) used to analyze

fluorescence intensity at the membrane. **D**, Distribution of fluorescence intensity from WT

(black) and Kv2.1/Kv2.2 DKO (red) neurons. Data represents the fluorescence intensity of 326

WT neurons from 1 mouse or 271 Kv2.1/Kv2.2 DKO neurons from 1 mouse. DRG from all levels

of the spinal cord were pooled. **E**, Kv2.1/Kv2.2 DKO data shown in D fit with a log normal

distribution (red fit). **F**, WT data shown in D fit with the Kv2.1/Kv2.2 DKO distribution (red fit)

where width and mean were constrained to the Kv2.1/Kv2.2 DKO distribution and amplitude

was unconstrained (equation 1). Red dotted line represents the mean of the Kv2.1/Kv2.2 DKO

distribution. Only WT data to the left of red dotted line was used for the fit. **G**, Percent of

neurons with detectable surface Kv2 protein, from an experiment where one WT mouse was

compared to one DKO mouse and an identical experiment where two WT mice were compared

to one DKO mouse (N=3 WT mice N=2 Kv2.1/Kv2.2 DKO mice).

 Figure 6

Figure 6 legend

DRG neurons have enriched Kv2.2 protein compared to neurons in the spinal cord. *A*, Anti-

Kv2.1 (magenta) and anti-Kv2.2 (green) immunofluorescence in a spinal cord section from the

2nd lumbar vertebra (left). Anti-Kv2.1 immunofluorescence (right top) and anti-Kv2.2

immunofluorescence (right bottom). Arrows show neurons in the spinal cord with punctate

anti-Kv2.1 immunofluorescence. Arrow heads show neurons in the spinal cord with anti-Kv2.2

immunofluorescence. Scale bars are 500 µm. *B*, Anti-Kv2.1 immunofluorescence from individual

neurons (circles) in the DRG and ventral horn normalized to the average fluorescence intensity

of neurons in the ventral horn. Diamonds to the right of data represent the average intensity in

DRG neurons of individual mice. Significant differences from 1 were calculated for individual

mice using Students t-test. N = 5 mice n = 295 in DRG and n = 200 in ventral horn. *C*, Identical

analysis shown in B with anti-Kv2.2 immunofluorescence.

Figure 7

Figure 7 legend

Kv2 channels form clusters on DRG neuron somas and stem axons that are distinct from Kv2

channel clusters on ventral horn neurons. ***A***, Z-projection of anti-Kv2.1 and anti-Kv2.2

immunofluorescence in a ventral horn neuron from the 1st lumbar vertebra of a 7 week old

male mouse. ***B***, Z-projection of anti-Kv2.1 and anti-Kv2.2 immunofluorescence in DRG neurons

from the same mouse and section of the spinal column as neuron shown in A. Inset is

enlargement of the Kv2.1 donut cluster in dotted box. ***C***, Z-projection of anti-Kv2.1 and anti-

Kv2.2 immunofluorescence in DRG neurons from the same mouse as A and B. Inset is

enlargement of the Kv2.2 donut cluster in dotted box. ***D***, Z-projection of anti-Kv2.1

immunofluorescence in DRG neuron from the 13th thoracic DRG of a 24 week old male mouse.

E, Z-projection of anti-Kv2.1 and anti-Kv2.2 immunofluorescence in DRG neuron from same

mouse in D. ***F***, Exemplar ROI for analyzing localization of Kv2 channel density relative to stem

axon. Same image as E. ROI line width is 1.24 μm. Numbers along line indicate approximate

distance from stem axon normalized to the midpoint of the line. ***G***, Anti-Kv2.1 (magenta) and

anti-Kv2.2 (green) immunofluorescence intensity along the ROI shown in F. Distance along the

line was normalized such that the stem axon is 0 and the midpoint of the line is 1. ***H***, Analysis of

the relative distance from the stem axon of the max anti-Kv2.1 or anti-Kv2.2

immunofluorescence. Dotted line represents the middle of neurons relative to the stem axon.

In all images arrows indicate asymmetrical clusters of Kv2 channels while arrowhead indicate

the apparent stem axons. Display settings are not identical between images. Scale bars are 20

μm.

Figure 8

Figure 8 legend

Kv2.2 channels are expressed in peripheral axons of DRG neurons at the paranodes and

juxtaparanodes. *A*, WT (top) and Kv2.2 KO (bottom) sections containing the DRG, peripheral

and central axons from the 1st lumbar DRG in age and sex matched 7 week old mice

immunolabeled for NF200 (white). Scale bar is 1 mm. *B*, High magnification z-projection of anti-

Kv2.2 and anti-NF200 immunofluorescence from box 1 in A of WT and Kv2.2 KO mice. Arrows

indicate myelinated axons which show prominent anti-Kv2.2 immunofluorescence. Scale bars

are 20 μm. *C*, High magnification z-projection of anti-Kv2.2 and anti-NF200

immunofluorescence from box 2 in A of WT and Kv2.2 KO mice. Arrows indicate myelinated

axons which show prominent anti-Kv2.2 immunofluorescence. Scale bars are 20 μm. *D*, High

magnification z-projection of anti-Kv2.2 immunofluorescence and MrgprD-GFP fluorescence in

the peripheral axons of the 12th thoracic DRG of a 13 week old MrgprD-GFP mouse. Arrow

indicates anti-Kv2.2 immunofluorescence on a GFP+ axon. Scale bar is 10 μm.

Figure 9

Figure 9 legend

Kv2.2 is expressed at the paranodes and juxtaparanodes of DRG neuron axons. **A**, Kv2.2 KO

(top) and WT (middle and bottom) sections containing the peripheral axons from the 12th

thoracic DRG in 28 week old mice immunolabeled for Kv2.2, Caspr and Kv1.2. Middle panels are

an exemplar of prominent Kv2.2 immunofluorescence in the paranodes of DRG neurons and

bottom panels are an exemplar of prominent Kv2.2 clusters in the juxtaparanodes. Scale bars

are 5 µm. **B**, Analysis of anti-Kv2.2 immunofluorescence intensity in the paranodes and

juxtaparanodes of age and sex matched WT and Kv2.2 KO mice. Individual points represent a

single paranode or juxtaparanode identified by anti-Caspr or anti-Kv1.2 immunofluorescence.

Figure 10

Figure 10 legend

Anti-Kv2 immunofluorescence intensities are non-uniform across DRG neuron subtypes. ***A***,

Exemplar z-projection of enrichment of anti-Kv2.1 (magenta) or anti-Kv2.2 (green)

immunofluorescence in neighboring neurons. DRG section is from a 10 week old male mouse.

Scale bar is 10 μm. **B**, Top images show anti-Kv2.1 (magenta) and anti-Kv2.2 (green)

immunofluorescence in DRG sections where subpopulation specific markers were used to

identify, from left to right, non-peptidergic nociceptors, peptidergic nociceptors, myelinated

neurons and proprioceptors. Fluorescence from specific markers is shown in bottom panels.

Arrows indicate four exemplar neurons that have clear positivity for each subpopulation

identified by fluorescence in lower panels. CGRP and NF200 subpopulations were identified

using anti-CGRP and anti-NF200 antibodies while MrgprD-GFP and PV-tdTomato

subpopulations were from transgenic mouse lines. Scale bars are 50 μm. **C**, Scatter plot of anti-

Kv2.1 and anti-Kv2.2 immunofluorescence of individual neurons. Each point represents one

neuron. Magenta circle highlights the subpopulation of neurons that have high anti-Kv2.1 but

low anti-Kv2.2 immunofluorescence while the green circle highlights the subpopulation of

neurons that have high anti-Kv2.2 but low anti-Kv2.1 immunofluorescence. **D**, Ranked anti-

Kv2.1 immunofluorescence (magenta points) or ranked anti-Kv2.2 immunofluorescence (green

points) of individual neurons from subpopulations shown in B. Only neurons that were positive

for each marker are shown. Each point represents one neuron. MrgprD population N = 4 mice,

CGRP population N = 3 mice, NF200 population N = 3 mice and PV population N = 2 mice.

Figure 11

 Figure 11 legend

 Kv2 channel expression and localization in human DRG neurons is similar to mice. *A*,

 Immunofluorescence from human DRG neurons labeled with anti-Kv2.1 and anti-Kv2.2

 antibodies. Autofluorescence attributed to lipofuscin is labeled in right panel while apparent

 Kv2.1 and Kv2.2 protein are labeled in left and middle panel respectively. Scale bar is 50 μm. *B*,

 Z-projection of anti-Kv2.1 (left) and anti-Kv2.2 immunofluorescence (right) of human DRG

 neuron somata. Arrows indicate asymmetric distribution of Kv2.1 and Kv2.2 clusters on neuron

 somata. Apparent lipofuscin autofluorescence was not present in these neurons. Scale bars are

 20 μm. *C*, Z-projection of anti-Kv2.1 (left) and anti-Kv2.2 immunofluorescence (right) of human

 DRG neuron somata. Insets show expansion of dotted line boxes which highlight donut shaped

 Kv2.1 and Kv2.2 clusters. Arrow in right image shows anti-Kv2.2 immunofluorescence on the

 apparent stem axon of DRG neuron. Scale bar is 20 μm. *D*, Z-projection of anti-Kv2.2 (upper

 left), anti-Caspr (upper right) and anti-Kv1.2 (bottom left) immunofluorescence of human DRG

 axon. Arrows in merge image (lower right) indicate juxtaparanodes and arrowheads indicate

 paranodes. Scale bar is 10 μm.

1008 Table 1

Figure	Genotype	Number of Mice (N)	Sections/plates Analyzed	Neurons Analyzed (n)	Age (Weeks)	Sex	Level of spinal column	Total Neurons analyzed
1	WT	1	1	124	11	M	L	115
2	WT	5	3,4,10,8,4	52,172,476,437,134,	13,7,7,7,7	M,M,F,M,M	L,T12,T12,T12,T12	1271
	Kv2.1 KO	5	4,7,5,4,5	232,124,236,293,239,	13,7,7,7,7	M,M,F,M,M	L,T12,T12,T12,T12	1124
3	WT	8	4,11,4,6,4,4,4,6	238,673,126,1227,706,748,602,572,	50,50,50,7,7,24,7,28	M,M,M,M,M,M,M,F	L,T12,T12,T12,T12,T12,L1,T12	4892
	Kv2.2 KO	8	3,6,7,4,4,4,4,5	359,400,209,853,913,719,729,745,	50,50,50,7,7,24,7,28	M,M,M,M,M,M,M,F	L,T12,T12,T12,T12,T12,L1,T12	4927
4	WT	10	7,5,6,9,8	444,268,167,669,600,	50,50,50,50,50	F,F,F,M,M	T12,T12,T12,T12,T12	3419
	Kv2.1 KO	7	5,6	298,324,	50,50	M,M	T12,T12	622
	WT	8	same as data in figure 3					
	Kv2.2 KO	8	same as data in figure 3					
5	WT	3	4,16,17,	104,379,328,	18,19,19	F,F,F	L,L,L	811
	Kv2.1 Kv2.2 DKO	2	5,14,	108,274,	18,19	F,F	L,L	382
6	WT DRG	5	1,1,1,1,1	139,51,30,40,38	10,10,10,8,8	M,M,F,F,M	T12,T8,L1,L1,L1	298
	WT Ventral Horn	5	1,1,1,1,1	75,51,23,24,28	10,10,10,8,8	M,M,F,F,M	T12,T8,L1,L1,L1	201
7	WT	2	2	N/A	7,24	M,M	L1,T12	N/A
8	WT	1	1	N/A	7	M	L1	N/A
	Kv2.2 KO	1	1	N/A	7	M	L1	N/A
	MrgprD-GFP	1	1	N/A	13	F	T12	N/A
9	WT	1	1	N/A	28	F	T12	N/A
	Kv2.2 KO	1	1	N/A	28	F	T12	N/A
10	WT	1	1	N/A	10	M	L5	N/A
	MrgprD-GFP	4	6,5,9,14	852,632,842,1392,	13,20,8,8	F,M,F,F	T12,L1,L1,L1	3718
	WT	5	7,4,6,7,4	482,464,592,641,166,	7,10,24,19,19	F,M,M,F,F	L1,T12,T12,T12,T12	2345
	PV-tdTomato	2	15,9	1978,1158,	8,8	F,M	L1,L1	3136
1-1	WT	1	1	N/A	10	M	L5	N/A
1-2	WT	1	1	N/A	28	M	L	N/A

1-3	WT	1	3	125	13	M	L	125
2-1	WT	1	33	5399	8	M	T11 and T12	5399
2-2	MrgprDGFP	1	1	N/A	13	F	T12	N/A
2-3	MrgprDGFP	4	6,15,9,14	5,568,769,051,487	14,20,8,8	F,M,F,F	L1,L1,L1,L1	3824
	WT	2	7,5	771,478	24,8	M,F	L1,L1	1249
6-1	WT DRG	4	1,1,1,1	25,30,28,35	7,7,8	M,F,F,F	L1,L1,L1,L1	118
	WT Ventral Horn	4	1,1,1,1	17,19,26,19	7,7,8	M,F,F,F	L1,L1,L1,L1	81
7-1	MrgprDGFP	1	1	N/A	14	F	T12	N/A
	WT	2	1,1	N/A	11,17	M,M	L,L	N/A
8-1	WT	1	1	N/A	7	F	T12	N/A
	Kv2.1 KO	1	1	N/A	7	F	T12	N/A

1009

Table 1 legend

Detailed information on mice used in each figure. In columns "Sections/Plates Analyzed",

"Neurons Analyzed", "Age", "Sex" and "Level of Spinal Column" information for each mouse is

separated by commas. In column "Level of Spinal Column" L1 refers to the first lumbar DRG and

T12 refers to the twelfth thoracic DRG and if only L is shown the specific DRG is not known but

the DRG is from the lumbar region.

1016

1017   Table 2

Antigen	Antibody name	Species/isotype	Manufaturer information	Concentration used	RRID	Figures
Kv2.1	K89/34	Mouse IgG1 mAb	NeuroMab	6.7 ug/mL	AB_10672253	1, 2, 3, 4,6-1, 7, 8-1, 10, 11, 11-1, 11-2
Kv2.2	Kv2.2c	Rabbit pAb	In-house Trimmer laboratory	1:100 purified from strip assay	AB_2801484	3, 4, 6-1, 7, 8, 9, 10, 11, 11-1, 11-2,11-3
Nav1.8	N134/12	Mouse IgG2a mAb	NeuroMab	1:5 tissue culture supernatant	AB_10672261	1,11-1
Kv2.1	K89/34R	Mouse IgG2a mAb	In-house Trimmer laboratory	1:5 tissue culture supernatant	AB_2315768	11-1
beta III Tubulin	NA	Rabbit pAb	Abcam ab18207	0.33 ug/mL	AB_444319	2, 8-1
Kv2.2	N372B/60	Mouse IgG2b mAb	NeuroMab	1 ug/mL	AB_2315868	1, 1-3, 6, 10
NF200	NA	chicken pAb	Abcam ab4680	0.2 ug/mL	AB_304560	3, 8, 10
Caspr	K65/35	Mouse IgG1 mAb	NeuroMab	1:5 tissue culture supernatant	AB_2877274	9
Kv1.2	K14/16	Mouse IgG2b mAb	NeuroMab	1:5 tissue culture supernatant	AB_2877295	9,11
CGRP	NA	Rabbit pAb	Immunostar	1:1000	AB_572217	10
DsRed	NA	Rabbit pAb	Takara Cat#632543	0.17 ug/mL	AB_2307319	10
Nav1.7	N68/6	Mouse IgG1 mAb	NeuroMab	1:5 tissue culture supernatant	AB_2184355	11-1,11-3

1018

1019

1020   Table 2 legend

1021   Detailed information on antibodies used throughout the manuscript. In column "Concentration

1022   Used" if only dilution is given concentration was unknown however, tissue culture supernatant

1023   concentrations of antibodies typically range between 15-30 µg/mL. **SUPPLEMENTAL FIGURES**

1024   **Figure 1-1**

1025

1026   **Figure 1-1 legend**

1027   Off target mouse anti-IgG1, IgG2b and IgG2a immunofluorescence increases with fixation time.

1028   Images of DRG sections labeled with fluorescently tagged antibodies which target IgG1 (top),

1029   IgG2b (middle) and IgG2a (bottom) after DRG were fixed in ice cold 4% PFA for 1, 3 and 12

1030   hours. DRG sections are from the same mouse. Scale bars are 100 μm.

1031

1032 **Figure 1-2**

1033

Figure 1-2 legend

Pre-incubation of mouse DRG sections in IgG H+L Fab fragments reduces off target secondary

antibody labeling. Representative images of DRG sections from the same DRG treated with

increasing concentrations (left to right) of IgG H+L Fab fragment and the same concentration of

secondary antibody used in experiments throughout this study. Images were taken with

identical imaging settings and are set to the same brightness and contrast. Scale bars are 100

μm.

Figure 1-3

1042

Figure 1-3 legend

Kv2.1 and Kv2.2 protein are enriched at the outer edge of DRG neuron somas relative to TRPV1.

A, Anti-Kv2.1, anti-Kv2.2 and anti-TRPV1 immunofluorescence from lumbar DRG neurons.

Prominent cytoplasmic anti-TRPV1 immunofluorescence was observed in a subset of small

diameter neurons. In merge image anti-Kv2.1, anti-Kv2.2 and anti-TRPV1 immunofluorescence

are magenta, green and blue respectively. Scale bar is 20 μm. **B**, Ratio of average anti-Kv2.1,

anti-Kv2.2 or anti-TRPV1 immunofluorescence from outer and inner ROIs for individual

neurons.

1051

1052   **Figure 2-1**

1053

Figure 2-1 legend

Kv2 antibodies used in knockout experiments are at saturating concentrations. **A**,

Concentration response of immunofluorescence from sections labeled with anti-Kv2.1 antibody

used in Figure 2. Blue line is a Hill fit of the data. 1:1 = 2 sections, 1:25 = 3 sections, 1:125 = 2

sections, 1:625 = 3 sections **B**, Concentration response of immunofluorescence from sections

labeled with anti-Kv2.2 antibody used in Figure 3. Blue line is a Hill fit of the data. 1:33 = 2

sections, 1:100 = 10 sections, 1:300 = 4 sections, 1:900 = 4 sections, 1:2700 = 3 sections

1061

1062   **Figure 2-2**

1063

Figure 2-2 legend

A method to sample neurons in DRG imaging data using watershed segmentation identifies the

outer region of neurons. **A**, Image of anti-Kv2.1, anti-Kv2.2, anti-βIII tubulin

immunofluorescence and MrgprD-GFP fluorescence. **B**, Same image as A with anti-Kv2.1

immunofluorescence channel removed as this is an example of processing data for Kv2.1 KO

analysis. *C*, Grayscale image of average fluorescence from all three channels shown in B.

Gaussian and median filters were applied to image to improve watershed segmentation. *D*,

Watershed segmentation of image in C using the MorphoLibJ Morphological Segmentation

Plugin in Fiji. *E*, Example of manually drawn boundary that encompasses the neuron somas in

the DRG so that only these watershed lines are selected. *F*, Selected ROIs from watershed

segmentation (yellow). ROIs were excluded based on roundness and size using the Analyze

Particles tool in Fiji. *G*, ROIs in F overlaid on DRG image showing that some ROIs are selecting

regions that do not contain neurons (arrow heads) or are selecting multiple neurons (arrows).

H, ROIs after processing using an in house R script which removes most, but not all, ROIs that

do not contain neurons (arrow heads) and ROIs that contain two neurons (arrows). Each

experiment performed was done alongside controls where the primary antibodies were

omitted and fluorescence from these control sections was used by the in house R script to

identify and remove ROIs that do not contain neurons. *I*, Example of automatically generated

annulus that encompasses the outer edge of the soma. Scale bars are 100 μm.

Figure 2-3

1084

Figure 2-3 legend

Method used to estimate percent of neurons expressing Kv2.1 and Kv2.2 reliably predicts the

percentage of neurons that express GFP in MrgprD-GFP mice. **A**, MrgprD-GFP (top) and WT

(bottom) DRG sections immunolabeled for BIII tubulin (white). Images were taken with identical

imaging settings and are set to the same brightness and contrast. Scale bars are 100 μm. **B**,

Distribution of fluorescence intensity from MrgprD-GFP (black) and WT (red) neurons. Data

represents the fluorescence intensity of 905 MrgprD-GFP neurons from 9 DRG sections from 1

mouse or 477 WT neurons from 5 DRG sections from 1 mouse. DRG sections were taken from 7

week old female mice and are from the 1st lumbar DRG. *C*, WT data shown in B fit with a log

normal distribution (red fit). *D*, MrgprD-GFP data shown in B fit with the WT distribution (red

fit) where width and mean were constrained to the WT distribution and amplitude was

unconstrained (equation 1). Only MrpgrD-GFP data to the left of the mean intensity of WT

neurons (red dotted line) was used for the fit. *E*, Percent of neurons with detectable GFP

protein of 4 mice (3 females 1 male). All DRG sections were taken from the 1st lumbar DRG.

Figure 6-1

Figure 6-1 legend

DRG neurons have enriched Kv2.2 protein compared to neurons in the ventral horn. *A*, Anti-

Kv2.1 (magenta) and anti-Kv2.2 (green) immunofluorescence in a spinal cord section from the

13th thoracic vertebra (left). Anti-Kv2.1 immunofluorescence (right top) and anti-Kv2.2

immunofluorescence (right bottom). Arrow heads show neurons in the spinal cord with anti-

Kv2.2 immunofluorescence. Scale bars are 500 µm. *B*, Anti-Kv2.1 immunofluorescence from

individual neurons (circles) in the DRG and ventral horn normalized to the average fluorescence

intensity of neurons in the ventral horn. Diamonds to the right of data represent the average

intensity in the DRG of individual mice. Significant differences from 1 were calculated for

individual mice using Students t-test. N = 4 mice n = 116 in DRG and n = 77 in ventral horn. *C*,

Identical analysis shown in B with anti-Kv2.2 immunofluorescence.

1112

1113   **Figure 7-1**

1114

Figure 7-1 legend

Kv2 channels are expressed on the stem axon of mouse DRG neurons. **A**, Z-projection with anti-

Kv2.1 and anti-Kv2.2 immunofluorescence on the stem axon of a neuron in the DRG of a

MrgprD-GFP mouse. **B**, Gallery of z-projected images of DRG neurons with anti-Kv2.1 and/or

anti-Kv2.2 immunofluorescence on stem axons. Arrows indicate stem axons. Scale bars are 10

μm

Figure 8-1

Figure 8-1 legend

Kv2.1 channels were not detected in peripheral axons of DRG neurons. *A*, WT (top) and Kv2.1

KO (bottom) sections containing the DRG and peripheral axons from the 12[th] thoracic DRG in

age and sex matched 7 week old mice immunolabeled for βIII tubulin (white). Scale bar is 500

μm. *B*, High magnification z-projection of anti-Kv2.1 and anti-βIII immunofluorescence from box

in A of WT and Kv2.1 KO mice. Scale bars are 20 μm.

1129

1130   **Figure 11-1**

1131

Figure 11-1 legend

Fluorescence from human DRG neurons labeled with ion channel targeting antibodies is distinct

from human DRG neurons where ion channel targeting antibodies were omitted. **A**, Top:

Immunofluorescence from human DRG section labeled with anti-Kv2.1, anti-Kv2.2 and anti-

Nav1.8 antibodies. Bottom: Fluorescence from human DRG section where the primary

antibodies are omitted. Arrows in top and bottom images indicate examples of

autofluorescence from apparent intracellular lipofuscin. Arrow heads in top image identify anti-

Nav1.8 immunofluorescence. Images on the right are fluorescence from each fluorescence

channel of the top and bottom images. Number next to target protein label represents

excitation wavelength. DRG sections from top and bottom images are from the same DRG.

Scale bars are 500 μm. **B**, Left: Immunofluorescence from human DRG section labeled with anti-

Nav1.7 antibody. Right: Fluorescence from human DRG section where the primary antibody has

been omitted. Number next to target protein label represents excitation wavelength. DRG

sections in left and right images are from the same DRG. Scale bars are 500 μm. **C**, Exemplar

manually drawn ROI to analyze fluorescence intensity in human DRG neurons that omits

apparent lipofuscin autofluorescence. Scale bar is 50 μm. **D**, Distribution of fluorescence

intensity of human DRG neurons labeled with an anti-Kv2.1 antibody (black) or when the anti-

Kv2.1 antibody was omitted (red). Data represents the fluorescence intensity of 293 neurons

labeled with anti-Kv2.1 antibody or 73 neurons where the anti-Kv2.1 antibody was omitted. **E**,

Distribution of fluorescence intensity of human DRG neurons labeled with anti-Kv2.2 antibody

(black) or when the anti-Kv2.2 antibody was omitted (red). Data represents the fluorescence

intensity of 293 neurons labeled with anti-Kv2.2 antibody or 73 neurons where the anti-Kv2.2

antibody was omitted. **F**, Distribution of fluorescence intensity of human DRG neurons labeled

with anti-Nav1.7 antibody (black) or when the anti-Nav1.7 antibody was omitted (red). Data

represents the fluorescence intensity of 99 neurons labeled with anti-Nav1.7 antibody or 99

neurons where the anti-Nav1.7 antibody was omitted. **G**, Distribution of fluorescence intensity

of human DRG neurons labeled with anti-Nav1.8 antibody (black) or when the anti-Nav1.8

antibody was omitted (red). Data represents the fluorescence intensity of 293 neurons labeled

with anti-Nav1.8 antibody or 73 neurons where the anti-Nav1.8 antibody was omitted. **H**,

Fluorescence intensity of human neurons labeled with both anti-Kv2.1 and anti-Kv2.2

antibodies. Individual points represent individual neurons.

Figure 11-2

1164

Figure 11-2 legend

Kv2 channels are enriched at the outer edge of human DRG neurons. *A*, Exemplar z-projection

of anti-Kv2.1 immunofluorescence enriched at the outer surface of a human DRG neuron. Scale

bar is 20 μm. *B*, Exemplar z-projection of anti-Kv2.2 immunofluorescence enriched at the outer

surface of a human DRG neuron. Scale bar is 20 μm. *C*, Exemplar z-projection of asymmetrical

anti-Kv2.2 immunofluorescence enriched at the outer surface of a human DRG neuron. Scale

bar is 20 μm. *D*, Exemplar z-projection of anti-Kv2.1 and anti-Kv2.2 immunofluorescence both

enriched at the outer surface of a human DRG neuron soma. Scale bar is 20 μm.

1173

1174 **Figure 11-3**

1175

1176 **Figure 11-3 legend**

1177 Z-projection of anti-Kv2.2 and anti-Nav1.7 immunofluorescence in a human DRG neuron soma

1178 and stem axon. Arrow in merge indicates the stem axon of the DRG neuron. Apparent lipofuscin

1179 autofluorescence is labeled in left panel. Scale bar is 20 μm.