Distinct cellular expression and subcellular localization of Kv2 voltage-gated K⁺ channel subtypes in

dorsal root ganglion neurons conserved between mice and humans

Robert G. Stewart[1], Miriam Camacena[1], Bryan A. Copits[2], Jon T. Sack[1,3]

Affiliations:

[1] Department of Physiology and Membrane Biology, University of California Davis,

Davis, CA 95616, USA.

[2] Washington University Pain Center, Washington University School of Medicine, St. Louis, MO 63110,

USA; Department of Anesthesiology, Washington University School of Medicine, St. Louis, MO 63110,

USA.

[3] Department of Anesthesiology and Pain Medicine, University of California Davis,

Davis, CA 95616, USA.

Corresponding author email: jsack@ucdavis.edu

Number of figures: 10

Number of supplemental figures: 15

Number of tables: 2

Conflict of interest statement: The authors declare no competing conflicts of interest.

Data availability: The data that support the findings of this study are available upon request from the

authors. Code used to analyze this data in ImageJ and R is available at https://github.com/SackLab/DRG-

Image-Processing

Running Title: Kv2 localization in DRG neurons

Keywords: Dorsal root ganglion, Voltage gated ion channels, Kv2.1, Kv2.2, Somatosensory neurons

Acknowledgments:

We thank James S. Trimmer (University of California, Davis) for generous provision of antibodies,

numerous discussions, and feedback on the manuscript. We thank Michael Ferns (University of

California, Davis) for feedback on the manuscript. We thank the human tissue donors and their families

for their generous donations to science. We thank Jiwon Yi, Zack Bertels, Maria Payne, and Jon Lemen

for performing hDRG extractions. GxTX was synthesized at the Molecular Foundry of the U.S.

Department of Energy under contract DE-AC02-05CH11231. This research was supported by U.S.

National Institutes of Health grant R01NS096317 to J.T. Sack and the University of California, Davis.

Author contributions: R. G. Stewart: Conceptualization, formal analysis, investigation, methodology,

visualization, writing-original draft, writing-reviewing and editing. M. Camacena: investigation. B. A.

Copits investigation, writing-reviewing and editing. J.T. Sack: Conceptualization, formal analysis, funding

acquisition, investigation, methodology, project administration, supervision, visualization, writing-

original draft, writing-reviewing and editing.

37 ABSTRACT

The distinct organization of Kv2 voltage-gated potassium channels on and near the cell body of brain

neurons enables their regulation of action potentials and specialized membrane contact sites.

Somatosensory neurons have a pseudounipolar morphology and transmit action potentials from

peripheral nerve endings through axons that bifurcate to the spinal cord and the cell body within ganglia

including the dorsal root ganglia (DRG). Kv2 channels regulate action potentials in somatosensory

neurons, yet little is known about where Kv2 channels are located. Here we define the cellular and

subcellular localization of the Kv2 paralogs, Kv2.1 and Kv2.2, in DRG somatosensory neurons with a

panel of antibodies, cell markers, and genetically modified mice. We find that relative to spinal cord

neurons, DRG neurons have similar levels of detectable Kv2.1, and higher levels of Kv2.2. In older mice,

detectable Kv2.2 remains similar while detectable Kv2.1 decreases. Both Kv2 subtypes adopt clustered

subcellular patterns that are distinct from central neurons. Most DRG neurons co-express Kv2.1 and

Kv2.2, although neuron subpopulations show preferential expression of Kv2.1 or Kv2.2. We find that Kv2

protein expression and subcellular localization is similar between mouse and human DRG neurons. We

conclude that the organization of both Kv2 channels is consistent with physiological roles in the somata

and stem axons of DRG neurons. The general prevalence of Kv2.2 in DRG as compared to central

neurons and the enrichment of Kv2.2 relative to detectable Kv2.1, in older mice, proprioceptors, and

axons suggest more widespread roles for Kv2.2 in DRG neurons.

Significance statement

The subcellular distribution of Kv2 voltage-gated potassium channels enable compartment-

specific modulation of membrane excitability and organization of membrane contact sites. Here

we identify subcellular distributions of the Kv2 paralogs, Kv2.1 and Kv2.2, in somatosensory

neurons that bear similarities to and distinctions from central neurons. The distribution of Kv2

channels is similar in mouse and human somatosensory neurons. These results identify unique

locations of Kv2 channels in somatosensory neurons that could enable roles in sensory

information processing.

INTRODUCTION

The subcellular localization of voltage-gated ion channels determines how electrical signals are propagated. The two members of the Kv2 family of voltage gated potassium channels, Kv2.1 and Kv2.2, are important for modulating electrical signals in mammalian somatosensory neurons (Bocksteins et al., 2009; Lee et al., 2020; Sun et al., 2022; Tsantoulas et al., 2014; Zheng et al., 2019), yet little is known about where Kv2 channels are localized in these neurons. In central neurons, Kv2 channels are sequestered to specific subcellular regions and identification of where these channels are has helped elucidate their functional roles (Bishop et al., 2015; Du, Tao-Cheng, Zerfas, & McBain, 1998; Irie, 2021; Jensen et al., 2017; Johnson et al., 2018; Kihira, Hermanstyne, & Misonou, 2010; Kirmiz, Vierra, Palacio, & Trimmer, 2018; Misonou, Mohapatra, Menegola, & Trimmer, 2005; Muennich & Fyffe, 2004; Romer et al., 2014; Scannevin, Murakoshi, Rhodes, & Trimmer, 1996; Trimmer, 1991; Vierra, O'Dwyer, Matsumoto, Santana, & Trimmer, 2021). Establishing where Kv2 channels are in somatosensory neurons likewise identifies sites where they could function.

Among vertebrates, the Kv2 family contains two conserved paralogs, Kv2.1 and Kv2.2, which can assemble into homo- or hetero-tetramers to form voltage gated K^+ channels (Blaine & Ribera, 1998; Kihira et al., 2010). In central neurons, Kv2.1 and Kv2.2 channels have unique cellular expression, subcellular localization, and show distinct physiological roles (Bishop et al., 2015; Newkirk et al., 2022). Among cortical neuron types, Kv2.1 is more broadly distributed than Kv2.2 (Bishop et al., 2015). Little is known of how Kv2.1 or Kv2.2 channels are distributed among somatosensory neuron subtypes. In central neurons, Kv2.1 and Kv2.2 channels localize in clusters on the cell soma, proximal dendrites, and axon initial segment (Bishop et al., 2015; King, Manning, & Trimmer, 2014; Sarmiere, Weigle, & Tamkun, 2008; Scannevin et al., 1996; Trimmer, 1991). Clustered Kv2 channels form endoplasmic reticulum-

plasma membrane junctions (Johnson et al., 2018; Kirmiz et al., 2018), where they perform

nonconducting functions which include mediating coupling of L-type calcium channels to ryanodine

receptors (Vierra et al., 2021), and Ca^{2+} uptake (Panzera et al., 2022). In somatosensory neurons it

remains unknown whether Kv2 channels form organized structures or are localized to specific

subcellular compartments. Identifying if channel expression is specific to neuron subtypes and within

specific subcellular domains could indicate whether Kv2.1 and Kv2.2 channels also have distinct

physiological roles in somatosensory neurons.

Kv2 channels are expressed and play important functional roles in DRG. Kv2 transcripts have been

detected in all classes of DRG neurons with a notable absence of Kv2.1 mRNA in proprioceptors (Usoskin

et al., 2015; Wangzhou et al., 2020; Zheng et al., 2019). Presence of Kv2.1 protein in DRG of mice and

rats has been identified by western blot and immunohistochemistry respectively (Sun et al., 2022;

Tsantoulas et al., 2012). Pharmacological inhibition of Kv2 channels enhances rat DRG neuron

responsiveness to sustained inputs (Tsantoulas et al., 2014). Kv2 conductances have been identified in

several mouse DRG neuron subtypes (Bocksteins et al., 2009; Regnier, Bocksteins, Van de Vijver,

Snyders, & van Bogaert, 2016; Zheng et al., 2019). Kv2 channels have also been implicated in

nociception. Kv2.1 and Kv2.2 mRNA transcript levels are decreased after peripheral axotomy of rat DRG

neurons (Tsantoulas et al., 2014). Suppression of Kv2.1 transcription by the epigenetic factor Cdyl is

implicated in regulating pain sensation (Sun et al., 2022). A missing piece of the puzzle of how Kv2

channels are involved in somatosensory neuron function is the location of the channels themselves.

Here we use antibodies against Kv2.1 and Kv2.2 to define their cellular expression and subcellular

localization in DRG neurons of both mice and humans.

MATERIALS AND METHODS

Mice

This study was approved by the UC Davis Institutional Animal Care and Use Committee and conforms to

guidelines established by the NIH. Mice were maintained on a 12 h light/dark cycle, and food and water

was provided *ad libitum.* Kv2.1 KO and Kv2.2 KO mice were from breeding Kv2.1+/- or Kv2.2+/-

heterozygous mice such that WT mice were from breeding pairs that generated KO mice. Kv2.1 and Kv2.2

DKO mice were generated from breeding Kv2.1+/- heterozygous and Kv2.2-/- homozygous mice. WT

controls for Kv2.1/Kv2.2 DKO experiments were C57BL/6J mice purchased from Jackson Laboratory (stock

#000664). The PV^{Ai14} mouse line was a generous gift from Dr. Theanne Griffith (University of California

Davis, Davis CA) and were a cross of $Rosa26^{Ai14}$ (stock #007914, MGI: J:155793) and *PVcre* (stock #008069,

MGI:J: 100886) mice. The $MrgprD^{GFP}$ mouse line was a generous gift from Dr. David Ginty (Harvard

University, Boston MA) (MGI: 3521853). Detailed information about the genotype, age, sex and level of

spinal column of mice used in each figure can be found in Table 1.

Human Tissue Collection

Human DRG were obtained in collaboration with Mid-America Transplant (St. Louis, MO) from three

donors. Donor #1 was a 59-year-old Caucasian male and the DRG was from the 2nd lumbar region (cause

of death: anoxia/stroke). Donor #2 was a 59-year-old black male and the DRG was from the 3rd lumbar

region (cause of death: hemorrhagic stroke). Donor #3 was a 58-year-old Caucasian female and the DRG

was from the 3rd lumbar region (cause of death: cerebrovascular/stroke). DRG were extracted less than

2 hours after aortic cross clamp and transported to the lab where they were dissected to remove the

dura, embedded in Optimal Cutting Temperature (OCT) compound, snap frozen, and stored at -80 °C

until use (Valtcheva et al., 2016). Human DRG were obtained from organ donors with full legal consent

for use of tissue for research in compliance with procedures approved by Mid-America Transplant.

Tissue Preparation

Mice were briefly anesthetized with 3-5% isoflurane and then decapitated. The spinal column was

dissected, and excess muscle tissue removed. The spinal column was then bisected in the middle of the

L1 vertebrae identified by the 13[th] rib and drop fixed for 1 hour in ice cold PFA: 4% formaldehyde

prepared fresh from paraformaldehyde in 0.1 M phosphate buffer (PB) pH adjusted to 7.4 with NaOH.

0.1 M PB buffer was diluted from a 0.4 M PB stock solution that was made by diluting 91.37 g of

$Na_2H_2PO_4$ and 20.98 g of $Na_2H_2PO_4 \bullet H_2O$ in 2 liters of milliQ water. The relatively short fixation period

was chosen because a longer fixation period increased the off target secondary antibody

immunofluorescence from mouse isotype secondary antibodies (Supplemental Figure 1). The spine was

washed 3× for 10 min each wash in PB and cryoprotected at 4 °C in 30% sucrose in PB for 24 hours. The

spine was cut into sections containing two vertebra per sample which were frozen in OCT (Fisher

cat#4585) and stored at -80 °C until sectioning. Vertebrae position relative to the 13[th] rib was recorded

for each frozen sample to determine the specific vertebrae position in the spinal cord. Samples were cut

into 30 μm sections on a freezing stage sliding microtome and were collected on Colorfrost Plus

microscope slides (Fisher cat#12-550-19). Slides were stored at -20 °C or immediately used for multiplex

immunofluorescence labeling.

Human DRG were cut into 30 μm sections on a freezing stage sliding microtome and were collected on

Colorfrost Plus microscope slides. Sections were briefly thawed to adhere to the slide but were

immediately returned to the cryostat kept at -20 °C. Slides were removed from the cryostat and

immediately transferred to freshly made 4% PFA pH 7.4 for 10 minutes. Slides were then placed in PB

solution for 10 minutes and immediately used for multiplex immunofluorescence labeling. Attempts at

immunolabeling Kv2 channels in human DRG tissue that was fixed in 4% PFA for 24 hours were not

successful.

Neuron Cell Culture

Cervical, thoracic and lumbar DRGs were harvested from mice and transferred to Hank's buffered saline

solution (HBSS) (Invitrogen). Ganglia were treated with collagenase (2 mg/ml; Type P, Sigma-Aldrich) in

HBSS for 15 min at 37°C followed by 0.05% Trypsin-EDTA (Gibco) for 2.5 min with gentle rotation. Trypsin

was neutralized with culture media (MEM, with L-glutamine, Phenol Red, without sodium pyruvate)

supplemented with 10% horse serum (heat-inactivated; Gibco), 10 U/ml penicillin, 10 µg/ml streptomycin,

MEM vitamin solution (Gibco), and B-27 supplement (Gibco). Serum-containing media was decanted and

neurons were triturated using a fire-polished Pasteur pipette in MEM culture media containing the

supplements listed above. Neurons were plated on laminin-treated (0.05 mg/ml, Sigma-Aldrich) 35 mm

glass bottom dishes (MatTek cat#P35G-1.5-7-C). Neurons were then incubated at 37°C in 5% CO_2. Neurons

were used for imaging experiments within 24 hours after plating.

Labeling Endogenous Kv2 in live cultured DRG neurons

Plates were removed from 37°C incubator 24 hours after plating. Neurons were washed once with 1 mL

neuronal external (NE) solution (3.5 KCl, 155 NaCl, 10 HEPES, 1.5 CaCl2, and 1 MgCl2, 10 mM glucose

adjusted to pH 7.4 with NaOH). Neurons were then incubated with NE solution supplemented with 1%

BSA for 5 minutes. Neurons were then washed once with NE solution and incubated in 5 µg/mL wheat

germ agglutinin conjugated to Alexa Fluor 405 (Biotium cat#29028-1) diluted in NE solution for 2.5

minutes. Neurons were washed twice with NE solution and imaged. NE solution was removed and then

neurons were incubated for 10 minutes in 100 nM GxTX-594 prepared as described (Thapa et al., 2021).

Neurons were then washed three times in NE solution and imaged again. The temperature of the

imaging chamber was 25 °C throughout the experiment. Experimenter was blinded to which culture

dishes contained DRG neurons from either WT or Kv2.1/Kv2.2 double knockout mice throughout the

experiment and image analysis.

Immunohistochemistry

A hydrophobic barrier was drawn around tissue sections mounted on slides as described above using a

hydrophobic barrier pen (Scientific Device cat#9804-02). Sections were incubated in 4% milk, 0.1 M

phosphate buffer (PB) and 0.2% Triton X-100 (vehicle) for 1 hour. Sections were then incubated in

vehicle containing 0.1 mg/mL IgG F(ab) polyclonal IgG antibody (Abcam cat#ab6668) for 1 hour. We

determined that a concentration of 0.1 mg/mL maximally blocked off-target secondary antibody labeling

when using isotype-specific anti-mouse secondary antibodies (Supplemental Figure 2). Sections were

washed 3× for 5 min each in vehicle and then incubated in vehicle containing primary antibodies for 1

hour. Primary antibodies and concentrations used are listed in Table 2. Sections were then washed 3×

for 5 min each in vehicle and then incubated in vehicle containing IgG-subclass-specific secondary Abs

for one hour (Thermo Fisher). Sections were then washed 3× for 5 min each in PB and mounted with

Prolong Gold (Thermo Fisher cat#P36930) and Deckglaser cover glass (Fisher Scientific cat#NC1776158).

All incubations and washes were done at room temperature with gentle rocking. Human tissue was

immunolabeled identically to mice with the exception that the step incubating sections in vehicle

containing 0.1 mg/mL IgG F(ab) polyclonal IgG antibody was omitted and sections were incubated in

primary antibodies for 2 hours instead of 1. Primary antibodies used in Figure 2 and Figure 3 were used

at saturating concentrations (Supplemental Figure 4).

Imaging

Images were acquired with an inverted scanning confocal and airy disk imaging system (Zeiss LSM 880

Airyscan, 410900-247-075) run by ZEN black v2.1. Laser lines were 405 nm, 488 nm, 543 nm and 633 nm.

Low-magnification images to image whole mouse DRG were acquired in confocal or airy disk imaging

mode with a 0.8 NA 20x objective (Zeiss 420650-9901) details in figure legends. Images containing both

the DRG and spinal cord were tile scan images acquired in confocal mode with the same 20x objective.

When imaging whole DRG for Figure 2 and Figure 3 the imaging plane was selected using fluorescence

from channels that did not contain anti-Kv2.1 or anti-Kv2.2 immunofluorescence. High-magnification

images were acquired in airy disk imaging mode with a 1.4 NA 63x oil objective (Zeiss 420782-9900-799).

Linear adjustments to contrast and brightness and average fluorescence intensity z-projections were

performed using ImageJ software.

Image Analysis

Images were analyzed using ImageJ software (Schindelin et al., 2012). A summary of automated analysis

for selecting neurons in DRG sections can be found in Supplemental Figure 5. The ImageJ plugin

MorphoLibJ was used for performing watershed segmentation of images (Legland, Arganda-Carreras, &

Andrey, 2016). Automatic generation of ROIs from watershed segmentation did not distinguish the

presence of nuclei and is thus expected to over-count larger diameter neurons. Unless stated otherwise

the region of DRG neurons used to analyze anti-Kv2 immunofluorescence is the outer edge

(Supplemental Figure 5 I). Immunofluorescence is defined as the raw pixel values from confocal images

and was not background subtracted. We noted that in smaller neurons the nucleus takes up a greater

percentage of the total volume and could skew fluorescence measurements if fluorescence in the entire

soma volume is measured. As Kv2 protein is not present in the nucleus and is enriched at the outer edge

(Figure 1) we found that measuring anti-Kv2 immunofluorescence at the outer edge of the neuron

reduces potential error that could be associated with soma diameter. For proteins that were not Kv2

channels we analyzed the immunofluorescence of the entire soma as these proteins did not exhibit the

same enrichment at the outer edge as Kv2 channels (Figure 9 B). Where specified, manual ROIs were

used in analysis otherwise automatically generated ROIs were used. When analyzing

immunofluorescence from human neurons, ROIs were drawn around areas within the neuronal soma

that do not contain apparent lipofuscin (Supplemental Figure 11 C). Fitting of imaging data was

performed using Igor Pro software version 8 (Wavemetrics, Lake Oswego, OR) that employs nonlinear

least square curve fitting via the Levenberg-Marquardt algorithm. Distributions of fluorescence intensity

from DRG neurons were fit with a log normal distribution:

$f(x) = A^{-\left[\frac{\ln\left(\frac{x}{x_0}\right)}{width}\right]^2}$ (Equation 1)

Where A = amplitude, x_0 = mean, and width = $\sqrt{2}$ times standard deviation. Concentration-effect

experiments in Supplemental Figure 4 were fit with the Hill equation:

$f(x) = base + \frac{max-base}{1+\left(\frac{x_{half}}{x}\right)^n}$ (Equation 2)

Code Accessibility

In-house Fiji macros and R script used to process imaging data are available for download at

https://github.com/SackLab/DRG-Image-Processing

Statistics

All statistical tests were performed in Igor Pro software version 8 (Wavemetrics, Lake Oswego, OR).

Independent replicates (n) are individual neurons while biological replicates (N) are individual mice. The

n and N values for each figure are listed in Table 1. Details of statistical tests are in the figure legends.

We did not observe a difference between males and females in detectable Kv2.1 or Kv2.2 protein, but

animal numbers were not sufficient for rigorous statistical comparison.

RESULTS

Kv2.1 and Kv2.2 immunolabeling is enriched at the apparent plasma membrane of DRG neuron somata

To assess the cellular expression and subcellular localization of Kv2.1 and Kv2.2 voltage gated potassium channels in DRG neurons we used a set of anti-Kv2.1 and anti-Kv2.2 antibodies that have previously been validated in brain tissue from Kv2 knockout mice (Bishop et al., 2015) to perform multiplex immunofluorescence labeling of DRG neurons. Anti-Kv2.1 (magenta) and anti-Kv2.2 (green) immunofluorescence is prominent in DRG neuron somata (Figure 1 A). In neurons we observed anti-Kv2.1 and anti-Kv2.2 immunofluorescence enriched at the outer edge of DRG neuron somata consistent with their plasma membrane localization (Figure 1 B arrows). This putative cell surface localization is distinct from prior reports of anti-Kv2 immunofluorescence in DRG (Tsantoulas et al., 2012) as well as that for other ion channel proteins such as Nav1.8 (Shields et al., 2012) and TRPV1 (Cho & Valtschanoff, 2008), all of which are primarily in the cytoplasm. Anti-Kv2.1 or anti-Kv2.2 immunofluorescence at the edge of neuron somata was more apparent when imaging thinner optical sections. To determine if enrichment of Kv2 immunolabeling at the outer edge of DRG neurons was an artifact of our immunohistochemistry or imaging protocols, we additionally labeled DRG sections with a knockout-validated antibody against Nav1.8. In the same DRG section labeled for Kv2 channels, cytoplasmic anti-Nav1.8 immunofluorescence was prominent in small to medium diameter lumbar DRG neurons, consistent with previous reports (He et al., 2010; Shields et al., 2012) (Figure 1 B). To quantify enrichment of anti-Kv2 versus anti-Nav1.8 immunofluorescence at the outer edge of DRG neurons, we manually drew regions of interest (ROIs) for each DRG soma. We generated a 2 μm wide annulus that encompassed the outer edge and a second 2 μm wide annulus just inside the first one (Figure 1 C arrows

1 and 2 respectively). By comparing the fluorescence intensity of immunofluorescence signals within

these annuli we found that the edge of DRG neuron somata is enriched for anti-Kv2.1 and anti-Kv2.2

immunofluorescence relative to that of Nav1.8 (ANOVA < 0.001) (Figure 1 D). Similarly, anti-Kv2.1 and

anti-Kv2.2 immunofluorescence is enriched at the edge of DRG neuron somata relative to anti-TRPV1

immunofluorescence (ANOVA < 0.001) (Supplemental Figure 3).

The majority of dorsal root ganglion neurons express Kv2 protein

To determine whether the anti-Kv2.1 immunofluorescence signal in DRG sections is specific and

dependent on the presence of Kv2.1 protein, we compared fluorescence intensities in DRG samples

prepared in parallel from age and sex matched wild-type (WT) mice and Kv2.1 knock-out (KO) mice. We

observed anti-Kv2.1 immunolabeling in WT DRG neurons that was absent from Kv2.1 KO, while

fluorescence corresponding to an antibody that targets βIII tubulin was similar in both WT and Kv2.1 KO

mice (Figure 2 A). To quantify fluorescence intensities of DRG sections, we manually drew ROIs around

neuron soma profiles with clearly visible nuclei and measured the fluorescence intensity at the outer

edge of profiles as shown in Figure 1 C. We define a profile as a slice of a DRG neuron in a histological

section (Coggeshall, 1992). We found that 99% of manually identified profiles from the WT mouse DRG

shown in Figure 2 A had anti-Kv2.1 immunofluorescence above the mean of Kv2.1 KO mice (Figure 2 B).

To reduce human bias in identification of profiles we used an automated method to generate ROIs. As

with the manual method, the automated method reliably identified neuronal profiles (Supplemental

Figure 5 I). However, the automated method did not distinguish profiles without nuclei which could lead

to overrepresentation of larger neurons (Coggeshall, 1992; Coggeshall & Lekan, 1996). Additionally, the

automated method occasionally selected ROIs which did not appear to be neurons (Supplemental Figure

5 H red arrows). Despite these limitations, the absolute reproducibility of the automated method when

scaled to identify thousands of neuronal profiles indicated that automated ROIs could provide a rigorous

means of identifying neuronal profiles for statistical analysis. In automatically generated ROIs from the

same WT mouse manually analyzed in Figure 2 B we observed a reduction in the mean anti-Kv2.1 but

not anti-βIII immunofluorescence in Kv2.1 KO mice (Figure 2 C and D). We found that 89% of

automatically generated ROIs had anti-Kv2.1 immunofluorescence above the mean of the paired Kv2.1

KO mouse. We manually identified that 9% of the automatically generated ROIs from the WT mouse did

not appear to be neurons, suggesting that 98% of profiles in this automated dataset could have anti-

Kv2.1 immunofluorescence above the mean of the Kv2.1 KO mouse, consistent with manual analysis.

We expanded the automated method to 5 pairs of age and sex matched WT and Kv2.1 KO mice, and

found that anti-Kv2.1 immunofluorescence was significantly reduced in each matched Kv2.1 KO mouse

relative to anti-Kv2.2 or anti-βIII tubulin immunofluorescence while no significant difference was

observed between anti-Kv2.2 and anti-βIII tubulin (Figure 2 E) (ANOVA p < 0.001). We identified that 93

± 6% (SD) of ROIs from the five WT mice had anti-Kv2.1 immunofluorescence above the mean of age and

sex matched Kv2.1 KO mice (Figure 2 F). This analysis confirms that anti-Kv2.1 immunofluorescence

reveals Kv2.1 protein at neuron surfaces.

We developed a method to estimate the fraction of ROIs that contain detectable Kv2.1 protein. The

histogram of fluorescence intensity in automatically generated ROIs from Kv2.1 KO DRG sections labeled

with an anti-Kv2.1 antibody had variability which could reasonably be fit with a log normal distribution

(equation 1) (Figure 2 G). We assumed a similar distribution of background immunofluorescence is also

present in neuron profiles of age and sex matched WT mice and fit the Kv2.1 KO distribution to the WT

histogram to estimate the fraction of ROIs without detectable Kv2.1 (Figure 2 H red gaussian). Using this

method, we compared Kv2.1 immunofluorescence in DRG sections of multiple age and sex matched WT

and Kv2.1 KO mice. The fitting indicated that 84 ± 9% (SD) of automatically generated ROIs from WT

mice have detectable Kv2.1 protein (N = 5 pairs of mice) (Figure 2 I). As approximately 9% of ROIs are

expected to not contain neurons these results suggest that greater than 90% of mouse DRG neurons

have detectable Kv2.1 protein. We further validated our automated methodology with a transgenic

MrgprD-GFP mouse line that expresses GFP in non-peptidergic nociceptors that comprise approximately

19-24% of all lumbar DRG neurons (Dirajlal, Pauers, & Stucky, 2003; Dong, Han, Zylka, Simon, &

Anderson, 2001; Wang & Zylka, 2009). In comparing the MrgprD-GFP mice to WT C57BL/6J mice we

found that 23 ± 15% (SD) of automatically generated profiles (N = 4 pairs of mice) had detectable GFP

(Supplemental Figure 6).

We compared anti-Kv2.2 immunofluorescence from an age and sex matched WT and Kv2.2 KO mouse by

drawing ROIs around neuron soma profiles with clearly visible nuclei. This identified that greater than

99% of WT profiles have anti-Kv2.2 immunofluorescence above the mean immunofluorescence of the

paired Kv2.2 KO mouse (Figure 3 B). Automatically generated ROIs for multiple age and sex matched WT

and Kv2.2 KO mice indicate that 91 ± 5% (SD) of ROIs from WT mice have detectable Kv2.2 protein (N = 8

pairs of mice) (Figure 3 F-H). Using the automated analysis, we found that anti-Kv2.2

immunofluorescence was significantly reduced in Kv2.2 KO mice relative to anti-Kv2.1 or anti-NF200

immunofluorescence in age and sex matched WT and Kv2.2 KO mice (ANOVA $p < 0.001$) while no

significant difference between anti-NF200 or anti-Kv2.1 immunofluorescence was observed (Figure 3 A-

E). These combined results suggest that more than 90% of neuron profiles contain detectable Kv2.1 or

Kv2.2 protein, raising the possibility that all mouse DRG neurons express Kv2 channels.

Kv2.1 immunolabeling decreases in older mice while Kv2.2 immunolabeling remains similar

Kv2.2 transcript in mouse DRG decreases during postnatal development (Regnier et al., 2016) and Kv2.1 protein expression in mouse brain decreases with age (Cotella et al., 2012; Regnier et al., 2016). To identify if Kv2 protein levels in DRG neurons vary between young adult and old mice we compared anti-Kv2.1 and anti-Kv2.2 immunofluorescence from samples subjected to identical immunolabeling and imaging protocols. We observed lower anti-Kv2.1 (Figure 4 A, B and C) in DRG sections from 50 week old mice relative to 7-16 week old mice. Comparisons with Kv2.1 KO mice indicate that 50 ± 23% (SD) of ROIs from 50 week old mice express detectable Kv2.1 protein, significantly less (p = 0.018) than the 84 ± 9% (SD) of ROIs with detectable Kv2.1 protein from 7-16 week old mice (Figure 4 D). In contrast, similar anti-Kv2.2 immunofluorescence was seen in samples from young adult and old mice (Figure 4 E, F and G). Comparisons with Kv2.2 KO mice indicate that 96 ± 3% of profiles from 50 week old mice express detectable Kv2.2 protein, a weakly significant increase (p = 0.03) from the 88 ± 3% (SD) of profiles with detectable Kv2.2 protein from 7-24 week old mice (Figure 4 H). We note that this weak statistical increase is underpowered due to the availability of only three 50 week old Kv2.2 KO mice.

Kv2 channels are expressed on the cell surface of acutely dissociated DRG neurons

To determine if Kv2 channels are present on neuron surfaces we labeled live acutely dissociated DRG neurons with a cell-impermeant fluorescent probe which binds Kv2 channels. The probe is a variant of the tarantula peptide guangxitoxin-1E conjugated to Alex Fluor 594 (GxTX-594) which binds to an extracellular site of Kv2.1 and Kv2.2 channels (Thapa et al., 2021). We applied GxTX-594 to live dissociated DRG neurons from WT mice and double knockout (DKO) mice lacking expression of both Kv2.1 and Kv2.2 protein (Figure 5). In WT DRG neurons, we observed fluorescence at the membrane

after application of 100 nM GxTX-594 that was not present before the addition of GxTX-594 (Figure 5 A).

Wheat germ agglutinin conjugated to Alexa 405 (WGA-405) was used to identify the surface membrane

of cultured DRG neurons and confirmed that the GxTX-594 fluorescence observed in WT neurons was

cell surface localized (Figure 5 A middle). The cell surface localized fluorescence of GxTX-594 observed in

WT neurons was not present in Kv2.1/Kv2.2 DKO neurons (Figure 5 B). To quantify the cell surface-

associated fluorescence signals of individual WT and Kv2.1/Kv2.2 DKO neurons, we used an automated

method to generate ROIs corresponding to the WGA-405 fluorescence and measured GxTX-594 labeling

within this ROI (Figure 5 C). This analysis confirmed a reduction in GxTX-594 fluorescence at the

membrane in cultured DRG neurons from Kv2.1/Kv2.2 DKO mice compared to WT mice (Figure 5 D).

Similar to the Kv2 immunofluorescence signals from KO mice (Figures 2 and 3), GxTX-594 fluorescence

from Kv2.1 /Kv2.2 DKO mice could be reasonably fit with a log normal distribution (equation 1) (Figure 5

E). Applying the same method used to estimate the fraction of neuronal profiles expressing detectable

Kv2.1 and Kv2.2 immunofluorescence we compared cultured DRG neurons of age and sex matched WT

and Kv2.1/Kv2.2 DKO mice and estimated that 76 ± 2.4% (SD) of DRG neurons have detectable cell

surface GxTX-594 fluorescence (Figure 5 F and G). This indicates that at least 76% of neurons express

Kv2 channels in their surface membrane, consistent with a report of Kv2 conductance in every DRG

neuron type assessed (Zheng et al., 2019) .

Anti-Kv2.2 immunofluorescence is enriched in DRG neurons relative to spinal cord neurons

In mammals, anti-Kv2.1 and anti-Kv2.2 immunofluorescence is apparent in central neurons of the brain

(Kihira et al., 2010; Bishop et al., 2015) and anti-Kv2.1 in the spinal cord (Bishop et al., 2015; Muennich &

Fyffe, 2004). We measured Kv2 immunofluorescence intensities in sections of mouse spinal column

which contain both the DRG and the spinal cord. Neurons in the ventral and dorsal horn of the spinal

cord exhibit anti-Kv2.1 immunofluorescence (Figure 6 A). We observed anti-Kv2.1 puncta on neurons in

the ventral horn (Figure 6 A inset) similar to that described in alpha motor neurons (Fletcher et al., 2017;

Romer et al., 2014). Fewer spinal cord neurons exhibit anti-Kv2.2 immunofluorescence than anti-Kv2.1

(Figure 6 A and Supplemental Figure 7 A arrow heads). Kv2.2 channel protein in the spinal cord of mice

has to our knowledge not been assessed, but anti-Kv2.2 immunofluorescence has been observed early

in development in the ventrolateral spinal cord in the frog *Xenopus laevis* (Gravagna, Knoeckel, Taylor,

Hultgren, & Ribera, 2008). We analyzed anti-Kv2 immunofluorescence from individual neurons in the

DRG and ventral horn of five mice and found that while anti-Kv2.1 immunofluorescence is similar

between the two regions (Figure 6 B), anti-Kv2.2 immunofluorescence is significantly higher in DRG

neurons relative to ventral horn neurons (Figure 6 C). Approximately 30% of neurons in the DRG have

anti-Kv2.2 immunofluorescence at a level higher than the brightest anti-Kv2.2 immunofluorescence

observed in the spinal cord. We repeated this analysis with a different anti-Kv2.2 antibody and found

similar results (Supplemental Figure 7).

Kv2 channels on DRG neuron somata and stem axons form clustered subcellular patterns with

similarities to and distinctions from central neurons.

In neurons throughout the brain (Trimmer, 1991, Scannevin et al., 1996, Bishop et al., 2015) and motor

neurons in the spinal cord (Muennich and Fyffe, 2004), Kv2 channels form punctate structures referred

to as clusters. In central neurons, Kv2 clusters localize to the soma, proximal dendrites and axon initial

segment (Lim, Antonucci, Scannevin, & Trimmer, 2000; Scannevin et al., 1996; Trimmer, 1991). At

clusters, Kv2 channels organize protein signaling complexes and endoplasmic reticulum-plasma

membrane junctions (Johnson et al., 2018; Kirmiz et al., 2018; Panzera et al., 2022; Vierra et al., 2021).

To determine whether Kv2 channels in DRG neurons form clusters similar to central neurons, we

compared *en face* z-projections of Kv2 clusters in DRG neurons and ventral horn neurons in the same

spinal cord section (Figure 7). Anti-Kv2.1 immunofluorescence in the ventral horn neurons of these mice

resembles previous reports of mice (Fletcher et al., 2017) and rats (Fletcher et al., 2017; Muennich &

Fyffe, 2004). In both DRG and spinal cord neurons, Kv2.1 and Kv2.2 channels form dense clusters (Figure

7 A and B). Kv2 channel clusters in DRG typically appeared smaller than Kv2 clusters in spinal cord.

However, we are not certain these observations in fixed tissue represent the in vivo arrangement, as

Kv2.1 channels in brain neurons can decluster in the period between sacrifice and fixation (Misonou et

al., 2005). In some DRG neurons, we observed donut shaped Kv2.1 and Kv2.2 clusters (Figure 7 B and C

insets) which resemble donut shaped Kv2 clusters in interneurons which harbor specialized protein

machinery in their center (Vullhorst et al., 2015).

DRG neurons are distinct from central neurons as they have a single stem axon that exits the cell soma

and bifurcates into the central and peripheral axon branches (Ha, 1970). We observed punctate

immunofluorescence for both Kv2.1 and Kv2.2 on the apparent stem axons of DRG neurons (Figure 7 B,

D and E arrow heads, Supplemental Figure 8). In DRG sections from the *MrgprD*[GFP] mouse line we

observed Kv2 clusters on the stem axon of GFP-expressing neurons. As *MrgprD*[GFP] marks non-peptidergic

nociceptors (Zheng et al., 2019) we confirm stem axon localization in this subpopulation of

unmyelinated neurons (Supplemental Figure 8 A).

While Kv2 clusters in central neurons and many DRG neurons are distributed evenly throughout the

soma, we found that DRG neurons asymmetrically distribute Kv2 clusters on the soma (Figure 7 B-E

arrows). To test if enrichment of Kv2 channel clusters is oriented to a specific region on the soma we

manually traced the surface of neurons whose apparent stem axons were visible (Figure 7 F). The

brightest anti-Kv2.1 and anti-Kv2.2 immunofluorescence is near the stem axon in 76 and 80% of DRG

neurons respectively (Figure 7 G and H). Asymmetric distribution of Kv2 channels is not frequently

observed in central neurons of mice (Bishop et al., 2018; Bishop et al., 2015; Romer et al., 2014)

suggesting a potentially unique role of Kv2 subcellular organization in DRG neurons.

Kv2.2 channels are expressed in peripheral axons of DRG neurons

To determine if anti-Kv2 immunofluorescence is detectible in DRG axons in regions beyond the stem

axon we immunolabeled samples containing both peripheral and central axons, using NF200 or βIII

tubulin as markers for DRG axons which target myelinated neurons or all DRG neurons respectively.

Neurons were not co-labeled with βIII tubulin and Kv2.2 targeting antibodies because they were of the

same isotype. We did not observe anti-Kv2.1 immunofluorescence in peripheral axons (Supplemental

Figure 9) from the same DRG section that had detectable anti-Kv2.1 immunofluorescence in neuron

somas (Figure 2). However, anti-Kv2.2 immunofluorescence is detectable in peripheral and central axons

of WT but not Kv2.2 KO mice (Figure 8 B and C). Anti-Kv2.2 immunofluorescence on myelinated

peripheral axons appears as discrete puncta or as larger bands which span the width of the axon (Figure

8 B arrows). We also observed anti-Kv2.2 immunofluorescence not colocalized with NF200, suggestive of

unmyelinated fibers (Figure 8 B arrow head). In *MrgprD^{GFP}* mice, anti-Kv2.2 immunofluorescence

colocalizes with axons expressing cytosolic GFP, identifying that non-peptidergic nociceptors also

express Kv2.2 protein along the peripheral axon (Figure 8 D). We further investigated the expression of

Kv2.2 in peripheral myelinated fibers by co-labeling peripheral axons with antibodies specific for CASPR

(Einheber et al., 1997) and Kv1.2 (Rasband et al., 1998) which identify the paranodal and juxtaparanodal

regions of nodes of Ranvier, respectively in myelinated fibers. We found punctate anti-Kv2.2

immunofluorescence associated with both CASPR and Kv1.2 in some WT DRG neuron axons that is not

present in Kv2.2 KO mice (Supplemental Figure 10 A). We analyzed anti-Kv2.2 immunofluorescence in

WT and Kv2.2 KO mice by manually drawing ROIs around CASPR and Kv1.2 immunofluorescence while

blinded to anti-Kv2.2 immunofluorescence and found that 26/34 CASPR and 20/31 Kv1.2

immunolabeled regions had anti-Kv2.2 immunofluorescence brighter than the brightest anti-Kv2.2

immunofluorescence measured in Kv2.2 KO mice (Supplemental Figure 10 B). These results indicate that

Kv2.2 but not Kv2.1 protein is detectable in axons of myelinated and unmyelinated somatosensory

neurons. However, these results do not identify if Kv2.2 channels are localized to specific subcellular

regions of somatosensory axons.

Anti-Kv2.1 and anti-Kv2.2 immunofluorescence is non-uniform across DRG neuron subtypes

Different voltage gated potassium channels make distinct contributions to outward potassium currents

in sensory neuron subtypes (Zheng et al., 2019). We observed subpopulations of DRG neurons that have

bright anti-Kv2.1 immunofluorescence but dim anti-Kv2.2 immunofluorescence and vice-versa (Figure 9

A and C, Figure 1 A, and Figure 6 A). We hypothesized that differences in Kv2.1 and Kv2.2 protein density

could potentially denote specific subtypes of sensory neurons. In order to test if relative density of Kv2.1

and Kv2.2 protein differs in subtypes of DRG neurons, we assessed anti-Kv2.1 and anti-Kv2.2

immunofluorescence in four DRG neuron subtypes. Neurons identified by genetically encoded markers

were GFP-expressing neurons from *MrgprD^{GFP}* mice which we refer to as non-peptidergic nociceptors

(Zheng et al., 2019), or tdTomato-expressing neurons from *PV^{Ai14}* mice which we refer to as

proprioceptors (Zheng et al., 2019). Neurons immunolabeled for CGRP we refer to as peptidergic

nociceptors (Zheng et al., 2019) and neurons immunolabeled for NF200 we refer to as myelinated DRG

neurons (Usoskin et al., 2015) (Figure 9 B and D). We quantified detectable Kv2.1 or Kv2.2 protein

density in individual ROIs by rank ordering the immunofluorescence intensity. Rank orders from multiple

mice were pooled with a value of 100% representing the brightest ROI in each mouse (Figure 9 D). The

rank percentiles of Kv2.1 and Kv2.2 have similar distributions in peptidergic and non-peptidergic

nociceptors. Many ROIs with the highest Kv2.1 or Kv2.2 rank percentile were NF200-positive myelinated

neurons. Proprioceptors, a subset of the myelinated neurons, appear to account for the subpopulation

of neurons with bright anti-Kv2.2 and dim anti-Kv2.1, and have high rank percentiles of Kv2.2 but low

rank percentiles of Kv2.1. (Figure 9 A, B, and C). Profiles with bright anti-Kv2.1 and dim anti-Kv2.2

immunofluorescence are mostly NF200-positive myelinated neurons of unknown subtype (Figure 9 A, B,

and C blue points). Overall, these results identify that the relative densities of Kv2.1 and Kv2.2 protein

are similar in DRG neuron subtypes except proprioceptors. These results are consistent with reports that

Kv2.1 and Kv2.2 mRNA transcript levels are similar in DRG neuron subtypes except proprioceptors. Our

finding that Kv2.1 protein density is anomalously low in proprioceptors mirrors the finding that Kv2.1

transcripts are also low in proprioceptors (Usoskin et al., 2015; Zheng et al., 2019).

Kv2 channel cellular expression and subcellular localization in human DRG neurons is similar to mice

Kv2 mRNA transcripts are present in human DRG neurons (Tavares-Ferreira et al., 2022; Wangzhou et

al., 2020), and we assessed whether anti-Kv2 immunofluorescence is associated with human DRG

neurons. As human Kv2.1 and Kv2.2 knockout controls are not available, we compared fluorescence in

DRG sections with anti-Kv2 primary antibodies to those lacking primary antibodies. Control samples

lacking primary antibodies revealed strong autofluorescence at all excitation wavelengths (Supplemental

Figure 11 A and B) consistent with other studies of human DRG neurons which have attributed this

autofluorescence to lipofuscin (Shiers et al., 2021). We found that most neurons have anti-Kv2.1 and

anti-Kv2.2 immunofluorescence brighter than controls lacking primary antibodies (Supplemental Figure

11 D and E) suggesting that Kv2 protein is present in human DRG neurons. As a control, we used the

same methods with anti-Nav1.7 and anti-Nav1.8 antibodies and found anti-Nav1.7 immunofluorescence

in nearly all human DRG neurons while anti-Nav1.8 was in a smaller fraction of DRG neurons, consistent

with previous reports of Nav1.7 protein and Nav1.8 transcript in human DRG (Shiers, Klein, & Price,

2020) (Supplemental Figure 11 A arrow heads, B, F and G). Similar to DRG neurons in mice, we observed

subpopulations of neurons that had bright anti-Kv2.1 immunofluorescence but dim anti-Kv2.2

immunofluorescence and conversely neurons that had bright anti-Kv2.2 immunofluorescence but dim

anti-Kv2.1 immunofluorescence (Figure 10 A and B). These observations were consistent in DRG from 3

human donors (Supplemental Figure 11 A and H and Supplemental Figure 12). We were unable to

identify if the population of human DRG neurons with low anti-Kv2.1 and high anti-Kv2.2 are

proprioceptors as we observed in mice; attempts to label parvalbumin in human neurons were

unsuccessful.

We assessed whether Kv2 protein subcellular localization in humans is similar to mice. We identified

neurons in all three human DRG samples where anti-Kv2.1 and anti-Kv2.2 immunofluorescence was

enriched at the outer edge of DRG neuron somata (Figure 10 A arrow and Supplemental Figure 13). In

some neurons from all three humans, Kv2 clusters exhibited asymmetric distribution on the neuronal

soma (Figure 10 B green arrow head, Supplemental Figure 13 A-E arrows). In one human we identified a

neuron with Kv2.2 immunofluorescence that was asymmetrically distributed near the stem axon

consistent with observations in mice (Figure 10 B green arrow). We identified Kv2.2

immunofluorescence on the stem axon of DRG neurons in all three humans using anti-NF200 or anti-

Nav1.7 as markers of the stem axon (Figure 10 B-C and Supplemental Figure 14 arrows). In two out of

three humans we also identified Kv2.1 immunofluorescence on the stem axon of human DRG neurons

(Figure 10 C inset and Supplemental Figure 14 B and D inset). To determine if Kv2.2 channels are present

in human DRG neuron axons, we labeled human DRG sections with the same anti-Kv2.2, anti-NF200,

anti-CASPR and anti-Kv1.2 antibodies that were used in mice. We found anti-Kv2.2 immunofluorescence

associated with anti-NF200, anti-CASPR and anti-Kv1.2 immunofluorescence in human DRG neuron

axons (Supplemental Figure 15) suggesting that Kv2.2 channels could contribute to electrical

propagation or other physiological functions in axons of human DRG neurons.

DISSCUSSION

Our results reveal overlapping yet distinct localization patterns of Kv2.1 and Kv2.2 in DRG neurons. Here,

we discuss the unique features of Kv2 localization in the DRG and how these could influence neuronal

physiology and somatosensation.

We identify differences between anti-Kv2.1 and anti-Kv2.2 immunofluorescence in DRG neurons. Kv2.2

mRNA is abundant in all DRG neuron subtypes identified, while Kv2.1 mRNA is low in proprioceptors and

abundant in other subtypes (Usoskin et al., 2015; Zheng et al., 2019). Consistent with these findings we

conservatively estimate that Kv2.1 and Kv2.2 protein is in at least 90% of mouse DRG neurons, though

Kv2.1 immunofluorescence is anomalously low in proprioceptors. Detection of Kv2.2 but not Kv2.1 in

DRG neuron peripheral and central axons suggests that Kv2.2 channels may be important for electrical

transmission in these axons. While detectable Kv2.2 remained constant in DRG neurons of older mice,

detectable Kv2.1 decreased, possibly contributing to the altered repolarization noted in DRG neurons

from aged mice (Scott, Leu, & Cinader, 1988). The differences between Kv2.1 and Kv2.2 in

somatosensory neurons are consistent with distinct roles for these Kv2 paralogs in somatosensory

neuron physiology, as seen in brain and pancreatic islets (Bishop et al., 2015; Johnston et al., 2008; Li et

al., 2013).

The expression and subcellular localization of Kv2 channels in DRG neurons are distinct from central

neurons. Kv2.1 is present in most brain neurons (Bishop et al., 2018; Bishop et al., 2015; Hwang, Fotuhi,

Bredt, Cunningham, & Snyder, 1993) while Kv2.2 is less broadly expressed (Bishop et al., 2015;

Hermanstyne et al., 2010; Hwang, Glatt, Bredt, Yellen, & Snyder, 1992; Johnston et al., 2008; Kihira et

al., 2010). In contrast to this, we find Kv2.2 expressed in at least 90% of mouse DRG neurons. We also

identify that Kv2.2 is enriched in the DRG relative to neurons in the spinal cord, suggesting that

abundant Kv2.2 protein expression is a distinct feature of DRG neuron physiology. In central neurons,

Kv2 channel clusters are typically evenly distributed across neuron somata (Bishop et al., 2015; Kihira et

al., 2010; Muennich & Fyffe, 2004) and asymmetric subcellular localization of Kv2 clusters on neuron

somata has to our knowledge not been described. We find that Kv2 channels are enriched at the plasma

membrane in a subcellular region near the stem axon of some DRG neurons. Differences in Kv2

localization between central and somatosensory neurons could arise from differences in the

extracellular environment surrounding DRG neuron somata. In central neurons, Kv2 channel clusters are

reported at cholinergic C-terminal synaptic sites, S-type synapses, and apposed to astrocytic end feet

(Du et al., 1998; Muennich & Fyffe, 2004). However, synapses have not been reported in the DRG and

DRG neurons are surrounded by satellite glia instead of astrocytes (Matsuda et al., 2005). Further

studies could investigate if Kv2 clusters are localized to extracellular structures in the DRG. Kv2.2 has

been identified in juxtaparanodes but not paranodes of axons in the osseous spiral lamina of the cochlea

(Kim & Rutherford, 2016) and previous work has also identified that Kv2.1 channels are necessary for

enabling ER Ca2+ uptake during electrical activity in both the soma and axon (Panzera et al., 2022). To

our knowledge, Kv2.2 immunofluorescence has not been previously reported in distal axons in the brain,

spinal cord or DRG. Our results indicate that Kv2.2 channels are present in myelinated and unmyelinated

axons of DRG neurons. The mechanism of Kv2.2 protein trafficking to the axon is unknown. Kv2.1

channels are trafficked to the axon initial segment through a distinct secretory pathway (Jensen et al.,

2017) allowing neurons to regulate precise compartment specific localization of Kv2.1 protein. A similar

mechanism for Kv2.2 localization in DRG axons could allow neurons to independently control Kv2.2

protein expression in the neuron soma and axon. The subcellular distribution of Kv2.1 and Kv2.2

subunits in DRG neurons suggests they could each distinctly modulate electrical signals. We note that

our observations could not distinguish whether immunofluorescence in DRG axons could represent

intracellular trafficking vesicles of Kv2.2 proteins, or whether axonal Kv2.2 is on the membrane surface

where it could modulate electrical signals.

Kv2 channels in DRG neurons may play important roles in somatosensation and nociception. Kv2

conductances can modulate action potential repolarization, afterhyperpolarization and repetitive firing

(Liu & Bean, 2014; Zheng et al., 2019). Kv2 conductances modulate electrical transmission in rat DRG

neurons (Tsantoulas et al., 2014). Knockdown of Kv2.1 increases hypersensitivity to painful stimuli and

regulation of Kv2.1 protein expression through the epigenetic factor Cdyl modulates neuronal

excitability and nociception (Sun et al., 2022). Consistent with these studies, our results show that Kv2

channels are localized to DRG neuron axons, where they could potentially influence electrical

transmission.

Regulatory pore-forming subunits of the Kv5, Kv6, Kv8, and Kv9 "silent subunit" families obligately

assemble into heteromeric channels with Kv2 subunits (Bocksteins, 2016) and have been implicated in

nociception. Women with decreased sensitivity to labor pain were identified to have a mutation in the

Kv6.4 silent subunit that disrupts Kv2.1/Kv6.4 heteromer expression and is proposed to modulate

nociceptor signaling (Lee et al., 2020). Downregulation of the Kv9.1 silent subunit after axotomy evokes

hyperexcitability of DRG neurons (Tsantoulas et al., 2012). While both Kv2 channels are broadly

expressed in DRG neurons, transcript levels of silent subunits in mice and humans show much more

neuron subtype specificity (Tavares-Ferreira et al., 2022; Zheng et al., 2019), raising the possibility of

molecularly distinct Kv2-containing channels in somatosensory neuron subtypes.

Previous studies have identified that voltage gated ion channel expression can be substantially different

between mice and human DRG, notably, in the expression of Nav1.7 (Shiers et al., 2020). However, we

find expression of Kv2 channels to be similar in mouse and human DRG. An overall similarity is

supported by the presence of detectable Kv2 channels in most neurons, consistent with transcriptomics

data from both mice and humans (Tavares-Ferreira et al., 2022; Zheng et al., 2019). The subcellular

localization of Kv2 channels also appears similar between mice and humans. These similarities include

enrichment of Kv2 channels at the outer edge of neurons, Kv2 clusters on the neuron soma as well as

presence of Kv2.1 and Kv2.2 on DRG neuron axons. These similarities suggest that the functional roles of

Kv2 channels could be similar in both mouse and human DRG neurons.

Overall, we find attributes of Kv2 channel localization in somatosensory neurons that could point to

their functional roles in sensory information processing. The broad expression of Kv2.2 among DRG

neuron types and expansive axonal localization of Kv2.2 suggest a key role for this channel in the

somatosensory nervous system. The specialized divergence of Kv2.1 and Kv2.2 suggest the functional

roles of Kv2 subtypes in DRG neurons are overlapping yet distinct.

REFERENCES

Bishop, H. I., Cobb, M. M., Kirmiz, M., Parajuli, L. K., Mandikian, D., Philp, A. M., . . . Trimmer, J.
S. (2018). Kv2 ion channels determine the expression and localization of the associated
AMIGO-1 cell adhesion molecule in adult brain neurons. *Front Mol Neurosci, 11*, 1.
doi:10.3389/fnmol.2018.00001
Bishop, H. I., Guan, D., Bocksteins, E., Parajuli, L. K., Murray, K. D., Cobb, M. M., . . . Trimmer, J.
S. (2015). Distinct cell- and layer-specific expression patterns and independent
regulation of Kv2 channel subtypes in cortical pyramidal neurons. *J Neurosci, 35*(44),
14922-14942. doi:10.1523/jneurosci.1897-15.2015
Blaine, J. T., & Ribera, A. B. (1998). Heteromultimeric potassium channels formed by members
of the Kv2 subfamily. *J Neurosci, 18*(23), 9585-9593. doi:10.1523/jneurosci.18-23-
09585.1998
Bocksteins, E. (2016). Kv5, Kv6, Kv8, and Kv9 subunits: No simple silent bystanders. *J Gen
Physiol, 147*(2), 105-125. doi:10.1085/jgp.201511507
Bocksteins, E., Raes, A. L., Van de Vijver, G., Bruyns, T., Van Bogaert, P. P., & Snyders, D. J.
(2009). Kv2.1 and silent Kv subunits underlie the delayed rectifier K+ current in cultured
small mouse DRG neurons. *Am J Physiol Cell Physiol, 296*(6), C1271-1278.
doi:10.1152/ajpcell.00088.2009
Cho, W. G., & Valtschanoff, J. G. (2008). Vanilloid receptor TRPV1-positive sensory afferents in
the mouse ankle and knee joints. *Brain Res., 1219*, 59-65.
doi:https://doi.org/10.1016/j.brainres.2008.04.043
Coggeshall, R. E. (1992). A consideration of neural counting methods. *Trends Neurosci, 15*(1), 9-
13. doi:10.1016/0166-2236(92)90339-a
Coggeshall, R. E., & Lekan, H. A. (1996). Methods for determining numbers of cells and
synapses: a case for more uniform standards of review. *J Comp Neurol, 364*(1), 6-15.
doi:10.1002/(SICI)1096-9861(19960101)364:1<6::AID-CNE2>3.0.CO;2-9
Cotella, D., Hernandez-Enriquez, B., Wu, X., Li, R., Pan, Z., Leveille, J., . . . Sesti, F. (2012). Toxic
role of K+ channel oxidation in mammalian brain. *J Neurosci, 32*(12), 4133-4144.
doi:10.1523/jneurosci.6153-11.2012
Dirajlal, S., Pauers, L. E., & Stucky, C. L. (2003). Differential response properties of IB(4)-positive
and -negative unmyelinated sensory neurons to protons and capsaicin. *J Neurophysiol,
89*(1), 513-524. doi:10.1152/jn.00371.2002
Dong, X., Han, S., Zylka, M. J., Simon, M. I., & Anderson, D. J. (2001). A diverse family of GPCRs
expressed in specific subsets of nociceptive sensory neurons. *Cell, 106*(5), 619-632.
doi:10.1016/s0092-8674(01)00483-4
Du, J., Tao-Cheng, J. H., Zerfas, P., & McBain, C. J. (1998). The K+ channel, Kv2.1, is apposed to
astrocytic processes and is associated with inhibitory postsynaptic membranes in
hippocampal and cortical principal neurons and inhibitory interneurons. *Neuroscience,
84*(1), 37-48. doi:https://doi.org/10.1016/S0306-4522(97)00519-8
Einheber, S., Zanazzi, G., Ching, W., Scherer, S., Milner, T. A., Peles, E., & Salzer, J. L. (1997). The
axonal membrane protein Caspr, a homologue of neurexin IV, is a component of the

septate-like paranodal junctions that assemble during myelination. *J Cell Biol, 139*(6),
1495-1506. doi:10.1083/jcb.139.6.1495

Fletcher, E. V., Simon, C. M., Pagiazitis, J. G., Chalif, J. I., Vukojicic, A., Drobac, E., . . . Mentis, G.
Z. (2017). Reduced sensory synaptic excitation impairs motor neuron function via Kv2.1
in spinal muscular atrophy. *Nat Neurosci, 20*(7), 905-916. doi:10.1038/nn.4561

Gravagna, N. G., Knoeckel, C. S., Taylor, A. D., Hultgren, B. A., & Ribera, A. B. (2008). Localization
of Kv2.2 protein in Xenopus laevis embryos and tadpoles. *J Comp Neurol, 510*(5), 508-
524. doi:10.1002/cne.21804

Ha, H. (1970). Axonal bifurcation in the dorsal root ganglion of the cat: a light and electron
microscopic study. *J Comp Neurol, 140*(2), 227-240. doi:10.1002/cne.901400206

He, X. H., Zang, Y., Chen, X., Pang, R. P., Xu, J. T., Zhou, X., . . . Liu, X. G. (2010). TNF-α
contributes to up-regulation of Nav1.3 and Nav1.8 in DRG neurons following motor fiber
injury. *Pain, 151*(2), 266-279. doi:10.1016/j.pain.2010.06.005

Hermanstyne, T. O., Kihira, Y., Misono, K., Deitchler, A., Yanagawa, Y., & Misonou, H. (2010).
Immunolocalization of the voltage-gated potassium channel Kv2.2 in GABAergic neurons
in the basal forebrain of rats and mice. *J Comp Neurol, 518*(21), 4298-4310.
doi:10.1002/cne.22457

Hwang, P. M., Fotuhi, M., Bredt, D. S., Cunningham, A. M., & Snyder, S. H. (1993). Contrasting
immunohistochemical localizations in rat brain of two novel K+ channels of the Shab
subfamily. *J Neurosci, 13*(4), 1569-1576. doi:10.1523/jneurosci.13-04-01569.1993

Hwang, P. M., Glatt, C. E., Bredt, D. S., Yellen, G., & Snyder, S. H. (1992). A novel K+ channel with
unique localizations in mammalian brain: molecular cloning and characterization.
Neuron, 8(3), 473-481. doi:10.1016/0896-6273(92)90275-i

Irie, T. (2021). Essential role of somatic Kv2 channels in high-frequency firing in cartwheel cells
of the dorsal cochlear nucleus. *eNeuro, 8*(3). doi:10.1523/eneuro.0515-20.2021

Jensen, C. S., Watanabe, S., Stas, J. I., Klaphaak, J., Yamane, A., Schmitt, N., . . . Misonou, H.
(2017). Trafficking of Kv2.1 channels to the axon initial segment by a novel
nonconventional secretory pathway. *J Neurosci, 37*(48), 11523-11536.
doi:10.1523/jneurosci.3510-16.2017

Johnson, B., Leek, A. N., Sole, L., Maverick, E. E., Levine, T. P., & Tamkun, M. M. (2018). Kv2
potassium channels form endoplasmic reticulum/plasma membrane junctions via
interaction with VAPA and VAPB. *Proc Natl Acad Sci U S A, 115*(31), E7331-e7340.
doi:10.1073/pnas.1805757115

Johnston, J., Griffin, S. J., Baker, C., Skrzypiec, A., Chernova, T., & Forsythe, I. D. (2008). Initial
segment Kv2.2 channels mediate a slow delayed rectifier and maintain high frequency
action potential firing in medial nucleus of the trapezoid body neurons. *J Physiol,*
586(14), 3493-3509. doi:10.1113/jphysiol.2008.153734

Kihira, Y., Hermanstyne, T. O., & Misonou, H. (2010). Formation of heteromeric Kv2 channels in
mammalian brain neurons. *J Biol Chem, 285*(20), 15048-15055.
doi:10.1074/jbc.M109.074260

Kim, K. X., & Rutherford, M. A. (2016). Maturation of NaV and Kv channel topographies in the
auditory nerve spike initiator before and after developmental onset of hearing function.
J Neurosci, 36(7), 2111-2118. doi:10.1523/jneurosci.3437-15.2016

King, A. N., Manning, C. F., & Trimmer, J. S. (2014). A unique ion channel clustering domain on
the axon initial segment of mammalian neurons. *J Comp Neurol, 522*(11), 2594-2608.
doi:10.1002/cne.23551
Kirmiz, M., Vierra, N. C., Palacio, S., & Trimmer, J. S. (2018). Identification of VAPA and VAPB as
Kv2 channel-interacting proteins defining endoplasmic reticulum-plasma membrane
junctions in mammalian brain neurons. *J Neurosci*. doi:10.1523/jneurosci.0893-18.2018
Lee, M. C., Nahorski, M. S., Hockley, J. R. F., Lu, V. B., Ison, G., Pattison, L. A., . . . Woods, C. G.
(2020). Human labor pain is influenced by the voltage-gated potassium channel Kv 6.4
subunit. *Cell Rep, 32*(3), 107941. doi:10.1016/j.celrep.2020.107941
Legland, D., Arganda-Carreras, I., & Andrey, P. (2016). MorphoLibJ: integrated library and
plugins for mathematical morphology with ImageJ. *Bioinformatics, 32*(22), 3532-3534.
doi:10.1093/bioinformatics/btw413
Li, X. N., Herrington, J., Petrov, A., Ge, L., Eiermann, G., Xiong, Y., . . . Zhou, Y. P. (2013). The role
of voltage-gated potassium channels Kv2.1 and Kv2.2 in the regulation of insulin and
somatostatin release from pancreatic islets. *J Pharmacol Exp Ther, 344*(2), 407-416.
doi:10.1124/jpet.112.199083
Lim, S. T., Antonucci, D. E., Scannevin, R. H., & Trimmer, J. S. (2000). A novel targeting signal for
proximal clustering of the Kv2.1 K+ channel in hippocampal neurons. *Neuron, 25*(2), 385-
397. doi:10.1016/s0896-6273(00)80902-2
Liu, P. W., & Bean, B. P. (2014). Kv2 channel regulation of action potential repolarization and
firing patterns in superior cervical ganglion neurons and hippocampal CA1 pyramidal
neurons. *J Neurosci, 34*(14), 4991-5002. doi:10.1523/jneurosci.1925-13.2014
Matsuda, S., Kobayashi, N., Terashita, T., Shimokawa, T., Shigemoto, K., Mominoki, K., . . .
Fujiwara, T. (2005). Phylogenetic investigation of Dogiel's pericellular nests and Cajal's
initial glomeruli in the dorsal root ganglion. *J Comp Neurol, 491*(3), 234-245.
doi:10.1002/cne.20713
Misonou, H., Mohapatra, D. P., Menegola, M., & Trimmer, J. S. (2005). Calcium- and metabolic
state-dependent modulation of the voltage-dependent Kv2.1 channel regulates
neuronal excitability in response to ischemia. *J Neurosci, 25*(48), 11184-11193.
doi:10.1523/jneurosci.3370-05.2005
Muennich, E. A., & Fyffe, R. E. (2004). Focal aggregation of voltage-gated, Kv2.1 subunit-
containing, potassium channels at synaptic sites in rat spinal motoneurones. *J Physiol,
554*(Pt 3), 673-685. doi:10.1113/jphysiol.2003.056192
Newkirk, G. S., Guan, D., Dembrow, N., Armstrong, W. E., Foehring, R. C., & Spain, W. J. (2022).
Kv2.1 potassium channels regulate repetitive burst firing in extratelencephalic
neocortical pyramidal neurons. *Cereb Cortex, 32*(5), 1055-1076.
doi:10.1093/cercor/bhab266
Panzera, L. C., Johnson, B., Quinn, J. A., Cho, I. H., Tamkun, M. M., & Hoppa, M. B. (2022).
Activity-dependent endoplasmic reticulum Ca(2+) uptake depends on Kv2.1-mediated
endoplasmic reticulum/plasma membrane junctions to promote synaptic transmission.
Proc Natl Acad Sci U S A, 119(30), e2117135119. doi:10.1073/pnas.2117135119
Rasband, M. N., Trimmer, J. S., Schwarz, T. L., Levinson, S. R., Ellisman, M. H., Schachner, M., &
Shrager, P. (1998). Potassium channel distribution, clustering, and function in

remyelinating rat axons. *J Neurosci, 18*(1), 36-47. doi:10.1523/jneurosci.18-01-
00036.1998

Regnier, G., Bocksteins, E., Van de Vijver, G., Snyders, D. J., & van Bogaert, P. P. (2016). The
contribution of Kv2.2-mediated currents decreases during the postnatal development of
mouse dorsal root ganglion neurons. *Physiol Rep, 4*(6). doi:10.14814/phy2.12731

Romer, S. H., Dominguez, K. M., Gelpi, M. W., Deardorff, A. S., Tracy, R. C., & Fyffe, R. E. W.
(2014). Redistribution of Kv2.1 ion channels on spinal motoneurons following peripheral
nerve injury. *Brain research, 1547*, 1-15. doi:10.1016/j.brainres.2013.12.012

Sarmiere, P. D., Weigle, C. M., & Tamkun, M. M. (2008). The Kv2.1 K+ channel targets to the
axon initial segment of hippocampal and cortical neurons in culture and in situ. *BMC*
Neurosci, 9, 112. doi:10.1186/1471-2202-9-112

Scannevin, R. H., Murakoshi, H., Rhodes, K. J., & Trimmer, J. S. (1996). Identification of a
cytoplasmic domain important in the polarized expression and clustering of the Kv2.1 K+
channel. *J Cell Biol, 135*(6 Pt 1), 1619-1632. doi:10.1083/jcb.135.6.1619

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., . . . Cardona, A.
(2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods, 9*(7),
676-682. doi:10.1038/nmeth.2019

Scott, B., Leu, J., & Cinader, B. (1988). Effects of aging on neuronal electrical membrane
properties. *Mech Ageing Dev, 44*(3), 203-214. doi:10.1016/0047-6374(88)90022-x

Shields, S. D., Ahn, H. S., Yang, Y., Han, C., Seal, R. P., Wood, J. N., . . . Dib-Hajj, S. D. (2012).
Nav1.8 expression is not restricted to nociceptors in mouse peripheral nervous system.
Pain, 153(10), 2017-2030. doi:10.1016/j.pain.2012.04.022

Shiers, S., Klein, R. M., & Price, T. J. (2020). Quantitative differences in neuronal subpopulations
between mouse and human dorsal root ganglia demonstrated with RNAscope in situ
hybridization. *Pain, 161*(10), 2410-2424. doi:10.1097/j.pain.0000000000001973

Shiers, S., Sankaranarayanan, I., Jeevakumar, V., Cervantes, A., Reese, J. C., & Price, T. J. (2021).
Convergence of peptidergic and non-peptidergic protein markers in the human dorsal
root ganglion and spinal dorsal horn. *J Comp Neurol, 529*(10), 2771-2788.
doi:10.1002/cne.25122

Sun, Z. W., Waybright, J. M., Beldar, S., Chen, L., Foley, C. A., Norris-Drouin, J. L., . . . Wang, Y.
(2022). Cdyl deficiency brakes neuronal excitability and nociception through promoting
Kcnb1 transcription in peripheral sensory neurons. *Adv Sci (Weinh), 9*(10), e2104317.
doi:10.1002/advs.202104317

Tavares-Ferreira, D., Shiers, S., Ray, P. R., Wangzhou, A., Jeevakumar, V., Sankaranarayanan, I., .
. . Price, T. J. (2022). Spatial transcriptomics of dorsal root ganglia identifies molecular
signatures of human nociceptors. *Sci Transl Med, 14*(632), eabj8186.
doi:10.1126/scitranslmed.abj8186

Thapa, P., Stewart, R., Sepela, R. J., Vivas, O., Parajuli, L. K., Lillya, M., . . . Sack, J. T. (2021).
EVAP: A two-photon imaging tool to study conformational changes in endogenous Kv2
channels in live tissues. *J Gen Physiol, 153*(11). doi:10.1085/jgp.202012858

Trimmer, J. S. (1991). Immunological identification and characterization of a delayed rectifier K+
channel polypeptide in rat brain. *Proc Natl Acad Sci U S A, 88*(23), 10764-10768.
doi:10.1073/pnas.88.23.10764

Tsantoulas, C., Zhu, L., Shaifta, Y., Grist, J., Ward, J. P., Raouf, R., . . . McMahon, S. B. (2012).
Sensory neuron downregulation of the Kv9.1 potassium channel subunit mediates
neuropathic pain following nerve injury. *J Neurosci, 32*(48), 17502-17513.
doi:10.1523/jneurosci.3561-12.2012
Tsantoulas, C., Zhu, L., Yip, P., Grist, J., Michael, G. J., & McMahon, S. B. (2014). Kv2 dysfunction
after peripheral axotomy enhances sensory neuron responsiveness to sustained input.
Exp Neurol, 251, 115-126. doi:10.1016/j.expneurol.2013.11.011
Usoskin, D., Furlan, A., Islam, S., Abdo, H., Lönnerberg, P., Lou, D., . . . Ernfors, P. (2015).
Unbiased classification of sensory neuron types by large-scale single-cell RNA
sequencing. *Nat Neurosci, 18*(1), 145-153. doi:10.1038/nn.3881
Valtcheva, M. V., Copits, B. A., Davidson, S., Sheahan, T. D., Pullen, M. Y., McCall, J. G., . . .
Gereau, R. W. t. (2016). Surgical extraction of human dorsal root ganglia from organ
donors and preparation of primary sensory neuron cultures. *Nat Protoc, 11*(10), 1877-
1888. doi:10.1038/nprot.2016.111
Vierra, N. C., O'Dwyer, S. C., Matsumoto, C., Santana, L. F., & Trimmer, J. S. (2021). Regulation
of neuronal excitation-transcription coupling by Kv2.1-induced clustering of somatic L-
type Ca(2+) channels at ER-PM junctions. *Proc Natl Acad Sci U S A, 118*(46).
doi:10.1073/pnas.2110094118
Vullhorst, D., Mitchell, R. M., Keating, C., Roychowdhury, S., Karavanova, I., Tao-Cheng, J. H., &
Buonanno, A. (2015). A negative feedback loop controls NMDA receptor function in
cortical interneurons via neuregulin 2/ErbB4 signalling. *Nat Commun, 6*, 7222.
doi:10.1038/ncomms8222
Wang, H., & Zylka, M. J. (2009). Mrgprd-expressing polymodal nociceptive neurons innervate
most known classes of substantia gelatinosa neurons. *J Neurosci, 29*(42), 13202-13209.
doi:10.1523/jneurosci.3248-09.2009
Wangzhou, A., McIlvried, L. A., Paige, C., Barragan-Iglesias, P., Shiers, S., Ahmad, A., . . . Price, T.
J. (2020). Pharmacological target-focused transcriptomic analysis of native vs cultured
human and mouse dorsal root ganglia. *Pain, 161*(7), 1497-1517.
doi:10.1097/j.pain.0000000000001866
Zheng, Y., Liu, P., Bai, L., Trimmer, J. S., Bean, B. P., & Ginty, D. D. (2019). Deep sequencing of
somatosensory neurons reveals molecular determinants of intrinsic physiological
properties. *Neuron, 103*(4), 598-616.e597. doi:10.1016/j.neuron.2019.05.039
833

Figure 1

Figure 1 legend

Kv2.1 and Kv2.2 protein are enriched at the outer edge of DRG neuron somas relative to

Nav1.8. Lumbar DRG section from an 11 week old male mouse labeled with antibodies which

target Kv2.1, Kv2.2 or Nav1.8. *A*, Anti-Kv2.1 (magenta) and anti-Kv2.2 (green)

immunofluorescence in a lumbar DRG section. Scale bar is 100 μm. *B*, Anti-Kv2.1, anti-Kv2.2 and

anti-Nav1.8 immunofluorescence from box shown in A. Arrows indicate prominent localization

of anti-Kv2 immunofluorescence at the edge of DRG neuron somas. In merge image anti-Kv2.1,

anti-Kv2.2 and anti-Nav1.8 immunofluorescence is magenta, green and blue respectively. Scale

bar is 20 μm. *C*, Representative ROIs that encompass the outer edge of DRG neurons (arrow 1)

and the region just inside the outer edge (arrow 2). *D*, Ratio of anti-Kv2.1, anti-Kv2.2 or anti-

Nav1.8 immunofluorescence from outer and inner ROIs for individual neurons from image in A.

Bars represent mean. One-way ANOVA $p < 0.001$. p values in figure represent post hoc Tukey's

test. N = 1 mouse, n = 124 neurons. Detailed information on mouse used can be found in table

1.

Figure 2

Figure 2 legend

Kv2.1 protein is detectable in mouse DRG neurons. *A*, WT (top) and Kv2.1 KO (bottom) DRG

sections taken from 7 week old female mice from the 13[th] thoracic DRG immunolabeled for

Kv2.1 (magenta) and BIII tubulin (white). Images were taken with identical imaging settings and

are set to the same brightness and contrast. Scale bars are 100 μm. *B*, Distribution of

fluorescence intensity from manual analysis of WT (black) and Kv2.1 KO (red) neurons. Dotted

lines represent mean. Data represent fluorescence intensity of 254 WT profiles from 10 DRG

sections from 1 mouse or 375 Kv2.1 KO profiles from 5 DRG sections from 1 mouse. Images

shown in A represent one section from WT and Kv2.1 KO mice used in this data set. *C*,

Distribution of fluorescence intensity from automated analysis of the same data set shown in B.

Dotted lines represent mean. Data represent fluorescence intensity of 476 WT or 576 Kv2.1 KO

profiles selected by automated analysis method. *D*, Distribution of BIII tubulin fluorescence

intensity from the same WT (black) and Kv2.1 KO (red) profiles shown in C. Dotted lines

represent mean. *E*, Mean fluorescence intensity of Kv2.1 KO neurons normalized to WT neurons

labeled with anti-Kv2.1, anti-Kv2.2 and anti-BIII tubulin antibodies. Each point represents one

Kv2.1 KO mouse normalized to one age and sex matched WT mouse which was stained

simultaneously and imaged with identical microscopy settings. The color of each point

represents the same mouse and purple points represent data from the mouse whose DRG

immunofluorescence data are shown in A, B, C and D. one-way ANOVA $p < 0.001$. p values in

figure represent post hoc Tukey's test. *F*, Percentage of ROIs with anti-Kv2.1

immunofluorescence above the mean immunofluorescence of 5 mice (1 female and 4 male).

Point colors correspond to the WT mice analyzed in E. All mice were compared to age and sex

matched Kv2.1 KO mice. N = 5 WT and 5 Kv2.1 KO mice. **G**, Kv2.1 KO data shown in B fit with a

log normal distribution (red fit). **H**, WT data shown in B fit with the Kv2.1 KO distribution (red

fit) where width and mean were constrained to the Kv2.1 KO distribution and amplitude was

unconstrained (equation 1). Red dotted line represents the mean of the Kv2.1 KO distribution.

Only WT data to the left of red dotted line was used for the fit. **I**, Percentage of ROIs with

detectable Kv2.1 protein of 5 mice (1 female and 4 males). Point colors correspond to the WT

mice analyzed in E. All mice were compared to age and sex matched Kv2.1 KO mice. N = 5 WT

and 5 Kv2.1 KO mice. Detailed information on each mouse used can be found in table 1.

Figure 3

Figure 3 legend

Kv2.2 protein is detectable in mouse DRG neurons. **A**, WT (top) and Kv2.2 KO (bottom) DRG

sections from the 13th thoracic DRG in 7 week old male mice immunolabeled for Kv2.2 (green)

and NF200 (white). Identical imaging and display settings. Scale bars are 100 μm. **B**,

Distribution of fluorescence intensities from manual analysis of WT (black) and Kv2.2 KO (red)

profiles. Dotted lines represent mean. Data represents fluorescence intensities from 241 WT

profiles from 11 DRG sections and 1 mouse or 130 Kv2.2 KO profiles from 6 DRG sections and 1

mouse. Images shown in A represent one section from WT and Kv2.1 KO mice used in this data

set. **C**, Distribution of fluorescence intensity from automated analysis of the same data set

shown in B. Dotted lines represent mean. Data represent fluorescence intensity of 673 WT or

400 Kv2.2 KO profiles selected by automated analysis method. **D**, Distribution of anti-NF200

immunofluorescence intensity from the same WT (black) and Kv2.2 KO (red) neurons shown in

B. Dotted lines represent mean. **E**, Mean fluorescence intensity of Kv2.2 KO ROIs normalized to

WT neurons labeled with anti-Kv2.1, anti-Kv2.2 and anti-NF200 antibodies. Each point

represents one Kv2.2 KO mouse normalized to one age and sex matched WT mouse which was

stained simultaneously and imaged with identical microscopy settings. The color of each point

represents the same mouse and purple points represent data from the male mouse whose DRG

immunofluorescence data are shown in A, B and C. Missing points in anti-Kv2.1 column are

because some sections were not labeled with anti-Kv2.1 antibodies. one-way ANOVA $p < 0.001$.

p values in figure represent Tukey's post hoc test. **F**, Kv2.2 KO data shown in B fit with a log

normal distribution (red fit). **G**, WT data shown in B fit with the Kv2.2 KO distribution (red fit)

where width and mean were constrained to the Kv2.2 KO distribution and amplitude was

unconstrained (equation 1). Red dotted line represents the mean of the Kv2.2 KO distribution.

Only WT data to the left of red dotted line was used for the fit. *H*, Percentage of ROIs with

detectable Kv2.2 protein of 8 mice (7 males and 1 female). Point colors correspond to the WT

mice analyzed in D. All mice were compared to age and sex matched Kv2.2 KO mice. N = 8 WT

and 8 Kv2.2 KO mice. Detailed information on each mouse used can be found in table 1.

 Figure 4

Figure 4 legend

Detectable Kv2.1 protein decreases in older mice while detectable Kv2.2 does not. *A*, DRG

sections from the 13[th] thoracic DRG in 7 week (left) and 50 week (right) old mice

immunolabeled for Kv2.1. Vertical bar on right is pseudo coloring key for pixel intensity.

Identical imaging and display settings. Scale bars are 100 µm. *B*, Distribution of fluorescence

intensities from 7 week old WT (black) and 7 week old Kv2.1 KO (red) ROIs generated by

automated method. 609 WT ROIs from 1 mouse. 367 Kv2.1 KO ROIs from 1 mouse. *C*,

Distribution of fluorescence intensities from 50 week old WT (black) and 50 week old Kv2.1 KO

(red) ROIs. 793 WT ROIs from 1 mouse. 378 Kv2.1 KO ROIs from 1 mouse. *D*, Percentage of ROIs

with detectable Kv2.1 protein in 7-16 week old and 50 week old mice. Data from 7-16 week old

mice is the same data in Figure 2. N = 4 mice 7 weeks old and 1 mouse 16 weeks old and N = 5

mice 50 weeks old. Detailed information on each mouse used can be found in table 1. *E*, DRG

sections from the 13th thoracic DRG in 7 week (left) and 50 week (right) old mice

immunolabeled for Kv2.2. Vertical bar on right is pseudo coloring key for pixel intensity.

Identical imaging and display settings. Scale bars are 100 µm. *F*, Distribution of fluorescence

intensities from 7 week old WT (black) and 7 week old Kv2.2 KO (red) ROIs generated by

automated method. 746 WT ROIs from 1 mouse. 717 Kv2.2 KO ROIs from 1 mouse. Data from

same mice shown in E. *G*, Distribution of fluorescence intensities from 50 week old WT (black)

and 50 week old Kv2.2 KO (red) ROIs generated by automated method. 671 WT ROIs from 1

mouse. 398 Kv2.2 KO ROIs from 1 mouse. Data from same mice shown in E. *H*, Percentage of

ROIs with detectable Kv2.2 protein in 7-24 week old and 50 week old mice. Data is the same

data from Figure 3 where mice were separated into a young group (7-24 weeks) and an old

group (50 weeks). N = 3 mice 7 weeks old, 1 mouse 24 weeks old and 1 mouse 25 weeks old. N

= 3 mice 50 weeks old. Detailed information on each mouse used can be found in table 1.

Figure 5

Figure 5 legend

Kv2 channels are expressed at the surface membrane of DRG neurons. *A*, Fluorescence of live

dissociated DRG neurons excited at 594 nm before (top left) and after (bottom left) application

of 100 nM GxTX-594. Fluorescence from membrane marker WGA-405 before (top middle) and

after application of 100 nM GxTX-594 (bottom middle). Merge image shows 594 excitation

fluorescence (magenta) and 405 excitation fluorescence (green). Scale bar 20 µm. *B*,

Dissociated DRG neurons from WT (top) and Kv2.1/Kv2.2 DKO (bottom) mice before and after

the application of GxTX-594, left panel and middle panel respectively. Arrows in middle panel

indicate location of surface membrane based on WGA-405 fluorescence. Right images are

merge of 594 excitation fluorescence (magenta) and 405 excitation fluorescence (green) after

application of 100 nM GxTX-594. Scale bars 100 µm. *C*, Example of WGA-405 fluorescence (left)

used in watershed segmentation (middle) to generate annulus ROI (right) used to analyze

fluorescence intensity at the membrane. *D*, Distribution of fluorescence intensity from WT

(black) and Kv2.1/Kv2.2 DKO (red) neurons. Data represents the fluorescence intensity of 326

WT neurons from 1 mouse or 271 Kv2.1/Kv2.2 DKO neurons from 1 mouse. DRG from all levels

of the spinal cord were pooled. *E*, Kv2.1/Kv2.2 DKO data shown in D fit with a log normal

distribution (red fit). *F*, WT data shown in D fit with the Kv2.1/Kv2.2 DKO distribution (red fit)

where width and mean were constrained to the Kv2.1/Kv2.2 DKO distribution and amplitude

was unconstrained (equation 1). Red dotted line represents the mean of the Kv2.1/Kv2.2 DKO

distribution. Only WT data to the left of red dotted line was used for the fit. *G*, Percentage of

neurons with detectable surface Kv2 protein, from an experiment where one WT mouse was

compared to one DKO mouse and an identical experiment where two WT mice were compared

to one DKO mouse (N=3 WT mice N=2 Kv2.1/Kv2.2 DKO mice). N = 3 WT mice and N = 2 DKO

mice. Detailed information on each mouse used can be found in table 1.

 Figure 6

Figure 6 legend

DRG neurons have enriched Kv2.2 protein compared to neurons in the spinal cord. **A**, Anti-

Kv2.1 (magenta) and anti-Kv2.2 (green) immunofluorescence in a spinal cord section from the

2nd lumbar vertebra (left). Anti-Kv2.1 immunofluorescence (right top) and anti-Kv2.2

immunofluorescence (right bottom). Arrows show neurons in the spinal cord with punctate

anti-Kv2.1 immunofluorescence. Arrow heads show neurons in the spinal cord with anti-Kv2.2

immunofluorescence. Scale bars are 500 μm. **B**, Anti-Kv2.1 immunofluorescence from individual

neuron profiles (circles) from multiple mice in the DRG and ventral horn normalized to the

average fluorescence intensity of neuron profiles in the ventral horn. Diamonds to the right of

data represent the average intensity of individual mice. Significant differences from 1 were

calculated for individual mice using Students t-test. N = 5 mice, n = 295 in DRG and n = 200 in

ventral horn. Detailed information on each mouse used can be found in table 1. **C**, Identical

analysis as in panel B but with anti-Kv2.2 immunofluorescence.

Figure 7

Figure 7 legend

Kv2 channels form clusters on DRG neuron somas and stem axons that are distinct from Kv2

channel clusters on ventral horn neurons. *A*, Z-projection of anti-Kv2.1 and anti-Kv2.2

immunofluorescence in a ventral horn neuron from the 1st lumbar vertebra of a 7 week old

male mouse. *B*, Z-projection of anti-Kv2.1 and anti-Kv2.2 immunofluorescence in DRG neurons

from the same mouse and section of the spinal column as neuron shown in A. Inset is

enlargement of the Kv2.1 donut cluster in dotted box. *C*, Z-projection of anti-Kv2.1 and anti-

Kv2.2 immunofluorescence in DRG neurons from the same mouse as A and B. Inset is

enlargement of the Kv2.2 donut cluster in dotted box. *D*, Z-projection of anti-Kv2.1

immunofluorescence in DRG neuron from the 13th thoracic DRG of a 24 week old male mouse.

E, Z-projection of anti-Kv2.1 and anti-Kv2.2 immunofluorescence in DRG neuron from same

mouse in D. *F*, Exemplar ROI for analyzing localization of Kv2 channel density relative to stem

axon. Same image as E. ROI line width is 1.24 μm. Numbers along line indicate approximate

distance from stem axon normalized to the midpoint of the line. *G*, Anti-Kv2.1 (magenta) and

anti-Kv2.2 (green) immunofluorescence intensity along the ROI shown in F. Distance along the

line was normalized such that the stem axon is 0 and the midpoint of the line is 1. *H*, Analysis of

the relative distance from the stem axon of the max anti-Kv2.1 or anti-Kv2.2

immunofluorescence. Dotted line represents the middle of neurons relative to the stem axon.

In all images arrows indicate asymmetrical clusters of Kv2 channels while arrow heads indicate

the apparent stem axons. Display settings are not identical between images. Scale bars are 20

μm.

1008

Figure 8

Figure 8 legend

Kv2.2 channels are expressed in peripheral axons of DRG neurons. *A*, WT (top) and Kv2.2 KO

(bottom) sections containing the DRG, peripheral and central axons from the 1st lumbar DRG in

age and sex matched 7 week old mice immunolabeled for NF200 (white). Scale bar is 1 mm. *B*,

High magnification z-projection of anti-Kv2.2 and anti-NF200 immunofluorescence from box 1

in A of WT and Kv2.2 KO mice. Arrows indicate myelinated axons which show prominent anti-

Kv2.2 immunofluorescence. Scale bars are 20 μm. *C*, High magnification z-projection of anti-

Kv2.2 and anti-NF200 immunofluorescence from box 2 in A of WT and Kv2.2 KO mice. Arrows

indicate myelinated axons which show prominent anti-Kv2.2 immunofluorescence. Scale bars

are 20 μm. *D*, High magnification z-projection of anti-Kv2.2 immunofluorescence and MrgprD-

GFP fluorescence in the peripheral axons of the 12th thoracic DRG of a 13 week old MrgprD-GFP

mouse. Arrow indicates anti-Kv2.2 immunofluorescence on a GFP+ axon. Scale bar is 10 μm.

1023

 Figure 9

 Figure 9 legend

 Anti-Kv2 immunofluorescence intensities are non-uniform across DRG neuron subtypes. ***A***,

 Exemplar z-projection of enrichment of anti-Kv2.1 (magenta) or anti-Kv2.2 (green)

immunofluorescence in neighboring neurons. DRG section is from a 10 week old male mouse.

Scale bar is 10 μm. **B**, Top images show anti-Kv2.1 (magenta) and anti-Kv2.2 (green)

immunofluorescence in DRG sections where subpopulation specific markers were used to

identify, from left to right, non-peptidergic nociceptors, peptidergic nociceptors, myelinated

neurons and proprioceptors. Fluorescence from specific markers is shown in bottom panels.

Arrows indicate four exemplar neurons that have clear positivity for each subpopulation

identified by fluorescence in lower panels. CGRP and NF200 subpopulations were identified

using anti-CGRP and anti-NF200 antibodies while MrgprD-GFP and PV-tdTomato

subpopulations were from transgenic mouse lines. Scale bars are 50 μm. **C**, Scatter plot of anti-

Kv2.1 and anti-Kv2.2 immunofluorescence of individual neuron profiles. Each point represents

one profile. Magenta circle highlights the subpopulation of profiles that have high anti-Kv2.1

but low anti-Kv2.2 immunofluorescence while the green circle highlights the subpopulation of

profiles that have high anti-Kv2.2 but low anti-Kv2.1 immunofluorescence. Blue points

represent myelinated DRG neuron profiles identified by NF200 immunofluorescence. **D**, Ranked

anti-Kv2.1 immunofluorescence (magenta points) or ranked anti-Kv2.2 immunofluorescence

(green points) of individual profiles from subpopulations shown in B. Only profiles that were

positive for each marker are shown. Each point represents one profile. MrgprD population N = 4

mice, CGRP population N = 3 mice, NF200 population N = 3 mice and PV population N = 2 mice.

Detailed information on each mouse used can be found in table 1.

1048

Figure 10

1050

Figure 10 legend

Kv2 channel expression and localization in human DRG neurons is similar to mice. **A**,

Immunofluorescence from human DRG neurons labeled with anti-Kv2.1 and anti-Kv2.2

antibodies. Autofluorescence attributed to lipofuscin is labeled in right panel while apparent

Kv2.1 and Kv2.2 protein are labeled in left and middle panel respectively. Scale bar is 50 μm. **B**,

Z-projection of anti-Kv2.2 (left) and anti-NF200 immunofluorescence (middle) of human DRG

neuron somata. Green arrow head indicates asymmetric distribution of Kv2.2 clusters on

neuron soma, green arrow indicates the apparent stem axon. Scale bar is 20 μm. **C**, Z-projection

of anti-Kv2.1 (upper left), anti-Kv2.2 (upper right) and anti-NF200 (lower left)

immunofluorescence of a human DRG neuron. Magenta and green arrows indicate Kv2.1 and

Kv2.2 respectively on the apparent stem axon. Inset shows expansion of dotted line boxes

which highlights Kv2.1 and Kv2.2 clusters on the apparent stem axon. Autofluorescence

attributed to lipofuscin is labeled in lower right panel. Scale bar is 50 µm. All images are from

donor #1. Detailed information on each donor can be found in the *Human Tissue Collection*

section of the methods.

1066

1067   **SUPPLEMENTAL FIGURES**
1068   **Supplemental Figure 1**

1069
1070   **Supplemental Figure 1 legend**
1071   Off-target mouse anti-IgG1, IgG2b, and IgG2a immunofluorescence increases with fixation time.
1072   Images of DRG sections labeled with fluorescently tagged antibodies which target IgG1 (top),
1073   IgG2b (middle) and IgG2a (bottom) after DRG were fixed in ice cold 4% PFA for 1, 3 and 12
1074   hours. DRG sections are from the same mouse. Scale bars are 100 µm.
1075

1076 **Supplemental Figure 2**

1077
Supplemental Figure 2 legend
Pre-incubation of mouse DRG sections in IgG H+L Fab fragments reduces off target secondary
antibody labeling. Representative images of DRG sections from the same DRG treated with
increasing concentrations (left to right) of IgG H+L Fab fragment and the same concentration of
secondary antibody used in experiments throughout this study. Images were taken with
identical imaging settings and are set to the same brightness and contrast. Scale bars are 100
μm.

Supplemental Figure 3

1086
Supplemental Figure 3 legend
Kv2.1 and Kv2.2 protein are enriched at the outer edge of DRG neuron somas relative to TRPV1.
A, Anti-Kv2.1, anti-Kv2.2 and anti-TRPV1 immunofluorescence from lumbar DRG neurons.
Prominent cytoplasmic anti-TRPV1 immunofluorescence was observed in a subset of small
diameter neurons. In merge image anti-Kv2.1, anti-Kv2.2 and anti-TRPV1 immunofluorescence
are magenta, green and blue respectively. Scale bar is 20 μm. ***B***, Ratio of average anti-Kv2.1,
anti-Kv2.2 or anti-TRPV1 immunofluorescence from outer and inner ROIs for individual
neurons.
1095
1096

1097   **Supplemental Figure 4**

1098
Supplemental Figure 4 legend
Kv2 antibodies used in knockout experiments are at saturating concentrations. ***A***,
Concentration response of immunofluorescence from sections labeled with anti-Kv2.1 antibody
used in Figure 2. Blue line is a Hill fit of the data. 1:1 = 2 sections, 1:25 = 3 sections, 1:125 = 2
sections, 1:625 = 3 sections ***B***, Concentration response of immunofluorescence from sections
labeled with anti-Kv2.2 antibody used in Figure 3. Blue line is a Hill fit of the data. 1:33 = 2
sections, 1:100 = 10 sections, 1:300 = 4 sections, 1:900 = 4 sections, 1:2700 = 3 sections
1106

 Supplemental Figure 5

Supplemental Figure 5 legend

A method to sample neurons in DRG imaging data using watershed segmentation identifies the outer region of neurons. ***A***, Image of anti-Kv2.1, anti-Kv2.2, anti-βIII tubulin immunofluorescence and MrgprD-GFP fluorescence. ***B***, Same image as A with anti-Kv2.1 immunofluorescence channel removed as this is an example of processing data for Kv2.1 KO analysis. ***C***, Grayscale image of average fluorescence from all three channels shown in B. Gaussian and median filters were applied to image to improve watershed segmentation. ***D***, Watershed segmentation of image in C using the MorphoLibJ Morphological Segmentation Plugin in Fiji. ***E***, Example of manually drawn boundary that encompasses the neuron somas in the DRG so that only these watershed lines are selected. ***F***, Selected ROIs from watershed segmentation (yellow). ROIs were excluded based on roundness and size using the Analyze

Particles tool in Fiji. **G**, ROIs in F overlaid on DRG image showing that some ROIs are selecting
regions that do not contain neurons (arrow heads) or are selecting multiple neurons (arrows).
H, ROIs after processing using an in-house R script which removes ROIs that do not contain
neurons (arrow heads) and ROIs that contain two neurons (arrows). This script did not remove
all ROIs that do not contain neurons (red arrows). Each experiment performed was done
alongside controls where the primary antibodies were omitted and fluorescence from these
control sections was used by the in-house R script to identify and remove ROIs that do not
contain neurons. **I**, Example of automatically generated annulus that encompasses the outer
edge of the soma. Scale bars are 100 μm.
1129

Supplemental Figure 6

1131

Supplemental Figure 6 legend

Method used to estimate percent of neurons expressing Kv2.1 and Kv2.2 reliably predicts the percentage of neurons that express GFP in MrgprD-GFP mice. **A**, MrgprD-GFP (top) and WT (bottom) DRG sections immunolabeled for BIII tubulin (white). Images were taken with identical imaging settings and are set to the same brightness and contrast. Scale bars are 100 µm. **B**, Distribution of fluorescence intensity from MrgprD-GFP (black) and WT (red) neurons. Data represents the fluorescence intensity of 905 MrgprD-GFP neurons from 9 DRG sections from 1 mouse or 477 WT neurons from 5 DRG sections from 1 mouse. DRG sections were taken from 7 week old female mice and are from the 1st lumbar DRG. **C**, WT data shown in B fit with a log normal distribution (red fit). **D**, MrgprD-GFP data shown in B fit with the WT distribution (red fit) where width and mean were constrained to the WT distribution and amplitude was unconstrained (equation 1). Only MrpgrD-GFP data to the left of the mean intensity of WT neurons (red dotted line) was used for the fit. **E**, Percent of neurons with detectable GFP protein of 4 mice (3 females 1 male). All DRG sections were taken from the 1st lumbar DRG.

1146 **Supplemental Figure 7**

1147

Supplemental Figure 7 legend

DRG neurons have enriched Kv2.2 protein compared to neurons in the ventral horn. **A**, Anti-

Kv2.1 (magenta) and anti-Kv2.2 (green) immunofluorescence in a spinal cord section from the

13[th] thoracic vertebra (left). Anti-Kv2.1 immunofluorescence (right top) and anti-Kv2.2

immunofluorescence (right bottom). Arrow heads show neurons in the spinal cord with anti-

Kv2.2 immunofluorescence. Scale bars are 500 μm. **B**, Anti-Kv2.1 immunofluorescence from

individual neurons (circles) in the DRG and ventral horn normalized to the average fluorescence
intensity of neurons in the ventral horn. Diamonds to the right of data represent the average
intensity in the DRG of individual mice. Significant differences from 1 were calculated for
individual mice using Students t-test. N = 4 mice n = 116 in DRG and n = 77 in ventral horn. *C*,
Identical analysis shown in B with anti-Kv2.2 immunofluorescence.
1159

1160 **Supplemental Figure 8**

1161

Supplemental Figure 8 legend

Kv2 channels are expressed on the stem axon of mouse DRG neurons. ***A***, Z-projection with anti-

Kv2.1 and anti-Kv2.2 immunofluorescence on the stem axon of a neuron in the DRG of a

MrgprD-GFP mouse. ***B***, Gallery of z-projected images of DRG neurons with anti-Kv2.1 and/or

anti-Kv2.2 immunofluorescence on stem axons. Arrows indicate stem axons. Scale bars are 10

μm

Supplemental Figure 9

1169

Supplemental Figure 9 legend

Kv2.1 channels were not detected in peripheral axons of DRG neurons. ***A***, WT (top) and Kv2.1

KO (bottom) sections containing the DRG and peripheral axons from the 12[th] thoracic DRG in

age and sex matched 7 week old mice immunolabeled for βIII tubulin (white). Scale bar is 500

μm. ***B***, High magnification z-projection of anti-Kv2.1 and anti-βIII immunofluorescence from box

in A of WT and Kv2.1 KO mice. Scale bars are 20 μm.

1176

1177

1178 **Supplemental Figure 10**

1179

Supplemental Figure 10 legend

Kv2.2 is expressed in myelinated fibers of DRG neuron axons. **A**, Kv2.2 KO (top) and WT (middle

and bottom) sections containing the peripheral axons from the 12th thoracic DRG in 28 week

old mice immunolabeled for Kv2.2, Caspr and Kv1.2. Middle panels are an exemplar of

prominent Kv2.2 immunofluorescence in CASPR labeled axons and bottom panels are an

exemplar of prominent Kv2.2 clusters in the Kv1.2 labeled axons. Scale bars are 5 µm. **B**,

Analysis of anti-Kv2.2 immunofluorescence intensity in CASPR and Kv1.2 labeled regions of age

and sex matched WT and Kv2.2 KO mice. Individual points represent single ROIs drawn around

anti-CASPR or anti-Kv1.2 immunofluorescence.

1189

1190

1191 **Supplemental Figure 11**

1192

Supplemental Figure 11 legend
Fluorescence from human DRG neurons labeled with ion channel targeting antibodies is distinct
from human DRG neurons where ion channel targeting antibodies were omitted. **A**, Top:
Immunofluorescence from human DRG section labeled with anti-Kv2.1, anti-Kv2.2 and anti-
Nav1.8 antibodies. Bottom: Fluorescence from human DRG section where the primary
antibodies are omitted. Arrows in top and bottom images indicate examples of
autofluorescence from apparent intracellular lipofuscin. Arrow heads in top image identify anti-
Nav1.8 immunofluorescence. Images on the right are fluorescence from each fluorescence
channel of the top and bottom images. Number next to target protein label represents
excitation wavelength. DRG sections from top and bottom images are from the same DRG.
Scale bars are 500 μm. **B**, Left: Immunofluorescence from human DRG section labeled with anti-
Nav1.7 antibody. Right: Fluorescence from human DRG section where the primary antibody has
been omitted. Number next to target protein label represents excitation wavelength. DRG
sections in left and right images are from the same DRG. Scale bars are 500 μm. **C**, Exemplar
manually drawn ROI to analyze fluorescence intensity in human DRG neurons that omits
apparent lipofuscin autofluorescence. Scale bar is 50 μm. **D**, Distribution of fluorescence
intensity of human DRG neurons labeled with an anti-Kv2.1 antibody (black) or when the anti-
Kv2.1 antibody was omitted (red). Data represents the fluorescence intensity of 293 neurons
labeled with anti-Kv2.1 antibody or 73 neurons where the anti-Kv2.1 antibody was omitted. **E**,

1212 Distribution of fluorescence intensity of human DRG neurons labeled with anti-Kv2.2 antibody
1213 (black) or when the anti-Kv2.2 antibody was omitted (red). Data represents the fluorescence
1214 intensity of 293 neurons labeled with anti-Kv2.2 antibody or 73 neurons where the anti-Kv2.2
1215 antibody was omitted. *F*, Distribution of fluorescence intensity of human DRG neurons labeled
1216 with anti-Nav1.7 antibody (black) or when the anti-Nav1.7 antibody was omitted (red). Data
1217 represents the fluorescence intensity of 99 neurons labeled with anti-Nav1.7 antibody or 99
1218 neurons where the anti-Nav1.7 antibody was omitted. *G*, Distribution of fluorescence intensity
1219 of human DRG neurons labeled with anti-Nav1.8 antibody (black) or when the anti-Nav1.8
1220 antibody was omitted (red). Data represents the fluorescence intensity of 293 neurons labeled
1221 with anti-Nav1.8 antibody or 73 neurons where the anti-Nav1.8 antibody was omitted. *H*,
1222 Fluorescence intensity of human neurons labeled with both anti-Kv2.1 and anti-Kv2.2
1223 antibodies. Individual points represent individual neurons. All images are from donor #2.
1224 Detailed information on each donor can be found in the *Human Tissue Collection* section of the
1225 methods.
1226

1227 **Supplemental Figure 12**

1228
Supplemental Figure 12 legend
Immunofluorescence from human DRG neurons from donor #2 ***A*** and donor #3 ***B*** labeled with
anti-Kv2.1 and anti-Kv2.2 antibodies. Autofluorescence attributed to lipofuscin is labeled in
right panels while apparent Kv2.1 and Kv2.2 protein are labeled in left and middle panels
respectively. Scale bars are 50 μm. Detailed information on each donor can be found in the
Human Tissue Collection section of the methods.
1235

1236 **Supplemental Figure 13**

1237
Supplemental Figure 13 legend
Kv2 channels are enriched at the outer edge of human DRG neurons. ***A-B***, Exemplar z-
projections of anti-Kv2.1 immunofluorescence enriched at the outer surface of human DRG
neurons. Arrows indicate asymmetric clusters. Images are from donor #2 Scale bars are 20 μm.
C-F, Exemplar z-projections of anti-Kv2.2 immunofluorescence enriched at the outer surface of
a human DRG neurons. Arrows indicate asymmetric clusters. Image in ***E*** is from donor #3 while
all other images are from donor #2. Scale bars are 20 μm. ***G***, Exemplar z-projection of anti-Kv2.1
and anti-Kv2.2 immunofluorescence both enriched at the outer surface of a human DRG neuron
soma. Image is from donor #2. Scale bar is 20 μm. Detailed information on each donor can be
found in the *Human Tissue Collection* section of the methods.
1248

 Supplemental Figure 14

Supplemental Figure 14 legend
A, Z-projection of anti-Kv2.2 and anti-Nav1.7 immunofluorescence in a human DRG neuron
soma and stem axon. Arrow in merge indicates the stem axon of the DRG neuron. Apparent
lipofuscin autofluorescence is labeled in merge. Image is from donor #2. Scale bar is 20 μm. **B,**
Z-projection of anti-Kv2.1 (magenta), anti-Kv2.2 (green) (right) and anti-NF200 (left)
immunofluorescence in a human DRG neuron soma and stem axon. Arrows in merge indicate
the stem axon of the DRG neuron. Image is from donor #1. Scale bar is 20 μm. **C,** Z-projection of
anti-Kv2.2 (left) and anti-NF200 (middle) immunofluorescence in a human DRG neuron soma
and stem axon. Arrow in merge indicates the stem axon of the DRG neuron. Apparent lipofuscin
autofluorescence is labeled in merge. Image is from donor #1. Scale bar is 50 μm. **D,** Z-
projection of anti-Kv2.1 (magenta), anti-Kv2.2 (green) (right) and anti-NF200 (left)
immunofluorescence in a human DRG neuron soma and stem axon. Arrows in merge indicate
the stem axon of the DRG neuron. Apparent lipofuscin autofluorescence is labeled in merge.
Image is from donor #3. Scale bar is 50 μm. Detailed information on each donor can be found in
the *Human Tissue Collection* section of the methods.
1267
1268
1269

1270 **Supplemental Figure 15**

1271
Supplemental Figure 15 legend
A, Z-projection of anti-Kv2.2 (left) and anti-NF200 (middle) immunofluorescence of human DRG.
Arrows in merge represent exemplar axons which have clear anti-Kv2.2 immunofluorescence.
Image is from donor #1. Scale bar is 50 μm. ***B***, Z-projection of anti-Kv2.2 (upper left), anti-CASPR
(upper right) and anti-Kv1.2 (bottom left) immunofluorescence of human DRG axon. Image is
from donor #2. Scale bar is 10 μm. Detailed information on each donor can be found in the
Human Tissue Collection section of the methods.
1279
1280

1281 **TABLE 1**

Figure	Genotype	Number of Mice (N)	Sections/plates Analyzed	ROIs Analyzed (n)	Age (Weeks)	Sex	Level of spinal column	Total ROIs analyzed
1	WT	1	1	124	11	M	L	124
2	WT	5	3,4,10,8,4	52,172,476,437,134,	16,7,7,7,7	M,M,F,M,M	L,T12,T12,T12,T12	1271
	Kv2.1 KO	5	4,7,5,4,5	232,124,576,293,239,	16,7,7,7,7	M,M,F,M,M	L,T12,T12,T12,T12	1124
3	WT	8	4,11,4,6,4,4,4,6	238,126,1227,673,706,748,602,572,	50,50,50,7,7,24,7,28	M,M,M,M,M,M,M,F	L,T12,T12,T12,T12,T12,L1,T12	4892
	Kv2.2 KO	8	3,6,7,4,4,4,4,5	359,209,853,400,913,719,729,745,	50,50,50,7,7,24,7,28	M,M,M,M,M,M,M,F	L,T12,T12,T12,T12,T12,L1,T12	4927
4	WT	10	7,5,6,9,8	444,268,167,669,600,	50,50,50,50,50	F,F,F,M,M	T12,T12,T12,T12,T12	3419
	Kv2.1 KO	7	5,6	298,324,	50,50	M,M	T12,T12	622
	WT	8	same as data in figure 3					
	Kv2.2 KO	8	same as data in figure 3					
5	WT	3	4,16,17,	104,379,328,	18,19,19	F,F,F	L,L,L	811
	Kv2.1 Kv2.2 DKO	2	5,14,	108,274,	18,19	F,F	L,L	382
6	WT DRG	5	1,1,1,1,1	139,51,30,40,38	10,10,10,8,8	M,M,F,F,M	T12,T8,L1,L1,L1	298
	WT Ventral Horn	5	1,1,1,1,1	75,51,23,24,28	10,10,10,8,8	M,M,F,F,M	T12,T8,L1,L1,L1	201
7	WT	2	2	N/A	7,24	M,M	L1,T12	N/A
8	WT	1	1	N/A	7	M	L1	N/A
	Kv2.2 KO	1	1	N/A	7	M	L1	N/A
	MrgprD-GFP	1	1	N/A	13	F	T12	N/A
9	WT	1	1	N/A	28	F	T12	N/A
	Kv2.2 KO	1	1	N/A	28	F	T12	N/A
10	WT	1	1	N/A	10	M	L5	N/A
	MrgprD-GFP	4	6,5,9,14	852,632,842,1392,	13,20,8,8	F,M,F,F	T12,L1,L1,L1	3718
	WT	5	7,4,6,7,4	482,464,592,641,166,	7,10,24,19,19	F,M,M,F,F	L1,T12,T12,T12,T12	2345
	PV-tdTomato	2	15,9	1978,1158,	8,8	F,M	L1,L1	3136
1-1	WT	1	1	N/A	10	M	L5	N/A
1-2	WT	1	1	N/A	28	M	L	N/A

1-3	WT	1	3	125	13	M	L	125
2-1	WT	1	33	5399	8	M	T11 and T12	5399
2-2	MrgprDGFP	1	1	N/A	13	F	T12	N/A
2-3	MrgprDGFP	4	6,15,9,14	5,568,769,051,487	14,20,8,8	F,M,F,F	L1,L1,L1,L1	3824
	WT	2	7,5	771,478	24,8	M,F	L1,L1	1249
6-1	WT DRG	4	1,1,1,1	25,30,28,35	7,7,8	M,F,F,F	L1,L1,L1,L1	118
	WT Ventral Horn	4	1,1,1,1	17,19,26,19	7,7,8	M,F,F,F	L1,L1,L1,L1	81
7-1	MrgprDGFP	1	1	N/A	14	F	T12	N/A
	WT	2	1,1	N/A	11,17	M,M	L,L	N/A
8-1	WT	1	1	N/A	7	F	T12	N/A
	Kv2.1 KO	1	1	N/A	7	F	T12	N/A

1282

Table 1 legend

Detailed information on mice used in each figure. In columns "Sections/Plates Analyzed", "ROIs Analyzed", "Age", "Sex" and "Level of Spinal Column" information for each mouse is separated by commas. "ROIs Analyzed" corresponds to ROIs generated by automated analysis. In column "Level of Spinal Column" L1 refers to the first lumbar DRG and T12 refers to the twelfth thoracic DRG and if only L is shown the specific DRG is not known but the DRG is from the lumbar region.

1291 **TABLE 2**

Antigen	Antibody name	Species/isotype	Manufaturer information	Concentration used	RRID	Figures
Kv2.1	K89/34	Mouse IgG1 mAb	NeuroMab	6.7 ug/mL	AB_10672253	1, 2, 3, 4,6-1, 7, 8-1, 10, 11, 11-1, 11-2
Kv2.2	Kv2.2c	Rabbit pAb	In-house Trimmer laboratory	1:100 purified from strip assay	AB_2801484	3, 4, 6-1, 7, 8, 9, 10, 11, 11-1, 11-2,11-3
Nav1.8	N134/12	Mouse IgG2a mAb	NeuroMab	1:5 tissue culture supernatant	AB_10672261	1,11-1
Kv2.1	K89/34R	Mouse IgG2a mAb	In-house Trimmer laboratory	1:5 tissue culture supernatant	AB_2315768	11-1
beta III Tubulin	NA	Rabbit pAb	Abcam ab18207	0.33 ug/mL	AB_444319	2, 8-1
Kv2.2	N372B/60	Mouse IgG2b mAb	NeuroMab	1 ug/mL	AB_2315868	1, 1-3, 6, 10
NF200	NA	chicken pAb	Abcam ab4680	0.2 ug/mL	AB_304560	3, 8, 10
Caspr	K65/35	Mouse IgG1 mAb	NeuroMab	1:5 tissue culture supernatant	AB_2877274	9
Kv1.2	K14/16	Mouse IgG2b mAb	NeuroMab	1:5 tissue culture supernatant	AB_2877295	9,11
CGRP	NA	Rabbit pAb	Immunostar	1:1000	AB_572217	10
DsRed	NA	Rabbit pAb	Takara Cat#632543	0.17 ug/mL	AB_2307319	10
Nav1.7	N68/6	Mouse IgG1 mAb	NeuroMab	1:5 tissue culture supernatant	AB_2184355	11-1,11-3

1292

1293

1294 **Table 2 legend**

1295 Detailed information on antibodies used throughout the manuscript. In column "Concentration

1296 Used" if only dilution is given concentration was unknown however, tissue culture supernatant

1297 concentrations of antibodies typically range between 15-30 µg/mL.