Ecology

Statistical Reports

Title: Mechanistic home range capture–recapture models for the estimation of population

density and landscape connectivity

Keita Fukasawa[1*] and Daishi Higashide[2]

1 Biodiversity Division, National Institute for Environmental Studies

16-2 Onogawa, Tsukuba, Ibaraki 305-8506, Japan

k.fukasawa37@gmail.com

2 Faculty of Bioresources and Environmental Sciences, Ishikawa Prefectural University

1-308 Suematsu, Nonoichi, Ishikawa 921-8836 Japan

dhigashide@gmail.com

* Corresponding author

Abstract

Spatial capture–recapture models (SCRs) provide an integrative statistical tool for analyzing animal movement and population patterns. Although incorporating home range formation with a theoretical basis of animal movement into SCRs can improve the prediction of animal space use in a heterogeneous landscape, this approach is challenging owing to the sparseness of recapture events.

In this study, we developed an advection–diffusion capture–recapture model (ADCR), which is an extension of SCRs incorporating home range formation with advection–diffusion formalism, providing a new framework to estimate population density and landscape permeability. we tested the unbiasedness of the estimator using simulated capture–recapture data generated by a step selection function. We also compared accuracy of population density estimates and home range shapes with those from SCR incorporating the least-cost path and basic SCR. In addition, ADCR was applied to real dataset of Asiatic black bear (*Ursus thibetanus*) in Japan to demonstrate the capacity of the ADCR to detect geographical barriers that constrain animal movements.

Population density and permeability of ADCR were substantially unbiased for simulated datasets. ADCR could detect envitonmental signals on connectivity more sensitively and could estimate population density, home range shapes and size variations better than the existing models. For the application to the bear dataset, ADCR could detect the effect of water body as a barrier of movement which is consistent with previous studies, whereas estimates by SCR with the least-cost path was difficult to interplet.

ADCR provides unique opportunities to elucidate both individual- and

population-level ecological processes from capture–recapture data. By offering a formal

link with step selection functions to estimate animal movement, it is suitable for

simultaneously modeling with capture–recapture data and animal movement data. This

study provides a basis for studies of the interplay between animal movement processes

and population patterns.

Key words: spatial capture–recapture, circuit theory, landscape resistance, abundance

estimation, mechanistic home range analysis, partial differential equation, step selection

function, advection–diffusion equation

Introduction

Anthropogenic activity leads to habitat loss and fragmentation, posing a major threat to

biodiversity. In a fragmented landscape, connectivity between habitat patches affects

population patterns and sustainability (Hanski et al. 1994). Functional connectivity in a

heterogeneous landscape is not only of general interest in basic ecology but also a key

factor affecting decision-making in wildlife management and conservation (Goodwin

and Fahrig 2002, Royle et al. 2018, Osada et al. 2019). Circuit theory and least-cost path

modeling are two major paradigms for evaluating landscape connectivity, with different

underlying behavioral assumptions and statistical properties (McRae et al. 2008,

Marrotte and Bowman 2017). Permeability, which is a measure of connectivity based on

circuit theory has explicit links with animal movement models, such as random walks and

step selection functions (McRae 2006, Duchesne et al. 2015), and models of animal gene

flow and movement have numerous applications (Dickson et al. 2019). Empirical

determination of connectivity parameters, such as effects of land cover types on

permeability, is important to improve objectivity, and the development of estimators is a

major challenge in ecology (Ovaskainen 2004, Ovaskainen et al. 2008, Beyer et al. 2016,

Signer et al. 2017, Osada et al. 2019).

Spatial capture–recapture models (SCRs) are powerful tools with the capacity

to estimate individual movement processes and population abundance in an integrative

manner (McClintock et al. 2022). An extention of SCR incorporating the least-cost path

distance in the detection function instead of Euclidian distance (Royle et al. 2013,

Sutherland et al. 2015, Fuller et al. 2016) can be applicable to estimate weights of

landscape factors on the cost path distance. Permeability has favorable properties for

practical applications, such as robustness to the resolution of landscape data (Marrotte

and Bowman 2017) and a straightforward link with movement models (Diniz et al.

2020). In particular, the process of home range formation considering permeability and

site fidelity has been studied extensively by step selection functions in movement ecology

(Thurfjell et al. 2014, Duchesne et al. 2015, Signer et al. 2017, Péron 2019, Potts and

Schlägel 2020). However, circuit theory-based SCRs are not as well-studied as SCRs

incorporating the least-cost path. Although circuit theory-based SCRs can provide

insight into the relationship between individual-level displacement and population-level

abundance patterns, the temporal sparseness of capture–recapture data limits the

application of SCR to complex movement processes (McClintock et al. 2022).

Incorporating mechanistic models of home range formation into SCR can enable

inferences about permeability as well as population abundance from capture–recapture

data in a computationally tractable manner. Mechanistic home range analysis (Moorcroft

and Lewis 2006, Moorcroft and Barnett 2008) is an analytical approach that ties the

process of animal movements to the resultant home range shape at an equilibrium state.

The approach has contributed to the development of advection–diffusion equations to

obtain flexible home ranges and likelihood-based modeling to estimate parameters

empirically. The advection–diffusion equations describe deterministic drift to a focal

point (e.g., an activity center) as advection terms and random walks as diffusion terms.

Circuit theory is analogous to a diffusion process with heterogeneous diffusivity (Hanks

and Hooten 2013, Yamaura et al. 2022), and the advection–diffusion equations with

heterogeneous diffusion coefficients in space are applicable to home ranges formed by

animal movement affected by barriers in a landscape. Although closed-form analytical

solutions of advection–diffusion equations rarely exist, computationally efficient ways

to obtain an equilibrium solution have been developed (Hirsch 2007). By integrating

mechanistic home range analysis into SCR, the simultaneous estimation of permeability

and population density is possible, without substantial computational costs. Hereafter,

we refer to this new SCR approach as the advection–diffusion capture–recapture model

(ADCR).

Although ADCRs and SCRs with the least-cost path (Royle et al., 2013;

Sutherland et al., 2015; Royle et al., 2018) differ with respect to connectivity

assumptions and cannot be directly compared, evaluations of the performance of

estimators are helpful for users. Applying statistical models to simulated data generated

from a mechanistic animal movement model is a straightforward way to test the

unbiasedness of estimators and robustness to deviation from assumptions (Dupont et al.

2022, Theng et al. 2022). Also, an application to real dataset is important to illustrate

how statistical models can yield realistic estimates even in situations where there is

unknown heterogeneity in the data generating process.

The aim of this study was to develop a formal framework for ADCR and to

compare the accuracy of the estimated abundance and home range shape with those

obtained by SCRs with the least-cost path. The identification of the optimal model is

outside of the scope of this paper. Rather, our aim is to establish the merit of the ADCR

under animal movement modeled by a step selection function, which is common in

movement ecology. We also compared density estimates with those of basic SCR

(Borchers and Efford 2008) as a null model of heterogeneous connectivity. We predict

that all three methods, ADCR, an SCR with the least-cost path, and basic SCR, will

accurately estimate population densities because SCR is robust to the violation of home

range assumptions (Efford 2019). We also predict ADCR will have higher predictive

ability of home range shapes than SCR with the least-cost path and basic SCR when

animal movement is driven by exploratory behavior in landscapes with heterogeneous

permeability. As an example of model application, we applied ADCR to estimate factors

affecting peameability for Asiatic black bear (*Ursus thibetanus*) in Japan.

Material and Methods

Model Framework

ADCR is an extension of SCR in which animal home ranges are modeled by

equilibrium solutions of the advection–diffusion equation, an Eulerian representation of

stochastic animal movement in a heterogeneous connectivity landscape. By integrating

the advection–diffusion equation with SCR, it is possible to simultaneously estimate

population density while inferring the effect of landscape connectivity on the skewed

shape of the home range, using capture–recapture data. Here, we describe the model

structure of ADCR.

Detection model and population process

In stationary SCR modeling, a detection model that describes the encounter of

an individual to a detector reflects the home range behavior of the animal. The home

range of individual is defined by its utilization distribution, which is a bivariate

probability distribution of an individual location in a space. In general, the model of

utilization distribution has latent variables of an activity center $\boldsymbol{\mu} = (\mu_x, \mu_y)$, where an

individual is attracted. For example, a basic SCR (Borchers and Efford 2008, Royle et al.

2009) often assumes a half-normal detection function, which implies that the utilization

distribution is a stationary bivariate normal distribution centered to the activity center.

To introduce the ADCR, which covers a more flexible class of utility distributions under

heterogeneous permeability landscapes, we adopted a detection model that describes an

explicit link between the encounter frequency of an individual to a detector and the

utilization distribution $p^*(\mathbf{s} \mid \boldsymbol{\mu})$, which defines the probability density at location $\mathbf{s} = (x,$

$y)$ of an individual with an activity center $\boldsymbol{\mu}$. Note that the integration of $p^*(\mathbf{s} \mid \boldsymbol{\mu})$ over $\mathbf{s}$

equals 1 because of the nature of probability distribution.

An SCR survey is conducted using an array of detectors in space. Researchers

may either iterate the survey on more than one occasion or treat the entire survey period

as a single occasion. When the SCR data are collected by count detectors, the encounter

frequency of the ith individual to the jth detector on the kth occasion is written as

$$\lambda_{ijk} = \exp(g_0)p^*(\mathbf{s}_j \mid \boldsymbol{\mu}_i), \qquad \text{(eqn. 1)}$$

where g_0 is a parameter of log detectability, $\boldsymbol{\mu}_i$ is the activity center of the ith individual,

and $\mathbf{s}_j$ is the location of the jth detector. The corresponding observed value, Y_{ijk}, follows

a probability distribution depending on λ_{ijk}, denoted by $p(Y_{ijk} \mid \boldsymbol{\mu}_i, \boldsymbol{\theta})$, where $\boldsymbol{\theta}$ is a vector

of parameters that affects λ_{ijk}, including g_0 and the parameters controlling p^*. The

number of detections follows a Poisson distribution $p(Y_{ijk} \mid \mu_i, \theta) = \text{Poisson}(Y_{ijk} \mid \lambda_{ijk})$ if

the conditional independence among detection events given λ_{ijk} is assumed. Although

we implemented an observation model for count detectors in this study, it can be easily

extended to a model of varying effort with various detector types (e.g., binary proximity

and multi-catch trap) as well as the basic SCR (Efford et al. 2013).

171   The population process of SCR describes the spatial pattern of activity centers

following a point process in a region $\mathcal{S}$ (Borchers and Efford 2008, Royle et al. 2009). $\mathcal{S}$

is often defined as a rectangular region that includes an array of detectors. In this study,

we applied a homogeneous Poisson process, the simplest form of the point process,

assuming that μ follows a uniform distribution over $\mathcal{S}$, $\mu \sim \text{Uniform}(\mathcal{S})$. Let ρ denote the

log population density, and the number of activity centers in an arbitrary region with

area A follows a Poisson distribution with expectation $\exp(\rho)A$. As is the case with the

basic SCRs (Borchers and Efford 2008), the model can be extended to an

inhomogeneous Poisson process that can accommodate a situation in which ρ varies in

space depending on the landscape covariates.

Home range formation

182   In ADCR, the utilization distribution is derived from a fine-scale movement

behavior that forms a stationary home range in a heterogeneous permeability landscape.

The Ornstein–Uhlenbeck process (Smouse et al. 2010) is a basic movement model

describing the home range behavior of an animal that is subject to drift toward the

activity center and a random walk. According to the circuit theory, permeability

corresponds to the step length of a random walk (McRae et al. 2008), and we considered

a situation in which permeability depends on landscape covariates. The

Ornstein–Uhlenbeck process converges to a probability distribution of individual

location at the equilibrium state (Smouse et al. 2010), and we applied it to utilization

distribution in ADCR.

To derive the probability distribution of an individual location, we introduced

an Eulerian approach that focuses on the flow of probability density in a space driven by

drift and random walk. Under the Ornstein–Uhlenbeck process, the probability

distribution of an individual location follows an advection-diffusion equation that

describes the evolution of the probability density over time in space (Méndez et al. 2013).

In our study, we consider a situation that the diffusion coefficient vary in space

(Ovaskainen 2004, Ovaskainen et al. 2008) as follows:

$$\frac{\partial p}{\partial t} = -c\left((\mu_x - x)\frac{\partial p}{\partial x} + (\mu_y - y)\frac{\partial p}{\partial y}\right) + D(x, y)\left(\frac{\partial^2 p}{\partial x^2} + \frac{\partial^2 p}{\partial y^2}\right),$$ (eqn. 2)

where $D(x, y) > 0$ is the spatially-varying diffusion coefficient and c is the intensity of

drift. The first term on the right-hand side is advection corresponding to drift toward the

activity center, and the second term is diffusion. The diffusion coefficient can be

interpreted as mean step length of diffusive movement per unit time (Potts and Schlägel

2020) meaning how easy an animal can go through the location. In terms of circuit

theory, it is equivalent to permeability or conductance (McRae 2006, McRae et al. 2008).

The intensity of drift is the strength of site fidelity; the large value indicates that an

animal is attracted toward the activity center strongly, resulting in small home range size.

To model the effect of landscape variables on permeability, we applied the log-linear

mode $\ln(D(x, y)) = \beta_0 + \boldsymbol{\beta}\mathbf{z}(x, y)$, where β_0 is the intercept, $\mathbf{z}(x, y)$ is a vector of the

landscape covariates at location (x, y), and $\boldsymbol{\beta}$ is the corresponding regression coefficients

that determine the dependence of permeability to landscape covariates. The intensity of

drift, c, takes a positive value, and $\ln(c)$ is denoted by the parameter α_0. To ensure that

the total mass of probability does not change with time, the reflecting boundary condition

is given for the space. Eqn. 2 describes the time evolution of the probability density of an

individual location, and we applied it to project the animal movement paths for the data

simulation (see **Simulation Study**).

Similar to mechanistic home range analysis (Moorcroft and Lewis 2006), the

home range in this system is defined as the equilibrium probability density distribution

$p^*(\mathbf{s} \mid \boldsymbol{\mu})$, which emerges when advection toward activity center and diffusion balances.

The $p^*(\mathbf{s} \mid \boldsymbol{\mu})$ is obtained by solving the following system:

$$-\left(\left(\mu_x - x\right)\frac{\partial p}{\partial x} + \left(\mu_y - y\right)\frac{\partial p}{\partial y} \right) + \frac{D(x,y)}{c}\left(\frac{\partial^2 p}{\partial x^2} + \frac{\partial^2 p}{\partial y^2}\right) = 0. \quad \text{(eqn.3)}$$

To make sure that $p^*(\mathbf{s} \mid \boldsymbol{\mu})$ is the probability distribution in the region $\mathcal{S}$, we imposed a

constraint of $\int_{\mathcal{S}} p \, d\mathbf{s} = 1$. Note that the parameters α_0 and β_0 are redundant under the

equilibrium assumption because $\ln(D(x, y)/c) = \beta_0 - \alpha_0 + \boldsymbol{\beta}\mathbf{z}(x, y)$. To ensure the

identifiability of the model, it was reparametrized as $\gamma_0 = \beta_0 - \alpha_0$. γ_0 is interpreted as the

logarithm of the relative intensity of diffusion and drift, which determines the average

home range size.

Although eqns. 2 and 3 imply that the landscape covariates are spatially

continuous, they are actually given by a discrete dataset in raster or vector format.

Accordingly, these equations can be numerically solved using a variety of discretization

techniques, including the finite difference method and finite element method (Hirsch

2007). In this study, the finite difference method was applied because it is suitable for

common situations in which landscape factors affecting the diffusion coefficient are

available as a raster dataset. In addition, grid cells of raster data were used to discretize

space for simplicity. Technical details for discretization and numerical derivation of $p^*(\mathbf{s} \mid$

$\boldsymbol{\mu}$) are shown in **Appendix S1**. Importantly, our home range specification is a

generalization of basic SCRs that assume a bivariate normal home range (Borchers and

Efford 2008, Royle et al. 2009), given that the solution of eqn. 3 when D(x, y) is

constant over space (i.e., $\boldsymbol{\beta} = \mathbf{0}$) is a bivariate normal distribution whose variance is

$\exp(\gamma_0)$ (Méndez et al. 2013). An example of derived home ranges with different βs in a

hypothetical landscape with a binary covariate for permeability is shown in Fig. 1,

which shows that the larger the effect of a landscape covariate, the more the home range

shape skews toward the habitat patch with high permeability.

Model estimation

We applied the maximum likelihood method to estimate the permeability

parameters and population density simultaneously from the capture–recapture data.

Although conditional likelihood maximization can also be used for parameter

estimation (Borchers and Efford 2008), full likelihood maximization was applied in this

study because it has the capacity to extend to more complex situations in which

population density varies across space. In this study, we applied a quasi-Newton

algorithm (Nocedal and Wright 2006) for maximizing log likelihood function. Technical

details in the model estimation was shown in **Appendix S2**.

Simulation study

Animal movement is temporally auto-correlated, and hence so are the locations of

detections. To test the capacity of ADCR to estimate parameters of connectivity and

population density correctly under a realistic movement process, we used a dataset

generated via Monte Carlo simulations of animal movement. The landscape is a

rectangular grid system of 50 × 50 (hence, the area is 2500), and animal location was

described as a discrete coordinate of grid cells. Arrays of count detectors were set to the

range, $x = [9, 41]$ and $y = [9, 41]$, with x and y axis spacing of 4 and 81 detectors. This

simulation setting mimics a standard spatial capture recapture survey with a buffer

outside the detector array to reduce boundary effects and with the distance between

detectors so that an inidividual is detected by multiple detectors. The spatial resolution

of the landscape grid was set coarsely to make the many iterative calculations in the

simulation tractable.

Simulations were iterated 100 times, and true parameter values and landscape

structure were randomly determined in each iteration. For each iteration, a

heterogeneous landscape was generated using the R package rflsgen (Justeau-Allaire et

al. 2022). Random patches were generated in the landscape by rflsgen with parameters

shown in Table S1. For covariate values, 0.5 and –0.5 were assigned to patch and

non-patch habitats, respectively. True parameter values were sampled from uniform

distributions for each iteration. The ranges of true parameters were set as log population

density $\rho = (-3, -1)$, log intensity of drift $\alpha_0 = (-1, 0)$ and coefficient of landscape

variable $\beta = (-2, 2)$. g_0 and β_0 were fixed to -3 and 1, respectively. At the beginning of

an iteration, the number of individuals in the landscape was determined by a Poisson

distribution with expected value of $\exp(\rho) \times 2500$ and activity centers were drawn from

a discrete uniform distribution. The mean population size was 367 (range: 112, 956).

The animal movement simulation was run for $t = 5000$ after 100 burn-in steps

for randomizing initial locations of individuals. For each time step, we calculated the

probability vector of individual locations for all cells given the location at the previous

step using the advection–diffusion equation with the explicit method (eqn. 4 in

Appendix S1), and the location was sampled from a categorical distribution. When there

was a detector in a cell in which an individual was located, the number of detection

events was simulated by a Poisson distribution with expectation $\exp(g_0)$. Then,

single-occasion capture–recapture data were obtained by censoring undetected

individuals. An example map of simulated movement and observations of an individual

is shown in Fig. 2. The mean number of individuals detected at least once was 274

(range: 77, 682) with the detection rate (= number of detected individuals at least once /

population size) of 0.745 (range: 0.631, 0.857). Mean number of detections was 2997

(range: 865, 7963). This simulation setting might generate a dataset with more

detections than a real capture-recapture survey, but it will allow us to detect the bias of

each estimator more sensitively.

We applied ADCR, SCR with the least-cost path, and basic SCR to the

simulated dataset to evaluate the accuracy of estimations. The basic SCR assumes that

detectability function is Gaussian function of the Euclidian distance from the activity

center $\ln(\lambda_{ijk}) = g_0 + g_1\|\mathbf{s}_j - \boldsymbol{\mu}_i\|^2$, implying the home range shape is two-dimentional

normal distribution. The formulation of SCR with the least-cost path is similar to basic

SCR, but the Euclidian distance is replaced by the least-cost path distance from the

activity center ($d_{lcp}(\mathbf{s}_j,\boldsymbol{\mu}_i; g_2)$), $\ln(\lambda_{ijk}) = g_0 + g_1 d_{lcp}(\mathbf{s}_j,\boldsymbol{\mu}_i; g_2)$. The log unit cost that an

individual move from a grid cell (**s**) to the adjacent cell (**s'**) is $g_2(z(\mathbf{s}) + z(\mathbf{s}')) / 2$, and

parameter g_2 determines the weight of a landscape covariate on cost (Royle et al. 2013,

Sutherland et al. 2015). Note that positive value of g_2 indicates the negative relationship

between the landscape connectivity and the landscape covariate, opposite to positive **β**

value of ADCR.

For the three estimators, the deviation of population density estimates from true

values was determined and root mean square error (RMSE) and mean bias error (MBE)

were calculated. To show how well the three estimators reproduce home range shapes,

the agreement between true (p^*_{true}) and predicted (p^*_{pred}) probability distributions of

individual locations was calculated by an overlapping coefficient defined by the volume

of overlapping region between two probability distributions (Inman and Bradley 1989).

The overlapping coefficient ranges from 0 (no overlap) to 1 (perfect overlap). To

evaluate the average performance over locations for activity center $\boldsymbol{\mu}$, we calculated the

overlapping coefficients averaged over $\boldsymbol{\mu}$ within the range of trap array x = [9, 41] and y

= [9, 41] for each model and iteration. As an indicator of home range size, we also

calculated 90% highest density areas of p^* whose $\boldsymbol{\mu}$s are within the range of trap array

and compared the means and SDs with the true values.

In order to demonstrate how well ADCR works under severe sparseness of

observations and biased detector deployments to a specific landscape element, we

conducted supplemental simulations with the following settings; true value of

detectability was set to $g_0 = -5$, and all the detectors in the landscape elements with low

permeability (i.e. covariate value = -0.5) were removed from the detector array of the

baseline simulation (range of detector quantities: 19 to 50). Other true parameters were

chosen within the range of the baseline simulations: log population density $\rho = -2$, log

intensity of drift $\alpha_0 = -0.5$, coefficient of landscape variable $\beta = 2$ and intercept of

diffusion coefficient $\beta_0 = 1$. Other settings, such as landscape structure, resolution,

number of time steps were identical to the baseline simulations. The number of

independent iterations was 100. The range of total population size was 296 to 386

(mean: 337). The number of individuals detected at least once ranged from 60 to 139

(mean: 96.8), and the detection rate was from 0.175 to 0.409 (mean: 0.287). The number

of detections ranged from 90 to 230 (mean: 161).

All models were estimated using maximum likelihood methods with R 4.1.3 (R

Core Team 2022). The movement simulation and log likelihood function of ADCR were

written largely in R language, with some subroutines in C++. The source code is

available online (https://github.com/kfukasawa37/adcrtest2). The log likelihood values

of SCR with the least-cost path were derived based on the R script by Sutherland et al.

(2015), with some modifications by Keita Fukasawa to deal with count detectors and

the negative effect of landscape on the cost function. The basic SCR was estimated

using the R package secr (Efford, 2022).

Case study

We used a capture-recapture dataset of Asiatic black bear population in eastern

Toyama prefecture, Japan, collected over three years (2013-2015) by authors. The data

was obtained using video-recording camera traps at 86 locations with a field design for

efficient photographing of chest marks (Higashide et al. 2013) and photograph-based

individual recognitions (Higashide et al. 2012). Note that all the camera locations were

in forest due to limitations of field implementations. Years were treated as occasions.

The dataset was constituted of 271 detections for 109 individuals with the locations and

year. For the details of the capture-recapture survey, see **Appendix S3**.

In Japan, human-bear conflicts are increasing (Sakurai and Jacobson 2011).

Understanding population status (Horino and Miura 2000) and space use (Ohnishi et al.

2019) of bears are crucial for developing plans to manage bear population and their

threat to human well-being. Population and landscape genetic studies suggested that

rivers (Saitoh et al. 2001, Ohnishi et al. 2007) and agricultural lands (Takahata et al.

2014, Ohnishi et al. 2019). To test the effect of rivers and agricultural lands as barriers

or corridors for black bears, we compiled 0.5km grid data of area ratio covered by water

surface and agricultural lands (Fig. S1). These areas were derived from the national

vegetation map created by Biodiversity Center of Japan, Ministry of Environment

(https://www.biodic.go.jp/index_e.html, accessed 24 May 2024) released in 2010s. We

estimated ADCR and SCR with the least-cost path including the two landscape

covariates and basic SCR. We compared the estimated log population density by the

three methods, as well as the comparison of the estimated effect of landscape by ADCR

and SCR with the least-cost path.

363   **Results**

For all iterations, estimations of ADCR converged successfully. All the estimates of

ADCR, including the population density and effect of landscape on connectivity, were

substantially unbiased. The true and estimated parameters were on the line y = x, on

average (Fig. 3). RMSE and MBE of g_0 were 0.0258 and 0.0109, respectively. The true

and predicted home ranges overlapped fairly well. The overlapping coefficients

marginalized over activity centers were (0.950, 0.9996) over 100 iterations, and the

mean was 0.988 (Fig. 4).

371   As predicted, population densities estimated by SCR with the least-cost path

and basic SCR were robust to true connectivity settings in our simulation experiments.

RMSEs and bias of log population density estimated by SCR with the least-cost path

and basic SCR were very similar to those of ADCR and were not sensitive to the true

effect of landscape on connectivity (Table 1). The coefficient of landscape estimated by

SCR with the least-cost path were linear with the true parameters of connectivity, on

average, but showed a slightly weaker relationship (Pearson's $r = -0.988$, Fig. S2) than

that of ADCR (Pearson's $r = 0.995$, Fig. 3). The estimates of γ_0 by ADCR were also

linear with the true values (Pearson's $r = 0.991$, Fig. 3). Although the mean home range

sizes measured by 90% highest density area were almost unbiased for the three

estimators (Fig. S3a), the heterogeneity in home range sizes were largely

underestimated by SCR with the least-cost path and basic SCR (Fig. S3b); MBE in

standard deviation of home range sizes were -5.38 for SCR with the least-cost path,

-15.3 for basic SCR and -0.0508 for ADCR.The overlapping coefficients of home range

predicted by SCR with the least-cost path (mean: 0.904; variance: 9.93×10^{-5}) and basic

SCR (mean: 0.883; variance: 2.23×10^{-3}) were lower than those for ADCR (mean: 0.998;

variance: 7.52×10^{-5}) over the range of true connectivity parameters. The predictive

performance of SCR with the least-cost path and basic SCR were decreased when the

connectivity strongly depends on landscape covariate, but SCR with the least-cost path

worked better than basic SCR (Fig. 4). For example of the range where the absolute

value of true landscape effect is [1.5, 2.0], the mean overlaps by SCR with the least-cost

path, basic SCR and ADCR were 0.892, 0.822 and 0.991, respectively. SCR with the

least-cost path tended to predict a more blurred and isotropic home range shape than the

true home range shape (Fig. S4).

Under the sceanio of severe data sparseness and biased detector alignment,

estimates of ADCR had large variances but were substantially unbiased; the mean ±

standard deviation of estimated log population density, γ_0 and effect of landscape were

-2.00±0.142, 1.41±0.425 and 2.12±1.10, respectively. In this scenario, accuracy of

density estimates by ADCR was much better than the other estimators (Fig. S5); the

MBEs of population density estimates by ADCR, SCR with the least cost path and basic

SCR were 0.00308, 0.0657 and -0.0373, respectively. ADCR could detect the correct

environmental signals on connectivity clearer than SCR with the least-cost path; the

95% confidence intervals (CIs) did not overlap 0 for 60 out of 100 iterations which is

outperformed SCR with the least-cost path (54 out of 100 iterations). Although the sign

(i.e. positive or negative) of landscape effect on connectivity was incorrect for two

iterations of ADCR and five iterations of SCR with the least-cost path out of 100

iterations, all of them were not significant with their 95% CI overlapping 0 (Table S2).

There were two iterations that SCR with the least-cost path estimated incorrect signs of

effects and ADCR could detect significant effects correctly.

For the applications on the black bear dataset, there were unexpected

discrepancies in the estimated effects of landscape factors on connectivity between

ADCR and SCR with the least-cost path (Fig. 5). The estimated coefficients (i.e.

parameter β) of agricultural land and water surface on the permeability of black bears

were 0.361 (SE: 0.113; 95% CI: 0.175, 0.547) and -6.41 (SE: 1.26; 95% CI -8.88, -3.94),

respectively. Inversely, the estimated effect of water surface on the cost by SCR with the

least-cost path (parameter g_2) was -25.5 (SE: 5.13; 95% CI: -35.6, -15.4), indicating

water surface was corridor for black bears (Fig. 5). Note that deletion of agricultural

land from the covariate did not affect the effect of water surface so much (-23.1, SE:

4.57). The effect of agricultural land by SCR with the least-cost path was not clear; the

maximum likelihood estimate was 0.0692 (SE: 0.0360; 95% CI: -0.00130, 0.140) (Fig.

5). Contrary to the predicted home ranges by ADCR showing skewed shapes avoiding

water surface (Fig. S6a), the home ranges predicted by SCR with the least-cost path

showed "leakage" of probability density into water surface (Fig. S6b). The log

population density estimates were similar among the methods but ADCR was the

highest: -1.45 (SE: 0.113; 95% CI: -1.67, -1.23) by ADCR, -1.56 (SE: 0.0996; 95% CI:

-1.76, -1.37) by SCR with the least-cost path and -1.59 (SE: 0.105; 95% CI: -1.79,

-1.37) by basic SCR.

428   **Discussion**

Individual movement is an important driver of spatial patterns of populations, and

incorporating movement processes in spatial capture–recapture analysis will facilitate

studies of the interplay between individual- and population-level ecological processes

(Royle et al. 2018, McClintock et al. 2022). In this study, we developed a formal

framework for a SCR model including mechanistic home range formation with an

explicit theoretical link to animal movement models. Through individual-based

simulations of animal movement and detection, we showed ADCR accurately estimated

the population density, effect of landscape on permeability and heterogeneous home

range sizes. For the application of models to black bear dataset, ADCR could detect

biologically-plausible environmental signals on the connectivity. These results suggest

that ADCR is an effective approach for estimating population density and random

walk-based landscape connectivity.

441   Although the integration of a least-cost path modeling with spatial capture

recapture analysis (Royle et al. 2013, Sutherland et al. 2015) was a great methodological

advancement for the simultaneous estimation of population density and ecological

distance, the model performance under the animal movement following step selection

function has not been studied well. Our simulation study demonstrated that ADCR

outperformed the SCR with the least cost path in terms of accracy in population density

estimation, successful detection of environmental signals on connectivity (i.e.

permeability or cost) and precise prediction of home range shapes and the

location-specific heterogeneity in home range sizes. The ability of ADCR to estimate

the effects of landscape variable on animal movement correctly (Fig. 3) is result from

the formal link between microscopic animal movement and the macroscopic utility

distribution, corresponding to Lagrangian and Eulerian descriptions of a system.

Previous study showed that SCR with the least-cost path is difficult to estimate accurate

cost on movement without an integration with GPS logging data (Dupont et al. 2022).

This might be due to lack of formal link between home range model and the underlying

movement processes.

For the case study of Asiatic black bears, the results of ADCR supported the

previous studies demonstrated rivers as physical barriers (Saitoh et al. 2001, Ohnishi et

al. 2007) and farmlands as corridors (Takahata et al. 2014), whereas SCR with the

least-cost path showed the unexpected estimates indicating water surface as corridor for

bears. Although our simulation study with sparse data and biased detector alignment

could not fully reproduce such an extraordinary discrepancy between estimators, there

were some cases that only ADCR could detect correct environmental signals on

connectivity (Table S2). We considered the unexpected behavior of SCR with the

least-cost path was estimation bias caused by combination of biased layout of detectors

toward forest and the model assumption that detection rate at the activity center is

constant irrespective of the home range size. This assumption implys that observations

do not reflect the ubiquitous tradeoff between exploration of large area and exploitation

of resources near the activity center (exploration-exploitation tradeoff, Eliassen et al.

2007, Berger-Tal et al. 2014, Berger-Tal and Saltz 2014), leading model vulnerability in

estimating of animal space use to sampling non-randomness. ADCR explicitly models

home ranges as probability distributions with sum-to-one constraint, and seemed to be

more suitable for general home range behavior. However, it may not be able to fully

handle a variety of the diverse animal movement processes in response to landscape

barriers such as the correlated random walk (Patlak 1953, Wang and Potts 2017) and

randomized shortest paths (Panzacchi et al. 2016, Long 2019). Extending the current

study, which integrated the utility distribution derived from the stochastic process of

animal movement with spatial capture-recapture models, to a variety of movement

models will lead to solve the problem.

The effect of a skewed home range on the accuracy of density estimation by

SCR has been studied extensively in ecology (Ivan et al. 2013, Royle et al. 2013,

Sutherland et al. 2015, Efford, 2019, Theng et al. 2022). The results of this study were

consistent with those of recent studies (Efford 2019, Theng et al. 2022) showing that

robustness of density estimation to deviations in assumptions on home range shapes

depends on independence of sampling design to animal space use. In addition, our study

showed that estimation bias of density by SCR with the least-cost path can be as large as

basic SCR (Fig. S5). Efford (2019) emphasized the importance of sampling design in

space, such as the use of a two-dimensional detector array and avoidance of detector

array alignment along landscape features, for the correct estimation of density.

Obviously, such sampling designs are also favorable for successful detection of

landscape effects on home range shape by ADCR.

In landscape ecological analyses, effects of choice of spatial resolution on the

results of study can often be problematic (Turner et al. 1989, Qi and Wu 1996). We

found that ADCR was robust to decreased spatial resolution of a landscape variable by

supplementary simulatios (Appendix S4, Fig. S7). This robustness is a favorable

characteristic of connectivity measures in circuit theory, which shares the same

theoretical foundation of random walk descriptions of animal movement (Hanks and

Hooten 2013).

The broad application of SCR in ecology can largely be attributed to its

flexibility to incorporate heterogeneity in ecological processes underlying spatial

capture recapture data, such as environment-dependent abundance and sex-biased

detection (Efford, 2022). Future advances in ADCR should focus on similar

functionality. In particular, incorporating heterogeneity in home range size (Efford et al.,

2016) as well as heterogeneity in connectivity is a major challenge. Also, the

generalization of ADCR for dealing with temporally open populations driven by

birth-death processes (Gardner et al. 2010, Efford & Schofield 2022) and integration

with GPS logging data (Dupont et al. 2022) would be helpful for getting more out of

capture–recapture data. The development of software tools that enable users to benefit

from the flexible nature of ADCR is an urgent task for expanding the application of the

method.

Spatial capture–recapture is a popular method for abundance analyses in

ecology and conservation (Iijima 2020), and large amounts of data have accumulated.

Moreover, recent technical advances in individual identification from environmental

DNA (Dugal et al. 2022) and machine observations, such as camera traps (Schneider et

al. 2022), will expand the applicability of capture–recapture approaches to broad taxa.

Such capture–recapture data have the potential to advance our understanding of the

process of animal home range formation. ADCR will be a useful tool for integrated

analyses of capture recapture data incorporating individual- and population-level

ecological processes.

Acknowledgements

Yutaka Osada provided valuable comments on an earlier version of the statistical model.

We thank Elise Zipkin and two anonymous reviewers for constructive comments in the

peer-review process. This study was supported by JSPS KAKENHI Grant Numbers

JP13247736, JP19153418, and by the Environment Research and Technology

Development Fund (JPMEERF20194005) of the Environmental Restoration and

Conservation Agency Provided by the Ministry of Environment of Japan.

Author contributions

KF conceived the ideas, designed the methodology, conducted the simulation study,

conducted field survey, and applied the model to actual dataset. DH conducted field

survey, and complied the dataset. KF led the writing and all authors contributed to the

revision of the manuscript.

Conflict of interest

The authors have no conflict of interest to be declared.

References

Berger-Tal, O., J. Nathan, E. Meron, and D. Saltz. 2014. The exploration-exploitation

dilemma: a multidisciplinary framework. PLOS ONE 9:e95693.

Berger-Tal, O., and D. Saltz. 2014. Using the movement patterns of reintroduced animals

to improve reintroduction success. Current Zoology 60:515–526.

Beyer, H. L., E. Gurarie, L. Börger, M. Panzacchi, M. Basille, I. Herfindal, B. Van

Moorter, S. R. Lele, and J. Matthiopoulos. 2016. 'You shall not pass!':

quantifying barrier permeability and proximity avoidance by animals. Journal of

Animal Ecology 85:43–53.

Borchers, D. L., and M. G. Efford. 2008. Spatially explicit maximum likelihood methods

for capture–recapture studies. Biometrics 64:377–385.

Dickson, B. G., C. M. Albano, R. Anantharaman, P. Beier, J. Fargione, T. A. Graves, M. E.

Gray, K. R. Hall, J. J. Lawler, P. B. Leonard, C. E. Littlefield, M. L. McClure, J.

Novembre, C. A. Schloss, N. H. Schumaker, V. B. Shah, and D. M. Theobald.

2019. Circuit‐theory applications to connectivity science and conservation.

Conservation Biology 33:239–249.

Diniz, M. F., S. A. Cushman, R. B. Machado, and P. De Marco Júnior. 2020. Landscape

connectivity modeling from the perspective of animal dispersal. Landscape

Ecology 35:41–58.

Duchesne, T., D. Fortin, and L.-P. Rivest. 2015. Equivalence between step selection

functions and biased correlated random walks for statistical inference on animal

movement. PLOS ONE 10:e0122947.

Dugal, L., L. Thomas, M. R. Jensen, E. E. Sigsgaard, T. Simpson, S. Jarman, P. F.

Thomsen, and M. Meekan. 2022. Individual haplotyping of whale sharks from

seawater environmental DNA. Molecular Ecology Resources 22:56–65.

Dupont, G., D. W. Linden, and C. Sutherland. 2022. Improved inferences about landscape

connectivity from spatial capture–recapture by integration of a movement model.

Ecology:e03544.

Efford, M. G. 2019. Non-circular home ranges and the estimation of population density.

Ecology 100:e02580.

Efford, M. G. 2022. secr: Spatially explicit capture-recapture models.

https://CRAN.R-project.org/package=secr.

Efford, M. G., D. L. Borchers, and G. Mowat. 2013. Varying effort in capture–recapture

studies. Methods in Ecology and Evolution 4:629–636.

Efford, M. G., D. K. Dawson, Y. V. Jhala, and Q. Qureshi. 2016. Density-dependent

home-range size revealed by spatially explicit capture–recapture. Ecography

39:676–688.

Efford, M. G., and M. R. Schofield. 2022. A review of movement models in open

population capture–recapture. Methods in Ecology and Evolution 13:2106–2118.

Eliassen, S., C. Jørgensen, M. Mangel, and J. Giske. 2007. Exploration or exploitation:

574   life expectancy changes the value of learning in foraging strategies. Oikos

575   116:513–523.

576   Fuller, A. K., C. S. Sutherland, J. A. Royle, and M. P. Hare. 2016. Estimating population

577   density and connectivity of American mink using spatial capture–recapture.

578   Ecological Applications 26:1125–1135.

579   Gardner, B., J. Reppucci, M. Lucherini, and J. A. Royle. 2010. Spatially explicit inference

580   for open populations: estimating demographic parameters from camera-trap

581   studies. Ecology 91:3376–3383.

582   Goodwin, B. J., and L. Fahrig. 2002. How does landscape structure influence landscape

583   connectivity? Oikos 99:552–570.

584   Hanks, E. M., and M. B. Hooten. 2013. Circuit theory and model-based inference for

585   landscape connectivity. Journal of the American Statistical Association

586   108:22–33.

587   Hanski, I., M. Kuussaari, and M. Nieminen. 1994. Metapopulation structure and

588   migration in the butterfly *Melitaea Cinxia*. Ecology 75:747–762.

589   Higashide, D., S. Miura, and H. Miguchi. 2012. Are chest marks unique to Asiatic black

590   bear individuals? Journal of Zoology 288:199–206.

Higashide, D., S. Miura, and H. Miguchi. 2013. Evaluation of camera-trap designs for

photographing chest marks of the free-ranging Asiatic black bear, *Ursus*

thibetanus. Mammal Study 38:35–39.

Hirsch, C. 2007. Numerical computation of internal and external flows: the fundamentals

of computational fluid dynamics. Elsevier.

Horino, S., and S. Miura. 2000. Population viability analysis of a Japanese black bear

population. Population Ecology 42:37–44.

Iijima, H. 2020. A review of wildlife abundance estimation models: comparison of

models for correct application. Mammal Study 45:177–188.

Inman, H. F., and E. L. Bradley. 1989. The overlapping coefficient as a measure of

agreement between probability distributions and point estimation of the overlap

of two normal densities. Communications in Statistics - Theory and Methods

18:3851–3874.

Ivan, J. S., G. C. White, and T. M. Shenk. 2013. Using simulation to compare methods for

estimating density from capture-recapture data. Ecology 94:817–826.

Justeau-Allaire, D., G. Blanchard, T. Ibanez, X. Lorca, G. Vieilledent, and P. Birnbaum.

2022. Fragmented landscape generator (flsgen): A neutral landscape generator

with control of landscape structure and fragmentation indices. Methods in

Ecology and Evolution 13:1412–1420.

Long, J. A. 2019. Estimating wildlife utilization distributions using randomized shortest

paths. Landscape Ecology 34:2509–2521.

Marrotte, R. R., and J. Bowman. 2017. The relationship between least-cost and resistance

distance. PLOS ONE 12:e0174212.

McClintock, B. T., B. Abrahms, R. B. Chandler, P. B. Conn, S. J. Converse, R. L. Emmet,

B. Gardner, N. J. Hostetter, and D. S. Johnson. 2022. An integrated path for spatial

capture–recapture and animal movement modeling. Ecology:e03473.

McRae, B. H. 2006. Isolation by resistance. Evolution 60:1551–1561.

McRae, B. H., B. G. Dickson, T. H. Keitt, and V. B. Shah. 2008. Using circuit theory to

model connectivity in ecology, evolution, and conservation. Ecology

89:2712–2724.

Méndez, V., D. Campos, and F. Bartumeus. 2013. Stochastic foundations in movement

ecology: anomalous diffusion, front propagation and random searches. Springer

Science & Business Media.

Moorcroft, P. R., and A. Barnett. 2008. Mechanistic home range models and resource

selection analysis: a reconciliation and unification. Ecology 89:1112–1119.

Moorcroft, P. R., and M. A. Lewis. 2006. Mechanistic home range analysis. Princeton

University Press, Princeton.

Nocedal, J., and S. Wright. 2006. Numerical optimization. Springer Science & Business

Media.

Ohnishi, N., T. Osawa, T. Yamamoto, and R. Uno. 2019. Landscape heterogeneity in

landform and land use provides functional resistance to gene flow in continuous

Asian black bear populations. Ecology and Evolution 9:4958–4968.

Ohnishi, N., T. Saitoh, Y. Ishibashi, and T. Oi. 2007. Low genetic diversities in isolated

populations of the Asian black bear (Ursus thibetanus) in Japan, in comparison

with large stable populations. Conservation Genetics 8:1331–1337.

Osada, Y., T. Kuriyama, M. Asada, H. Yokomizo, and T. Miyashita. 2019. Estimating

range expansion of wildlife in heterogeneous landscapes: a spatially explicit

state-space matrix model coupled with an improved numerical integration

technique. Ecology and Evolution 9:318–327.

Ovaskainen, O. 2004. Habitat-specific movement parameters estimated using

mark–recapture data and a diffusion model. Ecology 85:242–257.

Ovaskainen, O., H. Rekola, E. Meyke, and E. Arjas. 2008. Bayesian methods for

analyzing movements in heterogeneous landscapes from mark–recapture data.

Ecology 89:542–554.

Panzacchi, M., B. Van Moorter, O. Strand, M. Saerens, I. Kivimäki, C. C. St. Clair, I.

Herfindal, and L. Boitani. 2016. Predicting the continuum between corridors and

barriers to animal movements using Step Selection Functions and Randomized

Shortest Paths. Journal of Animal Ecology 85:32–42.

Péron, G. 2019. Modified home range kernel density estimators that take environmental

interactions into account. Movement Ecology 7:16.

Potts, J. R., and U. E. Schlägel. 2020. Parametrizing diffusion-taxis equations from

animal movement trajectories using step selection analysis. Methods in Ecology

and Evolution 11:1092–1105.

Qi, Y., and J. Wu. 1996. Effects of changing spatial resolution on the results of landscape

pattern analysis using spatial autocorrelation indices. Landscape Ecology

11:39–49.

R Core Team. 2022. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna.

Royle, J. A., R. B. Chandler, K. D. Gazenski, and T. A. Graves. 2013. Spatial

capture–recapture models for jointly estimating population density and landscape

connectivity. Ecology 94:287–294.

Royle, J. A., A. K. Fuller, and C. Sutherland. 2018. Unifying population and landscape

ecology with spatial capture–recapture. Ecography 41:444–456.

Royle, J. A., K. U. Karanth, A. M. Gopalaswamy, and N. S. Kumar. 2009. Bayesian

inference in camera trapping studies for a class of spatial capture–recapture

models. Ecology 90:3233–3244.

Saitoh, T., Y. Ishibashi, H. Kanamori, and E. Kitahara. 2001. Genetic status of fragmented

populations of the Asian black bear Ursus thibetanus in western Japan. Population

Ecology 43:221–227.

Sakurai, R., and S. K. Jacobson. 2011. Public perceptions of bears and management

interventions in Japan. Human-Wildlife Interactions 5:123–134.

Schneider, S., G. W. Taylor, and S. C. Kremer. 2022. Similarity learning networks for

animal individual re-identification: an ecological perspective. Mammalian

Biology 102:899–914.

Signer, J., J. Fieberg, and T. Avgar. 2017. Estimating utilization distributions from fitted

step-selection functions. Ecosphere 8:e01771.

Smouse, P. E., S. Focardi, P. R. Moorcroft, J. G. Kie, J. D. Forester, and J. M. Morales.

2010. Stochastic modelling of animal movement. Philosophical Transactions of

the Royal Society B: Biological Sciences 365:2201–2211.

Sutherland, C., A. K. Fuller, and J. A. Royle. 2015. Modelling non-Euclidean movement

and landscape connectivity in highly structured ecological networks. Methods in

Ecology and Evolution 6:169–177.

Takahata, C., S. E. Nielsen, A. Takii, and S. Izumiyama. 2014. Habitat selection of a large

carnivore along human-wildlife boundaries in a highly modified landscape. PLOS

ONE 9:e86181.

Theng, M., C. Milleret, C. Bracis, P. Cassey, and S. Delean. 2022. Confronting spatial

capture–recapture models with realistic animal movement simulations.

Ecology:e3676.

Thurfjell, H., S. Ciuti, and M. S. Boyce. 2014. Applications of step-selection functions in

ecology and conservation. Movement Ecology 2:4.

Turner, M. G., R. V. O'Neill, R. H. Gardner, and B. T. Milne. 1989. Effects of changing

spatial scale on the analysis of landscape pattern. Landscape Ecology 3:153–162.

Yamaura, Y., R. J. Fletcher Jr, S. J. Lade, M. Higa, and D. Lindenmayer. 2022. From

nature reserve to mosaic management: Improving matrix survival, not

permeability, benefits regional populations under habitat loss and fragmentation.

Journal of Applied Ecology 59:1472–1483.

697   **Figures and tables**

698 **Fig. 1**   Example of simulated utility distributions under various dependencies of

699   permeability on a hypothetical landscape variable. Black dots indicate cells

700		with high permeability; other cells have low permeability. The red X
701		indicates the activity center.
702	**Fig. 2**	Example of simulated movement and the detection of an individual. The
703		transparent solid line is the movement path. Open rectangles are traps, and
704		blue circles indicate detection counts. The background tile map indicates
705		zones with high (green) and low (pale red) permeability.
706	**Fig. 3**	True vs. estimated parameters by ADCR. Dots and error bars indicate
707		maximum likelihood estimates and 95% confidence intervals. The solid
708		black line is y = x.
709	**Fig. 4**	Overlap between true and predicted home ranges by ADCR, SCR with the
710		least-cost path and basic SCR.
711	**Fig. 5**	Estimated effect of landscape on (A) permeability by ADCR and (B) cost
712		by SCR with the least-cost path (SCR-LCP) for black bear dataset. Dots
713		and error bars are the maximum likelihood estimate and 95% CIs,
714		respectively.
715		

716 **Table 1** Root mean squared error (RMSE) and mean bias error (MBE) of log density for the three estimators for the baseline scenario.

717　　　　　　The values binned into three classes of the true landscape effect, β, were also shown.

	RMSE			MBE		
	ADCR	SCR with the least-cost path	basic SCR	ADCR	SCR with the least-cost path	basic SCR
Total	0.0788	0.0789	0.0822	-0.00606	0.00123	0.0125
$\beta < -0.5$	0.0752	0.0762	0.0795	-0.0205	-0.0102	-0.00296
$|\beta| <= 0.5$	0.0677	0.0676	0.0730	0.0130	0.0139	0.0333
$\beta > 0.5$	0.0890	0.0883	0.0906	-0.00499	0.00380	0.0133

 Supporting information

symbols mark the home range center. Black dots

are zones with high permeability.

Fig. S5 Estimated log population density by ADCR, SCR

with the least-cost path (SCR-LCP) and basic SCR

under the scenario of severe data sparseness (i.e. g_0

= -5) and biased detector alignment. Black dots are

maximum likelihood estimates (MLEs). Open

circles and and error bars indicate the mean and

SD of MLEs. Horizontal black lines are the true

value (-2.0).

Fig. S6 Examples of predicted home ranges by (a) ADCR

and (b) SCR with the least-cost path for the black

bear. Red x symbols mark the home range centers.

Black dots are cells contain water surface.

Fig. S7 Comparison between the estimates of baseline

simulations and 1/2 resolution scenarios for (a)

effects of landscape and (b) log population

densities by ADCR. Dots and error bars are

maximum likelihood estimates and 95% CI,

respectively. Glay lines are principal components

of parameter estimates.

Appendix S6 Supplementary table.

Table S1 Target parameter ranges for the neutral landscape

generator, rflsgen. The parameter names were

767   based on Justeau-Allaire et al. (2022). Parameters

768   not shown were set to default values.

769   **Table S2**   Contingency table on the correctness of sign (i.e.

770   positive or negative) of landscape effect on

771   connectivity estimated by ADCR and SCR with the

772   least-cost path for the simulation scenario of sparse

773   data and biased detector alignment. The

774   significance was determined by non-overlap of

775   95% confidence interval with 0.

Probability

Fig. 1 Example of simulated utility distributions under various dependencies of permeability on a hypothetical landscape variable. Black dots indicate cells with high permeability; other cells have low permeability. The red X indicates the activity center.

ß = 0 ß = 0.5 ß = 1 ß = 1.5 ß = 2 ß = 2.5

$\beta = 0$
(circular home range)

Large β
(strong dependency on the landscape variable)

Fig. 2 Example of simulated movement and the detection of an individual. The transparent solid line is the movement path. Open rectangles are traps, and blue circles indicate detection counts. The background tile map indicates zones with high (green) and low (pale red) permeability.

Fig. 3 True vs. estimated parameters by ADCR. Dots and error bars indicate maximum likelihood estimates and 95% confidence intervals. The solid black line is y = x.

Fig. 4 Overlap between true and predicted home ranges by ADCR, SCR with the least-cost path and basic SCR.

Fig. 5 Estimated effect of landscape on (A) permeability by ADCR and (B) cost by SCR with the least-cost path (SCR-LCP) for black bear dataset. Dots and error bars are the maximum likelihood estimate and 95% CIs, respectively.