Title: Mechanistic home range capture-recapture models for the estimation of population density

and landscape connectivity

Keita Fukasawa

Biodiversity Division, National Institute for Environmental Studies

16-2 Onogawa, Tsukuba, Ibaraki, 305-8506 Japan

k.fukasawa37@gmail.com

Running head: Advection-diffusion capture-recapture models

Abstract

1. Spatially explicit capture-recapture (SECR) models provide an integrative statistical tool for analysing animal movement and population patterns. Although incorporating home range formation with step selection functions into SECR can improve the prediction of animal space use in a heterogeneous landscape, this approach is challenging owing to the sparseness of recapture events.

2. In this study, I developed an advection-diffusion capture-recapture model (ADCR), which is an extension of SECR incorporating home range formation with advection-diffusion formalism, providing a new framework to estimate population density and landscape connectivity. I tested the unbiasedness of the estimator and robustness to the landscape resolution using simulated capture-recapture data generated by a step selection function. I also compared parameter estimates with those from a spatial capture-recapture model based on the least cost path (SCR-LCP) and basic SECR.

3. Population density, connectivity, and home range estimates of ADCR were unbiased over randomly determined sets of true parameters. Although the accuracy of density estimates by ADCR was nearly identical to those of existing models, home range shape could be predicted more accurately by ADCR than by SCR-LCP. ADCR was robust to the choice of landscape resolution, which is a favourable characteristic in random walk-based connectivity measures.

4. ADCR provides unique opportunities to elucidate both individual- and population-level ecological processes from capture-recapture data. By offering a formal link with step selection functions to estimate animal movement, it is suitable for simultaneously modelling with capture-recapture data and animal movement data. This study provides a basis for studies of the interplay between animal movement processes and population patterns.

mechanistic home range analysis, partial differential equation, step selection function, advection-diffusion equation

Introduction

Anthropogenic activity leads to habitat loss and fragmentation, posing a major threat to biodiversity. In a fragmented landscape, connectivity between habitat patches affects population patterns and sustainability (Hanski et al., 1994). Functional connectivity in a heterogeneous landscape is not only of general interest in basic ecology but also a key factor affecting decision-making in wildlife management and conservation (Goodwin & Fahrig, 2002; Royle et al., 2018; Osada et al., 2019). Circuit theory and least cost path (LCP) modelling are two major paradigms for evaluating landscape connectivity, with different underlying behavioural assumptions and statistical properties (McRae et al., 2008; Marrotte & Bowman, 2017). Connectivity based on circuit theory has explicit links with animal movement models, such as random walks and step selection functions (McRae, 2006; Duchesne et al., 2015), and models of animal gene flow and movement have numerous applications (Dickson et al., 2019). Empirical determination of connectivity parameters, such as conductance of land cover types, is important to improve objectivity, and the development of estimators is a major challenge in ecology (Ovaskainen, 2004; Ovaskainen et al., 2008; Beyer et al., 2016; Signer et al., 2017; Osada et al., 2019).

Spatially explicit capture-recapture models (SECRs) are powerful tools with the capacity to estimate individual movement processes and population abundance in an integrative manner (McClintock et al., 2022). Although random-walk-based connectivity measures have favourable properties for practical applications, such as robustness to the resolution of landscape data (Marrotte & Bowman, 2017) and a straightforward link with movement models (Diniz et al., 2020), they are not as well-studied as LCP-based estimators (Royle et al., 2013; Sutherland et al., 2015; Fuller et al., 2016). In particular, the process of home range formation considering

random-walk-based connectivity and site fidelity has been studied extensively by step selection

functions in movement ecology (Thurfjell et al., 2014; Duchesne et al., 2015; Signer et al., 2017;

Péron, 2019; Potts & Schlägel, 2020). The corresponding SECR estimator can provide insight

into the relationship between individual-level displacement and population-level abundance

patterns. However, the temporal sparseness of capture-recapture data limits the application of

SECR to complex movement processes (McClintock et al., 2022).

Incorporating mechanistic models of home range formation into SECR can enable

inferences about connectivity as well as population abundance from capture-recapture data in a

computationally tractable manner. Mechanistic home range analysis (Moorcroft & Lewis, 2006;

Moorcroft & Barnett, 2008) is an analytical approach that ties the process of animal movements

to the resultant home range shape. The approach has contributed to the development of

advection-diffusion equations (ADE) to obtain flexible home ranges and likelihood-based

modelling to estimate parameters empirically. ADEs describe deterministic drift to a focal point

(e.g., an activity centre) as advection terms and random walks as diffusion terms. Circuit theory is

analogous to a diffusion process with heterogeneous diffusivity (Hanks & Hooten, 2013), and

ADEs with heterogeneous diffusion coefficients in space are applicable to home ranges formed

by animal movement affected by landscape resistance. Although a closed-form analytical

solutions of ADEs rarely exist, computationally efficient ways to obtain a numerical solution

have been developed (Hirsch, 2007). By integrating mechanistic home range analysis into

SECR, the simultaneous estimation of connectivity and population density is possible, without

substantial computational costs. Hereafter, I refer to this new SECR approach as the

advection-diffusion capture-recapture model (ADCR).

Although ADCRs and LCP-based spatial capture-recapture models differ with respect

to connectivity assumptions (LCP-SCR, Royle et al., 2013; Sutherland et al., 2015; Royle et al.,

2018) and cannot be directly compared, evaluations of the performance of estimators are helpful

for users. Applying statistical models to simulated data generated from a mechanistic animal

movement model is a straightforward way to test the unbiasedness of estimators and robustness

to deviation from assumptions (Dupont et al., 2022; Theng et al., 2022). In particular, deviations

in the spatial resolution of landscapes can often alter the results of landscape ecology (Qi & Wu,

1996; Turner et al., 1989), and the robustness of models to differences in spatial resolution

between data generation and model estimation should be evaluated.

The aim of this study was to develop a formal framework for ADCR and to compare

the accuracy of the estimated abundance and home range shape with those obtained by

LCP-SCR under a random walk-based movement process. The identification of the optimal

model is outside of the scope of this paper. Rather, my aim is to establish the merit of the ADCR

under animal movement modelled by a step selection function, which is common in movement

ecology. I specifically focused on a common situation in which the resolution of landscape

factors is lower than that of the data generating process, and I tested the robustness of

connectivity estimates to a decreased resolution. I also compared density estimates with those of

basic SECR (Borchers & Efford, 2008) as a null model of heterogeneous connectivity. I predict

that all three methods, ADCR, LCP-SCR, and basic SECR, will accurately estimate population

densities because SECR is robust to the violation of home range assumptions (Efford 2019). I

also predict that ADCR will have an advantage in estimating the effect of landscape on

connectivity, as it is robust to the resolution of input data.

Material and Methods

Model Framework

The mathematical foundation of the model is a classic model of animal home range behaviour

described by the 2D Ornstein-Uhlenbeck (OU) process with variable diffusion coefficients in

space (Smouse et al., 2010). The OU process is a biassed random walk with limit $\Delta t \rightarrow 0$ and is

formalised by a stochastic differential equation for constant drift toward an activity centre, $\boldsymbol{\mu} = (\mu_x,$

$\mu_y)$, and a Brownian motion term as follows:

$d\mathbf{s}_t = c(\boldsymbol{\mu} - \mathbf{s}_t)dt + \sigma(\mathbf{s}_t)d\mathbf{W}_t$ (eqn. 1)

where $\mathbf{s}_t = (x_t, y_t)$ is the animal location at time t, $\mathbf{W}_t$ is the 2D independent Wiener process, c is the

intensity of drift, and $\sigma(\mathbf{s}_t)$ is the standard deviation of the Wiener process, corresponding to the

step length per unit time depending on the location. For the special case when $\sigma(\mathbf{s}_t)$ is constant

over space, the probability distribution of $\mathbf{s}_t$ conditional on the initial location $\mathbf{s}_0$ is a bivariate

normal distribution as follows (Dunn & Gipson, 1977):

$p(\mathbf{s}_t | \mathbf{s}_0) = N\left(\exp(-ct)\mathbf{s}_0 + \left(1 - \exp\left(-ct\right)\right)\boldsymbol{\mu}, \frac{\sigma^2}{2c}\left(1 - exp(-2ct)\right)\mathbf{I}\right).$ (eqn. 2)

For the limiting case of $t \rightarrow +\infty$, this distribution converges to a static bivariate normal distribution,

a default home range assumption for basic SECRs (Borchers & Efford, 2008; Royle et al., 2009).

In contrast to the above Lagrangian approach describing the microscopic displacement

of an individual, the Eulerian approach treats movement as the macroscopic flow of the

probability density of individual locations. It is described by the Fokker-Planck equation, a partial

differential equation that describes the time evolution of the probability density in space

(Moorcroft & Lewis, 2006). Let p denote the probability density of individual location; the

corresponding Fokker-Planck equation for eqn. 1 is as follows:

$\frac{\partial p}{\partial t} = -c\left(\left(\mu_x - x\right)\frac{\partial p}{\partial x} + \left(\mu_y - y\right)\frac{\partial p}{\partial y}\right) + D(x, y)\left(\frac{\partial^2 p}{\partial x^2} + \frac{\partial^2 p}{\partial y^2}\right)$ (eqn. 3)

where $D(x, y) > 0$ is the diffusion coefficient. The first term on the right-hand side is advection

corresponding to drift toward the activity centre, and the second term is random diffusion. The

diffusion coefficient is interpreted as conductance in terms of circuit theory (McRae, 2006;

McRae et al., 2008). To model the effect of environmental factors on connectivity, I applied the

log-linear model $\ln(D(x, y)) = \beta_0 + \boldsymbol{\beta}\mathbf{z}(x, y)$, where β_0 is the intercept, $\mathbf{z}(x, y)$ is a vector of

environmental variables at location (x, y), and $\boldsymbol{\beta}$ is the corresponding regression coefficients

determining the contributions of environmental factors to landscape connectivity. For the special

case when σ is constant over space, $D(x, y) = \sigma^2/2$. The intensity of drift, c, takes a positive value,

and $\ln(c)$ is denoted by parameter α_0. To ensure that the volume of p is constant over time, the

reflecting boundary condition is given for space.

Similar to mechanistic home range analysis (Moorcroft & Lewis, 2006), home range

in this system is defined as the equilibrium probability density distribution, $p^*(x, y; \boldsymbol{\mu})$, which is

the solution of p when $\delta p/\delta t = 0$ subject to the condition that $\int_S p \, d\mathbf{s} = 1$, where S is the spatial

domain. Note that α_0 and β_0 are redundant at the equilibrium state because $\ln(D(x, y)/c) = \beta_0 - \alpha_0$

$+ \boldsymbol{\beta}\mathbf{z}(x, y)$. To ensure the identifiability of the model, it was reparametrized by $\gamma_0 = \beta_0 - \alpha_0$. γ_0 is

interpreted as the logarithm of the relative intensity of diffusion and drift, which determines the

average home range size. Although the closed-form solution rarely exists when $D(x, y)$ varies in

space, it can be numerically solved by a variety of discretization techniques, such as the finite

difference method and finite element method (Hirsch, 2007). In this study, the finite difference

method was applied because it is suitable for common situations in which landscape factors

affecting the diffusion coefficient are available as a raster dataset. In this study, grid cells of raster

data were used to discretize space for simplicity. Technical details for discretization and

numerical derivation of p^* are shown in Appendix S1.

To estimate connectivity parameters and population density simultaneously from

capture-recapture data, the mechanistic home range analysis was combined with the

likelihood-based SECR framework. The likelihood of SECR is a product of the Poisson process

likelihood that k individuals were observed at least once and the conditional likelihood of

capture history given k individuals, which follows a multinomial distribution (Borchers &

Efford, 2008). Activity centres are latent variables included in both parts, and they were

marginalised over spatial domain S. For a simple case that the $\ln$(population density) ρ is

constant over space, the likelihood function is

$$L(\boldsymbol{\theta}, \rho) = \frac{\{exp(\rho)a(\boldsymbol{\theta})\}^k exp\left(-exp(\rho)a(\boldsymbol{\theta})\right)}{k!} \times \binom{k}{k_1, k_2, \ldots, k_C} \prod_{i=1}^{k} \frac{\int_S Pr(\omega_i|\boldsymbol{\mu}, \boldsymbol{\theta})d\boldsymbol{\mu}}{a(\boldsymbol{\theta})}$$

(eqn. 5)

where $\boldsymbol{\theta}$ is a parameter vector affecting the detection process including γ_0 and $\boldsymbol{\beta}$ mentioned

above, and k_1, k_2, …, k_C are frequencies of unique capture histories. $a(\boldsymbol{\theta}) = \int_S p_.(\boldsymbol{\mu}, \boldsymbol{\theta}) d\boldsymbol{\mu}$,

where $p_.(\boldsymbol{\mu}, \boldsymbol{\theta})$ is the probability that an individual whose activity centre is $\boldsymbol{\mu}$ is detected at least

once. $Pr(\boldsymbol{\omega}_i | \boldsymbol{\mu}, \boldsymbol{\theta})$ is the probability of capture history of the ith individual given $\boldsymbol{\mu}$. The

capture probability is the product of the likelihood of all detectors and occasions, which depends

on the detector type. For example, count data from detectors are often described by a Poisson

distribution (Distiller et al., 2020; Efford et al., 2013), Poisson($\omega_{ij} | \exp(g_0)E_j p^*(x_j, y_j | \boldsymbol{\mu}, \gamma_0, \boldsymbol{\beta})$),

where ω_{ij} is the detection number for the ith individual at the jth detector, g_0 is log catchability,

E_j is survey effort, and $p^*(x_j, y_j | \boldsymbol{\mu}, \gamma_0, \boldsymbol{\beta})$ is the probability density of individual location at

detector (x_j, y_j). If iterative surveys using binomial detectors are conducted for T occasions, the

frequency of detection ω_{ij} follows a binomial distribution, Binomial($\omega_{ij} | T$, 1 - exp(-exp(g_0)$p^*(x_j$,

$y_j | \boldsymbol{\mu}, \gamma_0, \boldsymbol{\beta})$)).

Model parameters ρ, g_0, γ_0, and $\boldsymbol{\beta}$ are estimated using a maximum likelihood method.

Although conditional likelihood maximisation can be used for parameter estimation in SECR

(Borchers & Efford, 2008), full likelihood maximisation was applied because it has the capacity

to deal with complex situations in which population density varies across space. Automatic

differentiation of likelihood functions (Auger-Méthé et al., 2017) is not currently available for

the model because p^* is derived using an iterative sparse linear solver, but a quasi-Newton

algorithm, such as the Broyden–Fletcher–Goldfarb–Shanno (BFGS, Nocedal & Wright, 2006)

algorithm with numerical differentiation of log likelihoods, works well.

Simulation study

To test the capacity of ADCR to estimate parameters of connectivity and population density

correctly, I used a dataset generated via Monte Carlo simulations of animal movement. The

landscape is a rectangular grid system of 50×50 (hence, the area is 2500), and animal location

was described as a discrete coordinate of grid cells. Arrays of count detectors were set to the

range, $x = [9, 41]$ and $y = [9, 41]$, with x and y axis spacing of 4 and 81 detectors.

Simulations were iterated 100 times, and true parameter values and landscape structure

were randomly determined in each iteration. For each iteration, a heterogeneous landscape was

generated using the R package rflsgen (Justeau-Allaire et al., 2022). Random patches were

generated in the landscape by rflsgen with parameters shown in Table S1. For covariate values,

0.5 and -0.5 were assigned to patch and non-patch habitats, respectively. True parameter values

were sampled from uniform distributions for each iteration. The ranges of true parameters were

set as follows: log population density ρ = (-3, -1), log intensity of drift α_0 = (-1, 0) and

coefficient of landscape variable β = (-2, 2). g_0 and β_0 were fixed to -3 and 1, respectively. At the

beginning of an iteration, the number of individuals in the landscape was determined by a

Poisson distribution with expected value of $\exp(\rho) \times 2500$ and activity centres were drawn from

a discrete uniform distribution.

The animal movement simulation was run for t = 5000 after 100 burn-in steps for

randomising initial locations of individuals. For each time step, we calculated the probability

vector of individual locations for all cells given the location at the previous step using the

advection-diffusion equation with the explicit method (eqn. 7 in Appendix S1), and the location

was sampled from a categorical distribution. When there was a detector in a cell in which an

individual was located, the number of detection events was simulated by a Poisson distribution

with expectation $\exp(g_0)$. Then, single-occasion capture-recapture data were obtained by

censoring undetected individuals. An example map of simulated movement and observations of

an individual is shown in Fig. 1.

I applied ADCR, LCP-SCR, and basic SECR to the simulated dataset to evaluate the

accuracy of estimations. For the three estimators, the deviation of estimates from true values

was determined and root mean square error (RMSE) and mean bias error (MBE) were

calculated. To show how well ADCR and LCP-SCR reproduce home range shapes, the

agreement between true (p^*_{true}) and predicted (p^*_{pred}) probability distributions of individual

locations was calculated by an overlapping coefficient (Inman & Bradley, 1989):

$$V(\boldsymbol{\mu}) = \iint_{\mathcal{S}} \; min\left(p^*_{pred}(x, y; \boldsymbol{\mu}), p^*_{true}(x, y; \boldsymbol{\mu})\right) dxdy$$

The overlapping coefficient ranges from 0 (no overlap) to 1 (perfect overlap). To evaluate the

average performance over locations for activity centre $\boldsymbol{\mu}$, I calculated the overlapping

coefficients marginalised over $\boldsymbol{\mu}$ within the range of trap array x = [9, 41] and y = [9, 41] for

each model and iteration.

To evaluate the robustness of ADCR and LCP-SCR to deviations in the landscape

resolution of data from the true state, I considered an additional scenario in which the available

landscape variable was restricted to 1/2 resolution. For the 1/2 resolution scenario, grid cells

with side length 2 were used for discretizing space in the model estimations, and the landscape

variable was averaged for each grid cell. I compared the density and connectivity estimates for

the 1/2 resolution scenario with estimates for the baseline scenario.

All models were estimated using maximum likelihood methods using R 4.1.3 (R Core

Team, 2022). The movement simulation and log likelihood function of ADCR was written

largely in R language, with some subroutines in C++. The source code is available online

(https://github.com/kfukasawa37/adcrtest2). The log likelihood values of LCP-SCR were

derived based on the R script by Sutherland et al. (2015), with some modifications by Keita

Fukasawa to deal with count detectors and the negative effect of landscape on the cost function.

The basic SECR was estimated using the R package secr (Efford, 2022).

232   **Results**

For all iterations, estimations of ADCR converged successfully. All parameters of ADCR,

including the population density and effect of landscape on connectivity, were estimated in an

unbiased way. The true and estimated parameters were on the line y = x, on average (Fig. 2).

RMSE and MBE of g_0 were 0.0258 and 0.0109, respectively. The true and predicted home

ranges overlapped fairly well. The overlapping coefficients marginalised over activity centres

were (0.950, 0.9996) over 100 iterations, and the mean was 0.988 (Fig. 3).

As predicted, population densities estimated by LCP-SCR and basic SECR were

robust to true connectivity settings in our simulation experiments. RMSEs and bias of log

population density estimated by LCP-SCR and basic SECR were very similar to those of ADCR

and were not sensitive to the true effect of landscape on connectivity (Table 1). Effects of

landscape on the LCP distance estimated by LCP-SCR were linear, on average, but showed a

slightly weaker   relationship with the true parameters of connectivity (Pearson's r = -0.988, Fig.

S1) than that of ADCR (Pearson's r = 0.995, Fig. 2). The overlapping coefficients of home range

predicted by LCP-SCR were lower than those for ADCR over the range of true connectivity

parameters, and the discrepancy between predicted and true home range was large when the true

effect of landscape on connectivity was far from 0 (Fig. 3). LCP-SCR tended to predict a more

blurred and isotropic home range shape than the true home range shape (Fig. S2).

ADCR was robust to mismatches in landscape resolution between the data generating

process and model estimation. The log population density and effect of landscape in the 1/2

resolution scenario were almost the same as those for the baseline scenario (Fig. 4ac, slopes of

principal axes were 0.999 and 1.05, respectively). Although the population density determined

by LCP-SCR was also robust to spatial resolution (Fig. 4b, slope of the principal axis was 1.00),

the effect of landscape on connectivity was weaker in the 1/2 resolution scenario than in the

baseline scenario (Fig. 4d, slope of principal axis was 0.912).

257   **Discussion**

Individual movement is an important driver of spatial patterns of populations, and incorporating

movement processes in spatial capture-recapture analysis will facilitate studies of the interplay

between individual- and population-level ecological processes (Royle et al., 2018; McClintock

et al., 2022). The integration of LCP modelling with spatial capture recapture analysis (Royle et

al., 2013; Sutherland et al., 2015) was a great methodological advancement for the simultaneous

estimation of population density and ecological distance, but accurate estimation of movement

parameters exclusively from capture-recapture data is still difficult (Dupont et al., 2022). In this

study, I developed a formal framework for a spatial capture-recapture model including

mechanistic home range formation with an explicit theoretical link to animal movement models.

Through individual-based simulations of animal movement and detection, ADCR accurately

estimated the population density and effect of landscape accurately, even when there was a

mismatch in spatial resolution between the data generating process and available data. This

property provides an advantage over existing estimators, suggesting that ADCR is an effective

approach for estimating population density and random walk-based landscape connectivity.

Landscape connectivity is an important determinant of animal space use, including

foraging and reproduction, and hence is related to survival and population dynamics. Predicting

connectivity is a focus of research in landscape ecology and crucial for conservation planning

(Dickson et al., 2019). Although circuit theory is a powerful tool for connectivity modelling in

heterogeneous landscapes, it is limited by the lack of a statistical tool for estimating relative

conductance values (Zeller et al., 2012). Estimated effects of landscape variables on diffusion

coefficients by ADCR can be directly applied to predict relative conductance for calculating

resistance distance. ADCR provides an important tool for objective reasoning in landscape

connectivity studies. Estimates obtained by ADCR were robust to different spatial resolutions,

which is a favourable characteristic of connectivity measures in circuit theory, which shares the

same theoretical foundation of random walk descriptions of animal movement (Hanks &

Hooten, 2013).

The effect of a skewed home range on the accuracy of density estimation by SECR

has been studied extensively in ecology (Ivan et al., 2013; Royle et al., 2013; Sutherland et al.,

2015; Efford, 2019; Theng et al., 2022). The results of this study were consistent with those of

recent studies (Efford 2019; Theng et al., 2022) showing that density estimates are generally

robust to deviations in assumptions on the distribution of home range shapes. Although some

studies (Royle et al., 2013; Sutherland et al., 2015) have reported that density estimates by basic

SECR can be biased under the specific spatial patterns of landscape connectivity, our results

indicated that the bias in density averaged over a randomly generated landscape was small.

Efford (2019) emphasised the importance of sampling design in space, such as the use of a

two-dimensional detector array and avoidance of detector array alignment along landscape

features, for the correct estimation of density. Obviously, such sampling designs are also

favourable for the precise estimation of landscape effects on home range shape by ADCR.

Besides the accurate estimation of population density, the integration of animal movement

modelling and capture-recapture can improve scale up from individual to population levels

(McClintock et al., 2022), and ADCR is useful for such a purpose.

Joint statistical modelling of animal movement trajectories obtained by GPS loggers

and spatial capture-recapture data is an important approach for understanding detailed animal

movement processes, while ensuring the generality of results (Dupont et al., 2022). Both ADCR

and the step selection function are based on biassed random walk models (Duchesne et al.,

2015), and ADCR can be applied to joint modelling, retaining consistency between the animal

movement model and resultant home range. Although the static home range assumption of

ADCR has an obvious advantage in the stability of estimates under a severe data deficiency,

intercepts of movement speed and site fidelity are aggregated into a single parameter and not

identifiable. Joint modelling of the step selection function and ADCR can overcome this

limitation by estimating these parameters separately.

The broad application of SECR in ecology can largely be attributed to its flexibility to

incorporate heterogeneity in ecological processes underlying spatial capture recapture data, such

as environment-dependent abundance and sex-biassed detection (Efford, 2022). Future advances

in ADCR should focus on similar functionality. In particular, incorporating heterogeneity in

home range size (Efford et al., 2016) as well as heterogeneity in connectivity is a major

challenge. Although I focused only on heterogeneous connectivity for simplicity, the

simultaneous estimation of environmental dependence for both home range size and

connectivity can be achieved by incorporating γ_0 varying with the location of the activity centre.

Also, the generalisation of ADCR for dealing with temporally open populations driven by

birth-death processes (Gardner et al., 2010; Efford & Schofield 2022) would be helpful for

analysing long-term capture-recapture data. The development of software tools that enable users

to benefit from the flexible nature of ADCR is an urgent task for expanding the application of

the method.

Similar to step selection analysis linked with advection-diffusion equations (Potts &

Schlägel, 2020), an important assumption of ADCR is that sites of animal detection do not show

substantial temporal autocorrelation. In general, this assumption would be valid for

capture-recapture data in which the number of detections per individual is not large. However,

our approach can be biassed if the detectability of individuals is quite high and animal steps are

highly correlated due to non-uniform turning angles. Modelling correlated movement with a

PDE formalism is still a challenging issue (Patlak, 1953; Wang & Potts, 2017), and future

research on this topic has the potential to improve home range representation under more

realistic situations.

Spatial capture-recapture is a popular method for abundance analyses in ecology and

conservation (Iijima, 2020), and large amounts of data have accumulated. Moreover, recent

technical advances in individual identification from environmental DNA (Dugal et al., 2022)

and machine observations, such as camera traps (Schneider et al., 2022), will expand the

applicability of capture-recapture approaches to broad taxa. Such capture-recapture data have

the potential to advance our understanding of the process of animal home range formation.

ADCR will be a useful tool for integrated analyses of capture recapture data incorporating

individual- and population-level ecological processes.

Acknowledgements

Yutaka Osada provided valuable comments on the earlier version of the statistical model. This study was supported by JSPS KAKENHI Grant Numbers JP13247736, JP19153418, and by the Environment Research and Technology Development Fund (JPMEERF20194005) of the Environmental Restoration and Conservation Agency Provided by the Ministry of Environment of Japan.

Conflict of interest

The author has no conflict of interest to be declared.

Author contributions

KF conceived the ideas, designed methodology, conducted simulation study and wrote the manuscript.

Data availability

No external data were used in this study.

References

Auger-Méthé, M., Albertsen, C. M., Jonsen, I. D., Derocher, A. E., Lidgard, D. C., Studholme, K. R., Bowen, W. D., Crossin, G. T., & Flemming, J. M. (2017). Spatiotemporal modelling of marine movement data using Template Model Builder (TMB). *Marine Ecology Progress Series*, *565*, 237–249. https://doi.org/10.3354/meps12019

Beyer, H. L., Gurarie, E., Börger, L., Panzacchi, M., Basille, M., Herfindal, I., Van Moorter, B., R.

Lele, S., & Matthiopoulos, J. (2016). 'You shall not pass!': Quantifying barrier

permeability and proximity avoidance by animals. *Journal of Animal Ecology*, *85*(1),

43–53. https://doi.org/10.1111/1365-2656.12275

Borchers, D. L., & Efford, M. G. (2008). Spatially explicit maximum likelihood methods for

capture–recapture studies. *Biometrics*, *64*(2), 377–385.

https://doi.org/10.1111/j.1541-0420.2007.00927.x

Dickson, B. G., Albano, C. M., Anantharaman, R., Beier, P., Fargione, J., Graves, T. A., Gray, M.

E., Hall, K. R., Lawler, J. J., Leonard, P. B., Littlefield, C. E., McClure, M. L., Novembre,

J., Schloss, C. A., Schumaker, N. H., Shah, V. B., & Theobald, D. M. (2019).

Circuit theory applications to connectivity science and conservation. *Conservation*

Biology, *33*(2), 239–249. https://doi.org/10.1111/cobi.13230

Diniz, M. F., Cushman, S. A., Machado, R. B., & De Marco Júnior, P. (2020). Landscape

connectivity modeling from the perspective of animal dispersal. *Landscape Ecology*,

35(1), 41–58. https://doi.org/10.1007/s10980-019-00935-3

Distiller, G. B., Borchers, D. L., Foster, R. J., & Harmsen, B. J. (2020). Using continuous-time

spatial capture–recapture models to make inference about animal activity patterns.

Ecology and Evolution, *10*(20), e6822. https://doi.org/10.1002/ece3.6822

Duchesne, T., Fortin, D., & Rivest, L.-P. (2015). Equivalence between step selection functions

and biased correlated random walks for statistical inference on animal movement. *PLOS*

ONE, *10*(4), e0122947. https://doi.org/10.1371/journal.pone.0122947

Dugal, L., Thomas, L., Jensen, M. R., Sigsgaard, E. E., Simpson, T., Jarman, S., Thomsen, P. F., &

Meekan, M. (2022). Individual haplotyping of whale sharks from seawater

environmental DNA. *Molecular Ecology Resources*, *22*(1), 56–65.

https://doi.org/10.1111/1755-0998.13451

Dunn, J. E., & Gipson, P. S. (1977). Analysis of radio telemetry data in studies of home range.

Biometrics, *33*(1), 85–101. https://doi.org/10.2307/2529305

Dupont, G., Linden, D. W., & Sutherland, C. (2022). Improved inferences about landscape

connectivity from spatial capture–recapture by integration of a movement model.

Ecology, e03544. https://doi.org/10.1002/ecy.3544

Efford, M. G. (2019). Non-circular home ranges and the estimation of population density.

Ecology, *100*(2), e02580. https://doi.org/10.1002/ecy.2580

Efford, M. G. (2022). *secr: Spatially explicit capture-recapture models* (R package version 4.5.6).

https://CRAN.R-project.org/package=secr

Efford, M. G., Borchers, D. L., & Mowat, G. (2013). Varying effort in capture–recapture studies.

Methods in Ecology and Evolution, *4*(7), 629–636.

https://doi.org/10.1111/2041-210X.12049

Efford, M. G., Dawson, D. K., Jhala, Y. V., & Qureshi, Q. (2016). Density-dependent home-range

size revealed by spatially explicit capture–recapture. *Ecography*, *39*(7), 676–688.

Efford, M. G., & Schofield, M. R. (2022). A review of movement models in open population

capture–recapture. *Methods in Ecology and Evolution*, *13*(10), 2106–2118.

https://doi.org/10.1111/2041-210X.13947

Fuller, A. K., Sutherland, C. S., Royle, J. A., & Hare, M. P. (2016). Estimating population density

and connectivity of American mink using spatial capture–recapture. *Ecological*

Applications, *26*(4), 1125–1135. https://doi.org/10.1890/15-0315

Gardner, B., Reppucci, J., Lucherini, M., & Royle, J. A. (2010). Spatially explicit inference for

open populations: Estimating demographic parameters from camera-trap studies.

Ecology, *91*(11), 3376–3383. https://doi.org/10.1890/09-0804.1

Goodwin, B. J., & Fahrig, L. (2002). How does landscape structure influence landscape

connectivity? *Oikos*, *99*(3), 552–570. https://doi.org/10.1034/j.1600-0706.2002.11824.x

Hanks, E. M., & Hooten, M. B. (2013). Circuit theory and model-based inference for landscape

connectivity. *Journal of the American Statistical Association*, *108*(501), 22–33.

https://doi.org/10.1080/01621459.2012.724647

Hanski, I., Kuussaari, M., & Nieminen, M. (1994). Metapopulation structure and migration in the

butterfly *Melitaea Cinxia*. *Ecology*, *75*(3), 747–762. https://doi.org/10.2307/1941732

Hirsch, C. (2007). *Numerical Computation of Internal and External Flows: The Fundamentals of*

Computational Fluid Dynamics. Elsevier.

Iijima, H. (2020). A review of wildlife abundance estimation models: comparison of models for

correct   application.   *Mammal   Study*,   *45*(3),   177–188.

https://doi.org/10.3106/ms2019-0082

Inman, H. F., & Bradley, E. L. (1989). The overlapping coefficient as a measure of agreement

between probability distributions and point estimation of the overlap of two normal

densities. *Communications in Statistics - Theory and Methods*, *18*(10), 3851–3874.

https://doi.org/10.1080/03610928908830127

Ivan, J. S., White, G. C., & Shenk, T. M. (2013). Using simulation to compare methods for

estimating   density   from   capture-recapture   data.   *Ecology*,   *94*(4),   817–826.

https://doi.org/10.1890/12-0102.1

Justeau-Allaire, D., Blanchard, G., Ibanez, T., Lorca, X., Vieilledent, G., & Birnbaum, P. (2022).

Fragmented landscape generator (flsgen): A neutral landscape generator with control of

landscape structure and fragmentation indices. *Methods in Ecology and Evolution*, *13*(7),

1412–1420. https://doi.org/10.1111/2041-210X.13859

Marrotte, R. R., & Bowman, J. (2017). The relationship between least-cost and resistance

distance. *PLOS ONE*, *12*(3), e0174212. https://doi.org/10.1371/journal.pone.0174212

McClintock, B. T., Abrahms, B., Chandler, R. B., Conn, P. B., Converse, S. J., Emmet, R. L.,

431   Gardner, B., Hostetter, N. J., & Johnson, D. S. (2022). An integrated path for spatial

432   capture–recapture and animal movement modeling. *Ecology*, e03473.

433   https://doi.org/10.1002/ecy.3473

434   McRae, B. H. (2006). Isolation by resistance. *Evolution*, *60*(8), 1551–1561.

435   https://doi.org/10.1111/j.0014-3820.2006.tb00500.x

436   McRae, B. H., Dickson, B. G., Keitt, T. H., & Shah, V. B. (2008). Using circuit theory to model

437   connectivity in ecology, evolution, and conservation. *Ecology*, *89*(10), 2712–2724.

438   https://doi.org/10.1890/07-1861.1

439   Moorcroft, P. R., & Barnett, A. (2008). Mechanistic home range models and resource selection

440   analysis: A reconciliation and unification. *Ecology*, *89*(4), 1112–1119.

441   https://doi.org/10.1890/06-1985.1

442   Moorcroft, P. R., & Lewis, M. A. (2006). *Mechanistic Home Range Analysis.* Princeton

443   University Press.

444   Nocedal, J., & Wright, S. (2006). *Numerical Optimization*. Springer Science & Business Media.

445   Osada, Y., Kuriyama, T., Asada, M., Yokomizo, H., & Miyashita, T. (2019). Estimating range

446   expansion of wildlife in heterogeneous landscapes: A spatially explicit state-space matrix

447   model coupled with an improved numerical integration technique. *Ecology and Evolution*,

9(1), 318–327. https://doi.org/10.1002/ece3.4739

Ovaskainen, O. (2004). Habitat-specific movement parameters estimated using mark–recapture

data and a diffusion model. *Ecology*, *85*(1), 242–257. https://doi.org/10.1890/02-0706

Ovaskainen, O., Rekola, H., Meyke, E., & Arjas, E. (2008). Bayesian methods for analyzing

movements in heterogeneous landscapes from mark–recapture data. *Ecology*, *89*(2),

542–554. https://doi.org/10.1890/07-0443.1

Patlak, C. S. (1953). Random walk with persistence and external bias. *The Bulletin of*

Mathematical Biophysics, *15*(3), 311–338. https://doi.org/10.1007/BF02476407

Péron, G. (2019). Modified home range kernel density estimators that take environmental

interactions into account. *Movement Ecology*, *7*(1), 16.

https://doi.org/10.1186/s40462-019-0161-9

Potts, J. R., & Schlägel, U. E. (2020). Parametrizing diffusion-taxis equations from animal

movement trajectories using step selection analysis. *Methods in Ecology and Evolution*,

11(9), 1092–1105. https://doi.org/10.1111/2041-210X.13406

Qi, Y., & Wu, J. (1996). Effects of changing spatial resolution on the results of landscape pattern

analysis using spatial autocorrelation indices. *Landscape Ecology*, *11*(1), 39–49.

https://doi.org/10.1007/BF02087112

R Core Team. (2022). *R: A language and environment for statistical computing* (4.1.3). R

Foundation for Statistical Computing. https://www.R-project.org/

Royle, J. A., Chandler, R. B., Gazenski, K. D., & Graves, T. A. (2013). Spatial capture–recapture

models for jointly estimating population density and landscape connectivity. *Ecology*,

94(2), 287–294. https://doi.org/10.1890/12-0413.1

Royle, J. A., Fuller, A. K., & Sutherland, C. (2018). Unifying population and landscape ecology

with spatial capture–recapture. *Ecography*, *41*(3), 444–456.

https://doi.org/10.1111/ecog.03170

Royle, J. A., Karanth, K. U., Gopalaswamy, A. M., & Kumar, N. S. (2009). Bayesian inference in

camera trapping studies for a class of spatial capture–recapture models. *Ecology*, *90*(11),

3233–3244. https://doi.org/10.1890/08-1481.1

Schneider, S., Taylor, G. W., & Kremer, S. C. (2022). Similarity learning networks for animal

individual re-identification: An ecological perspective. *Mammalian Biology*, *102*(3),

899–914. https://doi.org/10.1007/s42991-021-00215-1

Signer, J., Fieberg, J., & Avgar, T. (2017). Estimating utilization distributions from fitted

step-selection functions. *Ecosphere*, *8*(4), e01771. https://doi.org/10.1002/ecs2.1771

Smouse, P. E., Focardi, S., Moorcroft, P. R., Kie, J. G., Forester, J. D., & Morales, J. M. (2010).

Stochastic modelling of animal movement. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, *365*(1550), 2201–2211.

https://doi.org/10.1098/rstb.2010.0078

Sutherland, C., Fuller, A. K., & Royle, J. A. (2015). Modelling non-Euclidean movement and

landscape connectivity in highly structured ecological networks. *Methods in Ecology and*

Evolution, *6*(2), 169–177. https://doi.org/10.1111/2041-210X.12316

Theng, M., Milleret, C., Bracis, C., Cassey, P., & Delean, S. (2022). Confronting spatial

capture–recapture models with realistic animal movement simulations. *Ecology*, e3676.

https://doi.org/10.1002/ecy.3676

Thurfjell, H., Ciuti, S., & Boyce, M. S. (2014). Applications of step-selection functions in

ecology and conservation. *Movement Ecology*, *2*(1), 4.

https://doi.org/10.1186/2051-3933-2-4

Turner, M. G., O'Neill, R. V., Gardner, R. H., & Milne, B. T. (1989). Effects of changing spatial

scale on the analysis of landscape pattern. *Landscape Ecology*, *3*(3), 153–162.

https://doi.org/10.1007/BF00131534

Wang, Y.-S., & Potts, J. R. (2017). Partial differential equation techniques for analysing animal

movement: A comparison of different methods. *Journal of Theoretical Biology*, *416*,

52–67. https://doi.org/10.1016/j.jtbi.2017.01.002

Zeller, K. A., McGarigal, K., & Whiteley, A. R. (2012). Estimating landscape resistance to

movement: A review. *Landscape Ecology*, *27*(6), 777–797.

503   **Figures and tables**

504   **Figure 1**   Example of simulated movement and the detection of an individual. The

505   transparent solid line is the movement path. Open rectangles are traps, and blue

506   circles indicate detection counts. The background tile map indicates zones with

507   high (green) and low (pale red) permeability.

508   **Figure 2**   True vs. estimated parameters by ADCR. Dots and error bars indicate maximum

509   likelihood estimates and 95% confidence intervals. The solid black line is $y = x$.

510   **Figure 3**   Overlap between true and predicted home ranges by ADCR and SCR-LCP.

511   **Figure 4**   Comparison of estimates between baseline and 1/2 resolution scenarios: log

512   population density by (a) ADCR and (b) SCR-LCP, and effect of landscape on

513   connectivity by (c) ADCR and (d) SCR-LCP. Black and grey lines are $y=x$ and the

514   principal axis, respectively.

516 **Table 1** RMSE and MBE of log density for the three estimators for the baseline scenario. The values binned into three classes of the true landscape

517 effect, β, were also shown.

	RMSE			MBE		
	ADCR	LCP-SCR	basic SECR	ADCR	LCP-SCR	basic SECR
Total	0.0788	0.0789	0.0822	-0.00606	0.00123	0.0125
$\beta < -0.5$	0.0752	0.0762	0.0795	-0.0205	-0.0102	-0.00296
$|\beta| <= 0.5$	0.0677	0.0676	0.0730	0.0130	0.0139	0.0333
$\beta > 0.5$	0.0890	0.0883	0.0906	-0.00499	0.00380	0.0133

Supporting information

Appendix S1   Technical details in finite difference method.

Appendix S2   Supplementary figures.

Fig. S1   Relationship between the true effect of landscape on
connectivity and estimated effect of landscape on cost by
SCR-LCP. Error bars are 95% CIs.

Fig. S2   Example of true and predicted home ranges by ADCR and
SCR-LCP. Red x symbols mark the home range centre.
Black dots are zones with high permeability.

Appendix S3   Supplementary table.

Table S1   Target parameter ranges for the neutral landscape generator,
rflsgen.   The   parameter   names   were   based   on
Justeau-Allaire et al. (2022). Parameters not shown were
set to default values.

Fig. 1 Example of simulated movement and the detection of an individual. The transparent solid line is the movement path. Open rectangles are traps, and blue circles indicate detection counts. The background tile map indicates zones with high (green) and low (pale red) permeability.

Fig. 2 True vs. estimated parameters by ADCR. Dots and error bars indicate maximum likelihood estimates and 95% confidence intervals. The solid black line is y = x.

Fig. 3 Overlap between true and predicted home ranges by ADCR and SCR-LCP.

Fig. 4 Comparison of estimates between baseline and 1/2 resolution scenarios: log population density by (a) ADCR and (b) SCR-LCP, and effect of landscape on connectivity by (c) ADCR and (d) SCR-LCP. Black and grey lines are y=x and the principal axis, respectively.