A critical role of a eubiotic microbiota in gating proper immunocompetence in *Arabidopsis*

Authors: Bradley C. Paasch[1, 2, †], Reza Sohrabi[1, 3, †], James M. Kremer[3, †], Kinya Nomura[1, 2], Jennifer Martz[3], Brian Kvitko[4], James M. Tiedje[5], Sheng Yang He[1, 2, *]

Affiliations: [1]Department of Biology, Duke University, Durham, NC, USA; [2]Howard Hughes Medical Institute, Duke University, Durham, NC, USA; [3]Department of Energy Plant Research Laboratory, Michigan State University, East Lansing, MI, USA [4]Department of Plant Pathology, University of Georgia, Athens, GA, USA; [5]Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing, MI, USA.

[†]Denotes equal contribution
[*]Corresponding author

Abstract

Although many studies have shown that microbes can ectopically stimulate or suppress plant immune responses, the fundamental question of whether the entire preexisting microbiota is indeed required for proper development of plant immune response remains unanswered. Using a recently developed peat-based gnotobiotic plant growth system we found that *Arabidopsis* grown in the absence of a natural microbiota lacked age-dependent maturation of plant immune response and were defective in several aspects of pattern-triggered immunity. Axenic plants exhibited hypersusceptibility to infection by the foliar bacterial pathogen *Pseudomonas syringae* pv. *tomato* DC3000. Microbiota-mediated immunocompetence was suppressed by rich nutrient conditions, indicating a tripartite interaction between the host, microbiota, and abiotic environment. A synthetic microbiota composed of 48 culturable bacterial strains from the leaf endosphere of healthy *Arabidopsis* plants was able to substantially restore immunocompetence similar to plants inoculated with a soil-derived community. In contrast, a 52-member dysbiotic synthetic leaf microbiota overstimulated the immune transcriptome. Together, these results provide evidence for a causal role of a eubiotic microbiota in gating proper immunocompetence and age-dependent immunity in plants.

Main

The aboveground and belowground parts of land plants host a variety of microorganisms, which collectively constitute the plant microbiota, and microbiota members can reside on or inside plants and appear to be taxonomically conserved at the phylum level[1-8]. The broad conservation of plant microbiota suggests that plants likely have evolved mechanisms to select and maintain the abundance, composition, or function of microbiota to achieve homeostasis[9]. A correctly assembled microbiota (i.e., eubiotic microbiota) is likely essential for plant health and survival as recent studies have begun to reveal deleterious effects of dysbiotic microbiotas on plant health[10,11]. Although microbiota members have been shown to enhance the ability of plants to acquire nutrients, to grow and/or to resist abiotic and biotic stresses[1,2,12-14], due to poorly dissected microbe-microbe and microbe-plant interactions at the community level, it is not well understood how an entire indigenous plant microbiota might contribute to various plant functions.

Different members of the plant microbiota can form mutualistic, commensal or pathogenic interactions with plants. To protect against potentially harmful exploitations by microorganisms, plants have evolved cell surface and intracellular immune receptors that recognize evolutionarily conserved microbe-associated molecular patterns (PAMPs) or pathogen-derived effector proteins, resulting in pattern-triggered immunity (PTI) or effector-triggered immunity (ETI), respectively. While ETI appears to be specific for pathogens, PTI represents a basal line of plant defense against both pathogenic and nonpathogenic microbes and is required for maintaining a eubiotic phyllosphere microbiota in *Arabidopsis* to prevent dysbiosis[10,11]. PTI signaling is initiated upon perception of PAMPs by plasma membrane-localized pattern recognition receptors (PRRs)[15]. For example, a 22-amino-acid epitope derived from bacterial flagellin (flg22) is a well characterized elicitor of PTI and recognized by the PRR FLAGELLIN-SENSITIVE 2 (FLS2)[16]. FLS2 forms a complex with co-receptor BRASSINOSTEROID INSENSITIVE 1-ASSOCIATED RECEPTOR KINASE 1 (BAK1)[17]. Phosphorelays between FLS2, BAK1, BOTRYTIS-INDUCED KINASE 1 (BIK1) and a MAPK cascade initiate downstream PTI signaling events, including the production of reactive oxygen species (ROS), calcium fluxes , expression of a large suite of defense-related genes,

cell wall remodeling and stomatal closure[18-23]. Activation of PTI prior to an infection can
also result in enhanced pathogen resistance[24,25].

Age-related resistance (ARR) is a widely observed phenomenon in plants in
which young plants exhibit greater disease susceptibility compared to older plants[26,27].
This is observed across many flowering plants against a variety of pathogens[28]. In
Arabidopsis, for instance, the basal susceptibility of young plants to foliar bacterial
pathogen *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000 is greater compared to older
plants[29]. One hypothesis to explain ARR involves the growth-defense tradeoff concept:
in order to balance resource allocations during vigorous vegetative growth early in life,
young plants prioritize growth over defense[30,31]. Indeed, there is evidence of direct
molecular connections between plant growth and immunity[32-34] including common dual-
function signaling components as in the case of PTI and brassinosteroid-dependent
plant growth[35]. However, it is unclear whether molecular connections such as these are
a sole basis for ARR in plants. In the animal kingdom, development of gnotobiotic
animals such as germ-free mice led researchers to discover an important contribution of
endogenous microbiota in postnatal maturation of innate immune responses in newborn
animals[36,37]. This raises the possibility that plant microbiota may also contribute to the
maturation of plant immunity. However, it remains an open question whether age-
dependent immunity is entirely intrinsic to plant development or whether maturation of
PTI is, in part, the result of colonization of a microbiota. Furthermore, in animals,
presence of dysbiotic microbial communities can be linked to exaggerated immune
responses, which have debilitating clinical consequences[38]. It is not clear if dysbiotic
microbial communities in plants are associated with overactive immune responses.
Addressing these basic microbiome questions requires the development of proper
gnotobiotic plant growth systems and establishment of well characterized normal
(eubiotic) and dysbiotic microbial communities.

In a recent study, we reported two peat-based gnotobiotic plant growth systems,
FlowPot and GnotoPot[39], and two synthetic bacterial communities, a eubiotic community
from healthy *Arabidopsis* leaves and a dysbiotic community from leaves of the
Arabidopsis min7 fls2 efr cerk1 (*mfec*) quadruple mutant, which lacks the ability to
maintain a eubiotic endophytic bacterial community[10]. Here, we employed these tools to

address the questions regarding the role of the endogenous microbiome in the

development of ARR and a possible role of eubiosis in gating proper plant basal

immunity.

103   Results

104   Age-dependent PTI in conventionally grown plants

We began this project by characterizing possible maturation of PTI over time in

conventionally grown *Arabidopsis* plants. For this purpose, we performed the classical

flg22 protection assays using 2.5-week-old and 3.5-week-old *Arabidopsis* plants, which

were conventionally grown in a potting soil substrate in air-circulating growth chambers.

In 2.5-week-old plants, we observed a modest level of flg22-mediated resistance to

virulent *Pst* DC3000 in flg22-treated plants, compared to mock-treated plants. Older,

3.5-week-old plants, however, exhibited a significantly enhanced level of flg22-triggered

resistance, compared to 2.5-week-old plants (Fig. 1a). This result demonstrated age-

dependent development of PTI in soil-grown *Arabidopsis* plants, which is consistent with

a recent study showing that FLS2-dependent immunity increased in the first 6 days of

young seedling growth in agar medium without microbes[40].

117   Age-dependent PTI maturation requires microbiota colonization

Traditionally, age-related resistance has been attributed to developmental

transition processes[41]. We examined an additional hypothesis that the endogenous

microbiome might be involved in age-dependent PTI in plants. For this purpose, we

investigated the temporal maturation of flg22-mediated resistance in peat-based

gnotobiotic plant growth systems. Holoxenic (HO) plants colonized with a natural, soil-

derived microbial community ('MSU', which was harvested from agricultural soil located

at Michigan State University, East Lansing, Michigan; see Methods) and corresponding

axenic (AX) plants, which were mock-inoculated with autoclaved 'microbial community

derived from the same 'MSU' soil, were used. As shown in Fig. 1b, HO plants exhibited

progressively more robust flg22-mediated resistance against *Pst* DC3000 over time,

which is consistent with age-dependent PTI observed in plants grown conventionally in

potting soil (Fig. 1a). In contrast, AX plants mock-inoculated with the autoclaved

microbial community were greatly reduced in age-dependent flg22-mediated resistance

phenotype (Fig. 1c). *Arabidopsis* mutant *bak1-5 bkk1-1 cerk1* (*bbc*[42]; which is defective

in PTI signaling downstream of multiple PRRs, including the flg22 receptor FLS2, did

not show flg22-mediated resistance in HO plants at any age (Supplementary Fig. 1),

suggesting that microbiota-mediated age-dependent resistance requires canonical PTI

signaling co-receptors.

Next, we quantified induction of the PTI marker gene *FLG22-INDUCED*

RECEPTOR-LIKE KINASE 1 (*FRK1*) to further characterize age-dependent activation of

PTI in HO plants and apparent lack thereof in AX plants. While basal expression of

FRK1 was similar for both 3.5- and 5.5-week-old HO plants (Fig. 1d), flg22 induced a

higher level of *FRK1* expression in old HO plants than in younger HO plants (Fig. 1e).

Interestingly, basal expression of *FRK1* was lower in AX plants compared to either

young or old HO plants (Fig. 1d) and, notably, no significant age-dependent increase in

flg22-induced *FRK1* expression was observed in AX plants (Fig. 1e). Thus, the reduced

age-dependent maturation of PTI in AX is correlated with a lack of robust increase in

age-dependent expression of *FRK1* gene.

Axenic plants lack normal expression of immune-associated genes

To capture genome-wide gene expression in AX and HO plants beyond the

FRK1 marker gene, we conducted transcriptome analysis of AX and HO *Arabidopsis*

plants grown in the peat gnotobiotic system. To reduce possibility of community-specific

bias due to use of a single microbiota, microbial communities harvested from two

distinct soils were used: 'MSU', which was harvested a Alfisol soil type, and 'Malaka',

which was harvested from undisturbed grassland soil in Malaka Township, Iowa (see

Methods) and is a Mollisol soil. Principal component analysis (PCA) of RNA-seq gene

expression data revealed distinct expression patterns between HO plants and AX plants

(PC1, 26% variance; Fig. 2a). Using a $|\log_2 FC| \geq 1$ and false discovery rate (FDR) <

0.05 cutoff, we identified a total of 435 differentially expressed genes (DEGs) between

HO and AX plants across both microbiota inputs: 352 were depleted in AX plants and

83 were enriched in AX plants (Fig. 2b,c and Supplementary Table 1). Of the 352 DEGs

depleted in AX plants, 138 were depleted irrespective of the microbiota input source

(i.e., enriched in both HO plants colonized by the 'MSU' community and HO plants

colonized by the 'Malaka' community; Fig. 2d). Gene ontology (GO) term enrichment

analysis of these 138 'core' AX-depleted genes revealed an overrepresentation of terms

involved in plant immunity (Fig. 2e and Supplementary Table 2). The genes enriched in

AX plants did not display any significant GO term enrichment. Closer examination of

depleted DEGs in AX plants revealed numerous genes involved in PTI, SA-mediated

defense, and defense-associated metabolite biosynthesis (Fig. 2c and Supplementary

Table 1). These genes included *FRK1*; several leucine-rich repeat protein kinases such

as *IMPAIRED OOMYCETE SUSCEPTIBILITY 1* (*IOS1*), *AT1G51890, AT1G51790,*

AT1G51860 and *AT5G59680*; systemic immunity-associated genes *AZELAIC ACID*

INDUCED 1 (*AZI1*) and *AZI3; PATHOGENESIS RELATED 2* (*PR2*) and *PR4;*

glucosinolate biosynthesis genes such as *FAD-LINKED OXIDOREDUCTASE* (*FOX1*)

and the cytochrome P450 monooxygenases *CYP71A12 and CYP71B15*; and defense-

associated transcription factors *MYB15* and *WRKY 30* (Fig 2c and Supplementary

Table 1). Thus, consistent with the targeted *FRK1* gene expression analysis shown in

Fig. 1d, results from the transcriptome analysis using two independent soil-derived

microbiotas pointed to a broadly depleted PTI/SA defense gene expression in AX plants

compared to HO plants, which collectively contribute to induced and basal innate

immunity.

Axenic *Arabidopsis* is underdeveloped in PTI

In addition to depleted immune gene expression, we found AX plants exhibited

significantly lower levels of other PTI-associated immune responses compared to HO

plants. For example, 4.5-week-old AX plants exhibited significantly reduced flg22-

induced ROS production compared to HO plants both in the magnitude of maximum

ROS production (peak amplitude) and in the time to reach the maximum (Fig. 3a and

Supplementary Fig.2). Western blot analysis revealed that, despite possessing similar

levels of total MPK3 and MPK6 (Fig. 3b,c), less MPK was phosphorylated in AX plants

after the activation of PTI by treatment with flg22 (Fig. 3d). Although RT-qPCR analysis

consistently showed that both basal and flg22-induced expression of the *FLS2* receptor

gene is significantly reduced in AX plant leaf tissue compared to HO plant leaf tissue

(Fig. 3e), total FLS2 protein abundance was variable and only occasionally reduced in

AX plant leaves (Supplementary Fig. 3). In contrast, the co-receptor BAK1 protein was

consistently found in lower relative abundance in AX plants compared to HO plants (Fig.

3f). Additionally, quantification of the defense hormone salicylic acid (SA), which is

downstream of PTI signaling, revealed that AX plants possess lower basal levels of SA

compared to HO plants (Fig. 3g,h). Finally, AX plants were hypersensitive to infection by

the virulent foliar bacterial pathogen *Pst* DC3000, compared to HO plants (Fig. 3i).

Together, these studies demonstrate multiple compromised PTI immune phenotypes in

axenic plants.

A eubiotic leaf synthetic community (SynCom) substantially confers immunocompetence

We recently assembled a 48-member eubiotic SynCom (SynCom[Col-0]) composed

of endophytic bacteria from leaves of healthy *Arabidopsis* Col-0 plants [10]. To determine

to what extent an eubiotic SynCom derived from the leaf endosphere could restore

immunocompetence to AX plants, we compared the PTI phenotypes of Col-0 plants

grown with and without SynCom (SynCom[Col-0] vs. $MgCl_2$). Col-0 plants grown with the

'MSU' soil-derived microbiota were used as control. Again, we observed robust flg22-

induced production of ROS in HO plants inoculated with the 'MSU' soil-derived

microbiota. Interestingly, SynCom[Col-0]-inoculated plants produced similar total ROS as

HO plants (Fig. 4a and Supplementary Fig. 4). We next quantified flg22-induced *FRK1*

gene expression and observed that plants colonized by SynCom[Col-0] were restored in

basal and flg22-induced *FRK1* expression (Fig. 4b,c), which was again similar to what

was observed for HO plants (Fig. 1d,e). Taken together, these results suggest that a

leaf endosphere-derived bacterial SynCom can substantially restore immune

competence to AX plants similar to a natural soil-derived microbiota.

Impact of abiotic conditions on microbiota-mediated immune competence

During the development and optimization of the peat-based gnotobiotic system, we

noticed a correlation between levels of synthetic nutrients added during preparation and

microbiota-mediated restoration of immune competency. To systematically determine

the effects of nutrients on microbiota-mediated immunocompetence we measured flg22-

induced production of ROS in AX and HO plants along a nutrient gradient. Using the

same volume of liquid, GnotoPots were prepared with full strength Linsmaier and

Skoog[43] (1x LS) (see Methods for nutrient content), half strength (0.5x) LS and one

tenth strength (0.1x) LS. We observed a significant impact of nutrients on flg22-

mediated ROS production in HO plants. Decreasing nutrient strength significantly

increased ROS burst magnitude and shortened time to reach the maximum ROS

production (Fig.4d) in HO plants. At intermediate nutrient levels (0.5x LS), ROS burst

magnitude was moderately increased and time to reach the maximum was reduced

compared to higher (1x LS) nutrient concentrations, however the total ROS produced

was not significantly different (Fig. 4d and Supplementary Fig. 5). At low nutrient levels

(0.1x LS), ROS burst magnitude was increased, time to maximum shortened, and total

ROS increased (Fig. 4d and Supplementary Fig. 5). Nutrient concentration did not have

an effect on the timing of ROS production in AX plants, and only a marginal impact on

total ROS production was observed. Together, these findings clearly illustrated a

tripartite interaction between immunity, microbiota, and environment during microbiota-

mediated maturation of flg22-triggered immunity.

Dysbiotic microbiota overstimulates immune gene expression

Several recent reports have begun to show an important contribution of plant

immunity, including PTI and vesicular trafficking pathways, to maintaining microbiota

homeostasis in Arabidopsis leaves[1,10,11]. In particular, we were able to establish two

parallel leaf endosphere-derived bacterial SynComs: 48-member SynCom[Col-0] derived

from healthy Col-0 leaves and 52-member SynCom[mfec] derived from dysbiotic *mfec*

mutant leaves[10]. To investigate the impact of a normal (eubiotic) microbiota vs. a

dysbiotic microbiota on plant immunity, we examined the expression of several

immunity-associated marker genes (*FRK1*, *PR1* and *CYP71A12*) in plants colonized

with SynCom[mfec] or SynCom[Col-0] in comparison to AX plants in plate-based gnotobiotic

system. We found an intriguing gradient of expression of these genes, with the highest

expression observed in Col-0 plants colonized by SynCom^{mfec}, an intermediate level in

SynCom^{Col-0}-colonized plants and the lowest level in AX plants (Fig. 5a-c).

To gain a better understanding of plant transcriptional responses to eubiotic

microbiota vs. a dysbiotic microbiota, we performed RNA-seq analysis of Col-0 plants

colonized by SynCom^{mfec} and SynCom^{Col-0} grown in parallel in the GnotoPot system.

Colonization with SynCom^{Col-0} compared to SynCom^{mfec} resulted in 774 DEGs ($|\log_2FC|$

> 1 and FDR < 0.05) (Fig. 5d). GO term analysis of the 609 DEGs upregulated upon

colonization with SynCom^{mfec} vs. SynCom^{Col-0} showed an over-representation of GO

terms associated with biotic stress and immunity (Fig. 5e). Additionally, several

immunity pathways including the systemic acquired resistance, PTI signaling and

glucosinolate biosynthetic processes were upregulated. Further analysis showed that

several dysbiosis-associated genes were involved in pathogenesis-related processes

during biotic stresses which are associated with immunity, cell death and its regulation

(Fig. 5f). Collectively, our results showed that dysbiotic SynCom^{mfec} overstimulates

immune gene expression compared to eubiotic SynCom^{Col-0}.

269 Discussion

Here, we show that *Arabidopsis* plants grown without exposure to a microbiota

are greatly compromised in age-dependent immunity that occurs in plants colonized

naturally by microbiota. Axenically-grown plants exhibit significant defects in PTI and

are hypersusceptible to infection by a foliar bacterial pathogen. We also show that

immunocompetence can be restored by natural soil-derived microbiota as well as,

surprisingly, a 48-member eubiotic bacterial synthetic community (SynCom^{Col-0}) derived

from leaf endophytic bacteria. In contrast, a 52-member dysbiotic synthetic community

derived from leaf endophytic bacteria overstimulates immune gene expression. Finally,

our results show that the immune-modulation function of microbiota can be influenced

by environmental conditions. Together, these results have significant implications in the

formulation of a framework for explaining age-dependent immunity, microbiota-immunity

interplay and 'immunity-microbiome-environment' tritrophic interactions in plants.

With respect to age-dependent immunity, a previous study characterized the

ontogeny of flg22-triggered immunity in very young *Arabidopsis* seedlings (within six

days after germination) in axenic nutrient agar plates[40,44], providing insight into the

developmentally controlled maturation of immune responses immediately after

germination. Results presented here, however, provide evidence that flg22-triggered

immunity exhibits an age-dependent maturation period that extends through at least the

first 2-3 weeks of vegetative growth and that full-scale age-dependent immune

maturation requires exposure to microbiota. Importantly, microbiota-colonized HO plants

in peat-based gnotobiotic systems developed age-dependent PTI over time, mirroring

plants grown conventionally in potting soil. In contrast, development of age-dependent

PTI was greatly reduced in AX plants. The microbiota-mediated age-dependent

maturation bears striking conceptual parallels to that observed in germ-free mice in

which an important contribution of endogenous microbiota in postnatal maturation of

mammalian innate immunity is well recognized[36,37]. While ARR has typically been

proposed to be caused by developmental processes that antagonize immune

responses[41], results presented here revealed that microbiota-assisted immune

maturation is a previously unrecognized contributor that plays an important role in age-

dependent immune maturation in plants.

It should be pointed out that the discovery of a causal role of microbiota in age-

dependent immune maturation required the use of a gnotobiotic system capable of

growing plants with or without a natural or synthetic community of microbes. Because

agar plate, a commonly used gnotobiotic system, is not ideal for natural colonization of

plants by a complex microbial community due to artificial overgrowth of some microbes,

this has been achieved in this study by using FlowPot and GnotoPot gnotobiotic

systems with a peat-based substrate, which partially simulates the natural soil

substrate. We used FlowPots and GnotoPots interchangeably and some initial

experiments were repeated using both systems with similar results. For most

subsequent experiments we used GnotoPots, because they allowed plants to grow for a

longer duration compared to FlowPots[39]. An important realization during this study is

that peat-based plant gnotobiotic systems can be fine-tuned to simulate a range of

various abiotic conditions, such as nutrients. This was exciting to us because many of

the microbiota functions in nature seem to be context dependent. For example, excess

nutrients[45,46] have been shown to inhibit immune signaling in plants grown in

conventional potting soils. However, it was not known if immune-suppressing activities
of high nutrients is mediated in part by microbiota. In this study, fine-tuning the nutrient
conditions of GnotoPots enabled us to discover that high nutrient-mediated immune
suppression was most obvious in the presence of microbiota, suggesting an intricate
interplay between plant, microbiota, and nutrient conditions.

Recent studies began to illustrate the importance of immunity-microbiome
interplays in plants. For example, we and others have recently shown that PTI-
associated PRRs and ROS-generating RBOHD/F are essential components of a plant
genetic network in configuring a normal eubiotic leaf microbiota to prevent health-
damaging dysbiosis[1,10,11]. Similarly, bacterial members of both leaf and root microbiotas
either stimulate or suppress PTI-associated gene expression[47,48]. In this study, we
found that a synthetic community composed of 48 culturable *Arabidopsis* phyllosphere
bacteria (SynCom[Col-0]) was sufficient to restore immunocompetence in the leaves of AX
plants at a level similar to that conferred by a natural soil-derived microbial community.
This seems remarkable considering that most members of the leaf bacterial microbiota
live on the surfaces and less than 5% of leaf bacteria reside inside the leaves[10]. Results
presented here either suggest the importance of endophytic leaf bacteria in maturing
immune responses in *Arabidopsis* leaves or there are multiple functionally redundant
sub-communities of any given microbiota and each subcommunity can independently
confer immune maturation to plants.

Another important implication of the findings from this study is that not only do
Arabidopsis plants require a microbiota to properly develop PTI, but also that the
composition of the microbiota is important. We found that SynCom[Col-0], a eubiotic
microbiota derived from healthy *Arabidopsis* leaves, was sufficient to restore
immunocompetence to AX plants. In contrast, SynCom[mfec], a dysbiotic microbiota
derived from leaves of the *Arabidopsis mfec* quadruple mutant, overstimulated immune
gene expression (Fig. 5). This observation suggests that a healthy, eubiotic microbiota
is necessary to properly gate the plant immune system. We think that this is an
important observation because in human-microbiome interactions dysbiosis is
associated with autoimmune ailments such as inflammatory bowel disease, diabetes,
allergies and other health issues[49,50]. Thus, an intimate interplay between immunity and

microbiota appears to be core to host-microbiome interactions in both animal and plant

kingdoms. Deviations from a eubiotic microbiota could result in immunodeficiency (as in

the case of AX plants) or immune-overstimulation (as in the case of SynCom[mfec]-

inoculated plants). Thus, a eubiotic microbiota has a fundamental role in gating plant

immune response during growth and development.

Methods

Arabidopsis growth conditions

The following *Arabidopsis thaliana* genotypes were used in this study: Col-0,

bak1-5 bkk1-1 cerk1 mutant (*bbc*)[42]. Conventionally grown plants were grown using

potting soil composed of equal parts Suremix (Michigan Grower Products), medium

vermiculite, and perlite. The resulting potting soil was autoclaved once to eliminate

pests. Plants were grown in an air-circulating growth chamber with the following

conditions: 60% RH, 22°C, 12 h day/12 h night photoperiod cycle and provided with a

daytime photon flux of ~90-100 µmol m^{-2} s^{-1} and supplemented with 0.5x Hoagland

nutrient solution[51] as needed.

For peat-based gnotobiotic experiments, plants were grown in FlowPots or

GnotoPots. Methods for preparation and inoculation are described in[39]. Nutrients were

supplemented with buffered 0.5x LS liquid media (pH 5.7) (Caisson Labs), unless

indicated otherwise. Full strength LS contains 1900 mg/L KNO_3, 1650 mg/L NH_4NO_3,

332.2 mg/L $CaCl_2$, 200 mg/L MES buffer, 180.7 mg/L $MgSO_4$, 170 mg/L KH_2PO_4, 100

mg/L myo-inositol, 98 mg/L $KHCO_3$, 37.26 mg/L EDTA, 27.8 mg/L $FeSO_4 \cdot 7H_2O$, 16.9

mg/L $MnSO_4 \cdot H_2O$, 8.6 mg/L $ZnSO_4 \cdot 7H_2O$, 6.2 mg/L H_3BO_3, 0.83 mg/L KI, 0.4 mg/L

thiamine HCl, 0.25 mg/L $Na_2MoO_4 \cdot 2H_2O$, 0.025 mg/L $CoCl_2 \cdot 6H_2O$, and 0.025 mg/L

$CuSO_4 \cdot 5H_2O$. Soil for natural microbiota inoculation was harvested from a Miscanthus

plot at Michigan State University (42.716989, -84.462711; 'MSU' microbiota input). For

the transcriptome experiment using HO communities, a second natural microbiota input

was obtained from soil harvested from an undisturbed grassland in Malaka Township,

IA (41.836100, -93.007800; 'Malaka' microbiota input). For natural community

microbiota experiments, AX plants were mock-inoculated with an autoclaved soil slurry

(50 g soil/ L water) and HO plants were inoculated with the same unautoclaved soil

slurry. For experiments using synthetic communities, plants were inoculated as

previously described[10]. Briefly, individual microbiota members were cultured individually

on individual R2A (Sigma) plates before being pooled together in equal ratios (OD_{600}) in

10 mM $MgCl_2$. For GnotoPot assays bacterial suspensions were adjusted to a final

$OD_{600} = 0.04$ (~2×10^7 cfu/mL) and 1 mL was used to inoculate each GnotoPot. For

plate based assays 2 uL of bacterial suspension with final $OD_{600} = 0.01$ (~5×10^6

cfu/mL) was spotted directly onto seeds. AX plants for synthetic community experiments

were mock-inoculated with an equal volume of 10 mM $MgCl_2$.

Bacterial infection assays

For flg22 protection assays with conventional potting soil-grown *Arabidopsis*,

plants of the indicated ages were hand-infiltrated using a blunt-end syringe with 500 nM

flg22 and allowed to dry until no longer water-soaked in appearance. 16-24 h after

pretreatment with flg22 leaves were infiltrated with 5×10^7 cfu/mL *Pst* DC3000 using a

blunt-end syringe. Infected plants were partially covered with a clear plastic dome to

increase humidity. Bacterial populations were determined 24 h after infiltration.

For flg22 protection assays in gnotobiotic *Arabidopsis*, plants were grown in

FlowPots with 0.5x LS. Sow date was staggered, and plants of the indicated ages were

treated at the same time to allow direct comparison. Plants were pretreated with 100 nM

flg22 using a blunt-end syringe and allowed to dry until no longer water-soaked in

appearance. Control and flg22-treated plants were kept in microboxes with the lid on

overnight. Twenty-four h after flg22 pretreatment, plants were syringe-infiltrated with *Pst*

DC3000 at 1×10^6 cfu/mL. Infected plants were allowed to dry until no longer water-

soaked in appearance and then covered with a clear plastic dome to maintain high

humidity. Bacterial populations were determined 24 h after infiltration.

For disease assays (without flg22 pretreatment) in gnotobiotic *Arabidopsis*,

plants were grown in FlowPots with 0.5x LS and hand-infiltrated with *Pst* DC300 at

1×10^6 cfu/mL. Infected plants were allowed to dry then covered with a dome to increase

humidity. Bacterial populations were determined three days after infiltration.

Transcriptome analysis

For transcriptome experiments with natural community inputs, total RNA was extracted from whole rosettes of FlowPot-grown *Arabidopsis* inoculated with 'MSU' 'Malaka') soil-derived input microbiota or in the case of AX plants, mock-inoculated with a corresponding input microbiota that had been autoclaved. A biological replicate is defined as a pool of eight rosettes collected from four FlowPots within the same microbox. Three biological replicates per condition were collected, totaling six holoxenic and six axenic replicates. RNA was extracted using the RNeasy Plant Mini Kit (Qiagen) according to manufacturer's protocol with optional on-column DNase digestion. Purified RNA was eluted in TE buffer (Tris-HCl 10 mM, pH 7.5, EDTA 1 mM). RNA concentrations were determined using an ND-1000 NanoDrop spectrophotometer (Thermo Scientific) or by Qubit RNA HS fluorometric assay (Thermo Fisher). Total RNA samples were collected in 2.0 ml nucleic acid LoBind tubes (Eppendorf) and stored at -80°C. RNA was checked for quality using a Bioanalyzer 2100 (Agilent), and all samples were determined to have an RNA integrity (RIN) score of six or greater. Stranded sequencing libraries were prepared using the NuGEN Ovation RNA-SEQ System for Model Organisms (*Arabidopsis*) according to manufacturer's protocol (NuGEN). Library preparation and sequencing was performed by the Michigan State University Research Technology Service Facility (RTSF). Sequencing was performed on the HiSeq 2500 (Illumina) with a 1 x 50 bp single read stranded format using Illumina HiSeq SBS reagents (version 4). Base calling was done by Illumina Real Time Analysis (RTA version 1.18.64).

For transcriptome experiments with SynComs, plants were grown in GnotoPots under long day (16 h day/8h night) condition and sampled at day 26 after germination. RNA extraction was done as described above but samples were eluted in RNase/DNase-free water. RNA quality controls were performed using the Qubit (Thermo Fisher Scientific) and TapeStation (Agilent Technologies, Inc.). Stranded RNA-seq libraries were pooled and sequenced on the Illumina NovaSeq 6000 S1 to obtain 50-bp pair-end reads. Library preparation and sequencing was performed by the Sequencing and Genomic Technologies Core at Duke University's Center for Genomic and Computational Biology.

Raw transcriptome reads for both transcriptome experiments were processed on

the Duke Compute Cluster as follows: Read quality control was performed using

FastQC[52], adapter trimming and sequence mapping was achieved using Trimmomatic[53]

and STAR version 9.3.0[54]. Gene expression was quantified using the R package

Rsubreads version 2.8.2[55] . Differentially expressed genes (DEGs) were identified using

the R package DESeq2[56]. Read transformation and normalization for PCoA and

clustering of was done using the EdgeR package on the iDEP platform (version 1.0)[57].

Genes with differential expression were selected using $|\log_2 FC| > 1$ and FDR < 0.05

(calculated using default DESeq2 settings based on Benjamini–Hochberg corrected

Wald test) as selection criteria and GO analysis was performed using ShinyGO version

0.76.2[58] with an FDR cutoff of 0.05 and 4 gene per group selection criteria.

ROS burst assay

Leaf disks (4 mm in diameter) were taken from the center of leaves from plants of

various ages indicated and floated with abaxial side down in wells of a white 96-well

plate containing 200 µL sterile water in each well. Plates were covered with foil and leaf

disks were kept in sterile water overnight to attenuate wounding response. Twenty-four

h later, water was removed from wells and replaced with 100 µL of an immune-eliciting

solution containing 34 µg/mL luminol (Sigma), 20 µg/mL horseradish peroxidase

(Sigma), and 100 nM flg22. Luminescence was measured (total photon counting) over

60 min immediately after the addition of immune-eliciting solution using a SpectraMax L

microplate reader (Molecular Devices). Total ROS was calculated in Prism (GraphPad)

using 'Area under curve' analysis.

RT-qPCR analysis gene expression

For RT-qPCR analysis, plants of the indicated ages were treated with a foliar

spray of elicitor solution consisting of 100 nM flg22, 0.1% DMSO, and 0.025% Silwet-

L77 (Bioworld) or a mock solution that lacked flg22. Foliar sprays were applied ensuring

treatment solution came in contact with both the adaxial and abaxial side of leaves. For

gene expression analysis of plate grown plants, full rosette of 16-days old seedlings

were snipped and tissue from 5 plants were pooled to constitute one biological replicate.

Upon harvest, the above-ground tissue was excised and transferred to 2 mL screw-top

tubes before being frozen in liquid N_2 and stored at -80°C until further processing.

Total RNA was extracted from leaf tissues using either Trizol (Thermo Fisher)

and a Direct-zol RNA extraction kit (Zymo Research) or an RNeasy Plant Mini Kit

(QIAGEN) according to the manufacturer's instructions using the optional on-column

DNase treatment. cDNA synthesis was accomplished in 10 µL volumes with

SuperScript IV VILO master mix (Thermo Fisher) or M-MLV Reverse Transcriptase

(Thermo Fisher) according to the manufacturer's instructions using 640-1000 ng total

RNA as input. Upon synthesis, cDNA was diluted 10-fold and qPCR was performed in

duplicate on a minimum of three biological replicates in 10 µL reaction volumes

containing 5 µL SYBR Green PCR master mix, 0.25 µL of each primer, and 2 µL of

template cDNA on an ABI 7500 Fast (Applied Biosystems) or QuantStudio 3 (Applied

Biosystems) RT-qPCR system using the default settings. *PP2AA3* was used for

normalization. The primer sets used to quantify gene expression in this study are listed

in Supplemental Table 5.

Quantification of salicylic acid (SA) and glucosylated salicylic acid (SAG)

Plant hormones SA and SAG were extracted as previously described[59]. In brief,

100 mg of leaf tissue harvested from 4.5-week-old plants grown in FlowPots was frozen

and ground to fine powders with a TissueLyser (Qiagen) using two 45 s cycles at 28 Hz.

Frozen powders were resuspended in 1 mL extraction buffer containing 80% methanol,

0.1% formic acid, 0.1 mg/mL butylated hydroxytoluene, and 100 nM deuterated abscisic

acid (ABA-$^2H_6$) in water. Samples were extracted overnight at 4°C with gentle agitation.

The next day, samples were cleared by centrifugation at 12,000×*g* for 10 minutes,

filtered through a 0.2 µm PTFE membrane (Millipore), and transferred to autosampler

vials. Ten µL injections of prepared extracts were separated using an Ascentis Express

fused-core C18 column (2.1×50 m, 2.7 µm) heated to 50°C on an Acquity ultra

performance liquid chromatography system (Waters Corporation). A gradient of 0.15%

formic acid in water (solvent A) and methanol (solvent B) was applied over 2.5 minutes

at a flow rate of 0.4 mL/min. Separation consisted of a linear increase from A:B (49:1) to

100% B. Transitions from deprotonated molecules to characteristic product ions were

monitored for ABA-^{2}H$_6$ (m/z 269.1 > 159.1), SA (m/z 137.0 > 93.0), and SAG (m/z 299.1

> 137.0) on a Quattro Premier tandem mass spectrometer (Waters Corporation) in

negative ion mode. The capillary voltage, cone voltage, and extractor voltage were 3500

V, 25 V and 5 V, respectively. The flow rates were 50 L/h for the cone gas (N$_2$) and 600

L/h for the desolvation gas (N$_2$). ABA-^{2}H$_6$ served as the internal standard for hormone

quantification. Collision energies and source cone potentials were optimized using

QuanOptimize software. Masslynx v4.1 was used for data acquisition and processing.

Peaks were integrated and the analytes quantified based on standard curves

normalized to the internal standard.

Immunoblot analysis

Protein was extracted from leaves as previously described [17] with slight

modification. First, frozen leaf tissues were ground to fine powders with a TissueLyser

(Qiagen) using two 45 s cycles at 28 Hz. Powders were taken up into a protein

extraction buffer containing 50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 10% (v/v) glycerol,

1% (v/v) IGEPAL CA-630 (NP-40) (Sigma), 0.5% (w/v) sodium deoxycholate, 1

Complete EDTA-free Protease Inhibitor tablet (Roche) and incubated on ice for 15 min

with periodic inversion. Leaf lysates were cleared by centrifugation at 10,000×g for 5

min and total protein normalized via Bradford Assay (Biorad). Extracts were prepared

for SDS-PAGE with a 5x loading buffer containing 10% (w/v) sodium dodecyl sulfate,

20% glycerol, 0.2 M Tris-HCl (pH 6.8), and 0.05% bromophenol blue and gradually

denatured on a thermocycler using the following sequence: 37°C for 20 min, 50°C for 15

min, 70°C for 8 min, and 95 for 5 min. Protein was subsequently separated on NuPAGE

4-12% bis-tris gels (Thermo Fisher) for 2.5 h using 100 V. Proteins were then

transferred to PVDF using an iBlot 2 dry blotting system (Thermo Fisher), blocked in 3%

milk + 2% BSA and immunoblotted overnight at 4°C with antibodies specific to

Arabidopsis FLS2 (Agrisera), BAK1 (Agrisera), MPK3 (Sigma) or MPK6 (Sigma) at

concentrations recommended by the manufacturer. Blots for detecting phosphorylated

MAPK were blocked in 5% BSA and immunoblotted with a phosphor-p44/42 MAPK

(Erk1/2) (Thr202/Tyr204) antibody (Cell Signaling). Horseradish peroxidase-conjugated

anti-rabbit antibody produced in goat (Agrisera) was used as a secondary antibody and

the resulting proteins of interest were visualized with SuperSignal West

chemiluminescent substrate (Thermo Fisher). Ponceau S or Amido Black staining was

performed to verify equal loading.

536   Data availability

537   The RNA-seq raw sequencing and analyzed data have been deposited in the NCBI

538   Gene Expression Omnibus database under accession GSE218961 and GSE218962.

540   Code availability

541   The code used for RNA-seq raw data analysis can be found at:

542   https://github.com/rsohrabi/MIP_ms

544   References

1   Paasch, B. C. & He, S. Y. Toward understanding microbiota homeostasis in the

plant kingdom. *PLoS Pathogens* **17**, e1009472 (2021).

2   Sohrabi, R., Paasch, B. C., Liber, J. & He, S. Y. Phyllosphere microbiome.

Annual Review of Plant Biology **Forthcoming** (2022).

3   Bulgarelli, D., Schlaeppi, K., Spaepen, S., Van Themaat, E. V. L. & Schulze-

Lefert, P. Structure and functions of the bacterial microbiota of plants. *Annual*

Review of Plant Biology **64**, 807-838 (2013).

4   Lundberg, D. S. *et al.* Defining the core *Arabidopsis thaliana* root microbiome.

Nature **488**, 86-90 (2012).

5   Müller, D. B., Vogel, C., Bai, Y. & Vorholt, J. A. The plant microbiota: systems-

level insights and perspectives. *Annual Review of Genetics* **50**, 211-234 (2016).

6   Regalado, J. *et al.* Combining whole-genome shotgun sequencing and rRNA

gene amplicon analyses to improve detection of microbe–microbe interaction

networks in plant leaves. *The ISME journal* **14**, 2116-2130 (2020).

7   Lindow, S. E. & Brandl, M. T. Microbiology of the phyllosphere. *Applied and*

environmental microbiology **69**, 1875-1883 (2003).

8   Bai, Y. *et al.* Functional overlap of the Arabidopsis leaf and root microbiota.
Nature **528**, 364-369 (2015).

9   Fitzpatrick, C. R. *et al.* The plant microbiome: from ecology to reductionism and
beyond. *Annual Review of Microbiology* **74**, 81-100 (2020).

10   Chen, T. *et al.* A plant genetic network for preventing dysbiosis in the
phyllosphere. *Nature* **580**, 653-657 (2020).

11   Pfeilmeier, S. *et al.* The plant NADPH oxidase RBOHD is required for microbiota
homeostasis in leaves. *Nature Microbiology* **6**, 852-864 (2021).

12   Berendsen, R. L., Pieterse, C. M. & Bakker, P. A. The rhizosphere microbiome
and plant health. *Trends in Plant Science* **17**, 478-486 (2012).

13   Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T. & Singh, B. K. Plant–microbiome
interactions: from community assembly to plant health. *Nature Reviews*
Microbiology **18**, 607-621 (2020).

14   Mendes, R., Garbeva, P. & Raaijmakers, J. M. The rhizosphere microbiome:
significance of plant beneficial, plant pathogenic, and human pathogenic
microorganisms. *FEMS Microbiology Reviews* **37**, 634-663 (2013).

15   Zipfel, C. Plant pattern-recognition receptors. *Trends in Immunology* **35**, 345-351
(2014).

16   Chinchilla, D., Bauer, Z., Regenass, M., Boller, T. & Felix, G. The *Arabidopsis*
receptor kinase FLS2 binds flg22 and determines the specificity of flagellin
perception. *The Plant Cell* **18**, 465-476 (2006).

17   Chinchilla, D. *et al.* A flagellin-induced complex of the receptor FLS2 and BAK1
initiates plant defence. *Nature* **448**, 497-500 (2007).

18   Zhang, J. & Zhou, J.-M. Plant immunity triggered by microbial molecular
signatures. *Molecular Plant* **3**, 783-793 (2010).

19   Macho, A. P. & Zipfel, C. Plant PRRs and the activation of innate immune
signaling. *Molecular Cell* **54**, 263-272 (2014).

20   Couto, D. & Zipfel, C. Regulation of pattern recognition receptor signalling in
plants. *Nature Reviews Immunology* **16**, 537-552 (2016).

21   Li, B., Meng, X., Shan, L. & He, P. Transcriptional regulation of pattern-triggered
immunity in plants. *Cell Host & Microbe* **19**, 641-650 (2016).

22   Meng, X. & Zhang, S. MAPK cascades in plant disease resistance signaling.
Annual Review of Phytopathology **51**, 245-266 (2013).

23   Bigeard, J., Colcombet, J. & Hirt, H. Signaling mechanisms in pattern-triggered
immunity (PTI). *Molecular Plant* **8**, 521-539 (2015).

24   Zipfel, C. *et al.* Bacterial disease resistance in *Arabidopsis* through flagellin
perception. *Nature* **428**, 764-767 (2004).

25   Crabill, E., Joe, A., Block, A., Van Rooyen, J. M. & Alfano, J. R. Plant immunity
directly or indirectly restricts the injection of type III effectors by the
Pseudomonas syringae type III secretion system. *Plant physiology* **154**, 233-244
(2010).

26   Develey-Rivière, M. P. & Galiana, E. Resistance to pathogens and host
developmental stage: a multifaceted relationship within the plant kingdom. *New*
Phytologist **175**, 405-416 (2007).

27   Kus, J. V., Zaton, K., Sarkar, R. & Cameron, R. K. Age-related resistance in
Arabidopsis is a developmentally regulated defense response to *Pseudomonas*
syringae. The Plant Cell **14**, 479-490 (2002).

28   Panter, S. N. & Jones, D. A. Age-related resistance to plant pathogens.
Advances in Botanical Research **38**, 251-280 (2002).

29   Rusterucci, C. *et al.* Age-related resistance to *Pseudomonas syringae* pv. *tomato*
is associated with the transition to flowering in *Arabidopsis* and is effective
against *Peronospora parasitica. Physiological and Molecular Plant Pathology* **66**,
222-231 (2005).

30   Huot, B. *et al.* Dual impact of elevated temperature on plant defence and
bacterial virulence in *Arabidopsis. Nature Communications* **8**, 1-12 (2017).

31   He, Z., Webster, S. & He, S. Y. Growth–defense trade-offs in plants. *Current*
Biology **32**, R634-R639 (2022).

32   Wang, D., Pajerowska-Mukhtar, K., Culler, A. H. & Dong, X. Salicylic acid inhibits
pathogen growth in plants through repression of the auxin signaling pathway.
Current Biology **17**, 1784-1790 (2007).

33 Yang, D.-L. *et al.* Plant hormone jasmonate prioritizes defense over growth by
interfering with gibberellin signaling cascade. *Proceedings of the National*
Academy of Sciences **109**, E1192-E1200 (2012).

34 Lozano-Durán, R. *et al.* The transcriptional regulator BZR1 mediates trade-off
between plant innate immunity and growth. *eLife* **2**, e00983 (2013).

35 Yang, D.-H., Hettenhausen, C., Baldwin, I. T. & Wu, J. The multifaceted function
of BAK1/SERK3: plant immunity to pathogens and responses 1 to insect
herbivores. *Plant Signaling & Behavior* **6**, 1322-1324 (2011).

36 Hooper, L. V., Littman, D. R. & Macpherson, A. J. Interactions between the
microbiota and the immune system. *Science* **336**, 1268-1273 (2012).

37 Fulde, M. & Hornef, M. W. Maturation of the enteric mucosal innate immune
system during the postnatal period. *Immunological Reviews* **260**, 21-34 (2014).

38 Levy, M., Kolodziejczyk, A. A., Thaiss, C. A. & Elinav, E. Dysbiosis and the
immune system. *Nature Reviews Immunology* **17**, 219-232 (2017).

39 Kremer, J. M. *et al.* Peat-based gnotobiotic plant growth systems for *Arabidopsis*
microbiome research. *Nature Protocols* **16**, 2450-2470 (2021).

40 Zou, Y., Wang, S. & Lu, D. MiR172b-TOE1/2 module regulates plant innate
immunity in an age-dependent manner. *Biochemical and Biophysical Research*
Communications **531**, 503-507 (2020).

41 Carella, P., Wilson, D. C. & Cameron, R. K. Some things get better with age:
differences in salicylic acid accumulation and defense signaling in young and
mature Arabidopsis. *Frontiers in Plant Science* **5**, 775 (2015).

42 Xin, X.-F. *et al.* Bacteria establish an aqueous living space in plants crucial for
virulence. *Nature* **539**, 524-529 (2016).

43 Linsmaier, E. & Skoog, F. Organic growth factor requirements of tobacco tissue
cultures. *Physiologia Plantarum* **18** (1965).

44 Zou, Y. *et al.* Transcriptional regulation of the immune receptor FLS2 controls the
ontogeny of plant innate immunity. *The Plant Cell* **30**, 2779-2794 (2018).

45 Jeworutzki, E. *et al.* Early signaling through the Arabidopsis pattern recognition
receptors FLS2 and EFR involves Ca2+-associated opening of plasma
membrane anion channels. *The Plant Journal* **62**, 367-378 (2010).

46   Ding, S. *et al.* Nitrogen forms and metabolism affect plant defence to foliar and
653   root pathogens in tomato. *Plant, Cell & Environment* **44**, 1596-1610 (2021).

654   47   Ma, K.-W. *et al.* Coordination of microbe–host homeostasis by crosstalk with
655   plant innate immunity. *Nature Plants* **7**, 814-825 (2021).

656   48   Teixeira, P. J. *et al.* Specific modulation of the root immune system by a
657   community of commensal bacteria. *Proceedings of the National Academy of*
658   *Sciences* **118** (2021).

659   49   Blum, H. E. The human microbiome. *Advances in Medical Sciences* **62**, 414-420
660   (2017).

661   50   de Oliveira, G. L. V., Leite, A. Z., Higuchi, B. S., Gonzaga, M. I. & Mariano, V. S.
662   Intestinal dysbiosis and probiotic applications in autoimmune diseases.
663   *Immunology* **152**, 1-12 (2017).

664   51   Hoagland, D. R. & Arnon, D. I. The water-culture method for growing plants
665   without soil. *Circular. California Agricultural Experiment Station* **347** (1950).

666   52   Andrews, S.   (Babraham Bioinformatics, Babraham Institute, Cambridge,
667   United Kingdom, 2010).

668   53   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
669   sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

670   54   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-
671   21 (2013).

672   55   Liao, Y., Smyth, G. K. & Shi, W. The R package Rsubread is easier, faster,
673   cheaper and better for alignment and quantification of RNA sequencing reads.
674   *Nucleic Acids Research* **47**, e47-e47 (2019).

675   56   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
676   dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 1-21 (2014).

677   57   Ge, S. X., Son, E. W. & Yao, R. iDEP: an integrated web application for
678   differential expression and pathway analysis of RNA-Seq data. *BMC*
679   *Bioinformatics* **19**, 1-24 (2018).

680   58   Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for
681   animals and plants. *Bioinformatics* **36**, 2628-2629 (2020).

682 59   Zeng, W. *et al.* A genetic screen reveals *Arabidopsis* stomatal and/or apoplastic

683   defenses against *Pseudomonas syringae* pv. *tomato* DC3000. *PLoS Pathogens*

684   **7**, e1002291 (2011).

Acknowledgements

687   We thank undergraduates Caleigh Griffin, Trevor Ulrich, Franchesca Dion, and

688 Timothy Johnson and lab technician David Rhodes for assistance in gnotobiotic system

689 design and construction and for performing critical experiments that led to these

690 published results. We thank members of the He Lab for critical reading of the

691 manuscript. S.Y.H. is an Investigator at Howard Hughes Medical Institute.

Author information

694   These authors contributed equally: Bradley C. Paasch, Reza Sohrabi, James M.

695 Kremer.

Authors and Affiliations

Department of Biology, Duke University, Durham, NC, USA

Bradley C. Paasch, Reza Sohrabi, Kinya Nomura & Sheng Yang He

Howard Hughes Medical Institute, Duke University, Durham, NC, USA

Bradley C. Paasch, Reza Sohrabi, Kinya Nomura & Sheng Yang He

Department of Energy Plant Research Laboratory, Michigan State University, East

Lansing, MI, USA

James M. Kremer & Jennifer Martz

Department of Plant Pathology, University of Georgia, Athens, GA, USA

Brian Kvitko

Contributions

J.M.K., J.M.T., and S.Y.H. conceptualized the initial project.

B.C.P., R.S., J.M.K., J.M.T., and S.Y.H. designed the study and analyzed the data.

B.C.P. performed gnotobiotic plant assays.

R.S. performed dysbiosis experiments and RNA-seq analyses.

J.M.K. performed RNA-seq in FlowPots and preliminary gnotobiotic plant assays.

K.N. performed temporal assays in conventional and gnotobiotic plants.

J.M. assisted nutrient gnotobiotic assays.

B.K. performed temporal flg22 protection assays in conventionally grown plants.

B.C.P., R.S., and S.Y.H. wrote the manuscript with input from all authors.

Corresponding author

Correspondence to Sheng Yang He.

Email: shengyang.he@duke.edu

Ethics declarations

Competing interests

The authors declare no competing interests.

Figures and figure legends

Figure 1 | Age-dependent flg22-triggered immunity in *Arabidopsis*.

a, flg22 protection assay showing enhanced resistance against *Pst* DC3000 triggered

by pretreatment with 500 nM flg22 in 2.5-week-old and 3.5-week-old plants. Each

column represents bacterial titer 24 hours after inoculation as log transformed colony

forming units (cfu)/cm^2 and is the mean of six plants. Error bars indicate SD. Different

letters represent a significant difference ($p < 0.05$, two-way ANOVA with Tukey's HSD

post-hoc test). **b**, Age-dependent flg22 protection. Axenic (AX) or holoxenic (HO) plants

were treated 24 hours prior to inoculation with *Pst* DC3000 with either a water (mock) or

100 nM flg22 solution. Each column represents bacterial titer 24 hours after inoculation

as log transformed cfu/cm^2 and is the mean of 3 plants. Error bars indicate SD. Different

letters represent a significant difference ($p < 0.05$, two-way ANOVA with Tukey's HSD

post-hoc test). **c**, Relative protection displayed as fold change in bacterial cell counts

between flg22- and mock-treated samples. Error bars represent SD. Different letters

represent a significant different ($p < 0.05$, two-way ANOVA with Tukey's Honest

Significant Difference (HSD) post-hoc test). **d**, Basal and **e**, flg22-induced age-

dependent *FRK1* gene expression in 3.5-week-old and 5.5-week-old AX and HO plants.

Total RNA was extracted 4 h after treatment with a mock solution lacking flg22 for basal

expression or 100 nM flg22 for flg22-induced expression. Expression levels displayed

as relative to mock treated 3.5-week-old HO plants for both panels. *PP2AA3* was used

for normalization. Results represent the mean values ± SD of four plants. Different

letters represent a significant difference ($p < 0.05$, two-way ANOVA with Tukey's HSD

post-hoc test). Each experiment was repeated three independent times with similar

results.

Figure 2 | Axenic *Arabidopsis* plants are depleted in the basal expression of defense-related transcripts.

a, Principal component (PCA) analysis of genes expressed under AX and HO conditions using microbial communities from two different locations/soil types ('MSU': MI, Alfisol soil type and 'Malaka': IA, Mollisol soil type; 8 plants from one microbox pooled per biological rep, n = 3). **b**, Volcano plot of differentially expressed genes (DEGs). Colored regions represent significant differential expression with $|log_2FC| > 1$ and FDR < 0.05 (Benjamini-Hochberg corrected Wald Test) with the number of genes corresponding to each group indicated in parentheses. **c**, Heat map of DEGs generated using hierarchical clustering with euclidean distance and complete linkage. Label superscript indicates community used for inoculation of HO plants or mock inoculation of AX plants. A subset of the differentially regulated genes in HO and AX is shown on right. **d**, Venn diagram of upregulated DEGs showed 138 common genes in response to HO[MSU] and HO[Malaka] treatments. **e**, Gene ontology (GO) term enrichment (GO:BP biological process) analysis on 'core' depleted DEGs in AX plants. Top enriched GO terms displayed, ranked by significance (FDR adjusted hypergeometric test).

Figure 3 | Axenic *Arabidopsis* plants exhibit defects in PTI compared to colonized plants.

a, ROS burst dynamics induced by 100 nM flg22 in AX and HO plants in GnotoPots. Results represent the mean of eight plants ± SEM. **b,c**, Representative blots of total MPK3 (b) or MPK6 (c) proteins in 4.5 week old AX and HO plants. Protein was detected with MPK3 or MPK6-specific antibodies. **d**, Representative blot of phosphorylated MPK3/6 proteins detected using an α-p44/42-ERK antibody upon treatment with 100 nM flg22. Samples were taken at the indicated times after treatment. For panels (b-d), numbers below blots indicate band intensity relative to that of Ponceau S, quantified by ImageJ. **e**, Basal and flg22-induced expression of *FLS2* gene in AX and HO plant leaf tissue. Total RNA was extracted 1 h after treatment with 100 nM flg22 or mock solution. PP2AA3 was used for normalization. Bars represent the mean values ± SD of three biological samples, each consisting of three plants. Different letters represent a significant difference ($p < 0.05$, unpaired *t*-tests with Welch's correction). **f**, Total BAK1 protein detected in leaf lysates. Numbers below blot indicates band intensity relative to that of Amido Black, quantified by ImageJ. **g,h**, Total levels of salicylic acid (SA) (f) and glucosylated SA (g) in AX and HO plants. Each bar represents the mean values ± SD of eight biological samples, each consisting of at least three plants. (SA: $p = 0.003$, SAG: p

794   = 0.002; Student's t-test). **i**, *Pst* DC3000 populations in AX and HO plants. Each column

795   represents bacterial titer 3 days after inoculation as log transformed cfu/cm^2 and is the

796   mean of three plants. Error bars indicate SD ($p = 0.0006$, Student's *t*-test). Each

797   experiment was repeated three independent times with similar results.

Figure 4 | Natural microbiota and SynCom^{Col-0} restore

immunocompetence.

a, ROS burst dynamics induced by 100 nM flg22 in plants colonized by HO or

SynCom^{Col-0}. Results represent the mean of 12 plants ± SEM. **b**,**c**, Basal (b) and flg22-

induced (c) *FRK1* expression in AX (mock-inoculated with sterile $MgCl_2$) and

SynCom^{Col-0}-inoculated plants. Total RNA was extracted 3 h after treatment with a mock

solution lacking flg22 (b) or 100 nM flg22 (c). Results relative to basal expression in

SynCom^{Col-0}-inoculated plants. *PP2AA3* was used for normalization. Bars represent the

mean values ± SD of three plants. Different letters represent a significant difference.

Asterisks represents a significant difference ($p = 0.022$, $p = 0.0006$, Student's *t*-test). **d**,

ROS burst dynamics induced by 100 nM flg22 in AX and HO plants grown in GnotoPots

supplied with 0.1x, 0.5x, or 1x LS nutrient solution concentrations. Results represent the

mean of six plants ± SEM. Each experiment was repeated a minimum of two

independent times with similar results.

Figure 5 | Dysbiotic microbiota overstimulates immune gene expression.

a-c, Basal expression of defense-related genes *FRK1* (a), *CYP71A12* (b), *PR1* (c) in AX, SynCom[Col-0]- and SynCom[mfec]-inoculated plants grown in agar plates. *PP2AA3* was used for normalization. Bars represent the mean values ± SD of three biological samples, each consisting of five seedlings. Different letters represent a significant difference ($p < 0.05$, one-way ANOVA with Tukey's HSD post-hoc test). **d**, Volcano plot of genes differentially expressed in SynCom[Col-0]- and SynCom[mfec]-colonized plants. Colored regions represent significant differential expression with $|log_2FC| > 1$ and FDR < 0.05 (Benjamini-Hochberg corrected Wald Test) with the number of genes corresponding to each group indicated in parentheses. **e**, GO term enrichment for up-regulated DEGs in SynCom[mfec]-colonized plants compared to SynCom[Col-0]-colonized plants, ranked by significance (FDR adjusted hypergeometric test). **f**, Heat map for

831   selected genes from hierarchical clustering all DEGs. Gene descriptions are listed in

832   supplementary table 3.